High-resolution detection of copy number alterations in single cells with HiScanner

Yifan Zhao[1,2], Lovelace J Luquette[1], Alexander D Veit [1], Xiaochen Wang[3], Ruibin Xi[3], Vinayak V Viswanadham[1], Yuwei Zhang[1], Diane D Shao[4,5], Christopher A Walsh[4], Hong Wei Yang[6], Mark D Johnson[6]*, Peter J Park[1]*

1 Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
2 Program in Health Sciences & Technology, Harvard Medical School & Massachusetts Institute of Technology, Boston, MA, USA
3 Department of Probability and Statistics, School of Mathematical Sciences, Peking University, Beijing, China
4 Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA
5 Department of Neurology, Boston Children's Hospital, Boston, MA, USA
6 Department of Neurological Surgery, University of Massachusetts Chan Medical School, Worcester, MA 01655

*Correspondence to: mark.johnson3@umassmemorial.org, peter_park@hms.harvard.edu

Abstract

Improvements in single-cell whole-genome sequencing (scWGS) assays have enabled detailed characterization of somatic copy number alterations (CNAs) at the single-cell level. Yet, current computational methods are mostly designed for detecting chromosome-scale changes in cancer samples with low sequencing coverage. Here, we introduce HiScanner (High-resolution Single-Cell Allelic copy Number callER), which combines read depth, B-allele frequency, and haplotype phasing to identify CNAs with high resolution. In simulated data, HiScanner consistently outperforms state-of-the-art methods across various CNA types and sizes. When applied to high-coverage scWGS data from 65 cells across 11 neurotypical human brains, HiScanner shows a superior ability to detect smaller CNAs, uncovering distinct CNA patterns between neurons and oligodendrocytes. We also generated low-coverage scWGS data from 179 cells sampled from the same meningioma patient at two time points. For this serial dataset, integration of CNAs with point mutations revealed evolutionary trajectories of tumor cells. These findings show that HiScanner enables accurate characterization of frequency, clonality, and distribution of CNAs at the single-cell level in both non-neoplastic and neoplastic cells.

Introduction

Over the past decade, advances in single-cell omics technologies have provided an opportunity to interrogate cellular heterogeneity with unprecedented resolution. Single-cell RNA-sequencing has been the most popular[1,2], but significant progress has also been made for single-cell ATAC-sequencing for profiling open chromatin[3] and single-cell Hi-C for profiling three-dimensional folding[4,5]. Another technology that has advanced is single-cell whole-genome sequencing (scWGS), which can be used to uncover cell-to-cell variation in somatic mutations that may be masked in bulk sequencing.

Various scWGS approaches have been developed for detecting copy number alterations (CNAs), primarily in cancer cells. Examples include degenerate-oligonucleotide-primed PCR (DOP-PCR)[6], direct library preparation (DLP+)[7,8], and the now-defunct 10x Genomics chromium CNV assay[9]. However, the fraction of genome covered by these approaches is low (low breadth-of-coverage) and their main application was to detect clonal patterns of chromosomal arm-sized CNAs. Single-cell whole-genome amplification (scWGA) offers high breadth-of-coverage and could improve spatial resolution, but it has been hindered by the noise introduced during amplification. Earlier scWGA methods have involved exponential or quasi-linear amplification, generating highly uneven coverage across the genome, allelic bias (in which one allele is amplified at a higher rate), and frequent allelic dropout (where only one allele is amplified). Even if the cell is sequenced at high depth, these amplification artifacts can confound CNA detection by causing large fluctuations in the read depth ratio (RDR; ratio between observed and expected read count), yielding relatively few advantages over low-coverage (<1X, often <0.1X) scWGS.

Recent advances in scWGA chemistry[10,11] and declining sequencing costs have made high-coverage (>20X) scWGS more attractive for CNA detection. In particular, scWGA by primary template-directed amplification (PTA) achieves more uniform amplification across the genome[10,12]. We have already used the high-coverage PTA data to identify somatic single nucleotide variants in the brain, tracing cellular lineage across different tissues[13], providing a more precise estimate of age-associated mutational rate[12], and characterizing the regions more prone to mutagenesis in different cells[14]. PTA data also offer a potential to improve the resolution of CNA detection beyond the usual Mb-scale[15,16].

The use of BAF is a standard feature in best bulk CNA methods such as ASCAT[17] and Battenberg[18] for inferring allele-specific copy number state. Two recently developed single-cell callers, CHISEL[19] and Alleloscope[20], leverage phasing and BAF for identifying allele-specific CNAs in low-coverage scWGS. However, Alleloscope performs segmentation on bulk samples, not single cells, limiting its ability to detect rare CNAs which are critical to understanding the genomic architecture of non-cancerous cells and especially non-dividing cells. CHISEL is not tuned for the discovery of small CNAs (e.g., <5 Mb) potentially obscuring certain biological processes. SCOVAL[21] also utilizes BAF information, but it is primarily a validation approach and not a CNA caller. Current

single-cell CNA callers are not designed to leverage high-coverage data and are tailored
to the analysis of clonal CNAs in cancer cells.
[19,22–24]

In this work, we propose HiScanner (High-resolution Single-Cell Allelic copy Number
callER) for accurate, high-resolution allele-specific CNA detection in scWGS. HiScanner
leverages reference-based phasing and local sequence bias to efficiently estimate B-allele
frequency (BAF, the relative proportions of reads covering the two alleles at
heterozygous germline single nucleotide polymorphisms, gHETs) to complement read
depth-based CNA identification. Building upon our previously developed bulk CNA
caller framework[25], HiScanner performs read depth normalization at nucleotide
resolution, followed by joint segmentation of single-cell profiles using the Bayesian
information criterion (BIC). By simultaneously modeling RDR and BAF for individual
cells, HiScanner infers cell ploidy and integer allele-specific copy number states. It also
considers assay-specific allelic dropout patterns to find an optimal bin size, a nuanced yet
crucial parameter in CNA analysis. We demonstrate that HiScanner outperforms four
representative single-cell CNA methods—CHISEL (BAF-aware) and BAF-agnostic
methods (Ginkgo, SCOPE, SCYN) [19,22–24] — in real and simulated data with greater
sensitivity and precision at small size scales (200kb-5Mb). For biological discovery, we
analyzed two datasets. Application of HiScanner to a high-coverage scWGS dataset from
eleven neurotypical human brains uncovered a large number of previously undetected
small CNAs, revealing contrasting patterns between neuronal and glial cells. We also
sequenced 179 cells at low coverage (~0.5X on average) from biopsies of an initial
(n=88) and recurrent (n=91) meningioma tumor pair. HiScanner revealed phases and
timing of tumor progression by detecting subclone-defining CNAs, which were
subsequently corroborated by an orthogonal clustering based on somatic point
mutations[26]. Lastly, we introduce ViScanner, a web-based visualization tool built using
the HiGlass architecture[27], which offers[27] users the ability to navigate genomic data with
continuous zooming, allowing for user-friendly exploration of genomic regions at various
resolutions.
116
117

Results

Overview of the algorithm

HiScanner introduces three key improvements over existing single-cell CNA detection
methodologies to enhance the resolution of CNA discovery (Figure 1). First, optimal
resolution is inferred by determining the smallest permissible bin size based on the size of
amplification artifacts, rather than leaving the tuning of this critical parameter to the user.
Second, joint segmentation is performed across cells to avoid overfitting amplification
noise. Third, an efficient BAF estimator is implemented to mitigate the effects of phase-
switch errors and allelic dropout. In addition to these features, HiScanner corrects for bias
in read depth due to local GC content using a nucleotide-level model of GC bias[25]. These
improvements are described in more detail below.
130

Selecting optimal bin size based on allelic dropout events Variations in RDR across the
genome is the primary signal used to detect copy number changes, but scWGA introduces
considerable levels of uneven amplification along the genome that masquerade as CNAs.
A prevalent form of uneven amplification is allelic dropout, where failed amplification of
one allele results in an RDR decrease similar to a copy number loss (Supplementary
Figure 1a). Previous single-cell CNA callers partitioned the genome into large bins to
dampen the effect of segments lost to amplification, but this limits the size of CNAs that
can be detected. On the other hand, decreasing the bin size unmasks RDR fluctuations,
possibly inducing false positive CNA calls. A key insight exploited by HiScanner is that
the size distribution of allelic dropout events, which is usually bounded above and is
specific to assay-dependent amplicon length, can be used to determine a balanced bin size
that maximizes detection resolution and minimizes the effects of amplification noise
(Supplementary Figure 1b). Furthermore, because dropout sizes and frequencies can
differ considerably between scWGA protocols[28] (Supplementary Figure 1c), it is
important to fit the dropout size distribution on each dataset. For example, the average
size of dropout events in two cells amplified by PTA and MDA[29] differed by two orders
of magnitude (10^3 bp versus 10^5 bp, respectively, Supplementary Figure 1d). HiScanner
employs a two-state hidden Markov model (HMM) to detect dropout events using BAF at
gHETs (Supplementary Figure 1b). Subsequently, the predicted dropout events are used
to build a sample-specific dropout size distribution and the top 5^{th} percentile of dropout
sizes is used as the smallest permissible bin size.
152
Joint segmentation across multiple cells By leveraging signals from multiple cells,
HiScanner's joint segmentation promotes breakpoint sharing, thus improving detection
sensitivity. We extended our previous strategy[25] of merging adjacent bins with similar
copy number state (as determined by the BIC) to the multi-sample case by computing a
BIC from the joint breakpoint likelihood of a breakpoint across cells (Methods). This
merging approach retains information about bin position, avoids overfitting by penalizing
excessive segmentations, and amplifies signals of clonal CNAs. Notably, the clustering
approach proposed in CHISEL to simultaneously model BAF and RDR across all cells
does not consider the proximity of bins, making the method more susceptible to calling
false-positive breakpoints, especially in regions with complex structural variations or
relatively high amplification noise.
164
Accurate estimation of the B-allele frequency (BAF) BAF can provide additional
information for CNA inference where RDR is ambiguous; a shift in the BAF from the
expected value of 0.5 in diploid regions suggests an underlying CNA (for example, a
single copy gain shifts the BAF to 1/3 or 2/3, depending on which allele the BAF is
measured against). In bulk data, obtaining allele-specific read depth at gHET loci is an
effective way to compute BAF; however, in scWGS, uneven amplification between
homologous alleles (allelic bias) and dropout can hinder this approach. Furthermore, low-
coverage sequencing–which is more typical for single-cell CNA studies–reduces the
number of reads at gHETs, thereby increasing BAF noise with its undersampling. The
effects of both allelic bias and gHET undersampling on BAF estimation can be mitigated
by summing allele-specific read depth over bins. However, this requires phasing each
gHET to a haplotype, and phase switch errors, which can occur at rates of 1-2% [30,31] for

reference-based phasing approaches, confound bin-level BAF estimation by assigning
depth to the incorrect allele. CHISEL proposed a two-step process to calculate per-bin
BAF estimates: after an initial phasing based on an external reference panel,[32,33]
expectation-maximization (EM) is used for refinement, first aggregating phased gHETs
into haplotype blocks and then performing EM refinement of phase across cells.
Alleloscope also adopted an EM approach, but it operates on unphased, individual gHETs.
Although effective, the EM step used by both CHISEL and Alleloscope is
computationally expensive and is challenging to apply to high-resolution bins. Indeed,
CHISEL operates on ~5 Mb bins (by default) and Alleloscope only computes BAF on the
final copy number segments derived from bulk or pseudobulk.
187
HiScanner computes BAF by phasing gHETs using a reference-based method followed
by a computationally efficient, two-step procedure to reduce the effect of phase-switch
errors and dropout artifacts (Supplementary Figure 2; Supplementary Note). In the first
step, LOH is detected by exploiting GC bias, a source of non-biological fluctuations in
total read depth that is treated solely as a nuisance. In a segment without LOH, GC-
induced read depth changes—which are roughly evenly split between homologous alleles
due to similar nucleotide content—lead to a subtle but detectable difference in the
distributions of BAF before and after a haplotype-specific *shift* operation (Supplementary
Note). In regions affected by LOH, this shift procedure has little effect. Thus, HiScanner
predicts the presence of LOH in each segment by a Kolmogorov-Smirnov test of the BAF
and shifted BAF distributions. In the second step, peaks (local maximums) are called in
the per-bin BAF distribution of each segment. If the segment was not predicted to contain
LOH, then any BAF peak near 0 or 1 is interpreted as dropout artifacts and removed, and
the maximum of the remaining non-zero peak is the final segment BAF. Otherwise, the
BAF is set to 0. The efficiency of this procedure is critical for CNA analysis with high
resolution bins.
204
Visualization using ViScanner To enhance the accessibility of HiScanner output, we
developed a web-based visualization platform, ViScanner (Methods). Traditional single-
cell CNA analysis tools often generate static output, including mapping coordinates for
potential CNAs and genome-wide visualizations of CNAs. However, genome-scale plots
do not adequately support the exploration of small CNAs. We developed an interactive
React-based web application, ViScanner, which leverages the HiGlass toolkit[27] to provide
a user-friendly, dynamic resolution CNA browser. ViScanner allows users to scrutinize
HiScanner output with real-time zooming, akin to the user experience in web-based
services like Google Maps. Importantly, ViScanner ensures data security by executing
JavaScript locally in the user's browser, avoiding the need to upload data, even
temporarily, to any external server.
216
Comparison with other methods
To assess HiScanner's performance, we conducted benchmarking studies using both real
and simulated data. HiScanner was compared against four widely-used single-cell CNA
methods, Ginkgo[22], SCOPE[23], SCYN[24] and CHISEL[19]. Ginkgo is a popular web-based
application that identifies CNAs based on RDR and performs segmentation via circular
binary segmentation[34]. SCOPE is a normalization and copy number estimation method

that uses a Poisson latent factor model to account for technical biases and copy number
changes; SCYN employs SCOPE's normalization and employs dynamic programming
for efficient copy number segmentation. CHISEL is a more sophisticated algorithm that
combines phasing, BAF, and RDR to identify allele-specific CNAs. Our benchmarking
analysis focused on the detection of non-clonal CNAs (private to a single cell), which are
more difficult to identify than clonal CNAs (present in a subpopulation). Consequently,
methods designed to detect only clonal events, such as Alleloscope[20], were excluded to
avoid mischaracterizing their performance (Table 1).
231
We first evaluated the three callers on detecting non-clonal CNAs using high-coverage
simulated data. Previous benchmarking studies often simulated single-cell CNAs by
manipulating read counts within pre-determined CNA locations, thereby focusing
exclusively on RDR-based CNA callers[35,36]. However, allele-specific callers cannot be
properly assessed by this strategy because simulation at the level of read counts does not
preserve the haplotype structure. Moreover, although simulating read counts using
Lorenz curve sampling[35] can recapitulate read count variability associated with GC
content, it does not faithfully capture sampling and amplification noise characteristic of
scWGS sequencing data. To overcome these limitations, we adapted the synthetic diploid
X chromosome methodology[11] to simulate somatic CNAs. In this approach, sequencing
reads sampled from two male haploid X chromosomes from different individuals are
pooled to generate a diploid chromosome (Fig 2a). By sampling reads prior to pooling, it
is possible to insert copy gain or loss events specific to a single allele without disrupting
local haplotype structure. Further, one can explicitly control phasing accuracy and phase
switch rates, while preserving scWGS amplification artifacts.
247
We simulated four types of non-clonal CNAs with sizes ranging from 200 kb to 5 Mb
using the synthetic diploid X: single-copy losses, single-copy gains, two-copy gains, and
copy-neutral loss-of-heterozygosity (cnLOHs). Our analysis of four CNA types (200 kb
to 5 Mb) revealed distinct performance patterns across tools. For single-copy losses (Fig
2b,c), HiScanner and Ginkgo excelled at 1 Mb and above, with HiScanner achieving
slightly better sensitivity (~0.90 vs 0.85) while matching Ginkgo's high precision (~0.96).
CHISEL showed moderate performance (sensitivity 0.80, precision 0.51), while SCOPE
and SCYN demonstrated limited sensitivity despite reasonable precision. In single-copy
gain detection (Fig 2d,e), HiScanner outperformed others at smaller sizes (~0.65
sensitivity at 500kb), with Ginkgo following a similar but lower trajectory. Both achieved
strong performance (>0.90) above 2Mb, while CHISEL showed moderate results.
SCOPE and SCYN performed poorly across all sizes.
For two-copy gains (Fig 2f,g), performance differences narrowed above 1Mb, with
HiScanner and Ginkgo showing nearly identical high performance (~0.85 sensitivity,
~0.90 precision at 2.5Mb+). CHISEL matched their precision for larger events but lagged
in sensitivity. SCOPE and SCYN completely missed these events - after inspecting
SCOPE's normalized read depths (SCYN uses the same normalized profile as SCOPE),
we found that SCOPE's normalization algorithm interprets these amplifications as
technical errors and over-corrects for them. While this cautious approach makes sense

when analyzing hundreds of cells, it becomes problematic with just a few cells since
distinguishing real biological signal from technical artifacts becomes more challenging.

In cnLOH detection (Fig 2h,i), only HiScanner and CHISEL showed capability, with
CHISEL demonstrating sharp improvement around 1Mb (>0.90 precision and sensitivity
for larger events), while HiScanner showed gradual improvement. CHISEL's increased
sensitivity for large cnLOH events might be attributed to its joint clustering approach
which utilizes both RDR and BAF, whereas HiScanner's multi-sample segmentation is
based only on RDR. The other tools were not designed for cnLOH detection due to their
BAF-agnostic nature (Table 1). Overall, HiScanner demonstrated the most consistent
performance across CNA types, while other tools showed variable effectiveness
depending on the variant type and size, with all tools struggling with CNAs below 500kb.

278
Figure 2j illustrates a representative ~1Mb single copy loss. In this example, while all
five methods detected a change in the region, only HiScanner, Ginkgo, and SCOPE made
a precise call matching the ground truth boundaries. CHISEL correctly identified the loss
but predicted spurious copy number gains in the flanking regions, likely due to sensitivity
to coverage fluctuations. SCYN showed multiple errors, including a spurious loss within
the true CNA region and an incorrect gain outside of the region.
For the single copy gain (Fig 2k), the event is characterized by an increase in RDR and
shift in BAF. HiScanner made a clean detection of the gain, while CHISEL and Ginkgo
both identified the event but showed incorrect copy number states in the surrounding
regions. SCOPE and SCYN failed to detect the gain, maintaining a baseline copy number
state throughout the region. These examples illustrate HiScanner's superior ability to
make precise calls while avoiding false positives due to technical noise in both RDR and
BAF signals.

292
We next assessed the impact of bin size and sequencing depth on HiScanner's
performance across different CNA types. For bin size analysis, we compared 50KB,
100KB, and 500KB bins in simulated 30X data (Supplementary Figure 3a). While
smaller bins (50KB) showed improved sensitivity for detecting variants below 500kb,
they generally exhibited lower precision due to increased noise. The 100KB bin size
provided optimal balance between sensitivity and precision across most CNA types,
particularly for variants >1Mb. Larger bins (500KB) demonstrated reduced performance,
especially for smaller variants. Regarding sequencing depth, we evaluated 0.5X, 10X,
and 30X coverage (Supplementary Figure 3b). Higher sequencing depth (30X)
consistently improved both precision and sensitivity across all CNA types, though the
performance gain between 10X and 30X was modest. Low-coverage sequencing (0.5X)
showed substantially lower performance, particularly for single-copy events and smaller
variants, though it maintained reasonable accuracy for larger CNAs (>2Mb) and two-
copy gains. Notably, the performance difference between 10X and 30X coverage was less
pronounced for cnLOH events, suggesting that moderate coverage might be sufficient for
detecting these variants. These results validate that the default parameter used in our

benchmarking experiment (30X depth, 100KB bins) as providing optimal performance
while indicating that cost-effective alternatives (e.g., 10X coverage) might be suitable for
specific applications.
312
As an illustrative example using real data, we next applied the five methods to detect
CNAs in a single human neuron amplified by PTA and sequenced to 29X (Fig 3a). When
examining the genome-wide copy number profiles, HiScanner and Ginkgo showed
largely consistent patterns in detecting major copy number alterations across
chromosomes, revealing distinct loss events in chromosomes 2, 5, 6, 8, 11, 12, and 17.
These events span various sizes from 300 kb to 170 Mb (whole-chromosome), with some
potentially biallelic losses. SCYN and SCOPE demonstrated similar capabilities in
identifying these major events, though with slightly different segmentation patterns. In
contrast, CHISEL produced a notably different profile, suggesting an overall tetraploid
state with more fragmented segmentation, even when using 100kb bins. After parameter
adjustment with 5Mb bins, CHISEL showed improved stability but still maintained
higher baseline copy numbers compared to other methods.
325
The differences among these methods became more apparent when examining their
behavior in specific genomic contexts (Fig 3b). Near centromeric regions, Ginkgo and
CHISEL frequently reported focal amplifications with high copy numbers, suggesting
insufficient correction for the highly repetitive sequences in these regions. HiScanner,
however, maintained more stable signals across these regions, likely due to its use of
uniquely mappable positions and nucleotide-level normalization approach. SCYN and
SCOPE also showed robust performance in centromeric regions, avoiding the spurious
high copy number calls seen with other methods. The varying complexity of CNA calls
among methods is quantified in Fig 3c, where CHISEL showed the highest rate of
segmentation (10,322 breakpoints, or 0.38 per 100 kb). In comparison, HiScanner,
Ginkgo, SCYN, and SCOPE maintained more conservative segmentation patterns,
suggesting a better balance between sensitivity and specificity in CNA detection. When
examining the size distribution of detected CNAs and their BAF support (Fig 3d), we
observed that Ginkgo tended to call small copy number loss events that lacked
corresponding BAF evidence. HiScanner showed better concordance between copy
number calls and BAF signals, particularly for larger events.
342
343 To further demonstrate the performance differences among these methods, we examined
344 two specific genomic regions in detail: a diploid region with uneven read coverage (Fig
345 3e) and a single copy loss candidate (Fig 3f). In both cases, HiScanner maintained stable
346 copy number calls despite coverage fluctuations, while the other methods showed more
347 variable responses to these coverage differences.
348
In summary, our analysis of both simulated data and an aneuploid single neuron revealed
distinct characteristics of each caller. Ginkgo displayed high sensitivity to changes in
RDR but, without BAF information, lacked precision in detecting small CNAs and was

not capable of cnLOH detection. CHISEL tended to overfit when used with small initial
bin sizes and, in a single-sample context, produced an improbably large number of
breakpoints. Given that CHISEL is optimized for megabase-scale clonal CNA detection
(e.g., in cancer samples), this behavior is expected. SCYN showed limited sensitivity
across different CNA types in simulated data, and while it could identify major copy
number events in real data, it sometimes produced spurious calls in regions with coverage
fluctuations. SCOPE demonstrated more conservative calling patterns, maintaining
reasonable precision but with reduced sensitivity, particularly for smaller events and
subtle copy number changes. Overall, HiScanner maintained a balance between
sensitivity and precision in detecting CNAs at high resolution.
362
Cell-type-specific CNA patterns in neurotypical human brains

Previous studies using scWGS to investigate non-diseased human brains have reported
contradictory findings regarding the frequency of CNAs in neurons. A meta-analysis of
multiple low-coverage scWGS reported a linear decline of CNA neuron frequency with
age (R^2=0.92, range 0-40.0%) in brains aged 0.36 y.o. to 95 y.o.[15] Knouse et al. reported
a significantly lower frequency (2.2%; 95% CI 0.3%–7.9%) for large CNAs in human
brain cells[37]. More recently, Kalef-Ezra et al. applied PTA to human brain cells and found
33.8% to have at least one megabase-scale CNA[12,14]. These discrepancies highlight the
need for further research into the prevalence of CNAs in the human brain.
372
Using HiScanner, we analyzed previously published high-coverage PTA scWGS data
encompassing 65 single neurons and oligodendrocytes from eleven neurotypical human
brains spanning an age range from 0.4 to 83 years[12,14] (Fig 4a; Supplementary Table 1).
HiScanner was applied separately to each brain, as clonal CNAs should be specific to
each individual. Large-scale aneuploidy, characterized by multiple chromosome-arm
level losses, was clearly observed in two cells out of 65 (3.1%). These numbers are
concordant with Knouse et al[37] than the others.
380
Beyond these instances, HiScanner revealed an additional 614 smaller, mostly private
CNAs (314 private losses and 57 private gains) in the rest of the cells. Whereas previous
approaches were limited to a lower size bound of ~3 Mb on somatic CNA detection in
single cells[15,16,36], nearly all (609/614, 99.3%) of the CNAs identified by HiScanner were
smaller than 3 Mb (Fig 4b-c), with a median size of 612 Kb (range 300Kb-24.1Mb).
Using the size-specific sensitivities and specificities from our simulated data, we
estimated a false discovery rate of 26.3% among the 614 calls (Methods; Supplementary
Table 2). In total, 77% (24/31) of neurons and 53% (17/32) of oligodendrocytes
contained at least one CNA. We identified 72 clonal events (defined as CNAs sharing
identical start and end bin coordinates across multiple cells), with the majority (50/72)
being shared between more than two cells, demonstrating substantial clonal diversity.
Analysis of age-related patterns using linear mixed effects models revealed a weak
negative correlation between age and non-clonal CNA gains in oligodendrocytes
(Supplementary Figure 4a; slope=-0.020 CNA/year, p=0.010). No significant age-related
patterns were observed for neuronal gains (Supplementary Figure 4a; slope=-0.015
CNA/year, p=0.218) or for non-clonal losses in either cell type (Supplementary Figure

4b; neurons: slope=0.003 CNA/year, p=0.976; oligodendrocytes: slope=0.002, p=0.589),
suggesting that the relationship between CNAs and aging may be more complex than
described in prior literature[15].
400
Beyond these temporal patterns, our analysis revealed striking differences in CNA
profiles between cell types. Neurons contained both gains and losses, with losses being
more prevalent, whereas nearly all CNAs in oligodendrocytes were gains (Fig 4b-c). In
comparison to oligodendrocytes, the odds ratio of observing losses compared to gains in
neurons is 61 (p-value: 1e-23, Fisher's Exact Test). Classification into clonal vs non-
clonal categories revealed broad distinctions: only 5% of the losses in neurons are clonal,
whereas 78% of neuronal gains are clonal. Since neurons are typically non-proliferative,
this observation may suggest differential timings of gain and loss events: gains, which
were mostly shared, must have occurred early in development prior to terminal
differentiation into a post-mitotic neuron, whereas losses, which were mostly private,
must have occurred after development in the post-mitotic state. In oligodendrocytes, the
majority of CNAs are clonal gains (81% of gains being clonal and no losses), similar to
the high clonal fractions in neuronal gains. The differences between neuronal and
oligodendrocyte CNAs suggest differential timing and/or mutagenic processes.
415
For cnLOHs, we focused our analysis on events larger than 2Mb where our simulations
demonstrated high precision and sensitivity (~0.9). This analysis identified cnLOH events
in 8 neurons, with one neuron (5817PFC-B) showing an extensive pattern of 12 distinct
cnLOH events while the remaining neurons harbored 1-3 events each. Two of these
cnLOH events were clonal, appearing in pairs of neurons from the same individual, while
the rest were private. Notably, no cnLOH events were detected in oligodendrocytes,
similar to the cell-type specific pattern observed with single copy losses.
423
Analysis with Ginkgo yielded a fundamentally different view of neuronal and
oligodendrocyte CNAs, even though a similar range of CNAs were detected (Fig 4d-e).
Notably, Ginkgo called nearly a 10-fold more private neuronal losses than HiScanner and
a large number of private losses in oligodendrocytes whereas HiScanner detected almost
none. We assessed the quality of these calls by inspection of the BAF distribution in the
putative CNAs and found that a considerable proportion of Ginkgo's calls disagreed with
local BAF signals (Fig 4f-j): 97% (581/599) of CNAs called in oligodendrocytes and
66% (1410/2121) of CNAs called in neurons were BAF-discordant. This suggests that
Ginkgo's calls in PTA scWGS data should be filtered based on CNA size or be subjected
to BAF-based post-call filtering strategies[21] to ensure the reliability of small CNA calls.
In contrast, HiScanner's integration of BAF and RDR signals was effective in preventing
these types of errors.
436
Integrative CNA and SNV analysis in longitudinal meningioma scWGS

To demonstrate the applicability of HiScanner to low-coverage scWGS, we sequenced
PTA-amplified single cells from meningiomas, the most common primary intracranial
tumor, to an average depth of 0.5X (Supplementary Table 3). High-grade (grade II-III)
meningiomas are notorious for their high recurrence rate despite treatment, and the

effectiveness of postoperative radiotherapy in managing these recurrences is still a
subject of debate[38,39]. Characterizing the timing of oncogenic events and their evolution
in meningiomas may therefore prove helpful in assessing the impact of therapeutic
interventions and guiding clinical management strategies. We obtained tumor tissue
samples from initial and recurrent (grade II/III) meningiomas from a single patient. 88
and 91 single tumor cells were sorted from the grade II and grade III specimens,
respectively, totaling 179 cells (Fig 5a, Methods). All cells were amplified by PTA,
sequenced to 0.5X, and analyzed for CNAs by HiScanner.
450
Hierarchical clustering on the copy number profiles from both initial (grade II) and
recurrent (grade III) tumors revealed four main clusters (Fig 5b-d). Cluster 13, which is a
mixture of grade II/III cells, consists of cells that lack any large CNAs. Cluster 11 is
entirely composed of grade II cells and exhibits hallmarks of high-grade meningiomas
such as single-copy losses in chromosomal regions 1p and 22q (Fig 5e). Cluster 15 is
dominated by grade III cells which inherited all CNAs present in Cluster 11 along with
additional loss events on chromosomes 6q, 9p, and 11q (Fig 5f). Interestingly, one grade
II cell (BOS-EE06, Fig 5g) was assigned to Cluster 15, suggesting that it may be closely
related to the origin of a major grade III clone. If true, this would imply that the
additional copy number events found in grade III tumor cells likely occurred in the
natural course of tumor evolution rather than as a consequence of adjuvant therapy.
Lastly, Cluster 8 represents a small, distinct group of tetraploid cells from the grade III
tumor (Fig 5h), exhibiting a range of copy number states from two to six, including odd
copy numbers. Tetraploidy hints at a potential tumor progression pattern involving
whole-genome duplication, which is rarely reported in the meningioma literature;
however, further experiments (e.g., fluorescence in situ hybridization) would be essential
to validate this ploidy estimate. The heterogeneity of these clusters reflects the complex
nature of meningioma evolution and underscores the importance of single-cell profiling
in delineating tumor progression.
470
Next, we investigated whether cluster assignments derived from CNAs were consistent
with patterns of somatic SNVs (sSNVs). To this end, we employed a recently developed
somatic SNV-based cell clustering method, SECEDO[26]. Initially, applying SECEDO with
its default settings failed to differentiate between aneuploid and diploid cells, which are a
hallmark of meningiomas. However, upon refining the input by limiting to sSNV
locations identified in matched bulk WGS via GATK MuTect2[40] (Methods), SECEDO
yielded reasonable clustering results. SECEDO predicted three primary cell clusters
which closely corresponded to HiScanner's CNA-based clustering (Fig 5i, j): SNV
Cluster 4 aligned largely with CNA Cluster 11; sSNV Cluster 2 comprised a mix of cells
from CNA Clusters 8 and 13; and sSNV Cluster 4 was found to be a mix of CNA
Clusters 8 and 15. Notably, however, SECEDO's clustering did not distinguish diploid
and tetraploid grade III cells. A possible explanation for this is that the whole-genome
duplication preceded the majority of sSNV events. SECEDO also assigned three grade II
tumor cells to its grade III major cluster, suggesting that some sSNVs preceded CNAs in
the evolution of this tumor. To quantitatively compare the sSNV and CNA cluster
assignments, we computed the Adjusted Rand Index (ARI), a score ranging from 0
(random labeling) to 1 (perfect match). The two orthogonal clusterings produced an ARI

of 0.83, indicating a high level of agreement and again supporting the validity of
HiScanner's CNA calls.
490
491

Discussion

We developed a new single-cell CNA detection algorithm, benchmarked it on real and
synthetic data, and applied it to high-depth scWGS of non-diseased human brains and
low-depth scWGS of longitudinal meningioma tumor samples. In our benchmarking,
HiScanner achieved similar performance to a BAF-aware and BAF-agnostic single-cell
CNA caller for detecting large (3-5 Mb) CNAs but outperformed them in the task of
detecting small CNAs (<1 Mb) and cnLOH events. Small and cnLOH events are
characteristic of many mutagenic processes and detecting them accurately is essential.[41]
While our current model assumes independence between genomic bins for computational
efficiency, future work could explore approaches such as conditional random fields or
deep learning architectures to capture dependencies between adjacent genomic regions,
potentially improving sensitivity in challenging regions.
504
In the human brains, HiScanner detected many CNAs events smaller than the broadly
assumed ~3 Mb detection limit. Although the small sample size provided insufficient
power to ascertain an association between CNA burden and age, HiScanner revealed
previously undetected differences[12] in CNA patterns between cell types. Neuronal CNAs
were predominantly private losses, in line with previous studies in human neurons[42],
whereas oligodendrocytes mostly exhibited clonal gains. This finding suggests that
neurons likely acquired these losses post-mitotically via non-replicative mechanisms such
as nonhomologous end joining and microhomology-mediated end joining which can lead
to deletions or complex genomic rearrangements without the need for DNA synthesis[43].
Indeed, neurons frequently undergo double-stranded breaks and repair due to their
synaptic activity[44]. In contrast, the clonal gains found in both neurons and
oligodendrocytes may have occurred earlier in brain development through replicative
mechanisms such as microhomology-mediated break-induced replication and fork
stalling and template switching, which can induce duplications or deletions during DNA
replication[43]. While the clonal gains in post-mitotic neurons are most likely to have
occurred during development, the clonal gains in oligodendrocytes may have occurred
later in life in ancestral oligodendrocyte precursor cells, which remain proliferative.
Further investigation into the timing and mutational mechanisms of these CNAs with
more cells is warranted. It is also important to note that while we classify CNAs as clonal
when they share identical start and end bin coordinates across multiple cells, some of
these events could also potentially represent independent recurrent mutations rather than
true clonal expansions.
527
Our integrative analysis of sSNVs and CNAs provided a more detailed view of the clonal
origin and evolution of a meningioma tumor, highlighting the complex and dynamic
nature of cancer progression and the importance of single-cell profiling. This approach
identified an ancestor of the grade III major clone in the grade II tumor, suggesting that
the mutagenesis was not caused by radiotherapy. However, it remains possible that the
clonal expansion of the grade III clone was influenced by selective pressure induced by

radiotherapy. Timing the progression between tumor grades, and whether this occurs in
response to or because of therapy, is critical information for clinical decision making.
536
Several studies of somatic CNAs have delineated the evolutionary trajectories of
subpopulations of tumor cells, most notably in breast cancer where chromosome-scale
aneuploidy is common. A more comprehensive set of CNAs including smaller events will
allow similar analysis for other tumor types in which the large CNAs are less common.
The timing of SNV driver mutations with respect to the appearance of genome-doubling
or large CNAs can be inferred based on careful analysis of variant allele fractions in bulk
data[45]. With the growing availability of high-coverage scWGS and the accompany
development of methods for detecting SNVs, CNAs, and other variants, a timing analysis
at the single cell resolution will be possible, deepening our understanding of genomic
stability and evolution.
547

Methods

Ethics declaration

This study was performed in accordance with all relevant ethical regulations. The meningioma study was approved by the Brigham and Women's Hospital's Institutional Review Board.

Preprocessing of BAF

The GATK HaplotypeCaller "Best Practices" pipeline[40] with default parameters was used to call SNPs on bulk sequencing data from each individual. Germline heterozygous SNPs (gHETs) were defined as the variants passing the FILTER column flag, marked with a heterozygous '0/1' genotype, annotated non-zero population frequencies in the 1000 Genomes Project database[46] and listed in the Single Nucleotide Polymorphism Database[47]. SHAPEIT2[48] was applied to phase the gHETs using the 1000 Genomes Phase 3 integrated haplotype panel[49]. Only the gHETs that can be phased by SHAPEIT2 and have a variant allele frequency between 0.1 and 0.9 in bulk data were retained for subsequent analysis. BCFtools[50] was used to query the allelic-specific depth at each gHET site within each single cell BAM. Next, reads associated with the same haplotypes were aggregated to obtain bin-wise haplotype-specific read counts. The phased B-allele frequency (pBAF) was computed as the fraction of haplotype-B read count divided by the total read count within a given bin. Note that the pBAF here corresponds to individual bins and can be heavily influenced by local allelic imbalance. To ensure robust allelic copy number inference, we implemented additional per-segment refinement of pBAF after the segmentation step using the LOH test (Supplementary Notes).

Preprocessing of RDR

BICseq2-norm[25] with default parameters was used to perform GC correction and binning on single-cell and bulk sequencing data from each individual. Only uniquely mappable reads with a minimum mapping quality of 30 were considered and only mappable genomic loci, as determined by the 100mer mappability track from https://www.math.pku.edu.cn/teachers/xirb/downloads/software/BICseq2/BICseq2.html, were considered.

Allelic dropout analysis

Loss or under-amplification of one allele during whole-genome amplification, known as allelic dropout, is a well-documented phenomenon in single-cell sequencing assays[28]. In the sequencing data, dropout regions manifest with reduced RDRs and imbalanced allelic ratios, closely resembling regions with copy number losses. The characteristics and prevalence of dropout can be affected by several factors, including amplicon length and sequencing depths. Consequently, different single-cell assays may exhibit varying patterns of dropout, and these nuances should be accounted for when determining the optimal bin size for CNA analysis in single cells.

To determine the distribution of dropout sizes without relying on arbitrary thresholds, we
designed a hidden Markov model (HMM) with two distinct states—"Allelic dropout
(ADO)" and "non-ADO"—and used Gaussian distribution probabilities to describe them
(Supplementary Figure 1). This model assumes that CNAs are infrequent compared to
allelic imbalance by dropout in the selected single-cell genomes (in this study we used
oligodendrocytes). We used a binary representation of BAF data at gHETs on the
autosomes. A value of |BAF-0.5| less than 0.2 was marked as "0" to indicate no sign of a
homozygous state, while a value of 0.2 or greater was marked as "1" to indicate a
homozygous state. The HMM parameters are learned using the Baum-Welch algorithm
(implemented via the hmmlearn library in Python), making the approach adaptable to any
amplification method. To demonstrate this flexibility, we applied our framework to both
PTA and MDA data, using oligodendrocytes with minimal aneuploidy as training
examples (1278-Oligo-5 for PTA; 1278_ct_p1G9 for MDA). The trained HMMs were
then used to predict dropout events across additional cells (1278-Oligo-{7,8} for PTA;
1278_ct_{p1E3, p1E6, p2B9, p2E4, p2F5, p2G5} for MDA). For each amplification
method, we selected the 95th percentile of the observed dropout size distribution as the
minimum bin size, allowing the framework to automatically adapt to the characteristics of
different protocols.
609
Joint segmentation algorithm

We extended the single-sample segmentation algorithm implemented by BICseq2-seg[25]
to the multi-sample case. Within the original BIC-seq framework, the simplified Bayesian
Information Criterion (BIC) is defined as:

$$BIC(\lambda) = -2\sum_{j=0}^{m}[a_j(log(\frac{a_j}{a_j+b_j}) + b_j log(\frac{b_j}{a_j+b_j})] + (m+1)\lambda \log(N) \qquad (1)$$

where for every breakpoint j, a_j and b_j are the number of observed reads and the number
of expected reads between breakpoints β_j and β_{j+1}; $\lambda > 0$ is the tuning parameter; m is the
number of breakpoints; and $N = \sum_{j=0}^{m} a_j$ is the total number of reads in the given cell.
Recognizing that the joint log-likelihood of the short reads from G single-cell samples is
just the sum of the individual log-likelihoods, and assuming independence in cells'
segmentation patterns, the multi-sample form of the simplified BIC becomes:
621

622 $$BIC(\lambda) = -2\sum_{i=1}^{G}\sum_{j=0}^{m}[a_{kj}(log(\frac{a_{kj}}{a_{kj}+b_{kj}}) + b_{kj} log(\frac{b_{kj}}{a_{kj}+b_{kj}})] + (m+1)G\lambda \log(\sum_{k=1}^{G} N_k) \ (2).$$

623
While this formulation assumes independence between cells, which may be violated in
cases where cells share clonal origins (e.g., tumor samples), this approximation is
reasonable since (i) even in clonal populations, substantial cell-to-cell variation exists due
to both technical factors (e.g., amplification noise) and biological mechanisms (e.g.,
acquisition of new CNAs in subclones); and (ii), modeling the full dependency structure
between cells would be computationally expensive for large datasets.
630
Given an initial set of bins, HiScanner attempts to reduce the overall BIC by merging the
neighboring bins. HiScanner calculates the BIC for each possible merge of adjacent bins.
If the calculated BIC for a merged pair is lower than it would be if they remained
separate, HiScanner considers these bins for merging. Out of all possible merges,
HiScanner selects the pair of bins that, if merged, would result in the greatest reduction in

the BIC. The selected bins are then merged. HiScanner repeats this process, continuing to
merge bins pair by pair, until no more merges would result in a lower BIC.
638

Ploidy and copy number inference

We extended the likelihood-based method introduced in CHISEL[19] to infer ploidy and
allelic copy number states. The procedure has two main steps: identification of all
possible values of the scale factor γ, and finding the maximum likelihood estimate of
allelic copy number and γ based on BIC. The scale factor γ is a cell-specific constant
value and describes the linear relationship between the total copy number and RDR for
each bin. It follows that knowing the true copy number and RDR of one bin is sufficient
to compute the corresponding γ. We assume that within any cell, there exists a substantial
proportion of bins unaffected by CNAs, retaining balanced allelic copy numbers. The
allelic copy number of these bins takes the form {1,1} when there is no whole genome
duplication (WGD), {2,2} with one WGD, or {3,3} with two WGDs, etc. CHISEL's
global clustering approach allows for directly selecting the largest cluster with BAF≈ 0.5
as the "balanced" bin set, even when these bins are not contiguous. In contrast,
HiScanner's bottom-up segmentation restricts bin merging to spatially adjacent bins with
similar RDR and BAF values. In HiScanner, the "balanced" bins are identified by
requiring $0.45 < \text{BAF} < 0.55$. HiScanner employs a Gaussian Mixture Model (GMM) to
further segregate these "balanced" bins into clusters with similar RDR values. To
determine the best GMM, HiScanner conducts a parameter search for the number of
Gaussian components, iterating through $\Theta = \{1,...,t\}$, where t denotes the maximum
number of allowed WGD events. Only the set of bins in the largest cluster is chosen as
the set of "balanced" bins. The identification of candidate γ relies on these balanced bins,
where both alleles have equal copy numbers. We express the set of candidate γ values as:

$$\Gamma = \{\gamma = \frac{2\theta}{\frac{1}{|S|}\sum_{i \in S} x_i} : \theta \in \Theta\} \quad (3)$$

(i.e., Equation 38 in CHISEL[19]'s supplementary note) where Θ represents possible allele-
specific copy numbers, S is the set of balanced bins identified through BAF clustering, x_i
is the RDR of bin i, and |S| is the total number of balanced bins. The numerator 2θ
represents the expected total copy number for balanced bins, while the denominator
calculates the mean RDR across all balanced bins. This formulation systematically
identifies potential scale factors that map observed read depths to absolute copy numbers,
accounting for varying levels of genome duplication - from normal diploid state ($\theta=1$,
resulting in {1,1}), through single WGD ($\theta=2$, yielding {2,2}), to multiple WGD events
($\theta=3$ for {3,3}, and so on).
672

Next, HiScanner introduces a homozygosity testing framework that leverages local GC
bias and allelic imbalance to derive robust per-segment BAF estimates (Supplemental
Notes). The segment-wise RDR is simply the mean RDR of all bins in a given segment.
By modeling both RDR and BAF as Gaussian mixtures for each allelic copy number
state, HiScanner applies the maximum likelihood formulation in CHISEL[19]
(Supplementary Note 5.3) to find the most probable allelic copy number state for each
segment.
680

Simulation of male synthetic diploid X chromosomes

We follow a procedure similar to that described in Luquette *et al*[11] to construct synthetic diploid X (SDX) chromosomes from three single oligodendrocytes from two neurotypical male donors. For simplicity, here we denote the individuals as A and B, and the cells as OligoA, OligoB1, and OligoB2. We simulated three types of CNAs: single copy losses, single copy gains, and cnLOHs at 13 different sizes following a geometric progression from 200kb to 5Mb (Supplementary Data). For each CNA type and size, we randomly generated 100 start sites within chromosome X, after excluding pseudoautosomal regions and germline CNAs regions. For single copy losses, we removed reads within the randomly generated start and end sites from OligoB1. For single copy gains, we added reads mapped to CNA coordinates in OligoB2 to OligoB1. For cnLOH, we did not make changes to the read depth (only to BAF, described below), as both copies should be independently sampled in real data. After the read removal or addition, the modified OligoB1-reads and OligoA-reads were merged with SAMtools[50] to create the final SDX.

We next simulated gHETs and BAFs. To resemble a realistic SNP density, we randomly selected five positions in each 10 kb region to serve as simulated gHET loci. Only the reads uniquely mapping to the simulated gHETs are used to compute BAF. For cnLOH, only the reads from OligoA at CNA regions were used. Additional Poisson noise was added to allele-specific read depths at gHETs to represent sequencing errors. Furthermore, we simulated phase-switch errors that can occur during reference-based phasing by introducing an indicator variable i to determine whether to flip the allele-specific read depths. Specifically, i follows a binomial distribution with a probability of 0.01.

To evaluate performance across different sequencing depths and bin sizes, we created additional simulated datasets. For sequencing depth analysis, we downsampled our simulated 30X data to generate 10X and 0.5X coverage datasets. For bin size analysis, we tested performance using 50KB, 100KB, and 500KB bins on 30X data. All combinations of depth and bin size were evaluated using the same CNA types and sizes described above, allowing systematic assessment of these parameters' effects on detection sensitivity and precision.

Copy number calling with Ginkgo, SCOPE, SCYN and CHISEL

Ginkgo was run with default parameters with the "binning" parameter set to "`variable_100000_101_bwa`" and hg19 genome. Ginkgo's command line version was downloaded from https://github.com/robertaboukhalil/ginkgo. For SCOPE, we used the R package to process BAM files with a 100kb resolution, MAPQ threshold of 40, and GC/mappability correction using built-in hg19 tracks. Normalization was performed using `normalize_codex2_ns_noK` followed by `normalize_scope_foreach` with K=1 and T=1:7. Copy number segmentation was done using `segment_CBScs` with integer mode. SCYN was run with 100kb bins using SCOPE's normalized profiles as input, performing dynamic programming-based segmentation. For the single-neuron comparison, we applied CHISEL's complete pipeline to both single-cell PTA data and paired bulk sequencing data from the same individual. This was done separately for bin

sizes of 5 Mb and 100 kb. Regardless of bin size, we noticed a tendency for high ploidy predictions and overfitting, similar to what was reported previously[21]. To address this, we fine-tuned several parameters in the *chisel_calling* module to optimize the method performance. Specifically, we set the *sensitivity* parameter to 10, *maxploidy* to 3 and *k* to 10. In addition there was a strong GC bias in CHISEL-preprocessed RDR for 100 kb bins, likely due to the lack of GC-normalization in the CHISEL pipeline. To ensure fairness in the comparison of segmentation algorithms during the synthetic diploid experiment, we used the same RDR and BAF data preprocessed by HiScanner as input for both HiScanner's and CHISEL's segmentation algorithms. CHISEL's bioconda package was downloaded from https://github.com/raphael-group/chisel. We used bwa-mem[51] to align all the data presented in this manuscript to the "hg19/GRCh37+decoy" reference genome.

HiScanner analysis of single neurons and oligodendrocytes

We analyzed PTA-amplified scWGS data and paired bulk whole-genome sequencing from two public datasets of eleven neurotypical brains[12,14]. Each brain includes 2 to 3 neurons and 2 to 3 oligodendrocytes whose genomes were subject to high-depth sequencing (20-30X per cell). HiScanner was run separately for each brain, using the PTA and bulk sequencing data as input and a common bin size of 100kb, as determined by our allelic dropout size model. Two out of 33 neurons were found to be aneuploid, displaying extensive copy number losses. No oligodendrocytes were aneuploid. Given our focus on megabase-scale mosaic CNAs, we excluded the aneuploid neurons from subsequent analyses of CNA length and prevalence. Single-cell CNA calls that overlapped with bulk CNA calls were removed. To maintain consistency across samples from both sexes, our analysis was restricted to autosomal chromosomes.

We employed a linear mixed effects (LME) model framework to assess the relationship between CNA count and age across different cell types and CNA types. The statistical analyses were implemented using the *statsmodels* library in Python, using the *mixedlm* function from the *statsmodels.formula.api* module. We analyzed four combinations: gains in neurons, gains in oligodendrocytes, losses in neurons, and losses in oligodendrocytes. The dependent variable in our LME models was the CNA count (*cna_count*), with age as the predictor variable. To account for the within-individual correlation of CNA counts, we included random intercepts by specifying the individual's identifier (*indID*) as the grouping variable for random effects. This approach allowed us to model the intra-subject variability and adjust for the non-independence of the data. The formula for the LME models was set as *cna_count ~ age*, fitting each of the four specified datasets.

Estimation of false-discovery rate (FDR)

We computed mean FDR values for each CNA size and type assessed in the synthetic diploid experiment. Each CNA detected in the single neuron and oligodendrocyte data was matched with the FDR value from the same event type (e.g., gain or loss) with the largest size smaller than the CNA event. The final FDR estimate is then calculated as the mean of FDRs across all CNAs. This produces a conservative estimate of FDR since FDR increases as CNA size decreases.

Development of ViScanner

We developed a web-based visualization platform (https://parklab.github.io/ViScanner/) in order to visualize HiScanner output. It is a React-based web application hosted on Github. The interactive visualization is based on Higlass[27]. Custom plugin tracks, tailored to the output files of HiScanner, were created for HiGlass. When using ViScanner, the user's input files are loaded into the local browser cache and are not transferred to a server, thus enabling safe use with confidential data.

Meningioma data generation

The samples were collected from the operation room and directly frozen in -80°C after written informed consent was obtained under the auspices of a human subjects institutional review board protocol approved by the Brigham and Women's Hospital. The histological WHO Grade for each meningioma was determined by board-certified neuropathologists. The samples were shipped to Bioskryb, Inc. and stored at -80C until nuclei isolation. Nuclei were isolated using the NUC-V2 protocol on an S2 Genomics Singulator 100 machine. ResolveDNA reactions were run for 1 plate for each sample for whole-genome amplification. All cells passing quality control metrics (PTA yield) were then subjected to library preparation and sequenced on a NextSeq 1000 instrument targeting 8M paired-end reads at 2x150bp.

HiScanner analysis of the single-cell meningioma dataset

HiScanner was run on the entire set of initial and recurrent meningioma scWGS as well as paired tumor and blood bulk sequencing, with a bin size of 500kb. Hierarchical clustering was performed using R package *hclust* with complete linkage and canberra distance.

Somatic SNV analysis of the single-cell meningioma dataset

GATK4 Mutect2[40] with default parameters was run on tumor bulk separately for initial and recurrent samples, using the blood bulk as control. Only variants with "PASS" flag were kept. SECEDO was run on the entire set of initial and recurrent meningioma single cells, with pileup positions restricted to Mutect2-called somatic SNVs.

Data Availability

Newly generated scWGS data for 179 PTA-amplified single-cells and two matched tumor/blood bulk WGS of the initial and recurrent meningioma samples have been deposited in European Genome-Phenome Archive under dataset ID EGAD50000001254 [https://ega-archive.org/datasets/EGAD50000001254]. The ground truth CNA coordinates used in the simulation experiments are provided in the Supplemental Data file. Previously generated scWGS and matched bulk WGS of PTA neurons are available for download at dbGaP, accession number phs001485.v3.p1 [https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs001485.v3.p1]. scWGS of MDA and PTA oligodendrocytes are available for download at NIAGADs with accession number NG00162 [https://dss.niagads.org/datasets/ng00162/].

Code Availability

HiScanner is open-source software and is available on https://pypi.org/project/hiscanner/[52]. HiScanner output can be visualized through ViScanner (https://parklab.github.io/ViScanner/). Scripts to reproduce the results of the primary analyses will be made available on GitHub (https://github.com/parklab/HiScanner).

Acknowledgments

We extend our deepest gratitude to the donors and their families, whose generous contributions are pivotal to the advancement of science. This work was supported by R01HG012573 and R01CA269805 to PJP, F31CA264958 to VVV, and a Sontag Foundation grant to MDJ. We thank Teng Gao, Viktor Ljungström, Antuan Tran, and Dominika Maziec, for their invaluable input to this work.

Author contribution Statement

Study design: Yifan Z., L.J.L., V.V.V., P.J.P.; HiScanner algorithm and software development: Yifan. Z.; synthetic data generation and caller performance benchmark: Y.Z., L.J.L.; meningioma somatic SNV clustering analysis: Yifan. Z.; interpretation of non-diseased brain CNA call set: D.D.S., C.A.W.; generation and interpretation of meningioma data: H.W.Y., M.D.J.; implementation of multi-sample segmentation module: X.W., R.X.; implementation of ViScanner: A.D.V.; split-read-based validation: Yuwei Z., Yifan Z.; manuscript writing: Yifan. Z., P.J.P., L.J.L. with contributions from all authors.

Competing Interests Statement

P.J.P. is a member of the scientific advisory board for Bioskryb Genomics, Inc.

References

1. Svensson, V., Vento-Tormo, R. & Teichmann, S. A. Exponential scaling of single-cell RNA-seq in the past decade. *Nature Protocols* vol. 13 Preprint at https://doi.org/10.1038/nprot.2017.149 (2018).

2. Saliba, A. E., Westermann, A. J., Gorski, S. A. & Vogel, J. Single-cell RNA-seq: Advances and future challenges. *Nucleic Acids Research* vol. 42 Preprint at https://doi.org/10.1093/nar/gku555 (2014).

3. Buenrostro, J. D. *et al.* Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* **523**, (2015).

4. Ramani, V. *et al.* Massively multiplex single-cell Hi-C. *Nat Methods* **14**, (2017).

5. Nagano, T. *et al.* Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature* **502**, (2013).

6. Arneson, N., Hughes, S., Houlston, R. & Done, S. Whole-genome amplification by degenerate oligonucleotide primed PCR (DOP-PCR). *Cold Spring Harb Protoc* **3**, (2008).

7. Laks, E. *et al.* Clonal decomposition and DNA replication states defined by scaled single-cell genome sequencing. *Cell* **179**, 1207–1221 (2019).

8. Zahn, H. *et al.* Scalable whole-genome single-cell library preparation without preamplification. *Nat Methods* **14**, 167–173 (2017).

9. Andor, N. *et al.* Joint single cell DNA-seq and RNA-seq of gastric cancer cell lines reveals rules of in vitro evolution. *NAR Genom Bioinform* **2**, lqaa016 (2020).

10. Gonzalez-Pena, V. *et al.* Accurate genomic variant detection in single cells with primary template-directed amplification. *Proceedings of the National Academy of Sciences* **118**, e2024176118 (2021).

11. Luquette, L. J., Bohrson, C. L., Sherman, M. A. & Park, P. J. Identification of somatic mutations in single cell DNA-seq using a spatial model of allelic imbalance. *Nat Commun* **10**, 3908 (2019).

12. Luquette, L. J. *et al.* Single-cell genome sequencing of human neurons identifies somatic point mutation and indel enrichment in regulatory elements. *Nat Genet* **54**, 1564–1571 (2022).

13. Bizzotto, S. *et al.* Landmarks of human embryonic development inscribed in somatic mutations. *Science (1979)* **371**, (2021).

14. Ganz, J. *et al.* Contrasting somatic mutation patterns in aging human neurons and oligodendrocytes. *Cell* **187**, (2024).

15. Chronister, W. D. *et al.* Neurons with complex karyotypes are rare in aged human neocortex. *Cell Rep* **26**, 825–835 (2019).

16. McConnell, M. J. *et al.* Mosaic copy number variation in human neurons. *Science (1979)* **342**, 632–637 (2013).

17. Van Loo, P. *et al.* Allele-specific copy number analysis of tumors. *Proceedings of the National Academy of Sciences* **107**, 16910–16915 (2010).

18. Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994–1007 (2012).

19. Zaccaria, S. & Raphael, B. J. Characterizing allele-and haplotype-specific copy numbers in single cells with CHISEL. *Nat Biotechnol* **39**, 207–214 (2021).

20. Wu, C.-Y. *et al.* Integrative single-cell analysis of allele-specific copy number
alterations and chromatin accessibility in cancer. *Nat Biotechnol* **39**, 1259–1269
(2021).
21. Sun, C. *et al.* Mapping recurrent mosaic copy number variation in human neurons.
Nat Commun **15**, (2024).
22. Garvin, T. *et al.* Interactive analysis and assessment of single-cell copy-number
variations. *Nat Methods* **12**, 1058–1060 (2015).
23. Wang, R., Lin, D.-Y. & Jiang, Y. SCOPE: a normalization and copy-number
estimation method for single-cell DNA sequencing. *Cell Syst* **10**, 445–452 (2020).
24. Feng, X. *et al.* Scyn: single cell cnv profiling method using dynamic programming.
BMC Genomics **22**, 1–13 (2021).
25. Xi, R., Lee, S., Xia, Y., Kim, T.-M. & Park, P. J. Copy number analysis of whole-
genome data using BIC-seq2 and its application to detection of cancer
susceptibility variants. *Nucleic Acids Res* **44**, 6274–6286 (2016).
26. Rozhoňová, H. *et al.* SECEDO: SNV-based subclone detection using ultra-low
coverage single-cell DNA sequencing. *Bioinformatics* **38**, 4293–4300 (2022).
27. Kerpedjiev, P. *et al.* HiGlass: web-based visual exploration and analysis of
genome interaction maps. *Genome Biol* **19**, 125 (2018).
28. Zhang, C. Z. *et al.* Calibrating genomic and allelic coverage bias in single-cell
sequencing. *Nat Commun* **6**, (2015).
29. Dean, F. B. *et al.* Comprehensive human genome amplification using multiple
displacement amplification. *Proceedings of the National Academy of Sciences* **99**,
5261–5266 (2002).
30. Browning, B. L. & Browning, S. R. Genotype error biases trio-based estimates of
haplotype phase accuracy. *Am J Hum Genet* **109**, (2022).
31. Loh, P. R., Palamara, P. F. & Price, A. L. Fast and accurate long-range phasing in
a UK Biobank cohort. *Nat Genet* **48**, (2016).
32. Browning, B. L., Tian, X., Zhou, Y. & Browning, S. R. Fast two-stage phasing of
large-scale sequence data. *Am J Hum Genet* **108**, (2021).
33. Loh, P.-R. *et al.* Reference-based phasing using the Haplotype Reference
Consortium panel. *Nat Genet* **48**, 1443–1448 (2016).
34. Olshen, A. B., Venkatraman, E. S., Lucito, R. & Wigler, M. Circular binary
segmentation for the analysis of array-based DNA copy number data. *Biostatistics*
5, 557–572 (2004).
35. Mallory, X. F., Edrisi, M., Navin, N. & Nakhleh, L. Assessing the performance of
methods for copy number aberration detection from single-cell DNA sequencing
data. *PLoS Comput Biol* **16**, e1008012 (2020).
36. Knouse, K. A., Wu, J. & Amon, A. Assessment of megabase-scale somatic copy
number variation using single-cell sequencing. *Genome Res* **26**, 376–384 (2016).
37. Knouse, K. A., Wu, J., Whittaker, C. A. & Amon, A. Single cell sequencing
reveals low levels of aneuploidy across mammalian tissues. *Proc Natl Acad Sci U*
S A **111**, 13409–13414 (2014).
38. Keric, N. *et al.* Impact of postoperative radiotherapy on recurrence of primary
intracranial atypical meningiomas. *J Neurooncol* **146**, 347–355 (2020).
39. Chen, W. C. *et al.* Radiotherapy for meningiomas. *J Neurooncol* **160**, 505–515
(2022).

40.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297–1303
(2010).
41.   Steele, C. D. *et al.* Signatures of copy number alterations in human cancer. *Nature*
606, (2022).
42.   Cai, X. *et al.* Single-Cell, Genome-wide Sequencing Identifies Clonal Somatic
Copy-Number Variation in the Human Brain. *Cell Rep* **8**, (2014).
43.   Hastings, P. J., Lupski, J. R., Rosenberg, S. M. & Ira, G. Mechanisms of change in
gene copy number. *Nature Reviews Genetics* vol. 10 Preprint at
https://doi.org/10.1038/nrg2593 (2009).
44.   Suberbielle, E. *et al.* Physiologic brain activity causes DNA double-strand breaks
in neurons, with exacerbation by amyloid-β. *Nat Neurosci* **16**, (2013).
45.   Yang, L. *et al.* An enhanced genetic model of colorectal cancer progression history.
Genome Biol **20**, (2019).
46.   Consortium, 1000 Genomes Project & others. A global reference for human
genetic variation. *Nature* **526**, 68 (2015).
47.   Sherry, S. T., Ward, M. & Sirotkin, K. dbSNP—database for single nucleotide
polymorphisms and other classes of minor genetic variation. *Genome Res* **9**, 677–
679 (1999).
48.   Delaneau, O., Zagury, J.-F. & Marchini, J. Improved whole-chromosome phasing
for disease and population genetic studies. *Nat Methods* **10**, 5–6 (2013).
49.   Delaneau, O. & Marchini, J. Integrating sequence and array data to create an
improved 1000 Genomes Project haplotype reference panel. *Nat Commun* **5**, 3934
(2014).
50.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,
giab008 (2021).
51.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, (2009).
52.   Zhao, Y. HiScanner (HIgh-resolution Single-Cell Allelic copy Number callER) .
GitHub Preprint at https://doi.org/10.5281/zenodo.15108638 (2025).
965
966
967

Tables

Table 1. Qualitative comparison of five single-cell copy number callers (using default parameters). RDR: read-depth ratio. BAF: B-allele frequency. CBS: circular binary segmentation. BIC: Bayesian Information Criterion. HMM: hidden Markov model. *: Alleloscope models gHETs directly and does not perform binning.

	Input signal type(s)	GC correction	Bin size (kb)	Segmentation algorithm	Joint segmentation across cells	Integration of phasing and BAF
Ginkgo	RDR	Bin-level	500	CBS	No	N/A
SCOPE	RDR	Bin-level	500	CBS	Yes	N/A
SCYN	RDR	Bin-level	500	BIC-based	Yes	N/A
SCOVAL	RDR, BAF	Bin-level	500	CBS	Yes	Filtering step
HiScanner	RDR, BAF	Nucleotide-level	100	BIC-based	Yes	CNA calling step
CHISEL	RDR, BAF	Not performed	50000	K-means clustering	Yes	Segmentation step
Alleloscope	RDR, BAF	Not performed	N/A*	HMM	No	CNA calling step

Figure Legends

Figure 1. HiScanner algorithm overview. HiScanner starts with scWGS data as input. In the preprocessing step, gHETs are called from bulk and phased using population-level reference panels. Normalization is performed at the nucleotide level to remove GC bias from read depth signals. Subsequently, a two-state hidden Markov model is employed to estimate ADO event lengths, which are then used to find the optimal bin size by setting a cutoff at the top 5th percentile of the length distribution. For each cell, HiScanner aggregates allele- and haplotype-specific read counts per bin to compute BAF and performs semi-parametric regression to compute RDR. In the segmentation step, HiScanner jointly partitions the genomes across all cells using BIC to identify breakpoints. An LOH test is then used to further refine the BAF estimates within each segment (see Supp. Note for details). In the variant calling and interpretation step, for each cell, HiScanner simultaneously infers the cell-specific scale factor and allelic copy number for each segment by modeling the RDR and BAF as Gaussian distributions. A web-based visualization tool, ViScanner, is also provided, which enables interactive visualization of HiScanner CNA calls at any resolution. Other downstream analyses of HiScanner output include tasks such as phylogenetic inference and correlation of CNAs with SNVs and indels. HiScanner's key contributions are highlighted in light red boxes. Created with BioRender.com.

Figure 2. Method comparisons on simulated data. a. Construction of synthetic diploid X chromosomes. First, two single cells are obtained from different male donors. Next, reads are extracted from chromosome X, excluding reads from pseudoautosomal regions, and germline or somatic CNAs identified in the original cells are removed. This serves as a template for creating various non-clonal CNA conditions, including single copy loss (1|0), single copy gain (2|1), two copy gain (3|1), and cnLOH (2|0), after which the reads are merged to create a BAM file for each mutated synthetic diploid chromosome X. **b-i.** Evaluations of precision and sensitivity for five callers for simulated single copy losses, single copy gains, two copy gains, and cnLOHs. Shaded regions indicate 95% confidence interval bands calculated from bootstrap sampling. **j-k.** Examples illustrating caller discrepancies in detecting a simulated single copy loss (j) and a simulated single copy gain (k). Ground truth CNA boundaries are shown as red dashed lines.

Figure 3. Method comparison in a single, aneuploid PTA neuron. a. BAF track of single neuron 5823PFC-B using 100 kb mappable bins, and corresponding RDR tracks overlaid with total copy number calls reported by different methods: HiScanner, Ginkgo, CHISEL, and CHISEL after parameter optimization on 5 Mb bins, SCYN, SCOPE. **b.** RDR distribution for bins inside centromeric regions. CHISEL*: CHISEL on 5 Mb bins. **c.** Breakpoint density of each method's call set. **d.** Length distribution of BAF-concordant or BAF-discordant loss events called by HiScanner and Ginkgo. All HiScanner calls were BAF-concordant (i.e., y value=0 for the red bars). **e.** An example of a diploid region showing uneven read coverage. **f.** An example showing a 1 Mb loss on chromosome 13. The shaded region indicates the location of a CNA candidate.

1021
Figure 4. Application to somatic mosaicism in human brain cells. a. Experimental
workflow. Nuclei from neurons and oligodendrocytes were isolated from 11 human
brains, followed by whole-genome amplification, sequencing, and somatic CNA
detection using Ginkgo and HiScanner. Created with BioRender.com. **b-c.** Length
distribution of HiScanner CNA calls, categorized by type and clonality, in 31 neurons **(b)**
and 32 oligodendrocytes **(c)**. **d-e.** Length distribution of Ginkgo CNA calls. **f.** Example of
a diploid region with uneven coverage where a spurious loss is called by Ginkgo. **g.**
Example of two adjacent single copy losses, one of which is missed by Ginkgo. **h.**
Example of a 2Mb single copy loss. Shaded regions indicate positions where Ginkgo
calls are discrepant with HiScanner and BAF. **i-j.** Number of Ginkgo calls that are
concordant or discordant with BAF, categorized by length and cell types.
1033
1034
1035
Figure 5. Application to low-coverage scWGS of a recurrent meningioma tumor
pair. a. 179 single cells from a paired initial and recurrent meningioma tumor sample
were amplified by PTA and sequenced. **b-d.** Total copy number **(b),** RDR **(c)** and BAF **(d)**
matrices output by HiScanner. Cell ordering is determined by hierarchical clustering
applied to the copy number matrix. Each row represents one cell and each column
corresponds to one genomic bin (500 kb mappable). **e-h** Total copy number and BAF
profiles of four representative cells from the major clone of tumor grade II **(e)**, major
clone of tumor grade III **(f)** grade III, BOS-EE06 **(g)**, and tetraploid clone from tumor
grade III **(h)**. **i-j.** Scatter plots of the second and third eigenvectors of the Laplacian
similarity matrix output by SECEDO, colored by SECEDO-inferred SNV clusters **(i)** and
HiScanner-based CNA hierarchical clusters **(j).**
1047
1048
1049

HiScanner (High-resolution Single-Cell Allelic copy Number callER)

Input · **Preprocessing** · **Segmentation** · **Variant calling and interpretation**

a Brain dissection → Nuclei sorting → Amplification → scWGS data → HiScanner/Ginkgo — Copy number analysis

b Neurons (HiScanner)
c Oligodendrocytes (HiScanner)
d Neurons (Ginkgo)
e Oligodendrocytes (Ginkgo)

Legend: non-clonal gain, non-clonal loss, clonal gain, clonal loss

f 5219 5219-Neuron-4 14:53023886-55896222
g 5559 5559PFC-A 18:39925774-41267330

BAF at individual gHETs
Bin-level BAF estimate
Scaled RDR
HiScanner — Ginkgo

h 5559 5559PFC-A 20:29628891-34510991

i Neurons (Ginkgo)
j Oligodendrocytes (Ginkgo)

BAF concordant
BAF discordant