A single episode of sexual reproduction can prevent population extinction under multiple stressors

Yawako W. Kawaguchi[1], Masato Yamamichi[1,2,3,4,5*]

[1]Center for Frontier Research, National Institute of Genetics, 1111 Yata, Mishima, Shizuoka 411-8540, Japan

[2]Genetics Program, Graduate Institute for Advanced Studies, SOKENDAI, 1111 Yata, Mishima, Shizuoka 411-8540, Japan

[3]School of the Environment, The University of Queensland, St. Lucia, Brisbane, Queensland 4072, Australia

[4]Department of International Health and Medical Anthropology, Institute of Tropical Medicine, Nagasaki University, 1-12-4 Sakamoto, Nagasaki 852-8523, Japan

[5]Institute for Multidisciplinary Sciences, Yokohama National University, 79-5 Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

Corresponding author: *myamamichi@nig.ac.jp

ORCID

YWK: 0000-0002-0389-7744

MY: 0000-0003-2136-3399

Keywords

Adaptation, extinction, population biology, sex

Abstract

Recent studies have shown that organisms can adapt to changing environments through

rapid contemporary evolution. Although intraspecific genetic variation is necessary for

rapid evolution to occur, little is known how genetic variation has been produced and

maintained before rapid evolution. Here we show that a single episode of sexual

reproduction can produce a large amount of intraspecific trait variation that allows

population growth in degraded environments by laboratory experiments of a green alga,

Closterium peracerosum–strigosum–littorale complex. We observed population

dynamics of the alga under multiple stressors and found that high salinity and low pH

decreased population growth rates. By comparing parental and their hybrid F_1

populations, we observed larger variation in population growth rates of F_1 populations

(i.e., transgressive segregation) when pH was low. Interestingly, even when parental

populations had negative growth rates, some F_1 populations showed positive growth

rates in severe environmental conditions. Together with the recently obtained genomic

information of the alga, our results suggest that recombination and variation in the

number of gene copies might produce large genetic variation in the F_1 generation. This

will be an important step toward a better understanding of evolutionary rescue, where

rapid evolution prevents population extinction in changing environments.

Introduction

Traditionally, evolutionary biologists tended to think that evolution is a slow process (Darwin 1859), but studies in the last quarter century have shown that organisms can flexibly adapt to changing environments through rapid contemporary evolution (Thompson 1998, Hairston et al. 2005, Schoener 2011, Hendry 2017, Sanderson et al. 2022). One of the most important implications of this phenomenon is that rapid evolution may prevent population extinction due to environmental changes, a process called "evolutionary rescue" (reviewed in Kinnison and Hairston 2007, Alexander et al. 2014, Carlson et al. 2014, Bell 2017). Previous studies have demonstrated that rapid evolution can prevent population extinction in severe environments by using mathematical modeling (e.g., Gomulkiewicz and Holt 1995, Orr and Unckless 2014, Uecker and Hermisson 2016, Orive et al. 2017), laboratory experiments (e.g., Bell and Gonzalez 2009, Lachapelle and Bell 2012, Lindsey et al. 2013), and field observations (e.g., Vander Wal et al. 2013, Isanta-Navarro et al. 2021).

It is well known, however, that evolution has no foresight and organisms cannot "prepare" for novel environmental changes in the future by maintaining genetic variation. How is genetic variation for rapid evolution produced and maintained in populations before the environmental changes? In addition to *de novo* mutations and immigration, classical studies in evolutionary biology have revealed that sexual reproduction is an important source for genetic variation (Williams 1975, Maynard Smith 1978, Bell 1982, Hartfield and Keightley 2012), and several experimental studies have indeed demonstrated that sexual reproduction can promote evolutionary rescue (Lachapelle and Bell 2012, Petkovic and Colegrave 2019, 2023). However, previous studies tended to compare experimental population dynamics of lineages with and without sexual reproduction and showed that sexual reproduction can increase the speed of adaptive evolution (Kaltz and Bell 2002, Goddard et al. 2005, Morran et al. 2009, Gray and Goddard 2012, Lachapelle and Bell 2012, Petkovic and Colegrave 2019, 2023). Thus, it was not clear how a single event of sexual reproduction can affect the amount of variation and the resultant rapid evolution. Colegrave et al. (2002) examined the effects of a single sexual episode on growth in various carbon sources, but they did not consider evolutionary rescue.

Here we show how a single episode of sexual reproduction can produce a large amount of trait variation and it makes population growth possible in degraded

environments using laboratory experiments of a unicellular green alga, *Closterium*

peracerosum–strigosum–littorale complex (Fig. 1A). The alga is a freshwater species

belonging to the Zygnematophyceae, a group most closely related to land plants

(Wickett et al. 2014). It usually grows vegetatively (asexual reproduction), but it can

also mate with another cell and form a zygospore that has desiccation tolerance (sexual

reproduction: Fig. 1B). As we can easily induce sexual reproduction and obtain F_1

populations in laboratory experiments (Kawaguchi et al. 2023), we observed population

dynamics of the alga under multiple stressors: salinity and acidity (low pH). Freshwater

green algae go extinct under salinity stress (Lachapelle and Bell 2012, Petkovic and

Colegrave 2019, 2023), and some species of the genus *Closterium* has been reported to

prefer neutral or high pH water (Gough 1977, Renstrom 1979, Greenwood and Lowe

2006). By comparing parental and hybrid F_1 populations, we observed large variation in

population growth rates of F_1 populations. This can be seen as an example of

transgressive segregation, the formation of extreme phenotypes in hybrid populations

(Rieseberg et al. 1999). Interestingly, even under severe environmental conditions when

parental populations had negative growth rates, some F_1 populations showed positive

growth rates. This suggests the potential importance of sexual reproduction for

producing genetic variation within populations that can prevent population extinction in

deteriorating environments. We discuss the potential importance of gene copy number

variation and recombination for producing genetic variation based on recently

sequenced genomic information for the alga (Kawaguchi et al. 2023).

Materials and Methods

Base Population and Culture Conditions

Two strains of *C. psl.* complex, NIES-64 and NIES-65, were obtained from the National

Institute for Environmental Studies (NIES), Environment Agency (Ibaraki, Japan). The

strains are genetically close and sympatric (Kawaguchi et al. 2023). They are

heterothallic strains and they can form zygospores between cells of two mating types:

mating type plus (mt^+: NIES-65) and mating type minus (mt^-: NIES-64) (Tsuchikane

and Sekimoto 2019, Kawaguchi et al. 2023, Ohtaka and Sekimoto 2023). They

recognize each other through the sex pheromones and then form zygotes (Fig. 1B:

Sekimoto et al. 1994, Nojiri et al. 1995). Through sexual reproduction, haploid parental

populations produce diploid zygospores, and then haploid offspring (F_1) populations
arise via meiosis and recombination (Fig. 1B).
The base population used for the experiments was cultured in a nitrogen-
supplemented medium (CA medium). The CA medium contains the following reagents
in 99.9 mL of distilled water: 2 mg of calcium nitrate tetrahydrate ($Ca(NO_3)_2 \cdot 4H_2O$),
10 mg of potassium nitrate (KNO_3), 5 mg of ammonium nitrate (NH_4NO_3), 3 mg of
sodium beta-glycerophosphate pentahydrate (β-Na2glycerophosphate $\cdot$ $5H_2O$), 2 mg of
magnesium sulfate heptahydrate ($MgSO_4 \cdot 7H_2O$), 0.01 µg of vitamin B_{12}, 0.01 µg of
biotin, 1 µg of thiamine HCl, 0.1 mL of PIV metal solution, 0.1 mL of Fe (as EDTA; 1:1
molar), and 40 mg of HEPES. The PIV metal solution contains the following reagents in
100 mL of distilled water: 100 mg $Na_2EDTA \cdot 2H_2O$, 19.6 mg $FeCl_3 \cdot 6H_2O$, 3.6 mg
$MnCl_2 \cdot 4H_2O$, 1.04 mg $ZnCl_2$, 0.4 mg $CoCl_2 \cdot 6H_2O$ and 0.25 mg $Na_2MoO_4 \cdot 2H_2O$ (pH
7.2, 0% NaCl; Kasai et al. 2004). The culture conditions were maintained at 22 °C
under a 14 h light/10 h dark cycle. The light intensity was about 40 µmol photons m^{-2} s^{-1}.
122
Establishment of F_1 Populations
We grew cells from the parental strains, NIES-64 and NIES-65, vegetably for 10 days
and then they were washed two times with a nitrogen-depleted mating-inducing medium
(MI medium; Ichimura 1971). The MI medium contains the following reagents in 99.7
mL of distilled water: 4 mg of $MgSO_4 \cdot 7H_2O$, 5 mg of β-Na2glycerophosphate $\cdot$ $5H_2O$,
10 mg of $CaCl_2 \cdot 2H_2O$, 0.01 µg of vitamin B_{12}, 0.01 µg of Biotin, 1 µg of thiamine HCl,
0.3 mL of PIV metals, and 50 mg of Tris (hydroxymethyl) aminomethane. Subsequently,
the two strains were mixed and incubated in the MI medium at a concentration of
10,000 cells/ml for each strain. This mixture was exposed to 24 h continuous light for
10 days. The zygotes were then kept in the dark conditions for a week until they were
completely dried out (note that they have desiccation tolerance). After a week, we added
the CA medium to induce germination of the zygospores. Through these processes, we
established eight F_1 populations in total. Populations F_1-A through F_1-E germinated
from 1,000 zygospores each, whereas F_1-F germinated from 100 zygospores and F_1-G
and F_1-H germinated from 10,000 zygospores.
138
Culturing of Populations
The populations of NIES-64, NIES-65, and eight F_1 populations were cultured in six-

well plates at a density of 2,500 cells per 5 mL CA medium (Fig. S1). The culture

conditions varied across combinations of NaCl concentrations (0, 0. 05, 0.1, and 0.5%)

and pH levels (6.2, 7.2, and 8.2), resulting in a total of 12 different conditions. For

reference, 0.5% NaCl corresponds to 5g/L NaCl and ca. 0.0856 M whereas seawater is

about 3% NaCl (ca. 0.5 M). Each condition was replicated five times.

Measurement of Growth Rate

To calculate the population growth rate, algal biomass was measured on the first day

(day 0; the day the cells were introduced to each medium), as well as on days 1, 2, and 6,

using a Multiskan SkyHigh TCD (Thermo) at an absorbance of 684 nm as like previous

experimental studies on *Chlamydomonas* (Lachapelle and Bell 2012, Petkovic and

Colegrave 2019, 2023). As the pigment content variation of a cell may bias the

estimation of algal biomass, we used a wavelength where absorbance by the pigment is

small (Griffiths et al. 2011). We also checked the correlation between the absorbance

and the number of cells (Fig. S2). Before every measurement, the cells were agitated by

pipetting. To eliminate the effects of the background, the absorbance values of media

without cells were subtracted from all measurements.

The biomass dynamics was modeled using nonlinear least-squares fitting in R

(version 4.3.2) (R Core Team 2023) as follows:

$$N_t = N_0 e^{rt},$$ (1)

where N_t represents the algal biomass (absorbance values) on day t, r is the biomass

growth rate, t denotes the number of days, and N_0 is the algal biomass on day 0.

Hereafter we use "population growth" to represent biomass growth dynamics (Fig. S3).

To investigate whether the variation in growth rates in the F_1 populations is

significantly larger than those expected from the parental (NIES-64 and NIES-65)

populations, we calculated a mean and a standard deviation (SD) in population growth

rates of the two parental populations. Then we sampled F_1 individuals from a normal

distribution with the calculated mean and SD. This is an expected distribution of F_1

growth rates, and we calculated a SD of the distribution. The simulation was repeated

10,000 times to construct a null distribution of the SDs. We then examined whether the

observed SDs of the F_1 population significantly deviated from these simulated

distributions (whether $p < 0.01$ or not).

Results

We first analyzed how salinity and low pH affect population growth rates, r. To simplify

our analyses, we focused on conditions with 0.05%, 0.1%, and 0.5% NaCl and pH 6.2

and 7.2 in the main text (Fig. 2). See Fig. S3 and S4 for the entire dataset including

conditions with 0% NaCl and/or pH 8.2. We found that both stresses decreased

population growth rates significantly (two-way ANOVA: salinity; $F_{2,291} = 101$, $p <$

0.001, pH; $F_{1,291} = 46.5$, $p < 0.001$, Fig. 2, Table S1). In addition, the effects of stressors

were not additive (salinity × pH interaction; $F_{2,291} = 7.24$, $p < 0.001$, Fig. 2, Table S1),

indicating the importance of considering two stressors simultaneously. The results from

Fig. 2 were consistent with those based on the entire dataset (Table S2). As expected,

we found that combining two types of stressors can further decrease population growth

rates: for example, most populations showed positive growth rates in the condition with

0.05% NaCl and pH 6.2 (Fig. 2A, S4B) as well as that with 0.1% NaCl and pH 7.2 (Fig.

2E, S4G), but many of them showed negative growth rates in the condition with 0.1%

NaCl and pH 6.2 (Fig. 2B, S4C). Growth responses of the two parental populations to

stressors can be sometimes similar (Fig. 2B-C), but sometimes different (Fig. 2E).

Under the 0.1% NaCl and pH 7.2 condition (Fig. 2E, S4G), the NIES-64 strain showed

a significantly higher growth rate than the NIES-65 strain (Student's t-test: t-value =

3.74, d.f. = 7, $p = 0.00726$).

Under the stressful conditions, some F_1 populations exhibited larger variation

in growth rates compared with the parental ones (Fig. 2). Specifically, under conditions

of pH 6.2 with 0%, 0.05%, 0.1%, and 0.5% NaCl; pH 7.2 with 0%; and pH 8.2 with

0.5% NaCl, the F_1 populations showed significantly greater standard deviations in

growth rates than those expected from parental populations ($p < 0.01$, Fig. 2A-C and

S4A-E and L, see Materials and Methods section for details). For example, in the

condition with 0.05% NaCl and pH 6.2, the two parental populations showed very

similar positive growth rates whereas the F_1 populations exhibited large variation of

growth rates: the variation is so large that some of them are even negative (Fig. 2A,

S4B). Furthermore, some F_1 populations demonstrated positive growth rates even under

severe stress conditions where the parental populations showed negative growth rates

(0.1% NaCl and pH 6.2, Fig. 2B, S4C). Even in the severest condition (0.5% NaCl and

pH 6.2), some F_1 populations showed positive growth rates (Fig. 2C, S4D). We also

found that F_1 populations germinated from larger number of zygospores exhibited

higher growth rates in two conditions (Fig. S5D, S5F), but it is not always the case (see

Fig. S5 for details).

Discussion

Sexual reproduction has been a central enigma in evolutionary biology because of the

many costs of sex (Williams 1975, Maynard Smith 1978, Bell 1982, Hartfield and

Keightley 2012, Lehtonen et al. 2012). Recent studies have shown that sexual

reproduction is important for organisms to adapt to rapidly changing environments

(Colegrave 2002, Kaltz and Bell 2002, Goddard et al. 2005, Morran et al. 2009, Becks

and Agrawal 2010, Gray and Goddard 2012) and for preventing population extinction

(Lachapelle and Bell 2012, Petkovic and Colegrave 2019, 2023), but it was not clear

whether a single episode of sexual reproduction can produce large amount of genetic

variation. By using the unicellular green alga, *Closterium peracerosum–strigosum–*

littorale complex, we have shown large variation in population growth rates of F_1

populations (Fig. 2, S4). Even when parental populations show negative growth rates,

some F_1 populations show positive growth rates in severe environmental conditions

(e.g., the condition with 0.1% NaCl and pH 6.2, Fig. 2B, S4C). Our results indicate that,

although asexual reproduction is more efficient in population growth, sexual

reproduction is important for growth in stressful environments and for evolutionary

rescue.

Theory suggests that recombination due to sexual reproduction has two

antagonistic effects on evolutionary rescue: it generates and breaks up favorable gene

combinations (Uecker and Hermisson 2016, Orive et al. 2017). The former is related to

the Fisher-Muller hypothesis where sexual recombination can combine beneficial alleles

into the same genome (Fisher 1930, Muller 1932). Our results are consistent with the

two classical ideas: sexual reproduction tended to increase the variance in population

growth rates, and it resulted in positive (or negative) growth of some F_1 populations

even when their parental populations showed negative (or positive) growth (Fig. 2, S4).

Genome information recently obtained for this algal species (*C. psl.* complex) has

revealed that approximately 30% of the genes in the parental strains showed copy
number differentiation (Kawaguchi et al. 2023). Variation in the number of gene copies
may produce large variation in the F_1 generation through recombination (Hastings et al.
2009). Thus, it will be interesting to carefully consider how genetic architecture,
including structural variation, affects evolutionary rescue in future studies
(Gomulkiewicz et al. 2010, Yamamichi 2022).

When we regard population growth rate as a quantitative trait, it is possible to
consider our results as an example of transgressive segregation (Rieseberg et al. 1999).
Previous studies tended to focus on its effects on macroevolution such as speciation and
adaptive radiation (e.g., Kagawa and Takimoto 2018), but our results suggest that it may
affect rapid contemporary evolution as well as evolutionary rescue. Kawaguchi et al.
(2023) estimated the genome sizes of the F_1 populations between NIES-64 and NIES-65
and found that they were intermediate values between those of their parental
populations (Fig. 3a of Kawaguchi et al. 2023). This is consistent with the results on
population growth rates in, for example, Fig. 2E, but not consistent with the results
under low pH (Fig. 2A-C). Thus, it will be important to explore causes of the difference
between the estimated genome size and population growth rates in future studies. In this
context, it will be helpful to conduct linkage mapping to understand which genes are
responsible for population growth rates. Green algae are suitable as they grow as
haploids, and thus it should be easier to detect genes responsible for a quantitative trait
without considering allelic dominance.

We observed that the green alga grew very well in high pH medium (when pH
was 8.2, Fig. S3, S4), which is consistent with previous studies on some species of the
genus *Closterium* (Gough 1977, Renstrom 1979, Greenwood and Lowe 2006). On the
other hand, low pH reduced growth rates significantly as expected (when pH was 6.2,
Fig. 2, S3, S4). Interestingly, the F_1 populations tended to have larger variation in
growth rates in low pH environments (Fig. 2A-C). This may be seen as an example of
cryptic genetic variation, which has little or no effect on phenotype normally but
generates phenotypic variation under novel conditions (Paaby and Rockman 2014).
Previous studies on *Chlamydomonas* species have suggested that genes encoding heat-
shock proteins and plasma membrane H^+-ATPase were attributable to adaptation to low
pH environments (Gerloff-Elias et al. 2006, Hirooka et al. 2017). In the genus
Klebsormidium, belonging to streptophyte algae (as like *Closterium*), it has been

demonstrated that the ability to adapt to low pH conditions has evolved multiple times
independently (Škaloud et al. 2014). These previous studies, combined with our
findings, imply that adaptation to low pH environments may be related to cryptic
genetic variation that can promote rapid adaptation.

Under high salt stress, we also observed a notable reduction in population
growth rate (Fig. 2, S3, S4). Salt is known to impose osmotic and oxidative stresses by
disrupting the homeostasis of ions and inhibits growth by decreasing the rates of
photosynthesis (Husic and Tolbert 1986, Neelam and Subramanyam 2013) and promote
population extinction in *Chlamydomonas* (Lachapelle and Bell 2012, Petkovic and
Colegrave 2019, 2023). However, unlike low pH, salt stress alone did not result in
increased variation in growth rates of F_1 populations (Fig. 2D-F). It will be insightful to
compare genes responsible for pH and salt stresses and investigate how genetic
architecture, stressors, and recombination interact to produce transgressive segregation
and evolutionary rescue.

While previous studies tended to focus on a single environmental factor to
understand its deteriorating effect on population growth and evolutionary rescue
(Kinnison and Hairston 2007, Alexander et al. 2014, Carlson et al. 2014, Bell 2017),
recent studies proposed that multiple stressors are simultaneously affecting ecosystem
dynamics in the Anthropocene (Orr et al. 2020). Therefore, it will be important to
consider interacting multiple stressors in future studies on evolutionary rescue to deepen
our understanding on conservation of endangered species as well as management of
target species for hunting and fisheries. For archiving this goal, it will be essential to
utilize genome information to reveal the genetic architecture underlying adaptation. If
there are two or more stresses, more loci will affect phenotypic adaptation to the
degraded environments, and it will further make sexual recombination important for
producing combinations better adapted to the environmental changes.
299
300

301 **Author contributions**
302 Conceptualization: MY and YWK; Experiments and data analyses: YWK; Funding
303 acquisition: MY; Writing-original draft: MY; Writing-reviewing and editing: MY and
304 YWK.
305

Acknowledgments

C. psl. complex (NIES-64 and NIES-65) were provided by NIES through the National BioResource Project (NBRP) of the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan. We thank M. Aono, S. Shibasaki, and N. G. Hairston Jr. for their helpful comments and laboratory works.

Funding

YWK is supported by Japan Society for the Promotion of Science (JSPS) Grants-in-Aid for Scientific Research (KAKENHI) JP20J20877 and the National Institute of Genetics (NIG) Postdoctoral Research Fellowship. MY is supported by JSPS KAKENHI JP19K16223, JP20KK0169, JP21H02560, JP22H02688, and JP22H04983, Japan Science and Technology Agency (JST) Core Research for Evolutional Science and Technology (CREST) JPMJCR23N5, and Australian Research Council (ARC) Discovery Project DP220102040.

Conflict of interest

Authors declare that they have no competing interests.

References

Alexander, H. K., G. Martin, O. Y. Martin, and S. Bonhoeffer. 2014. Evolutionary rescue: linking theory for conservation and medicine. Evolutionary Applications **7**:1161-1179.

Becks, L., and A. F. Agrawal. 2010. Higher rates of sex evolve in spatially heterogeneous environments. Nature **468**:89-92.

Bell, G. 1982. The Masterpiece of Nature: The Evolution and Genetics of Sexuality. University of California Press, Berkeley, CA.

Bell, G. 2017. Evolutionary rescue. Annual Review of Ecology, Evolution, and Systematics **48**:605-627.

Bell, G., and A. Gonzalez. 2009. Evolutionary rescue can prevent extinction following environmental change. Ecology Letters **12**:942-948.

Carlson, S. M., C. J. Cunningham, and P. A. H. Westley. 2014. Evolutionary rescue in a changing world. Trends in Ecology & Evolution **29**:521-530.

Colegrave, N. 2002. Sex releases the speed limit on evolution. Nature **420**:664-666.

Colegrave, N., O. Kaltz, and G. Bell. 2002. The ecology and genetics of fitness in
Chlamydomonas. VIII. The dynamics of adaptation to novel environments after
a single episode of sex. Evolution **56**:14-21.

Darwin, C. 1859. On the Origin of Species by Means of Natural Selection, or the
Preservation of Favoured Races in the Struggle for Life. John Murray, London.

Fisher, R. A. 1930. The Genetical Theory of Natural Selection. The Clarendon Press,
Oxford, U.K.

Gerloff-Elias, A., D. Barua, A. Mölich, and E. Spijkerman. 2006. Temperature-and pH-
dependent accumulation of heat-shock proteins in the acidophilic green alga
Chlamydomonas acidophila. FEMS Microbiology Ecology **56**:345-354.

Goddard, M. R., H. C. J. Godfray, and A. Burt. 2005. Sex increases the efficacy of
natural selection in experimental yeast populations. Nature **434**:636-640.

Gomulkiewicz, R., and R. D. Holt. 1995. When does evolution by natural selection
prevent extinction? Evolution **49**:201-207.

Gomulkiewicz, R., R. D. Holt, M. Barfield, and S. L. Nuismer. 2010. Genetics,
adaptation, and invasion in harsh environments. Evolutionary Applications **3**:97-
108.

Gough, S. B. 1977. The growth of selected desmid (Desmidiales, Chlorophyta) taxa at
different calcium and pH levels. American Journal of Botany **64**:1297-1299.

Gray, J. C., and M. R. Goddard. 2012. Sex enhances adaptation by unlinking beneficial
from detrimental mutations in experimental yeast populations. BMC
Evolutionary Biology **12**:43.

Greenwood, J. L., and R. L. Lowe. 2006. The effects of pH on a periphyton community
in an acidic wetland, USA. Hydrobiologia **561**:71-82.

Griffiths, M. J., C. Garcin, R. P. van Hille, and S. T. L. Harrison. 2011. Interference by
pigment in the estimation of microalgal biomass concentration by optical density.
Journal of Microbiological Methods **85**:119-123.

Hairston, N. G., Jr., S. P. Ellner, M. A. Geber, T. Yoshida, and J. A. Fox. 2005. Rapid
evolution and the convergence of ecological and evolutionary time. Ecology
Letters **8**:1114-1127.

Hartfield, M., and P. D. Keightley. 2012. Current hypotheses for the evolution of sex
and recombination. Integrative zoology **7**:192-209.

Hastings, P. J., J. R. Lupski, S. M. Rosenberg, and G. Ira. 2009. Mechanisms of change
in gene copy number. Nature Reviews Genetics **10**:551-564.

Hendry, A. P. 2017. Eco-evolutionary Dynamics. Princeton University Press, Princeton,
NJ.

Hirooka, S., Y. Hirose, Y. Kanesaki, S. Higuchi, T. Fujiwara, R. Onuma, A. Era, R.
Ohbayashi, A. Uzuka, H. Nozaki, H. Yoshikawa, and S.-y. Miyagishima. 2017.
Acidophilic green algal genome provides insights into adaptation to an acidic
environment. Proceedings of the National Academy of Sciences of the United
States of America **114**:E8304-E8313.

Husic, H. D., and N. E. Tolbert. 1986. Effect of osmotic stress on carbon metabolism in
Chlamydomonas reinhardtii: accumulation of glycerol as an osmoregulatory
solute. Plant Physiology **82**:594-596.

Ichimura, T. 1971. Sexual cell division and conjugation-papilla formation in sexual
reproduction of *Closterium strigosum*. Pages 208-214 *in* Proceedings of the 7th
International Seaweed Symposium. The University of Tokyo Press, Tokyo.

Isanta-Navarro, J., N. G. Hairston, Jr., J. Beninde, A. Meyer, D. Straile, M. Möst, and D.
Martin-Creuzburg. 2021. Reversed evolution of grazer resistance to
cyanobacteria. Nature Communications **12**:1945.

Kagawa, K., and G. Takimoto. 2018. Hybridization can promote adaptive radiation by
means of transgressive segregation. Ecology Letters **21**:264-274.

Kaltz, O., and G. Bell. 2002. The ecology and genetics of fitness in *Chlamydomonas*.
XII. Repeated sexual episodes increase rates of adaptation to novel
environments. Evolution **56**:1743-1753.

Kasai, F., M. Kawachi, M. Erata, and M. M. Watanabe. 2004. NIES-Collection list of
strains seventh edition microalgae and protozoa. Res. Rep. NIES **182**:1-257.

Kawaguchi, Y. W., Y. Tsuchikane, K. Tanaka, T. Taji, Y. Suzuki, A. Toyoda, M. Ito, Y.
Watano, T. Nishiyama, and H. Sekimoto. 2023. Extensive copy number variation
explains genome size variation in the unicellular Zygnematophycean alga,
Closterium peracerosum–strigosum–littorale complex. Genome Biology and
Evolution **15**:evad115.

Kinnison, M. T., and N. G. Hairston, Jr. 2007. Eco-evolutionary conservation biology:
contemporary evolution and the dynamics of persistence. Functional Ecology
21:444-454.

Lachapelle, J., and G. Bell. 2012. Evolutionary rescue of sexual and asexual populations
406   in a deteriorating environment. Evolution **66**:3508-3518.
407   Lehtonen, J., M. D. Jennions, and H. Kokko. 2012. The many costs of sex. Trends in
408   Ecology & Evolution **27**:172-178.
409   Lindsey, H. A., J. Gallie, S. Taylor, and B. Kerr. 2013. Evolutionary rescue from
410   extinction is contingent on a lower rate of environmental change. Nature
411   **494**:463-467.
412   Maynard Smith, J. 1978. The Evolution of Sex. Cambridge University Press,
413   Cambridge, U.K.
414   Morran, L. T., M. D. Parmenter, and P. C. Phillips. 2009. Mutation load and rapid
415   adaptation favour outcrossing over self-fertilization. Nature **462**:350-352.
416   Muller, H. J. 1932. Some genetic aspects of sex. The American Naturalist **66**:118-138.
417   Neelam, S., and R. Subramanyam. 2013. Alteration of photochemistry and protein
418   degradation of photosystem II from *Chlamydomonas reinhardtii* under high salt
419   grown cells. Journal of Photochemistry and Photobiology B: Biology **124**:63-70.
420   Nojiri, T., T. Fujii, and H. Sekimoto. 1995. Purification and characterization of a novel
421   sex pheromone that induces the release of another sex pheromone during sexual
422   reproduction of the heterothallic *Closterium peracerosum-strigosum-littorale*
423   complex. Plant and cell physiology **36**:79-84.
424   Ohtaka, K., and H. Sekimoto. 2023. Zygnematophycean algae: Possible models for
425   cellular and evolutionary biology. Pages 59-68 *in* Seminars in Cell &
426   Developmental Biology. Elsevier.
427   Orive, M. E., M. Barfield, C. Fernandez, and R. D. Holt. 2017. Effects of clonal
428   reproduction on evolutionary lag and evolutionary rescue. The American
429   Naturalist **190**:469-490.
430   Orr, H. A., and R. L. Unckless. 2014. The population genetics of evolutionary rescue.
431   PLOS Genetics **10**:e1004551.
432   Orr, J. A., R. D. Vinebrooke, M. C. Jackson, K. J. Kroeker, R. L. Kordas, C. Mantyka-
433   Pringle, P. J. Van den Brink, F. De Laender, R. Stoks, M. Holmstrup, C. D.
434   Matthaei, W. A. Monk, M. R. Penk, S. Leuzinger, R. B. Schäfer, and J. J. Piggott.
435   2020. Towards a unified study of multiple stressors: divisions and common
436   goals across research disciplines. Proceedings of the Royal Society B:
437   Biological Sciences **287**:20200421.

Paaby, A. B., and M. V. Rockman. 2014. Cryptic genetic variation: evolution's hidden
substrate. Nature Reviews Genetics **15**:247-258.

Petkovic, N., and N. Colegrave. 2019. Sex increases the probability of evolutionary
rescue in the presence of a competitor. Journal of Evolutionary Biology
32:1252-1261.

Petkovic, N., and N. Colegrave. 2023. The effects of sex on extinction dynamics of
Chlamydomonas reinhardtii depend on the rate of environmental change.
Journal of Evolutionary Biology **36**:1783-1795.

R Core Team. 2023. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.

Renstrom, D. B. 1979. Culture and growth of *Closterium ehrenbergii* (Desmidiaceae).
Portland State University.

Rieseberg, L. H., M. A. Archer, and R. K. Wayne. 1999. Transgressive segregation,
adaptation and speciation. Heredity **83**:363-372.

Sanderson, S., M.-O. Beausoleil, R. E. O'Dea, Z. T. Wood, C. Correa, V. Frankel, L. D.
Gorné, G. E. Haines, M. T. Kinnison, K. B. Oke, F. Pelletier, F. Pérez-Jvostov, W.
D. Reyes-Corral, Y. Ritchot, F. Sorbara, K. M. Gotanda, and A. P. Hendry. 2022.
The pace of modern life, revisited. Molecular Ecology **31**:1028-1043.

Schoener, T. W. 2011. The newest synthesis: understanding the interplay of evolutionary
and ecological dynamics. science **331**:426-429.

Sekimoto, H., Y. Sone, and T. Fujii. 1994. Regulation of expression of the genes for a
sex pheromone by an inducer of the sex pheromone in the *Closterium*
peracerosum-strigosum-littorale complex. Planta **193**:137-144.

Škaloud, P., A. Lukešová, V. Malavasi, D. Ryšánek, K. Hrčková, and F. Rindi. 2014.
Molecular evidence for the polyphyletic origin of low pH adaptation in the
genus *Klebsormidium* (Klebsormidiophyceae, Streptophyta). Plant Ecology and
Evolution **147**:333-345.

Thompson, J. N. 1998. Rapid evolution as an ecological process. Trends in Ecology &
Evolution **13**:329-332.

Tsuchikane, Y., and H. Sekimoto. 2019. The genus *Closterium*, a new model organism
to study sexual reproduction in streptophytes. New Phytologist **221**:99-104.

Uecker, H., and J. Hermisson. 2016. The role of recombination in evolutionary rescue.
Genetics **202**:721-732.

Vander Wal, E., D. Garant, M. Festa-Bianchet, and F. Pelletier. 2013. Evolutionary

rescue in vertebrates: evidence, applications and uncertainty. Philosophical

Transactions of the Royal Society B: Biological Sciences **368**:20120090.

Wickett, N. J., S. Mirarab, N. Nguyen, T. Warnow, E. Carpenter, N. Matasci, S.

Ayyampalayam, M. S. Barker, J. G. Burleigh, M. A. Gitzendanner, B. R. Ruhfel,

E. Wafula, J. P. Der, S. W. Graham, S. Mathews, M. Melkonian, D. E. Soltis, P.

S. Soltis, N. W. Miles, C. J. Rothfels, L. Pokorny, A. J. Shaw, L. DeGironimo, D.

W. Stevenson, B. Surek, J. C. Villarreal, B. Roure, H. Philippe, C. W.

dePamphilis, T. Chen, M. K. Deyholos, R. S. Baucom, T. M. Kutchan, M. M.

Augustin, J. Wang, Y. Zhang, Z. Tian, Z. Yan, X. Wu, X. Sun, G. K.-S. Wong,

and J. Leebens-Mack. 2014. Phylotranscriptomic analysis of the origin and early

diversification of land plants. Proceedings of the National Academy of Sciences

of the United States of America **111**:E4859-E4868.

Williams, G. C. 1975. Sex and Evolution. Princeton University Press, Princeton, NJ.

Yamamichi, M. 2022. How does genetic architecture affect eco-evolutionary dynamics?

A theoretical perspective. Philosophical Transactions of the Royal Society B:

Biological Sciences **377**:20200504.

Figures

Fig. 1. Green alga, *Closterium peracerosum–strigosum–littorale* complex, and its life
cycle.

(A) A cell of *C. psl.* complex (NIES-65). (B) The life cycle of *C. psl.* complex, with
mating types "mt$^+$" (mating type plus) and "mt$^-$" (mating type minus). Chromosomes
from parental strains, mt$^-$ and mt$^+$, are colored by orange and red, respectively. The
symbols "n" and "2n" denote haploid and diploid cells, respectively. Haploid cells
proliferate via vegetative reproduction, whereas sexual reproduction between different
mating types produces diploid zygospores. Then, haploid cells (F$_1$) were produced from
the diploid zygospores through meiosis.

Fig. 2. Population growth rates under each stress condition.

Blue and orange points (and boxplots) denote parental (NIES-64 and NIES-65) and F_1 populations, respectively. Bars and boxes represent the median and interquartile range, respectively. Whiskers extend to 1.5 times the interquartile range. A, D: 0.05% NaCl. B, E: 0.1% NaCl. C, F: 0.5% NaCl. A-C: pH 6.2. D-F: pH 7.2. Under condition marked with an asterisk (A, B, C), the F_1 populations showed significantly larger variation in growth rates than the parental ones ($p < 0.01$). See Fig. S3 and S4 for more details and Fig. S5 for the effects of the founder population size.