Exogenous myristate fuels the growth of symbiotic arbuscular mycorrhizal fungi

but disrupts their carbon-phosphorus exchange with host plants

Hanwen Chen[1, #], Tian Xiong[1, #], Baoxing Guan[1, #], Jiaqi Huang[1, #], Danrui Zhao[1], Yao

Chen[1], Haoran Liang[1], Yingwei Li[1], Jingwen Wu[1], Shaoping Ye[2], Ting Li[2], Wensheng

Shu[1], Jin-tian Li[1], Yutao Wang[1, *]

[1]Guangdong Provincial Key Laboratory of Biotechnology for Plant Development,

Guangzhou Key Laboratory of Subtropical Biodiversity and Biomonitoring, School of

Life Sciences, South China Normal University, Guangzhou, 510631, China

[2]Guangzhou Institute of Forestry and Landscape Architecture, Guangzhou, 510405,

China

*Correspondence: School of Life Sciences, South China Normal University,

Guangzhou, 510631, China; Email: wangyutao@scnu.edu.cn (Yutao Wang); Tel./Fax:

+86 20 85211850

[#]: These authors contributed equally to this work.

ABSTRACT

Arbuscular mycorrhizal fungi (AMF) are obligate biotrophs that rely on host-

derived symbiotic carbohydrates. However, it remains unclear whether symbiotic AMF

can access exogenous non-symbiotic carbon sources, complicating our understanding

of their relationship with host plants. Here, we investigated the direct uptake of

exogenous $^{13}C_1$-labeled myristate by three symbiotic AMF species (*Rhizophagus*

irregularis, *R. intraradices*, and *R. diaphanous*) and assessed their growth responses

using AMF-carrot hairy root co-culture systems. Furthermore, we explored the

environmental distribution of myristate, and evaluated the impact of exogenous

myristate on the carbon-phosphorus exchange between *R. irregularis* and alfalfa or rice

in a greenhouse experiment. Symbiotic AMF can absorb exogenous myristate, as

evidenced by ^{13}C enrichment and transcriptional activation of fatty acid transport and

metabolism genes in AMF extraradical hyphae. Myristate is commonly present in

various soil and plant environments, and its application increased both intraradical and

extraradical fungal biomass, possibly linked to suppressed mycorrhizal-activated

defense responses in host roots. Unexpectedly, exogenous myristate reduced the

mycorrhizal phosphorus benefits for both alfalfa and rice and decreased their symbiotic

carbon allocation to root-colonizing AMF, although these effects varied with soil

phosphorus conditions. These findings provide new insights into understanding and

manipulating the nutritional interactions between AMF and host plants.

Keywords：arbuscular mycorrhizal fungi (AMF); carbon-phosphorus exchange;

exogenous non-symbiotic carbon; myristate; stable isotope labeling

INTRODUCTION

Arbuscular mycorrhizal fungi (AMF) are among the most widely distributed soil microorganisms, forming symbiotic relationships with approximately 70% of terrestrial vascular plants (Brundrett and Tedersoo, 2018). In this symbiosis, referred to as arbuscular mycorrhiza (AM), AMF develop an extensive extraradical hyphal network in the soil, which facilitates nutrient uptake for host plants—particularly phosphorus (P)—and enhances plant resilience to both biotic and abiotic stressors (Smith and Read, 2008). In return, host plants allocate 4−20% of their photosynthetically fixed carbon (C) to their fungal partners in the form of sugars and fatty acids (Helber et al., 2011; Jiang et al., 2017). This symbiotic C-P exchange, the cornerstone of AM symbiosis (Smith and Read, 2008), not only plays a pivotal role in the biogeochemical cycling of C and P elements but also acts as a key mechanism through which AMF influence plant community structure and productivity (Johnson, 2010; Wipf et al., 2019).

As one of the most ancient symbiotic relationships, AM has undergone extensive evolutionary processes (Foster and Kokko, 2006; Rich et al., 2021), with its establishment and development intricately regulated by a set of symbiotic genes and signaling molecules influenced by both biotic and abiotic factors (Maclean et al., 2017; Shi et al., 2021). For instance, in AM plants, the two pathways for P uptake—the direct P pathway via root hairs and epidermis and the mycorrhizal P pathway via AMF hyphae—are coordinately regulated by the same phosphate-sensing pathway (Shi et al., 2021; Das et al., 2022). Under P-deficiency conditions, host plants synthesize and release strigolactones into the soil, activating the mycorrhizal P pathway by stimulating the elongation and branching of AMF hyphae (Mori et al., 2016; Yu et al., 2022). Additionally, the extent of AMF colonization and the benefits conferred by AMF largely depend on the host plant species (genotypes), with some being "responsive plants" that

exhibit positive growth responses (e.g., alfalfa, maize) and others being "nonresponsive plants" that show minimal or no growth enhancement while exhibiting increased AM-activated defense responses (e.g., rice, wheat; Hata et al., 2010; Li et al., 2024).

AMF are considered obligate symbionts that rely on living host plants to complete their life cycle by producing daughter spores (Smith and Read, 2008), which presents challenges for basic research on AM and their applications in agriculture and other fields. To date, achieving a pure (plant-independent) culture of asymbiotic AMF has not been successful, partially due to their dependence on plant-derived symbiotic C (Bravo et al., 2017; Luginbuehl et al., 2017; Kameoka et al., 2019). Genomic and transcriptomic analyses have shown that AMF lack the genes required for the biosynthesis of long-chain fatty acids, and subsequent studies have demonstrated that fatty acids are transferred from host plants to AMF as their primary C source (Bravo et al., 2017; Jiang et al., 2017; Keymer et al., 2017; Luginbuehl et al., 2017). However, the mechanism underlying the allocation of plant-derived fatty acids to fungi in AM symbiosis remains unclear. In host plants, AMF colonization triggers the expression of lipid biosynthesis and transport genes (Gutjahr et al., 2012; Bravo et al., 2017; Keymer et al., 2017; Luginbuehl et al., 2017), with lipid transport to the AM symbiotic interface relying on the periarbuscular membrane-localized transporters stunted arbuscule (STR) and STR2 (Zhang et al., 2010; Gutjahr et al., 2012), and fungal uptake of plant-derived lipids mediated by fatty acid transporter 1 (RiFAT1) and RiFAT2 in the model AMF species *Rhizophagus irregularis* (Brands and Dörmann, 2022). This symbiotic C allocation is functionally linked to fungal P delivery within a framework of mutual nutritional rewards (Kiers et al., 2011), which have been disrupted during AMF-nonhost interactions due to the strong AM-activated defense responses in non-host roots (Wang et al., 2023).

Fatty acids and their derivatives, essential structural components and energy

sources for plant, animal, and microbial cells, are commonly found in soil, particularly

in the rhizosphere, where plant-derived fatty acids are made available through root

exudates and rhizodeposition (Swenson et al., 2018; Dai et al., 2022; Liu et al., 2024).

Following the discovery that AMF acquire fatty acids from their hosts during symbiosis,

the use of fatty acids in cultivating asymbiotic AMF has attracted considerable attention.

Kameoka et al. (2019) demonstrated that under asymbiotic conditions, exogenous

palmitoleic acid and several other fatty acids stimulated the hyphal elongation and

branching in *Rhisophagus irregularis* and *R. clarus,* but not in *Gigaspora margarita*,

resulting in the formation of infection-competent but morphologically and functionally

incomplete spores with reduced size, lower germination rates, and decreased infection

capacity compared to symbiotic spores. Additionally, Sugiura et al. (2020) showed that

exogenous myristate (tetradecanoic acid)—a common organic acid in various plant root

exudates (Li et al., 2017; Liu et al., 2024) and soil substrates (Swenson et al., 2018; Dai

et al., 2022)—can serve as a C source for the asymbiotic growth of *R. irregularis* and

R. clarus, promoting hyphal growth, the formation of branched absorbing structures,

and the production of morphologically incomplete, infection-competent spores.

Subsequent studies revealed that combining plant hormones with myristate enhanced

the efficiency of myristate as a C source in asymbiotic *R. clarus*, leading to the

generation of numerous infection-competent spores that were smaller than those

generated under symbiotic conditions (Sugiura et al., 2020; Tanaka et al., 2022). These

discoveries have significantly advanced the pure culture of AMF and, importantly,

identified myristate as the first external C source accessible to AMF under asymbiotic

conditions, providing exceptional opportunities for understanding and utilizing the

ecological functions of AM symbiosis in plant community regulation and global C

cycling (Rillig et al., 2020).

However, the effects of external myristate on the growth and development of

symbiotic AMF—the typical state of functioning AMF in nature—remain unclear.

Regarding sugars, another type of host-derived C source, symbiotic AMF may be

unable to absorb them additionally through extraradical hyphae (Bücking et al., 2008;

Helber et al., 2011; Sugiura et al., 2020). A recent study reported that while exogenous

myristate does not affect the extraradical biomass of symbiotic *R. irregularis* and *R.*

intraradices, it enhances their colonization rates by stimulating the growth and

development of extraradical hyphae. Based on this, the authors inferred that symbiotic

AMF may acquire exogenous myristate (Liu et al., 2023). To date, the effects of

exogenous myristate on AM symbiosis and their underlying mechanisms remain limited.

Specifically, whether symbiotic AMF can uptake exogenous myristate via extraradical

hyphae in the presence of plant-derived symbiotic C is yet to be determined. This

question is crucial for understanding the symbiotic mechanisms and ecological roles of

AM symbiosis (Rillig et al., 2020).

To address this significant knowledge gap, this study investigated the ability of

symbiotic AMF to absorb external non-symbiotic myristate and examined the effect of

exogenous myristate on symbiotic C-P exchanges in AM. Specifically, we aimed to

determine: (i) whether symbiotic AMF can take up exogenous non-symbiotic myristate,

(ii) how this additional non-symbiotic C source influences the symbiotic C allocation

from plants to AMF, and (iii) if and how fungal P transfer to the host plant is affected

by exogenous myristate. This work provides, to our knowledge, the strongest evidence

to date that symbiotic AMF can absorb non-symbiotic C sources like myristate in the

presence of plant-derived symbiotic carbohydrates. Furthermore, our findings reveal

that exogenous myristate may, unexpectedly, disrupt the C-P exchange in AM

symbiosis.

RESULTS

Tracing of ^{13}C in AMF extraradical hyphae (Exp. 1)

In the two-compartment AMF-carrot hairy root co-cultivation system (Exp. 1, Fig.

1a), exogenous myristate specifically induced the formation of branched absorbing

structures in *R. irregularis*, *R. intraradices*, and *R. diaphanus* in the HC approximately

two weeks after Myr$^+$ treatment (Fig. 1b). For *R. irregularis*, the ^{13}C/^{12}C ratio of

extraradical hyphae in HC (trial 1: 1.1504 ± 0.0575%, trial 2: 1.0919 ± 0.0001%) and

RC (trial 1: 1.1026 ± 0.0055%, trial 2: 1.0918 ± 0.0002%) under ^{13}C-Myr treatment

was slightly but significantly higher than that in the corresponding HC (trial 1: 1.0922

± 0.0003%, trial 2: 1.0908 ± 0.0003%) and RC (trial 1: 1.0923 ± 0.0002%, trial 2:

1.0908 ± 0.0002%) from the non-labeled control (^{12}C-Myr), respectively ($P < 0.05$, Fig.

1c). Similarly, for *R. intraradices* and *R. diaphanus*, the ^{13}C/^{12}C ratio of extraradical

hyphae in HC (*R. intraradices*: 1.0933 ± 0.0002%, *R. diaphanus*: 1.0922 ± 0.0005%)

and RC (*R. intraradices*: 1.0927 ± 0.0005%, *R. diaphanus*: 1.0930 ± 0.0004%) under

^{13}C-Myr treatment was slightly but significantly higher than those in HC (*R.*

intraradices: 1.0917 ± 0.0001%, *R. diaphanus*: 1.0908 ± 0.00003%) and RC (*R.*

intraradices: 1.0918 ± 0.00004%, *R. diaphanus*: 1.0908 ± 0.00003%) from the non-

labeled control ($P < 0.01$, Fig. 1c). Additionally, the calculated ^{13}C enrichment in

extraradical hyphae between HC and RC showed no significant difference for *R.*

irregularis and *R. diaphanus* (Fig. 1d, e), although it was higher in HC than in RC for

R. intraradices ($P < 0.05$), excluding the possibility that the ^{13}C enrichment in AMF

extraradical hyphae originated from ^{13}C-myristate retention on the hyphal surface from

the HC. Collectively, these results suggest that these three AMF species can take up a

small amount of exogenous myristate through their extraradical hyphae when

associated with host roots.

In HC, the calculated ^{13}C enrichments in extraradical hyphae of *R. irregularis* (trial

1: 0.5823 ± 0.4111‰, trial 2: 0.0107 ± 0.0014‰), *R. intraradices* (0.0163 ± 0.0031‰),

and *R. diaphanus* (0.0136 ± 0.0067‰) showed no significant differences in the ^{13}C-

Myr treatment (Fig. 1d). In RC, the ^{13}C enrichment in extraradical hyphae also showed

no significant differences (Fig. 1e), except for a slightly but significantly higher ^{13}C

enrichment in *R. diaphanus* (0.0217 ± 0.0031‰) compared to *R. intraradices* (0.0082

± 0.0037‰) and *R. irregularis* in trial 2 (0.0093 ± 0.0017‰, $P < 0.05$). These results

suggest a generally similar myristate absorption capacity among these AMF species

under the tested conditions.

Effects of myristate on AMF colonization and expression of fatty acid uptake and

metabolism marker genes (Exp. 1)

The expression of fatty acid uptake and metabolism marker genes in *R. irregularis*

extraradical hyphae were analyzed to investigate their uptake of myristate from HC at

the transcriptional level. The results showed that exogenous myristate significantly

increased the expression levels of the fungal fatty acid transporter gene *RiFAT2* ($P <$

0.05), while had no significant effect on *RiFAT1* (Fig. 2a). Additionally, exogenous

myristate significantly or marginally significantly increased the expression levels of the

fungal genes involved in fatty acid β-oxidation (*RiFAD1*, $P < 0.05$), palmitvaccenic acid

synthesis (*RiOLE1*, $P = 0.057$), and TCA cycle activity (*RiCIT1*, $P < 0.05$) (Fig. 2b).

These results suggest that exogenous myristate activates the expression of fungal genes

related to fatty acid uptake and metabolism in the extraradical hyphae of *R. irregularis*.

For AMF colonization intensity, exogenous myristate significantly increased the

hyphal and arbuscular colonization rates of *R. irregularis, R. intraradice*, and *R.*

diaphanus in carrot hairy roots ($P < 0.05$, Fig. 2c−e) but had no effect on vesicle

colonization.

Effects of exogenous myristate on AMF spore germination, root colonization, and

extraradical structure development (Exp. 2)

In the single-compartment *R. irregularis*-carrot hairy root co-cultivation system

(Exp. 2, Fig. 3a), exogenous myristate significantly promoted spore germination ($P <$

0.001, Fig. 3b), and enhanced hyphal branching and length ($P < 0.001$, Fig. 3c, d) before

visible physical contact between hyphae and hairy roots at four days after AMF

inoculation. Morphological examinations showed that exogenous myristate

significantly enhanced the colonization intensities of hyphae and arbuscules, but not

vesicles, in carrot hairy roots across various application levels (Fig. 3e). Exogenous

myristate also significantly increased the extraradical biomass of AMF (hyphae and

spore) at different application concentrations ($P < 0.05$, Fig. 3f).

Meanwhile, exogenous myristate significantly increased the density of AMF

spores from eight weeks post-AMF inoculation ($P < 0.05$, Fig. 3g), and the diameter of

newly produced AMF spores in media with exogenous myristate was identical to those

in media without myristate (Fig. 3h).

Environmental myristate levels and the effects of exogenous myristate on AMF

intraradical and extraradical biomass during symbiosis with alfalfa and rice (Exp.

3)

In Exp. 3 (Fig. 4a), our field survey revealed that myristate was extensively

detected in soil and plant samples from the paddy field, grassland, and woodland at a

$2.87–14.68$ mg kg^{-1} level (Fig. 4b). On average, the myristate level in plant (fresh leaf

or root) is mostly higher than that in soil (leaf litter, rhizosphere soil, or bulk soil) ($P <$

0.05, Fig. 4b). Additionally, the myristate levels in leaf litter and rhizosphere soil were

usually greater than in bulk soil ($P < 0.05$), suggesting that plants are significant source

of myristate in the soil. The myristate contents from the same sample type typically

varied across different habitats ($P < 0.05$).

In the pot experiment (Exp. 3), all typical AMF intraradical structures (hyphae,

arbuscules, and vesicles) were observed in the roots of both alfalfa and rice in the *R.*

irregularis-inoculated groups (AM), while no AMF colonization was detected in the

non-inoculated (NM) groups. Under low-P conditions, exogenous myristate

significantly increased arbuscule abundance in alfalfa roots ($P < 0.05$, Fig. 4c) but did

not significantly affect hyphae or vesicle abundance (Supporting Information Fig. S1a,

c). Exogenous myristate significantly enhanced arbuscule, hyphae, and vesicle

abundance in rice roots ($P < 0.05$, Fig. 4d, Supporting Information Fig. S1b, d).

However, under high-P conditions, exogenous myristate had no effect on AMF

abundance in either alfalfa or rice roots (Fig. 4, Supporting Information Fig. S1).

For intraradical AMF biomass, exogenous myristate significantly increased the

levels of the AMF signature fatty acid (NLFA 16:1ω5) in both alfalfa and rice roots

under low-P conditions ($P < 0.05$, Fig. 4e, f); however, it had no significant impact

under high-P conditions. In rice roots, NLFA 16:1ω5 levels were higher under high-P

than low-P conditions ($P < 0.05$). For extraradical AMF biomass, exogenous myristate

increased NLFA 16:1ω5 levels in the rhizosphere soils of alfalfa ($P < 0.05$) but not in

rice ($P = 0.14$) under low-P conditions (Fig. 4e, f). Under high-P conditions, myristate

had no effect on NLFA 16:1ω5 levels in the rhizosphere soil of either plant.

Effects of exogenous myristate on the transfer of symbiotic C to AMF (Exp. 3)

The analysis of ^{13}C-NLFA-isotope ratio mass spectrometry was performed to

investigate the impact of exogenous myristate on the allocation of symbiotic C from

plants to AMF. In AM plants, there was a notable enrichment of ^{13}C (δ^{13}C = 1.75 ±

0.6‰) in the AMF signature fatty acid (NLFA 16:1ω5) compared to NM plants ($\delta^{13}C =$

-24.8 ± 2.3‰), demonstrating the transfer of newly fixed ^{13}C from plants to AMF. The

calculated ^{13}C flow from alfalfa to root-colonizing AMF (0.3400–1.2642 μg) was over

ten times higher than that from rice (0.0261–0.0596 μg), highlighting a substantially

greater efficiency of plant-derived symbiotic C delivery to AMF in AM-responsive

alfalfa compared to non-responsive rice. In alfalfa, exogenous myristate reduced the

symbiotic C transfer to AMF (Myr effect: $P < 0.001$, Fig. 5a), although this effect was

not significant under high-P conditions. In rice, myristate also significantly influenced

the symbiotic C flow to AMF, with its effect significantly interacting with soil P level

(Myr × P: $P = 0.004$, Fig. 5b): myristate significantly increased the allocation of ^{13}C to

AMF under low-P conditions ($P < 0.05$) but decreased it under high-P conditions ($P <$

0.01).

At the transcriptional level, the qRT-PCR analysis revealed that the expressions of

the symbiotic lipid transfer marker genes *STR2* (in alfalfa and rice) and the fungal genes

responsible for symbiotic fatty acid uptake *(RiFAT1* and *RiFAT2)* or monosaccharide

uptake (*Monosaccharide Transporter 2* (*RiMST2*)) were exclusively detected in AM

roots, except for a low expression level of alfalfa *MsSTR2* occasionally observed in NM

roots (Fig. 5c–h). In AMF-inoculated alfalfa roots, exogenous myristate significantly

downregulated the expressions of the symbiotic C transfer genes *MsSTR2* in the plant

and *RiMST2* in AMF ($P < 0.05$), while having no effect on *RiFAT1* and *RiFAT2* in AMF

(Fig. 5c, g, e). In AMF-inoculated rice roots, the effects of myristate on the tested

symbiotic C transfer genes were largely influenced by soil P levels ($P < 0.05$). Under

low-P conditions, myristate significantly increased the expressions of plant *OsSTR2* and

fungal *RiFAT1* and *RiFAT2*, while decreasing the expression of fungal *RiMST2* ($P <$

0.05, Fig. 5d, f, h). Under high-P conditions, myristate significantly increased the

expression of *RiMST2* ($P < 0.05$) but had no effects on *OsSTR2*, *RiFAT1*, and *RiFAT2*.

In both plant species with AMF inoculation, soil P level showed significant effects on

STR, *RiFAT1*, and *MST2* genes ($P < 0.05$), with generally lower expression levels of

these genes under high-P compared to low-P conditions.

Effects of exogenous myristate on the mycorrhizal P pathway and mycorrhizal

growth response (Exp. 3)

In the pot experiments (Exp. 3), AMF inoculation significantly increased P

concentrations in both alfalfa and rice plants ($P < 0.001$, Fig. 6a, b). Under low-P

conditions, exogenous myristate significantly decreased the mycorrhizal P pathway in

both alfalfa and rice ($P < 0.05$, Fig. 6c, d). Under high-P conditions, while exogenous

myristate significantly decreased the mycorrhizal P response in alfalfa ($P < 0.05$), it had

no significant effect on rice.

At the transcriptional level, the expression of mycorrhizal P pathway marker genes

in plants (*Phosphate Transporter*, *MsPT8* or *OsPT11*) and AMF (*RiPT*) was detected

exclusively in the AM roots of rice and alfalfa, except for a low expression level of rice

OsPT11 observed in the NM root under low-P and Myr$^-$ conditions. This finding

confirmed that AM symbiosis activated the mycorrhizal P pathway at the transcriptional

level in both plant species (Fig. 6e, f). Exogenous myristate significantly influenced the

transcriptional responses of the mycorrhizal P pathway in both alfalfa and rice ($P <$

0.01). In alfalfa, exogenous myristate downregulated the expressions of *MsPT8* and

RiPT under both low- and high-P conditions ($P < 0.05$, Fig. 6e). In rice, exogenous

myristate decreased the expression levels of *OsPT11* and *RiPT* under low-P conditions

($P < 0.05$), but not under high-P conditions (Fig. 6f).

In alfalfa, AMF inoculation significantly increased plant biomass under both low-

and high-P conditions ($P < 0.05$, Supporting Information Fig. S2a), with mycorrhizal

growth responses significantly higher under high-P than low-P conditions ($P < 0.05$,

Supporting Information Fig. S2c). In rice, AMF inoculation had no effect on plant

biomass under low-P conditions, but reduced rice biomass under high-P conditions (P

< 0.05); mycorrhizal growth responses were significantly lower under high-P than low-

P conditions ($P < 0.05$, Supporting Information Fig. S2b, d). Exogenous myristate had

no effect on plant biomass or mycorrhizal growth response in either alfalfa or rice, while

soil P levels significantly influenced mycorrhizal growth response: high-P treatment

increased mycorrhizal growth response in alfalfa but decreased it in rice compared to

low-P conditions ($P < 0.01$, Supporting Information Fig. S2c, d).

Effects of exogenous myristate on the transcriptome of AM roots (Exp. 3)

Transcriptome analysis was performed to preliminarily explore the effects of

exogenous myristate on mycorrhizal roots at the transcriptional level. The RNA-seq

data identified 39,944 and 33,056 genes (transcripts) in alfalfa and rice, respectively.

Principal component analysis of gene expression profiles in mycorrhizal roots of alfalfa

and rice showed that the first and second principal components together explained 32.4%

and 44.1% of the variation, respectively (Supporting Information Fig. S3a, b). In both

alfalfa and rice roots, gene expression profiles with or without myristate application

could be partially separated under low-P conditions, but not under high-P conditions,

suggesting the importance of P status in myristate's effect.

Exogenous myristate induced a variety of differentially expressed genes (DEGs,

Myr$^+$ *vs.* Myr$^−$) in mycorrhizal alfalfa (LP: 308, HP: 137) and rice (LP: 590, HP: 855)

roots (Supporting Information Fig. S3c, d). Gene Ontology (GO) analysis of these

DEGs enriched several GO terms related to plant stress or defense, most of which were

downregulated by exogenous myristate (Fig. 7a, b). These included "defense response",

"response to salicylic acid", "peroxidase activity", etc., in rice, and "glycosyltransferase

activity" and "alternative oxidase activity" in alfalfa.

We subsequently analyzed the expression profiles of the component genes

associated with enriched GO terms that are related to defense or stress, using the RNA-

seq dataset (Fig. S4a, b). In alfalfa, exogenous myristate significantly downregulated

several key genes involved in reactive oxygen species balance (*alternative oxidase 2*

(*AOX2*), *AOX3*) and stress tolerance (*UDP-glycose flavonoid glycosyltransferase*

(*UFGT*), *UDP-glycosyltransferase* (*UGT*) *73C3*, *UGT73C6*) under low-P conditions,

as well as typical stress-responsive genes (*peroxidase* (*PRX*) *P7*, *cytochromes P450*

(*CYP*) *83B1*), and gene families such as *geraniol 8-hydroxylase-like* (*G8HL*),

CYP71D10, and *CYP736A12* under high-P conditions (Supplementary Fig. S4a). In rice,

exogenous myristate downregulated several typical defense-associated transcriptional

factors (*TGA2*, *TGA4*, *WRKY45*, *WRKY62*, *WRKY76*, and *bZIP79*) and marker genes

(*pathogenesis-related 5-like* (*PR5L*), *protein phosphatase 2C 35* (*OsPP2C35*), *1-*

aminocyclopropane-1-carboxylate synthase 2 (*ACS2*), *DWARF4*, *pyricularia oryzae*

resistance-ta (*PITA*), *mildew resistance locus O2* (*MLO2*), *NRR repressor homologue*

1 (*RH1*), and *chitinase 1* (*CHT1*)) under low-P conditions (Supplementary Fig. S4b).

However, most of these genes were upregulated following myristate application under

high-P conditions.

qRT-PCR verification of the effects of exogenous myristate on AM-activated

defense genes in mycorrhizal roots (Exp. 3)

To verify the impact of exogenous myristate on the AM-activated defense response,

qRT-PCR was performed to analyze the expression of defense marker genes in both

plant species. In alfalfa roots, exogenous myristate significantly decreased the

expression of *alternative oxidase 3 (MsAOX3)* and *UDP-glycosyltransferase 85A*

(*MsUGT85A*) (*P* < 0.05), while its effect on *peroxidase P7* (*MsPRXP7*) depended on

soil P level (Myr × P: $P < 0.05$): exogenous myristate significantly decreased *MsPRXP7*

expression under high-P conditions ($P < 0.05$) but had no effect under low-P conditions

(Fig. 7c). In rice roots, the effect of exogenous myristate on *chitinase 1* (*OsCHT1*) was

also dependent on soil P level (Myr × P: $P < 0.05$): it marginally decreased *OsCHT1*

expression under low-P conditions ($P = 0.053$) but increased it under high-P conditions

($P = 0.050$). Exogenous myristate significantly suppressed the expression of defense-

related transcriptional factor *OsTGA2* and *peroxidase 39* (*OsPRX39*) under low-P and

high-P conditions, respectively ($P < 0.05$, Fig. 7d). Overall, the qRT-PCR results largely

confirmed the inhibitory effects of myristate on AM-activated defense responses in

alfalfa and low-P rice.

DISCUSSION

AMF are obligate biotrophs that rely on host-derived symbiotic carbohydrates for

reproduction (Smith and Read, 2008). However, whether symbiotic AMF can uptake

external non-symbiotic organic C remains poorly understood (Rillig et al., 2020; Liu et

al., 2023). Under asymbiotic conditions, Sugiura et al. (2020) found that among eight

saturated or unsaturated fatty acids (C12 to C18) and two β-monoacylglycerols supplied,

only myristate led to an increase in the biomass of *R. irregularis*. In our study, we

observed slight yet statistically significant ^{13}C enrichments in the extraradical hyphae

of three AMF species from both hyphal and root compartments during the $^{13}C_1$-

myristate tracing experiment (Fig. 1c), indicating that the extraradical hyphae of the

tested AMF species can absorb external non-symbiotic myristate in the presence of

plant-derived symbiotic carbohydrates. Several additional observations further

supported this finding. First, exogenous myristate specifically induced the formation of

branched absorbing structures in AMF extraradical hyphae (Fig. 1b), which may serve

as preferential sites for myristate uptake (Bago et al., 1998; Sugiura et al., 2020).

Second, exogenous myristate activated the transcription of marker genes associated

with fatty acid uptake and metabolism in extraradical hyphae (Fig. 2). Third, myristate

application frequently enhanced both intraradical and extraradical AMF biomass (Figs.

2, 3, and 4), even when the symbiotic C supply from the plant was reduced (Figs. 4 and

5), accompanied by decreased mycorrhizal P responses (Fig. 6). In a priority study,

myristate application also increased colonization intensity, but not the extraradical

biomass of AMF in tomato roots (Liu et al., 2023). Overall, our findings indicate that

the extraradical hyphae of symbiotic AMF can uptake external non-symbiotic organic

C sources like myristate beyond those directly delivered by the host, at least under the

conditions studied.

Notably, myristate has been frequently detected in root exudates and soil substrates

from non-targeted metabolome analyses (Li et al., 2017; Swenson et al., 2018; Dai et

al., 2022; Liu et al., 2024), although its specific levels in various soil environments

remain largely unknown. Our field survey demonstrates that myristate is widely present

in soil, litter, and plant (Fig. 4b), with its environmental level (2.87−14.68 mg kg^{-1})

encompassing the myristate treatments applied in our greenhouse experiment (Exp. 3,

totaling 1.83 mg per 1000 ml or 300 ml soil substrate in rice and alfalfa, respectively).

Therefore, it is likely that the widespread AMF extraradical hyphae network may

absorb myristate from soil environments such as root exudates, decaying plants, or litter

in realistic environments. Further research is still needed to validate this hypothesis;

however, it effectively explains a long-observed but poorly understood phenomenon—

AMF often colonize and thrive in leaf litter and undecomposed plant material despite

their limitations in assimilating organic forms of nutrients (Went and Stark, 1968;

Zhang et al., 2018; Bunn et al., 2019)—as the myristate concentration in plant and leaf

litter is higher than that in soil (Fig. 4b).

During AM establishment, plant roots typically activate low-level defense

responses to limit excessive C consumption caused by over-colonization (Blilou et al.,

2000; Feng et al., 2019). In our study, myristate increased both intraradical and

extraradical biomass of the symbiotic AMF and promoted the formation of normal-

sized secondary spores (Figs. 2 and 3), unlike the small spores induced by myristate

under asymbiotic conditions (Sugiura et al., 2020). This suggests that plant roots supply

additional substances to AMF beyond myristate, the identification of which is essential

for the pure culture of AMF. Meanwhile, exogenous myristate transcriptionally

downregulated plant genes associated with AM-activated defenses under low-P

conditions (Fig. 7), suggesting that myristate may interfere with normal defense

mechanisms in host roots to facilitate AMF colonization. This effect contrast with AM

signals like strigolactones and quercetin, which enhance plant defenses (Sudheeran et

al., 2020; Kusajima et al., 2022). Further research is required to clarify myristate's role

in regulating AMF colonization and development.

Despite increased arbuscule abundance (Fig. 4)—where symbiotic nutritional

exchange occurs (Smith and Read, 2008)—exogenous myristate unexpectedly reduced

the mycorrhizal P response and downregulated plant and fungal mycorrhizal P

transporter gene expression in both alfalfa and rice roots (Fig. 6), indicating a

diminished mycorrhizal P benefit through inactivation of mycorrhizal P pathway

transcription. Meanwhile, exogenous myristate also reduced C supply to AMF from

both alfalfa and rice, except in low-P rice (Fig. 5). Together, these results suggest that

exogenous myristate disrupts the symbiotic C-P exchange, which is functionally

coupled in AM symbiosis (Kiers et al., 2011). This disruption could be attributed to

several factors. First, it may result from AMF partially shifting from utilizing symbiotic

C to absorbing external C (myristate, Fig. 1), thus reducing their competition for

symbiotic C at the AM interaction interface (Bitterlich et al., 2014). Correspondingly,

this decline in symbiotic C supply to AMF may decrease mycorrhizal P benefit, since

AMF can modulate P transport based on the rate at which plants supply C (Kiers et al.,

2011; Argüello et al., 2016). Second, exogenous myristate may interfere with AM

signaling, as implied by the interference in normal AM-activated defenses (Fig. 7),

thereby limiting nutrient exchange in the symbiosis. Third, AMF P transport proteins

expressed in arbuscular cells can re-uptake P from the symbiotic interface, creating P

competition between the plant and AMF (Xie et al., 2022). In our study, the increased

fungal biomass with exogenous myristate suggests a higher P demand by AMF,

reducing the feasibility of nutritional trade.

In contrast to its effect under high-P treatment, myristate increased, rather than

decreased, the allocation of rice-derived C to AMF under low-P conditions,

accompanied by enhanced expression of marker genes for fatty acid (but not sugar)

allocation from rice to AMF (Fig. 5b, f, g). These results highlight that myristate's

impact on AM symbiosis is influenced by soil P status, a significant regulatory factor

of the symbiosis by coordinating the direct P pathway and mycorrhizal P pathway in

AM plants (Shi et al., 2021; Das et al., 2022). The notable effects of soil P status on

myristate's impact on C-P exchange were also observed in alfalfa (Figs. 5 and 6), where

the unexpectedly higher AMF intraradical biomass (Fig. 4) and greater mycorrhizal

growth response (Fig. S2) in the high-P compared to the low-P treatment are likely due

to the inhibitory effects of extremely low-P environments on AMF colonization and

functioning (Janos, 2007; Wang et al., 2020). Our findings regarding the disruptive

effects of myristate on symbiotic C-P exchange, along with the broad yet varied

presence of myristate in different soil components (Fig. 4), imply that myristate may be

a critical influencing factor in AM symbiosis. Interestingly, myristate also exhibited

inhibitory effects on rhizospheric *Devosia*, which is responsible for promoting growth

and suppressing disease in *Arabidopsis* (Liu et al., 2024). Further research is required

to clarify the mechanisms underlying the disruptive effect of exogenous myristate on

symbiotic C-P exchange and its interaction with soil P.

Despite these encouraging findings, our work has several limitations. First, the

absorption of myristate by symbiotic AMF was observed only after exogenous

application under artificial conditions, which may not accurately reflect natural

environments. Second, our study focused exclusively on myristate and AMF species

from the genus *Rhizophagus*, although different AMF groups may respond to myristate

or other fatty acids in various ways, at least under asymbiotic conditions (Kameoka et

al., 2019; Sugiura et al., 2020). Third, our investigation into the mechanism by which

myristate disrupts C-P exchange in AM symbiosis remains preliminary, and the limited

P supply along with varied growth conditions between alfalfa and rice further

complicate the mechanistic analysis. Therefore, further research is needed to verify and

quantitatively assess the absorption of external myristate and other C sources by

symbiotic AMF under natural conditions, as well as to elucidate their regulatory

mechanisms and ecological significance in AM symbiosis.

CONCLUSIONS

In summary (Fig. 8), our study shows that symbiotic AMF can uptake external

non-symbiotic C sources such as myristate, which is commonly present in soil

components. Exogenous myristate enhanced both intraradical and extraradical biomass

of symbiotic AMF, possibly linked to the suppression of AM-activated defense

responses in host roots. Unexpectedly, exogenous myristate disrupted the C-P exchange

in AM symbiosis, resulting in reduced mycorrhizal P benefits for plants and decreased

allocation of symbiotic C to AMF, although these effects varied depending on soil P

conditions. These findings provide new insights into the nutritional interactions in AM

symbiosis, which are crucial for understanding and managing this ancient relationship

in a changing world.

MATERIALS AND METHODS

Biological materials

In vitro cultures of AMF species *Rhizophagus irregularis* (strain: MUCL 41833),

R. intraradices (strain: MUCL 49410), and *R. diaphanus* (strain: MUCL 49416)

associated with carrot (*Daucus carota*) hairy roots were generously supplied by the

Glomeromycota *In vitro* Collection (GINCO). Our study consisted of three

interconnected and complementary experiments.

Exp. 1: $^{13}C_1$-labeled myristate ($^{13}C_1$-Myr) tracing in two-compartment AMF-hairy

root co-culture systems

In Exp. 1, $^{13}C_1$-Myr tracing experiments were performed to determine whether

exogenous myristate can be directly absorbed by the extraradical hyphae of

Rhizophagus irregularis (two independent trials), *R. intraradices*, and *R. diaphanus*

associated with carrot hairy root cultivated in solid modified Strullu–Romand (MSR)

medium, utilizing a two-compartment Petri dish system (90 × 15 mm, see Fig. 1a) as

described by St-Arnaud et al. (1996), with minor modifications (see Supporting

Information Methods S1).

When dense AMF extraradical hyphae were observed in the hyphae compartment

(HC), 12–14 weeks later, 200 μL of 0.1 mM myristate with 98% $^{13}C_1$ labeling ($^{13}C_1$-

Myr) or unlabeled (^{12}C-Myr, control) myristate was added weekly to the HC until

harvest, two or three weeks later. At harvest, AMF extraradical hyphae from the HC and root compartment (RC) were collected separately by dissolving the medium in 10 mM sodium citrate, filtering through a 37 μm nylon mesh, and thoroughly rinsing five times with sodium citrate followed by ultrapure water. To check for possible contamination of ^{13}C-myristate from the HC to the RC, a small portion of the MSR medium from the RC was also collected randomly from three $^{13}C_1$-Myr and three ^{12}C-Myr (control) treatments, with roots and AMF hyphae removed using tweezers under a stereomicroscope, and then used for determination of ^{13}C enrichment. During the experiment, the MSR medium in the RC was not contaminated by ^{13}C-myristate from the HC, as evidenced by the identical $^{13}C/^{12}C$ ratios of the MSR medium collected in the RC in $^{13}C_1$-Myr (1.0966 ± 0.0012) and ^{12}C-Myr (1.0967 ± 0.0032) treatments.

The determination of ^{13}C enrichment in AMF extraradical hyphae and solid MSR medium was performed as previously described (Bao et al., 2019), with the details provided in the Supporting Information Methods S2.

Quantitative real-time PCR (qRT-PCR) targeting fungal marker genes and quantification of AMF colonization intensity (Exp. 1)

In Exp. 1, the expression of fungal marker genes related to fatty acid uptake (*RiFAT1* and *RiFAT2*, Brands and Dörmann, 2022) and metabolism—β-oxidation (*fatty acid degradation 1* (*RiFAD1*)), palmitvaccenic acid synthesis (*oleate desaturase-like enzyme 1* (*RiOLE1*), Brands et al., 2020), and the TCA cycle (*citrate synthase 1* (*RiCIT1*))—in extraradical hyphae of *R. irregularis* from the HC, with or without myristate addition, was assessed, using the elongation factor-1 beta gene (*RiEF-1β*) as the reference gene (Das et al., 2022). Relative gene expression levels were calculated using the comparative $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001), with four independent biological replicates and three technical replicates per biological replicate.

The collection of extraradical hyphae from the HC and qRT-PCR procedures is detailed

in the Supporting Information (Method S3, Table S1).

Carrot hairy roots in the RC from the two-compartment AMF-hairy root co-culture

system (Exp. 1) were also harvested using sterile tweezers and used to assess AMF

colonization intensity (see Supporting Information Methods S4).

Exp. 2: Effects of exogenous myristate on AMF development in single-

compartment AMF-hairy root co-culture systems

In Exp. 2, we assessed the effects of exogenous myristate at varying concentrations

on spore germination and the development of *R. irregularis* intraradical and extraradical

structures associated with carrot hairy roots in a single-compartment co-culture system.

Briefly, three 4-cm-long sterile carrot hairy root segments were cultivated in MSR

medium with or without *R. irregularis* inoculation (200 sterilized spores per Petri dish).

Potassium myristate solution (0, 0.1, 0.5, 1, 5, or 10 mM, 200 μL) was applied to the

plates weekly until harvest, with five biological replicates (plates) per treatment. The

germination rate of AMF spores, as well as extraradical hyphal branching and length,

were monitored under a stereomicroscope (M250FA, Leica, Germany) and calculated

using ImageJ at 4 days post-inoculation (Schneider et al., 2012). Spore density was

monitored every 2 weeks starting 8 weeks after AMF inoculation.

Carrot hairy roots and MSR medium were harvested at 14 weeks post-AMF

inoculation. The roots and solid culture medium were carefully separated using sterile

tweezers. The medium was weighed and dissolved in 10 mM sodium citrate, and the

AMF collection was preliminarily separated into hyphae and spores using a blender,

followed by filtration through a 250 μm and 37 μm nylon meshes, respectively, and

manual selection with tweezers under a stereomicroscope. Biomasses of the

extraradical hyphae and spores were oven-dried at 80°C for 48 hours and quantified

separately. To evaluate the development of AMF intraradical structures in the carrot

hairy roots, hyphal, vesicle, and arbuscular colonization intensities were assessed

(Supporting Information Methods S4).

Exp. 3: Environmental distribution of myristate and the effect of exogenous

myristate on AM symbiosis

In Exp. 3, a field survey and a pot experiment were conducted. The field survey

aimed to evaluate the environmental distribution of myristate in soil and plants. We

analyzed myristate concentrations in bulk soil (0-10 cm) and in plant leaves, roots, leaf

litter, and rhizosphere soil from the dominant species *O. sativa*, *Axonopus compressus*,

and *Morus alba* in a paddy field, grassland, and woodland, respectively. The procedures

regarding sample collection, pretreatment, and analysis of myristate were detailed in

Supporting Information Methods S5.

The pot experiment in Exp. 3 was conducted to assess the effect of exogenous

myristate on C-P nutritional exchanges in AM symbiosis between *R. irregularis* and

two phylogenetically distant plant species—the AM-responsive plant *Medicago sativa*

L. cv. Sanditi (alfalfa, dicot) and the AM-nonresponsive plant *Oryza sativa* L. cv.

Zhonghua 11 (rice, monocot) (Hata et al., 2010). This was achieved using ^{13}C-neutral

lipid fatty acid (NLFA)-isotope ratio mass spectrometry technology, along with

physiological and transcriptome analyses. The pot experiment employed a full factorial

design with four treatment factors: myristate addition (with or without 0.1 mM

myristate), AMF inoculation (with or without *R. irregularis*), host species (alfalfa or

rice), and soil P level (relatively low or high). Ten biological replicates were included

per treatment, totaling 160 pots. The experimental procedures for plant cultivation, P or

myristate treatment, plant harvest, and measurement of plant P concentration are

detailed in Supporting Information Methods S6.

RNA sequencing (RNA-seq) on alfalfa and rice roots and bioinformatics (Exp. 3)

In Exp. 3, alfalfa and rice roots inoculated with AMF, with or without myristate application at two P levels, were subjected to RNA sequencing (RNA-seq). Three biological replicates (roots from a single pot per replicate) were randomly selected from ten replicates per treatment, resulting in a total of 24 samples. The procedures for RNA-seq and bioinformatics followed our previous study (Wang et al., 2023), with details provided in Supporting Information Methods S7.

Quantitative real-time PCR (qRT-PCR, Exp. 3)

In Exp. 3, RNA extracted from alfalfa or rice roots was reverse-transcribed into cDNA using the QuantiTect Reverse Transcription Kit (Qiagen, Valencia, CA). qRT-PCR was performed, with three independent biological replicates (randomly selected from ten replicates) for each treatment and three technical replicates per biological replicate. The relative expression levels of plant and fungal genes involved in C delivery from plants to AMF (*MsSTR2*, *OsSTR2*, *RiFAT1*, *RiFAT2*, and *RiMST2*) (Brands and Dörmann, 2022; Das et al., 2022), the mycorrhizal P pathway (*MsPT8*, *OsPT11*, and *RiPT*) (Yang et al., 2012; Fiorilli et al., 2013), and AM-activated plant defense—specifically, reactive oxygen species balance (*MsAOX3*), stress responses (*MsUGT85A*, *MsPRXP7*, and *OsPRX39*), and typical defense marker (*OsCHT1* and *OsTGA2*) (Rehman et al., 2018; Suleman et al., 2021; Zhang et al., 2021)—were analyzed to characterize the impact of exogenous myristate on C-P nutritional exchange in AM symbiosis and the underlying mechanisms at the transcriptional level. The alfalfa *actin 2* (*MsACT2*), rice *cyclin 2* (*OsCyc2*), or *R. irregularis elongation factor-1 beta* (*RiEF-1β*) genes were used as reference genes (Yang et al., 2012; Das et al., 2022), with the procedures and primers provided in the Supporting Information (Method S3, Table S1).

¹³C labeling of plants, lipid extraction, measurement of AMF signature fatty acids,

and quantification of ¹³C enrichment and ¹³C flow (Exp. 3)

In Exp. 3, ¹³C labeling of plants, lipid extraction, measurement of AMF signature

fatty acids (NLFA 16:1ω5, Olsson et al., 2005), and quantification of ¹³C enrichment

and ¹³C flow to NLFA 16:1ω5 were performed as described in previous studies (Olsson

et al., 2005; Bao et al., 2019), with details provided in Supporting Information Methods

S8.

Statistical analysis

Mycorrhizal response was calculated based on plant biomass (mycorrhizal growth

response) and plant P concentration (mycorrhizal P response), respectively, following

Corrêa et al. (2014): Mycorrhizal response (%) = (AM − NM) / NM × 100%, where

AM denotes the observed value for the target parameter (biomass or P concentration)

in mycorrhizal plants, and NM represents the mean value of nonmycorrhizal plants.

Multiple comparisons among different treatments were conducted using one-way

analysis of variance (ANOVA) followed by Tukey's post hoc test (equal variances) or

Games-Howell's post hoc (unequal variances) test. A univariate two-way ANOVA,

followed by a post hoc Tukey test, was employed to assess the effects of myristate, P

level, and their interactions on AMF colonization intensity and plant P concentration.

The significant differences between AM and NM groups were determined using

Student's t test (for normally distributed data) or Mann-Whitney U-test (for non-

normally distributed data).

Author Contributions

Hanwen Chen: Investigation, Formal analysis, Methodology, Visualization. **Tian**

Xiong: Investigation, Formal analysis, Data curation, Methodology. **Baoxing Guan:**

Writing – original draft, Investigation, Formal analysis, Data curation, Visualization.

Jiaqi Huang: Investigation, Methodology, Resources. **Danrui Zhao:** Investigation,

Resources. **Yao Chen:** Investigation, Resources. **Haoran Liang:** Investigation,

Visualization. **Yingwei Li:** Investigation, Resources. **Jingwen Wu:** Investigation,

Resources. **Shaoping Ye:** Resources. **Ting Li:** Resources. **Wensheng Shu:** Supervision.

Jin-tian Li: Supervision. **Yutao Wang:** Conceptualization, Data curation, Writing –

original draft, Writing – review & editing, Funding acquisition, Supervision, Project

administration.

Acknowledgments

We thank the financial support from the National Key R&D Program of China

(Key special Project for Marine Environmental Security and Sustainable Development

of Coral Reefs 2022-702), the National Natural Science Foundation of China (nos.

32272203 and 31772397), the Fujian Province Science and Technology Plan Project

(no. 2024I1011), the Key-Area Research and Development Program of Guangdong

Province (2022B0202110001), the Agricultural and Social Development Project of

Guangzhou Municipal Science and Technology Bureau (no. 202206010058), and the

Guangdong Provincial Key Laboratory of Plant Resources (no. 2023PlantKFP03).

Data Availability

The RNA-seq data generated in this study have been deposited in the Sequence

Read Archive at the National Center for Biotechnology Information under BioProject

ID PRJNA1083641. The raw data are provided as a Source Data File.

Competing Interest

The authors declare that they have no competing interests.

References

Argüello A, O'Brien MJ, van der Heijden MGA, Wiemken A, Schmid B, Niklaus PA. 2016. Options of partners improve carbon for phosphorus trade in the arbuscular mycorrhizal mutualism. *Ecology Letters* **19**: 648–656.

Bago B, Azcón-Aguilar C, Goulet A, Piché Y. 1998. Branched absorbing structures (BAS): A feature of the extraradical mycelium of symbiotic arbuscular mycorrhizal fungi. *New Phytologist* **139**: 375–388.

Bao X, Wang Y, Olsson PA. 2019. Arbuscular mycorrhiza under water—Carbon–phosphorus exchange between rice and arbuscular mycorrhizal fungi under different flooding regimes. *Soil Biology and Biochemistry* **129**: 169–177.

Bitterlich M, Krügel U, Boldt-Burisch K, Franken P, Kühn C. 2014. The sucrose transporter SlSUT2 from tomato interacts with brassinosteroid functioning and affects arbuscular mycorrhiza formation. *The Plant Journal* **78**: 877–889.

Blilou I, Ocampo JA, García-Garrido JM. 2000. Induction of *Ltp* (lipid transfer protein) and *Pal* (phenylalanine ammonia-lyase) gene expression in rice roots colonized by the arbuscular mycorrhizal fungus *Glomus mosseae*. *Journal of Experimental Botany* **51**: 1969–1977.

Brands M, Cahoon EB, Dörmann P. 2020. Palmitvaccenic Acid (Δ11-cis-hexadecenoic acid) is synthesized by an OLE1-like desaturase in the arbuscular mycorrhiza fungus *Rhizophagus irregularis*. *Biochemistry* **59**: 1163–1172.

Brands M, Dörmann P. 2022. Two AMP-binding domain proteins from *Rhizophagus*

664   *irregularis* involved in import of exogenous fatty acids. *Molecular Plant-Microbe*

665   *Interactions* **35**: 464–476.

666  Bravo A, Brands M, Wewer V, Dörmann P, Harrison MJ. 2017. Arbuscular mycorrhiza-

667   specific enzymes FatM and RAM2 fine-tune lipid biosynthesis to promote

668   development of arbuscular mycorrhiza. *New Phytologist* **214**: 1631–1645.

669  Brundrett MC, Tedersoo L. 2018. Evolutionary history of mycorrhizal symbioses and

670   global host plant diversity. *New Phytologist* **220**: 1108–1115.

671  Bücking H, Abubaker J, Govindarajulu M, Tala M, Pfeffer PE, Nagahashi G, Lammers

672   P, Shachar-Hill Y. 2008. Root exudates stimulate the uptake and metabolism of

673   organic carbon in germinating spores of *Glomus intraradices*. *New Phytologist*

674   **180**: 684–695.

675  Bunn RA, Simpson DT, Bullington LS, Lekberg Y, Janos DP. 2019. Revisiting the

676   "direct mineral cycling" hypothesis: arbuscular mycorrhizal fungi colonize leaf

677   litter, but why? *The ISME Journal* **13**: 1891–1898.

678  Corrêa A, Cruz C, Pérez-Tienda J, Ferrol N. 2014. Shedding light onto nutrient

679   responses of arbuscular mycorrhizal plants: nutrient interactions may lead to

680   unpredicted outcomes of the symbiosis. *Plant Science* **221–222**: 29–41.

681  Dai H, Zhang X, Zhao B, Shi J, Zhang C, Wang G, Yu N, Wang E. 2022. Colonization

682   of mutualistic mycorrhizal and parasitic blast fungi requires OsRAM2-regulated

683   fatty acid biosynthesis in rice. *Molecular Plant-Microbe Interactions* **35**: 178–186.

684  Das D, Paries M, Hobecker K, Gigl M, Dawid C, Lam HM, Zhang J, Chen M, Gutjahr

685   C. 2022. PHOSPHATE STARVATION RESPONSE transcription factors enable

686   arbuscular mycorrhiza symbiosis. *Nature Communications* **13**: 1–12.

687  Feng F, Sun J, Radhakrishnan GV, Lee T, Bozsóki Z, Fort S, Gavrin A, Gysel K,

688   Thygesen MB, Andersen KR, Radutoiu S, Stougaard J, Oldroyd GED. 2019. A

689 combination of chitooligosaccharide and lipochitooligosaccharide recognition

690 promotes arbuscular mycorrhizal associations in *Medicago truncatula*. *Nature*

691 *Communications* **10**: 5047.

692 Fiorilli V, Lanfranco L, Bonfante P. 2013. The expression of *GintPT*, the phosphate

693 transporter of *Rhizophagus irregularis*, depends on the symbiotic status and

694 phosphate availability. *Planta* **237**: 1267–1277.

695 Foster KR, Kokko H. 2006. Cheating can stabilize cooperation in mutualisms.

696 *Proceedings of the Royal Society B: Biological Sciences* **273**: 2233–2239.

697 Gutjahr C, Radovanovic D, Geoffroy J, Zhang Q, Siegler H, Chiapello M, Casieri L,

698 An K, An G, Guiderdoni E, Kumar CS, Sundaresan V, Harrison MJ, Paszkowski

699 U. 2012. The half-size ABC transporters STR1 and STR2 are indispensable for

700 mycorrhizal arbuscule formation in rice. *The Plant Journal* **69**: 906–920.

701 Hata S, Kobae Y, Banba M. 2010. Interactions between plants and arbuscular

702 mycorrhizal fungi. *International Review of Cell and Molecular Biology* **281**: 1–48.

703 Helber N, Wippel K, Sauer N, Schaarschmidt S, Hause B, Requena N. 2011. A versatile

704 monosaccharide transporter that operates in the arbuscular mycorrhizal fungus

705 *Glomus sp* is crucial for the symbiotic relationship with plants. *The Plant Cell* **23**:

706 3812–3823.

707 Janos DP. 2007. Plant responsiveness to mycorrhizas differs from dependence upon

708 mycorrhizas. *Mycorrhiza* **17**: 75–91.

709 Jiang Y, Wang W, Xie Q, Liu N, Liu L, Wang D, Zhang X, Yang C, Chen X, Tang D,

710 Wang E. 2017. Plants transfer lipids to sustain colonization by mutualistic

711 mycorrhizal and parasitic fungi. *Science* **356**: 1172–1175.

712 Johnson NC. 2010. Resource stoichiometry elucidates the structure and function of

713 arbuscular mycorrhizas across scales. *New Phytologist* **185**: 631–647.

Kameoka H, Tsutsui I, Saito K, Kikuchi Y, Handa Y, Ezawa T, Hayashi H, Kawaguchi

M, Akiyama K. 2019. Stimulation of asymbiotic sporulation in arbuscular

mycorrhizal fungi by fatty acids. *Nature Microbiology* **4**: 1654–1660.

Keymer A, Pimprikar P, Wewer V, Huber C, Brands M, Bucerius SL, Delaux PM, Klingl

V, Röpenack-Lahaye EV, Wang TL, Eisenreich W, Dörmann P, Parniske M,

Gutjahr C. 2017. Lipid transfer from plants to arbuscular mycorrhiza fungi. *eLife*

6: e29107.

Kiers ET, Duhamel M, Beesetty Y, Mensah J, Franken O, Verbruggen E, Fellbaum CR,

Kowalchuk GA, Hart MM, Bago A, Palmer TM, West SA, Vandenkoornhuyse P,

Jansa J, Bücking H. 2011. Reciprocal rewards stabilize cooperation in the

mycorrhizal symbiosis. *Science* **333**: 880–882.

Kusajima M, Fujita M, Soudthedlath K, Nakamura H, Yoneyama K, Nomura T,

Akiyama K, Maruyama-Nakashita A, Asami T, Nakashita H. 2022. Strigolactones

modulate salicylic acid-mediated disease resistance in *Arabidopsis thaliana*.

International Journal of Molecular Sciences **23**: 5246.

Lanfranco L, Fiorilli V, Gutjahr C. 2018. Partner communication and role of nutrients

in the arbuscular mycorrhizal symbiosis. *New Phytologist* **220**: 1–10.

Li S, Xu C, Wang J, Guo B, Yang L, Chen J, Ding W. 2017. Cinnamic, myristic and

fumaric acids in tobacco root exudates induce the infection of plants by *Ralstonia*

solanacearum. *Plant and Soil* **412**: 381–395.

Li Y, Chen H, Gu L, Wu J, Zheng X, Fan Z, Pan D, Li JT, Shu W, Rosendahl S, Wang

Y. 2024. Domestication of rice may have changed its arbuscular mycorrhizal

properties by modifying phosphorus nutrition-related traits and decreasing

symbiotic compatibility. *New Phytologist* **243**: 1554–1570.

Liu X, Feng Z, Zhang W, Yao Q, Zhu H. 2023. Exogenous myristate promotes the

colonization of arbuscular mycorrhizal fungi in tomato. *Frontiers in Plant Science*

14: 1250684.

Liu Y, Zhang H, Wang J, Gao W, Sun X, Xiong Q, Shu X, Miao Y, Shen Q, Xun W,

Zhang R. 2024. Nonpathogenic *Pseudomonas syringae* derivatives and its

metabolites trigger the plant "cry for help" response to assemble disease

suppressing and growth promoting rhizomicrobiome. *Nature Communications* **15**:

1907.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-

time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **25**: 402–408.

Luginbuehl LH, Menard GN, Kurup S, Van Erp H, Radhakrishnan GV, Breakspear A,

Oldroyd GED, Eastmond PJ. 2017. Fatty acids in arbuscular mycorrhizal fungi are

synthesized by the host plant. *Science* **356**: 1175–1178.

MacLean AM, Bravo A, Harrison MJ. 2017. Plant signaling and metabolic pathways

enabling arbuscular mycorrhizal symbiosis. *The Plant Cell* **29**: 2319–2335.

Mori N, Nishiuma K, Sugiyama T, Hayashi H, Akiyama K. 2016. Carlactone-type

strigolactones and their synthetic analogues as inducers of hyphal branching in

arbuscular mycorrhizal fungi. *Phytochemistry* **130**: 90–98.

Olsson PA, van Aarle IM, Gavito ME, Bengtson P, Bengtsson G. 2005. ^{13}C

incorporation into signature fatty acids as an assay for carbon allocation in

arbuscular mycorrhiza. *Applied and Environmental Microbiology* **71**: 2592–2599.

Rehman HM, Nawaz MA, Shah ZH, Ludwig-Müller J, Chung G, Ahmad MQ, Yang

SH, Lee SI. 2018. Comparative genomic and transcriptomic analyses of Family-1

UDP glycosyltransferase in three *Brassica* species and *Arabidopsis* indicates

stress-responsive regulation. *Scientific Reports* **8**: 1875.

Rich MK, Vigneron N, Libourel C, Keller J, Xue L, Hajheidari M, Radhakrishnan GV,

Le Ru A, Diop SI, Potente G, Conti E, Duijsings D, Batut A, Le Faouder P, Kodama K, Kyozuka J, Sallet E, Bécard G, Rodriguez-Franco M, Ott T, Bertrand-Michel J, Oldroyd GED, Szövényi P, Bucher M, Delaux PM. 2021. Lipid exchanges drove the evolution of mutualism during plant terrestrialization. *Science* **372**: 864–868.

Rillig MC, Aguilar-Trigueros CA, Anderson IC, Antonovics J, Ballhausen MB, Bergmann JBM, Chaudhary VB, Deveautour C, Grünfeld L, Hempel S, Lakovic M, Lammel DR, Lehmann A, Lehmann J, Leifheit EF, Liang Y, Li E, Lozano YM, Manntschke A, Mansour I, Oviatt P, Pinek L, Powell JR, Roy J, Ryo M, Sosa-Hernández MA, Veresoglou SD, Wang D, Yang G, Zhang H. 2020. Myristate and the ecology of AM fungi: significance, opportunities, applications and challenges. *New Phytologist* **227**: 1610–1614.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* **9**: 671–675.

Shi J, Zhao B, Zheng S, Zhang X, Wang X, Dong W, Xie Q, Wang G, Xiao Y, Chen F, Yu N, Wang E. 2021. A phosphate starvation response-centered network regulates mycorrhizal symbiosis. *Cell* **184**: 5527–5540.

Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*. Academic press, London.

St-Arnaud M, Hamel C, Vimard B, Caron M, Fortin JA. 1996. Enhanced hyphal growth and spore production of the arbuscular mycorrhizal fungus *Glomus intraradices* in an in vitro system in the absence of host roots. *Mycological Research* **100**: 328–332.

Sudheeran PK, Ovadia R, Galsarker O, Maoz I, Sela N, Maurer D, Feygenberg O, Oren Shamir M, Alkan N. 2020. Glycosylated flavonoids: fruit's concealed antifungal arsenal. *New Phytologist* **225**: 1788–1798.

Sugiura Y, Akiyama R, Tanaka S, Yano K, Kameoka H, Marui S, Saito M, Kawaguchi

M, Akiyama K, Saito K. 2020. Myristate can be used as a carbon and energy source

for the asymbiotic growth of arbuscular mycorrhizal fungi. *Proceedings of the*

National Academy of Sciences of the United States of America **117**: 25779–25788.

Suleman M, Ma M, Ge G, Hua D, Li H. 2021. The role of alternative oxidase in plant

hypersensitive response. *Plant Biology* **23**: 415–419.

Swenson TL, Karaoz U, Swenson JM, Bowen BP, Northen TR. 2018. Linking soil

biology and chemistry in biological soil crust using isolate exometabolomics.

Nature Communications **9**: 19.

Tanaka S, Hashimoto K, Kobayashi Y, Yano K, Maeda T, Kameoka H, Ezawa T, Saito

K, Akiyama K, Kawaguchi M. 2022. Asymbiotic mass production of the

arbuscular mycorrhizal fungus *Rhizophagus clarus*. *Communications Biology* **5**:

43.

Wang XX, Li HB, Chu Q, Feng G, Kuyper TW, Rengel Z. 2020. Mycorrhizal impacts

on root trait plasticity of six maize varieties along a phosphorus supply gradient.

Plant and Soil **448**: 71–86.

Wang YT, Chen HW, Shao MS, Zhu T, Li SS, Olsson PA, Hammer EC. 2023.

Arbuscular mycorrhizal fungi trigger danger-associated peptide signaling and

inhibit carbon–phosphorus exchange with nonhost plants. *Plant Cell &*

Environment **46**: 2206–2221.

Went FW, Stark N. 1968. The biological and mechanical role of soil fungi. *Proceedings*

of the National Academy of Sciences of the United States of America **60**: 497–504.

Wipf D, Krajinski F, van Tuinen D, Recorbet G, Courty PE. 2019. Trading on the

arbuscular mycorrhiza market: from arbuscules to common mycorrhizal networks.

New Phytologist **223**: 1127–1142.

Xie X, Lai W, Che X, Wang S, Ren Y, Hu W, Chen H, Tang M. 2022. A SPX domain-
containing phosphate transporter from *Rhizophagus irregularis* handles phosphate
homeostasis at symbiotic interface of arbuscular mycorrhizas. *New Phytologist*
234: 650–671.

Yang S, Grønlund M, Jakobsen I, Grotemeyer MS, Rentsch D, Miyao A, Hirochika H,
Kumar CS, Sundaresan V, Salamin N, Catausan S, Mattes N, Heuer S, Paszkowski
U. 2012. Nonredundant regulation of rice arbuscular mycorrhizal symbiosis by
two members of the phosphate transporter1 gene family. *The Plant Cell* **24**: 4236–
4251.

Yu H, He Y, Zhang W, Chen L, Zhang J, Zhang X, Dawson W, Ding J. 2022. Greater
chemical signaling in root exudates enhances soil mutualistic associations in
invasive plants compared to natives. *New Phytologist* **236**: 1140–1153.

Zhang C, He J, Dai H, Wang G, Zhang X, Wang C, Shi J, Chen X, Wang D, Wang E.
2021. Discriminating symbiosis and immunity signals by receptor competition in
rice. *Proceedings of the National Academy of Sciences of the United States of*
America **118**: e2023738118.

Zhang L, Feng G, Declerk S. 2018. Signal beyond nutrient, fructose, exuded by an
arbuscular mycorrhizal fungus triggers phytate mineralization by a phosphate
solubilizing bacterium. *The ISME Journal* **12**: 2339–2351.

Zhang Q, Blaylock LA, Harrison MJ. 2010. Two *Medicago truncatula* half-ABC
transporters are essential for arbuscule development in arbuscular mycorrhizal
symbiosis. *The Plant Cell* **22**: 1483–1497.

Fig. 1 Uptake of exogenous myristate by extraradical hyphae (ERH) of arbuscular

mycorrhizal fungi (AMF) under symbiotic conditions in Exp. 1. **(a)** Schematic

illustration of the two-compartment AMF–carrot hairy root co-cultivation system and

the addition of myristate to the hyphae compartment (HC). Carrot hairy roots colonized

by *Rhizophagus irregularis*, *R. intraradices*, or *R. diaphanus* were grown in the root

compartment (RC). Twelve to fourteen weeks later, 0.1 mM of $^{13}C_1$-labeled ($^{13}C_1$-Myr),

non-labeled (^{12}C-Myr) myristate, or ddH$_2$O (Myr$^-$, control) was added weekly to the

HC with well-developed AMF extraradical hyphae (ERH) until harvest. **(b)** Branched

absorbing structures (BAS) of *R. irregularis*, *R. intraradices*, and *R. diaphanus* were

observed in the HC only when treated with exogenous myristate (Myr$^+$). **(c)** $^{13}C/^{12}C$

ratio of the harvested ERH from HC or RC of *R. irregularis*, *R. intraradices*, and *R.*

diaphanus when $^{13}C_1$-labeled ($^{13}C_1$-Myr) or non-labeled (^{12}C-Myr, background)

myristate was added to HC. Two independent experimental trials were performed for

R. irregularis. **(d, e)** Calculated ^{13}C enrichment in ERH of *R. irregularis*, *R.*

intraradices, and *R. diaphanus* in the HC and RC under ^{13}C-Myr treatment. Black

horizontal lines represent mean value from 2–5 biological replicates. *, **, and ***

indicate statistical significance between $^{13}C_1$-labeled and non-labeled treatments at the

0.05, 0.01, and 0.001 probability levels (Mann-Whitney U-test), respectively. Different

letters indicate significant difference at the 0.05 probability level (one-way ANOVA

followed by Games-Howell's post hoc test). Source data are provided as a Source Data

file.

Fig. 2 Effects of exogenous myristate on the expression of fungal fatty acid uptake and metabolism genes in the extraradical hyphae (ERH) and on root colonization by arbuscular mycorrhizal fungi (AMF) in Exp. 1. **(a, b)** Normalized expression (relative to *RiEF-1β*) of AMF genes involved in **(a)** fatty acid uptake (*RiFAT1* and *RiFAT2*) and **(b)** metabolism—β-oxidation (*RiFAD1*), palmitvaccenic acid synthesis (*RiOLE1*), and TCA cycle (*RiCIT1*)—in the ERH of *R. irregularis* collected from the hyphae compartment (HC) under myristate (Myr⁺) or ddH₂O (Myr⁻) treatments (n = 4). **(c–e)** Percent root colonization by **(c)** *R. irregularis* (n = 3–10), **(d)** *R. intraradices* (n = 9–22), and **(e)** *R. diaphanus* (n = 6–14) in carrot hairy roots from the root compartment (RC) when HC was treated with myristate (Myr⁺) or ddH₂O (Myr⁻). Two independent experimental trials were performed for *R. irregularis*. Means ± SE, *, **, and *** indicate statistical significance between Myr⁺ and Myr⁻ treatments at the 0.05, 0.01, and 0.001 probability levels, respectively. Source data are provided as a Source Data file.

Fig. 3 Effects of exogenous myristate on the development of intraradical and extraradical structures of AMF (*Rhizophagus irregularis*) in the single-compartment AMF-carrot hairy root co-cultivation system (Exp. 2). **(a)** Schematic illustration of the single-compartment *R. irregularis*-hairy root co-cultivation system and the addition (Myr⁺) or absence (Myr⁻) of myristate. **(b)** Spore germination rate, **(c)** extraradical hyphae (ERH) branching, and **(d)** ERH length in the *R. irregularis*-hairy root co-

cultivation system with (Myr$^+$) or without (Myr$^-$) myristate treatment at 4 days post-

inoculation (n = 11−12). **(e)** AMF percent root colonization in carrot hairy roots and **(f)**

biomass of the AMF extraradical spore and hypha (per gram of medium) at 14 weeks

post-inoculation (n = 3−6). **(g)** Dynamics of AMF spore density over time (per square

centimeter, n = 3−5); and **(h)** AMF spore diameter at harvest with 0, 0.1, 0.5, 1, 5, or

10 mM myristate treatment under symbiotic conditions (n = 5, calculation from > 900

spores in each replicate). Means ± SE; different letters indicate significant differences

at P < 0.05 level (One-way ANOVA followed by Tukey's post hoc test); *** indicates

statistical significance at the 0.001 probability level (Student's t-test). Source data are

provided as a Source Data file.

Fig. 4 Environmental myristate levels and the effects of exogenous myristate on arbuscule abundance, as well as the intraradical and extraradical biomass of arbuscular mycorrhizal fungi (AMF, *Rhizophagus irregularis*) associated with *Medicago sativa* or *Oryza sativa* under low (LP) or high (HP) phosphorus conditions (Exp. 3). **(a)** Schematic illustration of the field survey of environmental myristate levels and

experimental design in the pot experiment (Exp. 3). **(b)** Myristate concentration in bulk

soil, as well as in plant leaves, roots, leaf litter, and rhizosphere soil from the dominant

species *O. sativa*, *Axonopus compressus*, and *Morus alba* in the paddy field, grassland,

and woodland, respectively (n = 4). **(c, d)** Arbuscule abundance in *M. sativa* and *O.*

sativa roots treated with 0.1 mM myristate (Myr$^+$) or ddH$_2$O (Myr$^-$) (n = 10; vesicle

and hyphal abundance are shown in Supporting Information Fig. S1); **(e, f)** AMF

biomass, indicated by the AMF signature fatty acid (NLFA 16:1ω5), in AMF-colonized

roots and rhizosphere soil of *M. sativa* and *O. sativa* treated with myristate (Myr$^+$) or

ddH$_2$O (Myr$^-$) (n = 5). Means ± SE; different letters indicate significant differences at

$P < 0.05$ level (**b**: Student's *t*-test, **c−f**: one-way ANOVA followed by Tukey's post hoc

test). Source data are provided as a Source Data file.

Fig. 5 Effects of exogenous myristate on symbiotic carbon (C) allocation from *Medicago sativa* and *Oryza sativa* to arbuscular mycorrhizal fungi (AMF, *Rhizophagus irregularis*) and on the expression profiles of mycorrhizal C transportation marker genes. **(a, b)** Plant-derived ^{13}C flow to the AMF signature fatty acid (NLFA 16:1ω5) in AMF-colonized roots of *M. sativa* and *O. sativa* treated with 0.1 mM myristate (Myr$^+$) or ddH$_2$O (Myr$^-$) under low (LP) and high (HP) phosphorus conditions (n = 5). **(c−h)**

Normalized expression of plant and fungal (underlined) marker genes for mycorrhizal

C transport in AMF-colonized (AM) and non-colonized (NM) roots treated with 0.1

mM myristate (Myr$^+$) or ddH$_2$O (Myr$^-$) (n = 3). The mycorrhizal C transportation genes

include the plant symbiotic fatty acid transporter *STR2* **(c, d)**, the fungal fatty acid

transporters (*RiFAT1, RiFAT2*; **e, f**), and the monosaccharide transporter (*RiMST2*; **g,**

h). Normalized expression of *M. sativa*, *O. sativa*, and AMF genes was calculated

relative to *MsACT2*, *OsCyc2*, and *RiEF-1β*, respectively; n.d., not detected. Two-way

ANOVA in each histogram applies only to AM roots. Means ± SE; * and ** indicate

statistical significance at the 0.05 and 0.01 probability levels, respectively. Source data

are provided as a Source Data file.

Fig. 6 Effects of exogenous myristate on plant phosphorus (P) concentration, mycorrhizal P response, and expression of plant and fungal (underlined) marker genes for the mycorrhizal P pathway in *Medicago sativa* and *Oryza sativa*. **(a, b)** Plant P concentration (n = 8−10); **(c, d)** mycorrhizal P response (n = 8−10); and **(e, f)** relative expression of *MsPT8*, *OsPT11*, and *RiPT* in *M. sativa* and *O. sativa* roots under low (LP) and high (HP) phosphorus conditions with 0.1 mM myristate (Myr$^+$) or ddH$_2$O (Myr$^−$) treatments (n = 3). AM, inoculated with *Rhizophagus irregularis*; NM, non-inoculated; n.d., not detected. Normalized expression of *M. sativa*, *O. sativa*, and AMF genes was calculated relative to *MsACT2*, *OsCyc2*, and *RiEF-1β*, respectively. Two-

934   way ANOVA in each histogram (c−f) applies only to AM roots. Means ± SE; different

935   letters indicate significant differences at $P < 0.05$ level (one-way ANOVA followed by

936   Tukey's post hoc test). * and *** indicate statistical significance at the 0.05 and 0.001

937   probability levels (Student's *t*-test), respectively. Source data are provided as a Source

938   Data file.

Fig. 7 Effects of exogenous myristate on the global transcriptome and putative AM-activated defense genes in *Medicago sativa* and *Oryza sativa* roots inoculated with arbuscular mycorrhizal fungi (AMF, *Rhizophagus irregularis*) under low (LP) and high (HP) phosphorus conditions. **(a, b)** Significantly enriched gene ontology (GO) terms (P_{adj} < 0.05) for differentially expressed genes (DEGs) between myristate (Myr$^+$) and ddH$_2$O (Myr$^-$) treatments in RNA-seq. Underlined terms denote stress- or defense-related GO categories, with the expression profile of their component genes provided in Supporting Information (Fig. S4). **(c, d)** Quantitative real-time PCR analysis of the AM-activated defense gene expression—reactive oxygen species (ROS) balance (*MsAOX3*), stress responses (*MsUGT85A*, *MsPRXP7*, and *OsPRX39*), and defense responses *(OsCHT1* and *OsTGA2)* — between Myr$^+$ and Myr$^-$ treatments under LP and HP conditions (n = 3). Means ± SE; * indicates statistical significance between Myr$^+$

and Myr$^-$ treatments at the 0.05 probability level (Student's t-test). Source data are

provided as a Source Data file.

Fig. 8 A proposed model for direct uptake of exogenous myristate by symbiotic

arbuscular mycorrhizal fungi (AMF) and its regulatory effects on the arbuscular

mycorrhizal symbiosis. Symbiotic AMF can uptake exogenous myristate, possibly

through the fungal fatty acid transporters RiFAT2 in the extraradical hyphae (ERH), as

indicated by [13]C enrichment and transcriptional activation of fatty acid transport and

metabolism genes in AMF extraradical hyphae. Exogenous myristate increased both

intraradical and extraradical fungal biomass, possibly linked to the suppression of AM-

activated defense responses in host roots. Unexpectedly, as a non-symbiotic carbon

source, exogenous myristate disrupted the carbon (C)-phosphorus (P) exchange in AM

symbiosis, resulting in reduced mycorrhizal P benefits for plants and decreased

allocation of symbiotic C to AMF. Question mark or dashed arrow indicates putative or

unclear mechanisms. SA, salicylic acid; ROS, reactive oxygen species; PR,

pathogenesis-related; NLFA 16:1ω5, neutral lipid fatty acid 16:1ω5 (AMF marker).