Tap into non-symbiotic carbon? Exogenous myristate fuels the growth of symbiotic arbuscular mycorrhizal fungi but disrupts their carbon-phosphorus exchange with host plants

Hanwen Chen[1, #], Tian Xiong[1, #], Baoxing Guan[1, #], Jiaqi Huang[1], Danrui Zhao[1], Yao Chen[1], Haoran Liang[1], Yingwei Li[1], Jingwen Wu[1], Shaoping Ye[2], Ting Li[2], Wensheng Shu[1], Jin-tian Li[1], Yutao Wang[1, *]

[1]Guangdong Provincial Key Laboratory of Biotechnology for Plant Development, Guangzhou Key Laboratory of Subtropical Biodiversity and Biomonitoring, School of Life Sciences, South China Normal University, Guangzhou, 510631, China

[2]Guangzhou Institute of Forestry and Landscape Architecture, Guangzhou, 510405, China

*Correspondence: School of Life Sciences, South China Normal University, Guangzhou, 510631, China; Email: wangyutao@scnu.edu.cn (Yutao Wang); Tel./Fax: +86 20 85211850

[#]: These authors contributed equally to this work.

Abstract

Arbuscular mycorrhizal fungi (AMF) form symbiotic relationships with the roots of most land plants. In this association, known as arbuscular mycorrhiza (AM), the obligate biotrophic fungi acquire host-derived (symbiotic) carbohydrates and, in return, facilitate plant uptake of mineral nutrients such as phosphate. However, the capacity of symbiotic AMF to access external non-symbiotic carbon sources remains unclear, complicating our understanding of the nutritional exchanges in AM and their ecosystem functions. In this study, we explored the direct uptake of exogenous $^{13}C_1$-labeled myristate by symbiotic AMF (*Rhizophagus irregularis*) and their growth responses using a two-compartment and a single-compartment AMF-carrot hairy root co-culture system, respectively. Additionally, we assessed the impact of exogenous myristate on the carbon-phosphorus exchange between AMF and two host species in a greenhouse experiment, using ^{13}C-neutral lipid fatty acid-isotope ratio mass spectrometry technology, alongside physiological and transcriptome analyses. Our findings reveal that symbiotic AMF can absorb external nonsymbiotic myristate, at least under the conditions tested, as evidenced by ^{13}C enrichment and transcriptional activation of putative fatty acid transport and metabolism genes in extraradical hyphae. Exogenous myristate increased both intraradical and extraradical fungal biomass, potentially due to suppressed mycorrhizal defense responses in host roots. Intriguingly, exogenous myristate reduced the mycorrhizal phosphorus benefits for both *Medicago sativa* and *Oryza sativa* and decreased their carbon allocation to symbiotic AMF, although these effects were influenced by host identity and soil phosphorus conditions. These findings advance our understanding of the nutritional interactions in AM, which are essential for comprehending and manipulating this symbiotic relationship.

45   **Keywords** : arbuscular mycorrhizal fungi (AMF), carrot hairy root, carbon-

46   phosphorus exchange, *Medicago sativa* (alfalfa), non-symbiotic carbon (myristate),

47   *Oryza sativa* (rice)

Introduction

Arbuscular mycorrhizal fungi (AMF) are among the most widely distributed soil microorganisms, forming symbiotic relationships with approximately 70% of terrestrial vascular plants (Brundrett and Tedersoo, 2018). In this association, known as arbuscular mycorrhiza (AM), AMF develop an extensive extraradical hyphal network in the soil, facilitating nutrient uptake for host plants, particularly phosphorus (P), and enhancing plant resilience to both biotic and abiotic stressors (Smith and Read, 2008). In return, host plants allocate 4–20% of their photosynthetically fixed carbon (C) to their fungal partners in the form of sugars and fatty acids (Helber et al., 2011; Jiang et al., 2017). This symbiotic C-P exchange, which serves as the cornerstone of the AM symbiosis (Smith and Read, 2008), not only plays a pivotal role in the biogeochemical cycling of C and P elements but also acts as a key mechanism through which AMF influence plant community structure and productivity (Johnson, 2010; Wipf et al., 2019).

As one of the most ancient symbiotic relationships, AM has undergone extensive evolutionary processes (Foster and Kokko, 2006; Rich et al., 2021), with its establishment and development intricately regulated by a set of symbiotic genes and signaling molecules influenced by biotic and abiotic conditions (Maclean et al., 2017; Shi et al., 2021). For instance, in AM plants, the two pathways for P uptake—the direct P pathway (via root hair and epidermis) and mycorrhizal P pathway (via AMF hyphae)—are coordinately regulated by the same phosphate-sensing pathway in plants (Shi et al., 2021; Das et al., 2022). Under P deficiency conditions, host plants synthesize and release strigolactones into the soil, activating the mycorrhizal P pathway by stimulating the elongation and branching of AMF hyphae (Mori et al., 2016; Yu et al., 2022). Additionally, the extent of AMF colonization and AM benefits

largely depend on the host plant species, with some being "responsive plants" that

exhibit positive growth responses (e.g., alfalfa (*Medicago sativa* L.), maize (*Zea mays*

L.)) and others being "nonresponsive plants" that show minimal or no growth

enhancement (e.g., rice (*Oryza sativa* L.), wheat (*Triticum aestivum* L.)) (Hata et al.,

2010; Cope et al., 2022).

AMF are considered obligate symbionts that rely on living host plants to

complete their life cycle by producing daughter spores (Smith and Read, 2008), which

presents challenges for basic research on AM and their applications in agriculture and

other fields. Thus far, achieving a pure (plant-independent) culture of asymbiotic

AMF has not been successful, partially due to their dependence on plant-derived C

(Bravo et al., 2017; Luginbuehl et al., 2017; Kameoka et al., 2019). Genomic and

transcriptomic analyses reveal that AMF lack the genes required for the biosynthesis

of long-chain fatty acids, and subsequent studies have demonstrated that fatty acids

are transferred from host plants to AMF as their main C source (Bravo et al., 2017;

Jiang et al., 2017; Keymer et al., 2017; Luginbuehl et al., 2017). However, the

molecular mechanism underlying the allocation of plant-derived fatty acids to fungi in

AM symbiosis remains unclear. In plants, AMF colonization triggers the expression of

lipid biosynthesis and transport genes, such as *DIS*, *FatM*, *RAM2*, and *STR2* (Gutjahr

et al., 2012; Bravo et al., 2017; Keymer et al., 2017; Luginbuehl et al., 2017), with

lipid transport to the AM symbiotic interface relying on the periarbuscular membrane-

located transporters STR and STR2 (Zhang et al., 2010; Gutjahr et al., 2012) and

fungal uptake of plant-derived lipids mediated by RiFAT1 and RiFAT2 in the model

AMF species *Rhizophagus irregularis* (Brands and Dörmann, 2022).

Fatty acids and their derivatives, essential structural components and energy

sources for plant, animal, and microbial cells, are commonly found in soil,

particularly in the rhizosphere, where plant-derived fatty acids are made available

through root exudates and rhizodeposition (Swenson et al., 2018; Dai et al., 2022; Liu

et al., 2024). Following the discovery that AMF acquire fatty acids from their hosts

during symbiosis, the use of fatty acids in cultivating asymbiotic AMF has received

considerable attention. Kameoka et al. (2019) demonstrated that under asymbiotic

conditions, exogenous palmitoleic acid (C16:1 fatty acid) and several other fatty acids

stimulated the elongation and branching of *R. irregularis* hyphae, resulting in the

formation of infection-competent but morphologically and functionally incomplete

spores with reduced size, lower germination rates, and decreased infection capacity

compared to symbiotic spores. Additionally, Sugiura et al. (2020) showed that

exogenous myristate (tetradecanoic acid, C14:0 fatty acid), a common organic acid in

various plant root exudates (Fernández-Piñar et al., 2012; Luo et al., 2014; Li et al.,

2017; Liu et al., 2024) and soil substrates (Swenson et al., 2018; Dai et al., 2022), can

serve as a C and energy source for the asymbiotic growth of *R. irregularis,* promoting

hyphal growth, the formation of branched absorbing structures, and the production of

morphologically incomplete infection-competent spores. Subsequent studies revealed

that combining plant hormones (strigolactone and methyl jasmonate) with myristate

enhanced the efficiency of myristate as a C source in asymbiotic *R. clarus*, leading to

the generation of numerous infection-competent spores that were smaller than those

generated under symbiotic conditions (Sugiura et al., 2020; Tanaka et al., 2022).

These discoveries have significantly progressed the pure culture of AMF and,

importantly, identified myristate as the first external C source accessible to asymbiotic

AMF, providing exceptional opportunities for understanding and utilizing the

ecological functions of AMF in plant community regulation and global C cycling

(Rillig et al., 2020).

However, the effects of external myristate on the growth and development of

symbiotic AMF—the typical state of functioning AMF in nature—remain unclear.

Regarding sugars, the other type of host-derived C source, symbiotic AMF may be

unable to absorb them additionally through extraradical hyphae (Bücking et al., 2008;

Helber et al., 2011; Sugiura et al., 2020; but see Pfeffer et al., 1999). A recent study

reported that while exogenous myristate does not influence the extraradical biomass

of symbiotic AMF, it enhances AMF colonization by stimulating the growth and

development of extraradical hyphae. Based on this, the authors inferred that symbiotic

AMF may acquire exogenous myristate (Liu et al., 2023). To date, the effects of

exogenous fatty acids on AM symbiosis and their underlying mechanisms remain

largely unexplored. Specifically, the capacity of symbiotic AMF to uptake myristate

as an external non-symbiotic C source in the presence of plant-derived (symbiotic) C

is yet to be determined. This question is crucial for understanding the symbiotic

mechanisms and ecological roles of AM symbiosis (Rillig et al., 2020).

This study investigated the capacity of symbiotic AMF to absorb external non-

symbiotic myristate and examined the effect of exogenous myristate on symbiotic C-P

exchanges in AM. Specifically, we aimed to determine: (i) whether symbiotic AMF

can take up external non-symbiotic myristate, and if so, (ii) how this additional non-

symbiotic C source influences the symbiotic C allocation from plants to AMF, and (iii)

whether and how fungal P transfer to the host plant is influenced by exogenous

myristate. To address these questions, we performed a $^{13}C_1$-labeled myristate tracing

experiment to examine the direct uptake of exogenous myristate by extraradical

hyphae of AMF associated with carrot hairy roots using a two-compartment co-

culture system. Additionally, we assessed the impact of varying myristate

concentrations on spore germination and the intra- and extraradical growth of AMF

associated with carrot (*Daucus carota* var. *sativa* Hoffm.) hairy roots in a single-

compartment co-culture system. Furthermore, we conducted a greenhouse experiment,

using ^{13}C-neutral lipid fatty acid-isotope ratio mass spectrometry (^{13}C-NLFA-IRMS)

technology, and physiological and transcriptome analyses, to further investigate how

exogenous myristate influences the symbiotic interaction between AMF and two host

plants. This work provides, to our knowledge, the first solid evidence that symbiotic

AMF can absorb non-symbiotic C sources like myristate in the presence of plant-

derived carbohydrates. Moreover, our findings reveal that under symbiotic conditions,

exogenous myristate unexpectedly disrupts the C-P exchange in AM symbiosis.

Results

Tracing of ^{13}C in AMF extraradical hyphae and the effect of myristate on AMF

colonization and marker gene expression (Exp. 1)

In the two-compartment AMF-carrot hairy root co-cultivation system (Exp. 1,

Fig. 1a), exogenous myristate specifically induced the branched absorbing structures

of AMF in the HC two weeks after Myr$^+$ treatment (Fig. 1b). The ^{13}C/^{12}C ratio of the

MSR medium collected in RC in ^{13}C$_1$-Myr (1.0966 ± 0.0012) and ^{12}C-Myr (non-

labeled, 1.0967 ± 0.0032) treatments was identical, demonstrating that the exogenous

myristate added in HC did not diffuse directly into RC. The ^{13}C/^{12}C ratio of

extraradical hyphae collected in HC (1.0922 ± 0.0003%) and RC (1.0923 ± 0.0002%)

in ^{12}C-Myr treatment was identical and close to the natural ^{13}C/^{12}C ratio. In the ^{13}C$_1$-

Myr treatment, the ^{13}C/^{12}C ratio of extraradical hyphae in HC (1.1504 ± 0.0575%) and

RC (1.1026 ± 0.0055%) was significantly higher than that in the non-labeled control

($P < 0.05$, Fig. 1c), suggesting the uptake of ^{13}C$_1$-Myr by extraradical hyphae.

Exogenous myristate significantly increased the total, hyphal, and arbuscular

colonization rates in carrot hairy roots ($P < 0.05$, Fig. 1d) but had no effect on vesicle

colonization ($P > 0.05$).

qRT-PCR of the fungal marker genes in AMF extraradical hyphae collected in

HC showed that exogenous myristate increased the expression levels of the fungal

fatty acid transporter genes *RiFAT2* ($P < 0.05$) and *RiFAT1* ($P > 0.05$), as well as the

fungal genes involved in fatty acid β-oxidation (*RiFAD1*, $P < 0.05$), palmitvaccenic

acid synthesis (*RiOLE1*, $P = 0.057$), and TCA cycling (*RiCIT1*, $P < 0.05$) (Fig. 1e, f).

These results suggest that exogenous myristate activated the expression of fungal

genes related to fatty acid transportation and metabolism in the extraradical hyphae of

AMF.

Effects of exogenous myristate on AMF spore germination, root colonization and

extraradical structure development (Exp. 2)

In the single-compartment AMF-carrot hairy root co-cultivation system (Exp. 2,

Fig. 2a), exogenous myristate significantly promoted AMF spore germination ($P <$

0.001, Fig. 2b), and enhanced hyphal branching and length ($P < 0.001$) before

physical contact between AMF hyphae and hairy roots at four days post AMF

inoculation (Fig. 2c, d). Morphological examinations showed that exogenous

myristate significantly enhanced the colonization intensities of hyphae, vesicles and

arbuscules in carrot hairy roots across various application levels ($P < 0.05$, Fig. 2e),

with its promotional effects generally increasing at higher myristate concentrations.

Exogenous myristate also significantly increased the extraradical biomass of AMF

(hyphae and spore) at various application concentrations ($P < 0.05$, Fig. 2f).

Meanwhile, exogenous myristate significantly increased the density of AMF

spores from eight weeks post-AMF inoculation ($P < 0.05$, Fig. 2g), and the diameter

of newly produced AMF spores in mediums with exogenous myristate were identical

to those in mediums without myristate (Fig. 2h).

Effects of exogenous myristate on AMF colonization and biomass during

symbiosis with alfalfa and rice (Exp. 3)

In the pot experiment (Exp. 3), all typical AMF structures (hyphae, arbuscules,

and vesicles) were observed in the roots of both alfalfa and rice in AMF-inoculated

groups (AM), while no AMF colonization was detected in the non-inoculated (NM)

groups. Under low-P conditions, exogenous myristate significantly increased

arbuscule colonization in alfalfa roots ($P < 0.05$, Fig. 3a) but did not significantly

affect hyphae or vesicle colonization ($P > 0.05$, Supplementary Fig. S1a). Similarly,

exogenous myristate significantly enhanced arbuscule and vesicle colonization in rice

roots ($P < 0.05$, Fig. 3b, Supplementary Fig. S1b). However, under high-P conditions,

exogenous myristate showed no effect on AMF colonization in either alfalfa or rice

roots (Fig. 3, Supplementary Fig. S1).

The content of the AMF signature fatty acid (NLFA 16:1ω5) in AM roots (alfalfa:

143.2 ± 13.8 nmol g^{-1}; rice: 197.3 ± 11.9 nmol g^{-1}) was 10–20 times higher than in

their corresponding NM roots (alfalfa: 6.1 ± 0.5 nmol g^{-1}; rice: 17.3 ± 1.2 nmol g^{-1}),

suggesting that NLFA 16:1ω5 serves as a distinctive fatty acid marker for AMF in this

study. The levels of NLFA 16:1ω5 in plant roots corresponded with the patterns

observed in AMF colonization. Under low-P conditions, exogenous myristate

significantly increased the NLFA 16:1ω5 levels in both alfalfa and rice roots ($P < 0.05$,

Fig. 3c, d); however, it had no significant effect under high-P conditions. Similar

trends were observed in the rhizosphere soils: under low-P conditions, exogenous

myristate increased the NLFA 16:1ω5 levels in the rhizosphere soils of alfalfa ($P <$

0.05) but not rice ($P = 0.14$) (Fig. 3c, d). Under high-P conditions, myristate had no

effect on the NLFA 16:1ω5 levels in the rhizosphere soil of either plant.

Effects of exogenous myristate on the transfer of plant-derived C to AMF (Exp. 3)

The ^{13}C-NLFA-IRMS experiment investigated the effect of exogenous myristate on C transfer from plants to AMF. In AM plants, a significant enrichment of ^{13}C (δ^{13}C = 1.75 ± 0.6‰) in AMF signature fatty acid (NLFA 16:1ω5) was observed compared to NM plants (δ^{13}C = −24.8 ± 2.3‰), demonstrating the transfer of newly fixed ^{13}C from plants to AMF. In alfalfa, exogenous myristate significantly reduced the plant-derived C transfer to AMF ($P < 0.001$, Fig. 4a). In rice, myristate also significantly influenced the C flow to AMF, with its effect significantly interacting with soil P level (Myr × P: $P = 0.004$, Fig. 4b): myristate significantly increased the allocation of ^{13}C to AMF under low-P conditions ($P < 0.05$) but decreased it under high-P conditions ($P < 0.01$).

qRT-PCR analysis revealed that the expressions of the symbiotic lipid transfer marker genes *STR2* (in alfalfa and rice) and the fungal genes responsible for symbiotic fatty acid *(RiFAT1* and *RiFAT2)* or monosaccharide (*RiMST2*) uptake were mostly exclusively detected in AM roots, except for a low expression level of *STR2* occasionally detected in NM roots of alfalfa (Fig. 4c–h).

In AMF-inoculated alfalfa roots, exogenous myristate significantly downregulated the expressions of the symbiotic C transfer genes *MsSTR2* and *RiMST2* ($P < 0.05$, Fig. 4c, g), while having no effects on *MsFAT1* and *MsFAT2* (Fig. 4e). In AMF-inoculated rice roots, the effects of myristate on the tested symbiotic C transfer genes were mostly significantly influenced by soil P levels ($P < 0.05$). Under low-P conditions, myristate significantly increased the expressions of *OsSTR2*, *OsFAT1*, and *OsFAT2*, while decreasing the expression of *RiMST2* ($P < 0.05$, Fig. 4d, f, h). Under high-P conditions, myristate significantly increased the expressions of *RiMST2* ($P < 0.05$) but had no effects on *OsSTR2*, *OsFAT1*, and *OsFAT2* ($P > 0.05$). In

249   both plant species with AMF inoculation, soil P level showed significant effects on

250   *STR*, *RiFAT1*, and *MST2* genes ($P < 0.05$), with lower expression level of these genes

251   under high-P compared to low-P conditions.

Effects of exogenous myristate on the activity of mycorrhizal P pathway and the

MGR (Exp. 3)

In the pot experiments (Exp. 3), AMF inoculation significantly increased P

concentrations in both alfalfa and rice plants ($P < 0.001$, Fig. 5a, b). Under low-P

condition, exogenous myristate significantly decreased the mycorrhizal P pathway

(MPR) in both alfalfa and rice ($P < 0.05$, Fig. 5c, d). Under high-P conditions, while

exogenous myristate significantly decreased the MPR in alfalfa ($P < 0.05$), it had no

significant effect on the MPR in rice.

The expression of the mycorrhizal P pathway marker genes in plant (*MsPT8* or

OsPT11) and AMF (*RiPT*) was exclusively detected in the AM roots of rice and

alfalfa, except for a low expression level of *OsPT11* observed in the NM root of rice

under low-P and Myr$^-$ conditions. This observation confirmed that AM symbiosis

activated the mycorrhizal P pathway at the transcriptional level in both plant species

(Fig. 5e, f). Exogenous myristate significantly affected the transcriptional responses

of the mycorrhizal P pathway in both alfalfa and rice ($P < 0.01$). In alfalfa, the

expressions of *MsPT8* and *RiPT* were downregulated by exogenous myristate under

both low- and high-P conditions ($P < 0.05$, Fig. 5e). In rice, exogenous myristate

similarly decreased the expression levels of *OsPT11* and *RiPT* under low-P conditions

($P < 0.05$), but not under high-P conditions ($P > 0.05$).

In alfalfa, AMF inoculation significantly increased plant biomass under both

low- and high-P conditions ($P < 0.05$, Supplementary Fig. S2a), while the MGRs of

alfalfa were significantly higher under high-P conditions compared to low-P

conditions ($P < 0.05$, Supplementary Fig. S2c). In rice, AMF inoculation reduced the

plant biomass under high-P (but not low-P) conditions ($P < 0.05$), and the MGRs of

rice were significantly lower under high-P conditions compared to low-P conditions

($P < 0.05$, Supplementary Fig. S2b, d). Exogenous myristate had no effect on plant

biomass or MGR in either alfalfa or rice, whereas P levels significantly influenced

MGR, exhibiting contrasting effects between alfalfa and rice ($P < 0.001$,

Supplementary Fig. S2c, d).

Effects of exogenous myristate on the transcriptome of mycorrhizal roots (Exp. 3)

RNA-seq data encompassed the majority of plant genes (transcripts), with 39,944

and 33,056 genes in alfalfa and rice, respectively. Principal component analysis (PCA)

of gene expression profiles in mycorrhizal roots of alfalfa and rice revealed that the

first and second principal components together accounted for 32.4% and 44.1% of the

variation, respectively (Supplementary Fig. S3a, b). In both alfalfa and rice, the gene

expression profiles with or without myristate application were distinct under low-P

but not high-P conditions, suggesting the importance of P status in myristate's effect.

Exogenous myristate induced a number of differentially expressed genes (DEGs,

Myr^{+} *vs.* Myr^{-}) in mycorrhizal alfalfa (LP: 308, HP: 137) and rice (LP: 590, HP: 855)

roots (Supplementary Fig. S3c, d). Gene Ontology (GO) analysis of these DEGs

revealed the enrichment in several GO terms, most of which were associated with

plant stress or defense (Fig. 6a, b). These included "defense response", "response to

salicylic acid", "defense response to bacterium", "peroxidase activity", "reactive

oxygen species metabolic process", etc., in rice, and "glycosyltransferase activity",

"alternative oxidase activity", and "heme binding" in alfalfa.

We subsequently analyzed the expression profiles of the component genes

associated with these enriched GO terms using the RNA-seq dataset. In alfalfa,

exogenous myristate significantly downregulated several key genes involved in

reactive oxygen species (ROS) balance (*MsAOX2*, *MsAOX3*) and stress tolerance

(*MsUFGT*, *MsUGT73C3*, *MsUGT73C6*) under low-P conditions, as well as typical

stress-responsive genes (*MsPRXP7*, *MsCYP83B1*, and gene families such as *MsG8HL*,

MsCYP71D10, and *MsCYP736A12*) under high-P conditions (Fig. 6c). In rice,

exogenous myristate downregulated several typical defense-associated transcriptional

factors (*OsTGA2*, *OsTGA4*, *OsWRKY45*, *OsWRKY62*, *OsWRKY76*, and *OsbZIP79*)

and marker genes (*OsPR1a*, *OsPR5L*, *OsPP2C35*, *OsACS2*, *OsDWARF4*, *OsPITA*,

OsMLO2, *OsRH1*, and *OsCHT1*) under low-P conditions (Fig. 6d). However, most of

these genes were upregulated following myristate application under high-P conditions.

Discussion

AMF are obligate biotrophic organisms that rely on host-derived (symbiotic)

carbohydrates (Smith and Read, 2008). However, whether symbiotic AMF can take

up external non-symbiotic organic C remains poorly understood (Rillig et al., 2020).

Our study revealed ^{13}C enrichment in AMF extraradical hyphae from both HC and RC

during the $^{13}C_1$-myristate tracing experiment (Fig. 1a, b), indicating that the

extraradical hyphae of symbiotic AMF can absorb external non-symbiotic myristate in

the presence of plant-derived carbohydrates. This is further supported by two

additional observations in our study. First, exogenous myristate increased the

transcription of genes associated with fatty acid transport and metabolism in

extraradical hyphae (Fig. 1). Second, in our greenhouse experiment, myristate

application significantly enhanced intraradical and extraradical AMF biomass, even

when the symbiotic C supply from the plant was reduced (Figs. 3 and 4). These

findings indicate that the extraradical hyphae of symbiotic AMF can access external

(non-symbiotic) organic C sources like myristate beyond those directly delivered by

the host, at least under the conditions studied. Notably, myristate is a common root

exudate component in various host and non-AMF host plants (Fernández-Piñar et al.,

2012; Guelette et al., 2012; Li et al., 2017; Liu et al., 2024), and is abundant in litter

and soil substrates (Swenson et al., 2018; Dai et al., 2022). Yet AMF often colonize

and thrive in leaf litter and undecomposed plant material, a long-observed but poorly

understood phenomenon (Went and Stark, 1968; Bunn et al., 2019). Although further

research is required to determine how frequently symbiotic AMF encounter myristate

and to quantify its concentration in various microenvironments, our findings suggest

that in realistic environments, extraradical hyphae of symbiotic AMF could take up

myristate from sources such as root exudates, decaying plant/animal material, litter, or

other microbes, in addition to those symbiotic C source directly provided by the host

plant.

During AM establishment, plant roots activate mycorrhizal defense responses to

limit excessive C consumption caused by over-colonization (Blilou et al., 2000;

Siciliano et al., 2015; Feng et al., 2019). In our study, myristate increased both

intraradical and extraradical biomasses of the symbiotic AMF and promoted the

formation of normal-sized secondary spores (Figs. 2 and 3), unlike the small spores

induced by myristate under asymbiotic conditions (Sugiura et al., 2020). This suggests

that plant roots supply additional substances to AMF beyond myristate, the

identification of which is essential for the pure culture of AMF. Notably, exogenous

myristate transcriptionally downregulated plant genes associated with mycorrhizal

defenses under low-P conditions, including those involved in oxidative stress and

salicylic acid defense responses (Fig. 6), suggesting that myristate could interfere with

normal defense mechanisms in host roots to promote AMF colonization. This effect

differs from AM signals like strigolactones and quercetin, which enhance mycorrhizal

defenses in plants (Sudheeran et al., 2020; Kusajima et al., 2022). Further research is

needed to clarify myristate's role in regulating AMF colonization and development.

Despite enhanced mycorrhization, exogenous myristate unexpectedly reduced

the mycorrhizal P response (MPR) and downregulated mycorrhizal P transporter gene

expression in both alfalfa and rice, indicating a diminished mycorrhizal P benefit.

This could be attributed to several factors. Firstly, AMF can modulate P transport

based on the rate at which plants supply C (Kiers et al., 2011; Argüello et al., 2016).

Myristate, acting as an external non-symbiotic C source, likely lowered the "price" of

symbiotic C supplied by plants, leading to reduced P transfer. Secondly, exogenous

myristate may interfere with AM signaling, as suggested by decreased mycorrhizal

defenses, thereby limiting fungal P acquisition and delivery. This disruption might

partly explain why the reduction in the mycorrhizal P response induced by exogenous

myristate is primarily observed under low-P conditions. Thirdly, AMF P transport

proteins expressed in arbuscular cells can re-uptake P from the symbiotic interface,

creating nutrient competition between the plant and AMF (Xie et al., 2022). In our

study, the increased fungal biomass with exogenous myristate suggests a higher P

demand by AMF.

Meanwhile, external myristate also decreased C supply to AMF from both alfalfa

and rice, except in low-P rice. This reduction may be due to AMF shifting from

utilizing plant-derived C to absorbing external C (myristate), thereby reducing their

competition for plant C at the AM interactional interface (Bitterlich et al., 2014).

Furthermore, the decline in fungal P delivery to the host can be attributed to the

functional coupling of C-P exchange in AM symbiosis (Hammer et al., 2011; Kiers et

al., 2011). Notably, under low-P conditions, myristate increased the allocation of

plant-derived C to AMF in rice, underscoring the influence of both host identity and

soil P status on myristate's effect. Compared to rice, AM-responsive alfalfa exhibited

over tenfold higher ^{13}C transfer efficiency and *RiFAT* gene expression, along with

approximately fivefold higher fungal P transporter gene expression. However, the

increased rice-derived C under low-P conditions did not correspond to greater

mycorrhizal P benefits, suggesting an unrewarding allocation. Overall, this study

reveals that exogenous myristate disrupts the symbiotic C-P exchange. As one of the

oldest coevolving species relationships (Rich et al., 2021), AM symbiosis depends on

effective partner communication and reciprocal metabolic regulation for successful

integration (Lanfranco et al., 2018). Our findings emphasize the importance of

thoroughly understanding this ancient partnership when developing strategies to

enhance AM symbiosis.

Despite these encouraging findings, our work has several limitations. First, the

absorption of myristate by symbiotic AMF was observed following exogenous

myristate application, which may not fully represent realistic environmental

conditions. Second, only one AMF species, *R. irregularis*, was tested in our study,

though AMF species may respond to myristate (or other fatty acids) in varying ways,

at least under asymbiotic conditions (Kameoka et al., 2019; Sugiura et al., 2020).

Furthermore, our investigation into the mechanism by which myristate disrupts C-P

exchange in AM symbiosis remains relatively preliminary. Further research is

therefore needed to verify and quantitatively explore the absorption of non-symbiotic

myristate (or other C sources) by symbiotic AMF under natural conditions, as well as

its regulatory mechanisms and ecological significance to AM symbiosis.

Conclusion

In summary, our study demonstrates that symbiotic AMF can access external organic C sources like myristate in addition to those symbiotically allocated by their host plant. Exogenous myristate promoted the proliferation of both intraradical and extraradical biomasses of symbiotic AMF, potentially due to suppressed mycorrhizal defense responses in host roots. Notably, exogenous myristate disrupted the C-P exchange in AM symbiosis, leading to reduced mycorrhizal P benefits for plants and decreased plant-derived C allocation to AMF, although these effects varied depending on host identity and soil P conditions. These findings provide novel insights into the nutritional interactions in AM symbiosis, which are essential for understanding and managing this ancient relationship.

Materials and methods

Experimental design and biological materials

Three interconnected and complementary experiments were conducted in our

study. The AMF used in all experiments was *Rhizophagus irregularis* (strain: MUCL

41833), a well-studied 'model' AMF species. We investigated whether exogenous

myristate could be directly absorbed by *R. irregularis* (Exp. 1) and examined its effect

on the growth of *R. irregularis* (Exp. 2), where Ri T-DNA-transformed carrot

roots served as the AMF host. In Exp. 3, two phylogenetically distant species—

Medicago sativa L. cv. Sanditi (alfalfa, dicot) and *Oryza sativa* L. cv. Zhonghua 11

(rice, monocot)—were used as AMF hosts in a pot experiment to explore the

mechanisms underlying the effect of exogenous myristate on AM symbiosis.

Exp. 1: $^{13}C_1$-labeled myristate ($^{13}C_1$-Myr) tracing in a two-compartment AMF-

hairy root co-culture system

In Exp. 1, a $^{13}C_1$-My tracing experiment was performed to determine whether

exogenous myristate can be directly absorbed by AMF (*R. irregularis*) extraradical

hyphae associated with carrot hairy root in a two-compartment Petri dish system as

described by St-Arnaud et al. (1996) (90 ×15 mm, see Fig. 1a). The Petri dish was

divided into two compartments by a central plastic barrier. One compartment (root

compartment, RC) contained 25 mL of modified Strullu–Romand (MSR) solid

medium for the growth of carrot hairy root and AMF hyphae. The other compartment

(hyphal compartment, HC), into which only the AMF hyphae, and not roots, were

allowed to grow, contained 15 mL of solid MSR medium. Five 4 cm-long sterile

carrot hairy root segments were initially co-cultivated with 200 *R. irregularis* spores

in the RC in an incubator at 27°C in the dark. The Petri dishes were positioned at an

angle of approximately 45 degrees (with the HC underneath) in the incubator to

facilitate hyphal growth from the RC into the HC. The roots were routinely trimmed

as needed to prevent them from growing into the HC.

When dense AMF extraradical hyphae were observed in the HC (twelve weeks

later), 200 µL of 0.1 mM Myr with 98% $^{13}C_1$ labeling or unlabeled (^{12}C-Myr, control)

myristate was added weekly to the HC until harvest, four weeks later. During the

harvest, AMF extraradical hyphae from the HC and RC were collected separately by

dissolving the media in 10 mM sodium citrate and filtering through a 37 µm nylon

mesh. The carrot hairy roots in the RC were harvested using sterile tweezers.

The determination of ^{13}C enrichment in fatty acids followed our previous report

(Bao et al., 2019). The total ^{13}C enrichment of AMF hyphae from HC or RC were

sealed in tin capsules and analyzed using an elemental analyzer coupled to an IRMS

(EA-IRMS, Flash EA 2000 and Delta V Plus; Thermo Fisher Scientific, Bremen,

Germany). The ^{13}C:^{12}C ratio of CO_2 from the combusted samples was determined

with ± 0.2‰ precision and expressed as atom % of ^{13}C, using a sucrose standard

calibrated against the Vienna Pee Dee Belemnite standard.

Exp. 2: Effects of exogenous myristate on AMF development in a single-

compartment AMF-hairy root co-culture system

In Exp. 2, we evaluated the effects of exogenous myristate at varying

concentrations (0.1, 0.5, 1, 5, or 10 mM) on spore germination and the development

of AMF intraradical and extraradical structures associated with carrot hairy roots in a

single-compartment co-culture system. Briefly, three 4 cm-long sterile carrot hairy

root segments were cultivated in MSR medium with or without *R. irregularis*

inoculations (200 sterilized spores per Petri dish). Potassium myristate solution (0, 0.1,

0.5, 1, 5, or 10 mM, 200 µL) was applied to the plates weekly until harvest. The

germination rate of AMF spores, hyphal branching, and hyphal length were monitored

under a stereomicroscope (M250FA, Leica, Germany) and calculated using ImageJ at

4 days post-inoculation (Schneider et al., 2012). The spore density was dynamically

monitored every 2 weeks starting 8 weeks post-AMF inoculation. The carrot hairy

roots and culture medium were harvested at 14 weeks post-AMF inoculation. The

roots and solid culture medium were carefully separated using sterile tweezers. The

medium was weighed and dissolved in 10 mM sodium citrate, and the AMF collection

were preliminarily separated into hyphae and spores using a blender, followed by

filtration through a 250 µm and 37 µm nylon meshes, respectively, and manual

selection with tweezers under a stereoscopic microscope. Extraradical AMF biomass

was oven-dried at 80°C for 48 h and quantified.

Exp. 3: Effect of exogenous myristate on AM symbiosis in a pot experiment

In the pot experiment (Exp. 3), alfalfa and rice seeds were germinated for 14

days in sterilized sand before being transplanted into a substrate composed of

vermiculite, perlite, and sand (3:1:1, v/v/v). To account for potential biomass

differences between the two plant species, three alfalfa seedlings or one rice seedling

were planted per pot, containing either 300 or 1000 mL of substrate, respectively, with

or without *R. irregularis* inoculation (10 spores/mL of substrate). The plants were

grown in a greenhouse under a long-day photoperiod (16 h light/8 h dark) at

temperatures ranging from 20□30□. For P treatments, 10 mL Hoagland nutrient

solution with either relatively low or high P level for alfalfa (25 or 100 μM KH_2PO_4,

respectively, following He et al., 2020) and rice (100 or 400 μM KH_2PO_4, respectively,

following Shi et al., 2021) was applied twice a week. For myristate application, 5 mL

of potassium myristate solution (0.1 mM) or ddH_2O was applied near the plant roots

twice a week.

The plants were harvested eight weeks post-inoculation. They were carefully

washed with tap water and divided into shoot and root portions. A portion of the fresh

root samples was flash-frozen in liquid nitrogen and stored at $-80°C$ for molecular

analysis. The remaining plant samples were de-enzymed at $105°C$ for 30 min, dried at

$60°C$ for 7 days, and weighed. Plant P concentration was determined using the

molybdenum-antimony colorimetric method.

RNA extraction and Illumina sequencing (Exp. 3)

In Exp. 3, alfalfa and rice roots inoculated with AMF, with or without myristate

application at two P levels, underwent RNA sequencing (RNA-seq) using three

biological replicates (pots, totaling 24 pots). Total RNA was extracted from 100 mg of

root tissue using the TRIzol R Plus RNA Purification Kit (Takara, Dalian, China)

following the manufacturer's instructions. Library preparation was conducted with the

NebNext Ultra RNA Library Prep Kit (New England Biolabs, Ipswich, MA, USA).

Qualified libraries were sequenced on the Illumina HiSeq 4000 platform with 150-bp

paired-end reads. Fastp was employed for basic statistics and quality filtering of raw

data. Hisat2 was utilized to map the sequencing results to the rice reference genome

(IRGSP-1.0) or the alfalfa reference genome (Zhongmu NO.1, Shen et al., 2020).

Salmon quant was applied for gene expression quantification analysis.

Quantitative real-time PCR (qRT-PCR, Exp. 3)

In Exp. 3, RNA extracted from alfalfa or rice roots was reverse-transcribed into

cDNA using the QuantiTect Reverse Transcription Kit (Qiagen, Valencia, CA). qRT-

PCR was performed on a QuantStudio™ 5 Real-Time PCR System (Applied

Biosystems, CA, USA) with the SYBR ROX Mastermix Kit (Qiagen, Valencia, CA)

following the manufacturer's instructions. Three biological replicates (samples) were

utilized for each treatment, with three technical replicates per sample. The relative

expression levels of genes involved in C delivery from plants to AMF (*MsSTR2*,

OsSTR2, *RiFAT1*, *RiFAT2*, and *RiMST2*) and in the mycorrhizal P pathway (*MsPT8*,

OsPT11 and *RiPT*) were analyzed, using the alfalfa *actin 2* gene (*MsACT2*), rice cycle

2 gene (*OsCyc2*) or *R. irregularis* elongation factor-1 beta gene (*RiEF-1β*,) as

reference genes (Yang et al., 2012; Corrêa et al., 2014; Brands and Dörmann, 2022;

Das et al., 2022). Details of all the qRT-PCR primers used are provided in the

Supporting Information (Table S1).

^{13}C labeling of plants, lipid extraction and measurement of signature fatty acids

(Exp. 3)

In Exp. 3, the $^{13}CO_2$ labeling procedure was conducted as described in our

previous report, with minor modifications (Bao et al., 2019). Five days before harvest,

alfalfa and rice plants (five randomly selected replicates) were placed in a sealed

acrylic chamber containing $^{13}CO_2$ and incubated for 4 hours each.

The extraction and analysis of signature fatty acids were performed following the

method described by Olsson et al. (2005), with slight modifications based on Bao et al.

(2019). Fatty acid methyl esters were identified and quantified relative to an added

internal standard (fatty acid methyl ester 19:0). NLFA 16:1ω5 was used as a marker

for AMF in roots (Olsson and Lekberg, 2022).

Quantification of ^{13}C enrichment and excess ^{13}C in fatty acids (Exp. 3)

In Exp. 3, the ^{13}C enrichment in roots and fatty acid methyl esters was analyzed

using IRMS coupled to a Trace GC Ultra gas chromatograph via the ConFlow IV

interface (Thermo Fisher Scientific, Bremen, Germany), as described in our previous

study (Bao et al., 2019). The chromatographic conditions matched those specified for

lipid analysis, using helium (He) as the carrier gas. The effluent from the capillary

column passed through an Al tube filled with CuO wires at 860°C. The δ^{13}C values

were calibrated against the V-PDB using the CO_2 standard (RM 8542 (IAEA-CH-6,

sucrose)) and corrected for the additional non-labeled methyl group introduced during

methanolysis. The ^{13}C enrichment was calculated by subtracting the background ^{13}C

levels determined from five non-labeled controls (Olsson et al., 2005).

Quantification of AMF colonization intensity (Exps. 1, 2, and 3)

To assess AMF colonization intensity, 100 root segments (approximately 1 cm in

length) were treated with a 10% KOH solution at 95°C for 10 minutes, decolorized in

a 30% H_2O_2 solution at 25°C for 3 min, softened in a 1% HCl solution at 25°C for 3

min, and then stained with 0.05% Trypan Blue at 95°C for 15 min. The evenly

arranged root segments were examined under 400× magnification using a light

microscope (Nikon Eclipse 80i, Japan). The colonization rate was calculated using the

gridline intersect method (Giovannetti and Mosse, 1980).

Statistical analysis

Mycorrhizal response (MR) was calculated based on plant biomass and plant P

concentration, respectively, following Corrêa et al. (2014): MR (%) = (AM − NM) /

NM × 100%, where AM denotes the observed value for the target parameter (biomass

or P concentration) in mycorrhizal plants, and NM represents the mean values of

nonmycorrhizal plants. Multiple comparisons among different treatments were

conducted using one-way analysis of variance (ANOVA) followed by post hoc

Tukey's tests (assuming equal variances). A univariate two-way ANOVA, followed by

a post hoc Tukey test, was employed to assess the effects of myristate, P level, and

their interactions on AMF colonization intensity and plant P concentration. The

significant differences between AM and NM groups were determined using Student's t

test (for normally distributed data). For RNA-seq data, the DESeq2 R package was

utilized to conduct differentially expressed gene (DEG) analysis between plants with

(Myr$^+$) and without (Myr$^-$) myristate application (Love et al., 2014). The *p* values

obtained were adjusted using the Benjamini-Hochberg correction to control false

discovery rate. Genes (transcripts) with an adjusted P-value < 0.05 and $|\log_2(\text{fold}$

change$)| > 1$ identified by DESeq2 were considered as differentially expressed. The

clusterProfiler method was applied for Gene Ontology (GO) enrichment analysis of

DEGs (Wu et al., 2021), with GO terms showing an adjusted p-value < 0.05

considered significantly enriched.

Data accessibility

The RNA-seq data generated in this study have been deposited in the Sequence

Read Archive at the National Center for Biotechnology Information under BioProject

ID PRJNA1083641. All data required to evaluate the conclusions in the paper are

included in the paper and/or the Supplementary Materials.

Author contributions

Hanwen Chen: Writing – review & editing, Investigation, Formal analysis,

Methodology, Visualization. **Tian Xiong:** Investigation, Formal analysis, Data

curation, Methodology. **Baoxing Guan:** Writing – original draft, Investigation,

Formal analysis, Data curation, Visualization. **Jiaqi Huang:** Investigation, Resources.

Danrui Zhao: Investigation, Resources, Visualization. **Yao Chen:** Investigation,

Resources. **Haoran Liang:** Investigation, Visualization. **Yingwei Li:** Investigation,

Resources. **Jingwen Wu:** Investigation, Resources. **Shaoping Ye:** Investigation,

Resources. **Ting Li:** Investigation, Resources. **Wensheng Shu:** Supervision. **Jin-tian**

Li: Supervision. **Yutao Wang:** Writing – original draft, Writing – review & editing,

Conceptualization, Data curation, Funding acquisition, Supervision, Project

administration.

Acknowledgments

We thank the financial support from the National Key R&D Program of China

(Key special Project for Marine Environmental Security and Sustainable

Development of Coral Reefs 2022-702), the National Natural Science Foundation of

China (nos. 32272203 and 31772397), the Key-Area Research and Development

Program of Guangdong Province (2022B0202110001), the Agricultural and Social

Development Project of Guangzhou Municipal Science and Technology Bureau (no.

202206010058), and the Guangdong Provincial Key Laboratory of Plant Resources

(no. 2023PlantKFP03).

Competing interest

The authors declare that they have no known competing financial interests or

personal relationships that could have appeared to influence the work reported in this

paper.

References

Argüello A, O'Brien MJ, van der Heijden MGA, Wiemken A, Schmid B, Niklaus PA.

2016. Options of partners improve carbon for phosphorus trade in the arbuscular

mycorrhizal mutualism. *Ecology Letters* **19**: 648–656.

Bao X, Wang Y, Olsson PA. 2019. Arbuscular mycorrhiza under water—Carbon–

phosphorus exchange between rice and arbuscular mycorrhizal fungi under

different flooding regimes. *Soil Biology and Biochemistry* **129**: 169–177.

Bitterlich M, Krügel U, Boldt-Burisch K, Franken P, Kühn C. 2014. The sucrose

transporter SlSUT2 from tomato interacts with brassinosteroid functioning and

affects arbuscular mycorrhiza formation. *The Plant Journal* **78**: 877–889.

Blilou I, Ocampo JA, García-Garrido JM. 2000. Induction of Ltp (lipid transfer

protein) and Pal (phenylalanine ammonia-lyase) gene expression in rice roots

colonized by the arbuscular mycorrhizal fungus Glomus mosseae. *Journal of*

Experimental Botany **51**: 1969–1977.

Brands M, Dörmann P. 2022. Two AMP-binding domain proteins from *Rhizophagus*

irregularis involved in import of exogenous fatty acids. *Molecular Plant-*

Microbe Interactions **35**: 464–476.

Bravo A, Brands M, Wewer V, Dörmann P, Harrison MJ. 2017. Arbuscular

mycorrhiza-specific enzymes FatM and RAM2 fine-tune lipid biosynthesis to

promote development of arbuscular mycorrhiza. *New Phytologist* **214**: 1631–

1645.

Brundrett MC, Tedersoo L. 2018. Evolutionary history of mycorrhizal symbioses and

global host plant diversity. *New Phytologist* **220**: 1108–1115.

Bücking H, Abubaker J, Govindarajulu M, Tala M, Pfeffer PE, Nagahashi G,

Lammers P, Shachar-Hill Y. 2008. Root exudates stimulate the uptake and

metabolism of organic carbon in germinating spores of *Glomus intraradices*.

New Phytologist **180**: 684–695.

Bunn RA, Simpson DT, Bullington LS, Lekberg Y, Janos DP. 2019. Revisiting the

"direct mineral cycling" hypothesis: arbuscular mycorrhizal fungi colonize leaf

litter, but why? *The ISME Journal* **13**: 1891–1898.

Cope KR, Kafle A, Yakha JK, Pfeffer PE, Strahan GD, Garcia K, Subramanian S,

Bücking H. 2022. Physiological and transcriptomic response of *Medicago*

truncatula to colonization by high- or low-benefit arbuscular mycorrhizal fungi.

Mycorrhiza **32**: 281–303.

Corrêa A, Cruz C, Pérez-Tienda J, Ferrol N. 2014. Shedding light onto nutrient

responses of arbuscular mycorrhizal plants: nutrient interactions may lead to

unpredicted outcomes of the symbiosis. *Plant Science* **221–222**: 29–41.

Dai H, Zhang X, Zhao B, Shi J, Zhang C, Wang G, Yu N, Wang E. 2022. Colonization

of mutualistic mycorrhizal and parasitic blast fungi requires OsRAM2-regulated

fatty acid biosynthesis in rice. *Molecular Plant-Microbe Interactions* **35**: 178–

186.

Das D, Paries M, Hobecker K, Gigl M, Dawid C, Lam HM, Zhang J, Chen M,

Gutjahr C. 2022. PHOSPHATE STARVATION RESPONSE transcription factors

enable arbuscular mycorrhiza symbiosis. *Nature Communications* **13**: 1–12.

Feng F, Sun J, Radhakrishnan GV, Lee T, Bozsóki Z, Fort S, Gavrin A, Gysel K,

Thygesen MB, Andersen KR, Radutoiu S, Stougaard J, Oldroyd GED. 2019. A

combination of chitooligosaccharide and lipochitooligosaccharide recognition

promotes arbuscular mycorrhizal associations in *Medicago truncatula*. *Nature*

Communications **10**: 5047.

Fernández-Piñar R, Espinosa-Urgel M, Dubern JF, Heeb S, Ramos JL, Cámara M.

2012. Fatty acid-mediated signaling between two Pseudomonas species.

Environmental Microbiology Reports **4**: 417–423.

Foster KR, Kokko H. 2006. Cheating can stabilize cooperation in mutualisms.

Proceedings of the Royal Society B: Biological Sciences **273**: 2233–2239.

Giovannetti M, Mosse B. 1980. An evaluation of techniques for measuring vesicular-

arbuscular mycorrhizal infection in roots. *New Phytologist* **84**: 489–500.

Guelette BS, Benning UF, Hoffmann-Benning S. 2012. Identification of lipids and

lipid-binding proteins in phloem exudates from *Arabidopsis thaliana*. *Journal of*

Experimental Botany **63**: 3603–3616.

Gutjahr C, Radovanovic D, Geoffroy J, Zhang Q, Siegler H, Chiapello M, Casieri L,

An K, An G, Guiderdoni E, Kumar CS, Sundaresan V, Harrison MJ, Paszkowski

U. 2012. The half-size ABC transporters STR1 and STR2 are indispensable for

mycorrhizal arbuscule formation in rice. *The Plant Journal* **69**: 906–920.

Hammer EC, Pallon J, Wallander H, Olsson PA. 2011. Tit for tat? A mycorrhizal

fungus accumulates phosphorus under low plant carbon availability. *FEMS*

Microbiology Ecology **76**: 236–244.

Hata S, Kobae Y, Banba M. 2010. Interactions between plants and arbuscular

mycorrhizal fungi. *International Review of Cell and Molecular Biology* **281**: 1–

48.

Helber N, Wippel K, Sauer N, Schaarschmidt S, Hause B, Requena N. 2011. A

versatile monosaccharide transporter that operates in the arbuscular mycorrhizal

fungus *Glomus sp* is crucial for the symbiotic relationship with plants. *The Plant*

Cell **23**: 3812–3823.

Jiang Y, Wang W, Xie Q, Liu N, Liu L, Wang D, Zhang X, Yang C, Chen X, Tang D,

Wang E. 2017. Plants transfer lipids to sustain colonization by mutualistic

mycorrhizal and parasitic fungi. *Science* **356**: 1172–1175.

Johnson NC. 2010. Resource stoichiometry elucidates the structure and function of

arbuscular mycorrhizas across scales. *New Phytologist* **185**: 631–647.

Kameoka H, Tsutsui I, Saito K, Kikuchi Y, Handa Y, Ezawa T, Hayashi H, Kawaguchi

M, Akiyama K. 2019. Stimulation of asymbiotic sporulation in arbuscular

mycorrhizal fungi by fatty acids. *Nature Microbiology* **4**: 1654–1660.

Keymer A, Pimprikar P, Wewer V, Huber C, Brands M, Bucerius SL, Delaux PM,

Klingl V, Röpenack-Lahaye EV, Wang TL, Eisenreich W, Dörmann P, Parniske

M, Gutjahr C. 2017. Lipid transfer from plants to arbuscular mycorrhiza fungi.

eLife **6**: e29107.

Kiers ET, Duhamel M, Beesetty Y, Mensah J, Franken O, Verbruggen E, Fellbaum CR,

Kowalchuk GA, Hart MM, Bago A, Palmer TM, West SA, Vandenkoornhuyse P,

Jansa J, Bücking H. 2011. Reciprocal rewards stabilize cooperation in the

mycorrhizal symbiosis. *Science* **333**: 880–882.

Kusajima M, Fujita M, Soudthedlath K, Nakamura H, Yoneyama K, Nomura T,

Akiyama K, Maruyama-Nakashita A, Asami T, Nakashita H. 2022.

Strigolactones modulate salicylic acid-mediated disease resistance in

Arabidopsis thaliana. *International Journal of Molecular Sciences* **23**: 5246.

Lanfranco L, Fiorilli V, Gutjahr C. 2018. Partner communication and role of nutrients

in the arbuscular mycorrhizal symbiosis. *New Phytologist* **220**: 1–10.

Li S, Xu C, Wang J, Guo B, Yang L, Chen J, Ding W. 2017. Cinnamic, myristic and

fumaric acids in tobacco root exudates induce the infection of plants by

Ralstonia solanacearum. *Plant and Soil* **412**: 381–395.

He H, Wu M, Guo L, Fan C, Zhang Z, Su R, Q Peng, J Pang, Lambers H. 2020.

Release of tartrate as a major carboxylate by alfalfa (*Medicago sativa* L.) under

phosphorus deficiency and the effect of soil nitrogen supply. *Plant and Soil* **449**:

169–178.

Liu X, Feng Z, Zhang W, Yao Q, Zhu H. 2023. Exogenous myristate promotes the

colonization of arbuscular mycorrhizal fungi in tomato. *Frontiers in Plant*

Science **14**: 1250684.

Liu Y, Zhang H, Wang J, Gao W, Sun X, Xiong Q, Shu X, Miao Y, Shen Q, Xun W,

Zhang R. 2024. Nonpathogenic *Pseudomonas syringae* derivatives and its

metabolites trigger the plant "cry for help" response to assemble disease

suppressing and growth promoting rhizomicrobiome. *Nature Communications* **15**:

1907.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**: 550.

Luginbuehl LH, Menard GN, Kurup S, Van Erp H, Radhakrishnan GV, Breakspear A,

Oldroyd GED, Eastmond PJ. 2017. Fatty acids in arbuscular mycorrhizal fungi

are synthesized by the host plant. *Science* **356**: 1175–1178.

Luo Q, Sun L, Hu X, Zhou R. 2014. The variation of root exudates from the

hyperaccumulator *Sedum alfredii* under cadmium stress: metabonomics analysis.

PLoS ONE **9**: e115581.

MacLean AM, Bravo A, Harrison MJ. 2017. Plant signaling and metabolic pathways

enabling arbuscular mycorrhizal symbiosis. *The Plant Cell* **29**: 2319–2335.

Mori N, Nishiuma K, Sugiyama T, Hayashi H, Akiyama K. 2016. Carlactone-type

strigolactones and their synthetic analogues as inducers of hyphal branching in

arbuscular mycorrhizal fungi. *Phytochemistry* **130**: 90–98.

Olsson PA, van Aarle IM, Gavito ME, Bengtson P, Bengtsson G. 2005. ^{13}C

incorporation into signature fatty acids as an assay for carbon allocation in

arbuscular mycorrhiza. *Applied and Environmental Microbiology* **71**: 2592–2599.

Olsson PA, Lekberg Y. 2022. A critical review of the use of lipid signature molecules

for the quantification of arbuscular mycorrhiza fungi. *Soil Biology and*

Biochemistry **166**: 108574.

Pfeffer PE, Douds Jr DD, Becard G, Shachar-Hill Y. 1999. Carbon uptake and the

metabolism and transport of lipids in an arbuscular mycorrhiza. *Plant Physiology*

120: 587–598.

Rich MK, Vigneron N, Libourel C, Keller J, Xue L, Hajheidari M, Radhakrishnan GV,

Le Ru A, Diop SI, Potente G, Conti E, Duijsings D, Batut A, Le Faouder P,

Kodama K, Kyozuka J, Sallet E, Bécard G, Rodriguez-Franco M, Ott T,

Bertrand-Michel J, Oldroyd GED, Szövényi P, Bucher M, Delaux PM. 2021.

Lipid exchanges drove the evolution of mutualism during plant terrestrialization.

Science **372**: 864–868.

Rillig MC, Aguilar-Trigueros CA, Anderson IC, Antonovics J, Ballhausen MB,

Bergmann JBM, Chaudhary VB, Deveautour C, Grünfeld L, Hempel S, Lakovic

M, Lammel DR, Lehmann A, Lehmann J, Leifheit EF, Liang Y, Li E, Lozano

YM, Manntschke A, Mansour I, Oviatt P, Pinek L, Powell JR, Roy J, Ryo M,

Sosa-Hernández MA, Veresoglou SD, Wang D, Yang G, Zhang H. 2020.

Myristate and the ecology of AM fungi: significance, opportunities, applications

and challenges. *New Phytologist* **227**: 1610–1614.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of

image analysis. *Nature Methods* **9**: 671–675.

Shen C, Du H, Chen Z, Lu H, Zhu F, Chen H, Meng X, Liu Q, Liu P, Zheng L, Li X,

Dong J, Liang C, Wang T. 2020. The chromosome-level genome sequence of the

autotetraploid alfalfa and resequencing of core germplasms provide genomic

resources for alfalfa research. *Molecular Plant* **13**: 1250–1261.

Shi J, Zhao B, Zheng S, Zhang X, Wang X, Dong W, Xie Q, Wang G, Xiao Y, Chen F,

Yu N, Wang E. 2021. A phosphate starvation response-centered network

regulates mycorrhizal symbiosis. *Cell* **184**: 5527–5540.e18.

Siciliano I, Amaral Carneiro G, Spadaro D, Garibaldi A, Gullino ML. 2015. Jasmonic

acid, abscisic acid, and salicylic acid are involved in the phytoalexin responses of

rice to *Fusarium fujikuroi*, a high gibberellin producer pathogen. *Journal of*

Agricultural and Food Chemistry **63**: 8134–8142.

Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*. Academic press, London.

St-Arnaud M, Hamel C, Vimard B, Caron M, Fortin JA. 1996. Enhanced hyphal

growth and spore production of the arbuscular mycorrhizal fungus *Glomus intraradices* in an in vitro system in the absence of host roots. *Mycological Research* **100**: 328–332.

Sudheeran PK, Ovadia R, Galsarker O, Maoz I, Sela N, Maurer D, Feygenberg O, Oren Shamir M, Alkan N. 2020. Glycosylated flavonoids: fruit's concealed antifungal arsenal. *New Phytologist* **225**: 1788–1798.

Sugiura Y, Akiyama R, Tanaka S, Yano K, Kameoka H, Marui S, Saito M, Kawaguchi M, Akiyama K, Saito K. 2020. Myristate can be used as a carbon and energy source for the asymbiotic growth of arbuscular mycorrhizal fungi. *Proceedings of the National Academy of Sciences of the United States of America* **117**: 25779–25788.

Swenson TL, Karaoz U, Swenson JM, Bowen BP, Northen TR. 2018. Linking soil biology and chemistry in biological soil crust using isolate exometabolomics. *Nature Communications* **9**: 19.

Tanaka S, Hashimoto K, Kobayashi Y, Yano K, Maeda T, Kameoka H, Ezawa T, Saito K, Akiyama K, Kawaguchi M. 2022. Asymbiotic mass production of the arbuscular mycorrhizal fungus *Rhizophagus clarus*. *Communications Biology* **5**: 43.

Went FW, Stark N. 1968. The biological and mechanical role of soil fungi. *Proceedings of the National Academy of Sciences of the United States of America* **60**: 497–504.

Wipf D, Krajinski F, van Tuinen D, Recorbet G, Courty PE. 2019. Trading on the arbuscular mycorrhiza market: from arbuscules to common mycorrhizal networks. *New Phytologist* **223**: 1127–1142.

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, Fu X, Li

S, Bo X, Yu G. 2021. ClusterProfiler 4.0: A universal enrichment tool for

interpreting omics data. *The Innovation (Camb)* **2**: 100141.

Xie X, Lai W, Che X, Wang S, Ren Y, Hu W, Chen H, Tang M. 2022. A SPX domain-

containing phosphate transporter from *Rhizophagus irregularis* handles

phosphate homeostasis at symbiotic interface of arbuscular mycorrhizas. *New*

Phytologist **234**: 650–671.

Yang S, Grønlund M, Jakobsen I, Grotemeyer MS, Rentsch D, Miyao A, Hirochika H,

Kumar CS, Sundaresan V, Salamin N, Catausan S, Mattes N, Heuer S,

Paszkowski U. 2012. Nonredundant regulation of rice arbuscular mycorrhizal

symbiosis by two members of the phosphate transporter1 gene family. *The Plant*

Cell **24**: 4236–4251.

Yu H, He Y, Zhang W, Chen L, Zhang J, Zhang X, Dawson W, Ding J. 2022. Greater

chemical signaling in root exudates enhances soil mutualistic associations in

invasive plants compared to natives. *New Phytologist* **236**: 1140–1153.

Zhang Q, Blaylock LA, Harrison MJ. 2010. Two Medicago truncatula half-ABC

transporters are essential for arbuscule development in arbuscular mycorrhizal

symbiosis. *The Plant Cell* **22**: 1483–1497.

Fig. 1 Uptake of exogenous myristate by extraradical hyphae (ERH) of arbuscular

mycorrhizal fungi (AMF) under symbiotic conditions and its effects on

mycorrhization and expression of fungal marker genes for fatty acid uptake and

metabolism. **(a)** Schematic illustration of the two-compartment AMF-carrot hairy root

co-cultivation system and the addition of myristate to the hyphae compartment (HC).

Carrot hairy roots colonized by *Rhizophagus irregularis* were grown in the root

compartment (RC). Starting from the twelfth week, 0.1 mM of $^{13}C_1$-labeled ($^{13}C_1$-

Myr), non-labeled (^{12}C-Myr) myristate, or ddH$_2$O (Myr$^-$, control) was added weekly

into the HC with well-developed AMF hyphae until harvest. **(b)** Typical branched

absorbing structures (BAS) were observed exclusively in the HC treated with

exogenous myristate (Myr$^+$). **(c)** ^{13}C/^{12}C ratio of the harvested ERH from HC or RC

(n = 3–5) when ^{13}C$_1$-labeled (^{13}C$_1$-Myr) or non-labeled (^{12}C-Myr, background)

myristate was added to HC. The background ^{13}C/^{12}C ratio of extraradical hyphae in

HC (1.0922 ± 0.0003, n = 5) and RC (1.0923 ± 0.0002, n = 3) was identical. The

exogenous myristate added in HC cannot diffuse directly into RC, as evidenced by the

identical ^{13}C/^{12}C ratio of MSR medium collected in RC in ^{13}C$_1$-Myr (1.0966 ± 0.0012,

n = 3) and ^{12}C-Myr (1.0967 ± 0.0032, n = 3) treatments. **(d)** AMF percent root

colonization in carrot hairy roots from the RC when HC was treated with myristate

(Myr$^+$) or ddH$_2$O (Myr$^-$) (n = 3–10). **(e-f)** Normalized expression (relative to *RiEF-1β*)

of AMF genes potentially involved in fatty acid uptake (*RiFAT1* and *RiFAT2*) and

metabolism (β-oxidation (*RiFAD1*), palmitvaccenic acid synthesis (*RiOLE1*), and

TCA cycle (*RiCIT1*) in AMF ERH collected from HC with myristate (Myr$^+$) or

ddH$_2$O (Myr$^-$) treatments (n = 4). Means ± SE, *, **, and *** indicate statistical

significance at the 0.05, 0.01, and 0.001 probability levels (Student's *t*-test),

respectively.

Fig. 2 Effects of exogenous myristate on the development of intraradical and extraradical structures of arbuscular mycorrhizal fungal (AMF) in the single-compartment AMF-carrot hairy root co-cultivation system. **(a)** Schematic illustration of the single-compartment AMF-hairy root co-cultivation system and the addition of myristate (Myr⁺) or not (Myr⁻). **(b)** Spore germination rate, **(c)** hyphal branching, and **(d)** extraradical hyphal length in the co-cultivation system with (Myr⁺) or without (Myr⁻) myristate treatment at 4 days post-inoculation (dpi) (n = 11–12). **(e)** AMF percent root colonization in carrot hairy roots and **(f)** biomass of the AMF extraradical

structures (per gram of medium, n = 3−6). **(g)** Dynamics of AMF spore density over

time (per square centimeter, n = 3−5); and **(h)** AMF spore diameter at harvest with 0,

0.1, 0.5, 1, 5, or 10 mM myristate treatment under symbiotic conditions (n = 5,

calculation from > 900 spores in each replicate). Means ± SE; different letters indicate

significant differences at $P < 0.05$ level (Tukey's post hoc test); *** indicates

statistical significance at the 0.001 probability level (Student's *t*-test).

Fig. 3 Effects of exogenous myristate on arbuscular colonization and the intraradical

and extraradical biomass of arbuscular mycorrhizal fungi (AMF) associated with

Medicago sativa or *Oryza sativa* under low (LP) or high (HP) phosphorus conditions.

(a, b) Arbuscule colonization in *M. sativa* and *O. sativa* roots treated with 0.1 mM

myristate (Myr[+]) or ddH$_2$O (Myr[−]) (n = 10; Total, vesicle, and hyphal colonization

rates are provided in Supplementary Figure S1); **(c, d)** AMF biomass, indicated by the

AMF signature fatty acid (NLFA 16:1ω5), in AMF-colonized roots and rhizosphere

soil of *M. sativa* and *O. sativa* treated with myristate (Myr[+]) or ddH$_2$O (Myr[−]) (n = 5).

Means ± SE; different letters indicate significant differences at the *P* < 0.05 level

(Tukey's post hoc test).

Fig. 4 Effects of exogenous myristate on the symbiotic carbon (C) allocation from

Medicago sativa and *Oryza sativa* to arbuscular mycorrhizal fungi (AMF) and the

expression profiles of mycorrhizal C transportation genes. **(a, b)** Plant ^{13}C flow to the

AMF signature fatty acid (NLFA 16:1ω5) in AMF-colonized roots of *M. sativa* and *O.*

sativa treated with 0.1 mM myristate (Myr$^+$) or ddH$_2$O (Myr$^-$) under low (LP) and

high (HP) phosphorus conditions (n = 5). (**c–h**) Normalized expression of

mycorrhizal C transportation genes in AMF-colonized (AM) and non-colonized (NM)

roots treated with 0.1 mM myristate (Myr$^+$) or ddH$_2$O (Myr$^-$) (n = 3). The mycorrhizal

C transportation genes include the plant symbiotic fatty acid transportation gene *STR2*

(**c, d**), the fungal fatty acid transporters (*RiFAT1, RiFAT2*; **e, f**), and the

monosaccharide transporter (*RiMST2*; **g, h**). The normalized expression of *M. sativa*,

O. sativa, and AMF genes was calculated relative to *MsACT2*, *OsCyc2*, and *RiEF-1β*,

respectively. Two-way ANOVA in each histogram applies only to AM roots. Means ±

SE; * and ** indicate statistical significance at the 0.05 and 0.01 probability levels

(Student's *t*-test), respectively.

Fig. 5 Effects of exogenous myristate on plant phosphorus (P) concentration, mycorrhizal P response (MPR), and expression of mycorrhizal P pathway marker genes in *Medicago sativa* and *Oryza sativa*. **(a, b)** Plant P concentration (n = 10); **(c, d)** MPR (n = 10); and **(e, f)** relative expression of *MsPT8*, *OsPT11*, and *RiPT* (n = 3) of *M. sativa* and *O. sativa* roots under low (LP) and high (HP) phosphorus conditions with 0.1 mM myristate (Myr$^+$) or ddH$_2$O (Myr$^-$) treatments. AM, inoculated with *Rhizophagus irregularis*; NM, non-inoculated. The normalized expression of *M. sativa*, *O. sativa*, and AMF genes was calculated relative to *MsACT2*, *OsCyc2*, and *RiEF-1β*, respectively. Two-way ANOVA in each histogram (c–f) applies only to AM roots. Means ± SE; different letters indicate significant differences at $P < 0.05$ level

887   (Tukey's post hoc test). * and *** indicate statistical significance at the 0.05 and

888   0.001 probability levels (Student's t-test), respectively.

Fig. 6 Effects of exogenous myristate on the transcriptome of *Medicago sativa* and
Oryza sativa roots inoculated with arbuscular mycorrhizal fungi (AMF) under low
(LP) and high (HP) phosphorus conditions. **(a, b)** Significantly enriched gene
ontology (GO) terms ($P_{adj} < 0.05$) for the differentially expressed genes (DEGs)
between myristate (Myr^+) and ddH_2O (Myr^-) treatments. Underlines denote stress- or
defense-related GO terms. **(c, d)** Relative expression of genes associated with stress-
or defense-related GO terms between Myr^+ and Myr^- treatments. Underlined are
known stress- or defense-related transcription factors or marker genes. Genes
associated with multiple GO terms are marked in yellow. * indicate statistical
significance between Myr^+ and Myr^- treatments at the 0.05 probability level
(Benjamini-Hochberg approach).