Altering the redox status of *Chlamydia trachomatis* directly impacts its developmental

cycle progression

Vandana Singh and Scot P. Ouellette*

Department of Pathology, Microbiology, and Immunology, College of Medicine, University of

Nebraska Medical Center, Omaha, NE 68198

*Correspondence: scot.ouellette@unmc.edu

Keywords: *Chlamydia*, Alkyl hydroperoxide reductase subunit C (AhpC), CRISPRi,

Differentiation, Development, Oxidative stress, Peroxides, Reactive oxygen species (ROS),

ROS scavengers

ABSTRACT

Chlamydia trachomatis is an obligate intracellular bacterial pathogen with a unique

developmental cycle. It differentiates between two functional and morphological forms:

elementary body (EB) and reticulate body (RB). The signals that trigger differentiation from one

form to the other are unknown. EBs and RBs have distinctive characteristics that distinguish

them, including their size, infectivity, proteome, and transcriptome. Intriguingly, they also differ in

their overall redox status as EBs are oxidized and RBs are reduced. We hypothesize that

alterations in redox may serve as a trigger for secondary differentiation. To test this, we

examined the function of the primary antioxidant enzyme alkyl hydroperoxide reductase subunit

C (AhpC), a well-known member of the peroxiredoxins family, in chlamydial growth and

development. Based on our hypothesis, we predicted that altering the expression of *ahpC* would

modulate chlamydial redox status and trigger earlier or delayed secondary differentiation. To test

this, we created *ahpC* overexpression and knockdown strains. During *ahpC* knockdown, ROS

levels were elevated, and the bacteria were sensitive to a broad set of peroxide stresses.

Interestingly, we observed increased expression of EB-associated genes and concurrent higher

production of EBs at an earlier time in the developmental cycle, indicating earlier secondary

differentiation occurs under elevated oxidation conditions. In contrast, overexpression of AhpC

created a resistant phenotype against oxidizing agents and delayed secondary differentiation.

Together, these results indicate that redox potential is a critical factor in developmental cycle

progression. For the first time, our study provides a mechanism of chlamydial secondary

differentiation dependent on redox status.

Word count: 250

INTRODUCTION

All organisms that are exposed to oxygen are necessarily subjected to oxidative stress. Specifically, the process of metabolizing substrates in the presence of oxygen can generate reactive oxygen species (ROS), which are toxic at high enough concentrations. Thus, from bacteria to humans, systems have evolved to mitigate the accumulation of ROS. At the same time, host defense mechanisms have evolved to leverage ROS production as a means of limiting pathogen growth and survival. Not surprisingly, pathogens have co-evolved to resist these defense mechanisms. For example, many pathogens possess a variety of antioxidant enzymes, such as catalases, glutathione peroxidases, and peroxiredoxins, that help them both subvert ROS-mediated immune system assaults and mitigate metabolic ROS byproducts (Staerck et al., 2017; Wan et al., 2021).

Chlamydia is an obligate intracellular bacterium that has significantly reduced its genome size and content in adapting to obligate host dependence. *Chlamydia trachomatis*, the leading cause of bacterial sexually transmitted diseases and preventable infectious blindness, lacks homologs to catalases or glutathione peroxidases but does possess a homolog of Alkyl hydroperoxide reductase subunit C (AhpC). AhpC is a 2-cys peroxiredoxin and is widely conserved in prokaryotes (de Oliveira et al., 2021). Peroxiredoxins can scavenge hydrogen peroxide, peroxynitrite, and organic hydroperoxides (Parsonage et al., 2008; Poole & Ellis, 1996; Seaver & Imlay, 2001) and act as the primary scavenger in pathogens that lack both catalase and glutathione peroxidases (Mastronicola et al., 2014; Richard et al., 2011). Several studies from other bacterial systems have shown that AhpC has a significant role in ROS and RNI scavenging, virulence, and persistence (Cosgrove et al., 2007; Kimura et al., 2012; Oh & Jeon, 2014), and deletion of AhpC results in elevated level of ROS within the bacterium (Zhang et al., 2019). Though *Chlamydia* is dependent on its host for most of its energy requirements, it has some metabolic activities, such as a partial TCA cycle and oxidative phosphorylation (Gerard et al., 2002; Iliffe-Lee & McClarty, 1999), which can be a possible source of intracellular

ROS, to which the bacteria must adapt. There is a paucity of knowledge on how *C. trachomatis*

modulates oxidative stress. However, some studies explored the effects of redox changes on

chlamydial growth. Boncompain et al. reported that infection of *C. trachomatis* induced the

transient production of ROS by the host cell at a moderate level for the initial few hours of

infection only (Boncompain et al., 2010). Another study also found a similar observation about

ROS during infection, reporting that *Chlamydia* requires host-derived ROS for its growth (Abdul-

Sater et al., 2010). However, the key mechanisms *Chlamydia* employs to manage ROS-

mediated stress have not been characterized.

Chlamydia undergoes a complex developmental cycle that comprises two distinct

morphological forms, the elementary body (EB) and reticulate body (RB) (Abdelrahman &

Belland, 2005). The EB is the smaller (~0.3 µm), infectious, and nondividing form capable of

infecting susceptible host cells. Once internalized into a host-derived vacuole termed an

inclusion, the EB differentiates into the non-infectious, larger (~1 µm), and replicating RB - this

process is known as primary differentiation and represents the early phase of the developmental

cycle (Clifton et al., 2005). RBs replicate within the inclusion in the midcycle phase using an

asymmetric, MreB-dependent polarized division process (Abdelrahman et al., 2016; Lee et al.,

2020; Ouellette et al., 2022; Ouellette et al., 2020). In the late phase of the developmental cycle,

RBs asynchronously condense into EBs, and this process is termed secondary differentiation.

Despite these well-defined differences between chlamydial morphological forms, how

Chlamydia mechanistically differentiates between functional forms remains unclear.

Intriguingly, a recent study evaluated the redox potential of *Chlamydia* and demonstrated

that RBs are reduced, whereas EBs are oxidized (Wang et al., 2014) (Fig. 1A). This is

consistent with earlier studies that revealed differences in the crosslinking of outer membrane

proteins and type III secretion-related proteins in chlamydial developmental forms (Betts-

Hampikian & Fields, 2011; Caldwell et al., 1981; Everett & Hatch, 1995; Wang et al., 2014).

Taken together, these observations indicate that, during the developmental cycle, the redox

potential of *Chlamydia* is changing. However, whether redox changes in *Chlamydia* directly

affect developmental cycle progression is not characterized.

Based on the difference between redox status of EBs and RBs, we hypothesized that

changing redox conditions is a critical factor in the process of differentiation from one form to the

other. We named this the "redox threshold hypothesis". In this scenario, as soon as a given RB

has crossed an oxidative threshold, the activity of critical proteins is modified to trigger

differentiation to the EB (Fig. 1B). We used chlamydial transformants designed to overexpress

or reduce AhpC levels to explore the effects of altered redox potential in chlamydial growth and

development to test this hypothesis (Fig. 1C). This study establishes the role of AhpC as an

antioxidant in *Chlamydia,* as demonstrated by its ability to counteract different peroxide stresses

when overexpressed. Overexpression of AhpC had no negative effect on bacterial replication

but delayed the differentiation of RBs to EBs. In contrast, under conditions of *ahpC* knockdown,

the organism was highly sensitive to oxidizing conditions. Interestingly, this change in redox

potential caused earlier expression of EB-associated (late) genes and production of EBs,

leading to a shift in developmental cycle progression. This earlier activation of gene expression

related to secondary differentiation in *ahpC* knockdown was also observed when developmental

cycle progression was blocked by penicillin treatment. Taken together these data provide

mechanistic insight into chlamydial secondary differentiation and are the first to demonstrate

redox-regulated differentiation in *Chlamydia*.

RESULTS

Overexpression of *ahpC* has minimal impact on chlamydial growth

To study the effect of AhpC on chlamydial developmental cycle progression, we first generated

an *ahpC* overexpression (OE) strain using a plasmid encoding an anhydrotetracycline (aTc)

inducible *ahpC* (untagged) and transformed it into a *C. trachomatis* L2 strain lacking its

endogenous plasmid (-pL2). The same plasmid vector backbone (i.e., encoding mCherry in

place of *ahpC*) was used as an empty vector control (EV). HeLa cells were infected with these

transformants, and, at 10 hpi, expression of the construct was induced or not. Lacking an

antibody against AhpC, overexpression of *ahpC* was validated by reverse transcription-

quantitative PCR (RT-qPCR). Here, an approximate 1-log increase in transcripts of *ahpC* was

detected at 14 and 24 hpi in the induced strain in comparison to the uninduced strain (Fig. 2A).

Immunofluorescence analysis (IFA) was performed at 14 and 24 hpi to examine the organisms'

morphology and overall inclusion growth. For IFA, individual bacteria were labeled with an anti-

major outer membrane protein (MOMP) antibody, and DAPI was used to stain for DNA. IFA

imaging revealed that there was no observable impact on the morphology of the EV control

strain under the conditions tested. In contrast, we did note that overexpression of

ahpC increased the overall inclusion area (Fig. 2B and 2C). We next quantified the total number

of bacteria (i.e., both RBs and EBs) by measuring genomic DNA and observed a significant

increase in gDNA levels at 24 hpi in response to increased *ahpC* expression (Fig. 2D). Although

inclusions were larger in area and contained more total bacteria as assessed by gDNA levels

when overexpressing *ahpC*, recoverable inclusion forming units (IFUs: a measure of infectious

EBs) were significantly lower under these conditions at 24 hpi (Fig. 2E). However, by 48 hpi, the

difference between the induced and uninduced conditions, although still reduced, was not

statistically significant. The average number of IFUs was 5.30×10^7 and 1.56×10^9 for the *ahpC*

OE strain in the uninduced condition at 24 and 48 hpi, respectively. For the EV strain, the

average number of IFUs was 7.07×10^6 and 1.90×10^8 in the uninduced condition at 24 and 48

hpi, respectively. The inclusion forming unit analysis only measures viable EBs from a

population, suggesting that AhpC overexpression skews the ratio of RBs to EBs. To test this, we

calculated the ratio of IFUs to gDNA at 24 hpi, which revealed that the relative number of RBs is

higher (and EBs lower) as a result of overexpression of *ahpC* (Fig 2E). Overall, these data

support the hypothesis that overexpressing *ahpC* delays production of infectious EBs.

Overexpression of *ahpC* confers resistance to peroxides in *Chlamydia*

AhpC is an important peroxiredoxin involved in oxidative damage defense. Before determining

whether increased *ahpC* expression impacts chlamydial sensitivity to oxidizing agents, we first

sought to determine the response of *C. trachomatis* to inorganic or organic hydroperoxides such

as hydrogen peroxide (H_2O_2), cumene hydroperoxide (CHP), and tert-butyl hydroperoxide

(TBHP) as well as peroxynitrite (PN). To do this, we evaluated the sensitivity of HeLa cells,

infected or not with the empty vector (EV) control strain, to different concentrations of oxidizing

agents using a viability assay. Both uninfected and EV-infected HeLa cells tolerated up to 1 mM

oxidizing agents for 30 min (Fig. S1). Next, different concentrations (lower than 1 mM) of these

oxidants were tested on wild-type (WT) *C. trachomatis* L2/434/Bu (*Ctr* L2). HeLa cells were

infected with *Ctr* L2 and, at 16 hpi, were exposed to different concentrations of exogenous

oxidizing agents for 30 min only, before washing out the oxidizing agents and replacing the

media. Samples for IFU and IFA assays were then collected at 24 hpi to assess effects of the

oxidizing agents on growth of *Ctr* L2. 62.5 µM concentration of all three inorganic and organic

peroxides had no appreciable effect on IFUs, and inclusion size and morphology also remained

unaffected at this sublethal concentration. The concentrations of inorganic and organic

peroxides that resulted in a decrease in IFUs to ~50% of the untreated culture were 500 µM

(H_2O_2) and 250 µM (CHP and TBHP), respectively, with a concurrent decrease in inclusion size.

1 mM H_2O_2, CHP, and TBHP caused >90% reduction in IFUs and small inclusions. PN at 1 mM

concentration was less effective than other oxidizing agents and showed only a ~30% decline in

IFUs compared to the untreated culture (Fig. S2).

To further examine the antioxidant functions of AhpC, we next exploited our *ahpC*

overexpression strain to explore its capacity to protect chlamydiae from oxidizing agents. Both

ahpC OE and EV strains were used to infect HeLa cells and, at 10 hpi, expression of the

constructs was induced or not with 1 nM aTc. At 16 hpi, infected cells were exposed to various

concentrations of exogenous oxidizing agents for 30 min. At 24 hpi, IFU and IFA samples were

collected to measure chlamydial growth and assess chlamydial morphology, respectively. As

shown in Figure 3A, 3B, and S3, the *ahpC* overexpression strain showed increased resistance

to all oxidants tested as the mean IFUs and the size of inclusions were greater than that of the

uninduced but treated control. There was no change in bacterial growth and morphology in the

case of 62.5 µM concentrations of oxidizing agents, which was anticipated since we determined

this was a sublethal concentration. The resistant phenotype was more evident at the higher

concentrations of H_2O_2, CHP, and TBHP. Overexpression of *ahpC* also contributed to higher

resistance against PN. Conversely, the empty vector control strain behaved as expected (i.e.,

like WT) in the presence of oxidizing agents, with similar results observed in the presence or

absence of aTc (Fig. 3C, 3D, and S3). Collectively, these data demonstrate that the chlamydial

AhpC possesses antioxidant activity, as expected, and that increased *ahpC* expression is

protective for *Chlamydia* in the presence of increased oxidative stress.

Knockdown of *ahpC* negatively impacts chlamydial growth

To further define the function(s) of AhpC in the chlamydial developmental cycle, we used a

novel dCas12-based CRISPR interference (CRISPRi) strategy adapted for *Chlamydia* by our

lab (Ouellette et al., 2021). We generated an *ahpC* knockdown strain harboring the

pBOMBL12CRia with a crRNA targeting the *ahpC* 5' intergenic region and overlapping the ATG

start site (plasmid designated as pL12CRia(*ahpC*)). We used a strain carrying a pL12CRia

plasmid with a crRNA with no homology to any chlamydial sequence (i.e., non-targeting [NT]) to

serve as a negative control. To confirm the knockdown of *ahpC*, RT-qPCR was employed. Here,

we infected HeLa cells with the *ahpC* knockdown (*ahpC* KD) or NT strains and induced dCas12

expression or not at 10 hpi using 1 nM aTc. Nucleic acid samples were harvested at 14 hpi and

24 hpi. RT-qPCR analysis of the *ahpC* KD revealed approximately 90% reduction of *ahpC*

transcripts compared to the uninduced control, thus confirming the knockdown of *ahpC* (Fig.

4A). In the case of the NT control, there was no effect on *ahpC* transcripts. IFA of these strains

revealed noticeably smaller inclusions after blocking expression of *ahpC* as compared to the

uninduced sample or the NT conditions (Fig. 4B). We next measured total bacterial counts using

genomic DNA. Surprisingly, even though inclusions were smaller during *ahpC* knockdown, we

observed higher gDNA levels at 14 hpi compared to the uninduced control, followed by only a

small increase from 14 to 24 hpi (Fig. 4C). In contrast, the uninduced strain showed a

logarithmic increase in gDNA levels during this timeframe, similar to the NT strain (Fig. 4C). To

further explore the effect of reduced *ahpC* transcripts in *Chlamydia*, IFU assays were

performed. IFU analysis exhibited severely reduced progeny (>90%) in *ahpC* knockdown

compared to its respective uninduced control at both time points assessed (24 and 48 hpi) (Fig.

4D). In contrast, the NT strain showed less than a 50% reduction after inducing dCas12

expression at these timepoints consistent with prior observations that dCas12 expression

slightly delays developmental cycle progression (Hatch & Ouellette, 2023; Reuter et al., 2023).

The ratio of IFUs to gDNA at 24 hpi is significantly higher for the *ahpC* KD, suggesting a lower

number of RBs in proportion to EBs (Fig 4E). Taken together, these analyses indicate that

reducing *ahpC* levels and/or activity severely reduced chlamydial growth at 24 and 48 hpi.

To investigate whether *ahpC* knockdown resulted in increased ROS levels in the bacteria

(Zhang et al., 2019), the intracellular ROS levels were measured in infected cells in *ahpC*

knockdown conditions and compared to the uninduced control condition and uninfected cells.

We used the cell-permeable, fluorogenic dye CellROX Deep Red to measure ROS levels. This

dye remains non-fluorescent in a reduced state and exhibits bright fluorescence once oxidized

by ROS. ROS generation was measured in uninfected and *ahpC* KD-infected HeLa cells at 24

and 48 hpi. As shown in Figure 5A, *ahpC* knockdown resulted in significantly higher ROS than

the uninfected or infected but uninduced samples at both 24 hpi and 48 hpi time points. We also

performed live-cell microscopy for visualization of ROS generation in uninfected and *ahpC* KD-

infected HeLa cells at 24 and 40 hpi. These microscopy images revealed that *ahpC* KD resulted

in higher ROS in comparison to the uninduced and uninfected cells (Fig. 5B and S4).

Importantly, the signal was associated with chlamydial inclusions and not the host cell, thus

indicating that the measurements from Figure 5A were from the bacteria. These results together

indicate that AhpC knockdown results in increased ROS levels in *C. trachomatis*.

Knockdown of *ahpC* sensitizes *Chlamydia* to oxidizing agents

As overexpression of *ahpC* resulted in increased resistance to oxidizing agents, we next

explored whether knockdown of *ahpC* resulted in increased sensitivity to such agents. As

described above, *ahpC* KD and NT strains were used to infect HeLa cells and, at 10 hpi,

expression of dCas12 was induced or not with 1 nM aTc. Considering that the *ahpC* KD already

demonstrated reduced IFUs and inclusion sizes at 24 hpi, only sublethal concentrations (62.5

μM) of exogenous oxidizing agents were applied. At 24 hpi, IFA and IFU analyses were

performed to quantify the effect of the oxidants on chlamydial growth in the absence of *ahpC*

activity. Under conditions of *ahpC* knockdown, even sublethal concentrations (62.5 μM) of H_2O_2,

CHP, or TBHP further reduced the inclusion size, and the IFU data revealed a decrease from

>90% to <30% in comparison to the untreated control (Fig. 6A, 6B, and S5). Reduced

expression of *ahpC* also affected the survival of *Chlamydia* against peroxynitrite. Notably, there

was no significant change in the NT control strain in the uninduced and induced samples in

response to oxidizing agents as assessed by IFU and IFA (Fig. 6C, 6D, and S5). These data

further support that the chlamydial AhpC is a critical antioxidant enzyme in these bacteria.

Complementation restores the growth and resistance to low levels of peroxide stress of

the *ahpC* knockdown strain

To validate that the impaired growth, altered inclusion morphology, and enhanced sensitivity to

peroxides were due to the decreased level/activity of AhpC during knockdown, we generated a

complemented strain to restore *ahpC* expression during knockdown. For the construction of the

ahpC complementing plasmid, the *ahpC* gene was cloned and transcriptionally fused 3' to the

dCas12 in the pL12CRia(*ahpC*) knockdown plasmid. Here, the complementing *ahpC* allele is

also under the control of the aTc induced P$_{tet}$ promoter and is co-expressed with the aTc

inducible dCas12. Consequently, the *ahpC* knockdown effect is ablated, and the observed

phenotypes should be restored. After inducing *dCas12-ahpC* expression with aTc, the resultant

strain was verified by RT-qPCR. Increased transcripts for *ahpC* were quantified under these

conditions, indicating successful complementation of the knockdown effect (Fig. 7A). Of note,

the *ahpC* transcript levels remained elevated in comparison to the uninduced control at the 24

hpi time point. After confirming this strain, we measured genomic DNA to quantify total bacteria

and performed IFA and IFU assays to examine if complementation restored the phenotypes

observed during *ahpC* knockdown. These assays revealed normal inclusion morphology, gDNA

levels, and EB progeny production in the complemented strain, indicating successful

complementation of the knockdown phenotype (Fig. 6B, C, and D). Of note, a C-terminal 6xHis

tagged AhpC was not capable of complementing the knockdown phenotype (data not shown),

indicating a requirement for a free C-terminus in the function of AhpC in *Chlamydia*. Previous

studies in other bacteria have revealed that the C-terminal residues in AhpC play a crucial role

in the structural stability and enzymatic activity of AhpC (Dip et al., 2014; Feng et al., 2020; Wan

et al., 2021).

We next assessed whether complementation of the knockdown phenotype could also

restore the resistance to low levels of peroxide stress. As in previous experiments, a sublethal

concentration of oxidizing agents was added for 30' at 16 hpi after having induced *dCas12-*

ahpC expression at 10 hpi. Consistent with the growth parameters (Fig. 7B-D), the

complemented strain showed wild-type responses in these conditions (Fig. 7E, 7F, and S5).

These data indicate that the enhanced susceptibility of the *ahpC* knockdown strain to oxidizing

agents was due to the reduced levels of *ahpC* and not an indirect effect of knockdown.

We predicted that the growth defects resulting from higher production of ROS during

ahpC knockdown could be rescued by treating the *ahpC* KD with ROS scavengers. To test this

prediction, we utilized two characterized ROS scavengers, DMTU (*N,N'-dimethylthiourea*) and

α-tocopherol, which scavenge H_2O_2 and peroxyl radical, respectively (Hollander-Czytko et al.,

2005; Kiffin et al., 2006; Walch et al., 2015). First, the effect of ROS scavengers on uninfected

and infected HeLa cells was investigated using a viability assay. This assay revealed that 10

mM DMTU and 100 µM α-tocopherol had no adverse effects on uninfected or EV-infected HeLa

cells (data not shown). These same concentrations were tested on WT *Ctr* L2 infected HeLa

cells in untreated and 500 µM H_2O_2 treated conditions to test the potency of these scavengers

to rescue growth defects associated with this concentration of oxidizing agent. Scavengers were

added at 9.5 hpi and washed away at 16 hpi - the time of addition of H_2O_2 in the respective

samples. At 16.5 hpi, after 3 wash steps, scavengers were added again with fresh media, and,

at 24 hpi, bacterial growth and morphology were assessed using IFU and IFA assays. Neither

scavenger had a significant impact on chlamydial morphology or infectious progeny. In the case

of peroxide (500 µM H_2O_2) treated samples, scavengers restored IFUs from ~50% to ~100%,

and inclusion size was also recovered (Fig. S6A and B).

Next, the effect of the ROS scavengers under *ahpC* knockdown conditions was

assessed. In this experiment, scavengers were added at 9.5 hpi to provide the protective effect

of scavengers before reducing the activity of *ahpC*. At 10 hpi, knockdown was induced or not

with 1 nM aTc, and, at 14 hpi, scavengers were added again. At 24 hpi, IFU and IFA samples

were collected to examine bacterial growth and morphology to assess the effect of ROS

scavenging on the *ahpC* knockdown phenotype. As shown in Fig. 7G and 7H, both ROS

scavengers had a positive impact on restoring growth of the *ahpC* knockdown strain; inclusions

were larger, and IFUs were increased from <10% to >90% in the presence of scavengers in the

induced samples. These data provide compelling evidence that the adverse effects of *ahpC*

knockdown are due to increased ROS accumulation in this strain.

Chlamydial developmental cycle progression is altered by *ahpC* knockdown/

overexpression

Whereas the overexpression of AhpC delayed overall developmental progression and was

consistent with our hypothesis, the results with the *ahpC* knockdown strain appeared to refute

our hypothesis given the apparent decrease in IFUs and smaller inclusion sizes after

knockdown. To investigate the effects of changes in redox potential on chlamydial

developmental cycle progression, we performed a transcriptional analysis of well-characterized

late-cycle genes associated with secondary differentiation (*hctA*, *hctB*, *glgA*, *tsp,* and *omcB*)

(Barry et al., 1993; Belland et al., 2003; Brickman et al., 1993; Gehre et al., 2016; Newhall,

1987; Swoboda et al., 2023). In the uninduced samples, transcript levels of all the tested late

genes were higher at 24 hpi compared to 14 hpi, indicating their normal expression during the

late stage of the developmental cycle. In comparison to the uninduced control, under conditions

of *ahpC* knockdown, significantly higher expression of *hctA*, *hctB*, *omcB, tsp,* and *glgA* was

observed at the mid-developmental cycle timepoint of 14 hpi (Fig. 8A). At 24 hpi, expression of

these genes was comparable between the uninduced and induced conditions in the *ahpC*

knockdown strain. We subsequently performed RNA sequencing from the uninduced and

induced *ahpC* knockdown strain at 14 hpi. Experimental conditions were same as described for

the RT-qPCR study, and samples were processed as mentioned in materials and methods.

RNA-seq results were statistically analyzed by the UNMC Bioinformatics Core facility (Table

S2). We further categorized these genes based on significant difference (p<0.05) and fold-

change (≥1.5). Using these stringent parameters, we identified 161 up-regulated genes and 145

down-regulated genes in the *ahpC* knockdown conditions (Table S2). Here, we observed a

global increase in late gene transcripts at this mid-developmental cycle timepoint (Fig. 8B and

Table 1), indicating that increased oxidation results in earlier increases in late gene transcripts.

In contrast, the complementation strain showed no increase in the expression of these tested

late genes at 14 hpi, and *tsp* and *glgA* transcripts were reduced (Fig. S7A). Consistent with our

model, the overexpression of *ahpC* resulted in significantly lower expression of *hctA*, *hctB*,

omcB, tsp, and *glgA* at 14 and 24 hpi (Fig. S7B), suggesting a delayed transition to EBs.

Given the increase in late gene expression during *ahpC* knockdown at an earlier time

point in the developmental cycle than normal, we reasoned that the increase in oxidizing

conditions might prematurely trigger secondary differentiation and EB production. However,

complicating such an analysis is that fewer RBs are present to convert to EBs, which would

result in overall lower IFU yields as we had measured at 24 and 48 hpi (Fig. 4D). Nonetheless,

to assess EB production more rigorously at earlier time points in the developmental cycle, we

performed a growth curve analysis to precisely measure IFUs at two-hour intervals from 16 to

24 hpi. HeLa cells were infected, and knockdown was induced or not at 10 hpi with 1 nM aTc. At

the indicated time points, IFU samples were collected and quantified. Consistent with our

prediction, this experiment revealed higher IFUs (i.e., EBs) at 16 and 18 hpi during *ahpC*

knockdown compared to the uninduced samples (Fig. 8C). This difference was statistically

significant at 18 hpi. However, further EB production was stalled, with the uninduced strain

continuing to produce EBs such that, by 24 hpi, there were significantly higher EB yields under

these conditions (as noted in Fig. 4D). These data show that the phenotypic consequence of

higher expression at 14 hpi of genes functionally related to EBs is the concomitant earlier

production of EBs. These data underscore that reduced activity of *ahpC* causes earlier

secondary differentiation in *C. trachomatis*.

ahpC knockdown activates transcription of late-cycle genes when bacterial replication is

blocked

We detected earlier expression of EB-related genes during *ahpC* knockdown and were

curious if the *ahpC* knockdown condition could activate these genes under conditions when the

chlamydial developmental cycle is blocked. Pathogenic *Chlamydia* species undergo a polarized

cell division process, in which peptidoglycan is transiently synthesized only at the division

septum (Abdelrahman et al., 2016; Cox et al., 2020; Ouellette et al., 2020). As a result, during

penicillin treatment, division of RBs is blocked (Moulder, 1993; Moulder et al., 1956; Ouellette et

al., 2020). However, the bacteria continue to grow in size resulting in aberrantly enlarged RBs

(Barbour et al., 1982; Matsumoto & Manire, 1970; Ouellette et al., 2012), in which EB-related

genes are not transcribed and production of EBs is inhibited (Ouellette et al., 2006; Panzetta et

al., 2018).

To examine this, we generated an *ahpC* KD strain with spectinomycin resistance (*ahpC*

KD-spec) and validated its phenotype as being the same as the penicillin-resistant *ahpC* KD

strain*.* This new strain, *ahpC* KD-spec, was used to infect HeLa cells, and knockdown was

induced or not at 10 hpi with 1 nM aTc. At the same time, samples were treated or not with 1

unit per mL of penicillin (Pen). RNA samples were harvested at 16 and 24 hpi and processed for

RT-qPCR (Fig. S8B-D). IFA controls from these different conditions demonstrated the expected

phenotypes (Fig. S8A). For example, Pen treatment caused aberrantly enlarged RBs,

irrespective of *ahpC* KD, whereas *ahpC* KD itself caused smaller inclusions. Similarly, *ahpC* KD

resulted in reduced *ahpC* transcripts in the induced conditions irrespective of the presence of

Pen (Fig. S8B). Next, we quantified transcript levels for the late genes *hctA* and *hctB* in these

different conditions (Fig. S8C and D). As expected, in the uninduced and untreated condition,

both transcripts had higher expression at 24 hpi, indicating their regular developmental

expression during the late stage of the developmental cycle. Consistent with prior observations

(Ouellette et al., 2006), levels of these genes in uninduced + Pen conditions was low. As we

previously noted (Fig. 8A), the transcripts of *hctA* and *hctB* are higher in the induced than

uninduced samples at 16 hpi in the absence of Pen. Interestingly, in the induced + Pen

condition, these genes had higher expression at 24 hpi than the uninduced + Pen control. At 16

hpi, their expression was either higher or similar between both samples (i.e., induced + Pen and

uninduced + Pen). Collectively, these data support our observation that *ahpC* knockdown

activates the transcription of late-cycle genes – even under conditions where developmental

cycle   progression   is   blocked.

DISCUSSION

Many bacteria use redox sensing as a mechanism to control gene expression and

subsequent morphologic transitions. Typically, these organisms use a redox-sensitive

transcription factor, such as OxyR to sense these changes in redox conditions and alter gene

expression appropriately. For example, in *Pseudomonas aeruginosa*, OxyR is involved in

regulation of cellular metabolism, in addition to oxidative stress defense (Wei et al., 2012). In

Caulobacter crescentus, redox has been reported as a regulator of metabolism and cell cycle

progression (Hartl et al., 2020; Narayanan et al., 2015*).* Thus, the ability of bacteria to assess

oxidative stress (imbalanced level of ROS) and to make physiological adjustments by altering

gene expression patterns that favor their growth, and development is critical for their survival.

ROS generation is an inevitable condition for pathogens growing in aerobic conditions,

and resistance against oxidative stress is a key survival mechanism (Fang, 2011). Hence,

pathogens have evolved detoxifying proteins, such as peroxiredoxins, to eliminate ROS (Dip et

al., 2014; Wang et al., 2004; Yang et al., 2002). AhpC has been reported to have a crucial role in

bacterial physiology, survival, and virulence by scavenging ROS and RNI (Cosgrove et al.,

2007; Kimura et al., 2012; Loprasert et al., 2003; Oh & Jeon, 2014). In the process of

detoxification of peroxides and peroxynitrite, AhpC is converted into an oxidized dimer, requiring

alkyl hydroperoxide reductase AhpF or AhpD to regenerate its activity (Koshkin et al., 2004;

Poole & Ellis, 1996; Wong et al., 2017). *C. trachomatis* encodes AhpC (Ct603/CTL0866) but

lacks any annotated homologs of AhpF or AhpD. Therefore, it remains an open question how

AhpC activity is regulated. Some studies have mentioned *ahpC* as an iron-responsive gene in

Chlamydia (Brinkworth et al., 2018; Pokorzynski et al., 2019), but there is no detailed

investigation to date about the role of this crucial antioxidant in the growth and development of

Chlamydia. Our study is the first to characterize a function of AhpC in chlamydial biology.

An essential aspect of the scavenging activity of AhpC for many bacterial pathogens in

which it has been studied is that it works best with endogenous (i.e., low) levels of H_2O_2. These

bacteria, such as *Staphylococcus aureus* and *Yersinia pseudotuberculosis*, encode catalases,

which have high Km for hydrogen peroxide that serve a predominant role in scavenging

exogenous H_2O_2 (Cosgrove et al., 2007; Wan et al., 2021). Catalases can detoxify H_2O_2 at high

levels (millimolar levels) and are crucial in responding to external H_2O_2 stress (Mishra & Imlay,

2012). The *ahpC* mutants in these bacteria become sensitive to organic peroxides but resistant

to H_2O_2 due to higher catalase activity as a compensatory response to the lack of AhpC

(Antelmann et al., 1996; Mongkolsuk et al., 2000; Ochsner et al., 2000). Further, simultaneous

mutations in both catalase and *ahpC* displayed drastically enhanced sensitivity to all oxidizing

agents tested (Cosgrove et al., 2007; Ezraty et al., 2017; Seaver & Imlay, 2001). *C. trachomatis*

does not encode a catalase gene (Boncompain et al., 2014; Rusconi & Greub, 2013), and

hypersensitivity of *ahpC* knockdown to both inorganic and organic peroxides indicates the

absence of catalase and establishes that AhpC is the primary scavenger of ROS in *C.*

trachomatis. Contrary to previous studies, we did not observe significant effects of PN with

altered AhpC expression. One possible explanation may be our study's concentration (1 mM) of

PN. Higher concentrations of PN could not be used due to toxic effects on the host cell (data not

shown). The other possibility may be the presence of some other unknown mechanism(s) to

detoxify PN. For example, *Chlamydia* also encodes a superoxide dismutase (SOD) that may

prevent the accumulation of the necessary precursors that are needed for PN production. We

are currently investigating the function of the chlamydial SOD enzyme.

Resistance to different oxidants as a result of overexpression of *ahpC* (Fig. 3 and S3) is

consistent with studies in other bacteria (Sherman et al., 1996; Zuo et al., 2014), signifying that

AhpC is the principal defense mechanism against oxidative stress in *Chlamydia*. During *ahpC*

knockdown, the attenuated activity of AhpC severely affected the growth of *Chlamydia* (Fig. 4).

Notably, the observed morphology of inclusions during *ahpC* knockdown is strikingly similar to

Chlamydia treated with a high concentration (1 mM) of oxidizing agents used (Fig. 4 and Fig

S2), suggesting both conditions result in increased oxidative stress to the organism. This was

further supported by the increased sensitivity of *ahpC* KD to sublethal concentrations of

oxidants (Fig. 6 and S4), consistent with previous data from other bacteria (Cosgrove et al.,

2007; Seaver & Imlay, 2001; Storz et al., 1989; Zhang et al., 2019). In *ahpC* KD, the ROS level

was dramatically higher (Fig. 5 and S4), similar to previous studies in other bacterial systems

(Zhang et al., 2019). Further, the addition of ROS scavengers to the culture medium rescued the

negative phenotypes associated with *ahpC* knockdown (Fig. 7G and H). This strongly suggests

that, in the absence of AhpC, higher amounts of reactive oxygen species accumulate in

Chlamydia and that the negative impact on growth during *ahpC* knockdown is due to highly

oxidized conditions in the organism. Moreover, for the complete restoration of inclusion size,

multiple doses of scavengers were required throughout the experiment, further emphasizing that

the internal chlamydial environment is oxidized and/or susceptible to oxidation in the absence of

AhpC. These findings suggest that AhpC in *Chlamydia* is indispensable, having a functional role

even more extensive than previously reported in other bacteria.

In further support of this observation, we noticed upregulated transcripts of late-cycle

genes such as *hctA*, *hctB*, *glgA, omcA, omcB*, *dsbA*, and *tsp* at an earlier time (14 hpi) in the

chlamydial developmental cycle as a result of reduced AhpC activity (Fig. 8). Notably, in the

developmental cycle of *C. trachomatis*, only RBs are present at 14 hpi, and secondary

differentiation starts after 16 hpi (Abdelrahman & Belland, 2005). Expression of most of the late-

cycle genes occurs after this time in the developmental cycle (Belland et al., 2003). These late-

cycle genes are well characterized for their association with EBs. The *hctA* and *hctB* genes

encode histone-like proteins responsible for chromosomal condensation during the

differentiation of RBs into EBs. The *hctA* gene is among the first to be transcribed in the late

stage (Chiarelli et al., 2020). Tsp is a periplasmic protease thought to be crucial in the

degradation of RB-specific periplasmic proteins during secondary differentiation in *C.*

trachomatis (Swoboda et al., 2023). OmcA, OmcB, and DsbA are responsible for the rigid cell

wall and osmotic stability of the EBs (Christensen et al., 2019; Hatch, 1996; Newhall, 1987).

GlgA is key enzyme for glycogen accumulation in the inclusion lumen at late stages of the

developmental cycle (Gehre et al., 2016). One crucial point to consider is the redox status of

Chlamydia, 14 hpi is the time when reduced RBs predominate with virtually no oxidized EBs

detectable. However, we established that *ahpC* knockdown shifts the redox status of the

bacteria towards oxidation. Hence, the earlier and significantly higher detection of transcripts for

these late-cycle genes indicate earlier secondary differentiation in the *ahpC* knockdown.

Importantly, *ahpC* complementation during knockdown restored the regular developmental

expression of these late-cycle genes, further supporting that this phenotype was due to the

highly oxidized conditions created by reduced AhpC activity. The conclusion of these data is that

AhpC levels, and presumably activity, have a direct effect on secondary differentiation.

Secondary differentiation (differentiation from RBs to EBs) is an essential step for

chlamydial growth and survival, but there is a dearth of information regarding the mechanisms

of its regulation. EBs and RBs have significantly different proteomic repertoires (Saka et al.,

2011; Skipp et al., 2005; Skipp et al., 2016), and our group has identified critical functions for the

cytoplasmic ClpXP and ClpCP and periplasmic Tsp proteases during secondary differentiation

(Pan et al., 2023; Swoboda et al., 2023; Wood et al., 2022).The *tsp* gene, in addition to another

late-cycle gene, *hctB*, is transcriptionally regulated by sigma factor 28 (σ^{28}) (Douglas & Hatch,

2000; Hatch & Ouellette, 2023). Another sigma factor, σ^{54} has been linked to regulation of outer

membrane components, type III secretion system components, and other genes typically

expressed late in development (Hatch & Ouellette, 2023; Soules et al., 2020). In addition to the

two minor sigma factors, σ^{28} and σ^{54}, *Chlamydia* encodes one major sigma factor, σ^{66}, which is

transcriptionally regulated by the relative levels of RsbV1 (antagonist) and RsbW (anti-sigma

factor) (Thompson et al., 2015). This Rsb system senses ATP availability and has been

proposed to regulate σ^{66} (Thompson et al., 2015). A related study from Soules et al. (Soules et

al., 2020) showed that TCA intermediates act as ligands for RsbU, thereby linking the Rsb

system to the TCA cycle and ATP synthesis. Collectively, these data show that post-translational

mechanisms drive secondary differentiation in *Chlamydia*. However, no definitive "switch" that

triggers this step has been identified.

How might increasing oxidation regulate secondary differentiation? *Chlamydia* lacks an

identifiable ortholog of OxyR so some other mechanism must be employed. We speculate that

increasing ROS production from the metabolic activity of the RB will overcome the innate and

stochastic activity of AhpC to detoxify ROS. This could either directly alter the activity of specific

proteins or lead to the accumulation of damaged proteins that in turn triggers secondary

differentiation. These possibilities are not mutually exclusive. We propose a simple model to

explain this (Fig. 9). As secondary differentiation is asynchronous and RBs divide through an

asymmetric budding mechanism (Ouellette et al., 2020), AhpC as well as proteins impacted by

ROS levels are unevenly distributed between mother and daughter cell. This difference will lead

some RBs to breach an oxidative threshold sooner, allowing activation of late genes and

secondary differentiation earlier than other RBs. Consistent with this, the higher expression of

genes functionally related to EBs and EB production during *ahpC* KD at an early stage of the

chlamydial developmental cycle indicates earlier secondary differentiation as an outcome of

diminished activity of *ahpC* in *C. trachomatis*. Consequently, increased late gene transcription

should have the phenotypic effect of causing earlier production of EBs. Indeed, we quantified

more IFUs in the *ahpC* knockdown at 16 and 18 hpi compared to the uninduced control

condition. During *ahpC* KD, higher amounts of ROS accumulation in the bacterium create highly

oxidized conditions. In contrast, *ahpC* overexpression scavenges ROS at a greater level leading

to a more reducing environment. For the AhpC overexpression strain, the expression of late-

cycle genes in the induced conditions compared to the uninduced control at 24 hpi is

significantly lower with reduced production of EBs. These data support our hypothesis that the

developmental cycle is delayed in these conditions with a concomitant delay in achieving the

oxidative threshold, thus allowing RBs to continue to divide before committing to secondary

differentiation. Taken together, our data directly link oxidation state and secondary differentiation

in *Chlamydia*. Ongoing studies are focused on characterizing redox-sensitive chlamydial

proteins to understand which specific factors drive the shift from RBs to EBs (or EBs to RBs) in

C. trachomatis.

MATERIALS AND METHODS

Strains and cell culture

For chlamydial transformation, McCoy mouse fibroblast cells (kind gift of Dr. Harlan Caldwell (NIH/NIAID)) were used. Human cervix adenocarcinoma epithelial HeLa cells (kind gift of Dr. Harlan Caldwell (NIH/NIAID)) were used for RT-qPCR, immunofluorescence assays (IFA), inclusion forming unit assays (IFU), viability assays, oxidative stress, RNA-seq, ROS measurements, and ROS scavenger assays. Both cell types were routinely grown and passaged in Dulbecco's modified Eagle's medium (DMEM; Gibco, Waltham, MA) supplemented with 10% fetal bovine serum (FBS; Sigma, St. Louis, MO) and 10 µg/mL gentamicin (Gibco, Waltham, MA) at 37°C and 5% CO_2. All strains were verified to be Mycoplasma-negative using LookOut mycoplasma PCR detection kit (Sigma). For chlamydial transformations, *Chlamydia trachomatis* serovar L2 EBs lacking the endogenous pL2 plasmid (kind gift of Dr. Ian Clarke, University of Southampton) were used (Wang et al., 2011). Wild-type, density gradient-purified *Chlamydia trachomatis* 434/Bu (ATCC VR902B) EBs were used for sensitivity to oxidizing agents and ROS scavenger assays. Molecular biology reagents, oxidizing agents, CellROX Deep Red dye, and scavengers were purchased from Thermo Fisher unless otherwise noted.

Plasmid construction

The primers, gBlock gene fragments, plasmids, and bacterial strains used for molecular cloning are listed in Table S1 in the supplemental material. Constructs for chlamydial transformation were cloned using high-fidelity (HiFi) cloning system from New England BioLabs (NEB). Primers were designed using the NEBuilder online primer generation tool (https://nebuilderv1.neb.com). For the overexpression strain, the *ahpC* gene was amplified by PCR with Phusion DNA polymerase (NEB) using *C. trachomatis* serovar L2 434/Bu genomic DNA as a template. The PCR product was purified using a PCR purification kit (Qiagen, Hilden, Germany). The HiFi assembly reaction was performed as per the manufacturer's instructions in conjunction with the pBOMBDC plasmid digested with Fast Digest EagI and KpnI enzymes and dephosphorylated

with FastAP (Thermo Fisher, Waltham, MA). The HiFi reaction mix was transformed into *E. coli*

10-beta (NEB). Plasmids were first confirmed by restriction enzyme digestion, and final

verification of insert was performed using Sanger sequencing. A dCas12-based CRISPRi

approach was used for *ahpC* knockdown generation (Ouellette et al., 2021). The

pBOMBL12CRia plasmid having anhydrotetracycline (aTc) inducible catalytically dead dCas12

protein was used as vector. A crRNA targeting the 5' intergenic region of *ahpC* was designed

and ordered as a presynthesized DNA fragment. For *ahpC* knockdown construct, 2 ng of the

gBlock (Integrated DNA Technologies [IDT], Coralville, IA) listed in Table S1 was combined with

25 ng of BamHI-digested, alkaline phosphatase-treated pBOMBL12CRia(e.v.)::L2 in a HiFi

reaction according to the manufacturer's instructions (NEB). The plasmid was transformed into

NEB 10-beta cells and verified by Sanger sequencing prior to transformation into *Chlamydia*

trachomatis. To generate the complementation strain, the *ahpC* gene was amplified using

primers listed in Table S1 and fused with the *ahpC* knockdown construct (i.e., pBOMBL12CRia

(*ahpC*)) digested with the Fast Digest restriction enzyme SalI (Thermo Fisher) and alkaline

phosphatase-treated, using the HiFi reaction as mentioned above.

Chlamydial transformation

Chlamydial transformations were performed using a protocol described previously, with some

modifications (Mueller et al., 2017). One day before transformation, 1×10^6 McCoy cells were

seeded in one 6-well plate, and two wells were used per plasmid transformation. Briefly, for

each well of the 6-well plate, 2 µg of sequenced verified plasmids were incubated with 2.5×10^6

C. trachomatis serovar L2 without plasmid (-pL2) EBs in 50 µL Tris-CaCl$_2$ (10 mM Tris, 50 mM

CaCl2, pH 7.4) at room temperature for 30 min. McCoy cells were washed with 2 mL Hank's

Balanced Salt Solution (HBSS; Gibco), and 1 mL HBSS was added back into each well. 1 mL of

HBSS was added to each transformant mixture, and one well of a 6-well plate was infected

using this transformation solution. Cells were centrifuged at 400 × g for 15 min at room

temperature followed by 15 min incubation at 37°C. HBSS was aspirated and replaced with

antibiotic-free DMEM. At 8 hpi, 1 μg/mL of cycloheximide and 1 or 2 U/mL of penicillin G or 500

μg/mL spectinomycin were added to the culture media. The infection was passaged every 48 h

until a population of penicillin, or spectinomycin resistant, green fluorescent protein (GFP)

positive *C. trachomatis* was established. The chlamydial transformants were then serially diluted

to isolate clonal populations. These isolated populations were further expanded and frozen at -

80°C in a sucrose phosphate solution (2SP). To verify plasmid sequences, DNA was harvested

from infected cultures using the DNeasy kit (Qiagen) and transformed into NEB 10-beta for

plasmid propagation. Isolated plasmids were then verified by restriction digest and Sanger

sequencing.

Inclusion forming unit assay

Inclusion forming unit assay was performed to determine the infectious progeny (number of

EBs) from a primary infection based on inclusions formed in a secondary infection. *C.*

trachomatis transformants were infected into HeLa cells and induced or not with 1 nM aTc at 10

hpi. At 24 and 48 hpi, samples were harvested by scraping three wells of a 24-well plate in 2

sucrose-phosphate (2SP) solution and lysed via a single freeze-thaw cycle, serially diluted, and

used to infect a fresh HeLa cell monolayer and allowed to grow for 24 h. Samples were fixed

with methanol, stained with a goat antibody specific to *C. trachomatis* major outer membrane

protein (MOMP; Meridian Biosciences, Memphis TN) followed by staining with donkey antigoat

Alexa Fluor 594-conjugated secondary antibody (Invitrogen) and titers were enumerated using a

20x lens objective. All experiments were performed three times for three biological replicates.

Induced values were expressed as a percentage of the uninduced values, which was

considered as 100%.

Immunofluorescence assay

HeLa cells were cultured on glass coverslips in 24-well tissue cultures plates at 2 x 10^5

cells/well, infected with the relevant strains, and, at 10 hpi, samples were induced or not with 1

nM aTc. At 24 hpi, samples were fixed and permeabilized using 100% methanol. Organisms

were stained with anti-MOMP (Meridian Biosciences) primary antibody for all the strains and

primary mouse anti-Cpf1 (dCas12) (Sigma-Millipore) for knockdown or complementation

samples. Donkey anti-goat Alexa Fluor 594-conjugated secondary antibody (Invitrogen) was

used to visualize *Chlamydia* in all the samples, and donkey anti-mouse Alexa Fluor 488-

conjugated secondary antibody (Invitrogen) was used for dCas12 expression. DAPI (Invitrogen)

was used for visualization of host and bacterial cell DNA. These stained coverslips were

mounted on glass slides using ProLong glass antifade mounting media (Invitrogen) and imaged

using a 100x lens objective on a Zeiss Axioimager Z.2 equipped with Apotome.2 optical

sectioning hardware and X-Cite Series 120PC illumination lamp using a 2MP Axiocam 506

monochrome camera.

Nucleic acid extraction and RT-qPCR

HeLa cells were seeded in 6-well tissue culture plates, infected with the *C. trachomatis*

transformants, and induced or not at 10 hpi with 1 nM aTc. For each condition, triplicate wells

were used for simultaneous harvest of RNA (for transcript analysis by RT-qPCR), gDNA (to

normalize RT-qPCR data and quantification of gDNA), and IFA (to verify the morphological

changes in samples in the tested conditions). For RNA extraction, cells were rinsed with DPBS

twice and lysed with 1 mL TRIzol (Invitrogen/Thermo Fisher) per well as per manufacturer's

instructions. 200 μL of chloroform was added to extract the aqueous layer containing total RNA,

which was precipitated with isopropanol. A total of 10 μg of purified RNA was treated with

TURBO DNase (Invitrogen/Thermo Fisher) according to the manufacturer's instructions to

remove DNA contamination. DNA-free RNA was used for cDNA synthesis using random

nonamers (N9; NEB) and SuperScript III reverse transcriptase (Invitrogen/Thermo Fisher)

following the manufacturer's instructions. cDNA samples were diluted 10-fold with molecular

biology-grade water and stored at -80°C. A total of 2.5 μL of each diluted cDNA sample was

used per well of a 96-well qPCR plate. For each of three biological replicates, each sample was

analyzed in triplicate on a QuantStudio 3 system (Applied Biosystems/Thermo Fisher) using the

standard amplification cycle with melting curve analysis. For gDNA, one well of a 6-well plate

per condition was scraped in 500 µL DPBS, split in half (i.e., 250 µL), and frozen at -80°C. Each

sample was then thawed and frozen twice more for a total of three freeze/thaw cycles, and

gDNA was extracted using the DNeasy DNA extraction kit (Qiagen) according to the

manufacturer's guidelines. The isolated gDNA was quantified and diluted down to 5 ng/mL prior

to use in quantitative PCR (qPCR). A total of 2.5 µL of each diluted gDNA sample was mixed

with 10 µL of PowerUp SYBR green master mix in a 96-well qPCR plate and was analyzed on a

QuantStudio 3 system (Applied Biosystems). Each sample from each biological replicate was

tested in triplicate. For each primer set used, a standard curve of gDNA was generated against

purified *C. trachomatis* L2 genomic DNA, and the cDNA levels were normalized to gDNA levels

or 16S rRNA for analysis. All experiments were performed three times for three biological

replicates. At the same time, morphological differences were monitored in the IFA control, with

samples fixed with methanol at the time of harvesting of RNA/gDNA. Staining and imaging were

performed as described above.

Enrichment, library preparation, and statistical analyses of RNA-sequencing samples

Three biological replicates were prepared and processed for RNA isolation as mentioned above.

Enrichment of microbial RNA was done using MICROB*Enrich* and MICROB*Express*

(Invitrogen/ThermoFisher) kits as per the manufacturer's instructions. Samples were aliquoted,

stored at -80°C and submitted to the UNMC Genomics Core. Before library preparation, quality

control (QC) was done using Nanodrop and Fragment Analyzer (Advanced Analytical). Final

libraries were quantified using Qubit DS DNA HS Assay reagents in Qubit Fluro meter (Life

Technologies), and the size of the libraries was measured via Fragment Analyzer. Individual

RNA-seq libraries were prepared using 400 ng of total RNA by Illumina Ribo-Zero plus

Microbiome (Illumina, Inc. San Diego, CA) and were multiplexed and subjected to 100 bp paired

read sequencing to generate >20 million pairs of reads per sample using Midoutput NextSeq

300 cycle kit by the UNMC Genomics Core facility (Table S2). The original fastq format reads

were trimmed by fqtrim tool (https://ccb.jhu.edu/software/fqtrim) to remove adapters, terminal

unknown bases (Ns), and low quality 3' regions (Phred score < 30). The trimmed fastq files

were processed by FastQC (Andrews, 2010). *Chlamydia trachomatis* 434/Bu bacterial reference

genome and annotation files were downloaded from Ensembl

(http://bacteria.ensembl.org/Chlamydia_trachomatis_434_bu/Info/Index). Sequencing data were

analyzed by the Bioinformatics and Systems Biology Core (BSBC). The false discovery rate

(FDR) and Bonferroni adjusted p values were provided to adjust for multiple-testing problem.

Fold changes were calculated from the GLM, which corrects for differences in library size

between the samples and the effects of confounding factors. The trimmed fastq files were

mapped to *Chlamydia trachomatis* 434/Bu by CLC Genomics Workbench 23 for RNA-seq

analyses.

Data availability. The raw and processed RNA sequencing reads in fastq format have been

deposited in Gene Expression Omnibus (GEO: www.ncbi.nlm.nih.gov/geo/) under Accession

number GSE278846.

Viability assay

Cell viability assays were performed using PrestoBlue (Invitrogen/Thermo Fisher). HeLa cells

were seeded in 96 well plates and were infected or not with the indicated chlamydial strains

(e.g., pBOMBDC.ev). Samples were treated or not as per the experimental parameters (e.g.,

different concentrations of oxidizing agents or scavengers). Medium without cells was used as

the blank control. At the end point, 10% PrestoBlue (v/v) was added in the wells and incubated

at 37°C for 30 min while protecting the plate from light. Fluorescence was read at excitation 560

nm and emission 590 nm using an Infinite M200 Pro (Tecan). The treated values were

expressed as a percentage of the untreated values, which was considered as 100%. All

experiments were performed three times for three biological replicates.

Oxidizing agents' susceptibility testing

Susceptibility of host cells and chlamydiae to different oxidizing agents was tested in HeLa cells

infected with density gradient-purified *Chlamydia trachomatis* 434/Bu EBs or *C. trachomatis*

transformants. For *C. trachomatis* transformants, expression of the construct was induced or not

with 1 nM aTc at 10 hpi. At 16 hpi, different concentrations of oxidizing agents were added, and

samples were incubated at 37°C for 30 min. These samples were washed three times with

HBSS, fresh DMEM media was added in the wells, and cultures were allowed to grow until 24

hpi. At 24 hpi, inclusion forming unit assays and immunofluorescence analysis were performed

as described above.

ROS scavenging by chemical compounds

The scavenging capacity of different ROS scavengers was tested using HeLa cells infected with

Chlamydia trachomatis 434/Bu EBs or the *ahpC* KD strain. To examine the protective effect of

scavengers in the absence or presence of oxidative stress, *Chlamydia trachomatis* 434/Bu EBs

infected HeLa cells were used. In these samples, scavengers were added or not at 10 hpi in

respective wells. At 16 hpi, wells were washed three times with HBSS, and fresh DMEM

containing 500 µM H_2O_2 or not was added. After 30 min, wells were washed three times with

HBSS and fresh DMEM containing scavengers or not was added back in the respective wells,

which were incubated until 24 hpi prior to collecting samples for IFU assay and IFA analysis.

The effect of these scavengers was further tested in *ahpC* KD. HeLa cells were infected with

ahpC KD, scavengers were added or not at 9.5 hpi before induction at 10 hpi with 1 nM aTc. At

14 hpi, scavengers were added again in the respective wells, and, at 24 hpi, samples were

collected for IFU assay and IFA.

ROS detection by CellROX Deep Red

Intracellular ROS levels were measured in uninfected and *ahpC* KD-infected HeLa cells using

CellROX Deep Red dye (Invitrogen/Thermo Fisher). Samples were induced or not at 10 hpi with

1 nM aTc. Medium without cells was used as the blank control. At the end point, media was

removed, samples were washed three times with DPBS and incubated with CellROX Deep Red

dye for 30 min in the dark at 37°C. Fluorescence was read at excitation 644 nm and emission

665 nm using an Infinite M200 Pro (Tecan). Microscopic images were captured using live cells

at 24 or 40 hpi. Experimental conditions were same as mentioned above. All experiments were

performed three times for three biological replicates.

ACKNOWLEDGMENTS

We thank Dr. H. Caldwell (NIH/NIAID) for providing eukaryotic cell lines and Dr. I. Clarke

(University of Southampton) for providing the plasmidless strain of *C. trachomatis* serovar L2.

We thank Dr. Elizabeth A. Rucks for critical feedback. We thank the members of the

Rucks/Ouellette research group for thoughtful discussion of the material presented. Funding for

this work was provided by the National Institutes of Health (NIH/NIAID) grants R01AI170688

and R21AI178150 to SPO. Authors thank the UNMC Genomics Core Facility and Bioinformatics

and Systems Biology Core (BSBC) at UNMC for RNA-seq services. The UNMC Genomics Core

Facility receives partial support from the National Institute for General Medical Science (NIGMS)

INBRE - P20GM103427-19, as well as the National Cancer Institute (NCI) and The Fred &

Pamela Buffett Cancer Center Support Grant- P30CA036727. The BSBC receives support from

Nebraska Research Initiative (NRI) and NIH (2P20GM103427, 5P30CA036727, and

2U54GM115458).

Figure legends:

Fig. 1 Alterations in redox status of key proteins regulate and drive chlamydial differentiation.

(A) Key characteristics of chlamydial developmental forms. (B) Hypothetical model for triggering

secondary differentiation through oxidative stress (black angled line). Increasing oxidation of

critical protein(s) may lead to earlier differentiation whereas maintaining a reducing environment

may delay differentiation. (C) Schematic representation of experimental model for triggering

secondary differentiation through altered activity of AhpC. *ahpC* knockdown may lead to earlier

differentiation while overexpression of *ahpC* may delay differentiation.

Fig. 2 Overexpression of *ahpC* affects chlamydial growth and differentiation. (A) Transcriptional

analysis of *ahpC* in *ahpC* overexpression (*ahpC*) and empty vector (EV) control using RT-qPCR

following induction at 10 hpi with 1 nM aTc. RNA and genomic DNA (gDNA) were harvested at

14 and 24 hpi and processed as mentioned in materials and methods. Data are presented as a

ratio of cDNA to gDNA plotted on a log scale. ***$p < 0.0001$ vs uninduced sample by using two-

way ANOVA. Data represent three biological replicates. (B) Immunofluorescence assay (IFA) of

ahpC and EV at 14 and 24 hpi. Construct expression was induced or not at 10 hpi with 1 nM

aTc, and samples were fixed with methanol at 24 hpi then stained for major outer membrane

protein (MOMP - red) and DAPI (blue) to label DNA. Scale bars = 2 μm. Images were captured

using a Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Representative images of

three biological replicates are shown. (C) Impact of *ahpC* overexpression on inclusion area.

Inclusion area of *ahpC* overexpression and EV strains was measured using ImageJ.

Experimental conditions were the same as mentioned in section (B). The area of 50 inclusions

was measured per condition for each sample. ***$p < 0.001$ vs uninduced sample by using

ordinary one-way ANOVA. Data were collected from three biological replicates. (D) IFU assay of

ahpC overexpression and empty vector control. Expression of the construct was induced or not

at 10 hpi, and samples were harvested at 24 or 48 hpi for reinfection and enumeration. IFUs

were calculated as the percentage of uninduced samples. **$p < 0.001$ vs uninduced sample by

using multiple paired t test. Data represent three biological replicates. (E) Quantification of

genomic DNA (gDNA) determined by qPCR in *ahpC* overexpression and empty vector control.

Construct expression was induced or not at 10 hpi with 1 nM aTc, gDNA was harvested at 14

and 24 hpi, and ng gDNA were plotted on a log scale. ***$p < 0.0001$ vs uninduced sample by

using two-way ANOVA. Data represent three biological replicates. (F) Ratio of log10 IFUs and

log10 gDNA. IFU/ml from (E) was normalized with gDNA from (D). **$p < 0.001$ vs uninduced

sample by using multiple unpaired t test. Data represent three biological replicates.

Fig. 3 Higher expression of *ahpC* provides resistance to peroxides in *Chlamydia.* IFA of *ahpC*

(A) or EV (C) exposed to oxidizing agents. Construct expression was induced or not at 10 hpi

with 1 nM aTc, and samples were treated with three different concentrations of hydrogen

peroxide (H_2O_2) at 16 hpi for 30 min then fixed with methanol at 24 hpi and stained and imaged

as described in the legend of Fig. 2B. Representative images from three biological replicates

are shown. IFU analysis of *ahpC* (B) or EV (D) following treatment with oxidizing agents, CHP-

Cumene hydroperoxide, H_2O_2-Hydrogen peroxide, TBHP-Tert-butyl hydroperoxide, and PN-

Peroxynitrite. Samples were processed as described for (A) and (C), and IFUs harvested at 24

hpi. IFUs of treated samples were compared with respective untreated controls. ***$p < 0.0001$ vs

untreated sample by using two-way ANOVA. Data represent three biological replicates.

Fig. 4 Reduced levels of AhpC negatively impact chlamydial growth. (A) Transcriptional analysis

of *ahpC* in knockdown (*ahpC* KD) and non-target (NT) control using RT-qPCR following

induction at 10 hpi with 1 nM aTc. RNA and gDNA were harvested at 14 and 24 hpi. Quantified

cDNA was normalized to gDNA, and values were plotted on a log scale. ***$p <$ 0.0001 vs

uninduced sample by using two-way ANOVA. Data represent three biological replicates. (B) IFA

was performed to assess inclusion size and morphology using the same induction conditions as

in section (A). At 24 hpi, cells were fixed with methanol and stained using primary antibodies to

major outer membrane protein (MOMP), Cpf1 (dCas12), and DAPI. All images were acquired on

Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Bars, 2 µm. Representative images

of three biological replicates are shown. (C) Quantification of genomic DNA (gDNA) determined

by qPCR in *ahpC* KD and NT strains. dCas12 expression was induced or not at 10 hpi, and

gDNA was harvested at 14 and 24 hpi and plotted on a log scale. ***$p <$ 0.0001 vs uninduced

sample by using two-way ANOVA. Data represent three biological replicates. (D) IFU titers

following induction at 10 hpi with 1 nM aTc. IFUs were counted from 24 and 48 hpi samples and

calculated as percentage of uninduced samples. ***$p <$ 0.0001, **$p <$ 0.001 vs uninduced sample

by using multiple paired t test. Data represent three biological replicates. (E) Ratio of log10 IFUs

by log10 gDNA. IFU/ml from (D) was normalized with gDNA from (C). **$p <$ 0.001 vs uninduced

sample by using multiple unpaired t test. Data represent three biological replicates.

Fig. 5 Intracellular ROS levels were measured to investigate the function of AhpC in reducing

ROS. (A) HeLa cells were infected or not with *ahpC* knockdown, and construct was induced or

not at 10 hpi with 1 nM aTc. At 24 or 48 hpi, samples were washed with DPBS and incubated

with CellROX Deep red dye for 30 min in dark. ROS levels were measured at wavelengths of

640 nm (excitation) and 665 nm (emission). **$p <$ 0.001, *$p <$ 0.01 vs uninduced sample by using

two-way ANOVA. Data represent three biological replicates. (B) Microscopy images were

acquired using live cells on Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Scale

Bars= 10 µm. Representative images of three biological replicates are shown.

Fig. 6 *Chlamydia* is hypersensitive to oxidizing agents in *ahpC* knockdown condition. IFA of

ahpC KD (A) or NT (C) treated with 62.5 µM H_2O_2. dCas12 expression was induced or not at 10

hpi with 1 nM aTc, treated or not with H_2O_2 at 16 hpi for 30 min, and allowed to grow until 24 hpi.

Coverslips were fixed with methanol at 24 hpi and stained major outer membrane protein

(MOMP), Cpf1 (dCas12), and DAPI. Scale bars = 2 μm. Images were captured using a Zeiss

Axio Imager Z.2 with Apotome2 at 100x magnification. Representative images from three

biological replicates are shown. IFU analysis of *ahpC* KD (B) or NT (D) following treatment with

oxidizing agents, CHP-Cumene hydroperoxide, H_2O_2-Hydrogen peroxide, TBHP-Tert-butyl

hydroperoxide, or PN-Peroxynitrite. dCas12 expression was induced or not, and samples were

treated or not as mentioned in the legend of Fig. 3B. IFUs of treated samples were calculated

as percentage of respective untreated samples. ***p< 0.0001 vs untreated sample by using two-

way ANOVA. Data represent three biological replicates.

Fig. 7 Complementation of the phenotypes observed in the *ahpC* knockdown. (A) Confirmation

of complementation (comp) of *ahpC* knockdown by RT-qPCR. Samples were processed and

quantified as mentioned previously in the legend of Fig. 4A. Values were plotted on a log scale.

*p< 0.01 vs uninduced sample using ordinary one-way ANOVA. Data represent three biological

replicates. (B) IFA of comp strain was performed at 24 hpi, and staining and imaging were

performed as mentioned in the legend of Fig. 4B. Scale bars, 2 μm. Representative images

from three biological replicates are shown. (C) Genomic DNA quantitation was performed by

qPCR. Construct expression was induced or not at 10 hpi, and gDNA was harvested at 14 and

24 hpi and plotted on a log scale. Statistical analysis was calculated using ordinary one-way

ANOVA. Data represent three biological replicates. (D) IFU analysis of comp strain. Statistical

analysis was calculated using multiple paired t test. Data represent three biological replicates.

(E) IFA of comp strain following treatment with 62.5 μM H_2O_2. Samples were treated, stained,

and images acquired as mentioned in the legend of Fig. 6A. Scale bars = 2 μm. Representative

images from three biological replicates are shown. (F) IFU analysis of comp strain following

treatment with oxidizing agents. Experiments were performed as mentioned in the legend of Fig.

6B. IFUs were calculated as percentage of respective untreated samples. ***p< 0.0001 vs

untreated sample by using two-way ANOVA. Data represent three biological replicates. (G)

ahpC knockdown growth defect rescued by ROS scavengers. IFU analysis of *ahpC* knockdown

treated with or without scavengers, α-Tocopherol (100 µM) and DMTU (10 mM), as mentioned

in materials and methods. IFUs were calculated as a percentage of the untreated, uninduced

sample. ***$p<$ 0.0001 vs untreated, uninduced or induced sample by using two-way ANOVA.

Data represent three biological replicates. (H) IFA of *ahpC* knockdown treated with or without

scavengers. Experimental conditions were similar as in section (G). Staining and imaging were

performed as mentioned in Fig. 6A. Representative images from three biological replicates are

shown.

Fig. 8 Effects of *ahpC* knockdown/overexpression on chlamydial developmental cycle

progression. (A) RT-qPCR analysis of late cycle genes (*hctA*, *hctB*, *omcB*, *tsp*, and *glgA*) in

ahpC knockdown. Experimental conditions were the same as mentioned in the legend of Fig.

4A. Quantified cDNA was normalized to gDNA, and values were plotted on a log scale. ***$p<$

0.0001, **$p<$ 0.001, *$p<$ 0.01 vs uninduced sample by using two-way ANOVA. Data represent

three biological replicates. (B) Volcano plot of RNA-sequencing of *ahpC* knockdown.

Experimental conditions were the same as mentioned in the legend of Fig. 4A. The volcano plot

was prepared using GraphPad Prism software. The vertical dashed lines indicate a fold change

of 1.7 as compared to the respective uninduced control. The horizontal dashed line indicates

an p value of 0.05. The black dots represent genes not significantly different and grey dots

represent significantly altered but fold change lesser than 1.5. Blue and red spots represent

statistically significant altered genes with more than 1.5-fold change in transcription levels

between uninduced and induced samples. Red dots indicate ahpC and canonical late genes

also shown in Table 1. (C) One-step growth curve of *ahpC* knockdown. Samples were induced

or not with 1 nM aTc at 10 hpi and harvested at 16, 18, 20, 22, and 24 hpi. IFUs recovered are

displayed as log10 values. ***$p<$ 0.0001 vs uninduced sample by using two-way ANOVA. Data

represent three biological replicates.

Fig. 9 Altering the activity of AhpC in *Ctr* L2 impacts its developmental cycle progression. (Top)

In *Chlamydia*, secondary differentiation is asynchronous and RBs divide through an asymmetric

budding mechanism. In such conditions, either the mother or daughter cell may inherit more

oxidized proteins (represented by a darker shade), which can then impact whether a given RB

will divide again or undergo secondary differentiation. (Bottom) The black dots represent EBs,

the orange circles show RBs. The developmental cycle of wild-type *C. trachomatis* (*Ctr* L2) is

shown. In *ahpC* KD, highly oxidized conditions lead some RBs to cross the oxidative threshold

sooner, allowing activation of late genes and secondary differentiation earlier than other RBs. In

ahpC overexpression, a reducing environment results in a delay in achieving the oxidative

threshold, thus allowing RBs to continue to divide before committing to secondary

differentiation.

Table 1: List of canonical late genes significantly increased during *ahpC* knockdown at 14 hpi.

CT #	CTL #	Name	Function	Fold Change	Ref
CT046	CTL0302	*hct2 (hctB)*	Histone H1-like protein HC2	9.10	(Brickman et al., 1993)
CT578	CTL0841	*copB*	Needle tip; translocator	6.44	(Ouellette et al., 2005)
CT576	CTL0839	*scc2 (lcrH_1)*	Type III secretion chaperone (Low calcium response protein H)	5.52	(Ouellette et al., 2005)
CT579	CTL0842	*copD*	Needle tip; translocator	5.41	(Ouellette et al., 2005)
CT441	CTL0700	*tsp*	Carboxy-terminal processing protease	3.76	(Swoboda et al., 2023)
CT444	CTL0703	*omcA*	Small cysteine-rich outer membrane protein	3.63	(Hatch, 1996)

CT456	CTL0716	*tarP*	Translocated actin-recruiting phosphoprotein	3.22	(Clifton et al., 2004)
CT443	CTL0702	*omcB*	Large cysteine-rich periplasmic protein	2.94	(Newhall, 1987)
CT798	CTL0167	*glgA*	Glycogen synthase	2.26	(Belland et al., 2003; Gehre et al., 2016)
CT080	CTL0336	*ltuB*	Late transcription unit B protein	2.11	(Belland et al., 2003)
CT177	CTL0429	*dsbA*	Disulfide bond chaperone	1.91	(Christensen et al., 2019)
CT743	CTL0112	*hctA*	Histone H1-like protein HC1	1.81	(Barry et al., 1993)

Supplementary Information

Table S1 List of plasmids, strains, and primers used in this study.

Table S2 List of differentially expressed genes in *ahpC* knockdown RNA-seq

Fig. S1 Viability assay of uninfected or infected HeLa cells treated with oxidizing agents H_2O_2

(A), CHP (B), TBHP (C), and PN (D). HeLa cells were infected or not with empty vector control,

pBOMBDC.ev (EV), and treated or not with different concentrations of oxidizing agents at 16 hpi

for 30 min. At 24 hpi, an end point viability assay was performed using PrestoBlue as mentioned

in materials and methods. The treated values were expressed as a percentage of the untreated

values, which were considered as 100%. Data represent three biological replicates.

Fig. S2 Response of *C. trachomatis* L2 against oxidizing agents. (A) IFA of wild-type *Ctr* L2

exposed to oxidizing agents. Oxidizing agents' treatment, staining, and imaging were performed

as mentioned in materials and methods. Representative images from three biological replicates

are shown. (B) IFU analysis of *Ctr* L2 post-exposure with oxidizing agents. Conditions were

same as in section (A), and IFUs were harvested at 24 hpi. IFUs were calculated as percentage

of untreated samples. ***$p<$ 0.0001 vs untreated sample by using one way ANOVA. Data

represent three biological replicates.

Fig. S3 Overexpression of *ahpC* provides resistance to peroxides in *Chlamydia.* IFA of *ahpC* (A)

or EV (B) exposed to oxidizing agents (CHP, TBHP, and PN). Experiments were performed as

mentioned in the legend of Fig. 3A. Representative images from three biological replicates are

shown.

Fig. S4 Intracellular ROS levels in *ahpC* knockdown at 40 hpi. HeLa cells were infected or not

with *ahpC* knockdown, and construct was induced or not at 10 hpi with 1 nM aTc. At 40 hpi,

samples were washed with DPBS and incubated with CellROX Deep red dye for 30 min in dark.

Microscopic images of live cells on Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification

were captured. Scale Bars=10 µm. Representative images of three biological replicates are

shown.

Fig. S5 *Chlamydia* is hypersensitive to oxidizing agents as a result of reduced levels of *ahpC*.

IFA of *ahpC* KD (A), NT (B), comp (C) treated with CHP, TBHP, and PN. Experimental

conditions were the same as mentioned in the legend of Fig. 6A. Representative images from

three biological replicates are shown.

Fig. S6 Rescue of oxidative stress phenotype by ROS scavengers in *C. trachomatis* L2. (A) IFA

of wild-type *Ctr* L2 incubated or not with scavengers, α-Tocopherol (100 µM) and DMTU (10

mM), and treated or not with 500 µM H_2O_2 as mentioned in materials and methods.

Representative images from three biological replicates are shown. (B) IFU analysis of *Ctr* L2

grown under the same conditions as mentioned in the legend of Fig. 7G. IFUs were calculated

as percentage of untreated samples. ***$p<$ 0.0001, statistical analysis was performed using

ordinary one-way ANOVA. H_2O_2 treated sample and samples incubated with scavengers only

were compared to the untreated control. Samples treated with H_2O_2 and incubated with

scavengers were compared with sample treated with H_2O_2. Data represent three biological

replicates. Only significant differences are noted.

Fig. S7 Effects of *ahpC* knockdown/overexpression on chlamydial developmental cycle

progression. RT-qPCR analysis of late cycle genes (*hctA*, *hctB*, *omcB*, *tsp*, and *glgA*) in (A)

complementation, and (B) *ahpC* overexpression strain. Experimental conditions were the same

as mentioned in the legend of Fig. 7A, and 2A, respectively. Quantified cDNA was normalized to

gDNA, and values were plotted on a log scale. ***$p< 0.0001$, **$p< 0.001$, *$p< 0.01$ vs uninduced

sample by using two-way ANOVA. Data represent three biological replicates.

Fig. S8 Effect of penicillin treatment during *ahpC* knockdown. (A) IFA was performed to assess

inclusion size and morphology of *ahpC* KD-spec following induction (1 nM aTc) and penicillin

treatment (1 U/mL) at 10 hpi. At 24 hpi, cells were fixed with methanol and stained using primary

antibodies to major outer membrane protein (MOMP), Cpf1 (dCas12), and DAPI. All images

were acquired on Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Bars, 2 μm.

Representative images of three biological replicates are shown. Transcriptional analysis of (B)

ahpC, (C) *hctA*, and (D) *hctB* in *ahpC* KD-spec using RT-qPCR using the same conditions as in

section (A). RNA samples were harvested at 16 and 24 hpi. Quantified cDNA was normalized to

16SrRNA, and values were plotted on a log scale. ***$p< 0.0001$, **$p< 0.001$, *$p< 0.01$ vs

uninduced sample by using two-way ANOVA. Data represent three biological replicates.

REFERENCES

Abdelrahman, Y., Ouellette, S. P., Belland, R. J., & Cox, J. V. (2016). Polarized Cell Division of Chlamydia trachomatis. *PLoS Pathog*, *12*(8), e1005822. https://doi.org/10.1371/journal.ppat.1005822

Abdelrahman, Y. M., & Belland, R. J. (2005). The chlamydial developmental cycle. *FEMS Microbiol Rev*, *29*(5), 949-959. https://doi.org/10.1016/j.femsre.2005.03.002

Abdul-Sater, A. A., Said-Sadier, N., Lam, V. M., Singh, B., Pettengill, M. A., Soares, F., Tattoli, I., Lipinski, S., Girardin, S. E., Rosenstiel, P., & Ojcius, D. M. (2010). Enhancement of reactive oxygen species production and chlamydial infection by the mitochondrial Nod-like family member NLRX1. *J Biol Chem*, *285*(53), 41637-41645. https://doi.org/10.1074/jbc.M110.137885

Andrews, S. (2010). A quality control tool for high throughput sequence data. In. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/: Babraham Institute.

Antelmann, H., Engelmann, S., Schmid, R., & Hecker, M. (1996). General and oxidative stress responses in Bacillus subtilis: cloning, expression, and mutation of the alkyl hydroperoxide reductase operon. *J Bacteriol*, *178*(22), 6571-6578. https://doi.org/10.1128/jb.178.22.6571-6578.1996

Barbour, A. G., Amano, K., Hackstadt, T., Perry, L., & Caldwell, H. D. (1982). Chlamydia trachomatis has penicillin-binding proteins but not detectable muramic acid. *J Bacteriol*, *151*(1), 420-428. https://doi.org/10.1128/jb.151.1.420-428.1982

Barry, C. E., 3rd, Brickman, T. J., & Hackstadt, T. (1993). Hc1-mediated effects on DNA structure: a potential regulator of chlamydial development. *Mol Microbiol*, *9*(2), 273-283. https://doi.org/10.1111/j.1365-2958.1993.tb01689.x

Belland, R. J., Zhong, G., Crane, D. D., Hogan, D., Sturdevant, D., Sharma, J., Beatty, W. L., & Caldwell, H. D. (2003). Genomic transcriptional profiling of the developmental cycle of Chlamydia trachomatis. *Proc Natl Acad Sci U S A*, *100*(14), 8478-8483. https://doi.org/10.1073/pnas.1331135100

Betts-Hampikian, H. J., & Fields, K. A. (2011). Disulfide bonding within components of the Chlamydia type III secretion apparatus correlates with development. *J Bacteriol*, *193*(24), 6950-6959. https://doi.org/10.1128/JB.05163-11

Boncompain, G., Muller, C., Meas-Yedid, V., Schmitt-Kopplin, P., Lazarow, P. B., & Subtil, A. (2014). The intracellular bacteria Chlamydia hijack peroxisomes and utilize their enzymatic capacity to produce bacteria-specific phospholipids. *PLoS One*, *9*(1), e86196. https://doi.org/10.1371/journal.pone.0086196

Boncompain, G., Schneider, B., Delevoye, C., Kellermann, O., Dautry-Varsat, A., & Subtil, A. (2010). Production of reactive oxygen species is turned on and rapidly shut down in epithelial cells infected with Chlamydia trachomatis. *Infect Immun*, *78*(1), 80-87. https://doi.org/10.1128/IAI.00725-09

Brickman, T. J., Barry, C. E., 3rd, & Hackstadt, T. (1993). Molecular cloning and expression of hctB encoding a strain-variant chlamydial histone-like protein with DNA-binding activity. *J Bacteriol*, *175*(14), 4274-4281. https://doi.org/10.1128/jb.175.14.4274-4281.1993

Brinkworth, A. J., Wildung, M. R., & Carabeo, R. A. (2018). Genomewide Transcriptional Responses of Iron-Starved Chlamydia trachomatis Reveal Prioritization of Metabolic Precursor Synthesis over Protein Translation. *mSystems*, *3*(1). https://doi.org/10.1128/mSystems.00184-17

Caldwell, H. D., Kromhout, J., & Schachter, J. (1981). Purification and partial characterization of the major outer membrane protein of Chlamydia trachomatis. *Infect Immun*, *31*(3), 1161-1176. https://doi.org/10.1128/iai.31.3.1161-1176.1981

Chiarelli, T. J., Grieshaber, N. A., Omsland, A., Remien, C. H., & Grieshaber, S. S. (2020). Single-Inclusion Kinetics of Chlamydia trachomatis Development. *mSystems*, *5*(5). https://doi.org/10.1128/mSystems.00689-20

Christensen, S., Halili, M. A., Strange, N., Petit, G. A., Huston, W. M., Martin, J. L., & McMahon, R. M. (2019). Oxidoreductase disulfide bond proteins DsbA and DsbB form an active redox pair in Chlamydia trachomatis, a bacterium with disulfide dependent infection and development. *PLoS One*, *14*(9), e0222595. https://doi.org/10.1371/journal.pone.0222595

Clifton, D. R., Dooley, C. A., Grieshaber, S. S., Carabeo, R. A., Fields, K. A., & Hackstadt, T. (2005). Tyrosine phosphorylation of the chlamydial effector protein Tarp is species specific and not required for recruitment of actin. *Infect Immun*, *73*(7), 3860-3868. https://doi.org/10.1128/IAI.73.7.3860-3868.2005

Clifton, D. R., Fields, K. A., Grieshaber, S. S., Dooley, C. A., Fischer, E. R., Mead, D. J., Carabeo, R. A., & Hackstadt, T. (2004). A chlamydial type III translocated protein is tyrosine-phosphorylated at the site of entry and associated with recruitment of actin. *Proc Natl Acad Sci U S A*, *101*(27), 10166-10171. https://doi.org/10.1073/pnas.0402829101

Cosgrove, K., Coutts, G., Jonsson, I. M., Tarkowski, A., Kokai-Kun, J. F., Mond, J. J., & Foster, S. J. (2007). Catalase (KatA) and alkyl hydroperoxide reductase (AhpC) have compensatory roles in peroxide stress resistance and are required for survival, persistence, and nasal colonization in Staphylococcus aureus. *J Bacteriol*, *189*(3), 1025-1035. https://doi.org/10.1128/JB.01524-06

Cox, J. V., Abdelrahman, Y. M., & Ouellette, S. P. (2020). Penicillin-binding proteins regulate multiple steps in the polarized cell division process of Chlamydia. *Sci Rep*, *10*(1), 12588. https://doi.org/10.1038/s41598-020-69397-x

de Oliveira, M. A., Tairum, C. A., Netto, L. E. S., de Oliveira, A. L. P., Aleixo-Silva, R. L., Cabrera, V. I. M., Breyer, C. A., & Dos Santos, M. C. (2021). Relevance of peroxiredoxins in pathogenic microorganisms. *Appl Microbiol Biotechnol*, *105*(14-15), 5701-5717. https://doi.org/10.1007/s00253-021-11360-5

Dip, P. V., Kamariah, N., Nartey, W., Beushausen, C., Kostyuchenko, V. A., Ng, T. S., Lok, S. M., Saw, W. G., Eisenhaber, F., Eisenhaber, B., & Gruber, G. (2014). Key roles of the Escherichia coli AhpC C-terminus in assembly and catalysis of alkylhydroperoxide reductase, an enzyme essential for the alleviation of oxidative stress. *Biochim Biophys Acta*, *1837*(12), 1932-1943. https://doi.org/10.1016/j.bbabio.2014.08.007

Douglas, A. L., & Hatch, T. P. (2000). Expression of the transcripts of the sigma factors and putative sigma factor regulators of Chlamydia trachomatis L2. *Gene*, *247*(1-2), 209-214. https://doi.org/10.1016/s0378-1119(00)00094-9

Everett, K. D., & Hatch, T. P. (1995). Architecture of the cell envelope of Chlamydia psittaci 6BC. *J Bacteriol*, *177*(4), 877-882. https://doi.org/10.1128/jb.177.4.877-882.1995

Ezraty, B., Gennaris, A., Barras, F., & Collet, J. F. (2017). Oxidative stress, protein damage and repair in bacteria. *Nat Rev Microbiol*, *15*(7), 385-396. https://doi.org/10.1038/nrmicro.2017.26

Fang, F. C. (2011). Antimicrobial actions of reactive oxygen species. *mBio*, *2*(5). https://doi.org/10.1128/mBio.00141-11

Feng, X., Guo, K., & Gao, H. (2020). Plasticity of the peroxidase AhpC links multiple substrates to diverse disulfide-reducing pathways in Shewanella oneidensis. *J Biol Chem*, *295*(32), 11118-11130. https://doi.org/10.1074/jbc.RA120.014010

Gehre, L., Gorgette, O., Perrinet, S., Prevost, M. C., Ducatez, M., Giebel, A. M., Nelson, D. E., Ball, S. G., & Subtil, A. (2016). Sequestration of host metabolism by an intracellular pathogen. *Elife*, *5*, e12552. https://doi.org/10.7554/eLife.12552

Gerard, H. C., Freise, J., Wang, Z., Roberts, G., Rudy, D., Krauss-Opatz, B., Kohler, L., Zeidler, H., Schumacher, H. R., Whittum-Hudson, J. A., & Hudson, A. P. (2002). Chlamydia trachomatis genes whose products are related to energy metabolism are expressed differentially in active vs. persistent infection. *Microbes Infect*, *4*(1), 13-22. https://doi.org/10.1016/s1286-4579(01)01504-0

Hartl, J., Kiefer, P., Kaczmarczyk, A., Mittelviefhaus, M., Meyer, F., Vonderach, T., Hattendorf, B., Jenal, U., & Vorholt, J. A. (2020). Untargeted metabolomics links glutathione to bacterial cell cycle progression. *Nat Metab*, *2*(2), 153-166. https://doi.org/10.1038/s42255-019-0166-0

Hatch, N. D., & Ouellette, S. P. (2023). Identification of the alternative sigma factor regulons of Chlamydia trachomatis using multiplexed CRISPR interference. *mSphere*, *8*(5), e0039123. https://doi.org/10.1128/msphere.00391-23

Hatch, T. P. (1996). Disulfide cross-linked envelope proteins: the functional equivalent of peptidoglycan in chlamydiae? *J Bacteriol*, *178*(1), 1-5. https://doi.org/10.1128/jb.178.1.1-5.1996

Hollander-Czytko, H., Grabowski, J., Sandorf, I., Weckermann, K., & Weiler, E. W. (2005). Tocopherol content and activities of tyrosine aminotransferase and cystine lyase in Arabidopsis under stress conditions. *J Plant Physiol*, *162*(7), 767-770. https://doi.org/10.1016/j.jplph.2005.04.019

Iliffe-Lee, E. R., & McClarty, G. (1999). Glucose metabolism in Chlamydia trachomatis: the 'energy parasite' hypothesis revisited. *Mol Microbiol*, *33*(1), 177-187. https://doi.org/10.1046/j.1365-2958.1999.01464.x

Kiffin, R., Bandyopadhyay, U., & Cuervo, A. M. (2006). Oxidative stress and autophagy. *Antioxid Redox Signal*, *8*(1-2), 152-162. https://doi.org/10.1089/ars.2006.8.152

Kimura, A., Yuhara, S., Ohtsubo, Y., Nagata, Y., & Tsuda, M. (2012). Suppression of pleiotropic phenotypes of a Burkholderia multivorans fur mutant by oxyR mutation. *Microbiology (Reading)*, *158*(Pt 5), 1284-1293. https://doi.org/10.1099/mic.0.057372-0

Koshkin, A., Knudsen, G. M., & Ortiz De Montellano, P. R. (2004). Intermolecular interactions in the AhpC/AhpD antioxidant defense system of Mycobacterium tuberculosis. *Arch Biochem Biophys*, *427*(1), 41-47. https://doi.org/10.1016/j.abb.2004.04.017

Lee, J., Cox, J. V., & Ouellette, S. P. (2020). Critical Role for the Extended N Terminus of Chlamydial MreB in Directing Its Membrane Association and Potential Interaction with Divisome Proteins. *J Bacteriol*, *202*(9). https://doi.org/10.1128/JB.00034-20

Loprasert, S., Sallabhan, R., Whangsuk, W., & Mongkolsuk, S. (2003). Compensatory increase in ahpC gene expression and its role in protecting Burkholderia pseudomallei against reactive nitrogen intermediates. *Arch Microbiol*, *180*(6), 498-502. https://doi.org/10.1007/s00203-003-0621-9

Mastronicola, D., Falabella, M., Testa, F., Pucillo, L. P., Teixeira, M., Sarti, P., Saraiva, L. M., &
Giuffre, A. (2014). Functional characterization of peroxiredoxins from the human
protozoan parasite Giardia intestinalis. *PLoS Negl Trop Dis*, *8*(1), e2631.
https://doi.org/10.1371/journal.pntd.0002631

Matsumoto, A., & Manire, G. P. (1970). Electron Microscopic Observations on the Fine Structure
of Cell Walls of Chlamydia psittaci. *J Bacteriol*, *104*(3), 1332-1337.
https://doi.org/10.1128/jb.104.3.1332-1337.1970

Mishra, S., & Imlay, J. (2012). Why do bacteria use so many enzymes to scavenge hydrogen
peroxide? *Arch Biochem Biophys*, *525*(2), 145-160.
https://doi.org/10.1016/j.abb.2012.04.014

Mongkolsuk, S., Whangsuk, W., Vattanaviboon, P., Loprasert, S., & Fuangthong, M. (2000). A
Xanthomonas alkyl hydroperoxide reductase subunit C (ahpC) mutant showed an
altered peroxide stress response and complex regulation of the compensatory response
of peroxide detoxification enzymes. *J Bacteriol*, *182*(23), 6845-6849.
https://doi.org/10.1128/JB.182.23.6845-6849.2000

Moulder, J. W. (1993). Why is Chlamydia sensitive to penicillin in the absence of peptidoglycan?
Infect Agents Dis, *2*(2), 87-99. https://www.ncbi.nlm.nih.gov/pubmed/8162358

Moulder, J. W., Colon, J. I., Ruda, J., & Zebovitz, M. M. (1956). The effect of penicillin on
multiplying and nonmultiplying populations of sensitive and resistant strains of feline
pneumonitis virus. *J Infect Dis*, *98*(3), 229-238. https://doi.org/10.1093/infdis/98.3.229

Mueller, K. E., Wolf, K., & Fields, K. A. (2017). Chlamydia trachomatis Transformation and Allelic
Exchange Mutagenesis. *Curr Protoc Microbiol*, *45*, 11A 13 11-11A 13 15.
https://doi.org/10.1002/cpmc.31

Narayanan, S., Janakiraman, B., Kumar, L., & Radhakrishnan, S. K. (2015). A cell cycle-
controlled redox switch regulates the topoisomerase IV activity. *Genes Dev*, *29*(11),
1175-1187. https://doi.org/10.1101/gad.257030.114

Newhall, W. J. t. (1987). Biosynthesis and disulfide cross-linking of outer membrane
components during the growth cycle of Chlamydia trachomatis. *Infect Immun*, *55*(1),
162-168. https://doi.org/10.1128/iai.55.1.162-168.1987

Ochsner, U. A., Vasil, M. L., Alsabbagh, E., Parvatiyar, K., & Hassett, D. J. (2000). Role of the
Pseudomonas aeruginosa oxyR-recG operon in oxidative stress defense and DNA
repair: OxyR-dependent regulation of katB-ankB, ahpB, and ahpC-ahpF. *J Bacteriol*,
182(16), 4533-4544. https://doi.org/10.1128/JB.182.16.4533-4544.2000

Oh, E., & Jeon, B. (2014). Role of alkyl hydroperoxide reductase (AhpC) in the biofilm formation
of Campylobacter jejuni. *PLoS One*, *9*(1), e87312.
https://doi.org/10.1371/journal.pone.0087312

Ouellette, S. P., Abdelrahman, Y. M., Belland, R. J., & Byrne, G. I. (2005). The Chlamydia
pneumoniae type III secretion-related lcrH gene clusters are developmentally expressed
operons. *J Bacteriol*, *187*(22), 7853-7856. https://doi.org/10.1128/JB.187.22.7853-
7856.2005

Ouellette, S. P., Blay, E. A., Hatch, N. D., & Fisher-Marvin, L. A. (2021). CRISPR Interference To
Inducibly Repress Gene Expression in Chlamydia trachomatis. *Infect Immun*, *89*(7),
e0010821. https://doi.org/10.1128/IAI.00108-21

Ouellette, S. P., Fisher-Marvin, L. A., Harpring, M., Lee, J., Rucks, E. A., & Cox, J. V. (2022).
Localized cardiolipin synthesis is required for the assembly of MreB during the polarized
cell division of Chlamydia trachomatis. *PLoS Pathog*, *18*(9), e1010836.
https://doi.org/10.1371/journal.ppat.1010836

Ouellette, S. P., Hatch, T. P., AbdelRahman, Y. M., Rose, L. A., Belland, R. J., & Byrne, G. I.
(2006). Global transcriptional upregulation in the absence of increased translation in
Chlamydia during IFNgamma-mediated host cell tryptophan starvation. *Mol Microbiol*,
62(5), 1387-1401. https://doi.org/10.1111/j.1365-2958.2006.05465.x

Ouellette, S. P., Karimova, G., Subtil, A., & Ladant, D. (2012). Chlamydia co-opts the rod shape-
determining proteins MreB and Pbp2 for cell division. *Mol Microbiol*, *85*(1), 164-178.
https://doi.org/10.1111/j.1365-2958.2012.08100.x
Ouellette, S. P., Lee, J., & Cox, J. V. (2020). Division without Binary Fission: Cell Division in the
FtsZ-Less Chlamydia. *J Bacteriol*, *202*(17). https://doi.org/10.1128/JB.00252-20
Pan, S., Jensen, A. A., Wood, N. A., Henrichfreise, B., Brotz-Oesterhelt, H., Fisher, D. J., Sass,
P., & Ouellette, S. P. (2023). Molecular Characterization of the ClpC AAA+ ATPase in the
Biology of Chlamydia trachomatis. *mBio*, *14*(2), e0007523.
https://doi.org/10.1128/mbio.00075-23
Panzetta, M. E., Valdivia, R. H., & Saka, H. A. (2018). Chlamydia Persistence: A Survival
Strategy to Evade Antimicrobial Effects in-vitro and in-vivo. *Front Microbiol*, *9*, 3101.
https://doi.org/10.3389/fmicb.2018.03101
Parsonage, D., Karplus, P. A., & Poole, L. B. (2008). Substrate specificity and redox potential of
AhpC, a bacterial peroxiredoxin. *Proc Natl Acad Sci U S A*, *105*(24), 8209-8214.
https://doi.org/10.1073/pnas.0708308105
Pokorzynski, N. D., Brinkworth, A. J., & Carabeo, R. (2019). A bipartite iron-dependent
transcriptional regulation of the tryptophan salvage pathway in Chlamydia trachomatis.
Elife, *8*. https://doi.org/10.7554/eLife.42295
Poole, L. B., & Ellis, H. R. (1996). Flavin-dependent alkyl hydroperoxide reductase from
Salmonella typhimurium. 1. Purification and enzymatic activities of overexpressed AhpF
and AhpC proteins. *Biochemistry*, *35*(1), 56-64. https://doi.org/10.1021/bi951887s
Reuter, J., Otten, C., Jacquier, N., Lee, J., Mengin-Lecreulx, D., Lockener, I., Kluj, R., Mayer, C.,
Corona, F., Dannenberg, J., Aeby, S., Buhl, H., Greub, G., Vollmer, W., Ouellette, S. P.,
Schneider, T., & Henrichfreise, B. (2023). An NlpC/P60 protein catalyzes a key step in
peptidoglycan recycling at the intersection of energy recovery, cell division and immune
evasion in the intracellular pathogen Chlamydia trachomatis. *PLoS Pathog*, *19*(2),
e1011047. https://doi.org/10.1371/journal.ppat.1011047
Richard, D., Bartfai, R., Volz, J., Ralph, S. A., Muller, S., Stunnenberg, H. G., & Cowman, A. F.
(2011). A genome-wide chromatin-associated nuclear peroxiredoxin from the malaria
parasite Plasmodium falciparum. *J Biol Chem*, *286*(13), 11746-11755.
https://doi.org/10.1074/jbc.M110.198499
Rusconi, B., & Greub, G. (2013). Discovery of catalases in members of the Chlamydiales order.
J Bacteriol, *195*(16), 3543-3551. https://doi.org/10.1128/JB.00563-13
Saka, H. A., Thompson, J. W., Chen, Y. S., Kumar, Y., Dubois, L. G., Moseley, M. A., & Valdivia,
R. H. (2011). Quantitative proteomics reveals metabolic and pathogenic properties of
Chlamydia trachomatis developmental forms. *Mol Microbiol*, *82*(5), 1185-1203.
https://doi.org/10.1111/j.1365-2958.2011.07877.x
Seaver, L. C., & Imlay, J. A. (2001). Alkyl hydroperoxide reductase is the primary scavenger of
endogenous hydrogen peroxide in Escherichia coli. *J Bacteriol*, *183*(24), 7173-7181.
https://doi.org/10.1128/JB.183.24.7173-7181.2001
Sherman, D. R., Mdluli, K., Hickey, M. J., Arain, T. M., Morris, S. L., Barry, C. E., 3rd, & Stover,
C. K. (1996). Compensatory ahpC gene expression in isoniazid-resistant Mycobacterium
tuberculosis. *Science*, *272*(5268), 1641-1643.
https://doi.org/10.1126/science.272.5268.1641
Skipp, P., Robinson, J., O'Connor, C. D., & Clarke, I. N. (2005). Shotgun proteomic analysis of
Chlamydia trachomatis. *Proteomics*, *5*(6), 1558-1573.
https://doi.org/10.1002/pmic.200401044
Skipp, P. J., Hughes, C., McKenna, T., Edwards, R., Langridge, J., Thomson, N. R., & Clarke, I.
N. (2016). Quantitative Proteomics of the Infectious and Replicative Forms of Chlamydia
trachomatis. *PLoS One*, *11*(2), e0149011. https://doi.org/10.1371/journal.pone.0149011

Soules, K. R., Dmitriev, A., LaBrie, S. D., Dimond, Z. E., May, B. H., Johnson, D. K., Zhang, Y., Battaile, K. P., Lovell, S., & Hefty, P. S. (2020). Structural and ligand binding analyses of the periplasmic sensor domain of RsbU in Chlamydia trachomatis support a role in TCA cycle regulation. *Mol Microbiol*, *113*(1), 68-88. https://doi.org/10.1111/mmi.14401

Staerck, C., Gastebois, A., Vandeputte, P., Calenda, A., Larcher, G., Gillmann, L., Papon, N., Bouchara, J. P., & Fleury, M. J. J. (2017). Microbial antioxidant defense enzymes. *Microb Pathog*, *110*, 56-65. https://doi.org/10.1016/j.micpath.2017.06.015

Storz, G., Jacobson, F. S., Tartaglia, L. A., Morgan, R. W., Silveira, L. A., & Ames, B. N. (1989). An alkyl hydroperoxide reductase induced by oxidative stress in Salmonella typhimurium and Escherichia coli: genetic characterization and cloning of ahp. *J Bacteriol*, *171*(4), 2049-2055. https://doi.org/10.1128/jb.171.4.2049-2055.1989

Swoboda, A. R., Wood, N. A., Saery, E. A., Fisher, D. J., & Ouellette, S. P. (2023). The Periplasmic Tail-Specific Protease, Tsp, Is Essential for Secondary Differentiation in Chlamydia trachomatis. *J Bacteriol*, *205*(5), e0009923. https://doi.org/10.1128/jb.00099-23

Thompson, C. C., Griffiths, C., Nicod, S. S., Lowden, N. M., Wigneshweraraj, S., Fisher, D. J., & McClure, M. O. (2015). The Rsb Phosphoregulatory Network Controls Availability of the Primary Sigma Factor in Chlamydia trachomatis and Influences the Kinetics of Growth and Development. *PLoS Pathog*, *11*(8), e1005125. https://doi.org/10.1371/journal.ppat.1005125

Walch, M., Dotiwala, F., Mulik, S., Thiery, J., Kirchhausen, T., Clayberger, C., Krensky, A. M., Martinvalet, D., & Lieberman, J. (2015). Cytotoxic Cells Kill Intracellular Bacteria through Granulysin-Mediated Delivery of Granzymes. *Cell*, *161*(5), 1229. https://doi.org/10.1016/j.cell.2015.05.021

Wan, F., Feng, X., Yin, J., & Gao, H. (2021). Distinct H(2)O(2)-Scavenging System in Yersinia pseudotuberculosis: KatG and AhpC Act Together to Scavenge Endogenous Hydrogen Peroxide. *Front Microbiol*, *12*, 626874. https://doi.org/10.3389/fmicb.2021.626874

Wang, G., Conover, R. C., Benoit, S., Olczak, A. A., Olson, J. W., Johnson, M. K., & Maier, R. J. (2004). Role of a bacterial organic hydroperoxide detoxification system in preventing catalase inactivation. *J Biol Chem*, *279*(50), 51908-51914. https://doi.org/10.1074/jbc.M408450200

Wang, X., Schwarzer, C., Hybiske, K., Machen, T. E., & Stephens, R. S. (2014). Developmental stage oxidoreductive states of Chlamydia and infected host cells. *mBio*, *5*(6), e01924. https://doi.org/10.1128/mBio.01924-14

Wang, Y., Kahane, S., Cutcliffe, L. T., Skilton, R. J., Lambden, P. R., & Clarke, I. N. (2011). Development of a transformation system for Chlamydia trachomatis: restoration of glycogen biosynthesis by acquisition of a plasmid shuttle vector. *PLoS Pathog*, *7*(9), e1002258. https://doi.org/10.1371/journal.ppat.1002258

Wei, Q., Minh, P. N., Dotsch, A., Hildebrand, F., Panmanee, W., Elfarash, A., Schulz, S., Plaisance, S., Charlier, D., Hassett, D., Haussler, S., & Cornelis, P. (2012). Global regulation of gene expression by OxyR in an important human opportunistic pathogen. *Nucleic Acids Res*, *40*(10), 4320-4333. https://doi.org/10.1093/nar/gks017

Wong, C. F., Shin, J., Subramanian Manimekalai, M. S., Saw, W. G., Yin, Z., Bhushan, S., Kumar, A., Ragunathan, P., & Gruber, G. (2017). AhpC of the mycobacterial antioxidant defense system and its interaction with its reducing partner Thioredoxin-C. *Sci Rep*, *7*(1), 5159. https://doi.org/10.1038/s41598-017-05354-5

Wood, N. A., Swoboda, A. R., Blocker, A. M., Fisher, D. J., & Ouellette, S. P. (2022). Tag-Dependent Substrate Selection of ClpX Underlies Secondary Differentiation of Chlamydia trachomatis. *mBio*, *13*(5), e0185822. https://doi.org/10.1128/mbio.01858-22

Yang, K. S., Kang, S. W., Woo, H. A., Hwang, S. C., Chae, H. Z., Kim, K., & Rhee, S. G. (2002). Inactivation of human peroxiredoxin I during catalysis as the result of the oxidation of the

catalytic site cysteine to cysteine-sulfinic acid. *J Biol Chem*, *277*(41), 38029-38036. https://doi.org/10.1074/jbc.M206626200

Zhang, B., Gu, H., Yang, Y., Bai, H., Zhao, C., Si, M., Su, T., & Shen, X. (2019). Molecular Mechanisms of AhpC in Resistance to Oxidative Stress in Burkholderia thailandensis. *Front Microbiol*, *10*, 1483. https://doi.org/10.3389/fmicb.2019.01483

Zuo, F., Yu, R., Khaskheli, G. B., Ma, H., Chen, L., Zeng, Z., Mao, A., & Chen, S. (2014). Homologous overexpression of alkyl hydroperoxide reductase subunit C (ahpC) protects Bifidobacterium longum strain NCC2705 from oxidative stress. *Res Microbiol*, *165*(7), 581-589. https://doi.org/10.1016/j.resmic.2014.05.040