Altering the redox status of *Chlamydia trachomatis* directly impacts its developmental

cycle progression

Vandana Singh and Scot P. Ouellette*

Department of Pathology, Microbiology, and Immunology, College of Medicine, University of

Nebraska Medical Center, Omaha, NE 68198

*Correspondence: scot.ouellette@unmc.edu

Keywords: *Chlamydia*, Alkyl hydroperoxide reductase subunit C (AhpC), CRISPRi,

Differentiation, Development, Oxidative stress, Peroxides, Reactive oxygen species (ROS),

ROS scavengers

ABSTRACT

Chlamydia trachomatis is an obligate intracellular bacterial pathogen with a unique

developmental cycle. It differentiates between two functional and morphological forms:

elementary body (EB) and reticulate body (RB). The signals that trigger differentiation from one

form to the other are unknown. EBs and RBs have distinctive characteristics that distinguish

them, including their size, infectivity, proteome, and transcriptome. Intriguingly, they also differ in

their overall redox status as EBs are oxidized and RBs are reduced. We hypothesize that

alterations in redox may serve as a trigger for secondary differentiation. To test this, we

examined the function of the primary antioxidant enzyme alkyl hydroperoxide reductase subunit

C (AhpC), a well-known member of the peroxiredoxins family, in chlamydial growth and

development. Based on our hypothesis, we predicted that altering the expression of *ahpC* will

modulate chlamydial redox status and trigger earlier or delayed secondary differentiation. To test

this, we created *ahpC* overexpression and knockdown strains. During *ahpC* knockdown, ROS

levels were elevated, and the bacteria were sensitive to a broad set of peroxide stresses.

Interestingly, we observed increased expression of EB-associated genes and concurrent higher

production of EBs at an earlier time in the developmental cycle, indicating earlier secondary

differentiation occurs under elevated oxidation conditions. In contrast, overexpression of AhpC

created a resistant phenotype against oxidizing agents and delayed secondary differentiation.

Together, these results indicate that redox potential is a critical factor in developmental cycle

progression. For the first time, our study provides a mechanism of chlamydial secondary

differentiation dependent on redox status.

Word count: 250

INTRODUCTION

All organisms that are exposed to oxygen are necessarily subjected to oxidative stress. Specifically, the process of metabolizing substrates in the presence of oxygen can generate reactive oxygen species (ROS), which are toxic at high enough concentrations. Thus, from bacteria to humans, systems have evolved to mitigate the accumulation of ROS. At the same time, host defense mechanisms have evolved to leverage ROS production as a means of limiting pathogen growth and survival. Not surprisingly, pathogens have co-evolved to resist these defense mechanisms. For example, many pathogens possess a variety of antioxidant enzymes, such as catalases, glutathione peroxidases, and peroxiredoxins, that help them both subvert ROS-mediated immune system assaults and mitigate metabolic ROS byproducts (Staerck et al., 2017; Wan et al., 2021).

Chlamydia is an obligate intracellular bacterium that has significantly reduced its genome size and content in adapting to obligate host dependence. *Chlamydia trachomatis*, the leading cause of bacterial sexually transmitted diseases and preventable infectious blindness, lacks homologs to catalases or glutathione peroxidases but does possess a homolog of Alkyl hydroperoxide reductase subunit C (AhpC). AhpC is a well-known member of the peroxiredoxins family and is widely conserved in prokaryotes (de Oliveira et al., 2021). Peroxiredoxins can scavenge hydrogen peroxide, peroxynitrite, and organic hydroperoxides (Parsonage et al., 2008; Poole & Ellis, 1996; Seaver & Imlay, 2001) and act as the primary scavenger in pathogens that lack both catalase and glutathione peroxidases (Mastronicola et al., 2014; Richard et al., 2011). Several studies from other bacterial systems have shown that AhpC has a significant role in ROS and RNI scavenging, virulence, and persistence (Cosgrove et al., 2007; Kimura et al., 2012; Oh & Jeon, 2014), and deletion of AhpC creates highly oxidized conditions within the bacterium (Zhang et al., 2019). However, the function of AhpC in chlamydial biology has not been characterized.

Chlamydia is unique among pathogens in undergoing a complex developmental cycle

that comprises two distinct morphological forms, the elementary body (EB) and reticulate body

(RB) (Abdelrahman & Belland, 2005). The EB is the smaller (~0.3 μm), infectious, and non-

dividing form capable of infecting susceptible host cells. Once internalized into a host-derived

vacuole termed an inclusion, the EB differentiates into the non-infectious, larger (~1 μm), and

replicating RB - this process is known as primary differentiation and represents the early phase

of the developmental cycle (Clifton et al., 2005). RBs replicate within the inclusion in the mid-

cycle phase using an asymmetric, MreB-dependent polarized division process (Abdelrahman et

al., 2016; Lee et al., 2020; Ouellette et al., 2022; Ouellette et al., 2020). In the late phase of the

developmental cycle, RBs asynchronously condense into EBs, and this process is termed

secondary differentiation. Despite these well-defined differences between chlamydial

morphological forms, how *Chlamydia* mechanistically differentiates between functional forms

remains unclear.

Intriguingly, a recent study evaluated the redox potential of *Chlamydia* and demonstrated

that RBs are reduced, whereas EBs are oxidized (Wang et al., 2014). This is consistent with

earlier studies that revealed differences in the crosslinking of outer membrane proteins and type

III secretion-related proteins in chlamydial developmental forms (Betts-Hampikian & Fields,

2011; Caldwell et al., 1981; Everett & Hatch, 1995; Wang et al., 2014). Taken together, these

observations indicate that, during the developmental cycle, the redox potential of the bacteria is

changing. However, whether redox changes in the bacteria directly affect developmental cycle

progression is not characterized. Based on the different redox status of the chlamydial

developmental forms, we hypothesized that increasing the oxidation of the RB drives secondary

differentiation to the EB. We used chlamydial transformants designed to overexpress or reduce

AhpC levels to explore the effects of altered redox potential in chlamydial growth and

development to test this hypothesis. This study establishes the role of AhpC as an antioxidant in

Chlamydia, as demonstrated by its ability to counteract different peroxide stresses when

overexpressed. Overexpression of AhpC had no negative effect on bacterial replication but

delayed the differentiation of RBs to EBs. In contrast, under conditions of *ahpC* knockdown, the

organism was highly sensitive to oxidizing conditions. Interestingly, this change in redox

potential caused earlier expression of EB-associated (late) genes and production of EBs,

leading to a shift in developmental cycle progression. This earlier activation of gene expression

related to secondary differentiation in *ahpC* knockdown was also observed when developmental

cycle progression was blocked by penicillin treatment. Taken together these data provide

mechanistic insight into chlamydial secondary differentiation and are the first to demonstrate

redox-regulated differentiation in *Chlamydia*.

RESULTS

The redox threshold hypothesis and chlamydial secondary differentiation

The chlamydial EB and RB developmental forms differ in size, infectivity, and division capacity. In addition to these characteristics, both forms also differ in their redox status: EBs are oxidized and RBs are reduced (Fig. 1A). Based on this information, we hypothesized that changing redox potential is a critical factor in the process of differentiation from one form to the other. We named this the "redox threshold hypothesis". In this scenario, as soon as a given RB has crossed an oxidative threshold, the activity of critical proteins is modified to trigger differentiation to the EB (Fig. 1B). AhpC is an antioxidant enzyme, and several reports have indicated that ablation of *ahpC* creates highly oxidized conditions in bacteria (Feng et al., 2020; Zhang et al., 2019). Therefore, we hypothesize that AhpC, which maintains redox homeostasis in other bacteria, is a critical factor in mediating developmental cycle progression and that altering its activity will impact developmental cycle progression consistent with the redox threshold hypothesis (Fig. 1C).

Overexpression of *ahpC* has minimal impact on chlamydial growth

To study the effect of AhpC on chlamydial developmental cycle progression, we first generated an *ahpC* overexpression (OE) strain using a plasmid encoding an anhydrotetracycline (aTc)-inducible *ahpC* (untagged) and transformed it into a *C. trachomatis* L2 strain lacking its endogenous plasmid (-pL2). The same plasmid vector backbone (i.e., encoding mCherry in place of *ahpC*) was used as an empty vector control (EV). HeLa cells were infected with these transformants, and, at 10 hpi, expression of the construct was induced or not. Lacking an antibody against AhpC, overexpression of *ahpC* was validated by reverse transcription-quantitative PCR (RT-qPCR). Here, an approximate 1-log increase in transcripts of *ahpC* was detected at 14 and 24 hpi in the induced strain in comparison to the uninduced strain (Fig. 2A). Immunofluorescence analysis (IFA) was performed at 24 hpi to examine the organisms' morphology and overall inclusion growth. For IFA, individual bacteria were labeled with an anti-

major outer membrane protein (MOMP) antibody, and DAPI was used to stain for DNA. IFA

imaging revealed that there was no observable impact on the morphology of the EV control

strain under the conditions tested. In contrast, we did note that overexpression of

ahpC increased the overall inclusion area (Fig. 2B and 2C). We next quantified the total number

of bacteria (i.e., both RBs and EBs) by measuring genomic DNA and observed a significant

increase in gDNA levels at 24 hpi in response to increased *ahpC* expression (Fig. 2D). Although

inclusions were larger in area and contained more total bacteria as assessed by gDNA levels

when overexpressing *ahpC*, recoverable inclusion forming units (IFUs; a proxy for infectious

EBs) were significantly lower under these conditions at 24 hpi (Fig. 2E). However, by 48 hpi, the

difference between the induced and uninduced conditions, although still reduced, was not

statistically significant. The inclusion forming unit analysis only measures viable EBs from a

population, suggesting that AhpC overexpression skews the ratio of RBs to EBs. Overall, these

data support the hypothesis that overexpressing *ahpC* delays production of infectious EBs.

Overexpression of *ahpC* confers resistance to peroxides in *Chlamydia*

AhpC is an important peroxiredoxin involved in oxidative damage defense. Before determining

whether increased *ahpC* expression impacts chlamydial sensitivity to oxidizing agents, we first

sought to determine the response of *C. trachomatis* to inorganic or organic hydroperoxides such

as hydrogen peroxide (H_2O_2), cumene hydroperoxide (CHP), and tert-butyl hydroperoxide

(TBHP) as well as peroxynitrite (PN). To do this, we evaluated the sensitivity of HeLa cells,

infected or not with the empty vector (EV) control strain, to different concentrations of oxidizing

agents using a viability assay. Both uninfected and EV-infected HeLa cells tolerated up to 1 mM

oxidizing agents for 30 min (Fig. S1). Next, different concentrations (lower than 1 mM) of these

oxidants were tested on wild-type (WT) *C. trachomatis* L2/434/Bu (*Ctr* L2). HeLa cells were

infected with *Ctr* L2 and, at 16 hpi, were exposed to different concentrations of exogenous

oxidizing agents for 30 min only, before washing out the oxidizing agents and replacing the

media. Samples for IFU and IFA assays were then collected at 24 hpi to assess effects of the

oxidizing agents on growth of *Ctr* L2. 62.5 μM concentration of all three inorganic and organic

peroxides had no appreciable effect on IFUs, and inclusion size and morphology also remained

unaffected at this sublethal concentration. The concentrations of inorganic and organic

peroxides that resulted in a decrease in IFUs to ~50% of the untreated culture were 500 μM

(H_2O_2) and 250 μM (CHP and TBHP), respectively, with a concurrent decrease in inclusion size.

1 mM H_2O_2, CHP, and TBHP caused >90% reduction in IFUs and small inclusions. PN at 1 mM

concentration was less effective than other oxidizing agents and showed only a ~30% decline in

IFUs compared to the untreated culture (Fig. S2).

To further examine the antioxidant functions of AhpC, we next exploited our *ahpC*

overexpression strain to explore its capability to protect chlamydiae from oxidizing agents. Both

ahpC OE and EV strains were used to infect HeLa cells and, at 10 hpi, expression of the

constructs was induced or not with 1 nM aTc. At 16 hpi, infected cells were exposed to various

concentrations of exogenous oxidizing agents for 30 min. At 24 hpi, IFU and IFA samples were

collected to measure chlamydial growth and assess chlamydial morphology, respectively. As

shown in Figure 3A, 3B, and S3, the *ahpC* overexpression strain showed increased resistance

to all oxidants tested as the mean IFUs and the size of inclusions were greater than that of the

uninduced but treated control. There was no change in bacterial growth and morphology in the

case of 62.5 μM concentrations of oxidizing agents, which was anticipated since we determined

this was a sublethal concentration. The resistant phenotype was more evident at the higher

concentrations of H_2O_2, CHP, and TBHP. Overexpression of *ahpC* also contributed to higher

resistance against PN. Conversely, the empty vector control strain behaved as expected (i.e.,

like WT) in the presence of oxidizing agents, with similar results observed in the presence or

absence of aTc (Fig. 3C, 3D, and S3). Collectively, these data demonstrate that the chlamydial

AhpC possesses antioxidant activity, as expected, and that increased *ahpC* expression is

protective for *Chlamydia* in the presence of increased oxidative stress.

Knockdown of *ahpC* negatively impacts chlamydial growth

To further define the function(s) of AhpC in the chlamydial developmental cycle, we used a novel dCas12-based CRISPR interference (CRISPRi) strategy adapted for *Chlamydia* by our lab (Ouellette, 2018; Ouellette et al., 2021). We generated an *ahpC* knockdown strain harboring the pBOMBL12CRia with a crRNA targeting the *ahpC* 5' intergenic region (plasmid designated as pL12CRia(*ahpC*)). We used a strain carrying a pL12CRia plasmid with a crRNA with no homology to any chlamydial sequence (i.e., non-targeting [NT]) to serve as a negative control. To confirm the knockdown of *ahpC*, RT-qPCR was employed. Here, we infected HeLa cells with the *ahpC* knockdown (*ahpC* KD) or NT strains and induced dCas12 expression or not at 10 hpi using 1 nM aTc. Nucleic acid samples were harvested at 14 hpi and 24 hpi. RT-qPCR analysis of the *ahpC* KD revealed approximately 90% reduction of *ahpC* transcripts compared to the uninduced control, thus confirming the knockdown of *ahpC* (Fig. 4A). In the case of the NT control, there was no effect on *ahpC* transcripts. IFA of these strains revealed noticeably smaller inclusions after blocking expression of *ahpC* as compared to the uninduced sample or the NT conditions (Fig. 4B). We next measured total bacterial counts using genomic DNA. Surprisingly, even though inclusions were smaller during *ahpC* knockdown, we observed slightly higher gDNA levels at 14 hpi compared to the uninduced control, followed by only a small increase from 14 to 24 hpi (Fig. 4C). In contrast, the uninduced strain showed a logarithmic increase in gDNA levels during this timeframe. These effects were not detected for the NT strain (Fig. 4C). To further explore the effect of reduced *ahpC* transcripts in *Chlamydia*, IFU assays were performed. IFU analysis exhibited severely reduced progeny (>90%) in *ahpC* knockdown compared to its respective uninduced control at both time points assessed (24 and 48 hpi) (Fig. 4D). In contrast, the NT strain showed less than a 50% reduction after inducing dCas12 expression at these timepoints consistent with prior observations that dCas12 expression slightly delays developmental cycle progression (Hatch & Ouellette, 2023; Reuter et al., 2023).

Taken together, these analyses indicate that reducing *ahpC* levels and/or activity severely

disrupted chlamydial growth and development.

To investigate whether *ahpC* knockdown resulted in increased ROS levels in the bacteria

(Zhang et al., 2019), the intracellular ROS levels were measured in infected cells in *ahpC*

knockdown conditions and compared to the uninduced control condition and uninfected cells.

We used the cell-permeable, fluorogenic dye CellROX Deep Red to measure ROS levels. This

dye remains non-fluorescent in a reduced state and exhibits bright fluorescence once oxidized

by ROS. ROS generation was measured in uninfected and *ahpC* KD-infected HeLa cells at 24

and 48 hpi. As shown in Figure 4E, *ahpC* knockdown resulted in significantly higher ROS than

the uninfected or infected but uninduced samples at both 24 hpi and 48 hpi time points. These

results together indicate that AhpC plays an important role in counteracting oxidative stress by

reducing ROS in *C. trachomatis*.

Knockdown of *ahpC* sensitizes *Chlamydia* to oxidizing agents

As overexpression of *ahpC* resulted in increased resistance to oxidizing agents, we next

explored whether knockdown of *ahpC* resulted in increased sensitivity to such agents. As

described above, *ahpC* KD and NT strains were used to infect HeLa cells and, at 10 hpi,

expression of dCas12 was induced or not with 1 nM aTc. Considering that the *ahpC* KD already

demonstrated reduced IFUs and inclusion sizes at 24 hpi, only sublethal concentrations (62.5

μM) of exogenous oxidizing agents were applied. At 24 hpi, IFA and IFU analyses were

performed to quantify the effect of the oxidants on chlamydial growth in the absence of *ahpC*

activity. Under conditions of *ahpC* knockdown, even sublethal concentrations (62.5 μM) of H_2O_2,

CHP, or TBHP further reduced the inclusion size, and the IFU data revealed a decrease from

>90% to <30% in comparison to the untreated control (Fig. 5A, 5B, and S4). Reduced

expression of *ahpC* also affected the survival of *Chlamydia* against peroxynitrite. Notably, there

was no significant change in the NT control strain in the uninduced and induced samples in

response to oxidizing agents as assessed by IFU and IFA (Fig. 5C, 5D, and S4). These data

further support that the chlamydial AhpC is a critical antioxidant enzyme in these bacteria.

Complementation restores the growth and resistance to low levels of peroxide stress of

the *ahpC* knockdown strain

To validate that the impaired growth, altered inclusion morphology, and enhanced sensitivity to

peroxides were due to the decreased level/activity of AhpC, we generated a complemented

strain of *ahpC* knockdown. For the construction of the *ahpC* complementing plasmid, the *ahpC*

gene was cloned and transcriptionally fused 3' to the dCas12 in the pL12CRia(*ahpC*)

knockdown plasmid. Here, the complementing *ahpC* allele is also under the control of the aTc-

induced P$_{tet}$ promoter and is co-expressed with the aTc-inducible dCas12. Consequently, the

ahpC knockdown effect is ablated, and the observed phenotypes should be restored. After

inducing *dCas12-ahpC* expression with aTc, the resultant strain was verified by RT-qPCR.

Increased transcripts for *ahpC* were quantified under these conditions, indicating successful

complementation of the knockdown effect (Fig. 6A). Of note, the *ahpC* transcript levels

remained elevated in comparison to the uninduced control at the 24hpi time point. After

confirming this strain, we measured genomic DNA to quantify total bacteria and performed IFA

and IFU assays to examine if complementation restored the phenotypes observed during *ahpC*

knockdown. These assays revealed normal inclusion morphology, gDNA levels, and EB progeny

production in the complemented strain, indicating successful complementation of the

knockdown phenotype (Fig. 6B, C, and D). Of note, a C-terminal 6xHis tagged AhpC was not

capable of complementing the knockdown phenotype (data not shown), indicating a requirement

for a free C-terminus in the function of AhpC in *Chlamydia*. Previous studies in other bacteria

have revealed that the C-terminal residues in AhpC play a crucial role in the structural stability

and enzymatic activity of AhpC (Dip et al., 2014; Feng et al., 2020; Wan et al., 2021).

We next assessed whether complementation of the knockdown phenotype could also

restore the resistance to low levels of peroxide stress. As in previous experiments, a sublethal

concentration of oxidizing agents was added for 30' at 16 hpi after having induced *dCas12-*

ahpC expression at 10 hpi. Consistent with the growth parameters (Fig. 6B-D), the

complemented strain showed wild-type responses in these conditions (Fig. 6E, 6F, and S4).

These data indicate that the enhanced susceptibility of the *ahpC* knockdown strain to oxidizing

agents was due to the reduced levels of *ahpC* and not an indirect effect of knockdown.

We predicted that the growth defects resulting from higher production of ROS during

ahpC knockdown could be rescued by treating the *ahpC* KD with ROS scavengers. To test this

prediction, we utilized two characterized ROS scavengers, DMTU (*N,N'-dimethylthiourea*) and

α-tocopherol, which scavenge H_2O_2 and peroxyl radical, respectively (Hollander-Czytko et al.,

2005; Kiffin et al., 2006; Walch et al., 2015). First, the effect of ROS scavengers on uninfected

and infected HeLa cells was investigated using a viability assay. This assay revealed that 10

mM DMTU and 100 µM α-tocopherol had no adverse effects on uninfected or EV-infected HeLa

cells (data not shown). These same concentrations were tested on WT *Ctr* L2 infected HeLa

cells in untreated and 500 µM H_2O_2 treated conditions to test the potency of these scavengers

to rescue growth defects associated with this concentration of oxidizing agent. Scavengers were

added at 9.5 hpi and washed away at 16 hpi - the time of addition of H_2O_2 in the respective

samples. At 16.5 hpi, after 3 wash steps, scavengers were added again with fresh media, and,

at 24 hpi, bacterial growth and morphology were assessed using IFU and IFA assays. Neither

scavenger had a significant impact on chlamydial morphology or infectious progeny. In the case

of peroxide (500 µM H_2O_2) treated samples, scavengers restored IFUs from ~50% to ~100%,

and inclusion size was also recovered (Fig. S5A and B).

Next, the effect of the ROS scavengers under *ahpC* knockdown conditions was

assessed. In this experiment, scavengers were added at 9.5 hpi to provide the protective effect

of scavengers before reducing the activity of *ahpC*. At 10 hpi, knockdown was induced or not

with 1 nM aTc, and, at 14 hpi, scavengers were added again. At 24 hpi, IFU and IFA samples

were collected to examine bacterial growth and morphology to assess the effect of ROS

scavenging on the *ahpC* knockdown phenotype. As shown in Fig. 6G and 6H, both ROS

scavengers had a positive impact on restoring growth of the *ahpC* knockdown strain; inclusions

were larger, and IFUs were increased from <10% to >90% in the presence of scavengers in the

induced samples. These data provide compelling evidence that the adverse effects of *ahpC*

knockdown are due to increased ROS accumulation in this strain.

Chlamydial developmental cycle progression is altered by *ahpC* knockdown/

overexpression

Whereas the overexpression of AhpC delayed overall developmental progression and was

consistent with our hypothesis, the results with the *ahpC* knockdown strain appeared to refute

our hypothesis given the apparent decrease in IFUs and smaller inclusion sizes after

knockdown. To investigate the effects of changes in redox potential on chlamydial

developmental cycle progression, we performed a transcriptional analysis of well-characterized

late-cycle genes associated with secondary differentiation (*hctA*, *hctB*, *glgA*, *tsp,* and *omcB*)

(Belland et al., 2003; Gehre et al., 2016; Newhall, 1987; Ouellette et al., 2006; Swoboda et al.,

2023). In the uninduced samples, transcript levels of all the tested late genes were higher at 24

hpi compared to 14 hpi, indicating their normal expression during the late stage of the

developmental cycle. In comparison to the uninduced control, under conditions of *ahpC*

knockdown, significantly higher expression of *hctA*, *hctB*, *omcB, tsp,* and *glgA* was observed at

the mid-developmental cycle timepoint of 14 hpi (Fig. 7A). At 24 hpi, expression of these genes

was comparable between the uninduced and induced conditions in the *ahpC* knockdown strain.

In contrast, the complementation strain showed no increase in the expression of these tested

late genes at 14 hpi, and *tsp* and *glgA* transcripts were reduced (Fig. 7C). Consistent with our

model, the overexpression of *ahpC* resulted in significantly lower expression of *hctA*, *hctB*,

omcB, tsp, and *glgA* at 14 and 24 hpi (Fig. 7D), suggesting a delayed transition to EBs.

Given the increase in late gene expression during *ahpC* knockdown at an earlier time

point in the developmental cycle than normal, we reasoned that the increase in oxidizing

conditions might prematurely trigger secondary differentiation and EB production. However,

complicating such an analysis is that fewer RBs are present to convert to EBs, which would

result in overall lower IFU yields as we had measured at 24 and 48 hpi (Fig. 4D). Nonetheless,

to assess EB production more rigorously at earlier time points in the developmental cycle, we

performed a growth curve analysis to precisely measure IFUs at two-hour intervals from 16 to

24 hpi. HeLa cells were infected, and knockdown was induced or not at 10 hpi with 1 nM aTc. At

the indicated time points, IFU samples were collected and quantified. Consistent with our

prediction, this experiment revealed higher IFUs (i.e., EBs) at 16 and 18 hpi during *ahpC*

knockdown compared to the uninduced samples (Fig. 7B). This difference was statistically

significant at 18 hpi. However, further EB production was stalled, with the uninduced strain

continuing to produce EBs such that, by 24 hpi, there were significantly higher EB yields under

these conditions (as noted in Fig. 4D). These data show that the phenotypic consequence of

higher expression at 14 hpi of genes functionally related to EBs is the concomitant earlier

production of EBs. These data underscore that reduced activity of *ahpC* causes earlier

secondary differentiation in *C. trachomatis*.

ahpC knockdown activates transcription of late-cycle genes when bacterial replication is

blocked

We detected earlier expression of EB-related genes during *ahpC* knockdown and were curious

if the *ahpC* knockdown condition could activate these genes under conditions when the

chlamydial developmental cycle is blocked. Pathogenic *Chlamydia* species undergo a polarized

cell division process, in which peptidoglycan is transiently synthesized only at the division

septum (Abdelrahman et al., 2016; Cox et al., 2020; Ouellette et al., 2020). As a result, during

penicillin treatment, division of RBs is blocked (Moulder, 1993; Moulder et al., 1956; Ouellette et

al., 2020). However, the bacteria continue to grow in size resulting in aberrantly enlarged RBs

(Barbour et al., 1982; Matsumoto & Manire, 1970; Ouellette et al., 2012), in which EB-related

genes are not transcribed and production of EBs is inhibited (Ouellette et al., 2006; Panzetta et

al., 2018).

To examine this, we generated an *ahpC* KD strain with spectinomycin resistance (*ahpC*

KD-spec) and validated its phenotype as being the same as the penicillin-resistant *ahpC* KD

strain. This new strain, *ahpC* KD-spec, was used to infect HeLa cells, and knockdown was

induced or not at 10 hpi with 1 nM aTc. At the same time, samples were treated or not with 1

unit per mL of penicillin (Pen). RNA samples were harvested at 16 and 24 hpi and processed for

RT-qPCR (Fig. S6B-D). IFA controls from these different conditions demonstrated the expected

phenotypes (Fig. S6A). For example, Pen treatment caused aberrantly enlarged RBs,

irrespective of *ahpC* KD, whereas *ahpC* KD itself caused smaller inclusions. Similarly, *ahpC* KD

resulted in reduced *ahpC* transcripts in the induced conditions irrespective of the presence of

Pen (Fig. S6B). We next quantified transcript levels for the late genes *hctA* and *hctB* in these

different conditions (Fig. S6C and D). As expected, in the uninduced and untreated condition,

both transcripts had higher expression at 24 hpi, indicating their regular developmental

expression during the late stage of the developmental cycle. Consistent with prior observations

(Ouellette et al., 2006), transcript levels of these genes in uninduced+Pen conditions was low.

As we previously noted (Fig. 7A), the transcripts of *hctA* and *hctB* are higher in the induced than

uninduced samples at 16 hpi in the absence of Pen. Interestingly, in the induced+ Pen condition,

these genes had higher expression at 24 hpi than the uninduced+Pen control. At 16 hpi, their

expression was either higher or similar between both samples (i.e., induced+Pen and

uninduced+Pen). Collectively, these data support our observation that *ahpC* knockdown

activates the transcription of late-cycle genes – even under conditions where developmental

cycle progression is blocked.

DISCUSSION

Secondary differentiation (differentiation from RBs to EBs) is an essential step for chlamydial growth and survival, but there is a dearth of information regarding the mechanisms of its regulation. EBs and RBs have significantly different proteomic repertoires, and our group has identified critical functions for the cytoplasmic ClpXP and ClpCP and periplasmic Tsp proteases during secondary differentiation (Pan, Jensen et al., 2023; Swoboda et al., 2023; Wood et al., 2022). The *tsp* gene, in addition to another late-cycle gene, *hctB*, is transcriptionally regulated by sigma factor 28 (σ^{28}) (Hatch & Ouellette, 2023). Another sigma factor, σ^{54} has been linked to regulation of outer membrane components, type III secretion system components, and other genes typically expressed late in development (Hatch & Ouellette, 2023; Soules, LaBrie, et al., 2020) . In addition to the two minor sigma factors, σ^{28} and σ^{54}, *Chlamydia* encodes one major sigma factor, σ^{66}, which is transcriptionally regulated by the relative levels of RsbV1 (antagonist) and RsbW (anti-sigma factor) (Thompson et al., 2015). This Rsb system senses ATP availability and has been proposed to regulate σ^{66} (Thompson et al., 2015). A related study from Soules et al. (Soules, Dmitriev, et al., 2020) showed that TCA intermediates act as ligands for RsbU, thereby linking the Rsb system to the TCA cycle and ATP synthesis. Collectively, these data show that post-translational mechanisms drive secondary differentiation in *Chlamydia*. However, no definitive "switch" that triggers this step has been identified.

Aside from the clear morphological and functional differences between the EB and RB, they also differ in their redox status: EBs are more oxidized whereas RBs are more reduced (Wang et al., 2014). This same study noted that the overall redox status of the host cell was relatively unchanged throughout the developmental cycle, suggesting that changes in the bacteria were driven by endogenous processes (i.e., metabolism). Though *Chlamydia* is dependent on its host for most of its energy requirements, it has some metabolic activities, such as a partial TCA cycle and oxidative phosphorylation (Gerard et al., 2002; Iliffe-Lee & McClarty, 1999) that can be possible sources of intracellular ROS. We hypothesize that accumulating

redox stress from its metabolic activities serves as a signal to trigger secondary differentiation

from RBs to EBs. There is a paucity of knowledge on how *C. trachomatis* modulates oxidative

stress during disease pathogenesis. However, some studies explored the effects of redox

changes on *Chlamydia* growth. Boncompain et al. reported that infection of *C. trachomatis*

induced the transient production of ROS by the host cell at a moderate level for the initial few

hours of infection only (Boncompain et al., 2010). Another study also found a similar observation

about ROS during infection, reporting that *Chlamydia* requires host-derived ROS for its growth

(Abdul-Sater et al., 2010). Given that no studies have examined the role of chlamydial proteins

related to redox and how these proteins impact the growth of the pathogen, we initiated

experiments to investigate the function of a key antioxidant enzyme, AhpC, as a means of

testing our hypothesis.

Reactive oxygen species (ROS) generation is an inevitable condition for pathogens

growing in aerobic conditions, and resistance against oxidative stress (imbalanced level of

ROS) is a key survival mechanism (Fang, 2011). Hence, pathogens have evolved detoxifying

proteins, such as peroxiredoxins, to eliminate ROS (Dip et al., 2014; Wang et al., 2004; Yang et

al., 2002). AhpC is a 2-cys peroxiredoxin and has been reported to have a crucial role in

bacterial physiology, survival, and virulence by scavenging ROS and RNI (Cosgrove et al.,

2007; Kimura et al., 2012; Loprasert et al., 2003; Oh & Jeon, 2014). In the process of

detoxification of peroxides and peroxynitrite, AhpC is converted into an oxidized dimer, requiring

alkyl hydroperoxide reductase AhpF or AhpD to regenerate its activity (Koshkin et al., 2004;

Poole & Ellis, 1996; Wong et al., 2017). *C. trachomatis* encodes AhpC (Ct603) but lacks any

annotated homologs of AhpF or AhpD. Therefore, it remains an open question how AhpC

activity is regulated. Some studies have mentioned *ahpC* as an iron-responsive gene in

Chlamydia (Brinkworth et al., 2018; Pokorzynski et al., 2019), but there is no detailed

investigation to date about the role of this crucial antioxidant in the growth and development of

Chlamydia.

Using a broad set of methodologies, we created and characterized overexpression and

knockdown conditions of *ahpC*. Higher expression of *ahpC* resulted in lower EB production at

24 hpi coupled with a higher number of RBs and larger inclusion size (Fig. 2), suggesting

differentiation from RBs to EBs is delayed. The number of EBs is further decreased at 48 hpi,

but the difference is not statistically significant. This may be due to the highly oxidized

conditions in *Chlamydia* at 48 hpi (Wang et al., 2014) or the inability of *ahpC* overexpression to

generate sufficient reducing conditions to neutralize it completely. In our studies, *ahpC*

expression was induced at 10hpi, and adequate enzyme levels may not have been sustained

later during the developmental cycle. Consistent with studies in other bacteria (Sherman et al.,

1999; Zuo et al., 2014), overexpression of *ahpC* created resistance to different oxidants (Fig. 3

and S3), indicating that AhpC is the principal defense mechanism against oxidative stress in

Chlamydia. During *ahpC* knockdown, the attenuated activity of AhpC severely affected the

growth of bacteria (Fig. 4). Notably, the morphology of inclusions during *ahpC* knockdown is

strikingly similar to *Chlamydia* treated with a high concentration (1 mM) of oxidizing agents (Fig.

4 and Fig S2), indicating similar scenarios in both conditions. This was further supported by the

increased sensitivity of *ahpC* knockdown to sublethal concentrations of oxidants (Fig. 5 and S4).

Again, this observation is consistent with previous data from other bacteria (Cosgrove et al.,

2007; Seaver & Imlay, 2001; Storz et al., 1989; Zhang et al., 2019).

An essential aspect of the scavenging activity of AhpC for many bacterial pathogens in

which it has been studied is that it works best with endogenous (i.e., low) levels of H_2O_2. These

bacteria, such as *Staphylococcus aureus* and *Yersinia pseudotuberculosis*, encode catalases,

which have high Km for hydrogen peroxide and serve a predominant role in scavenging

exogenous H_2O_2 (Cosgrove et al., 2007; Wan et al., 2021). Catalases can detoxify H_2O_2 at high

levels (millimolar levels) and are crucial in responding to external H_2O_2 stress (Mishra & Imlay,

2012). The *ahpC* mutants in these bacteria became sensitive to organic peroxides but resistant

to H_2O_2 due to higher catalase activity as a compensatory response to the lack of AhpC

(Antelmann et al., 1996; Mongkolsuk et al., 2000; Ochsner et al., 2000). Further, simultaneous

mutations in both catalase and *ahpC* displayed drastically enhanced sensitivity to all oxidizing

agents tested (Cosgrove et al., 2007; Ezraty et al., 2017; Seaver & Imlay, 2001). *C. trachomatis*

does not encode a catalase gene (Boncompain et al., 2014; Rusconi & Greub, 2013), and

hypersensitivity of *ahpC* knockdown to both inorganic and organic peroxides indicates the

absence of catalase and establishes that AhpC is the primary scavenger of ROS in *C.*

trachomatis. We did not observe significant effects of PN with AhpC expression. One possible

explanation may be our study's concentration (1 mM) of PN. Higher concentrations of PN could

not be used due to toxic effects on the host cell. The other possibility may be the presence of

some other unknown mechanism(s) to detoxify PN. For example, *Chlamydia* also encodes a

superoxide dismutase (SOD) that may prevent the accumulation of the necessary precursors

that are needed for PN production. We are currently investigating the function of the chlamydial

SOD enzyme.

Previous studies in other bacterial systems indicated that ROS production significantly

increased after the deletion of *ahpC* (Zhang et al., 2019). Similarly, in *Chlamydia* during *ahpC*

knockdown, the ROS level was dramatically higher. Its ability to dissipate ROS produced by

exogenous oxidative stress was severely compromised (Fig. 4E). Further, the addition of ROS

scavengers to the culture medium rescued the negative phenotypes associated with *ahpC*

knockdown (Fig. 6G and H). This strongly suggests that, in the absence of AhpC, higher

amounts of reactive oxygen species accumulate in *Chlamydia* and that the negative impact on

growth during *ahpC* knockdown is due to highly oxidized conditions in the organism. Moreover,

for the complete restoration of inclusion size, multiple doses of scavengers were required

throughout the experiment, further emphasizing that the internal chlamydial environment is

oxidized and/or susceptible to oxidation in the absence of AhpC. These findings suggest that

AhpC in *Chlamydia* is indispensable, having a functional role even more extensive than

previously reported in other bacteria.

Interestingly, in our RT-qPCR study, late-cycle genes such as *hctA*, *hctB*, *glgA, omcB*,

and *tsp* are expressed at a higher level at an earlier time (14 hpi) in the chlamydial

developmental cycle as a result of reduced AhpC activity (Fig. 7). Notably, in the developmental

cycle of *C. trachomatis*, only RBs are present at 14 hpi, and secondary differentiation starts after

16 hpi (Abdelrahman & Belland, 2005). Expression of most of the late-cycle genes occurs after

this time in the developmental cycle (Belland et al., 2003). Among the tested late-cycle genes,

hctA and *hctB* encode histone-like proteins responsible for chromosomal condensation during

the differentiation of RBs into EBs. The *hctA* gene is among the first to be transcribed in the late-

stage (Chiarelli et al., 2020). The other three late genes, *tsp*, *omcB*, and *glgA*, are all well-

characterized late genes associated with secondary differentiation. Tsp is a periplasmic

protease thought to be crucial in the degradation of RB-specific periplasmic proteins during

secondary differentiation in *C. trachomatis* (Swoboda et al., 2023). OmcB is responsible for the

rigid cell wall and osmotic stability of the EBs (Mygind et al., 1998; Newhall, 1987). GlgA is a

non-essential secretory protein involved in glycogen metabolism and is responsible for glycogen

accumulation in the inclusion lumen at late stages of the developmental cycle (Gehre et al.,

2016; Sun et al., 2020). One crucial point to consider is the redox status of *Chlamydia*. 14 hpi is

when reduced RBs predominate with virtually no oxidized EBs detectable. However, we

established that *ahpC* knockdown shifts the redox status of the bacteria towards oxidation.

Hence, the earlier and significantly higher detection of transcripts for these late-cycle genes

indicate earlier secondary differentiation in the *ahpC* knockdown. Importantly, *ahpC*

complementation during knockdown restored the regular developmental expression of these

late-cycle genes, further supporting this phenotype was due to the highly oxidized conditions

created by reduced AhpC activity.

The higher expression of genes functionally related to EBs and EB production at an

early stage of the chlamydial developmental cycle indicates earlier secondary differentiation as

an outcome of diminished activity of *ahpC* in *C. trachomatis*. Consequently, increased late gene

transcription should have the phenotypic effect of causing earlier production of EBs. Indeed, we

quantified more IFUs (proxy for EBs) in the *ahpC* knockdown at 16 and 18 hpi compared to the

uninduced control condition. In contrast, for the AhpC overexpression strain, the expression of

these late-cycle genes in the induced conditions compared to the uninduced control at 24 hpi is

significantly lower, indicating reduced production of EBs. These data support our hypothesis

that the developmental cycle is delayed in the reduced environment. We propose a simple

model to explain these scenarios (Fig. 8). During *ahpC* KD, higher amounts of ROS

accumulation in the bacterium create highly oxidized conditions. As secondary differentiation is

asynchronous and RBs divide through an asymmetric budding mechanism (Ouellette et al.,

2020), ROS levels are unevenly distributed between mother and daughter cell. This difference

will lead some RBs to breach an oxidative threshold sooner, allowing activation of late genes

and secondary differentiation earlier than other RBs. In contrast, *ahpC* overexpression

scavenges ROS at a greater level leading to a more reducing environment. This results in a

delay in achieving the oxidative threshold, thus allowing RBs to continue to divide before

committing to secondary differentiation. Taken together, our data directly link oxidation state and

secondary differentiation in *Chlamydia*. Ongoing studies are focused on characterizing redox-

sensitive chlamydial proteins to understand which specific factors drive the shift from RBs to

EBs (or EBs to RBs) in *C. trachomatis*.

MATERIALS AND METHODS

Strains and cell culture

For chlamydial transformation, McCoy mouse fibroblast cells (kind gift of Dr. Harlan Caldwell (NIH/NIAID)) were used. Human cervix adenocarcinoma epithelial HeLa cells (kind gift of Dr. Harlan Caldwell (NIH/NIAID)) were used for RT-qPCR, immunofluorescence assays (IFA), inclusion forming unit assays (IFU), viability assays, oxidative stress, ROS measurements, and ROS scavenger assays. Both cell types were routinely grown and passaged in Dulbecco's modified Eagle's medium (DMEM; Gibco, Waltham, MA) supplemented with 10% fetal bovine serum (FBS; Sigma, St. Louis, MO) and 10 µg/mL gentamicin (Gibco, Waltham, MA) at 37^0C and 5% CO2. All strains were verified to be Mycoplasma-negative using LookOut mycoplasma PCR detection kit (Sigma). For chlamydial transformations, *Chlamydia trachomatis* serovar L2 EBs lacking the endogenous pL2 plasmid (kind gift of Dr. Ian Clarke, University of Southampton) were used (Wang et al., 2011). Wild-type, density gradient-purified *Chlamydia trachomatis* 434/Bu (ATCC VR902B) EBs were used for sensitivity to oxidizing agents and ROS scavenger assays. Molecular biology reagents, oxidizing agents, scavengers, and CellROX Deep Red dye were purchased from Thermo Fisher unless otherwise noted.

Plasmid construction

The primers, gBlock gene fragments, plasmids, and bacterial strains used for molecular cloning are listed in Table S1 in the supplemental material. Constructs for chlamydial transformation were cloned using high-fidelity (HiFi) cloning system from New England BioLabs (NEB). Primers were designed using the NEBuilder online primer generation tool (https://nebuilderv1.neb.com). For the overexpression strain, the *ahpC* gene was amplified by PCR with Phusion DNA polymerase (NEB) using *C. trachomatis* serovar L2 434/Bu genomic DNA as a template. The PCR product was purified using a PCR purification kit (Qiagen, Hilden, Germany). The HiFi assembly reaction was performed as per the manufacturer's instructions in conjunction with the pBOMBDC plasmid digested with Fast Digest EagI and KpnI enzymes and dephosphorylated

with FastAP (Thermo Fisher, Waltham, MA). The HiFi reaction mix was transformed into *E. coli*

10β (NEB). Plasmids were first confirmed by restriction enzyme digestion, and final verification

of insert was performed using Sanger sequencing. A dCas12-based CRISPRi approach was

used for *ahpC* knockdown generation (Ouellette et al., 2021). The pBOMBL12CRia plasmid

having anhydrotetracycline (aTc) inducible catalytically dead dCas12 protein was used as

vector. A crRNA targeting the 5' intergenic region of *ahpC* was designed and ordered as a

presynthesized DNA fragment. For *ahpC* knockdown construct, 2 ng of the gBlock (Integrated

DNA Technologies [IDT], Coralville, IA) listed in Table S1 was combined with 25 ng of BamHI-

digested, alkaline phosphatase-treated pBOMBL12CRia(e.v.)::L2 in a HiFi reaction according to

the manufacturer's instructions (NEB). The plasmid was transformed into NEB 10β cells and

verified by Sanger sequencing prior to transformation into *Chlamydia trachomatis*. To generate

the complementation strain, the *ahpC* gene was amplified using primers listed in Table S1 and

fused with the *ahpC* knockdown construct (i.e., pBOMBL12CRia (*ahpC*)) digested with the Fast

Digest restriction enzyme SalI (Thermo Fisher) and alkaline phosphatase-treated, using the HiFi

reaction as mentioned above.

Chlamydial transformation

Chlamydial transformations were performed using a protocol described previously, with some

modifications (Mueller et al., 2017). One day before transformation, 1×10^6 McCoy cells were

seeded in one 6-well plate, and two wells were used per plasmid transformation. Briefly, for

each well of the 6-well plate, 2 µg of sequenced verified plasmids were incubated with 2.5×10^6

C. trachomatis serovar L2 without plasmid (-pL2) EBs in 50 µL Tris-CaCl$_2$ (10 mM Tris, 50 mM

CaCl2, pH 7.4) at room temperature for 30 min. McCoy cells were washed with 2 mL Hank's

Balanced Salt Solution (HBSS; Gibco), and 1 mL HBSS was added back into each well. 1 mL of

HBSS was added to each transformant mixture, and one well of a 6-well plate was infected

using this transformation solution. Cells were centrifuged at 400 × g for 15 min at room

temperature followed by 15 min incubation at 37°C. HBSS was aspirated and replaced with

antibiotic-free DMEM. At 8 hpi, 1 µg/mL of cycloheximide and 1 or 2 U/mL of penicillin G or 500

µg/mL spectinomycin were added to the culture media. The infection was passaged every 48 h

until a population of penicillin or spectinomycin resistant, green fluorescent protein (GFP)

positive *C. trachomatis* was established. The chlamydial transformants were then serially diluted

to isolate clonal populations. These isolated populations were further expanded and frozen at -

80^{0}C in a sucrose phosphate solution (2SP). To verify plasmid sequences, DNA was harvested

from infected cultures using the DNeasy kit (Qiagen) and transformed into NEB 10β for plasmid

propagation. Isolated plasmids were then verified by restriction digest and Sanger sequencing.

Inclusion forming unit assay

Inclusion forming unit assay was performed to determine the infectious progeny (number of

EBs) from a primary infection based on inclusions formed in a secondary infection. *C.*

trachomatis transformants were infected into HeLa cells and induced or not with 1 nM aTc at 10

hpi. At 24 and 48 hpi, samples were harvested by scraping three wells of a 24-well plate in 2

sucrose-phosphate (2SP) solution and lysed via a single freeze-thaw cycle, serially diluted, and

used to infect a fresh HeLa cell monolayer and allowed to grow for 24 h. Samples were fixed

with methanol, stained with a goat antibody specific to *C. trachomatis* major outer membrane

protein (MOMP; Meridian Biosciences, Memphis TN) followed by staining with donkey antigoat

Alexa Fluor 594-conjugated secondary antibody (Invitrogen) and titers were enumerated using a

20x lens objective. All experiments were performed three times for three biological replicates.

Induced values were expressed as a percentage of the uninduced values, which was

considered as 100%.

Immunofluorescence assay

HeLa cells were cultured on glass coverslips in 24-well tissue cultures plates at 2 x 10^{5}

cells/well, infected with the relevant strains, and, at 10 hpi, samples were induced or not with 1

nM aTc. At 24 hpi, samples were fixed and permeabilized using 100% methanol. Organisms

were stained with anti-MOMP (Meridian Biosciences) primary antibody for all the strains and

primary mouse anti-Cpf1 (dCas12) (Sigma-Millipore) for knockdown or complementation

samples. Donkey anti-goat Alexa Fluor 594-conjugated secondary antibody (Invitrogen) was

used to visualize *Chlamydia* in all the samples, and donkey anti-mouse Alexa Fluor 488-

conjugated secondary antibody (Invitrogen) was used for dCas12 expression. DAPI (Invitrogen)

was used for visualization of host and bacterial cell DNA. These stained coverslips were

mounted on glass slides using ProLong glass antifade mounting media (Invitrogen) and imaged

using a 100x lens objective on a Zeiss Axioimager Z.2 equipped with Apotome.2 optical

sectioning hardware and X-Cite Series 120PC illumination lamp using a 2MP Axiocam 506

monochrome camera.

Nucleic acid extraction and RT-qPCR

HeLa cells were seeded in 6-well tissue culture plates, infected with the *C. trachomatis*

transformants, and induced or not at 10 hpi with 1 nM aTc. For each condition, triplicate wells

were used for simultaneous harvest of RNA (for transcript analysis by RT-qPCR), gDNA (to

normalize RT-qPCR data and quantification of gDNA), and IFA (to verify the morphological

changes in samples in the tested conditions). For RNA extraction, cells were rinsed with DPBS

twice and lysed with 1 mL TRIzol (Invitrogen/Thermo Fisher) per well as per manufacturer's

instructions. 200 µL of chloroform was added to extract the aqueous layer containing total RNA,

which was precipitated with isopropanol. A total of 10 µg of purified RNA was treated with

TURBO DNase (Invitrogen/Thermo Fisher) according to the manufacturer's instructions to

remove DNA contamination. DNA-free RNA was used for cDNA synthesis using random

nonamers (N9; NEB) and SuperScript III reverse transcriptase (Invitrogen/Thermo Fisher)

following the manufacturer's instructions. cDNA samples were diluted 10-fold with molecular

biology-grade water and stored at -80°C. A total of 2.5 µL of each diluted cDNA sample was

used per well of a 96-well qPCR plate. For each of three biological replicates, each sample was

analyzed in triplicate on a QuantStudio 3 system (Applied Biosystems/Thermo Fisher) using the

standard amplification cycle with melting curve analysis. For gDNA, one well of a 6-well plate

per condition was scraped in 500 μL DPBS, split in half (i.e., 250 μL), and frozen at -80°C. Each

sample was then thawed and frozen twice more for a total of three freeze/thaw cycles, and

gDNA was extracted using the DNeasy DNA extraction kit (Qiagen) according to the

manufacturer's guidelines. The isolated gDNA was quantified and diluted down to 5 ng/mL prior

to use in quantitative PCR (qPCR). A total of 2.5 μL of each diluted gDNA sample was mixed

with 10 μL of PowerUp SYBR green master mix in a 96-well qPCR plate and was analyzed on a

QuantStudio 3 system (Applied Biosystems). Each sample from each biological replicate was

tested in triplicate. For each primer set used, a standard curve of gDNA was generated against

purified *C. trachomatis* L2 genomic DNA, and the cDNA levels were normalized to gDNA levels

or 16S rRNA for analysis. All experiments were performed three times for three biological

replicates. At the same time, morphological differences were monitored in the IFA control, with

samples fixed with methanol at the time of harvesting of RNA/gDNA. Staining and imaging were

performed as described above.

Viability assay

Cell viability assays were performed using PrestoBlue (Invitrogen/Thermo Fisher). HeLa cells

were seeded in 96 well plates and were infected or not with the indicated chlamydial strains

(e.g., pBOMBDC.ev). Samples were treated or not as per the experimental parameters (e.g.,

different concentrations of oxidizing agents or scavengers). Medium without cells was used as

the blank control. At the end point, 10% PrestoBlue (v/v) was added in the wells and incubated

at 37^0 C for 30 min while protecting the plate from light. Fluorescence was read at excitation 560

nm and emission 590 nm using an Infinite M200 Pro (Tecan). The treated values were

expressed as a percentage of the untreated values, which was considered as 100%. All

experiments were performed three times for three biological replicates.

Oxidizing agents' susceptibility testing

Susceptibility of host cells and chlamydiae to different oxidizing agents was tested in HeLa cells

infected with density gradient-purified *Chlamydia trachomatis* 434/Bu EBs or *C. trachomatis*

transformants. For *C. trachomatis* transformants, expression of the construct was induced or not

with 1 nM aTc at 10 hpi. At 16 hpi, different concentrations of oxidizing agents were added, and

samples were incubated at 37^0C for 30 min. These samples were washed three times with

HBSS, fresh DMEM media was added in the wells, and cultures were allowed to grow until 24

hpi. At 24 hpi, inclusion forming unit assays and immunofluorescence analysis were performed

as described above.

ROS scavenging by chemical compounds

The scavenging capacity of different ROS scavengers was tested using HeLa cells infected with

Chlamydia trachomatis 434/Bu EBs or the *ahpC* KD strain. To examine the protective effect of

scavengers in the absence or presence of oxidative stress, *Chlamydia trachomatis* 434/Bu EBs

infected HeLa cells were used. In these samples, scavengers were added or not at 10 hpi in

respective wells. At 16 hpi, wells were washed three times with HBSS, and fresh DMEM

containing 500 µM H_2O_2 or not was added. After 30 min, wells were washed three times with

HBSS and fresh DMEM containing scavengers or not was added back in the respective wells,

which were incubated until 24 hpi prior to collecting samples for IFU assay and IFA analysis.

The effect of these scavengers was further tested in *ahpC* KD. HeLa cells were infected with

ahpC KD, scavengers were added or not at 9.5 hpi before induction at 10 hpi with 1nM aTc. At

14 hpi, scavengers were added again in the respective wells, and, at 24 hpi, samples were

collected for IFU assay and IFA.

ROS detection by CellROX Deep Red

Intracellular ROS levels were measured in uninfected and *ahpC* KD-infected HeLa cells using

CellROX Deep Red dye (Invitrogen/Thermo Fisher). Samples were induced or not at 10 hpi with

1 nM aTc. Medium without cells was used as the blank control. At 16 hpi, samples were treated

or not with 62.5 μM or 1 mM TBHP for 30 min. At the end point, media was removed, samples

were washed three times with DPBS and incubated with CellROX Deep Red dye for 30 min in

the dark at 37^0C. Fluorescence was read at excitation 644 nm and emission 665 nm using an

Infinite M200 Pro (Tecan). All experiments were performed three times for three biological

replicates.

ACKNOWLEDGMENTS

We thank Dr. H. Caldwell (NIH/NIAID) for providing eukaryotic cell lines and Dr. I. Clarke

(University of Southampton) for providing the plasmid less strain of *C. trachomatis* serovar L2.

We thank Dr. Elizabeth A. Rucks for critical feedback. We thank the members of the

Rucks/Ouellette research group for thoughtful discussion of the material presented. Funding for

this work was provided by the National Institutes of Health (NIH/NIAID) grants R01AI170688

and R21AI178150 to SPO.

Figure legends:

Fig. 1 Alterations in redox status of key proteins regulate and drive chlamydial differentiation.

(A) Key characteristics of chlamydial developmental forms. (B) Hypothetical model for triggering

secondary differentiation through oxidative stress (black angled line). Increasing oxidation of

critical protein(s) may lead to earlier differentiation whereas maintaining a reducing environment

may delay differentiation. (C) Schematic representation of experimental model for triggering

secondary differentiation through altered activity of AhpC. *ahpC* knockdown may lead to earlier

differentiation while overexpression of *ahpC* may delay differentiation.

Fig. 2 Overexpression of *ahpC* affects chlamydial growth and differentiation. (A) Transcriptional

analysis of *ahpC* in *ahpC* overexpression (*ahpC*) and empty vector (EV) control using RT-qPCR

following induction at 10 hpi with 1 nM aTc. RNA and genomic DNA (gDNA) were harvested at

14 and 24 hpi and processed as mentioned in materials and methods. Data are presented as a

ratio of cDNA to gDNA plotted on a log scale. ***$p< 0.0001$ vs uninduced sample by using two-

way ANOVA. Data represent three biological replicates. (B) Immunofluorescence assay (IFA) of

ahpC and EV at 24 hpi. Construct expression was induced or not at 10 hpi with 1 nM aTc, and

samples were fixed with methanol at 24 hpi then stained for major outer membrane protein

(MOMP - red) and DAPI (blue) to label DNA. Scale bars = 2 μm. Images were captured using a

Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Representative images of three

biological replicates are shown. (C) Impact of *ahpC* overexpression on inclusion area. Inclusion

area of *ahpC* overexpression and EV strains was measured using ImageJ. Experimental

conditions were the same as mentioned in section (B). The area of 50 inclusions was measured

per condition for each sample. ***p< 0.001 vs uninduced sample by using ordinary one-way

ANOVA. Data were collected from three biological replicates. (D) IFU assay of *ahpC*

overexpression and empty vector control. Expression of the construct was induced or not at 10

hpi, and samples were harvested at 24 or 48 hpi for reinfection and enumeration. IFUs were

calculated as the percentage of uninduced samples. **p< 0.001 vs uninduced sample by using

multiple paired t test. Data represent three biological replicates. (E) Quantification of genomic

DNA (gDNA) determined by qPCR in *ahpC* overexpression and empty vector control. Construct

expression was induced or not at 10 hpi with 1 nM aTc, gDNA was harvested at 14 and 24 hpi,

and ng gDNA were plotted on a log scale. ***p< 0.0001 vs uninduced sample by using two-way

ANOVA. Data represent three biological replicates.

Fig. 3 Higher expression of *ahpC* provides resistance to peroxides in *Chlamydia.* IFA of *ahpC*

(A) or EV (C) exposed to oxidizing agents. Construct expression was induced or not at 10 hpi

with 1 nM aTc, and samples were treated with three different concentrations of hydrogen

peroxide (H_2O_2) at 16 hpi for 30 min then fixed with methanol at 24 hpi and stained and imaged

as described in the legend of Fig. 2B. Representative images from three biological replicates

are shown. IFU analysis of *ahpC* (B) or EV (D) following treatment with oxidizing agents, CHP-

Cumene hydroperoxide, H_2O_2-Hydrogen peroxide, TBHP-Tert-butyl hydroperoxide, and PN-

Peroxynitrite. Samples were processed as described for (A) and (C), and IFUs harvested at 24

hpi. IFUs of treated samples were compared with respective untreated controls. ***p< 0.0001 vs

untreated sample by using two-way ANOVA. Data represent three biological replicates.

Fig. 4 Reduced levels of AhpC negatively impact chlamydial growth. (A) Transcriptional analysis

of *ahpC* in knockdown (*ahpC* KD) and non-target (NT) control using RT-qPCR following

induction at 10 hpi with 1 nM aTc. RNA and gDNA were harvested at 14 and 24 hpi. Quantified

cDNA was normalized to gDNA, and values were plotted on a log scale. ***p< 0.0001 vs

uninduced sample by using two-way ANOVA. Data represent three biological replicates. (B) IFA

was performed to assess inclusion size and morphology using the same induction conditions as

in section (A). At 24 hpi, cells were fixed with methanol and stained using primary antibodies to

major outer membrane protein (MOMP), Cpf1 (dCas12), and DAPI. All images were acquired on

Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Bars, 2 μm. Representative images

of three biological replicates are shown. (C) Quantification of genomic DNA (gDNA) determined

by qPCR in *ahpC* KD and NT strains. dCas12 expression was induced or not at 10 hpi, and

gDNA was harvested at 14 and 24 hpi and plotted on a log scale. ***p< 0.0001 vs uninduced

sample by using two-way ANOVA. Data represent three biological replicates. (D) IFU titers

following induction at 10 hpi with 1 nM aTc. IFUs were counted from 24 and 48 hpi samples and

calculated as percentage of uninduced samples. ***p< 0.0001, **p< 0.001 vs uninduced sample

by using multiple paired t test. Data represent three biological replicates. (E) Intracellular ROS

levels were measured to investigate the function of AhpC in reducing ROS. HeLa cells were

infected or not with *ahpC* knockdown, and knockdown was induced or not at 10 hpi with 1 nM

aTc. At 24 hpi, samples were washed with DPBS and incubated with CellROX Deep red dye for

30 min in dark. ROS levels were measured at wavelengths of 640nm (excitation) and 665nm

(emission). **p< 0.001, *p< 0.01 vs uninduced sample by using two-way ANOVA. Data

represent three biological replicates.

Fig. 5 *Chlamydia* is hypersensitive to oxidizing agents in *ahpC* knockdown condition. IFA of

ahpC KD (A) or NT (C) treated with 62.5 μM H_2O_2. dCas12 expression was induced or not at 10

hpi with 1 nM aTc, treated or not with H_2O_2 at 16 hpi for 30 min, and allowed to grow until 24 hpi.

Coverslips were fixed with methanol at 24 hpi and stained major outer membrane protein

(MOMP), Cpf1 (dCas12), and DAPI. Scale bars = 2 μm. Images were captured using a Zeiss

Axio Imager Z.2 with Apotome2 at 100x magnification. Representative images from three

biological replicates are shown. IFU analysis of *ahpC* KD (B) or NT (D) following treatment with

oxidizing agents, CHP-Cumene hydroperoxide, H_2O_2-Hydrogen peroxide, TBHP-Tert-butyl

hydroperoxide, or PN-Peroxynitrite. dCas12 expression was induced or not, and samples were

treated or not as mentioned in the legend of Fig. 3B. IFUs of treated samples were calculated as

percentage of respective untreated samples. ***$p< 0.0001$ vs untreated sample by using two-

way ANOVA. Data represent three biological replicates.

Fig. 6 Complementation of the phenotypes observed in the *ahpC* knockdown. (A) Confirmation

of complementation (comp) of *ahpC* knockdown by RT-qPCR. Samples were processed and

quantified as mentioned previously in the legend of Fig. 4A. Values were plotted on a log scale.

*$p< 0.01$ vs uninduced sample using ordinary one-way ANOVA. Data represent three biological

replicates. (B) IFA of comp strain was performed at 24 hpi, and staining and imaging were

performed as mentioned in the legend of Fig. 4B. Scale bars, 2 μm. Representative images from

three biological replicates are shown. (C) Genomic DNA quantitation was performed by qPCR.

Construct expression was induced or not at 10 hpi, and gDNA was harvested at 14 and 24 hpi

and plotted on a log scale. Statistical analysis was calculated using ordinary one-way ANOVA.

Data represent three biological replicates. (D) IFU analysis of comp strain. Statistical analysis

was calculated using multiple paired t test. Data represent three biological replicates. (E) IFA of

comp strain following treatment with 62.5 μM H_2O_2. Samples were treated, stained, and images

acquired as mentioned in the legend of Fig. 5A. Scale bars = 2 μm. Representative images from

three biological replicates are shown. (F) IFU analysis of comp strain following treatment with

oxidizing agents. Experiments were performed as mentioned in the legend of Fig. 5B. IFUs were

calculated as percentage of respective untreated samples. ***$p< 0.0001$ vs untreated sample by

using two-way ANOVA. Data represent three biological replicates. (G) *ahpC* knockdown growth

defect rescued by ROS scavengers. IFU analysis of *ahpC* knockdown treated with or without

scavengers, α-Tocopherol (100 µM) and DMTU (10 mM), as mentioned in materials and

methods. IFUs were calculated as a percentage of the untreated, uninduced sample. ***$p<$

0.0001 vs untreated, uninduced or induced sample by using two-way ANOVA. Data represent

three biological replicates. (H) IFA of *ahpC* knockdown treated with or without scavengers.

Experimental conditions were similar as in section (G). Staining and imaging were performed as

mentioned in Fig. 5A. Representative images from three biological replicates are shown.

Fig. 7 Effects of *ahpC* knockdown/overexpression on chlamydial developmental cycle

progression. RT-qPCR analysis of late cycle genes (*hctA*, *hctB*, *omcB*, *tsp*, and *glgA*) in (A)

ahpC knockdown, (C) complementation, and (D) *ahpC* overexpression strain. Experimental

conditions were the same as mentioned in the legend of Fig. 4A. Quantified cDNA was

normalized to gDNA, and values were plotted on a log scale. ***$p< 0.0001$, **$p< 0.001$, *$p< 0.01$

vs uninduced sample by using two-way ANOVA. Data represent three biological replicates. (B)

One-step growth curve of *ahpC* knockdown. Samples were induced or not with 1 nM aTc at 10

hpi and harvested at 16, 18, 20, 22, and 24 hpi. IFUs recovered are displayed as log10 values.

***$p< 0.0001$ vs uninduced sample by using two-way ANOVA. Data represent three biological

replicates.

Fig. 8 Altering the activity of AhpC in *Ctr* L2 impacts its developmental cycle progression. (Top)

In *Chlamydia*, secondary differentiation is asynchronous and RBs divide through an asymmetric

budding mechanism. In such conditions, either the mother or daughter cell may inherit more

oxidized proteins (represented by a darker shade), which can then impact whether a given RB

will divide again or undergo secondary differentiation. (Bottom) The black dots represent EBs,

the orange circles show RBs. The developmental cycle of wild-type *C. trachomatis* (Ctr L2) is

shown. In *ahpC* KD, highly oxidized conditions lead some RBs to cross the oxidative threshold

sooner, allowing activation of late genes and secondary differentiation earlier than other RBs. In

ahpC overexpression, a reducing environment results in a delay in achieving the oxidative

threshold, thus allowing RBs to continue to divide before committing to secondary

differentiation.

 Supplementary Information

 Table S1 List of plasmids, strains, and primers used in this study.

 Fig. S1 Viability assay of uninfected or infected HeLa cells treated with oxidizing agents H_2O_2

 (A), CHP (B), TBHP (C), and PN (D). HeLa cells were infected or not with empty vector control,

 pBOMBDC.ev (EV), and treated or not with different concentrations of oxidizing agents at 16 hpi

 for 30 min. At 24 hpi, an end point viability assay was performed using PrestoBlue as mentioned

 in materials and methods. The treated values were expressed as a percentage of the untreated

 values, which were considered as 100%. Data represent three biological replicates.

 Fig. S2 Response of *C. trachomatis* L2 against oxidizing agents. (A) IFA of wild-type *Ctr* L2

 exposed to oxidizing agents. Oxidizing agents' treatment, staining, and imaging were performed

 as mentioned in materials and methods. Representative images from three biological replicates

 are shown. (B) IFU analysis of *Ctr* L2 post-exposure with oxidizing agents. Conditions were

 same as in section (A), and IFUs were harvested at 24 hpi. IFUs were calculated as percentage

 of untreated samples. ***$p<$ 0.0001 vs untreated sample by using one way ANOVA. Data

 represent three biological replicates.

 Fig. S3 Overexpression of *ahpC* provides resistance to peroxides in *Chlamydia.* IFA of *ahpC* (A)

 or EV (B) exposed to oxidizing agents (CHP, TBHP, and PN). Experiments were performed as

 mentioned in the legend of Fig. 3A. Representative images from three biological replicates are

 shown.

 Fig. S4 *Chlamydia* is hypersensitive to oxidizing agents as a result of reduced levels of *ahpC*.

 IFA of *ahpC* KD (A), NT (B), comp (C) treated with CHP, TBHP, and PN. Experimental

 conditions were the same as mentioned in the legend of Fig. 5A. Representative images from

 three biological replicates are shown.

 Fig. S5 Rescue of oxidative stress phenotype by ROS scavengers in *C. trachomatis* L2. (A) IFA

 of wild-type *Ctr* L2 incubated or not with scavengers, α-Tocopherol (100 μM) and DMTU (10

 mM), and treated or not with 500 μM H_2O_2 as mentioned in materials and methods.

849 Representative images from three biological replicates are shown. (B) IFU analysis of *Ctr* L2

850 grown under the same conditions as mentioned in the legend of Fig. 6G. IFUs were calculated

851 as percentage of untreated samples. ***$p < 0.0001$, statistical analysis was performed using

852 ordinary one-way ANOVA. H_2O_2 treated sample and samples incubated with scavengers only

853 were compared to the untreated control. Samples treated with H_2O_2 and incubated with

854 scavengers were compared with sample treated with H_2O_2. Data represent three biological

855 replicates. Only significant differences are noted.

856 **Fig. S6** Effect of penicillin treatment during *ahpC* knockdown. (A) IFA was performed to assess

857 inclusion size and morphology of *ahpC* KD-spec following induction (1 nM aTc) and penicillin

858 treatment (1 U/mL) at 10 hpi. At 24 hpi, cells were fixed with methanol and stained using

859 primary antibodies to major outer membrane protein (MOMP), Cpf1 (dCas12), and DAPI. All

860 images were acquired on Zeiss Axio Imager Z.2 with Apotome2 at 100x magnification. Bars, 2

861 μm. Representative images of three biological replicates are shown. Transcriptional analysis of

862 (B) *ahpC*, (C) *hctA*, and (D) *hctB* in *ahpC* KD-spec using RT-qPCR using the same conditions

863 as in section (A). RNA samples were harvested at 16 and 24 hpi. Quantified cDNA was

864 normalized to 16SrRNA, and values were plotted on a log scale. ***$p < 0.0001$, **$p < 0.001$, *$p <$

865 0.01 vs uninduced sample by using two-way ANOVA. Data represent three biological replicates.

867 **REFERENCES**

868 Abdelrahman, Y., Ouellette, S. P., Belland, R. J., & Cox, J. V. (2016). Polarized Cell Division of
869   Chlamydia trachomatis. *PLoS Pathog*, *12*(8), e1005822.
870   https://doi.org/10.1371/journal.ppat.1005822
871 Abdelrahman, Y. M., & Belland, R. J. (2005). The chlamydial developmental cycle. *FEMS*
872   *Microbiol Rev*, *29*(5), 949-959. https://doi.org/10.1016/j.femsre.2005.03.002
873 Abdul-Sater, A. A., Said-Sadier, N., Lam, V. M., Singh, B., Pettengill, M. A., Soares, F., Tattoli, I.,
874   Lipinski, S., Girardin, S. E., Rosenstiel, P., & Ojcius, D. M. (2010). Enhancement of reactive
875   oxygen species production and chlamydial infection by the mitochondrial Nod-like family
876   member NLRX1. *J Biol Chem*, *285*(53), 41637-41645.
877   https://doi.org/10.1074/jbc.M110.137885

Antelmann, H., Engelmann, S., Schmid, R., & Hecker, M. (1996). General and oxidative stress
responses in Bacillus subtilis: cloning, expression, and mutation of the alkyl
hydroperoxide reductase operon. *J Bacteriol*, *178*(22), 6571-6578.
https://doi.org/10.1128/jb.178.22.6571-6578.1996

Barbour, A. G., Amano, K., Hackstadt, T., Perry, L., & Caldwell, H. D. (1982). Chlamydia
trachomatis has penicillin-binding proteins but not detectable muramic acid. *J Bacteriol*,
151(1), 420-428. https://doi.org/10.1128/jb.151.1.420-428.1982

Belland, R. J., Zhong, G., Crane, D. D., Hogan, D., Sturdevant, D., Sharma, J., Beatty, W. L., &
Caldwell, H. D. (2003). Genomic transcriptional profiling of the developmental cycle of
Chlamydia trachomatis. *Proc Natl Acad Sci U S A*, *100*(14), 8478-8483.
https://doi.org/10.1073/pnas.1331135100

Betts-Hampikian, H. J., & Fields, K. A. (2011). Disulfide bonding within components of the
Chlamydia type III secretion apparatus correlates with development. *J Bacteriol*, *193*(24),
6950-6959. https://doi.org/10.1128/JB.05163-11

Boncompain, G., Muller, C., Meas-Yedid, V., Schmitt-Kopplin, P., Lazarow, P. B., & Subtil, A.
(2014). The intracellular bacteria Chlamydia hijack peroxisomes and utilize their
enzymatic capacity to produce bacteria-specific phospholipids. *PLoS One*, *9*(1), e86196.
https://doi.org/10.1371/journal.pone.0086196

Boncompain, G., Schneider, B., Delevoye, C., Kellermann, O., Dautry-Varsat, A., & Subtil, A.
(2010). Production of reactive oxygen species is turned on and rapidly shut down in
epithelial cells infected with Chlamydia trachomatis. *Infect Immun*, *78*(1), 80-87.
https://doi.org/10.1128/IAI.00725-09

Brinkworth, A. J., Wildung, M. R., & Carabeo, R. A. (2018). Genomewide Transcriptional
Responses of Iron-Starved Chlamydia trachomatis Reveal Prioritization of Metabolic
Precursor Synthesis over Protein Translation. *mSystems*, *3*(1).
https://doi.org/10.1128/mSystems.00184-17

Caldwell, H. D., Kromhout, J., & Schachter, J. (1981). Purification and partial characterization of
the major outer membrane protein of Chlamydia trachomatis. *Infect Immun*, *31*(3),
1161-1176. https://doi.org/10.1128/iai.31.3.1161-1176.1981

Chiarelli, T. J., Grieshaber, N. A., Omsland, A., Remien, C. H., & Grieshaber, S. S. (2020). Single-
Inclusion Kinetics of Chlamydia trachomatis Development. *mSystems*, *5*(5).
https://doi.org/10.1128/mSystems.00689-20

Clifton, D. R., Dooley, C. A., Grieshaber, S. S., Carabeo, R. A., Fields, K. A., & Hackstadt, T. (2005).
Tyrosine phosphorylation of the chlamydial effector protein Tarp is species specific and
not required for recruitment of actin. *Infect Immun*, *73*(7), 3860-3868.
https://doi.org/10.1128/IAI.73.7.3860-3868.2005

Cosgrove, K., Coutts, G., Jonsson, I. M., Tarkowski, A., Kokai-Kun, J. F., Mond, J. J., & Foster, S. J.
(2007). Catalase (KatA) and alkyl hydroperoxide reductase (AhpC) have compensatory
roles in peroxide stress resistance and are required for survival, persistence, and nasal
colonization in Staphylococcus aureus. *J Bacteriol*, *189*(3), 1025-1035.
https://doi.org/10.1128/JB.01524-06

Cox, J. V., Abdelrahman, Y. M., & Ouellette, S. P. (2020). Penicillin-binding proteins regulate
multiple steps in the polarized cell division process of Chlamydia. *Sci Rep*, *10*(1), 12588.
https://doi.org/10.1038/s41598-020-69397-x

de Oliveira, M. A., Tairum, C. A., Netto, L. E. S., de Oliveira, A. L. P., Aleixo-Silva, R. L., Cabrera, V.
I. M., Breyer, C. A., & Dos Santos, M. C. (2021). Relevance of peroxiredoxins in
pathogenic microorganisms. *Appl Microbiol Biotechnol*, *105*(14-15), 5701-5717.
https://doi.org/10.1007/s00253-021-11360-5
Dip, P. V., Kamariah, N., Nartey, W., Beushausen, C., Kostyuchenko, V. A., Ng, T. S., Lok, S. M.,
Saw, W. G., Eisenhaber, F., Eisenhaber, B., & Gruber, G. (2014). Key roles of the
Escherichia coli AhpC C-terminus in assembly and catalysis of alkylhydroperoxide
reductase, an enzyme essential for the alleviation of oxidative stress. *Biochim Biophys*
Acta, *1837*(12), 1932-1943. https://doi.org/10.1016/j.bbabio.2014.08.007
Everett, K. D., & Hatch, T. P. (1995). Architecture of the cell envelope of Chlamydia psittaci 6BC. *J*
Bacteriol, *177*(4), 877-882. https://doi.org/10.1128/jb.177.4.877-882.1995
Ezraty, B., Gennaris, A., Barras, F., & Collet, J. F. (2017). Oxidative stress, protein damage and
repair   in   bacteria.   *Nat   Rev   Microbiol*,   *15*(7),   385-396.
https://doi.org/10.1038/nrmicro.2017.26
Fang,   F.   C.   (2011).   Antimicrobial   actions   of   reactive   oxygen   species.   *mBio*,   *2*(5).
https://doi.org/10.1128/mBio.00141-11
Feng, X., Guo, K., & Gao, H. (2020). Plasticity of the peroxidase AhpC links multiple substrates to
diverse disulfide-reducing pathways in Shewanella oneidensis. *J Biol Chem*, *295*(32),
11118-11130. https://doi.org/10.1074/jbc.RA120.014010
Gehre, L., Gorgette, O., Perrinet, S., Prevost, M. C., Ducatez, M., Giebel, A. M., Nelson, D. E., Ball,
S. G., & Subtil, A. (2016). Sequestration of host metabolism by an intracellular pathogen.
Elife, *5*, e12552. https://doi.org/10.7554/eLife.12552
Gerard, H. C., Freise, J., Wang, Z., Roberts, G., Rudy, D., Krauss-Opatz, B., Kohler, L., Zeidler, H.,
Schumacher,   H.   R.,   Whittum-Hudson,   J.   A.,   &   Hudson,   A.   P.   (2002).   Chlamydia
trachomatis genes whose products are related to energy metabolism are expressed
differentially   in   active   vs.   persistent   infection.   *Microbes Infect*,   *4*(1),   13-22.
https://doi.org/10.1016/s1286-4579(01)01504-0
Hatch, N. D., & Ouellette, S. P. (2023). Identification of the alternative sigma factor regulons of
Chlamydia trachomatis using multiplexed CRISPR interference. *mSphere*, *8*(5), e0039123.
https://doi.org/10.1128/msphere.00391-23
Hollander-Czytko, H., Grabowski, J., Sandorf, I., Weckermann, K., & Weiler, E. W. (2005).
Tocopherol content and activities of tyrosine aminotransferase and cystine lyase in
Arabidopsis   under   stress   conditions.   *J   Plant   Physiol*,   *162*(7),   767-770.
https://doi.org/10.1016/j.jplph.2005.04.019
Iliffe-Lee, E. R., & McClarty, G. (1999). Glucose metabolism in Chlamydia trachomatis: the
'energy   parasite'   hypothesis   revisited.   *Mol   Microbiol*,   *33*(1),   177-187.
https://doi.org/10.1046/j.1365-2958.1999.01464.x
Kiffin, R., Bandyopadhyay, U., & Cuervo, A. M. (2006). Oxidative stress and autophagy. *Antioxid*
Redox Signal, *8*(1-2), 152-162. https://doi.org/10.1089/ars.2006.8.152
Kimura, A., Yuhara, S., Ohtsubo, Y., Nagata, Y., & Tsuda, M. (2012). Suppression of pleiotropic
phenotypes of a Burkholderia multivorans fur mutant by oxyR mutation. *Microbiology*
(Reading), *158*(Pt 5), 1284-1293. https://doi.org/10.1099/mic.0.057372-0

Koshkin, A., Knudsen, G. M., & Ortiz De Montellano, P. R. (2004). Intermolecular interactions in
the AhpC/AhpD antioxidant defense system of Mycobacterium tuberculosis. *Arch*
Biochem Biophys, *427*(1), 41-47. https://doi.org/10.1016/j.abb.2004.04.017
Lee, J., Cox, J. V., & Ouellette, S. P. (2020). Critical Role for the Extended N Terminus of
Chlamydial MreB in Directing Its Membrane Association and Potential Interaction with
Divisome Proteins. *J Bacteriol*, *202*(9). https://doi.org/10.1128/JB.00034-20
Loprasert, S., Sallabhan, R., Whangsuk, W., & Mongkolsuk, S. (2003). Compensatory increase in
ahpC gene expression and its role in protecting Burkholderia pseudomallei against
reactive nitrogen intermediates. *Arch Microbiol*, *180*(6), 498-502.
https://doi.org/10.1007/s00203-003-0621-9
Mastronicola, D., Falabella, M., Testa, F., Pucillo, L. P., Teixeira, M., Sarti, P., Saraiva, L. M., &
Giuffre, A. (2014). Functional characterization of peroxiredoxins from the human
protozoan parasite Giardia intestinalis. *PLoS Negl Trop Dis*, *8*(1), e2631.
https://doi.org/10.1371/journal.pntd.0002631
Matsumoto, A., & Manire, G. P. (1970). Electron Microscopic Observations on the Fine Structure
of Cell Walls of Chlamydia psittaci. *J Bacteriol*, *104*(3), 1332-1337.
https://doi.org/10.1128/jb.104.3.1332-1337.1970
Mishra, S., & Imlay, J. (2012). Why do bacteria use so many enzymes to scavenge hydrogen
peroxide? *Arch Biochem Biophys*, *525*(2), 145-160.
https://doi.org/10.1016/j.abb.2012.04.014
Mongkolsuk, S., Whangsuk, W., Vattanaviboon, P., Loprasert, S., & Fuangthong, M. (2000). A
Xanthomonas alkyl hydroperoxide reductase subunit C (ahpC) mutant showed an altered
peroxide stress response and complex regulation of the compensatory response of
peroxide detoxification enzymes. *J Bacteriol*, *182*(23), 6845-6849.
https://doi.org/10.1128/JB.182.23.6845-6849.2000
Moulder, J. W. (1993). Why is Chlamydia sensitive to penicillin in the absence of peptidoglycan?
Infect Agents Dis, *2*(2), 87-99. https://www.ncbi.nlm.nih.gov/pubmed/8162358
Moulder, J. W., Colon, J. I., Ruda, J., & Zebovitz, M. M. (1956). The effect of penicillin on
multiplying and nonmultiplying populations of sensitive and resistant strains of feline
pneumonitis virus. *J Infect Dis*, *98*(3), 229-238. https://doi.org/10.1093/infdis/98.3.229
Mueller, K. E., Wolf, K., & Fields, K. A. (2017). Chlamydia trachomatis Transformation and Allelic
Exchange Mutagenesis. *Curr Protoc Microbiol*, *45*, 11A 13 11-11A 13 15.
https://doi.org/10.1002/cpmc.31
Mygind, P., Christiansen, G., Persson, K., & Birkelund, S. (1998). Analysis of the humoral immune
response to Chlamydia outer membrane protein 2. *Clin Diagn Lab Immunol*, *5*(3), 313-
318. https://doi.org/10.1128/CDLI.5.3.313-318.1998
Newhall, W. J. t. (1987). Biosynthesis and disulfide cross-linking of outer membrane components
during the growth cycle of Chlamydia trachomatis. *Infect Immun*, *55*(1), 162-168.
https://doi.org/10.1128/iai.55.1.162-168.1987
Ochsner, U. A., Vasil, M. L., Alsabbagh, E., Parvatiyar, K., & Hassett, D. J. (2000). Role of the
Pseudomonas aeruginosa oxyR-recG operon in oxidative stress defense and DNA repair:
OxyR-dependent regulation of katB-ankB, ahpB, and ahpC-ahpF. *J Bacteriol*, *182*(16),
4533-4544. https://doi.org/10.1128/JB.182.16.4533-4544.2000

Oh, E., & Jeon, B. (2014). Role of alkyl hydroperoxide reductase (AhpC) in the biofilm formation
of Campylobacter jejuni. *PLoS One*, *9*(1), e87312.
https://doi.org/10.1371/journal.pone.0087312

Ouellette, S. P. (2018). Feasibility of a Conditional Knockout System for Chlamydia Based on
CRISPR Interference. *Front Cell Infect Microbiol*, *8*, 59.
https://doi.org/10.3389/fcimb.2018.00059

Ouellette, S. P., Blay, E. A., Hatch, N. D., & Fisher-Marvin, L. A. (2021). CRISPR Interference To
Inducibly Repress Gene Expression in Chlamydia trachomatis. *Infect Immun*, *89*(7),
e0010821. https://doi.org/10.1128/IAI.00108-21

Ouellette, S. P., Fisher-Marvin, L. A., Harpring, M., Lee, J., Rucks, E. A., & Cox, J. V. (2022).
Localized cardiolipin synthesis is required for the assembly of MreB during the polarized
cell division of Chlamydia trachomatis. *PLoS Pathog*, *18*(9), e1010836.
https://doi.org/10.1371/journal.ppat.1010836

Ouellette, S. P., Hatch, T. P., AbdelRahman, Y. M., Rose, L. A., Belland, R. J., & Byrne, G. I. (2006).
Global transcriptional upregulation in the absence of increased translation in Chlamydia
during IFNgamma-mediated host cell tryptophan starvation. *Mol Microbiol*, *62*(5), 1387-
1401. https://doi.org/10.1111/j.1365-2958.2006.05465.x

Ouellette, S. P., Karimova, G., Subtil, A., & Ladant, D. (2012). Chlamydia co-opts the rod shape-
determining proteins MreB and Pbp2 for cell division. *Mol Microbiol*, *85*(1), 164-178.
https://doi.org/10.1111/j.1365-2958.2012.08100.x

Ouellette, S. P., Lee, J., & Cox, J. V. (2020). Division without Binary Fission: Cell Division in the
FtsZ-Less Chlamydia. *J Bacteriol*, *202*(17). https://doi.org/10.1128/JB.00252-20

Pan, S., Jensen, A. A., Wood, N. A., Henrichfreise, B., Brotz-Oesterhelt, H., Fisher, D. J., Sass, P., &
Ouellette, S. P. (2023). Molecular Characterization of the ClpC AAA+ ATPase in the
Biology of Chlamydia trachomatis. *mBio*, *14*(2), e0007523.
https://doi.org/10.1128/mbio.00075-23

Panzetta, M. E., Valdivia, R. H., & Saka, H. A. (2018). Chlamydia Persistence: A Survival Strategy
to Evade Antimicrobial Effects in-vitro and in-vivo. *Front Microbiol*, *9*, 3101.
https://doi.org/10.3389/fmicb.2018.03101

Parsonage, D., Karplus, P. A., & Poole, L. B. (2008). Substrate specificity and redox potential of
AhpC, a bacterial peroxiredoxin. *Proc Natl Acad Sci U S A*, *105*(24), 8209-8214.
https://doi.org/10.1073/pnas.0708308105

Pokorzynski, N. D., Brinkworth, A. J., & Carabeo, R. (2019). A bipartite iron-dependent
transcriptional regulation of the tryptophan salvage pathway in Chlamydia trachomatis.
Elife, *8*. https://doi.org/10.7554/eLife.42295

Poole, L. B., & Ellis, H. R. (1996). Flavin-dependent alkyl hydroperoxide reductase from
Salmonella typhimurium. 1. Purification and enzymatic activities of overexpressed AhpF
and AhpC proteins. *Biochemistry*, *35*(1), 56-64. https://doi.org/10.1021/bi951887s

Reuter, J., Otten, C., Jacquier, N., Lee, J., Mengin-Lecreulx, D., Lockener, I., Kluj, R., Mayer, C.,
Corona, F., Dannenberg, J., Aeby, S., Buhl, H., Greub, G., Vollmer, W., Ouellette, S. P.,
Schneider, T., & Henrichfreise, B. (2023). An NlpC/P60 protein catalyzes a key step in
peptidoglycan recycling at the intersection of energy recovery, cell division and immune
evasion in the intracellular pathogen Chlamydia trachomatis. *PLoS Pathog*, *19*(2),
e1011047. https://doi.org/10.1371/journal.ppat.1011047

Richard, D., Bartfai, R., Volz, J., Ralph, S. A., Muller, S., Stunnenberg, H. G., & Cowman, A. F. (2011). A genome-wide chromatin-associated nuclear peroxiredoxin from the malaria parasite Plasmodium falciparum. *J Biol Chem*, *286*(13), 11746-11755. https://doi.org/10.1074/jbc.M110.198499

Rusconi, B., & Greub, G. (2013). Discovery of catalases in members of the Chlamydiales order. *J Bacteriol*, *195*(16), 3543-3551. https://doi.org/10.1128/JB.00563-13

Seaver, L. C., & Imlay, J. A. (2001). Alkyl hydroperoxide reductase is the primary scavenger of endogenous hydrogen peroxide in Escherichia coli. *J Bacteriol*, *183*(24), 7173-7181. https://doi.org/10.1128/JB.183.24.7173-7181.2001

Sherman, D. R., Mdluli, K., Hickey, M. J., Barry, C. E., 3rd, & Stover, C. K. (1999). AhpC, oxidative stress and drug resistance in Mycobacterium tuberculosis. *Biofactors*, *10*(2-3), 211-217. https://doi.org/10.1002/biof.5520100219

Soules, K. R., Dmitriev, A., LaBrie, S. D., Dimond, Z. E., May, B. H., Johnson, D. K., Zhang, Y., Battaile, K. P., Lovell, S., & Hefty, P. S. (2020). Structural and ligand binding analyses of the periplasmic sensor domain of RsbU in Chlamydia trachomatis support a role in TCA cycle regulation. *Mol Microbiol*, *113*(1), 68-88. https://doi.org/10.1111/mmi.14401

Soules, K. R., LaBrie, S. D., May, B. H., & Hefty, P. S. (2020). Sigma 54-Regulated Transcription Is Associated with Membrane Reorganization and Type III Secretion Effectors during Conversion to Infectious Forms of Chlamydia trachomatis. *mBio*, *11*(5). https://doi.org/10.1128/mBio.01725-20

Staerck, C., Gastebois, A., Vandeputte, P., Calenda, A., Larcher, G., Gillmann, L., Papon, N., Bouchara, J. P., & Fleury, M. J. J. (2017). Microbial antioxidant defense enzymes. *Microb Pathog*, *110*, 56-65. https://doi.org/10.1016/j.micpath.2017.06.015

Storz, G., Jacobson, F. S., Tartaglia, L. A., Morgan, R. W., Silveira, L. A., & Ames, B. N. (1989). An alkyl hydroperoxide reductase induced by oxidative stress in Salmonella typhimurium and Escherichia coli: genetic characterization and cloning of ahp. *J Bacteriol*, *171*(4), 2049-2055. https://doi.org/10.1128/jb.171.4.2049-2055.1989

Sun, Z., Sun, Y., Li, Y., Luan, X., Chen, H., Wu, H., Peng, B., & Lu, C. (2020). Identification of HeLa cell proteins that interact with Chlamydia trachomatis glycogen synthase using yeast two-hybrid assays. *Mol Med Rep*, *21*(3), 1572-1580. https://doi.org/10.3892/mmr.2020.10947

Swoboda, A. R., Wood, N. A., Saery, E. A., Fisher, D. J., & Ouellette, S. P. (2023). The Periplasmic Tail-Specific Protease, Tsp, Is Essential for Secondary Differentiation in Chlamydia trachomatis. *J Bacteriol*, *205*(5), e0009923. https://doi.org/10.1128/jb.00099-23

Thompson, C. C., Griffiths, C., Nicod, S. S., Lowden, N. M., Wigneshweraraj, S., Fisher, D. J., & McClure, M. O. (2015). The Rsb Phosphoregulatory Network Controls Availability of the Primary Sigma Factor in Chlamydia trachomatis and Influences the Kinetics of Growth and Development. *PLoS Pathog*, *11*(8), e1005125. https://doi.org/10.1371/journal.ppat.1005125

Walch, M., Dotiwala, F., Mulik, S., Thiery, J., Kirchhausen, T., Clayberger, C., Krensky, A. M., Martinvalet, D., & Lieberman, J. (2015). Cytotoxic Cells Kill Intracellular Bacteria through Granulysin-Mediated Delivery of Granzymes. *Cell*, *161*(5), 1229. https://doi.org/10.1016/j.cell.2015.05.021

Wan, F., Feng, X., Yin, J., & Gao, H. (2021). Distinct H(2)O(2)-Scavenging System in Yersinia
pseudotuberculosis: KatG and AhpC Act Together to Scavenge Endogenous Hydrogen
Peroxide. *Front Microbiol*, *12*, 626874. https://doi.org/10.3389/fmicb.2021.626874

Wang, G., Conover, R. C., Benoit, S., Olczak, A. A., Olson, J. W., Johnson, M. K., & Maier, R. J.
(2004). Role of a bacterial organic hydroperoxide detoxification system in preventing
catalase inactivation. *J Biol Chem*, *279*(50), 51908-51914.
https://doi.org/10.1074/jbc.M408450200

Wang, X., Schwarzer, C., Hybiske, K., Machen, T. E., & Stephens, R. S. (2014). Developmental
stage oxidoreductive states of Chlamydia and infected host cells. *mBio*, *5*(6), e01924.
https://doi.org/10.1128/mBio.01924-14

Wang, Y., Kahane, S., Cutcliffe, L. T., Skilton, R. J., Lambden, P. R., & Clarke, I. N. (2011).
Development of a transformation system for Chlamydia trachomatis: restoration of
glycogen biosynthesis by acquisition of a plasmid shuttle vector. *PLoS Pathog*, *7*(9),
e1002258. https://doi.org/10.1371/journal.ppat.1002258

Wong, C. F., Shin, J., Subramanian Manimekalai, M. S., Saw, W. G., Yin, Z., Bhushan, S., Kumar, A.,
Ragunathan, P., & Grüber, G. (2017). AhpC of the mycobacterial antioxidant defense
system and its interaction with its reducing partner Thioredoxin-C. *Sci Rep*, *7*(1), 5159.
https://doi.org/10.1038/s41598-017-05354-5

Wood, N. A., Swoboda, A. R., Blocker, A. M., Fisher, D. J., & Ouellette, S. P. (2022). Tag-
Dependent Substrate Selection of ClpX Underlies Secondary Differentiation of Chlamydia
trachomatis. *mBio*, *13*(5), e0185822. https://doi.org/10.1128/mbio.01858-22

Yang, K. S., Kang, S. W., Woo, H. A., Hwang, S. C., Chae, H. Z., Kim, K., & Rhee, S. G. (2002).
Inactivation of human peroxiredoxin I during catalysis as the result of the oxidation of
the catalytic site cysteine to cysteine-sulfinic acid. *J Biol Chem*, *277*(41), 38029-38036.
https://doi.org/10.1074/jbc.M206626200

Zhang, B., Gu, H., Yang, Y., Bai, H., Zhao, C., Si, M., Su, T., & Shen, X. (2019). Molecular
Mechanisms of AhpC in Resistance to Oxidative Stress in Burkholderia thailandensis.
Front Microbiol, *10*, 1483. https://doi.org/10.3389/fmicb.2019.01483

Zuo, F., Yu, R., Khaskheli, G. B., Ma, H., Chen, L., Zeng, Z., Mao, A., & Chen, S. (2014).
Homologous overexpression of alkyl hydroperoxide reductase subunit C (ahpC) protects
Bifidobacterium longum strain NCC2705 from oxidative stress. *Res Microbiol*, *165*(7),
581-589. https://doi.org/10.1016/j.resmic.2014.05.040

1126