Interplay between a polerovirus and a closterovirus decreases aphid transmission of the polerovirus

Souheyla Khechmar [1,2], Quentin Chesnais [1,2], Claire Villeroy [1], Véronique Brault [1,2]*, Martin
Drucker [1,2]*

[1] SVQV, UMR 1131, INRAE Centre Grand Est, Colmar, France

[2] Université Strasbourg, Strasbourg, France

* Corresponding authors: martin.drucker@inrae.fr, veronique.brault@inrae.fr

Short title: Polerovirus-closterovirus interplay

Keywords: plant virus, virus co-infection, virus localization, aphid transmission, virus accumulation,
aphid behavior, plant-virus-virus-vector interactions.

ABSTRACT

Multi-infection of plant by viruses is very common and can change drastically infection parameters

such as virus accumulation, distribution and vector transmission. Sugar beet is an important crop that

is frequently co-infected by the polerovirus beet chlorosis virus (BChV) and the closterovirus beet

yellows virus (BYV), both vectored by the green peach aphid (*Myzus persicae*). These phloem-limited

viruses are acquired while aphids ingest phloem sap from infected plants. Here we found that co-

infection decreased transmission of BChV by ~50%, but had no impact on BYV transmission. The

drastic reduction of BChV transmission was not due to lower accumulation of BChV in co-infected

plants, nor to reduced phloem sap ingestion by aphids from these plants. Using the SABER-FISH *in*

situ hybridization technique on plants, we observed that 40% of the infected phloem cells were co-

infected and that co-infection caused redistribution of BYV in these cells. The BYV accumulation

pattern changed from distinct intracellular spherical inclusions in mono-infected cells to a diffuse form

in co-infected cells. There, BYV co-localized with BChV throughout the cytoplasm, indicative of

virus-virus interactions. We propose that BYV-BChV interactions could restrict BChV access to the

sieve tubes and reduce its accessibility for aphids and present a model of how co-infection could alter

BChV intracellular movement and/or phloem loading and reduce BChV transmission.

IMPORTANCE

Mixed viral infections in plants are understudied yet can have significant influences on disease

dynamics and virus transmission. We investigated how co-infection with two unrelated viruses, beet

chlorosis virus (BChV) and beet yellows virus (BYV), affects aphid transmission of the viruses in

sugar beet plants. We show that co-infection reduced BChV transmission by about 50% without

affecting BYV transmission, despite similar virus accumulation rates in co-infected and mono-infected

plants. Follow-up experiments examined the localization and intra-cellular distribution of the viruses,

leading to the discovery that co-infection caused a redistribution of BYV in the phloem vessels and

altered its repartition pattern within plant cells, suggesting virus-virus interactions. In conclusion, the

interplay between BChV and BYV affects the transmission of BChV but not BYV, possibly through

direct or indirect virus-virus interactions at the cellular level. Understanding these interactions could

be crucial for managing virus propagation in crops and preventing yield losses.

INTRODUCTION

Historically, studies on diseases in both plants and animals focused mainly on simple interactions

implicating only one host and one pathogen. However, in nature, a host is often subjected to a multitude

of pathogens simultaneously, a scenario referred to as co-infection, multi-infection or mixed infection,

and which might be the rule rather than the exception **(Abdullah et al., 2017; Dutt et al., 2022)**. The

development of molecular tools such as high throughput sequencing further highlighted the high

occurrence of mixed infections compared to mono-infection **(Dutt et al., 2022)**. Mixed infections can

be heterogeneous (*i.e.,* implying different pathogens such as bacteria, fungi or viruses), or

homogeneous (*i.e.,* with similar microorganisms) **(Devi et al., 2021)**. Multi-infection can occur through

the simultaneous inoculation of several pathogens or sequentially, causing co-infection or super-

infection, respectively. The outcome of a co-infection or super-infection will be affected by the intra-

host interactions between the different infectious agents as well as by their interaction with the host or,

in the case of vector-borne pathogens, with the vector **(Dutt et al., 2022)**.

Pathogen interactions, plant viruses in our case, within hosts can be synergistic, *i.e.,* the presence of

one pathogen facilitates the infection of one or more additional pathogens. For example, synergistic

interactions between tomato chlorosis virus (ToCV) and tomato infectious chlorosis virus (TICV) led

to an increased TICV accumulation in *Nicotiana benthamiana* **(Wintermantel et al., 2008)**.

Interactions between pathogens are considered neutral when they do not, or only marginally, interfere

with each other, as demonstrated for example for tomatos infected with three begomoviruses, tomato

yellow mottle virus, tomato leaf curl Sinaloa virus and tomato yellow leaf curl virus. There, the

accumulation of each virus was not strongly affected by the presence of the others although the triple

infection induced stronger symptoms **(Maliano et al., 2022)**. Finally, in antagonistic pathogen

interactions, the presence of one pathogen impacts the infection of the other pathogen negatively

(Abdullah et al., 2017; Dutt et al., 2022; Gautam et al., 2020; Syller & Grupa, 2016). An example

is the co-infection of two tobamoviruses, hibiscus latent Singapore virus (HLSV) and tobacco mosaic

virus (TMV), in *N. benthamiana* plants where HLSV levels decreased, compared to mono-infection,

whereas TMV levels remained almost unaltered **(Chen et al., 2012)**. As in this former example, and

unlike many synergistic interactions, antagonisms tend to occur mainly between closely related viruses,

inducing fitness costs for one or both competitors **(Syller & Grupa, 2016)**.

Virus-virus and virus-host interactions in a multi-infection context can impact different aspects of the

virus cycle, such as virus replication, virus movement and virus localization in plant cells or tissues, or

by interfering with plant defenses **(Mukasa et al., 2006; Singhal et al., 2021)**. The outcomes of the

viral multi-infection cannot only alter the fitness of co-existing viruses but have also a direct impact

on the fitness of the plant itself by affecting the severity of foliar symptoms, plant development, and

plant chemical and physiological profiles **(Moreno & López-Moya, 2020)**. Plant traits modified by

infection or multi-infection can have consequences on virus transmission because they can change the

nutritional quality of the plant or modify visual traits and production of volatiles that play important

roles in vector-plant interactions. These alterations of the multi-infected plants can therefore influence

vector behavior and subsequently virus transmission **(Mauck et al., 2012; Mauck et al., 2016;**

McLaughlin et al., 2022; Moreno & López-Moya, 2020). Studies on mixed viral infections with

regard to vector interaction and transmission are scarce. Ontiveros et al. reported that tomatos co-

infected with the begomovirus tomato yellow leaf curl virus (TYLCV) and the crinivirus ToCV were

more attractive for the whitefly vectors than ToCV mono-infected plants **(Ontiveros et al., 2022)**. Co-

infection with the reovirus raspberry latent virus and the closterovirus raspberry leaf mottle virus

changed aphid vector preferences for infected raspberry plants, but not their feeding behavior **(Lightle**

& Lee, 2014).

The viral and host factors mediating these synergistic or antagonistic effects have only been identified

in a few cases. For instance, the synergistic interaction between the crinivirus sweet potato feathery

mottle virus (SPCSV) and the potyvirus sweet potato chlorotic stunt virus (SPFMV) is mediated by the

SPCSV-encoded RNase3 protein **(Cuellar et al., 2009)**. RNase3 functions as a viral suppressor of

RNA silencing (VSR) that might target a specific host component by interfering with small-RNA

biogenesis **(Cuellar et al., 2015)**. Another example is the potyviral silencing suppressor HC-Pro that

increases the titer of potato leafroll virus (PLRV) in co-infection **(Savenkov & Valkonen, 2001)**. One

example on the host side is the salicylic acid-responsive gene PR-P6 whose expression correlates with

virus titer and symptoms severity in tomatos mono- or co-infected with TYLCV and ToCV **(Ontiveros**

et al., 2022).

Sugar beet (*Beta vulgaris*) is an economically important crop that is often co-infected in the field by

several viruses **(Hossain et al., 2021; Wintermantel, 2005)**, causing alone, or in combination, leaf

yellowing, yield reduction and low sugar content. Among the viruses detected are: the poleroviruses

beet western yellows virus (BWYV-USA) so far not reported in Europe, beet mild yellowing virus

(BMYV) and beet chlorosis virus (BChV), the closterovirus beet yellows virus (BYV) and a

potyvirus, beet mosaic virus (BtMV) all occurring worldwide **(Hossain et al., 2021; Stevens et al.,**

2004). The impact of multiple sugar beet infections on the severity of leaf yellowing has been studied
for BYV, BtMV and BWYV-USA **(Wintermantel, 2005)**. In this study, sugar beet lines were
inoculated with either one, two, or all three viruses. Faster appearance of symptoms and more severe
stunting were observed in mixed infections with BYV and BtMV, compared to mono-infections
(Wintermantel, 2005). Acceleration of symptom development and greater yield losses were observed
in sugar beets co-inoculated with BChV and BYV **(Hossain et al. 2021)**.

In this work, we focused on the co-infection of sugar beet with BYV and BChV. BYV is mainly
transmitted by the two aphid species *Myzus persicae* and *Aphis fabae* **(German-Retana et al., 1999)**.
The main vector species for BChV is *M. persicae* but this virus can also be transmitted efficiently by
Macrosiphum euphorbiae **(Kozłowska-Makulska et al., 2009; Limburg et al., 1997)**. BChV and
BYV use two different modes of transmission: circulative and persistent for BChV, and non-circulative
and semi-persistent for BYV. Viruses transmitted in the persistent and circulative mode are acquired
during vector feeding and pass through the insect's digestive tract to be released into the haemolymph.
From there, they reach the salivary glands where they accumulate and are inoculated, as a saliva
component, when aphids feed on new plants (Dáder et al., 2017). This transmission mode is
characterized by long phases of virus acquisition (hours to days) on the infected plant, and latency (the
time it takes for the virus to reach the aphid's salivary glands). The time required for aphids to inoculate
the virus is also long (several hours), and aphids remain viruliferous for the rest of their lives **(Bragard**
et al., 2013). In the case of semi-persistent transmission, the virus attaches to areas of the mouthparts
or anterior digestive tract. Maximum transmission is achieved after a few hours of acquisition phase,
and aphids transmit viruses for only a few hours to a few days **(Bragard et al., 2013; Brault et al.,**
2010; Jiménez et al., 2018). The two viruses differ also in their cellular tropism. BChV is restricted to
phloem tissues **(Boissinot et al., 2017; Esau & Hoefert, 1972; Mutterer et al., 1999)**. BYV is also a
phloem-limited virus, but can colonize epidermal and mesophyll cells at a late stage of infection **(Dolja,**
2003; Dolja & Koonin, 2013; Esau et al., 1967).

Here, we studied mixed viral infection of sugar beet and examined its effect on vector transmission.
Our results show that co-infection of sugar beet with BChV and BYV had no impact on BYV
transmission, but did decrease transmission of BChV by 50%. We present evidence that the decrease
in transmission is due to specific BYV-BChV interactions in co-infected cells and not to changes in
virus accumulation or vector-feeding behavior on co-infected plants.

MATERIALS AND METHODS

Plants, viruses and aphids

The virus-susceptible sugar beet variety "Auckland" was used in this study. Seeds were kindly

provided by the seed company SESVanderHave (Tienen, Belgium) via the Technical Institute of Sugar

Beet (ITB, Paris, France). One week after germination, plantlets were transplanted into individual pots

and grown in a climate chamber at 22 to 25°C with a day/night cycle of 16 h of light and 8 h of darkness.

The *Myzus persicae* biotype WMp2 (local code NL) was used in this study **(Reinink et al., 1989)**.

Aphids were maintained on Chinese cabbage (*Brassica rapa* subspc. *Pekinensis* var. Granaat) in a

growth chamber under controlled conditions at 20±1 °C with a day/night cycle of 16 h of light and 8 h

of darkness.

The BYV_BBRO_UK isolate used in this study was a kind gift from Prof Mark Varrelmann of the

Institut für Zuckerrübenforschung (IfZ, Göttingen, Germany) but was originally sampled in the UK

(Stevens et al., 1997). The isolate was maintained on sugar beet plants and propagated by aphid

transmission.

We used the BChV-2a isolate collected in East England (UK) with the GenBank accession number

AF352024 **(Hauser et al., 2002)**. BChV particles, prepared as described in **(Van den Heuvel et al.,**

1991), were maintained at -80 °C before being used as a virus source in an artificial medium for aphid

acquisition **(Bruyère et al., 1997)**.

For all experiments, we used the systemically infected third upper leaf at 21 days post-inoculation

(dpi).

Virus detection by DAS-ELISA

BChV and BYV were detected in leaves of sugar beets at 21 dpi by double-antibody sandwich enzyme-

linked immunosorbent assay (DAS-ELISA **(Adams & Clark, 1977)**) with turnip yellows virus

antibodies that recognize BChV as well (our observation) and beet yellows virus-specific antibodies,

respectively (Loewe Biochemica, Sauerlach, Germany). Samples were considered infected when the

$OD_{405\ nm}$ was twice the mean value of non-infected control plants (three technical replicates) plus three

times the standard deviation.

Establishing mixed infection in source plants

Non-viruliferous aphids were fed on BYV-infected plants, or purified BChV particles for 24 h, before

being transferred to 2-week-old healthy sugar beet plants for a 3-day inoculation access period. Aphids

were then removed manually. The third upper leaves (just below the two youngest leaves) were always

used as the virus sources. After aphid inoculation, the plants were maintained in a growth chamber

under controlled conditions (22-25 °C and 16/8 h light/dark photoperiod) for three weeks before being

analyzed by DAS-ELISA to verify infection.

Aphid transmission experiments

Aphids acquired viruses for 24 h from detached source leaves (three leaves from three independent

infected source plants that were placed on 1% agarose in petri dishes), followed by virus inoculation

feeding on 15 test plants using three aphids per plant during 72 h. Plants were treated three days later

with pirimicarb insecticide to kill aphids. After 3 weeks, plants were analyzed by DAS-ELISA, and

the transmission efficiency of each virus from co-infected plants was compared to that of

transmission of each virus from mono-infected plants.

Localization of viruses by SABER-FISH

For virus localization, we used signal amplification by exchange reaction fluorescent *in situ*

hybridization (SABER-FISH), a multiplexable nucleic acid-based detection method. It includes a signal

amplification step enabling highly sensitive target revelation. We used the protocol that was originally

developed for frozen retinal tissue as described by Kishi **(Kishi et al., 2019)** with some adaptations to

optimize the technique for plant tissues. For the detection of BYV and BChV, we designed 25 (BYV)

and 16 (BChV) primers with OligoMiner **(Beliveau et al., 2018)**. A TTT and a concatemer sequence

were added to the 3'-end of each primer (concatemer 27 for BChV and 26 for BYV, see **S1 Table**).

Concatemers were added to the 3'-end of the probes by *in vitro* primer exchange reaction (PER) to

obtain probes of 300-400 nucleotides length **(Kishi et al., 2019)**. The product of the PER reaction is a

probe consisting of two parts: a specific sequence that recognizes the target and the concatemer part

that enables target detection and amplification of the signal by binding a concatemer-specific

fluorescent imager oligonucleotide. After concatemer extension, probes were purified and concentrated

using MinElute PCR purification columns (Qiagen, Courtaboeuf, France). Probe concentrations after

purification were determined with a Nanodrop 2000 spectrometer (Thermo Fisher Scientific, Illkirch-

Grafenstedt, France) using ssDNA setting. A final concentration of 800 ng/µl of the probe was used for

hybridization.

To prepare paraffin-embedded tissues, sections of leaf samples were immersed in 10 ml ice-cold 4%

paraformaldehyde solution in PBS. Samples were fixed by applying a vacuum (~720 mm Hg)

overnight at 4 °C. After dehydration in a graded series of ethanol/water, followed by a graded series of

ethanol/xylene and xylene/paraffin, the samples were embedded in paraffin (Paraplast, Leica, Nanterre,

France). Sections of 10 µm were cut using a rotary microtome and immobilized on Superfrost+ slides.

After de-waxing and rehydration, the fixed tissue samples were processed for SABER-FISH as

described **(Kishi et al., 2019)**. Briefly, slides were first prehybridized in pre-hybridization buffer, then

hybridized with all PER probes simultaneously at 43 °C. This hybridization step was followed by a

short second hybridization step at 37 °C to bind the fluorescent imagers (with ATTO-565 or ATTO-

647N dyes conjugated to their 5' ends) to the probes. Slides were mounted in Fluoroshield medium

(Sigma, Saint-Quentin-Fallavier, France) and observed with a Zeiss Axio Imager M2 microscope

equipped with a Hamamatsu Orca-Flash 4.0LT black-and-white camera (Lordil, Lay-Saint-Christophe,

France). Images were acquired with a 5x, 20x or 40x objective and with Zeiss 43 HE or AHF Cy5 ET

bandpass filter sets for epifluorescence acquisition in the red or far red channels, respectively, or with

brightfield or DIC settings for transmission light acquisition. Images were processed with ZEN 2.5 or

Image J 1.54 software. Final figures were assembled with Microsoft PowerPoint or LibreOffice

Impress.

Relative virus quantification by one-step TaqMan RT-qPCR

Gene-specific primer pairs and TaqMan probes for BYV and BChV were designed in collaboration

with GEVES (Angers, France) using the Primer Express 3.0.1 primer design tool (Thermo Fisher

Scientific). Primer sets were designed to amplify between 75 to 200 nucleotides of the BYV and BChV

genomes **(Table 1)**. The specificity of the primer pairs was controlled. Each primer pair was evaluated

by a standard curve with six dilutions and three technical replicates of BChV- or BYV-infected plants.

Efficiency rates (E) of 95% (for BYV) and 94% (for BChV) were obtained with $R^2 = 0.99$ for both

viruses.

To select reference genes, we used a list of stably expressed plant genes previously established on

Arabidopsis thaliana infected by several viruses **(Zhang et al., 2023)**. Orthologous genes were

identified in *Beta vulgaris* using the KEGG database and primers compatible with TaqMan and SYBR

Green detection were designed using the IDT Primer Request tool available online

(https://eu.idtdna.com/pages/tools/primerquest). We tested the suitability and stability of the reference

genes by SYBR Green PCR. Of the nine tested reference genes, five were selected with an

amplification efficiency ranging from 90% to 99%, and with R^2 correlation values for the curves

ranging from 0.98 to 0.99 **(S2 Table)**. The expression stability of candidate genes was then verified on

five biological replicates per treatment (mono- or co-infected sugar beets) at 21 dpi. For each biological

replicate, three technical replicates were analyzed. Two hundred ng RNA were used to prepare

complementary DNA (cDNA) using M-MLV reverse transcriptase kit (Promega, Charbonnières-les-

Bains, France) and oligo(dT) nucleotides. qPCR for reference gene evaluation was performed using

Universal SYBR Green Supermix (Bio-Rad, Marnes-la-Coquette, France) with gene-specific primers

(20 μM) in a CFX96 Touch Deep Well Real-Time PCR detection system (Bio-Rad). Two μl of initial

cDNA were used as template in 10 μl qPCR reaction to evaluate the expression of reference genes. The

cycling conditions were 98 °C for 2 min, followed by 40 cycles of 15 sec at 98 °C, 20 sec at 65 °C,

with an increment ramp of 0.5 °C per sec. Data from these experiments were entered into geNorm,

(Vandesompele et al., 2002), Bestkeeper **(Pfaffl et al., 2004)** and Normfinder **(Andersen et al., 2004)**,

programs for gene stability validation **(Kozera & Rapacz, 2013; Liu et al., 2023)**.

Relative virus accumulation was measured on five biological replicates using AgPath-ID^{TM} One-step

RT-PCR Kit for TaqMan chemistry (Applied Biosystems by Thermo Fisher Scientific). The third upper

systemically infected leaf from each source plant was collected at 21 dpi. The whole leaf was

homogenized in liquid nitrogen. One hundred mg of ground tissue was used for total RNA extraction

using NucleoSpin RNA Plant Kit (Macherey-Nagel, Hoerdt, France). After spectrometric

quantification with a Nanodrop 2000, 30 ng of RNA were used in a 23 µL reaction volume. The cycling

conditions were 95 °C for 10 min, followed by 40 cycles of 15 sec at 95 °C, 1 min at 65 °C, using a

CFX384 Touch Real-Time PCR Detection System (Bio-Rad). The tubulin gene was used as a reference

gene and the relative accumulation was determined using the 2^{-DDCt} method **(Livak & Schmittgen,**

2001).

Table 1: Primer and probe sequences used for virus quantification by RT-qPCR.

Virus/sugar beet gene	Sequence (5'---3')	Primer orientation and Probe
BYV	5' - TAC TGT TCC AAA CCA GGT CCT TG - 3'	Forward
	5' - ROX - TTG CTT CTT TTT CAA CTC CAC CAC CCT GT - BHQ-2 - 3'	Probe
	5' - GTG CAA CGC AGT TCG AAA CTA A - 3'	Reverse

BChV	5' - GGG ACC ATG GCA CCA TCT T - 3'	Forward
	5' - VIC - TCC CTT ACC ACC GGA TAT TAC CCA ACT CCT - BHQ-1 - 3'	Probe
	5' - GTA ATC TGA CAG CTT TTT CTG AAG AGG - 3'	Reverse
Tub[a]	5' -GGCTTTCTTGCATTGGTACAC - 3'	Forward
	5' - FAM - CGAGGCTGAGAGCAACATGAACGA- BHQ -1 - 3'	Probe
	5' - CATCCTGGTACTGCTGGTATTC - 3'	Reverse

243   [a]tubulin

Aphid feeding behavior

We used the electrical penetration graph (EPG) system (Giga-8 DC-EPG amplifier, EPG Systems,
Wageningen, Netherlands) as described by Tjallingii **(Tjallingii, 1988)** to investigate the feeding
behavior of *M. persicae*. We created electrical circuits that each included one aphid and one plant by
tethering a thin gold wire (12.5 μm diameter and 2 cm long) on the insect's dorsum using conductive
silver glue (EPG Systems). To facilitate the tethering process, aphids were immobilized at the edge of
a pipette tip with negative pressure generated with a vacuum pump (model N86KN.18, KNF
Neuberger, Freiburg, Germany). Eight apterous aphids were connected to the Giga-8 DC-EPG
amplifier (EPG Systems) and each one was placed on the third upper leaf of an individual plant. The
circuit was closed by inserting a copper electrode into the plant substrate. The feeding behavior was
recorded for 8 h at a temperature of 20±1 °C and constant light inside a Faraday cage. Each plant and
aphid were used only once. Acquisition of the EPG waveforms were carried out with PROBE 3.5
software (EPG Systems). This software was also used to identify the different waveforms correlated
with the phases of *M. persicae* feeding behavior, as described **(Tjallingii & Hogen Esch, 1993)**. The
Excel workbook developed by Sarria et al. **(Sarria et al., 2009)** was used to calculate the parameters
of EPG data. For a given behavior, the parameters of occurrence and total duration were measured and
used to analyze the feeding behavior. We decided to select the following parameters for analysis
because they are relevant for the acquisition of phloem-limited viruses by aphids, *i.e.,* the total
durations and occurrences of stylet penetration, pathway phases, extracellular salivation phase, phloem
sap ingestion phases, and the time needed by the aphid to perform the first phloem phase.

Statistical analyses

Statistical analyses were made with R version 4.3.2 (https://www.r-project.org/).

Differences in virus accumulation between mono- and co-infections were determined using a Student

test.

We used Generalized Linear Models (GLMER, package: "lme4") with a likelihood ratio and chi-

squared (χ^2) test to assess whether co-infection affected virus transmission. Data on transmission rate

were analyzed following a binomial error distribution, and "session" was treated as random factor.

We used Generalized Linear Model (GLM) with the likelihood ratio and the chi-squared(χ^2) test to

assess whether mono or co-infection affected aphid feeding behavior. As feeding duration parameters

were not normally distributed, we used GLM using a gamma (link = "inverse") distribution, and

parameters related to frequency of penetration were modeled using GLM with Poisson (link =

"identity") distribution. The parameter "time to first phloem phase" was modeled using the Cox

proportional hazards (CPH) model and we treated cases where the given event did not occur as

censored. The assumption of the validity of proportional hazards was checked using the functions

"coxph" and "cox.zph", respectively (R packages: "survival" and "RVAideMemoire").

The fit of all generalized linear models was controlled by inspecting residuals and QQ plots. When a

significant effect was detected, to test for differences between treatments, a pairwise comparison using

estimated marginal means (package R: "emmeans") (*p*-value adjustment with Tukey method) at the

0.05 significance level was used.

RESULTS

Virus transmission by aphids from sugar beet co-infected with BChV and BYV

We wanted to know whether virus acquisition from BYV/BChV co-infected source plants would affect

the aphid transmission of each virus. For this, *Myzus persicae* aphids were allowed to acquire BChV

and/or BYV from mono- or co-infected sugar beets and then transferred onto test plants for virus

inoculation **(Fig. 1)**. The BYV transmission rate was not affected by sugar beet co-infection (aphid

transmission rate of 75% from mono-infected plants and 82% from BYV/BChV co-infected plants)

(Fig. 1A). BChV transmission was significantly reduced when aphids fed on BYV/BChV co-infected

plants compared to BChV mono-infected plants (30% *vs* 48%, respectively) **(Fig. 1B)**.

Aphids feeding on co-infected plants transmitted both viruses together at 19% transmission rate, which

was lower than expected, based on the virus transmission efficiency of each virus (36% expected co-

infection rate, calculated as the product of the transmission rates using mono-infected source plants).

BYV alone was present in 63% of the aphid-inoculated plants whereas BChV was detected alone in

11% of the infected plants **(S1 Fig.)**.

Figure 1. Aphid transmission efficiency of (A) BYV and (B) BChV from mono or co-infected plants. *Myzus persicae*
acquired virus from mono or co-infected leaves for 24 h. Then, three aphids were transferred per test plant for 72 h
inoculation. A DAS-ELISA assay was performed 3 weeks later to detect infection. The percentage of infected plants is
indicated (% Transmission). The difference was statistically significant for BChV but not for BYV transmission (GLMER,
BYV: p-value = 0.060; BChV: p-value< 0.001; n = 75, five independent experiments; df = 1;). NS = not significant, ***
= p-value < 0.001.

Aphid feeding behavior on mono-infected and co-infectd plants or co-infected with both viruses

Aphid feeding behavior plays an important role in virus transmission. To address whether aphid feeding behavior was impacted on sugar beet co-infected with BChV and BYV, compared to mono-infected sugar beet, we used the electrical penetration graph technique (EPG). Aphid total durations of probing time, intercellular pathway, intercellular salivation and the time spent to reach the phloem were the same for all conditions **(Fig. 2A)**. However, aphids spent significantly more time (~ one extra hour) ingesting phloem sap (E2 phase) on BYV mono-infected than on healthy or BChV-infected plants. The total number of all feeding phases were not affected by the different conditions except for the number of extracellular salivations (E1e) that was significantly lower on BYV infected plants than on BChV-infected plants, their number on healthy and BYV/BChV co-infected plants was intermediate (**Fig. 2B**).

Figure 2. Feeding behavior of *Myzus persicae* on healthy, BYV and BChV mono-infected or BYV/BChV co-infected sugar beets. (A-B) The behavior of individual aphids was recorded by electrical penetration graph (EPG) for 8 h on the third upper leaf. Selected EPG parameters are presented according to (A) duration or (B) occurrence. The histogram bars display means and standard errors (SEM). Different letters indicate significant differences between plant infection status as

tested by GLM (Generalized linear model) followed by pairwise comparisons using package R: emmeans (P<0.05 method:
Turkey). Statistical analysis of the duration of the events indicates a significant difference for the duration of phloem-sap
ingestion (E2) on BYV-infected *vs* healthy or BChV-infected plants (GLM, Df = 3, X^2 = 15.98, P = 0.001) but no differences
for the total duration of stylet penetrations (probing time) (GLM, Df = 3, X^2=2.585, P = 0.460), the total duration of pathway
phases (C) (GLM, Df = 3, X^2 = 6.962, P = 0.073), the total duration of intercellular salivation (E1e) (GLM, Df = 3, X^2 =
5.645, P = 0.130) and the time until first sap ingestion from the phloem (Cox, Df = 3 ,p= 0.526). Statistical analysis of the
occurrence of events revealed significant differences for the intercellular salivation between BYV and BChV infected plants
(GLM, Df = 3, X^2 = 8.581 ,P = 0.035) but no differences for the number of stylet insertions (GLM, Df = 3, X^2 = 4.725,P =
0.193), pathway phases (C) (GLM, Df = 3, X^2 = 4.954, P = 0.175), salivation phase (E1) (GLM, Df = 3, X^2 = 1.006, P =
0.800), and phloem sap ingestion (GLM, Df = 3, X^2 = 0.459, P = 0.928). n = 20-24 as indicated in the graphs.

BChV and BYV accumulation in mono- or co-infected plants

To address whether BChV and BYV accumulation could be impacted by co-infection, we quantified

virus titers by RT-qPCR on sugar beet mono- or co-infected with BYV and BChV. To account for

potential uneven virus distribution in the leaf, the entire leaf was ground and processed for RNA

extraction and virus quantification. A reference gene, tubulin, was selected for calibration, among six

tested, prior to virus quantification **(S2 Table)**. No significant differences in virus accumulation

between mono-infected plants or BYV/BChV co-infected plants were observed **(Fig. 3)**.

Figure 3. Relative accumulation of BYV and BChV in mono- and co-infected plants at 21 dpi. Virus accumulation in
source plants was measured by multiplex TaqMan RT-qPCR as described in Materials and Methods. Box plots show median
(black line), 25%-75% percentiles (box) and 10%-90% percentiles (whiskers), and outliers (dots). Differences were not
statistically significant (Student t-test, BYV: t = -0.90817, p-value = 0.390; BChV: t = 0.59456, p-value= 0.569; n = 5; df =
8). NS = not significant.

Tissue distribution of BChV and BYV in mono- and co-infected plants

To assess if co-infection affected tissue localization of BChV or BYV, SABER-FISH was performed

on mono- and co-infected plants at 21 dpi. In cross-sections of BYV mono-infected leaves,

fluorescence corresponding to BYV genomes was observed exclusively in phloem cells as expected

(Fig. 4B). The label was found in the phloem of primary (midribs) and secondary veins and the label

intensity, estimated by visual observations, was equal in both locations **(Fig. 4B)**. In BChV mono-

infected plants, fluorescence corresponding to BChV genomes was also observed, as expected, in the

phloem of primary and secondary veins **(Fig. 4C)**. However, the BChV label was less intense in the

primary veins than in the secondary veins. Co-infection altered the distribution of BYV and BChV

(Fig. 4D). Compared to mono-infection, less cells were labeled with BYV in midribs and more cells

in the secondary veins in co-infected plants. Compared to mono-infected plants, BChV was present in

more phloem cells of primary veins in co-infected plants, while the BChV label in secondary veins was

unchanged. Both viruses were still confined to phloem cells and no escape in non-phloem cells was

noticed **(Fig. 4D-F)**.

We estimated the percentage of co-infected cells and mono-infected cells in co-infected plants. Ten to

eleven sections from co-infected plants were observed in two independent experiments **(Table 2)**. We

noticed a slightly lower percentage of cells infected with BChV (63.2% of cells containing only BChV

or co-infected) compared to BYV (74.3% of cells containing only BYV or co-infected) **(Table 2)**.

Between 30-50% of all infected cells were co-infected, and similar percentages of cells were mono-

infected with BChV or BYV. This suggests that there is no exclusion between the two viruses at the

cellular level. The label intensity of each virus seemed to be similar in mono-infected and co-infected

cells, indicating that they did not interfere with each other's accumulation.

Figure 4. Tissue distribution of BChV and BYV in leaves mono-infected with BChV, BYV or co-infected with both
viruses. Leaves were processed three weeks after inoculation by SABER-FISH for detection of BChV and BYV. (A-D)
Representative images showing transversal sections of (A) healthy, (B) BYV mono-infected, (C) BChV mono-infected,
and (D) BChV/BYV co-infected leaves. The first column shows the BChV label (red), the second the BYV label (turquoise),
the third bright field acquisitions and the last column presents merges. (E-F) Magnifications of the regions outlined in (D)
showing virus distribution in (E) the midrib and (F) the leaf lamina. The first column shows image merges, the second the
BYV signal (red), and the last column the BChV label (red). Scale bars = 200 µm (A-D) and 50 µm (E-F).

Intracellular distribution of BYV and BChV in mono and co-infected plants

Cross sections of phloem tissue allow only a limited view of the phloem cell and sieve tube lumen

because their elongated shapes are arranged perpendicular to the section plane. To better assess if co-

infection affected intracellular distribution of BChV or BYV, SABER-FISH was performed on

longitudinal sections of infected plants at 21 dpi **(Fig. 4)**. In mono-infected plants, BYV label was

present in infected cells as spherical cytoplasmic inclusions that aligned like pearl chains, similarly as

reported before using light microscopy **(Esau, 1960) (Fig. 4A)**. In the case of BChV mono-infected

plants, the virus label was spread uniformly in the cytoplasm of infected cells **(Fig. 4B)**. Similar label

patterns were also found in immunofluorescence experiments **(S2 Fig.)**, suggesting that the FISH label

corresponded, at least partially, to virions.

Table 2: Number of cells infected by one or both viruses in sections of co-infected leaves.

Virus(es) observed	Exp. 1			Exp. 2		
	Plant #1	Plant #2	Plant #3	Plant #1	Mean[c]	%[d]
BChV	32[a] (33.0%[b])	13 (20.6%)	3 (6.3%)	22 (34.4%)	17.5 ± 12.4	25.7%
BYV	33 (34.0%)	19 (30.2%)	27 (56.3%)	21 (32.8%)	25 ± 6.3	36.8%
BYV+BChV	32 (33.0%)	31 (49.2%)	18 (37.5%)	21 (32.8%)	25.5 ± 7.0	37.5%

384 [a] Number of cells observed on 10 to 11 sections of each plant showing a fluorescent signal for the indicated virus. [b]
385 Percentage of cells labeled for the indicated virus/total number of labeled cells in 10 to 11 sections. [c] Mean number of cells
386 observed on 10 to 11 sections for 4 independent plants. [d] Percentage of cells containing one or both viruses/total number of
387 cells observed in the experiments.

Figure 5. Intra-cellular distribution of BYV and BChV in leaves of mono-infected plants. Leaves were processed three weeks after inoculation by SABER-FISH for detection of BYV and BChV. The images show longitudinal sections of (A) a leaf mono-infected with BYV and (B-C) a leaf mono-infected with BChV with virus label in the phloem in (B) probably a sieve tube and (C) probably in a sieve tube and in a companion cell (left and right arrows pointing from Phl, respectively). The first column represents BChV label (red), the second BYV label (turquoise), the third differential interference contrast images, and the last column presents image merges. Xyl, xylem; Phl, phloem. Scale bars = 20 µm.

In BYV/BChV co-infected plants, both viruses showed the same intracellular distribution as in mono-infected plants when the cells were infected by one virus only, *i.e.*, pearl-like inclusions for BYV and homogenous cytoplasmic distribution for BChV **(Fig. 6A-B)**. However, in co-infected cells, the BYV label was mostly dispersed in the cytoplasm and colocalized partially with the diffuse BChV label **(Fig. 6C-D)**.

Figure 6. Intra-cellular distribution of BYV and BChV in leaves of co-infected plants. Leaves were processed three
weeks after inoculation by SABER-FISH for detection of BChV (red) and BYV (turquoise). The images show longitudinal
sections of phloem with (A-B) mono-infected cells, (C) a cell mono-infected with BChV (arrow) and an adjacent cell co-
infected with both viruses (arrowheads), and (D) a co-infected phloem cell. The first column presents BChV label, the
second BYV label, the third differential interference contrast images, and the last column image merges. Scale bars = 20
µm.

DISCUSSION

Few studies have shown that viruses from various genera or families co-existing in multi-infection can modify vector transmission, and even less have addressed modifications at the cellular level. Additionally, to our knowledge, this is the first study examining the effect of co-infection by a polerovirus, (BChV) and a closterovirus (BYV), both phloem-limited and sharing the same aphid vector, on the transmission efficiency and other infection parameters. We found that co-infection of sugar beet with BYV and BChV reduced aphid transmission of BChV significantly, compared to transmission from mono-infected plants. No effect of the co-infection of sugar beet was observed on the transmission efficiency of BYV.

To find possible explanations for the reduction of BChV transmission from co-infected plants, we analyzed aphid feeding behavior, virus accumulation, and virus distribution in the leaves and the infected cells on mono- and co-infected plants.

Viral infections may impact aphid-feeding behavior, which can have a direct impact on virus acquisition and transmission **(Mauck et al., 2012; Mauck et al., 2016)**. The feeding behavior can be further modified in viral co-infections, but this has been hardly studied. One of the few published examples is that of raspberries co-infected with raspberry latent virus and raspberry leaf mottle virus. In this pathosystem, no changes in feeding behavior were observed, although aphid preferences for infected plants did change **(Lightle & Lee, 2014)**. Another example is the co-infection of melon plants with the potyvirus watermelon mosaic virus (WMV) and the crinivirus cucurbit yellow stunting disorder virus (CYSDV). There, WMV accumulation was ten times lower in co-infected plants, but its transmission rate by aphids remained stable. EPG analysis showed that aphids on co-infected melons performed longer duration of ingestion (more specifically, pulses of the subphase II-3) during intracellular test punctures in the epidermis and mesophyll, which is the feeding behavior associated with the acquisition of non-persistently transmitted viruses like WMV. The authors argued that the changed feeding behavior might compensate for the low WMV accumulation in co-infected plants **(Domingo-Calap et al., 2020)**. Thus, aphid feeding behavior could be a determining factor that affects virus acquisition and explains the lowered transmission of BChV from co-infected plants. Since BChV should be acquired from the phloem sap**,** longer phloem sap ingestion should increase acquisition **(Gray et al., 1991; Prado & Tjallingii, 1994)**. However, our results showed that despite aphids having a tendency to ingest phloem sap for longer durations on co-infected plants than on mono-infected ones, transmission of BChV dropped. This indicates that the decreased transmission of BChV cannot be

explained by aphid feeding behavior. Interestingly, for BYV, transmission from co-infected plants was

not affected although aphids tended to feed less on co-infected plants than on mono-infected plants.

We assume that under our experimental conditions, the 3 h sap ingestion observed for aphids feeding

on co-infected plants is sufficient to charge aphids maximally with BYV and that for this reason, we

did not observe a difference in transmission efficiency of BYV from co-infected or mono-infected

plants. This is in line with the results by Jiménez and co-workers (2018) who demonstrated that BYV

maximum acquisition was obtained after 3 to 6 h of phloem sap ingestion, with no significant increases

in transmission rates observed for longer sap ingestion periods **(Jiménez et al., 2018)**. Taken together,

we have no evidence that the altered aphid feeding behavior observed on co-infected sugar beets is the

cause of the lower BChV transmission from these plants.

Lower or higher virus accumulation in co-infected compared to mono-infected plants may also directly

affect transmission efficiency by modulating the amount of virus particles acquired by the vector. For

instance, vector transmission efficiency of two criniviruses, tomato chlorosis virus (ToCV) and tomato

infectious chlorosis virus (TICV), was linked to virus accumulation in mono- and co-infected tomatoes

(Wintermantel et al., 2008). However, an overall higher accumulation of one or both viruses in co-

infected plants does not necessarily lead to increased vector transmission. Rather, the virus titer

increase needs to occur in the aphid feeding sites (e.g., sieve tubes for polerovirus acquisition) **(Prado**

& Tjallingii, 1994). In this sense, a recent study showed that co-infection of tomato with a

begomovirus and a crinivirus resulted in a synergistic effect on the accumulation of both viruses, but

only increased the acquisition efficiency of the begomovirus **(Li et al., 2021)**. An increase in PLRV

accumulation in *Nicotiana clevelandii* was observed when the plants were co-infected with the potato

virus Y and the co-infection also induced phloem escape of PLRV **(Barker, 1987)**. However, aphid

transmission of PLRV and PVY was not addressed in this study. Our results showed that transmission

of BChV drops when plants are co-infected with BYV, although BChV levels were not statistically

different in mono- compared to co-infected plants. This indicates clearly that changes in aphid

transmission rates of BChV are not linked to virus titer.

Another reason to explain altered transmission might be changes in the tissue tropism of viruses in

plants that might modify virus accessibility to aphids. For instance, phloem escape of aphid-vectored

PLRV in tobacco was reported in plants co-infected with pea enation mosaic virus 2 (PEMV-2, **Ryabov**

et al., 2001) or PVY **(Barker, 1987)**. The changed tissue specificity correlated with PLRV becoming

mechanically transmissible in PLRV/PEMV-2 co-infections**.** In our observations, we did not find

evidence of a phloem escape of BChV or BYV in co-infected plants, ruling out altered tissue tropism

as the reason for reduced BChV transmission from co-infected plants. What we did observe in co-

infected plants, though, was that BChV localized equally in the phloem of primary and secondary veins,

while BYV was more present in those of secondary veins. However, since both viruses are acquired

preferentially from sieve tube sap, it is unlikely that the tissue relocalization impacted virus acquisition

greatly.

We found that in co-infected phloem cells (representing about 40% of all infected cells), the

intracellular distribution of BYV was modified. BYV was present as spherical, cytoplasmic inclusions

aligning in chains in cells of mono-infected plants. BChV was spread uniformly in the cytoplasm in

cells of mono-infected plants. In co-infected plants, both viruses displayed the same intracellular

localization when cells were infected by only one virus. This was different in co-infected cells. While

BChV maintained its diffuse distribution, the typical BYV inclusions disappeared and BYV label was

diffuse in the cytoplasm and colocalized partially with BChV. We take this as evidence that the two

viruses interact directly or indirectly with each other. Since no other notable differences were detected

in co-infected *vs* mono-infected plants, virus-virus interactions in co-infected cells likely caused the

drop in BChV transmission. At this stage, the exact mechanism of interaction between BChV and BYV

is unknown. Because BChV should be acquired predominantly from the phloem sap and only to a small

extent from phloem cells **(Prado & Tjallingii, 1994)**, we propose that co-infection limits BChV release

from companion cells into the sieve tubes **(Fig. 7)**. This would result in a lower BChV accessibility to

aphids, compared to BChV mono-infected plants. This could be achieved by alterations of the

intracellular virus trafficking or modifications of the cell wall or plasmodesmata. Several BYV and

BChV proteins are known to be involved in virus movement and interaction with plasmodesmata and

are therefore good candidates for further studies **(Boissinot et al., 2020; Dolja, 2003)**.

Our results show that the percentage of co-infected cells and those mono-infected with BYV or BChV

were in the same range (about 30%) in co-infection. This suggests that co-infection is a random event

and that the viruses do not exclude each other. Further, there was no alteration in the accumulation of

either virus, indicating that there is no competition for plant resources and that they do not interfere

with each other's replication. This is indicative of neutralism. This assumption is also reinforced by

the fact that BYV/BChV co-infection, compared to infection with BYV alone, does not have a more

severe impact on sugar beet root mass (https://www.itbfr.org/publications/fiches-bioagresseurs/les-

jaunisses-virales-et-leurs-pucerons-vecteurs/, accessed on 4 Avril 2024). Note that the same sugar beet

cultivar was used in both studies.

In conclusion, our results shed some light on the mode of interaction between two unrelated viruses, a

polerovirus and a closterovirus, a combination that has never been studied before. Except for the

antagonism observed for BChV transmission by aphids, the outcome of the sugar beet co-infection

with BChV and BYV is fairly neutral for all other parameters analyzed. The reasons for BChV

decreased aphid transmission by aphid are unknown but the subcellular localization of both viruses in

co-infected plants strongly suggests interactions between them that are worth exploring. Finally, the

alteration of aphid transmission of BChV from co-infected plants might have consequences at the

epidemiological level. For example, it might explain partially the huge changes in incidences of the

two viruses from one year to another **(Hossain et al., 2021)**.

Figure 7. Model for explaining the reduction of BChV transmission from BChV/BYV co-infected plants. In mono-
infected plants (middle and right cell), BChV and BYV particles are dispatched from infected companion cells through
plasmodesmata into the sieve tube lumen and acquired by aphids ingesting phloem sap from the sieve tubes. In co-infected
plants, roughly one-third of the infected phloem cells are co-infected. Co-infection does not affect virus accumulation
measurably, and the two viruses might even accumulate to similar levels in mono-infected and co-infected cells, as
presented in the schema (left cell). BYV and BChV interact in co-infected cells directly or indirectly. This inhibits the
passage of BChV towards and/or through plasmodesmata and BChV release into the phloem sap, whereas phloem sap
loading with BYV remains unchanged. Consequently, the BChV load in the sieve tube is significantly lower, while that of
BYV is not changed, and aphids acquire and transmit less BChV from the sieve tubes of co-infected plants than from those
of mono-infected plants. Two scenarios are thinkable that are not mutually exclusive: Co-infected phloem cells inhibit
BChV release totally, and the BChV particles in the phloem sap origin from BChV mono-infected cells, or BYV infection
interferes with phloem loading from all companion cells. In the former case, BYV operates only in co-infected cells; the
latter case requires the movement of a signal or factor from BYV-infected cells to uninfected and BChV-infected cells.

AUTHORS' CONTRIBUTIONS

SK: Conceptualization, methodology, validation, formal analysis, investigation, writing – original draft preparation, writing – review & editing, visualization

QC: Methodology, formal analysis, writing – review & editing

CV: Resources

VB: Conceptualization, methodology, investigation, writing – original draft preparation, writing – review & editing, supervision, project administration, funding acquisition

MD: Conceptualization, methodology, investigation, writing – original draft preparation, writing – review & editing, visualization, supervision, project administration, funding acquisition

ACKNOWLEDGMENTS

Plants were produced by the experimental unit of INRAE Grand Est-Colmar (UEAV). We thank Samia Djanane for help with RT-qPCR, Amandine Velt for help with probe design and Danaé Brun for help with virus identification.

FUNDING

This work was financed by the PNRI project ProViBe awarded to VB and the ANR PRCI project ManiVir (ANR-23-CE20-0049-01) awarded to MD. SK received additional funding from INRAE SPE department. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

DATA AVAILABILITY

All relevant data are within the manuscript and its Supporting Information files.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

SUPPLEMENTS

Figure S1. Aphid transmission efficiency of BYV and BChV from co-infected plants.

Figure S2. Tissue distribution of BChV and BYV in roots mono-infected with BChV or BYV.

Table S1. Primers with concatemer sequences used for Primer Exchange Reaction.

Table S2. Amplification efficacy (%) of reference genes, and the associated correlation coefficient (R^2).

Data S1. Transmission experiments.

Data S2. EPG experiments.

554   **Data S3.** RT-qPCR virus accumulation.

555   **Data S4.** Co-infection cell count.

REFERENCES

Abdullah, A. S., Moffat, C. S., Lopez-Ruiz, F. J., Gibberd, M. R., Hamblin, J., & Zerihun, A. (2017). Host–Multi-Pathogen Warfare : Pathogen Interactions in Co-infected Plants. *Frontiers in Plant Science*, *8*, 1806. https://doi.org/10.3389/fpls.2017.01806

Adams, A. N., & Clark, M. F. (1977). Characteristics of the Microplate Method of Enzyme-Linked Immunosorbent Assay for the Detection of Plant Viruses. *Journal of General Virology*, *34*(3), 475-483. https://doi.org/10.1099/0022-1317-34-3-475

Andersen, C. L., Jensen, J. L., & Ørntoft, T. F. (2004). Normalization of Real-Time Quantitative Reverse Transcription-PCR Data : A Model-Based Variance Estimation Approach to Identify Genes Suited for Normalization, Applied to Bladder and Colon Cancer Data Sets. *Cancer Research*, *64*(15), 5245-5250. https://doi.org/10.1158/0008-5472.CAN-04-0496

Barker, H. (1987). Invasion of Non-phloem Tissue in Nicotiana clevelandii by Potato Leafroll Luteovirus Is Enhanced in Plants also Infected with Potato Y Potyvirus. *Journal of General Virology*, *68*(4), 1223-1227. https://doi.org/10.1099/0022-1317-68-4-1223

Beliveau, B. J., Kishi, J. Y., Nir, G., Sasaki, H. M., Saka, S. K., Nguyen, S. C., Wu, C., & Yin, P. (2018). OligoMiner provides a rapid, flexible environment for the design of genome-scale oligonucleotide in situ hybridization probes. *Proceedings of the National Academy of Sciences*, *115*(10). https://doi.org/10.1073/pnas.1714530115

Boissinot, S., Brault, V., & Herrbach, E. (2020). Luteovirids. In *Encyclopedia of Life Sciences* (p. 327-339). John Wiley & Sons, Ltd. https://doi.org/10.1002/9780470015902.a0029206

Boissinot, S., Pichon, E., Sorin, C., Piccini, C., Scheidecker, D., Ziegler-Graff, V., & Brault, V. (2017). Systemic Propagation of a Fluorescent Infectious Clone of a Polerovirus Following Inoculation by Agrobacteria and Aphids. *Viruses*, *9*(7), 166. https://doi.org/10.3390/v9070166

Bragard, C., Caciagli, P., Lemaire, O., Lopez-Moya, J. J., MacFarlane, S., Peters, D., Susi, P., & Torrance, L. (2013). Status and Prospects of Plant Virus Control Through Interference with Vector Transmission. *Annual Review of Phytopathology*, *51*(1), 177-201. https://doi.org/10.1146/annurev-phyto-082712-102346

Brault, V., Uzest, M., Monsion, B., Jacquot, E., & Blanc, S. (2010). Aphids as transport devices for plant viruses. *Comptes Rendus. Biologies*, *333*(6-7), 524-538. https://doi.org/10.1016/j.crvi.2010.04.001

Bruyère, A., Brault, V., Ziegler-Graff, V., Simonis, M. T., Van den Heuvel, J. F., Richards, K., Guilley, H., Jonard, G., & Herrbach, E. (1997). Effects of mutations in the beet western yellows virus readthrough protein on its expression and packaging and on virus accumulation, symptoms, and aphid transmission. *Virology*, *230*(2), 323-334. https://doi.org/10.1006/viro.1997.8476

Chen, Z., Tan, J. Y., Wen, Y., Niu, S., & Wong, S.-M. (2012). A Game-Theoretic Model of Interactions between Hibiscus Latent Singapore Virus and Tobacco Mosaic Virus. *PLoS ONE*, *7*(5), e37007. https://doi.org/10.1371/journal.pone.0037007

Cuellar, W. J., Galvez, M., Fuentes, S., Tugume, J., & Kreuze, J. (2015). Synergistic interactions of begomoviruses with *Sweet potato chlorotic stunt virus* (genus *Crinivirus*) in sweet potato (*Ipomoea batatas* L .). *Molecular Plant Pathology*, *16*(5), 459-471. https://doi.org/10.1111/mpp.12200

Cuellar, W. J., Kreuze, J. F., Rajamäki, M.-L., Cruzado, K. R., Untiveros, M., & Valkonen, J. P. T. (2009). Elimination of antiviral defense by viral RNase III. *Proceedings of the National Academy of Sciences*, *106*(25), 10354-10358. https://doi.org/10.1073/pnas.0806042106

Dáder, B., Then, C., Berthelot, E., Ducousso, M., Ng, J. C. K., & Drucker, M. (2017). Insect transmission of plant viruses : Multilayered interactions optimize viral propagation. *Insect Science*, *24*(6), 929-946. https://doi.org/10.1111/1744-7917.12470

Devi, P., Khan, A., Chattopadhyay, P., Mehta, P., Sahni, S., Sharma, S., & Pandey, R. (2021). Co-infections as Modulators of Disease Outcome : Minor Players or Major Players? *Frontiers in Microbiology*, *12*, 664386. https://doi.org/10.3389/fmicb.2021.664386

Dolja, V. V. (2003). *Beet yellows virus* : The importance of being different. *Molecular Plant Pathology*, *4*(2), 91-98. https://doi.org/10.1046/j.1364-3703.2003.00154.x

Dolja, V. V., & Koonin, E. V. (2013). The closterovirus-derived gene expression and RNA interference vectors as tools for research and plant biotechnology. *Frontiers in Microbiology*, *4*. https://doi.org/10.3389/fmicb.2013.00083

Domingo-Calap, M. L., Moreno, A. B., Díaz Pendón, J. A., Moreno, A., Fereres, A., & López-Moya, J. J. (2020). Assessing the Impact on Virus Transmission and Insect Vector Behavior of a Viral Mixed Infection in Melon. *Phytopathology®*, *110*(1), 174-186. https://doi.org/10.1094/PHYTO-04-19-0126-FI

Dutt, A., Andrivon, D., & Le May, C. (2022). Multi-infections, competitive interactions, and pathogen coexistence. *Plant Pathology*, *71*(1), 5-22. https://doi.org/10.1111/ppa.13469

Esau, K. (1960). Cytologic and histologic symptoms of beet yellows. *Virology*, *10*, 73-85. https://doi.org/10.1016/0042-6822(60)90007-6

Esau, K., Cronshaw, J., & Hoefert, L. L. (1967). Relation of beet yellows virus to the phloem and to movement in the sieve tube. *The Journal of Cell Biology*, *32*(1), 71-87. https://doi.org/10.1083/jcb.32.1.71

Esau, K., & Hoefert, L. L. (1972). Ultrastructure of sugarbeet leaves infected with beet western yellows virus. *Journal of Ultrastructure Research*, *40*(5-6), 556-571. https://doi.org/10.1016/S0022-5320(72)80043-1

Gautam, S., Gadhave, K. R., Buck, J. W., Dutta, B., Coolong, T., Adkins, S., & Srinivasan, R. (2020). Virus-virus interactions in a plant host and in a hemipteran vector : Implications for vector fitness and virus epidemics. *Virus Research*, *286*, 198069. https://doi.org/10.1016/j.virusres.2020.198069

German-Retana, S., Candresse, T., & Martelli, G. (1999). CLOSTEROVIRUSES (*CLOSTEROVIRIDAE*). In A. Granoff & R. G. Webster (Éds.), *Encyclopedia of Virology (Second Edition)* (p. 266-273). Elsevier. https://doi.org/10.1006/rwvi.1999.0053

Gray, S. M., Power, A. G., Smith, D. S., Seaman, A. J., & Altman, N. S. (1991). Aphid Transmission of Barley Yellow Dwarf Virus : Acquisition Access Periods and Virus Concentration Requirements. *Phytopathology*, *81*, 539-545. https://doi.org/10.1094/Phyto-81-539

Hauser, S., Stevens, M., Beuve, M., & Lemaire, O. (2002). Biological properties and molecular characterization of beet chlorosis virus (BChV). *Archives of Virology*, *147*(4), 745-762. https://doi.org/10.1007/s007050200023

Hossain, R., Menzel, W., Lachmann, C., & Varrelmann, M. (2021). New insights into virus yellows
distribution in Europe and effects of beet yellows virus, beet mild yellowing virus, and beet
chlorosis virus on sugar beet yield following field inoculation. *Plant Pathology*, *70*(3),
584-593. https://doi.org/10.1111/ppa.13306

Jiménez, J., Tjallingii, W. F., Moreno, A., & Fereres, A. (2018). Newly Distinguished Cell Punctures
Associated with Transmission of the Semipersistent Phloem-Limited Beet Yellows Virus.
Journal of Virology, *92*(21), e01076-18. https://doi.org/10.1128/JVI.01076-18

Kishi, J. Y., Lapan, S. W., Beliveau, B. J., West, E. R., Zhu, A., Sasaki, H. M., Saka, S. K., Wang,
Y., Cepko, C. L., & Yin, P. (2019). SABER amplifies FISH : Enhanced multiplexed imaging
of RNA and DNA in cells and tissues. *Nature Methods*, *16*(6), 533-544.
https://doi.org/10.1038/s41592-019-0404-0

Kozera, B., & Rapacz, M. (2013). Reference genes in real-time PCR. *Journal of Applied Genetics*,
54(4), 391-406. https://doi.org/10.1007/s13353-013-0173-x

Kozłowska-Makulska, A., Beuve, M., Syller, J., Szyndel, M. S., Lemaire, O., Bouzoubaa, S., &
Herrbach, E. (2009). Aphid transmissibility of different European beet polerovirus isolates.
European Journal of Plant Pathology, *125*(2), 337-341. https://doi.org/10.1007/s10658-009-
9474-7

Li, J., Wang, J., Ding, T., & Chu, D. (2021). Synergistic Effects of a Tomato chlorosis virus and
Tomato yellow leaf curl virus Mixed Infection on Host Tomato Plants and the Whitefly
Vector. *Frontiers in Plant Science*, *12*, 672400. https://doi.org/10.3389/fpls.2021.672400

Lightle, D., & Lee, J. (2014). Raspberry viruses affect the behavior and performance of
Amphorophora agathonica in single and mixed infections. *Entomologia Experimentalis et
Applicata*, *151*(1), 57-64. https://doi.org/10.1111/eea.12170

Limburg, D. D., Mauk, P. A., & Godfrey, L. D. (1997). Characteristics of Beet Yellows Closterovirus
Transmission to Sugar Beets by *Aphis fabae*. *Phytopathology®*, *87*(7), 766-771.
https://doi.org/10.1094/PHYTO.1997.87.7.766

Liu, W., Wei, T., & Wang, X. (2023). Plant reoviruses hijack autophagy in insect vectors. *Trends in
Microbiology*, *31*(12), 1251-1261. https://doi.org/10.1016/j.tim.2023.06.008

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods*, *25*(4), 402-408.
https://doi.org/10.1006/meth.2001.1262

Maliano, M. R., Rojas, M. R., Macedo, M. A., Barboza, N., & Gilbertson, R. L. (2022). The invasion
biology of tomato begomoviruses in Costa Rica reveals neutral synergism that may lead to
increased disease pressure and economic loss. *Virus Research*, *317*, 198793.
https://doi.org/10.1016/j.virusres.2022.198793

Mauck, K., Bosque-Pérez, N. A., Eigenbrode, S. D., De Moraes, C. M., & Mescher, M. C. (2012).
Transmission mechanisms shape pathogen effects on host–vector interactions : Evidence from
plant viruses. *Functional Ecology*, *26*(5), 1162-1175. https://doi.org/10.1111/j.1365-
2435.2012.02026.x

Mauck, K. E., De Moraes, C. M., & Mescher, M. C. (2016). Effects of pathogens on sensory-
mediated interactions between plants and insect vectors. *Current Opinion in Plant Biology*,
32, 53-61. https://doi.org/10.1016/j.pbi.2016.06.012

McLaughlin, A. A., Hanley-Bowdoin, L., Kennedy, G. G., & Jacobson, A. L. (2022). Vector
acquisition and co-inoculation of two plant viruses influences transmission, infection, and
replication in new hosts. *Scientific Reports*, *12*(1), 20355. https://doi.org/10.1038/s41598-
022-24880-5

Moreno, A. B., & López-Moya, J. J. (2020). When Viruses Play Team Sports : Mixed Infections in
Plants. *Phytopathology*, *110*(1), 29-48. https://doi.org/10.1094/PHYTO-07-19-0250-FI

Mukasa, S. B., Rubaihayo, P. R., & Valkonen, J. P. T. (2006). Interactions between a crinivirus, an
ipomovirus and a potyvirus in coinfected sweetpotato plants. *Plant Pathology*, *55*(3),
458-467. https://doi.org/10.1111/j.1365-3059.2006.01350.x

Mutterer, J. D., Stussi-Garaud, C., Michler, P., Richards, K. E., Jonard, G., & Ziegler-Graff, V.
(1999). Role of the beet western yellows virus readthrough protein in virus movement in
Nicotiana clevelandii. *Journal of General Virology*, *80*(10), 2771-2778.
https://doi.org/10.1099/0022-1317-80-10-2771

Ontiveros, I., López-Moya, J. J., & Díaz-Pendón, J. A. (2022). Coinfection of Tomato Plants with
Tomato yellow leaf curl virus and Tomato chlorosis virus Affects the Interaction with Host
and Whiteflies. *Phytopathology*, *112*(4), 944-952. https://doi.org/10.1094/PHYTO-08-21-
0341-R

Pfaffl, M. W., Tichopad, A., Prgomet, C., & Neuvians, T. P. (2004). Determination of stable
housekeeping genes, differentially regulated target genes and sample integrity : BestKeeper –
Excel-based tool using pair-wise correlations. *Biotechnology Letters*, *26*(6), 509-515.
https://doi.org/10.1023/B:BILE.0000019559.84305.47

Prado, E., & Tjallingii, W. F. (1994). Aphid activities during sieve element punctures. *Entomologia
Experimentalis et Applicata*, *72*(2), 157-165. https://doi.org/10.1111/j.1570-
7458.1994.tb01813.x

Reinink, K., Dieleman, F. L., Jansen, J., & Montenarie, A. M. (1989). Interactions between plant and
aphid genotypes in resistance of lettuce to Myzus persicae and Macrosiphum euphorbiae.
Euphytica, *43*(3), 215-222. https://doi.org/10.1007/BF00023056

Ryabov, E. V., Fraser, G., Mayo, M. A., Barker, H., & Taliansky, M. (2001). Umbravirus gene
expression helps potato leafroll virus to invade mesophyll tissues and to be transmitted
mechanically between plants. *Virology*, *286*(2), 363-372.
https://doi.org/10.1006/viro.2001.0982

Sarria, E., Cid, M., Garzo, E., & Fereres, A. (2009). Excel Workbook for automatic parameter
calculation of EPG data. *Computers and Electronics in Agriculture*, *67*(1), 35-42.
https://doi.org/10.1016/j.compag.2009.02.006

Savenkov, E. I., & Valkonen, J. P. T. (2001). Potyviral Helper-Component Proteinase Expressed in
Transgenic Plants Enhances Titers of Potato Leaf Roll Virus but Does Not Alleviate Its
Phloem Limitation. *Virology*, *283*(2), 285-293. https://doi.org/10.1006/viro.2000.0838

Singhal, P., Nabi, S. U., Yadav, M. K., & Dubey, A. (2021). Mixed infection of plant viruses :
Diagnostics, interactions and impact on host. *Journal of Plant Diseases and Protection*,
128(2), 353-368. https://doi.org/10.1007/s41348-020-00384-0

Stevens, M., Hallsworth, P. B., & Smith, H. G. (2004). The effects of *Beet mild yellowing virus* and
Beet chlorosis virus on the yield of UK field-grown sugar beet in 1997,1999 and 2000. *Annals
of Applied Biology*, *144*(1), 113-119. https://doi.org/10.1111/j.1744-7348.2004.tb00323.x

Stevens, M., Hull, R., & Smith, H. G. (1997). Comparison of ELISA and RT-PCR for the detection
of beet yellows closterovirus in plants and aphids. *Journal of Virological Methods*, *68*(1),
9-16. https://doi.org/10.1016/s0166-0934(97)00103-1

Syller, J., & Grupa, A. (2016). Antagonistic within-host interactions between plant viruses :
Molecular basis and impact on viral and host fitness. *Molecular Plant Pathology*, *17*(5),
769-782. https://doi.org/10.1111/mpp.12322

Tjallingii, W. F. (1988). Electrical recording of stylet penetration activities. In *Aphids, their biology,*
natural enemies and control (p. 95-108). Elsevier Science Publishers.

Tjallingii, W. F., & Hogen Esch, T. (1993). Fine structure of aphid stylet routes in plant tissues in
correlation with EPG signals. *Physiological Entomology*, *18*(3), 317-328.
https://doi.org/10.1111/j.1365-3032.1993.tb00604.x

Van den Heuvel, J. F. J. M., Boerma, T. M., & Peters, D. (1991). Transmission of Potato leafroll
virus from plants and artificial diets by Myzus persicae. *Phytopathology*, *81*, 150-154.
https://doi.org/10.1094/Phyto-81-150

Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., & Speleman, F.
(2002). Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. *Genome Biology*, *3*(7), research0034.1.
https://doi.org/10.1186/gb-2002-3-7-research0034

Wintermantel, W. M. (2005). Co-infection of *Beet mosaic virus* with Beet Yellowing Viruses Leads
to Increased Symptom Expression on Sugar Beet. *Plant Disease*, *89*(3), 325-331.
https://doi.org/10.1094/PD-89-0325

Wintermantel, W. M., Cortez, A. A., Anchieta, A. G., Gulati-Sakhuja, A., & Hladky, L. L. (2008).
Co-Infection by Two Criniviruses Alters Accumulation of Each Virus in a Host-Specific
Manner and Influences Efficiency of Virus Transmission. *Phytopathology®*, *98*(12),
1340-1345. https://doi.org/10.1094/PHYTO-98-12-1340

Zhang, G., Zhang, Z., Wan, Q., Zhou, H., Jiao, M., Zheng, H., Lu, Y., Rao, S., Wu, G., Chen, J., Yan,
F., Peng, J., & Wu, J. (2023). Selection and Validation of Reference Genes for RT-qPCR
Analysis of Gene Expression in Nicotiana benthamiana upon Single Infections by 11
Positive-Sense Single-Stranded RNA Viruses from Four Genera. *Plants*, *12*(4), 857.
https://doi.org/10.3390/plants12040857

754
755

SUPPLEMENTS (Figures S1-S2, Tables S1-S2)

Figure S1. Aphid transmission efficiency of BYV and BChV from co-infected plants. *Myzus persicae* acquired virus
from co-infected leaves for 24 h. Then, three aphids were transferred per test plant for 72 h inoculation. A DAS-ELISA
assay was performed 3 weeks later to detect infection. The percentage of expected and observed co-infected plants is
indicated. Difference was statistically significant (p-value= 0.02865; n=75, five independent experiments; df=1; chi-
squared(χ2)).

Figure S2. Tissue distribution of BChV and BYV in roots mono-infected with BChV or BYV. Sugar beet roots were
processed three weeks after inoculation for immunofluorescence detection of BChV and BYV. (A-B) show longitudinal
sections of (A) BYV mono-infected, (B) BChV mono-infected roots. The first column shows the virus label, the second
and the third columns show autofluorescence and the last column presents merges. The white arrows in the image merges
indicate the virus label. Scale bars = 200 µm (A-B).

Table S1. Primers with concatemer sequences (red) used for Primer Exchange Reaction.

BYV	
Primer1	CTGTCGGGCACAATGGGGCTTGCACTTTATAAACCTA
Primer2	CATGTCTCGGCGGGTGTCCACGACTTTATAAACCTA
Primer4	GGGTTGTCGGTAGAAGCACTCAAGTGCACACGTTTATAAACCTA
Primer5	GCACGACCTCGGTGCAGGCTCGTTTATAAACCTA
Primer6	GGGAACTGACAGAGAGGACTTCAGTTGGGAACACGTTTATAAACCTA
Primer7	CTGTAAGAGCCGACCAGCTGGAAGAGTCTTCGTTTATAAACCTA
Primer8	GGGCGCGAGTTAGGTTTCGATGTAACACCTAACGTTTATAAACCTA
Primer9	CGCGAGGAGTTTGTTGCCTCCTGCGTTTATAAACCTA
Primer10	CGCGCGCCTGAGCTTCAACAAGTTGTTCGTTTATAAACCTA
Primer11	GTTCGCCCGGAGTCACCATGGTGAGGTTTATAAACCTA
Primer12	GCCATGTCGATAGCGAGAGTTAAACGTTAAAGGCAGGGTTTATAAACCTA
Primer13	CACTCGAGTGTAAGAGACCGAACTGGGTTTCAACCTATTCATACTTTATAAACCTA
Primer14	CCATTTCGGCGTGCGTGGGAATTTGTGCATTTATAAACCTA
Primer15	CTACGAAAGGAGCCGCCACTTCAGCCAGTTCTTTATAAACCTATTTATAAACCTA
Primer16	CTTCCACTCCACTCATGTTTATTGTGAATTCACCGAGAGCCGTTTATAAACCTA
Primer17	GGAACAGTTTGGGCAACGAACGAGCTATCACTATGTAAGCTTTATAAACCTA
Primer18	CGGCGCGAACACTGGGTCCGAGTTTATAAACCTA
Primer19	GCGGCGTGGCCGCAGTGACTACTTTATAAACCTA
Primer 20	GAGCGGCGTGGGCGAGGTAACATAACCTTTATAAACCTA
Primer21	CGAGAGACTCGTCTTCCGCGCACAGGTTTATAAACCTA
Primer 22	GCTGACACGTTCGTAGACTGATTCGATCGCCTTGTTTATAAACCTA
Primer23	CCACTCTGCACGGCCTCCCCCATATAGAAGTTTATAAACCTA

Primer 24	CCAGCAGACGGGAACCGGCCGTTTTATAAACCTA
Primer25	CGGTCTGCACACGTGGTAATCTGTGCTCCTTTTATAAACCTA
Hairpin	Hairpin.26 : AATAAACCTAGGGCCTTTTGGCCCTAGGTTTATTTAGGTTTATTTTTTT
Imager	Fluor26 /ATTO647N/TTTAGGTTTATT TAGGTTTATT

BChV	
Primer 1	GTGCCATGGTCCCAAGAGCTCCAAATAAGATTAGGGTTTCATCATCAT
Primer 2	TGTAGCCTCTTTGCAAGACGTGTATTACAAGTTCTTTGTAGCTGTTTCATCATCAT
Primer 3	AGGCCATCAATGAGGCCACTATTATAGCCAGGCTTTCATCATCAT
Primer 4	GGGTTTCATTGGTACAGCCCATGTAGTGAACCCCTTTCATCATCAT
Primer 5	CGCGCTCATCACCATCCTTGCCGTTTCATCATCAT
Primer 6	GGTCTCTCTATGCAAACTCTCTTCATCGTGGTGAACGCTTTCATCATCAT
Primer 7	TCGTTTTGAAGCAAATCTCAAAGTTCATTCACTGTCCTCCTGCTTTTCATCATCAT
Primer 8	CAAGTGTCTTTCGCTCGTCACTAGTAGTTCGCTGTTTTCATCATCAT
Primer 9	GGTATCAACTTGGTGGACCAGTTATTGAAAACTTCTTGTGGTTCCTCACTTTCATCATCAT
Primer 10	GGACTTCCATGTCATCTTCGAGCATCCAGTCCGCTACTTTCATCATCAT
Primer 11	CGTAATCGTTTTGTTGTGTGGTTGTTGTTTCTGGTCAAGCGATTGCTTTCATCATCAT
Primer 12	GCTTGATCCTCCTGAATTAGTTTTGTGGTTGGACTGGAATGACCTTTCATCATCAT
Primer 13	GGTCTCCACGATTCAACCAATTCCGGCGTTGGATTAAAATTTCCTTTCATCATCAT
Primer 14	GGAGTTTTAAGGTCTTGTCTTTGACCGGTTTTGAGTTTCTCGGAGTCTTTCATCATCAT
Primer 15	CCAGTGACGGTTGAGAAAACACTCCCATTGATGGGTTTCATCATCAT
Primer 16	TGCGTGTGAACCGTCGTCATGACTTATGAGCTTTCATCATCAT
Hairpin	Hairpin.27 : ACATCATCATGGGCCTTTTGGCCCATGATGATGTATGATGATGTTTTTTTT
Imager	Fluor 27 /5ATTO550N/TTATGATGATGTATGATGATGT

772
773

774 **Table S2. Amplification efficacy (%) of reference genes, and the associated correlation coefficient (R^2).**

775 Efficacy expected values according to literature (De Keyser et al., 2013; Broeders et al., 2014; Veselenak et al., 2015;
776 Shehata et al., 2019; Stephan, Tilmes, et Hülskamp, 2019) and according to R^2 value.
777
778 Abbreviations: *PP2A*: Protein phosphatase 2A; *NbUBC*: Ubiquitin-conjugating enzyme 3; *NbTUB*: β-Tubulin; *NbLip*:
779 Lipopyl synthase 2, mitochondria; *GAPDH*: Glyceraldehyde 3-phosphate dehydrogenase.

Gene	Efficacy (%)	R^2
Expected values	**80-120**	**≥ 0.980**
PP2A	101.5	0.996
NbUBC	101.8	0.995
NbTUB	106.5	0.997
NbLip	91	0.997
GAPDH	97.6	0.991