Driving the evolution of C4 photosynthesis in order

Yong-yao Zhao[1], Yao xin Zhang[123], Xin qi Yu[1], Xin-guang Zhu[12]

1.Center of Excellence for Molecular Plant Sciences, Chinese Academy of Sciences
2.State Key Laboratory for Plant Molecular Genetics
3.University of Chinese Academy of Sciences

C4 photosynthesis is an efficient photosynthetic type (High water use efficiency, High nitrogen
use efficiency, High light use efficiency). Although occurring in only 3% plant species, C4
photosynthesis contributes 23% of terrestrial gross primary productivity. C4 photosynthesis has
been considered evolved from C3 photosynthesis, although a mass of theories and hypotheses
have been made, how does it drive C4 evolution remain unclear. Here, Flaveria genus, the leading
model used for C4 evolutionary study, was used to explain the evolutionary driver for C4
photosynthesis through eco-physiological, biochemical and anatomical analysis. Precipitation was
negatively associated with C4 metabolic enzyme activity and C4 flux during C4 evolution. Results
showed an increasing temperature and precipitation in the earlier evolutionary stage, meanwhile
vein density increasing. Decreased precipitation and risen temperature drove the emergence of
C2 photosynthesis and the transition from C2 to partial C4 cycle. There was a positive relationship
between CO2 compensation point, stomata density and precipitation, indicating decreasing
precipitation enhance photorespiration through changing stomata during C4 evolution. This study
proposed that different step during the evolution of C4 photosynthesis required different drivers
and enhanced water source should be needed in the initial step.

Introduction
Photosynthesis undergone evolution, resulting in different photosynthetic type. C4
photosynthesis is one of these types of photosynthesis, which evolved from the C3
photosynthesis. Compared to its C3 ancestor, the plant with C4 photosynthesis has higher water
use efficiency, nitrogen use efficiency and light use efficiency (Vogan and Sage 2011; Taylor et al.
2010). Although occurring in only 3% plant species, C4 photosynthesis contributes 23% of
terrestrial gross primary productivity(Still et al. 2003; Sage et al. 2012). The evolution of C4
photosynthesis go through multiple stages (Sage et al. 2012; Sage 2004; Schluter and Weber 2016;
Brautigam and Gowik 2016). First stage is preconditioning, when plants have C3 photosynthesis
(Sage et al. 2012). At this stage, plants acquired new cis-elements (Akyildiz et al. 2007); gene
duplication occurs (Akyildiz et al. 2007; Monson 2003); vein density increases (Marshall et al.
2007; Mckown and dengler 2007) and the number of plastids in the bundle sheath
increases(Sage et al. 2013). The second stage is the formation of proto-kranz, when plants still
perform C3 photosynthesis (Sage et al. 2013), the number of plastids continue to increase (Stata
et al. 2019; Sage et al. 2012) and the chloroplast and mitochondria localize to the centripetal wall
of bundle sheath cells, which can be thought to be the initial step of C2 photosynthesis
(Hattersley 1989; Sage et al. 2012). The third stage is the formation of C2 photosynthesis (Type I
species) (Schulze et al. 2013; Sage et al. 2012; Edwards and KU 1987). C2 photosynthesis is
characterize as the transfer of glycine decarboxylase (GDC) and photosynthetic apparatus from
mesophyll cell to bundle sheath cell(Schulze et al. 2013; Stata et al. 2019; Sage et al. 2013). The

released CO2 from photorespiration by GDC can moderately concentrate CO2 in the bundle
sheath (Keerberg et al. 2014; Sage et al. 2012). The fourth stage is the enhancement of C2
photosynthesis and the induction of C4 cycle (Type II species) (Schluter and Weber 2016; Sage et
al. 2018). The induction of C4 cycle is proposed as a mechanism to resolve the imbalance of
ammonia between mesophyll cell and bundle sheath cell (Mallmann et al. 2014). The fifth stage
is the formation of full C4 cycle (C4-like) (Stata et al. 2019). For this stage, mesophyll remains
Rubisco(Taniguchi et al. 2021; Monson et al. 1987; Stata et al. 2019; Sage et al. 2018). The final
stage is the optimization of C4 cycle (C4 species) (Sage et al. 2012), and mesophyll cell no longer
contains Rubisco (Stata et al. 2019).
The evolution of C4 photosynthesis was considered to be driven by high photorespiration that is
caused by the low atmosphere CO2 concentration, drought, salt and heat (Sage et al. 2012;
Brautigam and Gowik 2016; Sage 2004; Edwards and Smith 2010; Bromham and Bennett 2014).
The low atmosphere CO_2 concentration, drought and salt could decrease the ratio of intercellular
CO_2 to O_2 concentration, thereby enhancing photorespiration (Sage et al. 2012). Increasing
temperature enhance photorespiration by reducing CO2 relative to O2 in the chloroplast
stroma(Ogren 1984) and decrease the specificity of Rubisco for CO2 relative to O2(Ogren 1984;
Sage et al. 2012). Previous studies have showed that the decrease in atmosphere CO2
concentration and the appearance of dry, salty and hot environment are correlated with C4
evolution(Christin et al. 2008; Edwards and Smith 2010; Edwards and Still 2008; Bromham and
Bennett 2014), which suggests they are the drivers for C4 evolution. However, the above
researches may not be comprehensive enough. Given the multiple steps and the complexity
during C4 evolution, it is possible that different phase during C4 evolution is under different
driving force, rather than the same driving force persistently along the C4 evolution, which has
been few studied. Each step of C4 evolution should obtains fitness for plants (Heckmann et al.
2013; Sage et al. 2012). The increase in vein density is an important characteristics in initial stage
during C4 evolution(Sage et al. 2012; Sedelnikova et al. 2018; Kumpers et al. 2017; Sage et al.
2014) (Mckown and dengler 2007; Marshall et al. 2007; Williams et al. 2013), which should be to
prevent loss of water in the mesophyll and maintain stomatal open, since evaporation is
enhanced at this time(Brautigam and Gowik 2016; Sage et al. 2018) (Sage et al. 2012). Therefore,
available water content in this stage might be increased(Brautigam and Gowik 2016). This was
inconsistent with the notions that the origin of C4 photosynthesis promoted by drier
condition(Sage et al. 2018; Edwards and Smith 2010).Since C2 photosynthesis and C4
photosynthesis could largely reduce photorespiration(Sage 2004; Schluter and Weber 2016; Sage
et al. 2012), the formation of C2 photosynthesis and the transition from C2 to partial C4 might be
driven by the conditions which could enhance photorespiration(Schluter and Weber 2016; Sage
et al. 2018; Sage et al. 2012), which remains fewer studies.
Flaveria genus is the model system to study C4 evolution(Sage et al. 2012; Sage 2017; Schluter
and Weber 2016; Mallmann et al. 2014; Sage et al. 2013; Vogan and Sage 2011), and the
establishment of current C4 evolution pyramid model is largely based on the studies in the
Flaveria genus (Sage et al. 2013; Vogan and Sage 2011; Sage et al. 2012). Flaveria genus contain
numerous species that represent evolutionary step towards C4 photosynthesis, which includes
multiple intermediate stage during the evolution of C4 photosynthesis(Mckown and dengler 2007;
Sage et al. 2012; Schulze et al. 2013; Ku et al. 1991; Heckmann et al. 2013). Flaveria is the most
young genus among the known C4 linage, which suggested that it might be going through the

evolution(Schulze et al. 2013). Flaveria species has the close relationship, which make it ideal to
perform the comparison study (Christin et al. 2011; Sage et al. 2012). The ecology of the habitat
for the intermediate species between C3 and C4 species and C4 plants could reflect the
environmental conditions drive the evolution of C4 photosynthesis(Sage et al. 2012; Sage et al.
2018). Therefore, the ecology of Flaveria genus enable us to finely study the driving force of C4
evolution. In this study, we collected the ecological data of the C4 evolutionary model genus
Flaveria, associated with the biochemical, anatomical and physiological data, finally trying to
interpret the drivers and mechanism of C4 evolution. This study attempted to answer:1) how
does the conditions promote C4 evolution? 2) how does plant obtain fitness during C4 evolution?
98
Result
1, Most of Flaveria are distributed inland, and a few are distributed along the coast
The locations of Flaveria species were mapped to the America by downloading the longitude and
latitude of each Flaveria samples from Global Biodiversity Information Facility (GBIF) (Fig 1A,B).
We found most of Flaveria species are located inland (Fig 1A,B). A few species (bro, flo, lin) are
distributed along the coast (Fig 1B), which was consistent with the previous reports(Powell 1978).
With evolution, intermediates gradually spread to location of higher latitude (Fig 1B). The C4
species widespread and distribute all over the world (Fig 1A).
107
2, The temperature and precipitation of Flaveria habitat
The temperature and precipitation of the habitat for each Flaveria species were obtained from
the database worldclim(Fick and Hijmans 2017). The results from physiological, molecular
experiments and phylogenetic tree indicated the direction of C4 evolution could be well
described by species's order of phylogenetic tree towards C4 photosynthesis(Sage et al. 2012;
Sage et al. 2018; Schluter and Weber 2016). Along the evolutionary way towards C_4
photosynthesis (Ordinal species towards C4 photosynthesis in the phylogenetic tree), annual
mean temperature, mean temperature of warmest quarter, mean temperature of wettest
quarter and max temperature of warmest month significantly increased in the F. rob (C3) and F.
son (Type I) (Fig 2A, Supplementary fig 2, Supplementary fig 3, Supplementary file), compared
with F. Pri. However, it significantly decreased in the F.ang, compared with F.son (Type I) (Fig 2a,b,
supplementary file). For the mean temperature of wettest quarter, mean temperature of
warmest quarter, max temperature of warmest month, compared with F. son, the habitat of
species with C2 photosynthesis (F. ang, F. chl, F. pub, F. opp, F. ano, F. ram) have the lower mean
temperature (Fig 2A, supplementary fig 2D). The habitat of species F. flo and F. lin have the higher
mean temperature compared to other intermediate species (Fig 2a,b, table 1). Max temperature
of warmest month for F. son is extremely high (Fig 2d).
For the annual precipitation, precipitation of wettest quarter, precipitation of warmest quarter
and precipitation of wettest month, along the direction of C_4 evolution, the precipitations
significantly increased in the F. rob (C3), compared with front species(Fig 2B, Supplementary fig 2,
supplementary fig 3). However, it significantly decreased in F. son (Type I), and largely decrease in
the F. ang (Type I) (Fig 2A, Supplementary fig 2, Supplementary fig 3, Supplementary file). The
precipitation of warmest quarter is greatly higher than precipitation of coldest quarter, and
similar to precipitation of wettest quarter (Fig 2B, Supplementary fig 1). Expect for F. flo (Type II)
and F. lin (Type II), the precipitation of the species with C2 photosynthesis decrease persistently

along the direction of C4 evolution (Fig 2d,e,f, Supplementary fig 2b). In the latter of C4 evolution,
the precipitation was a bit of increase, and the variation was large (Fig 2d,e,f, Supplementary fig
2b).
Along the phylogenetic tree towards C4 photosynthesis, the initial increase in precipitation and
temperature (F. pri to F. rob) coincided with the initial appearance of the higher vein density(Fig 2,
supplementary fig 4).
139
3, The association between precipitation (or temperature) and C4 metabolism related traits
during the transition from proto-kranz to C4-like.
For the clade A, we found the relationship between (PEPC, PPDK and MDH activity) and
precipitation is described as one in which (PEPC, PPDK and MDH activity) decreases with
precipitation in the form of a negative power function, which can be described as a linear
relationship between log enzyme activity and log precipitation (Fig 3 A,B,C). The relationship
between C4 flux and precipitation can be described as the linear function (Fig 3D). For the clade B,
the relationship between log enzyme activity and log precipitation also can be described as the
linear function (Fig 3 I,J,K). But the linear relation between C4 flux and precipitation in clade B is
weaker(Fig 3L). The linear relation between log enzyme activity and log temperature is weaker
than that between Log enzyme activity and Log precipitation, whether in clade A or clade B (Fig
3E,F,M,N,G,H,O,P).
152
4, The change of temperature and precipitation between C3 and C2, C2 and initial C4, initial C4
and developed C4.
Previous study has reported that C3-C4 species can represent the transition between C3 and C4
photosynthesis(Heckmann et al. 2013), and F. ram has the enhanced C4 cycle flux, increased C4
cycle related enzyme activity and rised gene expression (Adachi et al. 2023; Mallmann et al. 2014;
Vogan and Sage 2011). According to the phylogenetic tree, F. ang was closer to F.ram as
compared with F.son, therefore, among all Flaveria species, F.son most likely to represent the
initial appearance of C2 photosynthesis in Flaveria; F. ram and F. ano most likely to represent the
initial appearance of C4 photosynthesis in clade A and clade B, respectively; F. Pal and F. Bro most
likely to represent the initial appearance of the fully functional C4 cycle for clade A and clade B.
The comparison of F. rob and F. son suggested that the first decreased precipitation along the
evolutionary way towards C4 photosynthesis concurrently occurred with the increased mean
temperature (Fig 4A,F). Precipitation of F.ano and F.ram is lower than that of ang, and
Temperature of ram and ano is higher than that of F. ang(Fig 4B,G,C,H). Precipitation of pal is
lower than that of F. ram, and precipitation of F. bro is lower than that of F. flo (Fig 4D,I,E,J).
However, temperature of F.Pal and F. Bro showed the inconsistent change, F. Pal is same as F. ram,
and F. bro is lower than F. flo (Fig 4D,I,E,J). Interestingly, compared with their front species, we
observed that the decreased precipitation for the species habitat (F. son, F.ram, F.flo, F.pal and
F.bro) concurrently appeared with the decreased stomatal density, decreased inhibition of O2
and CO2 compensation point, decreased maximum stomatal conductance (Supplementary Fig
4E,F,G,H). Although precipitation of F. flo and F. lin were increased compared to other species
with C2 photosynthesis (Fig 3b), they also have the decreased stomatal density, maximum
stomatal conductance and increased stomatal size, which might attribute to their salty habitat.
176

5, The association between precipitation, stomata and CO_2 compensation point.

Previous work found there is a difference in stomata in Flaveria genus(Zhao et al. 2022). Since

stomata have long been thought to be associated with water content, we checked the

relationship between stomata and precipitation in Flaveria genus. We found there was a positive

and linear relationship between the stomata, CO2 compensation point and precipitation during

the transition between proto-kranz and C4-like in Flaveria clade A and clade B(Fig 5A,B,C,D,E,F).

The linear relationship between stomata, CO2 compensation point and precipitation in Flaveria

clade B was weaker than that in clade A (Fig 5A,B,C,F,G,H). The relation between temperature

and stomata (or temperature and CO2 compensation point) was not apparent as the relation

between these two parameters and precipitation(Fig 5D,I,E,J).

Discussion

Evolutionary conditions along the path towards C4 photosynthesis.

The heat and drought have long been thought to be associated with the evolution of C4

photosynthesis(Sage et al. 2018; Taylor et al. 2012). Discovering the environmental conditions for

the habitats of C4 and C3-C4 species could contribute to explain the drivers for C4 evolution(Sage

et al. 2018; Sage et al. 2012; Edwards and Smith 2010). In this study, we examined the data for

precipitation and temperature of the diverse Flaveria species. The precipitation and temperature

of different time-scale was shown (Fig 2a, b, supplementary fig 1, 2, 4). We noticed that the

trajectory trend in the wettest quarter mostly coincides with the warmest quarter(Fig 2,

supplementary fig 2) and mean temperature of wettest quarter was overall higher than mean

temperature of driest quarter (Supplementary fig 2D, Supplementary fig 1E), suggesting that the

habitat of Flaveria is raining and hot during the same period. Sufficient water content is

necessary to maintain high photosynthetic activity for plants, and the description from previous

work indicated that Flaveria species perform higher photosynthesis during summer (Powell 1978).

Therefore, this period should approximately be the summer in Mexico, and Flaveria have higher

photosynthetic activity during this time as compared with the rest of time (Powell 1978)

(Fig2a,b,d,e, supplementary fig 1). In the perspective of long-term, higher photosynthetic activity

indicating higher photorespiratory flux in C3 plants (Previous work has proved that there is no

significant difference in the Sc/o between different Flaveria species(Kubien et al. 2008), and Ci

between the C3 and type I species has no significant difference(Sage et al. 2013)), thus C3 and

initial intermediate Flaveria species in this period would have the potential of high

photorespiratory flux which has long been thought to be the driver force for the origin of C4

plants(Sage et al. 2012; Sage et al. 2018). Due to high photosynthetic activity and high

photorespiratory flux, this period for Flaveria is valuable to explain the driving of C4 evolution (Fig

2a,b,c,d,e,f).

The evolutionary selective pressure on photosynthesis should act on the period that

photosynthesis function during a year(Sage et al. 2018), therefore the driest and coldest season

should not be considered (Supplementary fig 1a,b,c,d,e,f). Since Flaveria genus is the leading

model for C4 evolutionary study, and the habitat ecology for C4 species and the intermediate

species that branch the lineage of C3 and C4 species could reflect the environmental conditions

drive the evolution of C4 photosynthesis, three type of rainy and warm periods suggested that

the conditions promote C4 evolution step by step and in order. Specifically, the initial stage during

C4 evolution need rising precipitation and temperature, and later stage need decreased

precipitation (Fig 2a,b,c,d,e,f, supplementary fig 2,a,b). Heat previously is considered to be the
factor that promote the origin of C4 photosynthesis(Sage et al. 2018), however, our study
proposed that, in addition to increasing temperature, precipitation increasing is a vital factor to
launch the evolution of C4 photosynthesis (Fig 2a,b,c,d,e,f). Previous study proposed it is possible
that heat occurred with drought simultaneously to enhance photorespiration and then promote
the evolution of C4 photosynthesis(Sage et al. 2012), our study indicated that this process indeed
exist (Fig 2a,b,c,d,e,f), and specifically, this process is the appearance of C2 photosynthesis (son)
and partial C4 (F. ram, F. ano). Although the precipitation of the Type II species F. flo and F. lin was
high (Fig 2), they inhabit contain saline region (Fig 1, table 1), which even under high
precipitation may experience strong osmotic stress. The habitat of C4 species was various,
indicating that adaptation extent of C4 plants will be wider (Fig1, table1).
232
The role of water content and temperature
Some studies have proposed that drought is driver force for C4 evolution(Sage 2004; Zhou et al.
2018; Sage et al. 2018; Schluter and Weber 2016; Edwards and Smith 2010). Some studies
suggested that heat is the crucial environment to drive the C4 evolution(Sage et al. 2018; Sage et
al. 2012). In this study, our result proposed that drought play a more important role than heat
during C4 evolution. During the transition between C3 and C4 photosynthesis, precipitation
decline step by step, however, there is no clear trend for temperature (Fig 2). Whether the clade
A or Clade B, there showed a negative association between core C4 enzyme activity and
precipitation (Fig 3), and there is a strong negative relationship between precipitation and core
C4 enzyme activity in clade A (Fig3). However, the relationship isn't apparent for temperature (Fig
3). These suggested that, in terms of the whole evolutionary process, drought might be the main
driver for the transition from C3 to C4 photosynthesis rather than heat.
Initial step is key for C4 evolution(Sage et al. 2012). For this step, the increased vein density and
increased number of chloroplasts in bundle sheath is the important characters during the C4
evolution(Sage et al. 2012). The increased vein density has been deducted to be used to maintain
enough water content in the leaf and avoid the reduce of stomata aperture, since the
evaporation might be increased in this stage(Sage et al. 2012; Sage et al. 2018). In agreement
with this statement, our data showed that, when was at the stage of vein density increasing,
there was no change in stomatal development and temperature was indeed increased (Fig 2A,
supplementary fig 4). The increased temperature could increase vapor pression between leaf and
air (supplementary fig 5B), hence enhance the evaporation(Sage et al. 2018); while no change in
the stomatal development indicate that water shortage in leaf might not happen, since water
shortage could alter stomatal development(Franks et al. 2009; Taylor et al. 2012). Our data
showed that precipitation was increased in this stage, which matched with the increased vein
density, together ensure the enough water content in the leaf (Fig 2B). Due to the sufficient
supply of water, stomatal aperture can be maintained. Maintaining stomatal open could not only
maintain intercellular CO2 concentration, but also could help alleviate the increase of leaf
temperature through transpiration, which benefits for maintaining efficient function of
photosynthesis. The increase in vein density inevitably decreases proportion of mesophyll in a
leaf, decreasing the photosynthetic capacity of the leaf. In theory, this should create a strong
evolutionary pressure to select for the plants with increased photosynthetic capacity of BSC to
offset the loss of photosynthetic capacity. Indeed, the relative number and size of chloroplast of

vascular bundle sheath in F. rob increase(Sage et al. 2013). Besides, the light intensity of F. rob
habitat is higher than that of F. pri (supplementary fig 5A), which further enhances the
photosynthesis of bundle sheath. The size of vascular bundle sheath in F. rob decrease compared
with F. pri(Sage et al. 2013; Mckown and dengler 2007), the size of mesophyll cell decrease
compared with F. pri(Mckown and dengler 2007), and thickness of leaf become thin(Mckown and
dengler 2007), these all enhancing the intensity of the light that cast vascular bundle sheath,
rising the probability of the selection for bundle sheath with higher photosynthetic capacity.
272
Different evolutionary step promoted by different driver force
A mass of researches focused on the driver force of C4 evolution. For example, decreasing
atmosphere CO2 concentration has been thought to be associated to the driving force of C4
evolution(Christin et al. 2008; Concentrations and Paleogene). Drought, salt and heat is also
thought to be the conditions benefit for C4 evolution(Sage et al. 2018; Sage 2004; Bromham and
Bennett 2014) (Edwards and Still 2008; Edwards and Smith 2010; Pearcy and Ehleringer 2006;
Sage et al. 2012). However, most of these studies suggest that C4 evolution promoted by single
ecological condition in the entire evolutionary process. Since C4 photosynthesis is the complex
traits, we proposed that single and same ecological driver is difficult to achieve the C4
photosynthesis. Since Flaveria species is an excellent representative for the different step during
C4 evolution, in this work, from the results of the study on Flaveria genus, which supported that
the different step of the C4 evolution is driven by the different driver mechanism.
After the initial step of C4 evolution, the promotion of C2 photosynthesis is considered to be a
crucial step during C4 evolution (Sage et al. 2012; Schluter and Weber 2016). Even though, there
is a viewpoint that the way to C4 evolution could be turned into the way to C2 evolution
(Brautigam and Gowik 2016). Our data indicated increasing temperature and decreasing
precipitation simultaneously promote the emergence of C2 photosynthesis (Fig 3). The increase
in temperature and decrease in precipitation could largely enhance photorespiration, promoting
the selection of the transfer of GDC from mesophyll to bundle sheath (photorespiration CO2
pump), which could improve the fitness of plants. Previous work has proposed that drought
enhance photorespiration through close stomata(Sage et al. 2012). In this study, we provided the
evidence that stomata play a role in the process that decreased precipitation enhanced
photorespiration, supported by the relation between stomatal density and CO2 compensation
point (Fig 5). The photorespiratory CO2 pump could largely weaken the photorespiration(Sage et
al. 2018; Sage et al. 2012; Schluter and Weber 2016; Ku et al. 1991), which indicated by the
largely decrease in CO2 compensation point and O2 inhibition(Sage et al. 2013; Ku et al. 1991).
The gradual enhancement of C2 photosynthesis mainly associate to the decreasing precipitation
(Fig 2a,b,c,d,e,f, Fig 3). The gradual enhancement of C2 photosynthesis has long been considered
to cause the imbalance of nitrogen between mesophyll cell and bundle sheath cell, which induce
the partial C4 cycle(Mallmann et al. 2014). In case of the progress from the C2 photosynthesis to
partial C4 photosynthesis, precipitation decreases and temperature increases, just like the former
step (Fig 4 B,C,G,H). After the induce of partial C4 photosynthesis, the formation of fully
functional C4 photosynthesis was drived by inconsistent conditions for Clade A and Clade B.
Precipitation was significantly decreased both in clade A and clade B, but there is no difference in
temperature in clade A(Fig 4), even though, temperature was decreased in clade B(Fig 4). These
results suggested that precipitation was the driving force rather than temperature at this

transitional stage. Lacking of photosynthetic capacity and growing under high temperature,
plants suffering the carbon limitation in the early phase during C4 evolution, suggesting the main
driving force for C4 evolution at this phase is carbon limitation. After the preconditioning for C2
photosynthesis, plants suffering water limitation and water limitation lead to deceasing of carbon
acquirement, which gradually established C2 photosynthesis and further partial C4
photosynthesis. Therefore, the main driving reason for C4 evolution at these phases is the water
limitation.
316
317
Methods
1, Obtain the ecological data of Flaveria species.
First, the geographical distribution of Flaveria was acquired from Global Biodiversity Information
Facility (GBIF) database using the rgibf software package. Using ArcMAP software, the elevation
data of the respective locations and 19 Bioclimatic factors, including Mean Diurnal Range, annual
mean Temperature, Isothermality, Temperature Seasonality, Max Temperature of Warmest
Month, Temperature Annual Range, Mean Temperature of Wettest Quarter, Mean Temperature
of Warmest Quarter, Annual Precipitation, Precipitation of Wettest Month, Precipitation of
Warmest Quarter, Precipitation of Wettest Quarter, Precipitation of Coldest Quarter, Min
Temperature of Coldest Month, Precipitation of Driest Quarter, Mean Temperature of Coldest
Quarter, Precipitation of Driest Month, Precipitation Seasonality (Coefficient of Variation),
Precipitation of Coldest Quarter, were acquired from the Worldclim database based on the
geographical distribution information of the Flaveria. Specifically, downloading the data from the
worldclim at first. Secondly, preparing the file for the information of sampling location. Thirdly,
importing the layers file of climate factor. Fourthly, importing the information of sampling
location. Fifth, generating the new layers. Lastly, exporting the file that including sample and the
data of geographical factors. All bioclimatic variables are averages from 1970 to2000. The
elevation data came from SRTM elevation data. All data from the worldclim database have a
maximum spatial resolution of 30 arc-seconds. The water vaper pression and solar radiation were
also extracted from the World clim database(Fick and Hijmans 2017) based on the geographical
distribution information of Flaveria.
2,Obtain the geology, anatomical and physiological data of Flaveria species.
The anatomical and physiological data for Flaveria was acquired from previous research. The data
of veinal data was from (Mckown and dengler 2007). The data of CO2 compensation point and
O2 inhibition is from (Ku et al. 1991). The data related to stomata is from (Zhao et al. 2022). The
C4 metabolic enzyme activity data is from (Adachi et al. 2023) (supplementary table S7). The data
in the graph was digitized by Origin 2018. The growth habitat is aquired from(Powell 1978;
Kocacinar et al. 2008).
3, Map the Flaveria species.
We mapped the location of Flaveria species using the software google earth Pro according the
latitude and longtitude which is extracted from Global Biodiversity Information Facility (GBIF).
4, Statistical analysis, regression analysis and Graph drawing.
The graph was plotted and the regression was performed by using graphpad prism 8. The
statistic test was performed by graphpad prism 8 and WPS office. The statistic test for the
difference in precipitation or temperature between all Flaveria species was performed by

one-way ANOVA and Tukey's multiple test (supplementary file). The rest of other statistic test was
performed through T-test. The precipitation of the wettest quarter and the C4 metabolic enzyme
activity took the logarithm with 2 as the base, and then they were performed linear regression by
graphpad prism 8. The precipitation of the wettest quarter and the C4 metabolic enzyme activity
was directly performed linear regression using graphpad prism 8. The equations for the
regression were showed in supplementary file. The type of the statistic test used in each analysis
was stated in the supplementary file.

Table 1 The habitat of Flaveria species

Species (photosynthetic type)	Growth habitat	Geographic location
Flaveria pringlei (C3)	Rather arid and gypseous habitats, scattered localities, waste places, cultivated fields, roadsides	Southern Mexico
Flaveria cronquistii (C3)	Rocky limestone (possibly gypseous) areas	Southern Mexico
Flaveria robusta (C3)	Gypsum and slate, sandy clay, or open rocky slopes, weedy at roadsides	Western Mexico
Flaveria sonorensis (Type I)	Close to warm mineral springs	North-western Mexico
Flaveria angustifolia (Type I)	Sandy or loamy soils in open fields	Southern Mexico
Flaveria pubescens (Type I)	Scattered localities, gypsum soils	Central Mexico
Flaveria oppositifolia (Type I)	high gypsum plains or other areas with dilute gypsum, saline, or rarely loamy soil, usually in moist areas or in disturbed habitats at roadside	Central Mexico
Flaveria linearis (Type II)	Near the coast and on offshore islands	Western Florida, USA
Flaveria floridana (Type II)	Restricted to saline, white sand near the beaches of brackish marshes	Western Florida, USA
Flaveria ramosissima (Type II)	Sandy-clay bottoms, along dry washes, fields and roadsides, possibly in gypseous soils,	Southern Mexico
Flaveria anomala (Type II)	High gypsum plains, often in disturbed sites	Central Mexico
Flaveria palmeri (C4-like)	Gypsum flats and knolls, or clay-loam soils probably with dilute gypsum, bolsons near lagunas, distributed areas near roads and fields	Northern Mexico
Flaveria brownie (C4-like)	Saline, sandy, and marshy areas of the coastal flats and islands of the lower Texas Gulf Coast	North-eastern Mexico
Flaveria vaginata (C4-like)	Arid and gypseous soils, scrub vegetation, cultivated fields	South-central Mexico

Flaveria bidentis (C4)	A weed in moist places on waste ground, near rivers, streams, and barrancas, in fields, pastures, disturbed ground, near streets and sidewalks in towns, claygravel, or sand	Africa, Caribbean, South America, southern USA
Flaveria trinervia (C4)	a variety of habitats, usually saline, gypseous, and disturbed, usually near ephemeral or permanent water sources	Africa, Caribbean, Central/South America, Mexico, USA
Flaveria chloraefolia (Type I)	Permanent or ephemeral saline marshy soils, springs, streams, roadsides	North Mexico

The information of habitat of Flaveria species was from (Powell 1978; Kocacinar et al. 2008)

Figure legend

Fig 1 Distribution of the Flaveria species

A, The location of Flaveria species was mapped to the world map. The main habitat of Flaveria is

Mexico which was displayed. B, The location of Flaveria species except C4 species was mapped to

the world map. C,D, The latitude and longitude of Flaveria species. The latitude and longitude

was acquired from the GBIF. Different colour indicate different photosynthetic type.

Fig 2 The environment conditions of the habitat for Flaveria species showed rhythmic change

A, mean temperature of warmest quarter of Flaveria species. B, precipitation of wettest quarter

of Flaveria species. C,mean temperature of wettest quarter for F. rob and F. pri. D, precipitation of

wettest quarter for F.rob and F.pri. E, the change in vein density in the C3 and intermediate

species. Different colour indicates different photosynthetic types. Orange indicates C3, grey

indicates intermediate species, green indicates C4-like, blue indicates C4. Black lines indicate

median.

Fig 3 The association between precipitation (temperature) and C4 enzyme activity or C4 flux

The precipitation indicating precipitation of the wettest quarter. The temperature indicating

mean temperature of the wettest quarter. (A-H, clade A). A, the relation between Log PEPC

activity per chl and Log precipitation. B, the relation between Log PPDK activity per chl and Log

precipitation. C, the relation between Log MDH activity per chl and Log precipitation.D, the

relation between C4 flux and precipitation. E, the relation between Log PEPC activity per chl and

Log temperature. F, the relation between Log PPDK activity per chl and Log temperature. G, the

relation between Log MDH activity per chl and Log temperature. H, the relation between C4 flux

and log temperature. (I-P, clade B). I, the relation between Log PEPC activity per chl and log

precipitation. J, the relation between Log PPDK activity per chl and log precipitation. The change

trend of precipitation of wettest quarter. K, the relation between Log MDH activity per chl and log

precipitation. L, the relation between C4 flux and precipitation. M, the relation between log PEPC

activity per chl and log temperature. N, the relation between log PPDK activity per chl and log

temperature. O, the relation between log MDH activity per chl and log temperature. P, the

relation between C4 flux and temperature. (a), mean temperature of wettest quarter (b), O2

inhibition (c), CO2 compensation point (d), vein density (e), distance between vein (f), Maximum

stomatal conductance (gsmax) (g) and stomatal density (SD) (h) from C3 to C4 photosynthesis in

Flaveria genus. Different colour indicates different photosynthetic types. Orange indicates C3,

purple indicates Type I, yellow indicates Type II, green indicates C4-like, blue indicates C4.

Fig 4 Precipitation and temperature during the transition from proto-kranz to C4-like

A, precipitation of the wettest quarter during the transition from proto-kranz(F.rob) to C2(F.son).

B, precipitation of the wettest quarter during the transition from C2 (F.son) to partial C4 (F.ano) in

clade B. C, Precipitation of the wettest quarter during the transition from proto-kranz(F.son) to

partial C4 in clade A. D, Precipitation of the wettest quarter during the transition from partial C4

to fully functional C4 photosynthesis. E, Precipitation of the wettest quarter during the transition

from partial C4 to fully functional C4 photosynthesis. F, Mean temperature of the wettest quarter

during the transition from proto-kranz (F.rob) to C2(F.son). G, Mean temperature of the wettest

quarter during the transition from C2 (F.son) to partial C4 (F.ano) in clade B. H, Mean

temperature of the wettest quarter during the transition from C2(F.son) to partial C4 (F.ram) in

clade A. I, Mean temperature of the wettest quarter during the transition from partial C4 to fully

functional C4 photosynthesis (F.pal) in clade A. J, Mean temperature of the wettest quarter

during the transition from partial C4 (F.ano) to fully functional C4 photosynthesis (F.bro) in clade

B. The C4 evolution was launched by increased temperature and increased precipitation. The

latter of C4 evolution was promote by increased temperature and decreased precipitation.

Fig 5 The association between stomatal density, CO2 compensation point, precipitation and

temperature

(A-E, clade A) A, The relation between stomatal density and precipitation. B, the relation

between CO2 compensation point and precipitation. C, The relation between stomatal density

and CO2 compensation point. D, The relation between stomatal density and temperature. E, the

relation between CO2 compensation point and temperature. (F-J, clade B) F, The relation

between stomatal density and precipitation. G, the relation between CO2 compensation point

and precipitation. H, The relation between stomatal density and CO2 compensation point. I, the

relation between stomatal density and temperature. J, the relation between CO2 compensation

point and temperature.

Fig 6, Schematic diagram for the driver process of C4 evolution

The schematic diagram for the conditions driving the evolutionary process for C4 photosynthesis.

The raindrop number denotes the precipitation. The thermometer denotes the temperature. The

arrows denote the direction of C4 evolution. The cycle composed by white and black denotes the

main driving element that promote the corresponding stage of C4 evolution. The driving reason

in C3 and proto-kranz was mainly carbon limitation, and the driving reason in the latter stage was

mainly water limitation. More and more black indicating the carbon limitation was increasing

stepwise, and more and more white indicating the water limitation was increasing stepwise.

Supplementary fig 1, Temperature and precipitation of habitat for Flaveria species in driest and

coldest period.

A, precipitation of driest quarter. B, precipitation of coldest quarter. C, precipitation of driest

month. D, Mean temperature of driest quarter. E, Mean temperature of coldest quarter. F, Min

temperature of coldest month. Orange indicates C3, purple indicates Type I, yellow indicates Type

II, green indicates C4-like, blue indicates C4. Black lines indicate median.

Supplementary fig 2, Temperature and precipitation of habitat for Flaveria species in wettest and

warmest period. Orange indicates C3, purple indicates Type I, yellow indicates Type II, green

indicates C4-like, blue indicates C4. Black lines indicate median.

Supplementary fig 3, annual temperature and precipitation of habitat for Flaveria species

a, Annual mean temperature. b, Annual precipitation. Orange indicates C3, purple indicates Type

I, yellow indicates Type II, green indicates C4-like, blue indicates C4. Black lines indicate median.

Supplementary fig 4, The change trend for the evolution towards C4 photosynthesis.

A, Precipitation of wettest quarter. B, Mean temperature of wettest quarter. C, Vein density. D,

Distance between vein. E, O2 inhibition. F, CO2 compensation point. G, Gs_{max}. H, stomatal density

(SD).

Supplementary fig 5, The difference in solar radiation and water vaper pression between F. pri

and F. rob.

A, The difference in solar radiation between F.pri and F.rob. B, The difference in water vaper

pression between F.pri and F.rob.

Table 1 The habitat for Flaveria species

Supplementary file The statistics test and regression analysis

Adachi S, Stata M, Martin DG, Cheng S, Liu H, Zhu X-G, Sage RF (2023) The Evolution of C4

Photosynthesis in Flaveria (Asteraceae): Insights from the Flaveria linearis Complex. Plant physiology

191 (1):233-251. doi:10.1093/plphys/kiac467

Akyildiz M, Gowik U, Engelmann S, Koczor M, Streubel M, Westhoff P (2007) Evolution and function of

a cis-regulatory module for mesophyll-specific gene expression in the C_4 dicot Flaveria trinervia. Plant

Cell 19 (11):3391-3402. doi:10.1105/tpc.107.053322

Brautigam A, Gowik U (2016) Photorespiration connects C3 and C4 photosynthesis. J Exp Bot 67

(10):2953-2962. doi:10.1093/jxb/erw056

Bromham L, Bennett TH (2014) Salt tolerance evolves more frequently in C4 grass lineages. Journal of

Evolutionary Biology 27 (3):653-659. doi:10.1111/jeb.12320

Christin PA, Besnard G, Samaritani E, Duvall MR, Hodkinson TR, Savolainen V, Salamin N (2008)

Oligocene CO2 decline promoted C4 photosynthesis in grasses. Curr Biol 18 (1):37-43.

doi:10.1016/j.cub.2007.11.058

Christin PA, Osborne CP, Sage RF, Arakaki M, Edwards EJ (2011) C(4) eudicots are not younger than C(4)

monocots. J Exp Bot 62 (9):3171-3181. doi:10.1093/jxb/err041

Concentrations MDiACD, Paleogene Dt <science.1110063.pdf>.

Edwards EJ, Smith SA (2010) Phylogenetic analyses reveal the shady history of C4 grasses. Proc Natl

Acad Sci U S A 107 (6):2532-2537. doi:10.1073/pnas.0909672107

Edwards EJ, Still CJ (2008) Climate, phylogeny and the ecological distribution of C4 grasses. Ecol Lett 11

(3):266-276. doi:10.1111/j.1461-0248.2007.01144.x

Edwards Ge, KU Msb (1987) Biochemistry of C3-C4 Intermediates. The Biochemistry of Plants, 10

Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land

areas. International Journal of Climatology 37 (12):4302-4315. doi:10.1002/joc.5086

Franks PJ, Drake PL, Beerling DJ (2009) Plasticity in maximum stomatal conductance constrained by

negative correlation between stomatal size and density: an analysis using Eucalyptus globulus. Plant,

cell & environment 32 (12):1737-1748. doi:10.1111/j.1365-3040.2009.002031.x

Hattersley RHBaPW (1989) Leaf Anatomy of C_3-C_4 Species as Related to Evolution of C_4
Photosynthesis.pdf. Plant physiology 91:1543-1550

Heckmann D, Schulze S, Denton A, Gowik U, Westhoff P, Weber AP, Lercher MJ (2013) Predicting C4
photosynthesis evolution: modular, individually adaptive steps on a Mount Fuji fitness landscape. Cell
153 (7):1579-1588. doi:10.1016/j.cell.2013.04.058

Keerberg O, Parnik T, Ivanova H, Bassuner B, Bauwe H (2014) C_2 photosynthesis generates about 3-fold
elevated leaf CO_2 levels in the C_3-C_4 intermediate species Flaveria pubescens. J Exp Bot 65
(13):3649-3656. doi:10.1093/jxb/eru239

Kocacinar F, McKown AD, Sage TL, Sage RF (2008) Photosynthetic pathway influences xylem structure
and function in Flaveria (Asteraceae). Plant, cell & environment 31 (10):1363-1376.
doi:10.1111/j.1365-3040.2008.01847.x

Ku MS, Wu J, Dai Z, Scott RA, Chu C, Edwards GE (1991) Photosynthetic and photorespiratory
characteristics of flaveria species. Plant physiology 96 (2):518-528. doi:10.1104/pp.96.2.518

Kubien DS, Whitney SM, Moore PV, Jesson LK (2008) The biochemistry of Rubisco in Flaveria. Journal
of Experimental Botany 59 (7):1767-1777. doi:10.1093/jxb/erm283

Kumpers BM, Burgess SJ, Reyna-Llorens I, Smith-Unna R, Boursnell C, Hibberd JM (2017) Shared
characteristics underpinning C4 leaf maturation derived from analysis of multiple C3 and C4 species of
Flaveria. J Exp Bot 68 (2):177-189. doi:10.1093/jxb/erw488

Mallmann J, Heckmann D, Brautigam A, Lercher MJ, Weber AP, Westhoff P, Gowik U (2014) The role of
photorespiration during the evolution of C_4 photosynthesis in the genus Flaveria. Elife 3:e02478.
doi:10.7554/eLife.02478

Marshall DM, Muhaidat R, Brown NJ, Liu Z, Stanley S, Griffiths H, Sage RF, Hibberd JM (2007) Cleome,
a genus closely related to Arabidopsis, contains species spanning a developmental progression from
C(3) to C(4) photosynthesis. Plant J 51 (5):886-896. doi:10.1111/j.1365-313X.2007.03188.x

Mckown A, dengler N (2007) Key innovations in the evolution of Kranz anatomy and C4 vein pattern in
Flaveria (Asteraceae). American Journal of Botany 94:382–399

Monson, William S. Schuster, Ku MSB (1987) Photosynthesis in Flaveria brownil A.M. Powell. Plant
physiology 85:1063-1067

Monson RK (2003) Gene duplication, neofunctionalization, and the evolution ofC4 photosynthesis. Int
J Plant Sci

Ogren DBJaWL (1984) The CO2_O2 specificity of ribulose 1,5-bisphosphate carboxylase oxygenase_
dependence on ribulosebisphosphate concentration, pH and temperature. planta 161:308-313

Pearcy RW, Ehleringer J (2006) Comparative ecophysiology of C3 and C4 plants. Plant, cell &
environment 7 (1):1-13. doi:10.1111/j.1365-3040.1984.tb01194.x

Powell AM (1978) Systematics of Flaveria (Flaveriinae--Asteraceae). Annals of the Missouri Botanical
Garden 65:590-636

Sage RF (2004) The evolution of C_4 photosynthesis. New Phytologist 161:341-270.
doi:10.1046/j.1469-8137.2004.00974.x

Sage RF (2017) A portrait of the C4 photosynthetic family on the 50th anniversary of its discovery:
species number, evolutionary lineages, and Hall of Fame. J Exp Bot 68 (2):4039-4056.
doi:10.1093/jxb/erx005

Sage RF, Khoshravesh R, Sage TL (2014) From proto-Kranz to C4 Kranz: building the bridge to C4
photosynthesis. J Exp Bot 65 (13):3341-3356. doi:10.1093/jxb/eru180

Sage RF, Monson RK, Ehleringer JR, Adachi S, Pearcy RW (2018) Some like it hot: the physiological
ecology of C4 plant evolution. Oecologia 187 (4):941-966. doi:10.1007/s00442-018-4191-6

Sage RF, Sage TL, Kocacinar F (2012) Photorespiration and the evolution of C_4 photosynthesis. Annu
Rev Plant Biol 63:19-47. doi:10.1146/annurev-arplant-042811-105511

Sage TL, Busch FA, Johnson DC, Friesen PC, Stinson CR, Stata M, Sultmanis S, Rahman BA, Rawsthorne
S, Sage RF (2013) Initial events during the evolution of C4 photosynthesis in C3 species of Flaveria.
Plant physiology 163 (3):1266-1276. doi:10.1104/pp.113.221119

Schluter U, Weber AP (2016) The Road to C4 Photosynthesis: Evolution of a Complex Trait via
Intermediary States. Plant Cell Physiol 57 (5):881-889. doi:10.1093/pcp/pcw009

Schulze S, Mallmann J, Burscheidt J, Koczor M, Streubel M, Bauwe H, Gowik U, Westhoff P (2013)
Evolution of C4 photosynthesis in the genus flaveria: establishment of a photorespiratory CO2 pump.
Plant Cell 25 (7):2522-2535. doi:10.1105/tpc.113.114520

Sedelnikova OV, Hughes TE, Langdale JA (2018) Understanding the Genetic Basis of C_4 Kranz Anatomy
with a View to Engineering C_3 Crops. Annu Rev Genet 52:249-270.
doi:10.1146/annurev-genet-120417-031217

Stata M, Sage TL, Sage RF (2019) Mind the gap: the evolutionary engagement of the C4 metabolic
cycle in support of net carbon assimilation. Curr Opin Plant Biol 49:27-34.
doi:10.1016/j.pbi.2019.04.008

Still CJ, Berry JA, Collatz GJ, DeFries RS (2003) Global distribution of C3and C4vegetation: Carbon cycle
implications. Global Biogeochemical Cycles 17 (1):6-1-6-14. doi:10.1029/2001gb001807

Taniguchi YY, Gowik U, Kinoshita Y, Kishizaki R, Ono N, Yokota A, Westhoff P, Munekage YN (2021)
Dynamic changes of genome sizes and gradual gain of cell-specific distribution of C4 enzymes during
C4 evolution in genus Flaveria. Plant Genome:e20095. doi:10.1002/tpg2.20095

Taylor SH, Franks PJ, Hulme SP, Spriggs E, Christin PA, Edwards EJ, Woodward FI, Osborne CP (2012)
Photosynthetic pathway and ecological adaptation explain stomatal trait diversity amongst grasses.
New Phytol 193 (2):387-396. doi:10.1111/j.1469-8137.2011.03935.x

Taylor SH, Hulme SP, Rees M, Ripley BS, Woodward FI, Osborne CP (2010) Ecophysiological traits in C_3
and C_4 grasses: a phylogenetically controlled screening experiment. New Phytol 185 (3):780-791.
doi:10.1111/j.1469-8137.2009.03102.x

Vogan PJ, Sage RF (2011) Water-use efficiency and nitrogen-use efficiency of C_3-C_4 intermediate
species of Flaveria Juss. (Asteraceae). Plant, cell & environment 34 (9):1415-1430.
doi:10.1111/j.1365-3040.2011.02340.x

Williams BP, Johnston IG, Covshoff S, Hibberd JM (2013) Phenotypic landscape inference reveals
multiple evolutionary paths to C_4 photosynthesis. Elife 2:e00961. doi:10.7554/eLife.00961

Zhao YY, Lyu MA, Miao F, Chen G, Zhu XG (2022) The evolution of stomatal traits along the trajectory
toward C4 photosynthesis. Plant physiology 190 (1):441-458. doi:10.1093/plphys/kiac252

Zhou H, Helliker BR, Huber M, Dicks A, Akcay E (2018) C_4 photosynthesis and climate through the lens
of optimality. Proc Natl Acad Sci U S A 115 (47):12057-12062. doi:10.1073/pnas.1718988115

Clade A

CladeB

Clade A   Clade B

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

The trajectory towards C4

C3 Type I Type II C4-like C4

Supplementary Figure 4

Supplementary Figure 5