1   The driving process for the evolutionay trajectory towards C_4 photosynthesis
2

3   Yong-Yao Zhao[1], Yao Xin Zhang[123], Xin qi Yu[1], Xin-Guang Zhu[12]
4
5   1. State Key Laboratory of Carbon Capture, Center of Excellence for Molecular Plant Sciences,
6   Chinese Academy of Sciences
7   2. State Key Laboratory for Plant Molecular Genetics
8   3. University of Chinese Academy of Sciences
9
10   Author for correspondence:
11   Xin-Guang Zhu
12   zhuxg@cemps.ac.cn

13
14

Compared to the plants with C_3 photosynthesis, the plants with C_4 photosynthesis has higher
light, water, and nitrogen use efficiencies. Though only 3% of higher plants have C_4
photosynthesis, C_4 photosynthesis contribute 23% of terrestrial gross primary productivity.
Though much effort has been made, the driving process for the evolutionary trajectory from C_3
towards C_4 photosynthesis remain elusive. Here, Flaveria genus, the model for C_4 evolutionary
researches, was used to study the driving procedure through a combined analysis of
eco-physiological, biochemical and anatomical data. Results showed that precipitation was
negatively associated with C_4 metabolic enzyme activity and C_4 flux from proto-Kranz to C_4-like.
An increased temperature and precipitation concurrent with increased vein density were observed
during the earlier evolutionary stage. Then, decreased precipitation and increased temperature
accompanied the emergence of C_2 photosynthesis and further transition from C_2 to developing C_4
photosynthesis. There was a positive relationship between CO_2 compensation point, stomata
density and precipitation, suggesting decreasing precipitation enhance photorespiration through
closing stomata during evolution. As the driving force from proto-Kranz to C_4-like, drought is
more important than heat during C_4 evolution. This study indicated that the stages during the
evolution towards C_4 photosynthesis required different driving force and increased water
availability should be needed in the initial phase.

32
33   **Key words:**
34   Evolution, C_4 photosynthesis, intermediate plants, precipitation, temperature, driving process
35
Introduction
Photosynthesis undergone evolution, which results in different photosynthetic type, and C_4
photosynthesis is one of these photosynthetic types(Sage 2004; Langdale 2011). Compared to C_3
relative, the plant with C_4 photosynthesis has higher water use efficiency, nitrogen use efficiency
and light use efficiency (Vogan and Sage 2011; Taylor et al. 2010). Although only accompanying 3%
plant species, C_4 photosynthesis contributes 23% of terrestrial gross primary productivity (Still et
al. 2003; Sage et al. 2012). The evolution from C_3 to C_4 photosynthesis went through multiple
stages (C_4 evolution) (Sage et al. 2012; Sage 2004; Schluter and Weber 2016; Brautigam and
Gowik 2016). The first stage of C_4 evolution is preconditioning in which C_3 plants acquired new

cis-elements (Akyildiz et al. 2007).For this stage, gene duplication occurred (Akyildiz et al. 2007;
Monson 2003), vein density increased (Marshall et al. 2007; Mckown and dengler 2007) and the
number of plastids in the bundle sheath increases (Sage et al. 2013). The second stage is
proto-Kranz. In this stage, plants still perform C_3 photosynthesis (Sage et al. 2013) while the
number of plastids continued to increase (Stata et al. 2019; Sage et al. 2012) and chloroplast and
mitochondria localized to the centripetal wall of bundle sheath cells. The third stage is C_2
photosynthesis (Type I) (Schulze et al. 2013; Sage et al. 2012; Edwards and KU 1987), which is
characterize as the transfer of glycine decarboxylase P subunit (GDCP) and photosynthetic
apparatus from mesophyll cell to bundle sheath cell (Schulze et al. 2013; Stata et al. 2019; Sage et
al. 2013). The released CO_2 from photorespiration can be moderately concentrated in the bundle
sheath cell (Photorespiratory CO_2 pump) (Keerberg et al. 2014; Sage et al. 2012). The fourth stage
is inducing and developing C_4 photosynthesis (Type II) (Schluter and Weber 2016; Sage et al.
2018; Nakamura et al. 2013; Mallmann et al. 2014). The establishment of C_2 photosynthesis could
cause the imbalance of ammonia between mesophyll cell and bundle sheath cell, and the induction
of C_4 metabolism is proposed as a mechanism to resolve this trouble (Mallmann et al. 2014). The
fifth stage is functionally developed C_4 photosynthesis (C_4-like) (Stata et al. 2019). For this stage,
though there are still some Rubisco in mesophyll (Taniguchi et al. 2021; Monson et al. 1987; Stata
et al. 2019; Sage et al. 2018), various types of indicators at this stage are all close to plants with
ultimate C_4 photosynthesis (Vogan and Sage 2011; Ku et al. 1991; Adachi et al. 2023). The final
stage is the optimization (C_4 photosynthesis) (Sage et al. 2012), and mesophyll cell contain few
Rubisco (Stata et al. 2019). These stages in the evolutionary transition between C_3 and C_4
photosynthesis are called intermediate stage, and the plants in these stages are intermediate plants
(or intermediate species).

The evolution from C_3 towards C_4 photosynthesis is considered to be driven by enhancing
photorespiration, which is promoted by the low atmosphere CO_2 concentration, drought, salt and
heat (Sage et al. 2012; Brautigam and Gowik 2016; Sage 2004; Edwards and Smith 2010;
Bromham and Bennett 2014). The low atmosphere CO_2 concentration, drought and salt might
decrease the ratio of CO_2 to O_2 concentration, thereby enhancing photorespiration (Sage et al.
2012). Increasing temperature enhances photorespiration by reducing CO_2 relative to O_2 in the
chloroplast stroma (Ogren 1984) and decreases the specificity of Rubisco for CO_2 relative to O_2
(Ogren 1984; Sage et al. 2012). Previous studies have shown that C_4 evolution are correlated with
the decrease in atmosphere CO_2 concentration and the appearance of dry, salty and hot
environment   (Christin et al. 2008; Edwards and Smith 2010; Edwards and Still 2008; Bromham
and Bennett 2014), which indicates they may be driving force for C_4 evolution. However, all these
studies assume that the entire process of C_4 evolution were under a single driver force. Given C_4
evolution includes multiple phases and the evolution is complicate, it is possible that these phases
of C_4 evolution might be under different driving force rather than the same driving force
persistently along the evolutionary road towards C_4 photosynthesis (Heckmann et al. 2013; Sage et
al. 2012).

Plants obtain fitness in each step of C_4 evolution (Heckmann et al. 2013; Sage et al. 2012). The
increase in vein density is an important characteristics during the initial stage of C_4 evolution
(Sage et al. 2012; Sedelnikova et al. 2018; Kumpers et al. 2017; Sage et al. 2014) (Mckown and
dengler 2007; Marshall et al. 2007; Williams et al. 2013), which might help supply water to the
mesophyll and maintain stomatal openning, since evaporation might be enhanced due to heat at

this time (Brautigam and Gowik 2016; Sage et al. 2018) (Sage et al. 2012). Therefore, available
water content in the environment for this stage may be increased to ensure the fitness of plants
(Brautigam and Gowik 2016). However, this contradicts with the notion that the origin of C_4
photosynthesis was promoted by dry environment (Sage et al. 2018; Edwards and Smith 2010;
Sedelnikova et al. 2018). Since C_2 photosynthesis and C_4 photosynthesis could largely reduce
photorespiration (Sage 2004; Schluter and Weber 2016; Sage et al. 2012), the formation of C_2
from C_3 photosynthesis and the transition from C_2 to developing C_4 photosynthesis may be driven
by the conditions which could enhance photorespiration (Schluter and Weber 2016; Sage et al.
2018; Sage et al. 2012),but the driving forces during these transitions also remain undefined.
Flaveria genus is a widely used system to study C_4 evolution (Sage et al. 2012; Sage 2017;
Schluter and Weber 2016; Mallmann et al. 2014; Sage et al. 2013; Vogan and Sage 2011;
Nakamura et al. 2013), and the studies on Flaveria genus has laid the foundation for the pyramid
model of C_4 evolution (Sage et al. 2013; Vogan and Sage 2011; Sage et al. 2012). The
photosynthetic types of Flaveria species span entire evolutionary stage towards C_4
photosynthesis(Sage et al. 2012; Sage et al. 2014; Schluter and Weber 2016; R.K. Monson 1986) .
Flaveria is also the youngest genus among the known C_4 linage and it's species has the close
relationship (Schulze et al. 2013). These make Flaveria genus an ideal model to study C_4 evolution
(Christin et al. 2011; Sage et al. 2012). The ecology for the habitats of intermediate species and C_4
species could reflect the environmental conditions that drive C_4 evolution (Sage et al. 2012; Sage
et al. 2018), therefore, the ecology for Flaveria habitats enable us to finely study the driving force
in the evolutionary road towards C_4 photosynthesis. In this study, we collected the ecological,
biochemical, anatomical and physiological data for different species of Flaveria genus to explore
driving process for the evolutionary trajectory towards C_4 photosynthesis and how these different
evolutionary phases help plants gain fitness.
113
Results
1. Most of Flaveria species are distributed inland and a few species are distributed along the coast
The habitats of Flaveria species were mapped to the North and South America continents with
the longitude and latitude of each Flaveria samples downloaded from Global Biodiversity
Information Facility (GBIF) (Fig 1A, B). Species were ordered based on the amount of C_4 flux
(Fig 1B, C, supplementary fig 1). We found most Flaveria species are located inland (Fig 1A, B).
A few species (*F.bro*, *F.flo*, *F.lin*) are distributed along the coast (Fig 1B), which was consistent
with the previous reports (Powell 1978). During evolution, intermediate species gradually spread
to locations with higher latitude (Fig 1B). Compared with the intermediate species, the distribution
of C_4 species is more widespread (Fig 1B, C), indicating that adaptive capacity of C_4 plants was
enhanced (Fig1).
125
2. The temperature and precipitation of Flaveria growth habitats
The temperature and precipitation of the habitats for each Flaveria species were obtained from
the worldclim database (Fick and Hijmans 2017). The results from physiological, anatomic,
molecular and computational modeling experiments indicate the direction of C_4 evolution could be
described by species's order towards C_4 photosynthesis in phylogenetic tree (Figure 2A) (Mckown
and dengler 2007; Nakamura et al. 2013; Gowik et al. 2011; Sage et al. 2012; Sage et al. 2018;
Schluter and Weber 2016). Along the evolution towards C_4 photosynthesis (Ordinal species

towards C_4 photosynthesis of the phylogenetic tree in Figure 2A), the annual mean temperature,
mean temperature of warmest quarter, mean temperature of wettest quarter and max temperature
of warmest month first increased and then decreased, though difference in some species habitats is
not significant (Figure 2A, supplementary fig 2, supplementary fig 3, supplementary fig 4,
supplementary file). Mean temperature of warmest quarter, mean temperature of wettest quarter
and max temperature of warmest month increased significantly in the habitat of *F. rob* (C_3) and *F.*
son (Type I) (Fig 2A, B, supplementary fig 2, supplementary fig 3, supplementary file), compared
with those of *F. pri* (C_3). However, these evironmental conditions significantly decreased in the *F.*
ang(Type I), compared with *F. son* (Type I) (Fig 2A, B, supplementary fig2, supplementary fig3,
supplementary file). For the mean temperature of wettest quarter, mean temperature of warmest
quarter and max temperature of warmest month, compared with *F. son*(Type I), the habitat of
species with C_2 photosynthesis (*F. ang*, *F. chl*, *F. pub*, *F. opp*, *F. ano*, *F. ram*) have a lower mean
temperature, but the results of statistic test for some species is not significant or in margin (Fig 2A,
supplementary fig 3D, supplementary file). The habitats of species *F. flo*(Type II) and *F. lin*(Type
II) had significantly higher mean temperature compared to most of other intermediate species (Fig
2A, supplementary file). Max temperature of warmest month for *F. son*(Type II) habitats is
extremely high (Fig 2d, supplementary file).
For the precipitation of wettest quarter, precipitation of warmest quarter and precipitation of
wettest month, along C_4 evolution, they significantly increased in the *F. rob* (C_3) habitat, compared
with those in *F. pri* habitat (Fig 2A, supplementary fig 3, supplementary file). Compared with *F.*
rob (C_3)habitat, precipitation significantly decreased in the *F. ang* (Type I) habitat (Fig 2A,
supplementary fig 3, supplementary file). Precipitation decreased in *F. son* (Type I) habitat, and
though the difference was not statistically significant (Fig 2A, supplementary fig 3, supplementary
file). Precipitation of warmest quarter was higher than precipitation of coldest quarter, and similar
to precipitation of wettest quarter (Fig 2A, supplementary fig 2, supplementary figure 3,
supplementary file). Expect *F. flo* (Type II) and *F. lin* (Type II), precipitation of the habitat of
species with C_2 photosynthesis decreased persistently along C_4 evolution, though difference for
the habitats of some species is not significant (Fig 2A, supplementary fig 3, supplementary file).
In the latter of C_4 evolution, the initial increase in precipitation and temperature (*F. pri* to *F. rob*)
coincided with the initial appearance of the higher vein density (Fig 2A, B, C, supplementary fig 3,
supplementary 4). Although the precipitation of the Type II species *F. flo* and *F. lin* habitat was
high (Fig 2, supplementary fig 3, supplementary file), they inhabited the region including saline
land (Table 1), probably experiencing strong osmotic stress.
166
3. The association between precipitation (or temperature) and C_4 metabolism related traits during
the transition from proto-Kranz to C_4-like
For the clade A, the activities of C_4 metabolism related enzymes PEPC, PPDK and MDH
showed a negative correlation with precipitation, and there showed a linear relationship between
log enzyme activity and log precipitation (Fig 3 A, B, C, supplementary file). The negative
relationship between C_4 flux and precipitation can be described as a linear function (Fig 3D). For
the clade B, the relationship between log enzyme activity and log precipitation also can be
described as a linear function (Fig 3 I, J, K, supplementary file). However, the linear relation
between C_4 flux and precipitation in clade B is weaker (Fig 3L, supplementary file). The linear
relation between log enzyme activity and log temperature is weaker than that between Log

enzyme activity and Log precipitation, regardless of clade A or clade B (Fig 3E, F, M, N, G, H, O,
P, supplementary file).

4. The change in temperature and precipitation between the transition of stages.
The comparison of *F. rob* and *F. son* suggests that incipiently decreased precipitation along the
evolution towards C_4 photosynthesis concurrently occurred with increased mean temperature (Fig
4A, F). Precipitation levels of *F.ano* and *F.ram* were significantly lower than that of *F. ang*, and
temperature values of *F. ram* and *F. ano* were significantly higher than that of *F. ang* (Fig 4B, G, C,
H). Precipitation of *F. pal* was significantly lower than that of *F. ram*, and precipitation of *F. bro* is
significantly lower than that of *F. flo* (Fig 4D,I,E,J). However, temperature for *F.pal* and *F. bro*
showed the inconsistent change, temperature for *F. pal* was same as that of *F. ram*, and
temperature of *F. bro* was significantly lower than that of *F. flo* (Fig 4D,I,E,J). Furthermore,
compared with their more ancient species, we observed that the decreased precipitation for the
species habitat (*F. son*, *F.ram*, *F.flo*, *F.pal* and *F.bro*) concurrently appeared with the decreased
stomatal density, decreased inhibition of O_2 and CO_2 compensation point, decreased maximum
stomatal conductance (Supplementary Fig 5E, F, G, H). Although the precipitation of *F. flo* and *F.*
lin were increased compared to other species with C_2 photosynthesis (Fig 2A, supplementary fig3,
supplementary fig4), they also had the decreased stomatal density, maximum stomatal
conductance(Supplementary fig 5), which might be related to their salty habitats (table1).

5. The association between precipitation, stomata and CO_2 compensation point.
Previous work found there is a difference in stomata density between species at different
evolutionary stages in the Flaveria genus(Zhao et al. 2022). Furthermore, there was a positive and
linear relationship between stomata density, CO_2 compensation point and precipitation during the
transition between proto-Kranz and C_4-like species in Flaveria clade A and clade B (Fig 5A, B, C,
D, E, and F). The linear relationship between stomata, CO_2 compensation point and precipitation
in Flaveria clade B were weaker than that in clade A (Fig 5A, B, C, F, G, H, Supplementary file).
The relation between temperature and stomata density or CO_2 compensation point was not
apparent as the relationships between these two parameters and precipitation (Fig 5D, I, E, J).

Discussion
Evolutionary conditions along the path towards C_4 photosynthesis
Heat and drought have long been thought to be associated with C_4 evolution (Sage et al. 2018;
Taylor et al. 2012). Exploring environmental conditions of the habitats for C_4 and intermediate
species could contribute to elucidate the drivers for C_4 evolution (Sage et al. 2018; Sage et al.
2012; Edwards and Smith 2010). In this work, we examined precipitation and temperature in
diverse Flaveria species during different periods (Fig 2A, supplementary fig 2, 3, 4). We noticed
that the quantity and trend of wettest quarter mostly coincided with those of warmest quarter (Fig
2A, supplementary fig 3), mean temperature of wettest quarter was overall higher than mean
temperature of driest quarter (Supplementary fig 3D, Supplementary fig 2E) and precipitation of
warmest quarter was overall higher than precipitation of coldest quarter (Supplementary fig 3C
supplementary fig 2B). These results suggest that rain and heat were in the same season for the
habitat of Flaveria, and sufficient water content is necessary to maintain high photosynthetic
activity for plants, which is in line with previous work that indicates *Flaveria* species perform

high photosynthesis during summer with heat in Mexico (Powell 1978). Because there is no
apparent change in Rubisco Sc/o (Kubien et al., 2008) and similar Ci/Ca between C_3 and type I
intermediate species (Adachi et al., 2023), the higher photosynthetic activity would indicate also a
higher photorespiratory flux in C_3 plants and initial intermediate species. Besides, under such
conditions, photorespiration, long regarded as the driving factor of C_4 evolution, might further be
enhanced by heat (Sage et al. 2012; Sage et al. 2018). Furthermore, any evolutionary selective
pressure on photosynthesis should act in the periods with active photosynthesis (Sage et al. 2018),
i.e. the driest and coldest season should not be considered for the study on the evolutionary
trajectory towards C_4 photosynthesis (Supplementary fig 2). Therefore, the periods with rain and
heat for Flaveria genus is valuable to explore the driving process of C_4 evolution (Fig 2A,
supplementary fig 3).
We found that the initial stage of C_4 evolution was associated with high precipitation and
temperature, and later stage was associated with decreased precipitation (Fig 2A, Fig 4,
supplementary fig 3). Heat was previously considered to be a major factor that promoted the
origin of C_4 photosynthesis (Sage et al. 2018). Here, our results suggest that in addition to
increasing temperature, the increase in precipitation was also a vital factor to initiate the evolution
towards C_4 photosynthesis (Fig 2, Supplementary fig 3). Previous study proposed it is possible
that heat occurred with drought simultaneously to enhance photorespiration and then promote C_4
evolution (Sage et al. 2012). Here, data from this study indicates that this process indeed existed,
and it was at the transition from C_3 to C_2 (*F. son*) and the transition from C_2 to developing C_4
photosynthesis (*F. ram*, *F. ano*) (Fig 4).
242
The evolutionary mechanism during the initial phase of C_4 evolution
The initial stage of C_4 evolution was featured as having increased vein density and increased
number of chloroplasts in bundle sheath (Sage et al. 2012). The increased vein density can help
maintain sufficient water content in leaf and avoid the reduction of stomata aperture, since the
evaporation might be increased at this stage (Sage et al. 2012; Sage et al. 2018). In agreement with
this, our data showed that, when at the stage of vein density increasing, there was no change in
stomatal development, and temperature was indeed increased (Fig 2A,B,C,D, supplementary fig 5).
The increased temperature could increase vapor pressure between leaf and air (supplementary fig
5B), hence enhanced the evaporation (Sage et al. 2018); while no change in the stomatal
development indicates that water shortage in leaf did not occur (supplementary fig 5), since
long-term water shortage could alter stomatal development (Franks et al. 2009; Taylor et al. 2012).
Importantly, our data also show that precipitation was increased at this stage, which matched with
the increased vein density and ensured sufficient water supply to the leaf (Fig 2A, B, C). Increased
precipitation and keeping stomatal opening could not only maintain intercellular CO_2
concentration, but also could help alleviate the increase of leaf temperature to increase the
acquirement of carbon. Therefore, the initial stage of C_4 evolution was promoted by high
temperature and precipitation (Fig 2).
The increase in vein density could decrease volume of mesophyll in a leaf (Sedelnikova et al.
2018; Mckown and dengler 2007), which inevitably decreases the photosynthetic capacity of the
leaf. In theory, this should create a strong evolutionary pressure to select for the increased
photosynthetic capacity of BSC to offset the loss of photosynthetic capacity. The relative number
and size of chloroplast in bundle sheath/ area of BSC of *F. rob* was indeed potentially increased

compared to those of *F.pri*. Besides, the light intensity of *F. rob* habitat was higher than that of *F.*
pri (supplementary fig 5A). The size of bundle sheath in *F. rob* decreased compared with *F. pri*
(Sage et al. 2013; Mckown and dengler 2007), the size of mesophyll cell decreased compared with
F. pri (Mckown and dengler 2007), and thickness of leaf become less (Mckown and dengler 2007);
all these enhanced the level of light that bundle sheath experiences, consistent with the notion that
selection pressure for bundle sheath with higher photosynthetic capacity might exist at this stage.
271

Drought as a more important factor than heat for driving C_4 evolution
Some studies have proposed that drought is a primary driving force for C_4 evolution (Sage 2004;
Zhou et al. 2018; Sage et al. 2018; Schluter and Weber 2016; Edwards and Smith 2010), and other
studies suggested that heat is the critical environment factor driving C_4 evolution (Sage et al. 2018;
Sage et al. 2012). In this study, we proposed that drought played a more important role than heat
during C_4 evolution (Fig 3). From proto-Kranz to C_4-like, precipitation decline step by step,
however, there is no clear trend of change for temperature (Fig 2). Regardless of clade A or Clade
B, we found a negative association between core C_4 enzyme activity and precipitation (Fig 3) and
positive association between CO_2 compensation point and precipitation (Fig 5). In particular, for
clade A, there was a strong negative relationship between precipitation and activities of core C_4
enzymes (Fig3). However, the relationship wasn't apparent for temperature (Fig 3). These results
suggest that the gradual enhancement of C_2 photosynthesis and the evolution towards C_4
photosynthesis mainly associate to the decreasing precipitation (Fig 2A,B,C,D,E,F, Fig 3).
Therefore, drought rather than heat, might be the major driver for the transition from C_3 to C_4
photosynthesis
287

Different evolutionary drivers at the transition of the stages in C_4 evolution
The environmental drivers for the evolution towards C_4 photosynthesis has been intensively
studied. For example, decreasing atmosphere CO_2 concentration has been thought as a driving
force of C_4 evolution (Christin et al. 2008; Concentrations and Paleogene). Drought, salt and heat
were also thought to as conditions promoting C_4 evolution (Sage et al. 2018; Sage 2004; Bromham
and Bennett 2014) (Edwards and Still 2008; Edwards and Smith 2010; Pearcy and Ehleringer
2006; Sage et al. 2012). These studies assume that entire C_4 evolutionary process is promoted by a
single ecological condition. However, here, our results suggested that environmental conditions
promoted C_4 evolution in an orderly and stepwise manner(Fig 6), and different selection pressure
drove the transition of the stages during the evolutionary way towards C_4 photosynthesis (Fig 6).
Besides the initialization of C_4 evolution, the evolution towards C_2 photosynthesis is also
considered to be a crucial step during C_4 evolution (Sage et al. 2012; Schluter and Weber 2016).
Previous studies have reported that Flaveria species with C_2 photosynthesis can represent a
transition between C_3 and C_4 photosynthesis during the evolution (Heckmann et al. 2013;
Mallmann et al. 2014; Schluter and Weber 2016; Sage et al. 2012), and *F. ram* and *F. ano* have
enhanced C_4 cycle flux, C_4 cycle related enzyme activity and gene expression levels (Adachi et al.
2023; Mallmann et al. 2014; Vogan and Sage 2011; R.K. Monson 1986). According to the
phylogenetic tree, *F. ang* is closer to *F.ram* as compared with *F.son* (Fig 2A), therefore, among all
Flaveria species, *F.son* most likely represents the emergence of C_2 photosynthesis in Flaveria.
Similarly, *F. ram* and *F. ano* most likely represent the emergence in clade A and clade B,
respectively, and *F. pal* and *F. bro* most likely represent the emergence of a fully functional C_4

cycle for clade A and clade B, respectively (Fig 2A). Therefore, the increased temperature and
decreased precipitation simultaneously further promoted the emergence of C_2 photosynthesis (Fig
4A, F). Increasing temperature and decreasing precipitation could enhance photorespiration and
thus promote the formation of photorespiratory CO_2 pump which improve the fitness of these
intermediate plants. The photorespiratory CO_2 pump could largely decrease the photorespiration
(Sage et al. 2018; Sage et al. 2012; Schluter and Weber 2016; Ku et al. 1991), indicated by the
large decrease in CO_2 compensation point and O_2 inhibition (Sage et al. 2013; Ku et al. 1991).
Previous study proposed photorespiration promote C_4 evolution by decreasing stomatal
conductance under drought (Sage et al. 2012). We provided the evidence that there is a positive
relationship between stomatal density, CO_2 compensation point and precipitation (Fig 5), which
suggested that the concurrent evolution of decreased stomata density and also emergence of $C_{2\ and}$
$_{C4}$ photosynthesis during C_4 evolution (Fig 5). The enhancement of C_2 photosynthesis has long
been considered to cause the imbalance of nitrogen between mesophyll cell and bundle sheath cell,
which induces C_4 cycle (Mallmann et al. 2014). In case of the evolutionary process from C_2
photosynthesis to developing C_4 photosynthesis, precipitation decreases and temperature increases
(Fig 4B,C,G,H), which could further drive this procedure. The formation of fully functional C_4
photosynthesis was driven by inconsistent conditions for Clade A and Clade B, and precipitation
have played a dominated role in this step. There is no significant difference in temperature of the
habitats, which suggested that the temperature was not a deciding factor controlling this transition.
328
In summary, this study analyzed the data of environmental ecology, biochemistry and anatomy
for different species of Flaveria genus to reveal driving process of the evolution from C_3 towards
C_4 photosynthesis, and results indicated plants suffered carbon limitation in the early phase of C_4
evolution and the primary driving force for this phase is carbon limitation (Fig 6). Later, plants
suffered water limitation leading to decease in acquirement of carbon, therefore, the primary
driving force for these phases of C_4 evolution is water limitation (Fig 6). For a long time, most of
studies proposed that environmental condition causing the carbon restriction through enhancing
photorespiration (drought, heat, salty and low atmosphere CO_2 concentration) drives the evolution
towards C_4 photosynthesis(Christin et al. 2008; Edwards and Smith 2010; Schluter and Weber
2016; Sage 2004; Sage et al. 2012; Sage et al. 2018). Our study proposed these studies are not
comprehensive and suggested that the steps of the evolution from C_3 to C_4 photosynthesis were
driven by different environmental factors which sequentially promoted C_4 evolution, in contrast to
the current notion that the entire process of C_4 evolution is driven by same environmental factor
(Fig 6). The initial evolution was promoted by high temperature and precipitation, while later
decreased precipitation further promoted the transition between C_2 and C_4 photosynthesis(Fig 6).
More important, the increase in vein density and the activation of bundle sheath cell represent a
starting point of the evolution from C_3 towards C_4 photosynthesis(Christin et al. 2012; Mckown
and dengler 2007; Sage et al. 2012), therefore, high humidity and temperature together may be the
most important condition facilitated the initial evolution towards C_4. Therefore, the origin of C_4
evolution is in the habitat with high humidity and temperature, rather than drought, heat, salty and
low atmosphere CO_2 concentration (Bromham and Bennett 2014; Christin et al. 2008; Sage et al.
2018).
351
Methods

1. Obtain the ecological data of Flaveria species.

First, the geographical distribution of Flaveria was acquired from Global Biodiversity
Information Facility (GBIF) database using the rgibf software package. Using ArcMAP software,
we acquired from the Worldclim database the elevation data of the respective locations and 19
Bioclimatic factors, including Mean Diurnal Range, annual mean Temperature, Isothermality,
Temperature Seasonality, Max Temperature of Warmest Month, Temperature Annual Range, Mean
Temperature of Wettest Quarter, Mean Temperature of Warmest Quarter, Annual Precipitation,
Precipitation of Wettest Month, Precipitation of Warmest Quarter, Precipitation of Wettest Quarter,
Precipitation of Coldest Quarter, Min Temperature of Coldest Month, Precipitation of Driest
Quarter, Mean Temperature of Coldest Quarter, Precipitation of Driest Month, Precipitation
Seasonality (Coefficient of Variation), Precipitation of Coldest Quarter based on the geographical
distribution information of the Flaveria. Specifically, we downloaded the data from the worldclim
at first. Secondly, we prepared the file for the information of sampling location. Thirdly, we
imported the layers file of climate factor. Fourthly, we imported the information of sampling
location. Fifth, generating the new layers. Lastly, we exported the file that including sample and
the data of geographical factors. All bioclimatic variables were averaged from 1970 to 2000. The
elevation data came from SRTM elevation data. All data from the worldclim database have a
maximum spatial resolution of 30 arc-seconds. The water vaper pressure and solar radiation were
also extracted from the World clim database(Fick and Hijmans 2017) based on the geographical
distribution information of Flaveria.

2. Obtain the anatomical and physiological data of Flaveria species.

The anatomical and physiological data for Flaveria was acquired from previous research. The
data of vein data was from (Mckown and dengler 2007). The data of CO_2 compensation point and
O_2 inhibition is from (Ku et al. 1991). The data related to stomata is from (Zhao et al. 2022). The
C_4 metabolic enzyme activity data is from (Adachi et al. 2023) (supplementary table S7). The data
in the graph was digitized with Origin 2018. The growth habitat was acquired from (Powell 1978;
Kocacinar et al. 2008).

3. Map the Flaveria species

We mapped and marked the locations for the habitats of different Flaveria species to a world
map using the software google earth Pro (version 7.3) based on the Latitude and longitude   from
Global Biodiversity Information Facility (GBIF). Coordinates were plotted as placemarks, with
customized icons to distinguish between different species to facilitate the visualization and
analysis of distribution patterns and geographical clusters. Google Earth Pro, version 7.3. (2020).
[Software]. Google LLC. Available at: https://www.google.com/earth/.

4. Statistical analysis, regression analysis and Graph drawing.

The graph was plotted and the regression analysis was performed by using GraphPad prism 8.
The statistic test was performed by GraphPad prism 8 and WPS office. The statistic test for the
difference in precipitation or temperature between all Flaveria species was performed by one-way
ANOVA and Tukey's multiple test (supplementary file). The rest of other statistic test was
performed through Student's t test. We used the $\log_2$ value of the precipitation of wettest quarter
and $\log_2$ value of precipitation of C_4 metabolic enzyme activities, and linear regression between

them were performed with GraphPad prism 8.

Table 1 The habitat of Flaveria species

Species (photosynthetic type)	Growth habitat	Geographic location
Flaveria pringlei (C3)	Rather arid and gypseous habitats, scattered localities, waste places, cultivated fields, roadsides	Southern Mexico
Flaveria cronquistii (C3)	Rocky limestone (possibly gypseous) areas	Southern Mexico
Flaveria robusta (C3)	Gypsum and slate, sandy clay, or open rocky slopes, weedy at roadsides	Western Mexico
Flaveria sonorensis (Type I)	Close to warm mineral springs	North-western Mexico
Flaveria angustifolia (Type I)	Sandy or loamy soils in open fields	Southern Mexico
Flaveria pubescens (Type I)	Scattered localities, gypsum soils	Central Mexico
Flaveria oppositifolia (Type I)	high gypsum plains or other areas with dilute gypsum, saline, or rarely loamy soil, usually in moist areas or in disturbed habitats at roadside	Central Mexico
Flaveria linearis (Type II)	Near the coast and on offshore islands	Western Florida, USA
Flaveria floridana (Type II)	Restricted to saline, white sand near the beaches of brackish marshes	Western Florida, USA
Flaveria ramosissima (Type II)	Sandy-clay bottoms, along dry washes, fields and roadsides, possibly in gypseous soils,	Southern Mexico
Flaveria anomala (Type II)	High gypsum plains, often in disturbed sites	Central Mexico
Flaveria palmeri (C$_4$-like)	Gypsum flats and knolls, or clay-loam soils probably with dilute gypsum, bolsons near lagunas, distributed areas near roads and fields	Northern Mexico
Flaveria brownie (C$_4$-like)	Saline, sandy, and marshy areas of the coastal flats and islands of the lower Texas Gulf Coast	North-eastern Mexico
Flaveria vaginata (C$_4$-like)	Arid and gypseous soils, scrub vegetation, cultivated fields	South-central Mexico
Flaveria bidentis (C$_4$)	A weed in moist places on waste ground, near rivers, streams, and barrancas, in fields, pastures, disturbed ground, near streets and sidewalks in towns, claygravel, or sand	Africa, Caribbean, South America, southern USA

Flaveria trinervia (C₄)	a variety of habitats, usually saline, gypseous, and disturbed, usually near ephemeral or permanent water sources	Africa, Caribbean, Central/South America, Mexico, USA
Flaveria chloraefolia (Type I)	Permanent or ephemeral saline marshy soils, springs, streams, roadsides	North Mexico

The information of habitat of Flaveria species was from (Powell 1978; Kocacinar et al. 2008)

Figure legends

Fig 1 Distribution of the Flaveria species

(A) The location of Flaveria species was mapped to the world map. The main habitat of Flaveria is

Mexico which was displayed. (B) The location of Flaveria species except C_4 species was mapped

to the world map. (C) and (D) The latitude and longitude of different Flaveria species. The latitude

and longitude were acquired from GBIF. Different colors indicate different photosynthetic types.

Fig 2 The environment conditions of the habitats for Flaveria species

(A) The average temperatures of the warmest quarter for different Flaveria species (above panel).

The precipitation of the wettest quarter for different Flaveria species (below panel). (B) The

average temperatures of the wettest quarter for *F. rob* and *F. pri.* (C) The precipitation of wettest

quarter for *F.rob* and *F.pri*. (D) The changes in vein density in the C_3 and intermediate species.

Different colors indicate different photosynthetic types. Orange color represents C_3, grey color

represents intermediate species, green color represents C_4-like, blue color represents C_4 species.

Black lines represent median values.

Fig 3 The association between precipitation and temperature with C_4 enzyme activities or C_4 flux

Here the precipitation represents the precipitation of the wettest quarter. The temperature

represents the average temperature of the wettest quarter. A-H, clade A. (A) The relationship

between Log PEPC activity per chl and Log precipitation. (B) The relationship between Log

PPDK activity per chl and Log precipitation. (C) The relationship between Log MDH activity per

chl and Log precipitation. (D) The relationship between C_4 flux and precipitation. (E) The

relationship between Log PEPC activity per chl and Log temperature. (F) The relationship

between Log PPDK activity per chl and Log temperature. (G) The relationship between Log MDH

activity per chl and Log temperature. (H) The relationship between C_4 flux and Log temperature.

I-P, clade B. (I) The relationship between Log PEPC activity per chl and log precipitation. (J) The

relationship between Log PPDK activity per chl and log precipitation. (K). The relationship

between Log MDH activity per chl and log precipitation. (L). The relationship between C_4 flux

and precipitation. (M) The relationship between log PEPC activity per chl and log temperature. (N)

The relationship between log PPDK activity per chl and log temperature. (O) The relationship

between log MDH activity per chl and log temperature. (P) The relationship between C_4 flux and

temperature.

Fig 4 Precipitation and temperature during the transition from proto-Kranz to C_4-like

(A) Precipitation of the wettest quarter during the transition from proto-Kranz (*F. rob*) to C_2 (*F. son*). (B) Precipitation of the wettest quarter during the transition from C_2 (*F. son*) to partial C_4 (*F. ano*) in clade B. (C) Precipitation of the wettest quarter during the transition from proto-Kranz (*F.son*) to partial C_4 in clade A. (D) Precipitation of the wettest quarter during the transition from partial C_4 to fully functional C_4 photosynthesis. (E) Precipitation of the wettest quarter during the transition from partial C_4 to fully functional C_4 photosynthesis. (F) Mean temperature of the wettest quarter during the transition from proto-kranz (*F.rob*) to C_2 (*F.son*). (G) Mean temperature of the wettest quarter during the transition from C_2 (*F. son*) to partial C_4 (*F. ano*) in clade B. (H) Mean temperature of the wettest quarter during the transition from C_2 (*F. son*) to partial C_4 (*F. ram*) in clade A. (I) Mean temperature of the wettest quarter during the transition from partial C_4 to fully functional C_4 photosynthesis (*F. pal*) in clade A. (J) Mean temperature of the wettest quarter during the transition from partial C_4 (*F. ano*) to fully functional C_4 photosynthesis (*F. bro*) in clade B. The C_4 evolution was launched by increased temperature and increased precipitation. The latter of C_4 evolution was promoted by increased temperature and decreased precipitation.

Fig 5 The association between stomatal density, CO_2 compensation point, precipitation and temperature.A-E, clade A.
(A) The relationship between stomatal density and precipitation. (B) The relationship between CO_2 compensation point and precipitation. (C) The relation between stomatal density and CO_2 compensation point. (D) The relationship between stomatal density and temperature. (E) The relationship between CO_2 compensation point and temperature. F-J, clade B. (F) The relationship between stomatal density and precipitation. (G) The relationship between CO_2 compensation point and precipitation. (H) The relation between stomatal density and CO_2 compensation point. (I) The relation between stomatal density and temperature. (J) The relation between CO_2 compensation point and temperature.

Fig 6. Schematic diagram for the driver process of C_4 evolution.
The schematic diagram for the conditions driving the evolutionary process for C_4 photosynthesis. The raindrop number denotes the precipitation. The thermometer denotes the temperature. The arrows denote the direction of C_4 evolution. The cycle composed by white and black denotes the main driving element that promote the corresponding stage of C_4 evolution. The driving reason in C_3 and proto-Kranz was mainly carbon limitation, and the driving reason in the latter stage was mainly water limitation. More and more black indicating the carbon limitation was increasing stepwise, and more and more white indicating the water limitation was increasing stepwise.

Supplementary fig 1. The C_4 flux for Flaveria.
The data of C_4 flux was from table 1 in (Vogan and Sage 2011)

Supplementary fig 2. Temperature and precipitation of habitat for Flaveria species in driest and coldest period.
(A) Precipitation of the driest quarter. (B) Precipitation of the coldest quarter. (C) Precipitation of the driest month. (D) Mean temperature of the driest quarter. (E) Mean temperature of coldest quarter. (F) Min temperature of coldest month. Orange, C_3; Purple, Type I; Yellow, Type II; Green, C_4-like; Blue, C_4. Black lines indicate median.

Supplementary fig 3. Temperature and precipitation of habitat for Flaveria species in wettest and
warmest period.
(A) Max temperature of warmest month. (B) Precipitation of westest month. (C) Precipitation of
warmest quarter. (D) Mean temperature of wettest quarter. Orange indicates C_3, purple indicates
Type I, yellow indicates Type II, green indicates C_4-like, blue indicates C_4. Black lines indicate
median.
490

491 Supplementary fig 4. Annual temperature and precipitation of habitats for Flaveria species
492 (A) Annual mean temperature. (B) Annual precipitation. Orange, C_3; Purple, Type I; Yellow, Type
493 II; Green, C_4-like; Blue, C_4. Black lines indicate median.
494

495 Supplementary fig 5. The change trend for the evolution towards C_4 photosynthesis.
496 (A) Precipitation of wettest quarter. (B) Mean temperature of wettest quarter. (C) Vein density. (D)
497 Distance between vein. (E) O_2 inhibition. (F) CO_2 compensation point. (G) Gs_{max} (H) Stomatal
498 density (SD).
499

Supplementary fig 6. The difference in solar radiation and water vaper pressure between *F. pri* and
F. rob.
A, The difference in solar radiation between *F.pri* and F.rob. B, The difference in water vaper
pressure between *F.pri* and *F.rob*.
Table 1 The habitat for Flaveria species
Supplementary file 1: The statistics test and regression analysis
506

509   Acknowledgements
510   Thanks for the research group, and the postdoc support in shanghai.
511

Author contribution
YYZ and ZGZ designed the study. YXZ and YYZ extracted the climate data and the latitude
and longitude data, and YYZ analyzed these climate data. XQY and YYZ mapped and marked the
coordinates of Flaveria genus location and do the visualization. YYZ extracted the physiological
and anatomic data from the previous works. YYZ analyzed all these data, and did the statistic test
and regression analysis. YYZ plotted the graphs. YYZ and XGZ wrote the paper.
518

519   Competing interests
520   The authors declare no competing financial interests.
521

522   References
523   Adachi S, Stata M, Martin DG, Cheng S, Liu H, Zhu X-G, Sage RF (2023) The Evolution of C4
524   Photosynthesis in Flaveria (Asteraceae): Insights from the Flaveria linearis Complex. Plant
525   physiology 191 (1):233-251. doi:10.1093/plphys/kiac467
526   Akyildiz M, Gowik U, Engelmann S, Koczor M, Streubel M, Westhoff P (2007) Evolution and

function of a cis-regulatory module for mesophyll-specific gene expression in the C_4 dicot Flaveria trinervia. Plant Cell 19 (11):3391-3402. doi:10.1105/tpc.107.053322

Brautigam A, Gowik U (2016) Photorespiration connects C3 and C4 photosynthesis. J Exp Bot 67 (10):2953-2962. doi:10.1093/jxb/erw056

Bromham L, Bennett TH (2014) Salt tolerance evolves more frequently in C4 grass lineages. Journal of Evolutionary Biology 27 (3):653-659. doi:10.1111/jeb.12320

Christin P-A, Osborne CP, Chatelet DS, Columbus JT, Besnard G, Hodkinson TR, Garrison LM, Vorontsova MS, Edwards EJ (2012) Anatomical enablers and the evolution of C4 photosynthesis in grasses. Proceedings of the National Academy of Sciences 110 (4):1381-1386. doi:10.1073/pnas.1216777110

Christin PA, Besnard G, Samaritani E, Duvall MR, Hodkinson TR, Savolainen V, Salamin N (2008) Oligocene CO2 decline promoted C4 photosynthesis in grasses. Curr Biol 18 (1):37-43. doi:10.1016/j.cub.2007.11.058

Christin PA, Osborne CP, Sage RF, Arakaki M, Edwards EJ (2011) C(4) eudicots are not younger than C(4) monocots. J Exp Bot 62 (9):3171-3181. doi:10.1093/jxb/err041

Concentrations MDiACD, Paleogene Dt <science.1110063.pdf>.

Edwards EJ, Smith SA (2010) Phylogenetic analyses reveal the shady history of C4 grasses. Proc Natl Acad Sci U S A 107 (6):2532-2537. doi:10.1073/pnas.0909672107

Edwards EJ, Still CJ (2008) Climate, phylogeny and the ecological distribution of C4 grasses. Ecol Lett 11 (3):266-276. doi:10.1111/j.1461-0248.2007.01144.x

Edwards Ge, KU Msb (1987) Biochemistry of C3-C4 Intermediates. The Biochemistry of Plants, 10

Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology 37 (12):4302-4315. doi:10.1002/joc.5086

Franks PJ, Drake PL, Beerling DJ (2009) Plasticity in maximum stomatal conductance constrained by negative correlation between stomatal size and density: an analysis using Eucalyptus globulus. Plant, cell & environment 32 (12):1737-1748. doi:10.1111/j.1365-3040.2009.002031.x

Gowik U, Bräutigam A, Weber KL, Weber APM, Westhoff P (2011) Evolution of C_4 Photosynthesis in the Genus Flaveria: How Many and Which Genes Does It Take to Make C_4? The Plant Cell 23 (6):2087-2105. doi:10.1105/tpc.111.086264

Heckmann D, Schulze S, Denton A, Gowik U, Westhoff P, Weber AP, Lercher MJ (2013) Predicting C4 photosynthesis evolution: modular, individually adaptive steps on a Mount Fuji fitness landscape. Cell 153 (7):1579-1588. doi:10.1016/j.cell.2013.04.058

Keerberg O, Parnik T, Ivanova H, Bassuner B, Bauwe H (2014) C_2 photosynthesis generates about 3-fold elevated leaf CO_2 levels in the C_3-C_4 intermediate species Flaveria pubescens. J Exp Bot 65 (13):3649-3656. doi:10.1093/jxb/eru239

Kocacinar F, McKown AD, Sage TL, Sage RF (2008) Photosynthetic pathway influences xylem structure and function in Flaveria (Asteraceae). Plant, cell & environment 31 (10):1363-1376. doi:10.1111/j.1365-3040.2008.01847.x

Ku MS, Wu J, Dai Z, Scott RA, Chu C, Edwards GE (1991) Photosynthetic and photorespiratory characteristics of flaveria species. Plant physiology 96 (2):518-528.

doi:10.1104/pp.96.2.518

Kumpers BM, Burgess SJ, Reyna-Llorens I, Smith-Unna R, Boursnell C, Hibberd JM (2017)
Shared characteristics underpinning C4 leaf maturation derived from analysis of multiple C3
and C4 species of Flaveria. J Exp Bot 68 (2):177-189. doi:10.1093/jxb/erw488

Langdale JA (2011) C4 cycles: past, present, and future research on C4 photosynthesis.
Plant Cell 23 (11):3879-3892. doi:10.1105/tpc.111.092098

Mallmann J, Heckmann D, Brautigam A, Lercher MJ, Weber AP, Westhoff P, Gowik U (2014)
The role of photorespiration during the evolution of C_4 photosynthesis in the genus Flaveria.
Elife 3:e02478. doi:10.7554/eLife.02478

Marshall DM, Muhaidat R, Brown NJ, Liu Z, Stanley S, Griffiths H, Sage RF, Hibberd JM
(2007) Cleome, a genus closely related to Arabidopsis, contains species spanning a
developmental progression from C(3) to C(4) photosynthesis. Plant J 51 (5):886-896.
doi:10.1111/j.1365-313X.2007.03188.x

Mckown A, dengler N (2007) Key innovations in the evolution of Kranz anatomy and C4
vein pattern in Flaveria (Asteraceae). American Journal of Botany 94:382–399

Monson, Willlam S. Schuster, Ku MSB (1987) Photosynthesis in Flaveria brownil A.M.
Powell. Plant physiology 85:1063-1067

Monson RK (2003) Gene duplication, neofunctionalization, and the evolution ofC4
photosynthesis. Int J Plant Sci

Nakamura N, Iwano M, Havaux M, Yokota A, Munekage YN (2013) Promotion of cyclic
electron transport around photosystem I during the evolution of NADP-malic enzyme-type C4
photosynthesis in the genus Flaveria. New Phytol 199 (3):832-842. doi:10.1111/nph.12296

Ogren DBJaWL (1984) The CO2_O2 specificity of ribulose 1,5-bisphosphate carboxylase
oxygenase_ dependence on ribulosebisphosphate concentration, pH and temperature. planta
161:308-313

Pearcy RW, Ehleringer J (2006) Comparative ecophysiology of C3 and C4 plants. Plant, cell
& environment 7 (1):1-13. doi:10.1111/j.1365-3040.1984.tb01194.x

Powell AM (1978) Systematics of Flaveria (Flaveriinae--Asteraceae). Annals of the Missouri
Botanical Garden 65:590-636

R.K. Monson BdM, M.S.B. Ku and G.E. Edwards (1986) Co-function of C3- and
C4-photosynthetic pathways in C3, C4 and C3-C4 intermediate Flaveria species. Planta

Sage RF (2004) The evolution of C_4 photosynthesis. New Phytologist 161:341-270.
doi:10.1046/j.1469-8137.2004.00974.x

Sage RF (2017) A portrait of the C4 photosynthetic family on the 50th anniversary of its
discovery: species number, evolutionary lineages, and Hall of Fame. J Exp Bot 68
(2):4039-4056. doi:10.1093/jxb/erx005

Sage RF, Khoshravesh R, Sage TL (2014) From proto-Kranz to C4 Kranz: building the bridge
to C4 photosynthesis. J Exp Bot 65 (13):3341-3356. doi:10.1093/jxb/eru180

Sage RF, Monson RK, Ehleringer JR, Adachi S, Pearcy RW (2018) Some like it hot: the
physiological ecology of C4 plant evolution. Oecologia 187 (4):941-966.
doi:10.1007/s00442-018-4191-6

Sage RF, Sage TL, Kocacinar F (2012) Photorespiration and the evolution of C_4
photosynthesis. Annu Rev Plant Biol 63:19-47. doi:10.1146/annurev-arplant-042811-105511

Sage TL, Busch FA, Johnson DC, Friesen PC, Stinson CR, Stata M, Sultmanis S, Rahman BA,

Rawsthorne S, Sage RF (2013) Initial events during the evolution of C4 photosynthesis in C3
species of Flaveria. Plant physiology 163 (3):1266-1276. doi:10.1104/pp.113.221119

Schluter U, Weber AP (2016) The Road to C4 Photosynthesis: Evolution of a Complex Trait
via Intermediary States. Plant Cell Physiol 57 (5):881-889. doi:10.1093/pcp/pcw009

Schulze S, Mallmann J, Burscheidt J, Koczor M, Streubel M, Bauwe H, Gowik U, Westhoff P
(2013) Evolution of C4 photosynthesis in the genus flaveria: establishment of a
photorespiratory CO2 pump. Plant Cell 25 (7):2522-2535. doi:10.1105/tpc.113.114520

Sedelnikova OV, Hughes TE, Langdale JA (2018) Understanding the Genetic Basis of C_4
Kranz Anatomy with a View to Engineering C_3 Crops. Annu Rev Genet 52:249-270.
doi:10.1146/annurev-genet-120417-031217

Stata M, Sage TL, Sage RF (2019) Mind the gap: the evolutionary engagement of the C4
metabolic cycle in support of net carbon assimilation. Curr Opin Plant Biol 49:27-34.
doi:10.1016/j.pbi.2019.04.008

Still CJ, Berry JA, Collatz GJ, DeFries RS (2003) Global distribution of C3and C4vegetation:
Carbon cycle implications. Global Biogeochemical Cycles 17 (1):6-1-6-14.
doi:10.1029/2001gb001807

Taniguchi YY, Gowik U, Kinoshita Y, Kishizaki R, Ono N, Yokota A, Westhoff P, Munekage YN
(2021) Dynamic changes of genome sizes and gradual gain of cell-specific distribution of C4
enzymes during C4 evolution in genus Flaveria. Plant Genome:e20095.
doi:10.1002/tpg2.20095

Taylor SH, Franks PJ, Hulme SP, Spriggs E, Christin PA, Edwards EJ, Woodward FI, Osborne
CP (2012) Photosynthetic pathway and ecological adaptation explain stomatal trait diversity
amongst grasses. New Phytol 193 (2):387-396. doi:10.1111/j.1469-8137.2011.03935.x

Taylor SH, Hulme SP, Rees M, Ripley BS, Woodward FI, Osborne CP (2010) Ecophysiological
traits in C_3 and C_4 grasses: a phylogenetically controlled screening experiment. New Phytol
185 (3):780-791. doi:10.1111/j.1469-8137.2009.03102.x

Vogan PJ, Sage RF (2011) Water-use efficiency and nitrogen-use efficiency of C_3-C_4
intermediate species of Flaveria Juss. (Asteraceae). Plant, cell & environment 34
(9):1415-1430. doi:10.1111/j.1365-3040.2011.02340.x

Williams BP, Johnston IG, Covshoff S, Hibberd JM (2013) Phenotypic landscape inference
reveals multiple evolutionary paths to C_4 photosynthesis. elife 2:e00961.
doi:10.7554/eLife.00961

Zhao YY, Lyu MA, Miao F, Chen G, Zhu XG (2022) The evolution of stomatal traits along the
trajectory toward C4 photosynthesis. Plant physiology 190 (1):441-458.
doi:10.1093/plphys/kiac252

Zhou H, Helliker BR, Huber M, Dicks A, Akcay E (2018) C_4 photosynthesis and climate
through the lens of optimality. Proc Natl Acad Sci U S A 115 (47):12057-12062.
doi:10.1073/pnas.1718988115

Clade A

CladeB

Clade A

Clade B