Genetically encoded reporters of actin filament organization in living cells and tissues

Carla Silva Martins[1,11], François Iv[1,11], Shashi Kumar Suman[2], Thomas C. Panagiotou[3], Clara Sidor[4], María Ruso-López[5], Camille N. Plancke[2], Shizue Omi[1], Maxime Gomes[1], Alexander Llewellyn[1], Sourish Reddy Bandi[1], Laurie Ramond[1], Federica Arbizzani[6], Caio Vaz Rimoli[1], Frank Schnorrer[4], François Robin[2], Andrew Wilde[3,7], Loïc LeGoff[1], Jean-Denis Pedelacq[8], Stéphanie Cabantous[9], Sergio A. Rincon[5], Cristel Chandre[10], Sophie Brasselet[1,*], Manos Mavrakis[1,12,*]

[1]Institut Fresnel, CNRS UMR7249, Aix Marseille Univ, Centrale Med, 13013 Marseille, France
[2]Sorbonne Université, CNRS, Institut de Biologie Paris Seine, Laboratoire de Biologie du Développement/UMR7622, 75005 Paris, France
[3]Department of Molecular Genetics, University of Toronto, Toronto, ON M5S 1M1, Canada
[4]Aix Marseille Univ, CNRS, IBDM, Turing Centre for Living Systems, 13009 Marseille, France
[5]Instituto de Biología Funcional y Genómica and Departamento de Microbiología y Genética, Consejo Superior de Investigaciones Científicas (CSIC) / Universidad de Salamanca, Salamanca 37007, Spain
[6]Institut Curie, PSL University, CNRS UMR 144, 75005 Paris, France
[7]Department of Biochemistry, University of Toronto, Toronto, ON M5S 1M1, Canada
[8]Institut de Pharmacologie et de Biologie Structurale, IPBS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France
[9]Centre de Recherche en Cancérologie de Toulouse (CRCT), Inserm, Université Paul Sabatier-Toulouse III, CNRS, 31037 Toulouse, France
[10]CNRS, Aix Marseille Univ, I2M, Marseille 13009, France
[11]These authors contributed equally
[12]Lead contact

*Correspondence: sophie.brasselet@fresnel.fr (S.B.), manos.mavrakis@fresnel.fr (M.M.)

SUMMARY

The cytoskeletal protein actin is crucial for cell shape and integrity throughout eukaryotes. Actin filaments perform essential biological functions, including muscle contraction, cell division and tissue morphogenesis. These diverse activities are achieved through the ability of actin filaments to be arranged into diverse architectures, but a detailed appreciation of the dynamic organizational state of the actin filaments has been hindered by available tools. Fluorescence polarization microscopy is uniquely placed for measuring actin organization by exploiting the sensitivity of polarized light excitation to the orientation of fluorophores attached to actin filaments. By engineering constrained fluorescent protein fusions to widely used actin localization reporters, we have succeeded in developing novel genetically-encoded reporters for non-invasive, quantitative measurements of actin filament organization in living cells by fluorescence polarization microscopy. We show examples of actin organization measurements in living mammalian cells in culture, as well as *in vivo* in fission yeast, *C. elegans* and *Drosophila*.

KEYWORDS

actin filaments, molecular organization, molecular order, molecular orientation, filament alignment, fluorescence polarization microscopy, morphogenesis, GFP mobility

INTRODUCTION

Actin is one of the most abundant and conserved proteins throughout eukaryotes, including structural homologs in bacteria. Actin monomers polymerize through non-covalent association to generate filaments. These filaments perform a wide range of essential biological functions, such as muscle contraction, cell division, cell adhesion, cell motility, tissue morphogenesis and intracellular pathogen movement (Pollard and Cooper, 2009). These various activities are achieved through dozens of actin binding proteins which configure actin into a diverse set of organization states (Pollard, 2016). The precise geometrical organization of actin filament assemblies, i.e. how actin filaments are physically oriented in space and to what extent they are aligned to each other (collectively referred to as filament *organization*, hereafter), is crucial to biological function (Blanchoin et al., 2014; Fletcher and Mullins, 2010; Heisenberg and Bellaiche, 2013; Lappalainen et al., 2022; Paluch and Heisenberg, 2009). Typically, multiple actin-binding proteins populate actin filament (F-actin) assemblies in cells, and predicting actin filament organization and its effect on function remains elusive.

Much progress has been made in defining the proteome of the actin cytoskeleton and their dynamics, but a detailed appreciation of the dynamic organizational state of the actin filaments themselves has been hindered by available tools. Cryo-electron tomography emerges as a very powerful method to visualize the nanoscale organization of actin filaments in cells (Chakraborty et al., 2020), but it is not applicable to living samples. The most widely used approach for imaging actin filaments in living cells uses fluorescence microscopy and either one of the following three actin filament-binding probes: (1) small (~1 kDa) organic fluorescent dye conjugates of the F-actin-binding drugs phalloidin (Barak et al., 1980; Wulf et al., 1979) and jasplakinolide (Lukinavicius et al., 2014); such examples are AlexaFluor488 (AF488)-phalloidin and silicon rhodamine (SiR)-jasplakinolide, the latter known with the misleading name "SiR-actin", (2) GFP fusions to G-actin (Ballestrem et al., 1998; Choidas et al., 1998; Doyle and Botstein, 1996; Verkhusha et al., 1999; Westphal et al., 1997), and (3) GFP fusions to actin-binding peptides or protein domains (for example, the actin-binding domain of moesin (Edwards et al., 1997)). The intensity and distribution of the fluorescent pixels in the images inform us on the relative localization and levels of filamentous actin within the cell. However, how actin filaments are organized at a given image pixel, within the optical resolution of the microscope (typically ~200 nm), cannot be deduced from the fluorescence intensity alone, nor from the pattern of fluorescent pixel distribution. Quantitative imaging tools that enable real-time measurements of filament organization are currently limiting.

Fluorescence polarization microscopy (hereafter, polarimetry) is ideally placed for measuring actin filament organization in living cells by exploiting the sensitivity of polarized light excitation to the orientation of fluorophores attached to actin filaments. Fluorescence is maximized when polarized light is aligned with the excitation dipoles of the fluorophores (Forkey et al., 2000). Thus, an ensemble of fluorophores will be excited more efficiently in preferred directions depending on the organization of the fluorophores. By measuring the modulation of fluorescence induced by the rotation of light polarization in the sample plane, we can extract independently two angles per image pixel: the mean orientation of the fluorophores within the focal volume (Figure 1A, angle rho, ρ) and the in-plane projection of the angle explored by these fluorophores, (Figure 1A, angle psi, ψ) (Brasselet et al., 2013). If the fluorophores are linked to actin filaments in a constrained manner, the measured parameters reflect directly the molecular-scale organization of the labeled filaments in living cells at each pixel location, with ρ their mean orientation, and ψ their degree of alignment; the higher the filament alignment, the lower the ψ (see methods for details).

Fluorophore conjugates to phalloidin and jasplakinolide have been used successfully for polarimetry in fixed cells (Juanes et al., 2019; Loison et al., 2018;

Mavrakis et al., 2014; Spira et al., 2017; Swaminathan et al., 2017). However, their use
in living cells is far from ideal given that these drugs stabilize actin filaments (Bubb et al.,
1994; Cramer, 1999; Dancker et al., 1975; Wieland, 1977). Introducing these drugs into
tissues is also experimentally challenging, while controlling their intracellular
concentration spatiotemporally is practically impossible, rendering their use in living
tissues very limited. The goal of our study is to extend the potential of polarimetry to living
cells and tissues by generating minimally perturbative, genetically-encoded, fluorescent
protein-based reporters for live-cell measurements of actin filament organization. To this
end, we tailor GFP fusions to widely used F-actin localization reporters by constraining
the mobility of the GFP in order to render them usable for organization measurements by
polarimetry. We use stress fibers in cultured mammalian cells as a model system of
known F-actin organization to identify constrained GFP fusions that report faithfully the
orientation and alignment of actin filaments in live cells. We further validate the
functionality and use of the novel reporters in three genetically tractable *in vivo* model
systems, the fission yeast *Schizosaccharomyces pombe*, the nematode *Caenorhabditis*
elegans and the fruit fly *Drosophila melanogaster*.

124
125 **RESULTS**
126
127 **Widely used genetically encoded actin localization reporters are not suitable for**
128 **organization measurements with polarimetry.**
129
For the sake of illustrating how we display and interpret polarimetry measurements in
cells throughout our study, Figure 1B shows an example of polarimetry in fixed
mammalian cells stained with AF488-phalloidin or SiR-jasplakinolide. We focus on the
analysis of pixels containing stress fibers (SFs), which electron microscopy (EM) has
shown to form bundles of actin filaments highly parallel to each other (Heuser and
Kirschner, 1980; Svitkina et al., 1995). The angles ρ and ψ are represented as orientation
and organization maps, respectively. The distribution of ψ angles for multiple SFs in tens
of cells is shown in Figure 1C. The analysis reveals three key findings. First, AF488 and
SiR fluorophore dipoles are, on the average, parallel and perpendicular to actin filaments,
respectively, in line with previous reports (Juanes et al., 2019; Loison et al., 2018;
Mavrakis et al., 2014; Mehta et al., 2016; Rimoli et al., 2022; Swaminathan et al., 2017;
Valades Cruz et al., 2016). Second, actin filaments are oriented along the axis of the SFs:
SFs oriented differently in the cell show distinct ρ stick colors according to their precise
orientation, allowing for an easy visual tracking of changes in mean filament orientation
(Figure 1B). Third, actin filaments within SFs are highly aligned. Single AF488
fluorophores wobble by ~90-100° with a tilt angle of ~20° off the actin filament axis (Rimoli
et al., 2022; Valades Cruz et al., 2016) thus measured ψ of ~120-130° (Figure 1C and
1D,iii) reflect that the contained filaments at the image pixels of SFs are highly parallel to
each other. The visual inspection of the color-coded ψ maps reveals that all SFs share
very similar filament organization despite their different orientations (Figure 1B). Given
the precision of only a few degrees for ρ and ψ angle measurements (Kress et al., 2013),
the very narrow distribution of ρ and ψ angles within a region of interest in a given SF (for
example, standard deviation=3-5° for ROI5 of AF488, Figure 1B) further reveals a very
homogeneous population of orientations and aperture angles explored at an image pixel,
in full line with what one expects from EM. This analysis showcases the potential of
polarimetry for organization measurements in cells: even though one cannot resolve
individual filaments, their molecular-scale organization at a given image pixel is
detectable and can be quantified as shown. This said, the reason why such
measurements are possible in these examples is because AF488 and SiR are sufficiently
constrained in their conjugates with phalloidin and jasplakinolide, respectively.

To assess the usability of available genetically-encoded F-actin localization reporters for organization measurements in living cells using polarimetry, we measured actin filament orientation and alignment on stress fibers (SFs) of live U2OS cells expressing either of five widely used F-actin-binding EGFP fusions: the F-actin binding peptides Lifeact (Riedl et al., 2008) and F-tractin9-52 (Johnson and Schell, 2009), the actin-binding domain of human Utrophin Utr1-261 (Burkel et al., 2007), the synthetic actin-binding Affimer, Affimer6 (Lopata et al., 2018), and human non-muscle beta G-actin (Ballestrem et al., 1998). The measured ψ angles on SFs were very high (>160°) in all cases (Figure 1E). Given that actin filaments in stress fibers are highly aligned to each other, the high ψ values cannot result from disordered actin filaments, but they rather reflect the high rotational mobility of EGFP in the respective fusions (Figure 1D,i), rendering the latter not suitable for organization measurements.

Design of GFP-based reporters with constrained GFP mobility

Tailoring the available F-actin localization reporters for organization measurements requires that GFP mobility is constrained. We reasoned that there are three main sources that contribute to the flexibility of the EGFP. The first source is the presence of an amino acid linker between the actin-binding moiety (ABD) and the fused EGFP. All the widely used EGFP fusions tested above have been generated by standard restriction-ligation cloning using the multiple cloning sites of the EGFP-C1 and EGFP-N1 vectors from Clontech (Zhang et al., 1996) and thus inevitably introducing several amino acid residues in between the EGFP terminus and the ABD. Although the presence of such a linker is reasonably assumed to be important for minimizing interference with protein folding and interactions with actin-binding proteins, we hypothesized that it also contributes to the rotational mobility of the fused EGFP. The second source is the flexibility of the terminus of the ABD to which EGFP is fused. Thus, shortening or removing altogether the linker and/or flexible stretches from the terminus of the ABD could be promising approaches for constraining the GFP, assuming that protein folding, F-actin binding and interactions with actin-binding partners are not compromised.

The third source of flexibility are the N- and C-termini of EGFP themselves. The crystal structures of GFP (Ormo et al., 1996; Yang et al., 1996), EGFP (Royant and Noirclerc-Savoye, 2011) and practically all fluorescent protein (FP) variants with the same termini show that the C-terminus after the end of the β11 strand comprises 11 residues that are unstructured and most of which are absent from the respective crystal structures due to their flexibility (Figure 1F,G). The N-terminal stretch preceding the β1 strand comprises 12 residues, with residues 5-9 forming a 3_{10} helix (in red in Figure 1F,G), while the first 4 amino acid residues are similarly unstructured and often not visible in the crystal structures. Shortening of the N-terminus by 4-6 residues has indeed been shown to constrain the mobility of GFPS65T and EGFP in C-terminal fusions to septins and nucleoporins (DeMay et al., 2011a; Kampmann et al., 2011; Vrabioiu and Mitchison, 2006). However, more extensive truncation of the termini has not been explored to date. This is probably due to early studies with wild-type GFP, GFPS65T or EGFP that showed that removing more than six N-terminal or more than nine C-terminal residues significantly impairs or abolishes GFP fluorescence altogether (Dopf and Horiagon, 1996; Kim and Kaang, 1998; Li et al., 1997).

Prompted by the recent finding that better-folded variants of GFP tolerate more extensive terminal truncations (Raghunathan et al., 2012), we made a side-by-side comparison of truncation mutants of EGFP and the exceptionally stable superfolder GFP (sfGFP) (Pedelacq et al., 2006) to determine to what extent we can shorten their termini without compromising fluorescence. To this end, we expressed truncation mutants in U2OS cells and assessed their fluorescence both by fluorescence-activated cell sorting (FACS) and by spinning disk confocal fluorescence microscopy imaging. Both FACS and

imaging confirmed the sensitivity of EGFP to terminal truncations, and showed, at the same time, that sfGFP missing either the entire N- (ΔN12) or C- terminus (ΔC11), or both N- and C-termini (ΔN12ΔC11), retains its fluorescence and is usable for fluorescence imaging (Figure 1G,H).

We hypothesized that engaging both GFP termini could also constrain GFP mobility. Intramolecular GFP fusions, with GFP placed within a protein structure, for example within a loop, or between a transmembrane and extracellular domain, could reduce its rotational mobility. Such constructs have notably been used to constrain GFP mobility in fusions with the integral membrane proteins integrin (Swaminathan et al., 2017) and the major histocompatibility complex class I (MHC I) protein (Kress et al., 2011). Alternatively, circularly permuted GFP (cpGFP) (Akemann et al., 2001; Baird et al., 1999; Topell et al., 1999) with the original termini connected via an ABD (Figure 1J) could also act to constrain GFP mobility. Given that better-folded variants of GFP also behave much better in intramolecular fusions in terms of functionality (Bendezu et al., 2015; Osawa and Erickson, 2005) and that they can be beneficial for the folding and stability of circular permutants (Carlson et al., 2010), we considered the use of sfGFP as the best choice for GFP fusion engineering. To suppress GFP dimerization-related artifacts, we further introduced the V206K mutation to generated monomeric sfGFP (msfGFP) (Costantini et al., 2012; Cranfill et al., 2016; Zacharias et al., 2002) which we used for all subsequent screening (Figure 1I).

Besides the use of full-length msfGFP for fusions with ABDs, we employed two additional approaches for intramolecular G-actin fusions. Instead of the full-length GFP, we used the 16-residue GFP fragment, β11, that we complemented with co-expressed GFP1-10 in a bipartite split-GFP complementation assay-like manner (Cabantous et al., 2005) (Figure 1K). We finally employed the tetracysteine-biarsenical system, which uses among the smallest genetically encoded tags for fluorescent labeling and which resembles conceptually the bifunctional rhodamine approach (Figure 1K). To this end, we genetically fused a short peptide sequence containing 4 cysteines to G-actin. Upon addition of the membrane-permeable, nonfluorescent biarsenical dye FlAsH (fluorescein arsenical hairpin binder) and its specific binding to the tetracysteine motif, the dye becomes fluorescent *in situ* in live cells (Griffin et al., 1998; Martin et al., 2005). Both the increased functionality due to the small size of the genetic tag (Andresen et al., 2004) and the rigidity of the peptide-fluorophore complex suggested by nuclear magnetic resonance (NMR) (Madani et al., 2009) prompted us to generate such fusions for G-actin. Terminal and intramolecular fusions of G-actin with this approach have been successfully used to visualize F-actin in mammalian and fission yeast cells (Chen et al., 2012; Martin et al., 2005).

Taking together the above-mentioned points, we embarked on a screen of ABD-GFP fusions using the widely used ABDs tested earlier, namely Lifeact, Utrophin Utr1-261, F-tractin9-52, Affimer6, and human non-muscle beta G-actin. To constrain GFP mobility, we generated terminal, intramolecular and circularly permuted GFP fusions without linker sequences, with shortened GFP termini and shortened termini of the ABD. We expressed fusions transiently in U2OS cells and made polarimetry measurements on SFs, using the latter as a model structure of known actin filament organization. Given that SFs comprise highly aligned actin filaments and if GFP mobility is sufficiently constrained, we expect to measure mean fluorophore dipole orientations (ρ angle values) that are either parallel or perpendicular to the SF axis. The smaller the measured ψ angles are on SFs, the more constrained will be the GFP mobility of the respective fusion and thus the larger the range of changes in F-actin organization that we will be able to detect. With the intention to provide a rational for constraining GFP mobility, we chose to show and discuss the results of all constructs and not only the ones with constrained GFP mobility.

Engineering actin filament organization reporters using Lifeact.

The N-terminal 17 amino acids of the budding yeast actin-binding protein Abp140 are
known as the actin-binding peptide "Lifeact" (Riedl et al., 2008) and are used widely for
labeling F-actin in live cells. The original reporter used a C-terminal EGFP (Riedl et al.,
2008) but N-terminal EGFP fusions have also been used successfully (Courtemanche et
al., 2016). To constrain GFP mobility in fusions with Lifeact, we generated C-terminal
fusions (Figure 2A, L1 to L37), N-terminal fusions (Figure 2A, L38 to L53), as well as
circular permutants with Lifeact connecting the original N- and C-termini (Figure 2A, L54
to L72). To remove flexible stretches, we did not include linkers in the fusions, and we
further shortened the N- or/and the C-termini of GFP. At the onset of this study, the crystal
structure of Lifeact bound to F-actin (Figure S1C,D and (Belyy et al., 2020; Kumari et al.,
2020) was not yet known. Circular dichroism and NMR spectroscopies had suggested
that Lifeact forms a helix between residues 2 and 10 (Riedl et al., 2008), though a helix
from position 3 to 17, i.e. essentially encompassing the entire Lifeact sequence, was most
commonly found with the secondary structure prediction server Robetta (predictions not
shown). These findings, the fact that the very C-terminus of Lifeact was not reported to
be conserved in fungi (Riedl et al., 2008), and our quest for the minimal actin-binding
stretch of Lifeact, prompted us to also shorten Lifeact on either or both its termini, aiming
at removing flexible stretches that are not essential for actin binding.

We screened the generated fusions with respect to their fluorescent levels, their
capacity to bind F-actin, their localization, and the extent to which GFP is constrained
based on polarimetry measurements (Figures 2A and S1A). Extensive shortening of the
msfGFP termini did not affect fluorescence of terminal fusions, in line with the results of
our truncation screen. The fluorescence of cpGFP fusions, on the other hand, was
expected to depend on the length and composition of the linker connecting the original
termini. Earlier studies suggested that linkers comprising at least 20 residues are needed
to connect the GFP barrel ends, i.e. in between the end of β11 and the beginning of β1
strands, to allow for stable cpGFP folding and fluorescence, and this using flexible
glycine-rich linkers (Akemann et al., 2001; Baird et al., 1999; Topell et al., 1999). Not
surprisingly, fluorescence was severely compromised or absent for cpGFP fusions when
using 11-16 residue-linkers.

The capacity of the fusions to bind F-actin, using SF labeling as a readout,
depended, as expected, on the length of Lifeact. Shortening of Lifeact on either or both
termini when fused to full-length msfGFP showed that (a) Val3 is essential for binding F-
actin, (b) Gly2 is not needed *per se* for binding F-actin but that it contributes to the latter,
and (c) the six C-terminal residues are also not essential for F-actin binding; these
residues are also the least conserved ones among fungi (Figures 2A and S1B-G).
Importantly, the proximity of either terminus of Lifeact to the terminus of GFP, notably
when combining shortened Lifeact with shortened GFP termini, was critical for F-actin
binding. This was particularly evident in cpGFP fusions with shortened GFP termini,
whereby F-actin binding was compromised despite the presence of full-length Lifeact,
suggesting that a minimum of flexibility is required on either side of the actin-binding
moiety to allow for F-actin binding. This latter effect proved eventually to be the bottleneck
for constraining efficiently GFP in cpGFP fusions given that even moderate shortening of
the GFP termini compromised F-actin binding.

Under our low-level expression conditions (see methods for promoter details), all
fusions labeled all types of SFs, notably dorsal, ventral, peripheral, perinuclear actin cap
and arc SFs, including focal adhesions (FAs), as well as mitochondrial actin (Figure S1H).
To exclude that the additional localization of fusions to arc nodes reflected GFP
dimerization-related artifacts (Figure S1I), we compared fusions bearing one, two or all
three GFP monomerizing mutations (Figure 1I); they all localized to arc nodes excluding
such a scenario.

Terminal GFP fusions proved to be the most efficient way to constrain GFP mobility (Figures 2A-C and S1A). Polarimetry measurements showed that removing only the linker between full-length Lifeact and full-length GFP did not have any effect on GFP mobility (Figures 2A and S1A). The additional removal of seven residues from the N-terminus of GFP or ten residues from its C-terminus, was necessary to start constraining its mobility. Combining shorter GFP termini with shorter Lifeact termini also proved determinant, notably in N-terminal Lifeact fusions whereby the removal of a single residue, Gly2, from Lifeact, reduced ψ angle values by several tens of degrees (compare L42 with L45 in Figure 2A,C). Altogether, polarimetry measurements showed that we succeeded to immobilize GFP both in N- and C-terminal fusions with Lifeact. We chose to focus on fusions L22 (LifeactΔC4-msfGFPΔN7) and L45 (msfGFPΔC9-LifeactΔN2) as the best performing reporters for further functional characterization and F-actin organization measurements in live cells and tissues.

Engineering actin filament organization reporters using the Utrophin Calponin Homology Domain.

The N-terminal 261 amino acids of human utrophin contain an F-actin-binding calponin homology domain known as Utr-CH or Utr261 (Burkel et al., 2007), which is widely used for visualizing F-actin in live cells and tissues. The original GFP fusion is N-terminal to the Utr-CH domain (Burkel et al., 2007) but C-terminal EGFP fusions have also been used successfully (Belin et al., 2014). To constrain GFP mobility in fusions with Utr261, we generated N-terminal fusions (Figure S2A, U1 to U24), C-terminal fusions (Figure S2A, U25 to U42), as well as circular permutants with Utr261 connecting the original N- and C-termini (Figure S2A, U43 to U63). We did not include linkers in the fusions, and we further shortened the N- or/and the C-termini of GFP. The structure of the utrophin calponin homology domain bound to F-actin (Kumari et al., 2020) was not available at the time of the beginning of this study, but the biochemical and structural data concerning the N-terminal residues 28-261 of human utrophin (Keep et al., 1999; Moores and Kendrick-Jones, 2000) provided already key insights. Although the N-terminal 27 residues of utrophin maximize its affinity for F-actin, they are dispensable for F-actin binding (Moores and Kendrick-Jones, 2000). The susceptibility of these same residues to degradation further suggested that they are not part of a compact structure and might be inherently flexible (Moores and Kendrick-Jones, 2000). This observation aligns with their partial disorder in their complex with F-actin (Kumari et al., 2020). A truncation mutant of Utr261, Utr230-EN (Belin et al., 2013), was also shown to bind cytoplasmic actin filaments. These findings prompted us to also shorten Utr261 on either or both termini.

We screened the generated utrophin fusions along the same lines as for the Lifeact fusions (Figures S2B-G). All constructs were fluorescent except the three cpGFP fusions with the shortest N- and C-termini of GFP, suggesting again that a minimum of flexibility is needed to allow connecting the original N- and C-termini of GFP while binding to F-actin. Shortening of Utr261 on either or both termini in terminal fusions with full-length msfGFP confirmed that the N-terminal 27 residues of Utr261 are indeed dispensable for binding F-actin, and further showed that the C-terminal 31 or 39 residues are also not required for binding F-actin (Figures S2A-G). Importantly, the proximity of residues 29-32 in the N-terminal helix of Utr261 to the C-terminus of GFP, notably when combining shortened Utr261 with shortened GFP termini, was critical for F-actin binding (Figures S2A and S2E), in line with the structure of the UtrCH-F-actin complex (Kumari et al., 2020). The localization of all fusions was similar to the one of Lifeact fusions (Figure S2C-F).

Terminal GFP fusions proved to be the most efficient way to constrain GFP mobility (Figures 3A-C and S2A,B). As expected from the results of the Lifeact fusions, removing only the linker between full-length Utr261 and full-length GFP did not have any effect on

GFP mobility. The additional removal of nine residues from the N-terminus of GFP and of the N-terminal 27 residues of Utr261 were necessary to start constraining its mobility. It is noteworthy that the additional removal of a tenth residue from the GFP C-terminus was sufficient to reduce ψ angle values by several tens of degrees (compare U13 with U20 in Figure 3A,B). Interestingly, all C-terminal GFP fusions, including combinations of the shortest C-terminus of Utr261 and the shortest N-terminus of GFP, and the recently reported construct UG7 (Nakai et al., 2019), were flexible, reflecting most likely inherent flexibility in the very C-terminus of Utr261. We decided to focus on fusion U20 (msfGFPΔC10-Utr28-222) as the best performing reporter for further functional characterization and F-actin organization measurements in live cells and tissues.

Engineering actin filament organization reporters using F-tractin.

The N-terminal residues 9-52 of the rat enzyme inositol triphosphate 3-kinase A (ITPKA), also known as F-tractin-P (P for prototype), were shown to contain an F-actin-binding domain, a C-terminal GFP fusion of which is widely used as a reporter of F-actin localization in live cells (Johnson and Schell, 2009; Yi et al., 2012) (Figure 3D and S3A). A slightly shorter peptide, N9-40, has been shown to retain F-actin binding, was given the name F-tractin (Yi et al., 2012), and is used interchangeably with F-tractin-P for visualizing F-actin in live cells (John Hammer, National Institutes of Health, Bethesda, MD, USA, personal correspondence). To constrain GFP mobility in fusions with F-tractin and F-tractin-P, we generated C-terminal fusions (Figure S3C, F1 to F22), N-terminal fusions (Figure S3C, F23 to F27), as well as circular permutants with the F-tractin peptide connecting the original N- and C-termini (Figure S3C, F28 to F33). We did not include linkers in the fusions, and we further shortened the N- or/and the C-termini of GFP. The structure of F-tractin or F-tractin-P, alone or in complex with F-actin, has not been solved to date. Secondary structure prediction of F-tractin-P using the program JPred3 in the original article suggested that F-actin binding resides in a putative α-helix comprising residues ~30-40 (Johnson and Schell, 2009). Secondary structure prediction using multiple programs, including JPred4, PHD, Phyre2, RaptorX and AlphaFold, suggests that the glycine- and proline-rich N-terminal ~30 residues are unstructured, with residues ~30-50 predicted to form a helix (Figure S3B). These predictions prompted us to also shorten F-tractin on either of its termini.

We screened the generated F-tractin fusions similarly to the Lifeact and utrophin fusions (Figures S3C-E). Unlike the previous screens, all constructs were now fluorescent, including cpGFP fusions with the shortest N- and C-termini of GFP, reflecting most likely the fact that the N-terminus of F-tractin is unstructured and does not bind F-actin thus imposing less constraints for connecting the original termini of cpGFP. Shortening of F-tractin and combinations thereof with shortened GFP termini showed (a) that the N-terminal residues 9-14 are dispensable for F-actin binding and (b) that residues 37-40 are critical for F-actin binding (Figure S3C and S3E). All fusions localized similarly to Lifeact and utrophin fusions (Figure S3E).

C-terminal GFP fusions of F-tractin proved to be the most efficient way to constrain GFP mobility (Figures 3D-F and S3C,D). The additional removal of seven residues from the N-terminus of GFP was necessary to constrain its mobility. In line with our results from the Lifeact and utrophin screens, it was very striking that the additional removal of a single residue from the C-terminus of F-tractin was sufficient to reduce ψ angle values by several tens of degrees (compare F5 with F11 in Figure 3D,E). Interestingly, all C-terminal GFP fusions of F-tractin-P, including combinations with the shortest N-termini of GFP, were flexible; we observed the same behavior for all N-terminal fusions of F-tractin, including combinations with the shortest C-termini of GFP. We interpret both observations as reflecting the inherent flexibility of the respective F-tractin termini. The fusion F11 (F-

tractinN9-39-msfGFPΔN7) turned out to be the best performing reporter for F-actin organization measurements in live cells.

Engineering actin filament organization reporters using Affimer6.

Affimers, originally named Adhirons (Tiede et al., 2014), are synthetic, non-antibody-based protein binders that can be engineered to bind specific proteins of interest with high affinity and specificity. Among the recently developed Affimers is the F-actin-binding Affimer, Affimer6 (Lopata et al., 2018), an N-terminal GFP fusion of which can be used to monitor F-actin localization in live cells. To constrain GFP mobility in fusions with Affimer6, we generated C-terminal fusions (Figure S3F, Af1 to Af11), N-terminal fusions (Figure S3F, Af12 to Af16), as well as circular permutants with the Affimer scaffold connecting the original N- and C-termini (Figure S3F, Af17 to Af25). We did not include linkers in the fusions, and we further shortened the N- or/and the C-termini of GFP. The structure of the Affimer scaffold revealed a very compact fold, with hardly any flexible unstructured residues at its termini (Tiede et al., 2014) (Figure S3I). Thus, we only attempted to shorten its C-terminus to remove potentially flexible residues that could contribute to GFP mobility in C-terminal GFP fusions. F-actin binding is not expected to be affected since the actin-binding loops are far from the C-terminus (Tiede et al., 2014).

We screened the generated Affimer6 fusions in a similar manner to the other ABD fusions (Figure S3F-I). All constructs were fluorescent apart from the three cpGFP fusions with the shortest N- and C-termini of GFP: the highly compact structure of the Affimer scaffold does most likely not provide the flexibility needed to connect the original N- and C-termini of GFP in these cpGFP fusions. F-actin binding was compromised only for shortened Affimer C-termini combined with highly shortened GFP termini; it is possible that the GFP in these fusions adopts a position that interferes with the actin-binding loops. All Affimer6 fusions localized in an indistinguishable manner from the other ABD fusions (Figure S3H).

Fusing full-length Affimer6 C-terminally to the shortest N-terminus of GFP proved to be the most efficient way to constrain GFP mobility (Figures 3G-I and S3F,G). It was again remarkable that the presence of an additional single residue at the N-terminus of GFP was sufficient to increase ψ angle values by several tens of degrees (compare Af6 with Af7 in Figure S3F,G). N-terminal GFP fusions of Affimer6 with C-terminally truncated GFP also constrained GFP mobility, but much less efficiently, as was the recently reported Affimer6-based construct POLArISact (Sugizaki et al., 2021). We decided to use fusion Af7 (Affimer6-msfGFPΔN12) as the best performing reporter for further functional characterization and F-actin organization measurements in live cells and tissues. The ψ values we obtained with this reporter are the lowest ones among all reporters, reflecting a highly constrained GFP, and making it the most sensitive one for detecting small changes in actin filament organization.

Engineering red FP-based actin filament organization reporters.

Having succeeded to generate GFP-based reporters for the different ABDs, we naturally embarked on the making of the red FP fusion counterparts. The fact that robustly folding GFPs tolerate much more extensive terminal truncations without losing fluorescence (Figure 1G,H) prompted us to undertake a similar approach. Thus, we compared side by side the sensitivity of the widely used red FP, mApple (Cranfill et al., 2016; Shaner et al., 2008), and of superfolder Cherry 2 (sfCherry2) (Feng et al., 2017) to terminal truncations. Both FACS and imaging corroborated our results with green FPs: whereas mApple is sensitive to N-terminal truncations, sfCherry2 tolerates missing its entire N- (ΔN12) or C-terminus (ΔC4) (Figure S4A,B), making the latter the best choice for constraining mobility in its fusions to actin-binding domains.

We decided to generate selectively sfCherry2-based terminal fusions for Lifeact and Affimer6. The length and composition of the termini of sfCherry2 are not the same as for GFP necessitating a minimum of screening, but the results from our previous screens helped narrow down our efforts to a limited set of constructs. F-actin binding was, as expected, impaired by the proximity of shortened FP termini to shortened Lifeact. Indeed, the best performing constructs were the ones combining full-length Affimer6 and Lifeact with the most extensively truncated sfCherry2 termini (Figure 3J,K), very similarly to the respective best performing GFP-based ones. The fusions Af30 (Affimer6-sfCherry2ΔN12) and L81 (sfCherry2ΔC4-Lifeact) were the best performing ones and are ideally placed for F-actin organization measurements in live cells and tissues.

Engineering actin filament organization reporters using G-actin.

N-terminal GFP fusions of G-actin, with a flexible linker in between the GFP and the G-actin, are widely used for monitoring actin localization in live cells and tissues. Early studies using such fusions to *Dictyostelium discoideum* actin (Westphal et al., 1997), to human non-muscle beta G-actin (Ballestrem et al., 1998) and to the *Drosophila* non-muscle G-actin Act5C (Verkhusha et al., 1999) showed that such fusions are functional, as long as their expression levels are kept low; the choice of an appropriate promoter to keep GFP-actin levels low was shown to be critical for their functionality (Ballestrem et al., 1998). An important finding was that C-terminal tagging of G-actin, even with tags as small as a hexahistidine tag or a dodecapeptide, impairs its incorporation into actin filaments (Brault et al., 1999; Chen et al., 2012). *Drosophila* expressing such C-terminally tagged G-actin have flight muscle with virtually no detectable sarcomeric organization and are flightless, but N-terminal fusions with the same tags restore sarcomeric organization and flight capacity (Brault et al., 1999). Thus, to constrain GFP mobility in fusions with human non-muscle beta G-actin, we generated exclusively N-terminal fusions. We did not include linkers in these fusions, and we additionally shortened the C-termini of GFP (Figure 4A, A1 to A5). We also generated an N-terminal fusion with a tetracysteine peptide (Figure 4A, A6).

As an alternative to N-terminal fusions and to maximize our chances to constrain GFP mobility, we also considered generating intramolecular GFP fusions. (Bendezu et al., 2009) succeeded to generate a fully functional intramolecular mCherry fusion of the bacterial actin homolog MreB by inserting mCherry right before helix 7 of MreB (van den Ent et al., 2001). Very interestingly, a second study by (Chen et al., 2012) succeeded to generate a functional intramolecular fusion of fission yeast actin by inserting a dodecapeptide-based tetracysteine tag into Ser233-Ser234 of the loop following helix 7 (h7, hereafter), and use FlAsH labeling to monitor F-actin in fission yeast cells. Motivated by these studies, we chose to engineer intramolecular fusions by inserting either full-length GFP (Figure 4A, A7 to A23), the GFP strand β11 (Figure 4A, A24 to A37), or a tetracysteine peptide (Figure 4A, A38 to A47), either before or after h7 (Figure 4B). We used the exact same insertion site after h7 as in the study by (Chen et al., 2012). Fluorescence imaging showed that only fusions into the loop following h7 incorporated into actin filaments (A8, A25, A40 and A41 in Figure 4A). We thus focused on this insertion site for subsequent screening.

As a proof of principle that such intramolecular fusions are functional, we compared the functionality of intramolecular full-length msfGFP fusions ("iGFP" for short) for the two vertebrate non-muscle actin isoforms, namely human beta- and gamma-actin isoforms, using the exact same insertion site after h7, like for construct A8 (Figure 4A). The two isoforms have distinct localizations and functions (Baranwal et al., 2012; Chen et al., 2017; Dugina et al., 2009; Shum et al., 2011), and intramolecular GFP fusions would provide new less-perturbative tools to study actin isoform function given that they differ only by four amino acids at their very N-terminus. To this end, we generated stable

inducible HeLa cell lines expressing the iGFP-actin isoform fusions (Figure S4C-M). Quantification of mitotic and multinucleated cells upon depletion of the endogenous isoforms using isoform-specific siRNAs showed that our iGFP fusions rescued these defects (Figure S4C-F). The localization of the iGFP fusions during mitosis and in nondividing cells recapitulated their distinct distributions (Figure S4I-M), corroborating previous studies (Chen et al., 2017; Dugina et al., 2009) and confirming further their functionality. Motivated by these results and in order to constrain fluorophore mobility in intramolecular fusions with human beta G-actin, we generated constructs that did not include linkers and where the N- and C-termini of GFP were shortened (Figure 4A). Three out of four serines within our insertion site are highly conserved across actin sequences (Figure 4B,C): we also generated constructs with differences in the exact number and position of these serines to establish possible effects on fluorophore mobility.

We screened the generated G-actin fusions with regard to their fluorescence, their capacity to integrate into F-actin, their localization and the extent to which GFP or FlAsH is constrained based on polarimetry measurements (Figures 4A-E and S4N,O). All terminal and intramolecular fusions with GFP and tetracysteine peptides were fluorescent. However, the only intramolecular β11-based fusion that was fluorescent was the one that included linkers (fusion A25 in Figure 4A): we reasoned that the absence of linkers and further shortening in the subsequent β11-based fusions did not provide the flexibility needed for the complementation of β11 with GFP1-10 (Cabantous et al., 2005). The absence of linkers and the additional shortening of the GFP C-terminus in terminal fusions did not compromise binding to F-actin, but seemed to enrich less these fusions in myosin-II containing SFs (Figures 4A and S4O). Myosin-II interacts with the actin N-terminus providing a possible explanation for the latter observation. The absence of linkers in intramolecular GFP fusions did not compromise binding to F-actin, either. F-actin binding upon additional shortening of both GFP N- and C-termini in intramolecular fusions depended on the extent of shortening, as well as the exact number and position of the serine residues encompassing the insertion site. As expected, the use of a truncated CMV promoter for low-level expression was critical, with the widely used full-strength CMV promoter leading systematically to aggregation (Figure S4O). All G-actin fusions localized similarly to the other ABD-GFP fusions (Figure S4O). Interestingly, intramolecular tetracysteine peptide fusions labeled additionally nuclear F-actin (A41 in Figure S4O), showing nuclear F-actin bundles morphologically very similar to ones detected with the use of a nuclear actin chromobody (Lamm et al., 2020) and an actin-NLS-FLAG construct (Kokai et al., 2014; Lawson et al., 2022). The small size of these peptides compared to the size of GFP could possibly explain this difference: GFP fusions were typically excluded from the nucleus, consistent with such an explanation.

Terminal and intramolecular GFP fusions without linkers and with extensively shortened GFP termini were most efficient to constrain GFP mobility (Figures 4A,D-E and S4N). We note that GFP mobility is much less constrained than in fusions with actin-binding peptides and domains: we attribute this increased mobility to the inherent flexibility of the actin N-terminus and of the insertion site loop in terminal and intramolecular fusions, respectively. Despite this difference, constrained GFP fusions directly to G-actin can still be useful for studies focusing on G-actin itself, notably actin isoforms. The fusions A4 (msfGFPΔC10-actin) and A18 (actin-h7-msfGFPΔN7ΔC11-SSSSactin) are the best performing ones for F-actin organization measurements in live cells.

Comparison of the localization of our organization reporters to different F-actin populations, notably different types of SFs, in U2OS cells did not show any significant differences. At the low-expression levels we used and during the transient expression of the reporters, we did not detect any evident signs of perturbation in terms of SF biogenesis, maintenance or organization. Given that different actin-binding probes have

different affinities for F-actin, and that cells and tissues express tens of actin-binding proteins to regulate the dynamics, localization and specific geometries of actin assemblies related to specific functions, it is expected that these same probes behave differently in different cellular contexts. Thus, to gain deeper insights into the functionality of our organization reporters, we chose to test them in a more physiological context, notably in three different genetically tractable *in vivo* model systems: the fission yeast *S. pombe*, the nematode *C. elegans* and the fruit fly *Drosophila melanogaster*.

Actin filament organization reporters in fission yeast.

We generated transgenic fission yeast expressing selected reporters. We focused on the best performing reporters from our screen on SFs, in particular L22, L45, U20 and Af7, and included the respective constructs with unconstrained GFP for comparison, i.e. L2, L38, U7 and Af1, respectively. Cells expressing these reporters under the control of the promoter of the actin cytoskeleton regulator Cdc42, looked healthy, with no signs of vacuolation, and were rod-shaped, suggesting the absence of major polarity defects. We started by comparing the localization of the reporters in populations of actively dividing yeast cells (Figure S5A-D). Actin in fission yeast is found in three distinct structures: actin patches, actin cables and the cytokinetic actomyosin ring (Figure S5B). Although all reporters labeled all three structures, there were dramatic differences in their respective enrichments, as assessed by the differences in labeling intensities of the different structures (Figure S5A,B), as well as in the numbers of patches and cables labeled by each reporter (quantification in Figure S5C,D). Patches and, to a lesser extent, rings, were labeled well with Lifeact constructs, whereas cables and rings, and, to a lesser extent, patches, were most prominent with Affimer6 constructs (Figure S5A-D). Utrophin labeling seemed altogether very inefficient. We attribute these striking differences in localization to differences in the affinities of the reporters for the different actin structures.

To assess the functionality of the reporters, we examined their effect in three different contexts: the timing of the different stages of cytokinesis (Figure 5A,B), the growth of cells under sensitized conditions, notably the presence of the Arp2/3 complex inhibitor CK666 and the G-actin sequestering drug latrunculin A (LatA) (Figure 5C), and finally genetic interactions with a profilin mutant (Figure 5D). Quantification of the timing of cytokinesis stages showed that cytokinesis proceeded overall similarly for all reporters with respect to a control strain expressing only cytokinetic markers, with cells completing cytokinesis within ~40-50 minutes in all cases (see mean times for each stage in the legend of Figure 5B). We observed the most important delays for the Affimer6 constructs, which showed increased maturation and constriction times, and for some of the Lifeact constructs, notably L22, which took longer for the assembly (Figure 5B).

The dilution assays to assess cell viability and growth in the presence of CK666 and LatA led to three observations. First, there was no difference in cell growth upon expression of the reporters with DMSO as a vehicle control, reflecting their overall nonperturbative character. Second, cell growth in the presence of CK666 was slightly impaired upon expression of L22 and the effect was even more significant for the original Lifeact construct, which is expressed at a higher level, from the actin promoter (Huang et al., 2012). Third, the inability of cells to grow in the presence of LatA was significantly reversed for the Affimer6 constructs, as well as the original Lifeact construct and L22 (Figure 5C). The main effect of CK666 is on actin patches, for the formation of which Arp2/3 is essential. The CK666 results thus suggest that the original Lifeact and, to a lesser extent, L22 interfere with branched actin formation. LatA, on the other hand, at the low concentration used, impacts primarily the formation of cables, which are not essential for growth, and of cytokinetic rings which are essential for cell division. The LatA results thus reveal a stabilization effect on the rings upon expression of the Affimer6 constructs, the original Lifeact construct and L22. The increased cytokinetic ring maturation and

constriction times for Affimer6 and increased ring assembly time for L22 we observed could well be explained by the same mechanisms revealed from the drug assays. Finally, dilution assays in the presence of a thermosensitive profilin mutant did not show any significant difference for the reporters, suggesting that they do not interfere with actin nucleation and polymerization *per se* (Figure 5D). Altogether, these results show that our reporters are largely nonperturbative, however specific reporters can contribute to mild perturbations, notably in the context of sensitized conditions.

As an example of the applicability of the reporters *in vivo*, we chose to measure actin filament organization in the constricting cytokinetic ring of live dividing fission yeast cells. We focused on the Affimer6 constructs Af1 and Af7 and the Lifeact constructs L2 and L22 which were most efficient in labeling the cytokinetic ring (Figure S5A,B). Measurements of mean actin filament orientation (ρ angles) and alignment (ψ angles) in constricting rings resembled the behavior of the respective constructs on SFs. The unconstrained GFP fusions Af1 and L2 displayed very high ψ values, whereas Af7 and L22 fusions bearing constrained GFP showed statistically significantly lower ψ values (Figure 5E-G). The measured values were comparable to the ones on SFs, suggesting highly aligned actin filaments, with filament alignment persisting throughout ring constriction (Figure S5G). Considering that Af7 and L22 dipoles are parallel to actin filaments (Figures 2B and 3G), the quantification of ρ angle distribution with respect to the division plane showed that actin filaments are parallel to the constricting ring axis (Figure S5E,F). Our results are fully consistent with recent electron cryotomography of dividing fission yeast that revealed straight, overlapping actin filaments running nearly parallel to each other and to the membrane (Swulius et al., 2018). The combination of the organization reporters with mutants promises to further our understanding of the mechanisms that drive actomyosin remodeling in cell division.

Actin filament organization reporters in *C. elegans*.

Next, we sought to test selected reporters in the context of animal morphogenesis, notably *C. elegans* embryonic elongation (Figure S6A). To this end, we generated transgenic *C. elegans* expressing the organization reporters L22, L45 and Af7 under the control of an epidermal promoter (Figure S6B). To assess their functionality, we quantified embryonic lethality by scoring egg hatching in comparison with wild-type embryos and embryos expressing the original Lifeact-GFP reporter. The measured lethality was comparable to wild-type embryos and lower than the one of Lifeact-GFP expressing embryos (Figure 6A). In a second functionality test, we filmed living elongating embryos and quantified embryonic length until hatching: embryonic elongation proceeded similarly for all strains (Figure 6B), again confirming the minimally perturbative character of the reporters.

Elongation of the *C. elegans* embryo proceeds along its anterior-posterior axis, increasing in length about fourfold and decreasing in circumference about threefold, with practically no cell divisions nor cell rearrangements (Priess and Hirsh, 1986). The epidermis comprises three major cell types, namely dorsal, ventral and lateral seam cells. Whereas actin filaments progressively form circumferential bundles in dorsal and ventral epidermal cells (DV cells), actin filaments in seam cells have been reported to stay rather disorganized throughout elongation (Gally et al., 2009; Vuong-Brender et al., 2017). Recent work evidenced an interplay of stress anisotropy in seam cells and stiffness anisotropy in DV cells as critical for elongation through a circumferential squeezing-like mechanism (Vuong-Brender et al., 2017), prompting us to quantify actin filament organization in DV and seam cells as a function of elongation.

The localization of L22, L45 and Af7 in the elongating epidermis was in line with phalloidin staining (Priess and Hirsh, 1986) and the localization of a GFP fusion to the actin-binding domain of the spectraplakin VAB-10 (Gally et al., 2009; Vuong-Brender et

al., 2017), showing the characteristic circumferential bundles in DV cells and a fuzzier mesh-like distribution in seam cells (Figure S6C,D,F). Considering that Af7 and L22 dipoles are parallel to actin filaments (Figures 2B and 3G) and that L45 dipoles are perpendicular to actin filaments (Figure 2C), quantification of actin filament organization in the circumferential bundles of DV cells in >2-fold elongated (>2F) embryos showed that actin filaments are parallel to the circumferential bundles (Figure 6I,J and Figure S6D-G), as expected, with ψ values comparable to the ones on SFs, suggesting highly aligned actin filaments (Figure 6C). Strikingly, quantification of actin filament orientation with respect to the DV/seam boundary showed that the polarization of actin filament orientation was already present in 1.5-fold elongated (1.5F) embryos before the formation of distinct circumferential bundles (Figure 6E,F and Figure S6C), with actin filament alignment increasing as bundles form (Figure 6D). What was, however, even more unexpected was that actin filament organization in seam cells was also polarized: seam cells in 1.5F embryos contained already actin filaments oriented perpendicular to the DV/seam boundary and with regions of actin filament alignment. As elongation proceeds, this polarization becomes progressively more pronounced, with actin filaments oriented essentially perpendicular to the DV/seam boundary, just like in DV cells, and with regions of high actin filament alignment (Figure 6D-J and Figure S6E,G). The experimental demonstration of the polarization of the actin cytoskeleton during *C. elegans* embryonic elongation, a first of its kind, reveals the potential of the new organization reporters, which, combined with mutants and biophysical measurements, promise to uncover the mechanisms that drive animal morphogenesis.

Actin filament organization reporters in *Drosophila*.

We finally chose to characterize selected organization reporters in *Drosophila*. To this end, we generated transgenic *Drosophila* expressing reporters with the inducible GAL4/ UAS expression system, allowing us to express them in the early embryo, the wing and the indirect flight muscle using respective tissue-specific promoters. Actin filament organization measurements in the actomyosin rings of living cellularizing embryos recapitulated earlier polarimetry measurements in fixed phalloidin-stained embryos (Mavrakis et al., 2014) confirming the arrangement of highly aligned filaments following the contour of the associated membrane front (Figure 7A and Figure S7A). Apicolaterally to this front and at the basal-most part of the lateral membranes, a basal adherens junction has been reported to form (Hunter and Wieschaus, 2000). In this junction, actin filaments were found to be highly aligned to each other, following the junction contour in line with a belt-like arrangement (Figure 7B).

We assessed the functionality of the reporters in two systems that depend on the integrity of the actin cytoskeleton, namely the adult wing and the adult flight muscle. To measure their effect on wing growth, we compared the wing area in wings expressing the reporters with the one in wings bearing only the wing-specific promoter (control), wings expressing the original Lifeact-GFP, and wings expressing a dominant-negative form of the insulin receptor expected to reduce wing growth. Wing growth was indeed reduced by ~46% for the latter, but was largely comparable, within less than 8% of change, among the control and the other strains (Figure S7B). As a second functionality test, we performed flight tests to compare the flight ability of strains expressing different reporters with the one of strains bearing only the muscle-specific promoter (control) and strains expressing two widely used actin localization reporters: the actin-binding domain of moesin (GFP-GMA) and Lifeact-GFP (Figure 7C). Although all reporters localized largely as expected in the muscle sarcomeres (Figure S7C), flight tests showed significant differences among the strains. The flight ability of flies expressing L45 and U20 was comparable to the one of control and GFP-GMA flies. However, flies expressing either L22 or the original Lifeact-GFP, performed very poorly, with Af7 expressing flies being

entirely flightless (Figure 7C); the development of the flight muscle in the latter strain was
altogether impaired (Figure S7C). The muscle-specific promoter being active in all
embryonic, larval and adult muscle, we attributed this dramatic effect to the expression of
the reporters throughout muscle development, and reasoned that their limited expression
after muscle development could be less perturbative. In support of this scenario, and with
the use of the temperature-sensitive GAL80ts system to express the reporters only in a
narrow time window of a few days in adult muscle, the flight ability of all strains was
improved, including the strains expressing the original Lifeact-GFP, L22, and Af7 (Figure
7C). Muscle morphology and sarcomere localization were comparable to the control in all
cases, including now for Af7 (Figure S7D). Importantly, despite their limited temporal
expression, the fluorescence levels of the reporters rendered them usable for polarimetry
measurements in the flight muscle.
To this end, we dissected live flight muscle expressing L22, L45 and Af7, and the
original Lifeact-GFP for comparison, and measured actin filament organization in the
respective myofibrils. Not surprisingly, ψ values in the original Lifeact-GFP were too high
to render this fusion usable for organization measurements, but L22, L45 and Af7
behaved as expected, with Af7 further displaying the lowest ψ values i.e., the highest
filament alignment, from all systems (Figure 7D), in line with the crystal-like arrangement
of actin filaments in sarcomeres reported by EM (Loison et al., 2018). Considering the
GFP dipole orientations of L22, L45 and Af7 with respect to actin filaments, actin filament
orientation and alignment maps revealed the expected high order within and across
different myofibrils in the muscle (Figure 7E-G). The potential of the reporters to quantify
actin filament organization in live muscle, combined with the genetically-tractable
character of *Drosophila* and its use as model for studying muscle biogenesis and
pathophysiology, promises to contribute novel insights into the regulation of muscle
structure and function.
767
768 **DISCUSSION**
769
Despite the established contribution of the actin cytoskeleton to cell and tissue integrity
and function in a wide range of biological processes, and its direct link to animal
pathophysiology, measuring its organization in real time in living cells remains a
challenge. Polarimetry exploits the interaction between fluorophore dipoles and polarized
light for probing molecular organization. However, fluorophore mobility must be
constrained to allow for such measurements. Our novel genetically-encoded reporters
bearing constrained fluorescent proteins enable the quantification of actin filament
orientation and alignment, and dynamical changes thereof, in living cells. We succeeded
in engineering constrained GFP fusions to widely used actin localization reporters,
namely Lifeact, the Utrophin calponin homology domain, F-tractin, Affimer6 and G-actin
itself, thus providing the possibility to correlate any biological process of interest with live
measurements of actin filament organization by polarimetry. We further provide reference
angle values for each reporter in SFs that can be used for comparison and interpretation
of organization measurements in any F-actin population of interest with no need for any
a priori knowledge of its organization.
Even though ensemble measurements cannot resolve individual filaments, their
molecular-scale organization at a given image pixel is detectable and can be quantified
by polarimetry. Moreover, the mesh size of the actin network is typically on the order of
100 nm or less (Chugh and Paluch, 2018), making it so far only attainable in a fixed cell
context with EM and single molecule localization microscopy. Thus, the capacity to obtain
quantitative measurements of actin filament orientation and alignment per image pixel in
a living cellular context is of significant added value. Cells continuously and dynamically
remodel actin filaments in order to accomplish specific biological functions, and do so
through the complex regulation of actin filament architecture and dynamics. Bottom-up

approaches with purified proteins are key for assigning specific functions, for example filament branching, to distinct actin-binding proteins (Mullins and Hansen, 2013). However, whether and how these assigned functions account for F-actin organization in cells, in the presence of tens of other actin interactors, can now be formally tested by combining live cell organization measurements with mutants or treatments affecting one or several interactors.

The use of the reporters in the context of cell and tissue morphogenesis, including during embryogenesis and organogenesis, promises to uncover how specific geometries of actin filaments contribute to function. Organization measurements with the novel reporters in the presence of mutants will further provide new insights into the developmental regulation of F-actin organization. The genetically-encoded character of the reporters renders them also fully compatible with animal disease models, enabling *in vivo* studies on the role of any gene of interest in F-actin organization in the context of pathophysiology, for example in muscle development, maintenance and repair, in health and disease. Being able to measure changes in filament organization in real-time will additionally help generate accurate biophysical models with experimentally testable predictions regarding how F-actin organization impacts function.

The novel F-actin organization reporters are compatible with all fluorescence microscopy techniques routinely used for live ensemble imaging that can be coupled with polarized fluorescence imaging, such as transmission polarized microscopy (Abrahamsson et al., 2015; DeMay et al., 2011b), wide-field (Vrabioiu and Mitchison, 2006), confocal (Kress et al., 2013) and spinning disk confocal microscopy (Wang et al., 2013), total internal reflection microscopy (Sund et al., 1999) and two-photon microscopy (Benninger et al., 2005; Ferrand et al., 2014; Gasecka et al., 2009; Lazar et al., 2011). The reporters are also compatible with super-resolution imaging techniques, either based on polarized structured illumination microscopy employing standard FPs (Hafi et al., 2014; Zhanghao et al., 2019), or based on single molecule orientation and localization microscopy using photoactivatable or photoconvertible FPs (Brasselet and Alonso, 2023). It is important to note that the introduction of a single point mutation to sfGFP or of a few point mutations to sfCherry2 are sufficient to generate the respective photoactivatable versions (Feng et al., 2017; Slocum and Webb, 2017). Given that these mutations are within the barrel structure, we do not expect them to alter the mobility of the FP. Our strategy can thus be simply adapted to the context of single-molecule organization measurements using, for example, single-particle tracking PALM coupled to polarization splitting (Rimoli et al., 2022; Valades Cruz et al., 2016) or Point Spread Function engineering (Brasselet and Alonso, 2023). The ultimate choice of the technique will depend on the desired spatial and temporal resolution, optical sectioning and imaging depth, as well as on the requirement for capturing the actin filament organization dynamics under study. Thus, the use and interpretation of the organization measurements using the reporters must take into account the specificities and limitations of each technique. All measurements in this study have used a spinning disk confocal microscope with polarized excitation for second-scale measurements of F-actin organization. If higher temporal resolution is needed, for example for monitoring fluctuating filaments or rapidly remodeling cells, a wide-field microscope with polarization splitting would allow for subsecond-scale measurements using the exact same reporters (Kampmann et al., 2011; Mattheyses et al., 2010; Mehta et al., 2016; Nordenfelt et al., 2017; Swaminathan et al., 2017; Vrabioiu and Mitchison, 2006). Our open-source software, PyPOLAR, for analyzing polarimetry data generated by a range of different techniques has been designed explicitly for use by biologists and biophysicists, and promises to further facilitate the use of the novel molecular tools.

The genetically-encoded character of the reporters makes them particularly useful for live cell and tissue studies, both *in vitro* and *ex vivo*, as well as *in vivo* in the context of genetically-tractable model organisms, as exemplified by their use in fission yeast,

C. elegans and *Drosophila*. The use of specific promoters and inducible expression systems can control their spatial and temporal expression, as well as their levels. As expected for any actin-binding protein, there is an inevitable risk for competition and interference with endogenous actin-binding interactors, more so the higher the levels of the reporters. The regulation of their expression is thus a key advantage, as shown by the mitigation of the effects we observed on *Drosophila* flight capacity and flight muscle integrity by Af7 when lowering its levels. In addition to the controls and functional readouts we provide in our study for the specific models we use, additional or different readouts related to a given question under study might be important to consider.

As already reported in the literature and also shown in our study, different actin-binding probes have different affinities for F-actin and for distinct actin filament assemblies depending on the density and geometry of the latter, as well as on the presence of competing actin-binding proteins (tropomyosins, myosins, formins). The availability of constrained GFP fusions to five widely used F-actin localization reporters, including G-actin itself, increases the chances that one is able to label and consequently measure the organization of the F-actin pools of interest. Constrained Cherry fusions for F-actin organization measurements provide additional experimental flexibility, notably for two-color imaging with GFP fusions to any protein of interest while measuring F-actin organization. Our red FP fusions are also compatible with the standard CFP/YFP-like donor/acceptor pairs used for FRET-based force measurements (Gayrard and Borghi, 2016), enabling a direct correlation between F-actin organization and mechanical properties.

Finally, our results from the engineering of constrained FP fusions to ABDs informed us at different levels with respect to the mechanisms of FP immobilization. First, shortening of the N- and C-termini in terminal fusions seems to be the most efficient way to constrain sfGFP and sfCherry2 mobility without compromising their fluorescence. Second, shortening of the FP termini might compromise F-actin binding if the fused ABD terminus is shortened at the same time; our results suggest that two residues between the end of the FP barrel and the ABD are typically needed so as not compromise F-actin binding. Third, circularly permuted sfGFP fusions can constrain GFP mobility, but not nearly as efficiently as terminal fusions because cpGFP fusions tolerate poorly the shortening of the GFP termini or/and the shortening of the ABD: fluorescence or/and F-actin binding are rapidly compromised upon such shortening because of the flexibility required to connect the original termini of cpGFP. Fourth, FP immobilization ultimately depends on the flexibility of the terminus of the fused ABD or of the secondary structure of the ABD (e.g. loop) to which it is fused: if the ABD terminus or the secondary structure element is inherently flexible, even the shortest FP will remain mobile. Even though our results relate primarily to sfGFP and sfCherry2 fusions to ABDs, the above-mentioned conclusions can be rationally applied to other FPs, and fusions to proteins other than actin-binding ones, to generate new tools for measuring any protein organization by polarimetry. Structural and cell biology approaches employing FP-based sensors to probe protein proximity, protein-protein interactions and mechanical forces, notably the ones using FRET, are also likely to benefit from using constrained FP fusions.

ACKNOWLEDGMENTS

We thank Louwrens van Dellen (I. Fresnel, Marseille, France) for the development of the image acquisition software. This research has received funding from the French National Research Agency (ANR) grants Equipex+ IDEC (France 2030 investment plan ANR-21-ESRE-0002), 3DPolariSR (ANR-20-CE42-0003), SEPTIMORF (ANR-17-CE13-0014), SEPTISS (ANR-22-CE13-0039), and from the France-BioImaging infrastructure (ANR-10-INBS-04). This research has further received funding from the Excellence Initiative of Aix-Marseille University – A*Midex (A*Midex grant NEUROPOL), a French

"Investissements d'Avenir" program, the Fondation pour la Recherche Médicale (FRM grant ING20150531962) and SATT Sud-Est.

AUTHOR CONTRIBUTIONS

C.S.M. : Methodology, Investigation, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing; F.I. : Investigation; S.K.S. : Investigation, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing; T.C.P. : Investigation, Validation, Visualization, Formal analysis; C.S. : Investigation, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing; M.R.-L. : Investigation; C.N.P. : Investigation; S.O. : Investigation; M.G. : Investigation; A.L. : Investigation; S.R.B. : Investigation; L.R. : Investigation; F.A. : Investigation; C.V.R. : Investigation; F.S. : Writing – review & editing, Supervision, Resources; F.R. : Writing – review & editing, Supervision, Resources; A.W. : Writing – review & editing, Supervision, Resources; L.L. : Investigation, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing; J.-D.P. : Investigation, Visualization, Writing – original draft, Writing – review & editing; S.C. : Investigation, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing, Resources; S.A.R. : Investigation, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing, Supervision, Resources ; C.C. : Software; S.B. : Conceptualization, Methodology, Investigation, Software, Writing – original draft, Writing – review & editing, Supervision, Resources, Project administration, Funding acquisition; M.M. : Conceptualization, Methodology, Investigation, Software, Validation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing, Supervision, Resources, Project administration, Funding acquisition

DECLARATION OF INTERESTS

The authors declare no competing interests.

STAR methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources, reagents, and software should be directed to and will be fulfilled by the lead contact, Manos Mavrakis (manos.mavrakis@fresnel.fr).

Materials availability

All plasmids and strains generated in this study are available upon request.

Data and code availability

The datasets supporting the current study have not been deposited in a public repository but are available from the lead contact upon request. The codes and softwares developed and used in this study are open source and available on GitHub under a BSD license; the software identifiers are listed in the Key Resources Table and the links are provided in the respective method details sections.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell lines and cell culture

U2OS osteosarcoma cells were used for the screening of actin organization reporters with respect to their localization and usability for polarimetry measurements. U2OS cells were from ATCC (HTB-96). Cells were maintained in McCoy's medium (Thermo Fisher, Cat#16600082) supplemented with 10% fetal bovine serum (Dominique Dutscher, Cat# S181H), 100 U/mL penicillin and 100 µg/mL streptomycin antibiotics (Sigma-Aldrich, Cat#P4333) in a humidified atmosphere at 37°C containing 5% CO_2. Transfections were performed 16 h prior to live imaging using FuGENE HD Transfection Reagent (Promega, Cat#E2311), following the manufacturer's instructions. To obtain single cells for imaging, 25×10^3 U2OS cells were typically seeded into a 24-well glass bottom plate (Cellvis, Cat#P24-1.5H-N) a day prior to the day of transfection, for allowing an optimal number of cells to attach and spread. A total of 0.2 µg of DNA and a 4:1 ratio of FuGENE HD (µL) : DNA (µg) were used per reaction. To minimize overexpression, the amount of DNA for pCMV plasmids was reduced to 50 ng, leading to a 16:1 ratio of FuGENE HD (µL) : DNA (µg). Cells were imaged 16h post-transfection.

HeLa cells were used for the characterization of intramolecular GFP (iGFP)-beta and -gamma actin fusions. Stable HeLa cell lines with regulated expression of either iGFP-beta actin or iGFP-gamma actin were generated with the Flp-In system (Life Technologies) using HeLa cells that contained a single FRT site according to the manufacturer's instructions (Renshaw et al., 2014). The resulting cell lines were cultured in Dulbecco's Modified Eagle Medium (Wisent Bio Products, Cat#319-030 CL) supplemented with 10% fetal bovine serum (Wisent Bio Products, Cat#080-650), 1% penicillin/streptomycin (Invitrogen Cat#15140122), 5 µg/mL blasticidin (BioShop Canada Inc, Cat#BLA477), and 2 µg/mL puromycin (BioShop Canada Inc, Cat#PUR333). Expression of GFP fusion proteins were induced by addition of 0.25 µg/mL doxycycline to the growth media for 24 h before either fixation or harvesting. Cells were maintained in a Forma Series II incubator (Thermo Scientific) at 37°C in a 5% CO_2 atmosphere.

Fission yeast strains, maintenance and genetics
Standard *Schizosaccharomyces pombe* media and genetic manipulations were used (Moreno et al., 1991). All strains used in the study were isogenic to wild-type 972 and their genotypes are described in Table S3. The generation of transgenic strains is described in the method details section. Strains from genetic crosses were selected by random spore germination and replica in plates with appropriate supplements or drugs. Transformations were performed using the lithium acetate-DMSO method as described in (Bahler et al., 1998). Drop assays (Figures 5C,D) were performed by serial dilutions of 1/4 from a starting sample of optical density of 1.0 of the indicated strains, which were plated on YE5S medium supplemented with the corresponding drug and incubated for 3 days at 28°C unless stated differently.

***C. elegans* strains, maintenance and genetics**
Bristol strain N2 was used as the wild-type strain. *C. elegans* strains used in this study and their genotypes are listed in the Key Resources Table and were reared using standard methods (Brenner, 1974). The generation of transgenic strains is described in the method details section. The strains were grown at 20°C and fed Escherichia coli OP50. The EG6699 (unc-119(ed3) III) strain, used as the host strain of FBR193, FBR195

994 and FBR196 strains generated in this study, was grown at 15°C and fed E. coli HB101
995 before injection (Hochbaum et al., 2010).
996

Drosophila strains, maintenance and genetics

Drosophila melanogaster strains used in this study and their genotypes are listed in the
Key Resources Table. The generation of the transgenic strains UASp–L22, UASp–L45,
UASp–U20, and UASp–Af7 is described in the method details section. Fly stocks were
grown and maintained at 25°C on semi-defined medium
(https://bdsc.indiana.edu/information/recipes/germanfood.html). The GAL4/UASp
expression system was used to drive expression in the *Drosophila* embryo using GAL4
expressed under the control of the maternal alphaTub67C promoter (mat–α-tub–GAL4)
(BDSC_80361). Crosses were maintained at 25°C. The GAL4/UAS expression system
was used to drive the expression of actin organization reporters in *Drosophila* indirect
flight muscles. Mef2-GAL4 (BDSC_27390) or tub-GAL80ts ; Mef2-GAL4 (BDSC_7108,
BDSC_27390) females were crossed to males of the following genotypes: w[1118]
(BDSC_3605), UAS–GFP-GMA (BDSC_31776), UAS–Lifeact-EGFP (BDSC_35544),
UASp–L22, UASp–L45, UASp–U20, and UASp–Af7. Crosses with the Mef2-GAL4 driver
were grown at 25°C. Crosses with the tub-GAL80ts ; Mef2-GAL4 driver were grown at
18°C (no GAL4 activity): a few days after eclosion, adults were transferred to a permissive
temperature of 31°C for 5 days prior to flight tests or fixation/staining or to a permissive
temperature of 25°C for 5 days prior to live polarimetry. The GAL4/UAS expression
system was used to drive the expression of actin organization reporters in the *Drosophila*
wing. nub-GAL4 (BDSC_86108) females were collected within 2 days and combined to
create a uniform population. Eight of these females were crossed to three males of the
following genotypes: UAS–dInR-DN (BDSC_8253), sqh–Lifeact-EGFP, UASp–L22,
UASp–L45, UASp–U20, and UASp–Af7. The rearing temperature was maintained at
25°C, and the tubes were flipped daily.
1021

METHOD DETAILS

1023

Generation of mammalian expression plasmids for screening actin organization reporters

All constructs were designed in silico with SnapGene (Dotmatics) and are listed in Table
S1. To drive expression of the constructs in mammalian cells, we used the immediate
early enhancer and promoter of human cytomegalovirus (CMV promoter, 508 base pairs),
as well as a truncated version (CMV$_{trunc}$, 54 base pairs) for low-level expression; the latter
was originally generated for reduced expression of EGFP-beta-actin (Watanabe and
Mitchison, 2002). Addgene plasmids #31502 and #54759 were used to obtain the
CMV$_{trunc}$ and CMV backbones, respectively. All constructs screened for actin organization
reporters were driven by CMV$_{trunc}$ apart from the ones labeled with an asterisk in Figures
2A, 3A, 3D, 3G, 4A, S2A, S3A, S3F and the iGFP-beta- and -gamma constructs (Figure
S4E-O). Fluorescent protein constructs screened for fluorescence in the truncation
screens (Figures 1G,H and S4A,B) were driven by CMV. Lifeact-mEGFP and EGFP-beta-
actin cDNAs were a gift from Yannick Hamon (CIML, France). FtrN9-52-mEGFP and
EGFP-Affimer6 cDNAs were a gift from John Hammer (NIH/NHLBI, USA) and Michelle
Peckham (University of Leeds, UK), respectively. Beta- and gamma-actin cDNAs were a

gift from Boris Hinz (University of Toronto, Canada). Synthetic genes for sfGFP, msfGFP, human beta-actin and GFP11ΔC8-FtrN10-52-GFP1-10 were from Eurofins Genomics (Germany). Fluorescent protein fusions were generated using monomeric (A206K) EGFP (mEGFP) (Zacharias et al., 2002; Zhang et al., 1996), monomeric (V206K) superfolder GFP (msfGFP) (Costantini et al., 2012; Cranfill et al., 2016; Pedelacq et al., 2006; Zacharias et al., 2002), monomeric Apple (mApple) (Cranfill et al., 2016; Shaner et al., 2008) and superfolder Cherry2 (sfCherry2) (Feng et al., 2017). To optimize the intramolecular self-association of β11 with the GFP1-10 moiety in our circular permutants, we have used the sfGFP-evolved sequences, GFP1-10 OPT and GFP11 M3, which have been optimally engineered to work in bipartite split-GFP complementation assays (Cabantous et al., 2005). All constructs were generated with seamless cloning (In-Fusion HD Cloning Plus Kit from Takara Bio, Cat. # 638910) using NheI/BamHI (or AflII/BamHI for iGFP constructs) linearized plasmid backbones and the oligonucleotide primer sequences listed in Table S2. Primers were Cloning Oligo (<60 bp) or EXTREmer (>60 bp) synthesis and purification quality from Eurofins Genomics (Germany). Restriction enzymes were FastDigest enzymes from Thermo Scientific. All plasmids were verified by sequencing (Eurofins Genomics, Germany) after each cloning step.

We note the following with respect to residue numbering of EGFP/sfGFP in our study: although the valine following the initiating methionine is typically numbered 1a to maintain correspondence between EGFP/sfGFP and wild-type GFP numbering (Tsien, 1998), we number this valine as 2 in this study to facilitate the naming of N-terminal truncations in the screen. As a result, the last residue of EGFP/sfGFP is 239, an N-terminally truncated msfGFP mutant missing the first six residues (ΔN6) starts with ELFTGV..., and a C-terminally truncated msfGFP mutant missing the last nine residues (ΔC9) ends with ...AAGI.

Screening of actin organization reporters in live U2OS cells
Confocal fluorescence microscopy and image processing
For live cell imaging, right before microscopy and due to the absence of CO_2 control on our microscope setup, the culture medium was exchanged by Leibovitz medium (Thermo Fisher, Cat#21083027) supplemented with 10% fetal bovine serum and antibiotics. Cells were kept at 37°C in a heating chamber (OkoLab, Cat#H301-TUNIT-BL). Fluorescence images were acquired using a custom spinning disk microscope (detailed in the Polarimetry methods section) with a Nikon Plan Apo ×100/1.45 NA oil immersion objective lens, 488- 561- and 641-nm laser lines and an iXon Ultra 888 EMCCD camera. Z-stacks were acquired with a Δz interval of 0.5 µm. Exposure times were in the range of 0.5–2.0 s depending on the exact condition.

Images were processed with the open-source image processing software ImageJ/Fiji. The images displayed in Figure 1H and Figure S4B are maximum intensity projections of two consecutive z-planes displayed with the same intensity range to allow for intensity comparison between the FP truncation mutants. All the other shown images are maximum intensity projections of two consecutive z-planes contrasted manually in order to optimize the image display.

Polarimetry measurements in live U2OS cells

Polarimetry stacks using 36 polarization angles were recorded in focal planes containing peripheral stress fibers with a typical exposure time of 0.1-0.2 s per polarized image (see details for the optical setup and signal processing in the Polarimetry methods section). Typically, a minimum of five fields of views containing single cells was acquired per experimental condition. Polarimetry stacks were systematically registered using the StackReg plugin for ImageJ to correct for x and y axis drift during acquisition. To select peripheral SF-associated pixels for analysis, binary masks of SF segments were generated using the open source tool FilamentSensor 0.2.3 (Eltzner et al., 2020), freely available at http://www.filament-sensor.de/. A pre-processing tab in the FilamentSensor software requires adjustment in the contrast and removal of standalone pixels, followed by the optional application of filters. A standard and optimized preprocessing was using Laplace filter, 8 neighbors and factor 4; Gaussian filter, sigma 1; Cross correlation filter, size 10 and zero 30%; and a directed Gaussian filter, sigma 8. The binarization method chosen was by area, and filament detection parameters were typically chosen as follows: minimum mean value 25, sigma 2, minimum standard deviation 5, minimum filament length 20, minimum angle difference 20, tolerance 5%. The final selection was done manually, and only identified filaments that were colocalizing with peripheral SFs were used to generate the binary masks for selecting the pixels for polarimetry analysis.

Polarimetry data were analyzed according to the framework defined by (Kress et al., 2013) to obtain the ρ and ψ angle per image pixel. Analysis and data representation, including color-coded stick representations of the measured angles per pixel were done with the Polarimetry software which is a Matlab App Designer standalone application. The source code is available at https://github.com/cchandre/Polarimetry.git, and the desktop app can be freely obtained at https://www.fresnel.fr/polarimetry under a BSD license. The Matlab-based Polarimetry software is the precursor of the Python-based app PyPOLAR used for the analysis of the yeast, *Drosophila* and *C. elegans* polarimetry data (see respective methods sections). The distributions of the ψ angles are represented in box plots with overlaid data points. Each data point represents a single actin fiber. If more than one fibers were identified in the same field of view, measurements for each fiber are shown as distinct datapoints, which results in more than one measurements per field of view. On each box, the central mark indicates the median, and the left and right edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers. Boxplots were generated with custom-written Matlab code (see *Data post-processing with Matlab* in https://www.fresnel.fr/polarimetry). The number of measurements for each construct, the respective median ψ values and the statistical test used in GraphPad Prism to evaluate differences are mentioned in the respective legend.

Polarimetry measurements in fixed U2OS cells

U2OS cells were fixed for 20 min with 4% paraformaldehyde (Electron Microscopy Sciences, Cat#15714) in 37°C-prewarmed cytoskeleton buffer (10 mM MES pH 6.1, 150 mM NaCl, 5 mM EGTA, 5 mM $MgCl_2$, 5 mM glucose), followed by 2 × 5 min wash steps in phosphate-buffered saline (PBS) solution. Cells were subsequently incubated with 0.165 µM Alexa Fluor 488-phalloidin (Thermo Fisher Scientific, Cat# A12379) or 0.165 µM SiR-actin (Spirochrome, Cat#SC006) in PBS containing 0.1% saponin and 1% IgG-free/protease free bovine serum albumin (Jackson ImmunoResearch, Cat#001-000-161)

for 1 h at RT. Coverslips were mounted with 15 µL Fluoromount (F4680; Sigma-Aldrich) for image acquisition.

Polarimetry stacks using 18 polarization angles were recorded in focal planes containing ventral and peripheral stress fibers with a typical exposure time of 0.1-0.2 s per polarized image. The pixels of the stress fibers for analysis were selected by a combination of intensity thresholding and manual selection of the region to analyze. Analysis and data representation, including color-coded stick representations of the measured angles per pixel were done with Polarimetry or PyPOLAR softwares. The distributions of the ψ angles are represented in box plots with overlaid data points as described above for measurements in live cells. The number of measurements for each dye and the respective median ψ values are mentioned in the respective legend.

Flow cytometry analysis of truncation mutants of fluorescent protein variants

U2OS cells were seeded at a density of 5×10^4 cells per well in 24-well plates. After 24 h, cells were transfected with 0.5 µg of corresponding plasmid DNAs in 50 µL of Jet Prime Buffer mixed with 1 µL of jetPRIME reagent (Polyplus, Cat#101000046). Twenty-four hours after transfection, cells were trypsinized, resuspended in PBS supplemented with 2% fetal bovine serum, then transferred in 96-well conical bottom plates. Cells were fixed with 3.7% paraformaldehyde for 10 min and resuspended in 1% BSA, PBS buffer. Green (mEFP, msfGFP) and red (mApple, sfCherry2) fluorescence were collected on 10,000 cells using a MACSQuant VYB flow cytometer (Miltenyi Biotec). The fluorescence threshold was defined based on the background fluorescence of untransfected U2OS cells. The percentage of green and red positive fluorescence was analyzed with FlowJo® software (BD Biosciences). The fluorescence of full-length constructs was normalized to 100% for each independent experiment. Bar graphs of the measured fluorescence were prepared with GraphPad Prism. The mean values and the statistical test used to evaluate differences are indicated in the respective legend.

Characterization of intramolecular GFP (iGFP)-beta and -gamma actin fusions

siRNA treatment and rescue experiments

HeLa cells grown to 40% confluency in 6-well plates were transfected with 100 pmol double-stranded siRNA targeting *ACTB* (sequence #1: AAAUAUGAGAUGCGUUGUUACAGGA; sequence #2: UCCUGUAACAACGCAUCUCAUAUUUGG) or *ACTG1* (sequence #1: GCAUGGGUUAAUUGAGAAUAGAAAT; sequence #2: AUUUCUAUUCUCAAUUAACCCAUGCAG) using Lipofectamine 2000 (Invitrogen, Cat#11668019) following manufacturer's instructions. For rescue experiments, siRNA-resistant transgenes were expressed 24 h after siRNA transfection and either fixed or harvested 48 h post-siRNA transfection. All siRNAs were obtained from IDT (Integrated DNA Technologies).

Cell harvesting, SDS-PAGE, and western blotting

HeLa cells grown to near confluency in 6-well dishes were harvested by scraping in 100 µL RIPA lysis buffer (50 mM Tris pH 7.4, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 1 mM PMSF, 2 µg/mL aprotinin, 2 µg/mL leupeptin) on ice. Lysates were spun at 20,800 *g* for 20 min, after which supernatants were collected, mixed

with SDS-PAGE sample buffer, and boiled at 95°C for 5 min. Samples were run on a 10% polyacrylamide gel and subsequently transferred onto nitrocellulose membranes (Bio-Rad). Membranes were blocked for 1 h in 5% skim milk powder in TBST (TBS with 0.0025% Tween-20), before incubation with either mouse anti-gamma actin antibody (Bio-Rad, Cat#MCA5776GA; dilution 1:200), mouse anti-beta actin antibody (Bio-Rad, Cat#MCA5775GA; dilution 1:200), or mouse anti-acetylated alpha tubulin antibody (Santa Cruz, Cat#sc-23950; dilution 1:2000) for 1 h. Following secondary antibody incubation, membranes were developed with chemiluminescent solutions (Thermo) for 1-2 min at room temperature and visualized using a Bio-Rad MP Imager (Bio-Rad).

Multinucleation and mitotic staging assays
Stable iGFP-actin HeLa cells were seeded onto glass coverslips in 6-well dishes. At roughly 40% confluency, cells were transfected with 100 pmol of either control, beta, or gamma actin-targeting siRNA (Integrated DNA Technologies) with Lipofectamine 2000 (Invitrogen) following manufacturer's instructions. Where indicated, 24 h post-transfection cells were induced to express either iGFP-beta or -gamma actin by the addition of 0.25 µg/mL doxycycline. Cells were fixed 48 h post-transfection as described below. For multinucleation assays, cells were classified as either mono- or multi-nucleate by manually scoring the number of nuclei, as reported by Hoechst staining, within each cell boundary, as reported by acetylated alpha-tubulin staining, omitting cells fixed mid-division. For mitotic staging experiments, cells were manually classified as mitotic if the following features were observed: i) condensed chromosomes by Hoechst staining and/or b) an intercellular bridge as reported by acetylated alpha-tubulin staining. Early mitotic cells were further classified into either 'prophase' or 'metaphase' populations based on the organization of their condensed chromosomes, with 'metaphase' cells exhibiting sharp alignment with the metaphase plate, and 'prophase' cells exhibiting chromosomal rosettes or otherwise unaligned chromosomes. Data was entered into GraphPad Prism to generate bar graphs and perform statistical tests; the number of cells scored for each condition and details of statistical tests performed are described in the respective figure legends.

Immunofluorescence
To visualize iGFP-tagged actins, HeLa cells were fixed with 3.7% paraformaldehyde in PHEM buffer (60 mM PIPES pH 7.0, 25 mM HEPES, 10 mM EGTA, 4 mM MgSO$_4$·7 H$_2$O) for 10 min at room temperature and permeabilized by 0.2% Triton-X-100 in PBS for 10 min. Coverslips were blocked for 1 h in 3% BSA in PBS. Where indicated, iGFP-expressing cells were probed with anti-vinculin antibody (Sigma-Aldrich, Cat#V9131, 1:100) for 16 h at 4°C. Coverslips were subsequently incubated with either Alexa 594 or Alexa 647 conjugated goat-anti mouse secondary antibody (Invitrogen, 1:400) for 1 h. For experiments visualizing endogenous beta actin and either iGFP gamma actin or vinculin, cells were fixed with 3.0% paraformaldehyde in PHEM buffer for 30 min at 37°C, followed by a second fixation for 5 min in -20°C methanol. Coverslips were blocked with 3% BSA in PBS overnight at 4°C, before incubation for 1 h with mouse anti-beta actin antibody (Bio-Rad, Cat#MCA5775GA; dilution 1:600), either with or without anti-vinculin antibody (Sigma-Aldrich, Cat#V9131, 1:100), diluted in 1% BSA in PBS. Coverslips were

subsequently incubated with either Alexa 594 or Alexa 647 conjugated goat-anti mouse secondary antibody (Invitrogen, 1:400) for 1 h.

For multinucleation and mitotic staging experiments, cells were fixed with -20°C methanol for 10 min, blocked with 3% BSA in PBS, and subsequently stained with mouse anti-acetylated alpha tubulin antibody (Santa Cruz, sc-23950, 1:400) for 1 h. Coverslips were then incubated with Alexa 594 conjugated goat-anti mouse secondary antibody (Invitrogen, 1:400) for 1 h.

Prior to mounting with Mowiol (Polyvinyl alcohol 4-88, Fluka), coverslips were incubated in 1 µg/mL Hoechst 33258 (Sigma) for 10 min and rinsed in ddH$_2$O. Cells were visualized with either a PerkinElmer UltraView spinning disk confocal scanner mounted on a Nikon TE2000-E with a 60x/1.4 NA oil-immersion objective lens and 1.515 immersion oil at room temperature or a Leica SP8 scanning confocal microscope with a 63x/1.4 NA oil-immersion objective lens and Leica Type F immersion oil. Images were acquired using METAMORPH software (v.7.7.0.0; Molecular Devices) driving an electron multiplying charge-coupled device (CCD) camera (ImagEM, Hammamatsu) or LAS-X software (v.1.4.4; Leica) driving HyD detectors. Z sections (0.2 µm apart) were acquired to produce a stack that was then imported into AutoQuant X3 (Media Cybernetics) for 3D deconvolution (5 iterations). Single Z-slices were generated in ImageJ (v2.1.0). Images were overlaid in Adobe Photoshop (v23.0.2) involving adjustments to brightness and contrast.

Characterization of fission yeast strains expressing actin organization reporters
Generation of fission yeast strains

msfGFP-tagged actin organization reporters were expressed from the fission yeast *leu1+* locus under the control of the *cdc42+* promoter using the integrative vector pJK148 (Keeney and Boeke, 1994). Briefly, around 500 bp from the *cdc42+* promoter were amplified by PCR and cloned into the pJK148 vector using the *Sac*I and *Xba*I sites, creating pSRP12. The *adh1+* terminator was amplified by PCR and cloned into pSRP12 using the *BamH*I and *Sal*I sites, creating pSRP14. Finally, the fragments coding for each of the actin reporters fused to msfGFP were obtained by digestion with *Nhe*I and *BamH*I from the respective mammalian expression plasmids and cloned into pSRP14, between the *cdc42+* promoter and the *adh1+* terminator, creating the plasmids pSRP16 to pSRP23, respectively (see Table S1). All oligos used are listed in Table S2. Plasmids were linearized by *Nru*I digestion, before transformation of a wild-type strain. Genetic crosses were performed to combine the actin reporter-expressing strains to strains expressing the proper marker to check cytokinesis dynamics, microtubule organization or to the profilin mutant thermosensitive strain, *cdc3-319* (see Table S3).

Microscopy and image analysis

For imaging, fission yeast cells were grown at 28°C (32°C for cells shown in Figure S5A) in YE5S medium to exponential growth. For time-lapse imaging, 300 µL of early log-phase cell cultures were placed in a well from a µ-Slide 8 well (Ibidi, Cat#80821) previously coated with 10 µL of 500 µg/mL soybean lectin (Sigma-Aldrich, Cat#L1395). Cells were left for 1 min to attach to the bottom of the well and culture media was removed carefully. Then, cells were washed three times with the same media and finally 300 µL of fresh

media were added (Cortes et al., 2012), before incubation in the microscope chamber at
the same temperature at which cells had been cultured.
Time-lapse images shown in Figure 5A are maximum intensity projections obtained from
z-stacks of 7 slices at 1 µm interval every 2 minutes, acquired using an Olympus IX81
spinning disk confocal microscope with Roper technology controlled by Metamorph 7.7
software (Molecular Devices), equipped with a 100X/1.40 Plan Apo oil lens, a Yokogawa
confocal unit, an EVOLVE CCD camera (Photometrics) and a laser bench with 491-561
nm diode. Exposure time for green or red channels was 0.5 s.
Time-lapse images shown in Figure S5A are maximum intensity projections obtained from
z-stacks of 7 slices at 0.3 µm interval every 6 minutes, acquired using a Dragonfly 200
Nikon Ti2-E spinning disk confocal microscope controlled by Fusion software (Andor),
equipped with a 100X/1.45 Plan Apo oil lens, an Andor confocal unit, an sCMOS Sona
4.2B-11 camera (Andor) and a laser bench with 405-561 nm diode (Andor). Exposure
time was 0.3 s for the green channel and 0.2 s for the red channel. Microscopy images
shown in Figure S5B are maximum intensity projections obtained from z-stacks of 7 slices
at 0.3 µm interval, acquired using the same microscope setup from Nikon. Exposure time
was 0.35 s. For the sake of comparison, images in Figure S5A and S5B are displayed
using the same intensity range with Metamorph 7.7.
Quantification of the time for acto-myosin ring assembly, maturation and constriction was
performed by analyzing the time between the initial recruitment of myosin cortical nodes
and their compaction into a tight ring, the time until the ring starts to constrict and the time
until the myosin signal disappears after final constriction, respectively. Scatter dot plots
of the measured times were prepared with GraphPad Prism. The number of cells used
for each strain, the mean measured times and the statistical test used to evaluate
differences, the latter performed with GraphPad Prism, are indicated in the respective
legend. Actin patch and actin cable number per cell were quantified from maximum
intensity projections obtained from z-stacks of 7 slices at 1 µm from cells in G2 phase
(around 10 µm long). Scatter dot plots of the measured actin cables and patches were
prepared with GraphPad Prism. The number of cells used for each strain and the mean
numbers of cables and patches are indicated in the respective legend.
1297
Polarimetry measurements in the cytokinetic ring of live fission yeast
For live polarimetry measurements, strains co-expressing Af1, Af7, L1 or L22 and an
mCherry-tubulin marker (strains SR3.51, SR3.54, SR3.57 and SR3.58 in Table S3) were
incubated at 25°C in YE5S medium. 1 mL of exponentially growing cells were harvested
by centrifugation for 60 s at 800 g, most of the supernatant was discarded and 1 mL of
the cells was deposited onto a 2% YE5S agar pad at the center of a polydimethylsiloxane
slide chamber prepared as described in (Costa et al., 2013).
Three large field-of-view images (66 x 66 µm) typically containing 5-10 dividing cells per
image, were collected for each strain. Before each polarimetry measurement, a two-color
z stack was acquired ($\Delta z = 1.0$ µm) to image both GFP fusions and microtubules; the
distribution of the latter was used in addition to the morphology of the actomyosin ring to
confirm that cells were undergoing cytokinesis. A polarimetry stack using 18 polarization
angles was then recorded for each position within a z stack ($\Delta z = 1.0$ µm) for the GFP
channel, and thus allowed to obtain polarimetry images throughout the cytokinetic rings,
containing both tangential-most views with the ring parallel to the xy plane, and more

equatorial views showing cross-sections that appear as spots on either side of the ring. An exposure time of 0.5 s was used per polarized image. To minimize bias in the measured orientations due to the contribution of off-plane orientations we focused on the tangential-most views for the analysis (Figure 5E-G). One tangential view of the ring was analyzed per cell; in a few cells where both tangential views were present in the respective z planes, both were analyzed. Equatorial views were used for measuring the diameter of the constricting rings (Figure S5G).

Polarimetry stack images were first processed with the open-source image processing software ImageJ/Fiji. Images within each polarimetry stack were registered using the StackReg plugin to correct for drift during the acquisition. The z planes containing the tangential-most views of the ring were identified for each cell. The pixels of the cytokinetic ring for analysis were selected by a combination of intensity thresholding and manual selection of the region to analyze. Each region of interest contained typically 40-80 analyzed pixels i.e. 40–80 color-coded sticks in the tangential-most view of the cytokinetic ring per cell (Figure 5E-F). Analysis and data representation, including color-coded stick representations of the measured angles per pixel and polar histograms were done with PyPOLAR. The source code is available at https://github.com/cchandre/Polarimetry.git, and the desktop app can be freely obtained at https://www.fresnel.fr/polarimetry under a BSD license. GraphPad Prism was used to generate scatter plots of the quantified ψ angle distributions per strain; the number of cells measured for each strain, the respective median values and the statistical test used to evaluate differences are mentioned in the respective legend. Considering that Af7 and L22 dipoles are parallel to actin filaments, in order to assess the extent to which the measured actin filament orientations were more parallel or more perpendicular with respect to the ring axis, the ring axis angle in each cell was used as the reference angle in the "reference angle" tool in PyPOLAR to normalize the angle distributions from 0°–180° to 0°–90° and generate 0°–90° polar histograms, with 0° and 90° defining orientations parallel and perpendicular to the ring axis, respectively (Figure S5E-F).

Characterization of *C. elegans* strains expressing actin organization reporters
Plasmid construction for generation of transgenic animals

Sequences encoding actin organization reporters were codon optimised for optimal expression in the worms using the *C. elegans* codon adapter web tool (Redemann et al., 2011) and synthetized by GENEWIZ. Plasmids pFBR101, pFBR102 and pFBR105 (see Table S1) were constructed in two steps from pML36 (kind gift from Michel Labouesse lab), which contained a pCFJ151 backbone (ttTi5605 insertion homology arms) with a dpy-7 promoter for epidermal cell expression and a unc-54 3'UTR (universal 3'UTR for optimal expression). Briefly, the plasmid pML36 was opened and amplified by PCR using custom made oligos (Sigma-Aldrich). The sequences encoding organization reporters were also amplified by PCR and joined using overlapping ends into opened pML36 plasmid using the NEBuilder HiFi DNA Assembly Cloning kit (New England Biolabs, Cat#E5520S). All PCR reactions were carried out by Phusion High-Fidelity DNA Polymerase (ThermoFisher Scientific, Cat#F531L). Primers were custom made by Sigma-Aldrich. The sequences of all oligos are listed in Table S2. The final plasmids were verified by DNA sequencing (Eurofins Genomics, Germany).

Transgenic worm construction by MosSCI method

Worm MosSCI transgenesis was performed by direct microinjection as described in (Frokjaer-Jensen et al., 2008). Briefly, the injection mix was injected in the arm of both gonads in the young hermaphrodite animal of EG6699 strain. The injection mix contained a cocktail of pJL43.1 (50 ng/mL), pCJF90 (2.5 ng/mL), pCFJ104 (5 ng/mL), and an expression clone (50 ng/mL) in DNase/RNase-free water. All plasmids used for injection were purified by HiSpeed Plasmid Midi kit (Qiagen, Cat#12643). After injection, transgene insertion screening was performed as described at http://www.wormbuilder.org. Transgenic animals were verified by PCR genotyping and DNA sequencing.

Worm embryonic growth and lethality tests

A few young hermaphrodite animals were picked and fed on freshly seeded Escherichia coli OP50 for 2h. After 2 h, all animals were transferred on fresh OP50 plates and laid eggs were counted. This process was redone until sufficient number (>1000) of embryos were achieved and counted. After 12-16 h, all previously scored embryo-containing plates were recounted for unhatched embryos (dead eggs) and hatched larvae. For this experiment, strains were grown at 20°C. The embryonic lethality for each strain, scored as the percentage of unhatched embryos, is shown in Figure 6A.

Embryonic growth rate was measured by imaging embryos by differential interference contrast (DIC) microscopy on a Leica DM6000 microscope until they hatch as larvae. Briefly, embryos were collected by dissecting gravid hermaphrodites in M9 medium and mounted on a 5% agarose pad for imaging. Z-stack images were acquired with 40-50 planes per embryo and a Δz interval of 1 μm, and with a 10 min interval for 12-14 h at 20°C. Embryonic length was measured manually with the segmented line tool in the ImageJ/Fiji software and growth curves plotted with Microsoft Excel software (Figure 6B). The number of embryos measured for each strain is indicated in the respective legend.

Polarimetry measurements in live C. elegans embryos

More than >30 young gravid hermaphrodite animals were picked and fed on freshly seeded OP50 E. coli overnight. Next day, mixed stage embryos were picked and mounted on a 5% agarose pad in M9 medium for imaging. Temporary hypoxic conditions were created by adding OP50 E. coli, preventing embryonic muscle activity that usually starts around 1.7-fold. Embryonic stages were evaluated by brightfield microscopy. Length measurements were subsequently performed using ImageJ/Fiji. Polarimetry stacks using 18 polarization angles were recorded in the epidermis of 1.5-fold, 1.5-2-fold and >2-fold stage embryos. The pixels containing dorsal and ventral epidermal cells (DV cells) and seam cells were selected by a combination of intensity thresholding and manual selection of the region to analyze. Analysis and data representation, including color-coded stick representations of the measured angles per pixel and histograms were done with PyPOLAR. GraphPad Prism was used to generate scatter plots of the quantified ψ angle distributions per strain and per developmental stage; the number of embryos for each strain and for each stage and the respective median ψ values are mentioned in the respective legend. To assess how the measured actin filament orientations in DV and seam cells distribute with respect to the DV/seam boundary for each developmental stage, the ρ angle distributions were normalized with respect to the DV/seam boundary from 0°–180° to 0°–90° to generate 0°–90° polar histograms. The DV/seam boundary for

each embryo was drawn manually with the freehand line selection tool or the elliptical
selection tool in Fiji and converted to a mask which was then used with the "edge
detection" tool in PyPOLAR to define the boundary as the reference for the normalization
of the angles. Considering that Af7 and L22 dipoles are parallel to actin filaments and that
L45 dipoles are perpendicular to actin filaments, the more perpendicular mean actin
filament orientations are to the boundary, the closer the angle values are to 90° (for Af7
and L22) or to 0° (for L45) (Figure 6F,H,J and Figure S6E,G).
1412
Generation of *Drosophila* expressing selected actin organization reporters
Drosophila transgenics
Selected actin organization reporters were subcloned into pUASp plasmids for generating
Drosophila transgenics. The respective mammalian expression plasmids were used as
templates to subclone the reporters into *KpnI*/*BamH*I linearized UASp vectors using
seamless cloning (In-Fusion HD Cloning Plus Kit, Takara Bio, Cat#638910). All primers
were Cloning Oligo synthesis and purification quality from Eurofins Genomics and are
listed in Table S2. Restriction enzymes were FastDigest enzymes from Thermo Fisher
Scientific. All plasmids were verified by sequencing (Eurofins Genomics) after each
cloning step. Midipreps of each UASp construct DNA were sent to BestGene Inc.
(California, USA) for injections into *D. melanogaster* w1118 embryos and generation of
the transformants UASp–L22, UASp–L45, UASp–U20, and UASp–Af7 (see Key
Resources Table and Table S1).
1426
Preparation of live *Drosophila* embryos for polarimetry measurements
mat–α-tub–GAL4 females were crossed to UASp–Af7 males, and F2 embryos were
collected and prepared for imaging following standard procedures (Mavrakis, 2016).
Briefly, F1 progeny was placed in embryo collection cages with fresh yeasted apple juice
agar plates. For live imaging, cellularizing F2 embryos were dechorionated with 50%
bleach, washed with water, transfered onto a heptane glue-coated round coverslip,
covered with halocarbon oil 200 (Tebubio, Cat#25073) and mounted in an Attofluor cell
chamber (ThermoFisher Scientific, Cat#A7816).
1435
Polarimetry measurements in live cellularizing Drosophila embryos
Polarimetry stacks using 18 polarization angles were recorded in focal planes through
actomyosin rings at the invaginating membrane front (Figure 7A) or in focal planes
apicolaterally to the former through the basal adherens junctions (Figure 7B). An
exposure time of 0.2 s was used per polarized image. The pixels of the actomyosin rings
or basal adherens junctions for analysis were selected by a combination of intensity
thresholding and manual selection of the region to analyze. Analysis and data
representation, including color-coded stick representations of the measured angles per
pixel and polar histograms were done with PyPOLAR. Considering that Af7 dipoles are
parallel to actin filaments, in order to assess how the measured actin filament orientations
distribute with respect to the actomyosin ring contour, the "edge detection" tool in
PyPOLAR was used in combination with intensity thresholding to isolate the ring contour-
associated pixels and normalize the angle distributions from 0°–180° to 0°–90° and
generate 0°–90° polar histograms, with 0° and 90° defining orientations parallel and
perpendicular to the ring contour, respectively (Figure S7A).

Characterization of actin organization reporters in the *Drosophila* flight muscle

Flight tests

Flight tests were performed as described in (Schnorrer et al., 2010). 3 to 7 day-old males were dropped into a 1 m long / 15 cm in diameter plexiglass cylinder with marked sections. Landing in the different sections depends on their flight ability, which was thereby scored (top 40-cm section: wild type, middle 40-cm section: impaired flight ("weak flier" in Figure 7C), bottom 20-cm section: flightless) (see cartoon in Figure 7C). For each genotype, flight assays were performed three times with a minimum of ten males per assay. The total number of flies scored for each genotype is mentioned in the respective figure legend. GraphPad Prism was used to generate bar graphs of the quantified flight ability per genotype and the mean percentages are mentioned in the respective legend.

Preparation of fixed adult flight muscles

Head, wings and abdomen were cut off the thorax of anaesthetized adult flies with fine scissors, and the thoraxes were fixed for 30 min in 4% paraformaldehyde in PBST (PBS + 0.3% Triton X-100). After three 10 min washes in PBST, thoraxes were placed on double-sided tape and cut sagittally dorsal to ventral with a microtome blade (Feather C35). The thorax halves were placed in PBST with Alexa 568-phalloidin (Invitrogen, 1:500) and incubated overnight at 4°C on a shaker. Hemithoraxes were then washed 3 times 10 min in PBST at room temperature and mounted in Vectashield with 2 spacer coverslips on each side.

Preparation of live flight muscles

Flight muscles were dissected, mounted in Schneider medium (no fixation), and imaged within 30 min following dissection. After removal of the head, abdomen and wings, a first incision was performed through the cuticle with sharp forceps (Dumont #5 forceps, Fine Science Tools, Cat#11252-20) at the median plane. The thorax was then gently pulled open into two halves, which were then fully disconnected through cutting of the ventral connective tissues using fine dissection scissors (Fine Science Tools, Cat#15009-08). The dissection resulted in relatively intact flight muscles still attached to the tendon cells of the thorax. Samples were mounted in Schneider medium using two coverslip spacers and imaged immediately.

Polarimetry measurements in live Drosophila flight muscle

The polarimetry analysis shown was from flight muscle expressing the reporters throughout muscle development with the Mef2-GAL4 driver apart from flight muscle expressing Af7, for which Af7 was expressed only transiently after muscle development with the tub-GAL80ts ; Mef2-GAL4 driver. One hemithorax per animal was used for polarimetry measurements, with 4-7 hemithoraces measured for each strain. Polarimetry stacks using 18 polarization angles were recorded in 1-7 different fields of view for each hemithorax. Ten myofibrils were analyzed per field of view (red-outlined boxes in Figure 7E-G). The pixels containing individual myofibrils within each field of view were selected by a combination of intensity thresholding and manual selection of the region to analyze. Each myofibril contained typically 3,000-10,000 analyzed pixels. Analysis and data representation, including color-coded stick representations of the measured angles per

pixel and histograms were done with PyPOLAR. The histograms shown in Figure 7E-G are from single myofibrils. GraphPad Prism was used to generate scatter plots of the quantified ψ angle distributions per strain; the number of myofibrils measured for each strain, the respective median values and the statistical test used to evaluate differences are mentioned in the respective legend.

Characterization of actin organization reporters in the *Drosophila* wing

For wing analysis, we anesthetized 50 young adult males from the progeny (using CO_2) and removed one wing from each fly with fine tweezers. These wings were then directly mounted between a slide and a coverslip using UV-cured optical adhesive (Thorlabs, Cat#NOA63). Images of the wings were captured using a digital microscope (Dino-Lite). For wing size analysis, we utilized landmarks within the wing vein pattern to measure specific distances. For the long axis, we measured the distance between the proximal end of L5 and the distal end of L3, following the nomenclature from (De Celis, 2003). For the short axis, we measured the distance between the distal end of L5 and the intersection of the opposite side of the wing with a line perpendicular to the long axis, passing through the distal end of L5. These two distances are represented in Figure S7B. MATLAB was used to generate box plots of the quantified wing area. The number of wings for each genotype, the respective median values of $L_L \cdot L_S$ and the statistical test used to evaluate differences are mentioned in the respective legend.

Protein structures and protein sequence alignments

Cartoon representations of protein structures were generated with the open-source software PyMOL (Schrödinger). The structure shown for cpGFP1-10/11 in Figure 1J corresponds to the structure of circular permutated red fluorescent protein Kate (PDB 3RWT) and is used to illustrate the design principle of our circular permutants. The PDB IDs for the remaining structures are as follows: 2B3P (sfGFP, Figure 1F, J, K), 1GFL (wild-type GFP, Figure 1I), 7AD9 (Lifeact-F-actin complex, Figure S1C, D), 1QAG (Utrophin, Figure S2G), 4N6T (Adhiron/Affimer, Figure S3I) and 5JLF (F-actin-tropomyosin complex, Figure 4B). The structure of F-tractin (Figure S3E) was generated using the AlphaFold database at EMBL-EBI. The multiple sequence alignment of Lifeact sequences (Figure S1B) was generated with Clustal Omega (EMBL-EBI). Graphical representations illustrating the conservation of residues for Lifeact, F-tractin and G-actin (Figures S1B, S3D, 4C) were generated using the WebLogo application (University of California, Berkeley). Interface areas were analyzed using PISA calculations as implemented on the EMBL-EBI server and visually inspected using PyMOL.

Polarimetry
Optical setup

Fluorescence images were acquired with a confocal spinning disk unit (CSU-X1-M1, Yokogawa) connected to the side-port of an inverted microscope (Eclipse Ti2-E, Nikon) using a x2 magnifier (Yokogawa), a Nikon Plan Apo ×100/1.45 NA oil immersion objective lens and an EMCCD camera with 1024×1024 pixels, 13×13 µm pixel size (iXon Ultra 888, Andor) resulting in an image pixel size of 65 nm. Z-stacks were acquired using a piezo stage (P-736, PI). The lateral position of the sample was controlled with a translation piezo stage (U-780, PI). The spinning disk is equipped with a multiline dichroic mirror

(Di01-T405/488/568/647-13x15x0.5, Semrock) and an emission filter wheel with filters
adapted to the studied emission: band pass 540/80 for EGFP/sfGFP and AF488 (FF01-
540/80-25, Semrock), band pass 593/46 for sfCherry2 (FF01-593/46-25, Semrock), and
long pass 655 for SiR-actin (Et655lp, Chroma). The laser excitation is provided by
polarized continuous lasers (488-, 561- and 641-nm laser lines, Sapphire, Coherent)
combined with a set of dichroic mirrors, each of the laser being used separately with a
power of typically 0.5 mW at the entrance of the spinning disk. The laser beams are sent
into an electro-optic modulator (EOM) (Pockels cell, No 28-NP; Quantum Technology)
followed by a quarter wave plate (WPQ05M-488; Thorlabs) to create a linear rotating
polarization. The voltages sent to the Pockels cell to provide known polarization rotations
are determined in a preliminary calibration step, using a polarimeter based on the quarter
wave plate method, as described in (Wang et al., 2013). As the whole optical path involves
reflections on mirrors and transmission through a dichroic mirror, the polarization after
the Pockels cell system is likely to be deformed. Polarization distortion compensation of
the spinning disk dichroic mirror is provided by placing an identical dichroic mirror (Di01-
T405/488/568/647-13x15x0.5, Semrock) in the path of the laser line just after the quarter
wave plate, such that s and p polarization components are exchanged at the first and
second dichroic transmissions. This configuration ensures minimization of the
polarization ellipticity and diattenuation produced by the dichroic mirror. The remaining
distorsions are characterized following the procedure of (Wang et al., 2013), using a
polarimeter based on the quarter wave plate method. The beam is then expanded using
a 10× telescope (BE10, Thorlabs) and sent directly to the microlens array of the CSU by
reflection on a second entrance mirror. The microlens and pinhole arrays of the CSU
disks rotate synchronously at a speed of 1,800 rpm, while the EMCCD and EOM are
synchronized to ensure a fast stack recording for a given number of incident polarization
(Wang et al., 2013). Exposure times are in the range of 0.1-0.5 s, and 18 polarization
angles are typically measured per polarimetry stack, which leads to a few seconds per
polarimetry stack.
1571
Signal processing
Fluorescence is generated from the coupling of fluorophore dipoles with the incident
linearly polarized electric field denoted $E(\alpha)$, whose orientation is an angle α with the
horizontal axis X of the sample plane. Inside the confocal volume, each fluorescent
molecule exhibits an absorption dipole vector μ_{abs} with an orientation (θ, ϕ) in the
macroscopic sample frame. The recorded fluorescence intensity from a single molecule
is proportional to the absorption probability $P_{abs}(\theta, \phi) = |\mu_{abs}(\theta, \phi) \cdot E(\alpha)|^2$. The total
intensity from an ensemble of molecules in the focal volume is therefore the sum of the
intensities from all single molecules present in this volume, whose size is typically 300
nm laterally and 600 nm longitudinally. This results in an averaged intensity: $I(\alpha) =$
$\iint |\mu_{abs}(\theta, \phi) \cdot E(\alpha)|^2 \sin\theta \, d\theta d\phi$ (Kress et al., 2013). The intensity is thus maximized
when the absorption dipoles of the molecules are aligned with the electric field. We
assume that the orientations explored by molecular dipoles are constrained within an
angular cone of total aperture angle ψ, oriented in the sample plane along the direction ρ
relative to X, the horizontal axis of the sample plane. Physically, ψ is related to a
« molecular order » quantity, which determines the degree of angular variations present
within the focal spot at a given pixel position, averaged over time and space. Note that

when fluorescent molecules are attached to actin with a degree of angular fluctuations due to their linker to actin, ψ encompasses three contributions : (1) the mean tilt angle ξ of the molecule with respect to the actin filament axis, (2) the angular fluctuations of the molecule due to its linker flexibility, and (3) the static organization of the actin filaments. The mean orientation ρ, on the other hand, determines the mean direction of the molecules. Therefore when the molecules are attached to actin in a constrained manner (i.e. angular fluctuations are not isotropic), in an assembly of aligned filaments, ρ can take two values : either $\rho = 0°$ when the tilt angle of the molecules ξ is close to the filament axis with $\xi < 45°$, or $\rho = 90°$ when the molecules are away from the filament axis with $\xi > 45°$. Thus, the angles ρ and ψ quantify the full information on the molecular organization at each pixel of an image. We note that the measurements performed in this work are limited to a projection of the fluorophores' distribution in the sample plane, which is imposed by the manipulation of light polarization in this plane. This 2D projection leads to an overestimation of the order angle ψ when the cone distribution is tilted more than 45° out of plane (Kress et al., 2013).

The angles ρ and ψ are deduced from the measurement of the intensity modulation $I(\alpha)$, which takes the form (Kress et al., 2013; Wang et al., 2013): $I(\alpha) = a_0 + a_2(\rho, \psi) \cos 2\alpha + b_2(\rho, \psi) \sin 2\alpha$. The coefficients $a_2(\rho, \psi)$ and $b_2(\rho, \psi)$, which can be directly related to the parameters (ρ, ψ) (see below), are deduced from the decomposition $I(\alpha)$ into circular functions $(\cos 2\alpha, \sin 2\alpha)$. In practice, when several input polarization angles α_k are used in a polarimetry stack (typically, for 18 polarization angles, $\alpha_k = 0, 10°, ..., 170°$), we use the operations $a_2 = \frac{2}{a_0} \sum I(\alpha_k) \cos 2\alpha_k$ and $b_2 = \frac{2}{a_0} \sum I(\alpha_k) \sin 2\alpha_k$, using $a_0 = \sum I(\alpha_k)$.

To retreive the angular parameters (ρ, ψ) from the measured quantities $a_2(\rho, \psi)$ and $b_2(\rho, \psi)$, the following method is used to account for polarization distorsions (Kress et al., 2013): The presence of polarization distorsions is modelled in the intensity equation $I(\alpha) = \iint |\mu_{abs}(\theta, \phi) \cdot E(\alpha)|^2 \sin \theta \, d\theta d\phi$ by including a distorted $E(\alpha) = (\cos \alpha, \gamma \sin \alpha \, e^{i\delta})$, with γ a diattenuation factor, which produces an energy loss between the s and p polarization components of an electric field, and δ a birefringence factor, which produces a phase difference between the s and p components. In this model, the polarization distorsions are supposed to originate from an equivalent phase plate whose axes coincide with the horizontal and vertical directions of the sample, which is reasonable considering that all reflections in the optical path involve s and p directions along these axes. Including these distorsions allows the construction of a map of the dependence of both a_2 and b_2 parameters, as fuctions of (ρ, ψ). Without any distorsions, these maps take the form of disks from which (ρ, ψ) can be unambiguously determined by the one-to-one relationship between (ρ, ψ) and (a_2, b_2) (Kress et al., 2013). In the presence of distorsions, the disks are deformed but the relation stays unambiguous, therefore it is possible to find (ρ, ψ) from the measurement of (a_2, b_2), using a minimization method in the $(\rho, \psi) \, vs \, (a_2, b_2)$ lookup table for instance. Finally, the parameters (ρ, ψ) extracted from the $(\rho, \psi) \, vs \, (a_2, b_2)$ disk analysis are represented in a single polarimetry image that combines molecular order and orientation, superimposed to the fluorescence intensity image built from the total intensity $\sum I(\alpha_k)$.

In experimental measurements, the $I(\alpha)$ modulation is affected by noise, which impacts the determination of the (ρ, ψ) parameters. The precision on the determination of (ρ, ψ) increases as the inverse square of the total intensity. It has been shown that above 5000

photons per pixel (which is typically the case for GFP imaging), the precision reaches about 1° for ρ and 3° for ψ, except at extreme high-order conditions ($\psi \sim 0°$) where the precision in ψ reaches 5° (Kress et al., 2013).

We note that the reasoning for the dependence of absorption probability on the fluorophore dipole orientation is similar to that for the dependence of emission probability: the polarized emission scheme exploited in fluorescence anisotropy and polarization emission analysis (Chen et al., 2020; Forkey et al., 2000) is not exploited in this study, but could be similarly applied (Kampmann et al., 2011; Mattheyses et al., 2010; Mehta et al., 2016; Nordenfelt et al., 2017; Swaminathan et al., 2017; Vrabioiu and Mitchison, 2006). Finally, the angle ψ used in this work can also be directly related to other quantities used to define molecular orientational organization, in particular the generic, distribution-independent order parameter used in aligned structures such as lipid membranes and liquid-crystalline polymers (Jahnig, 1979).

QUANTIFICATION AND STATISTICAL ANALYSIS

The quantification method for each experiment is described in the respective method details section. The statistical details of the experiments, including the exact value of n and what n represents, the definition of center, dispersion and precision measures (mean, median, SD, SEM) in the plots and graphs, the software and statistical test used to evaluate statistical significance of differences, and the definition of statistical significance are mentioned in the method details sections and respective Figure legends.

REFERENCES

Abrahamsson, S., McQuilken, M., Mehta, S.B., Verma, A., Larsch, J., Ilic, R., Heintzmann, R., Bargmann, C.I., Gladfelter, A.S., and Oldenbourg, R. (2015). MultiFocus Polarization Microscope (MF-PolScope) for 3D polarization imaging of up to 25 focal planes simultaneously. Opt Express *23*, 7734-7754.

Akemann, W., Raj, C.D., and Knöpfel, T. (2001). Functional characterization of permuted enhanced green fluorescent proteins comprising varying linker peptides. Photochem Photobiol *74*, 356-363.

Andresen, M., Schmitz-Salue, R., and Jakobs, S. (2004). Short tetracysteine tags to beta-tubulin demonstrate the significance of small labels for live cell imaging. Mol Biol Cell *15*, 5616-5622.

Bahler, J., Wu, J.Q., Longtine, M.S., Shah, N.G., McKenzie, A., 3rd, Steever, A.B., Wach, A., Philippsen, P., and Pringle, J.R. (1998). Heterologous modules for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe. Yeast *14*, 943-951.

Baird, G.S., Zacharias, D.A., and Tsien, R.Y. (1999). Circular permutation and receptor insertion within green fluorescent proteins. P Natl Acad Sci USA *96*, 11241-11246.

Ballestrem, C., Wehrle-Haller, B., and Imhof, B.A. (1998). Actin dynamics in living mammalian cells. J Cell Sci *111 (Pt 12)*, 1649-1658.

Barak, L.S., Yocum, R.R., Nothnagel, E.A., and Webb, W.W. (1980). Fluorescence staining of the actin cytoskeleton in living cells with 7-nitrobenz-2-oxa-1,3-diazole-phallacidin. Proc Natl Acad Sci U S A *77*, 980-984.

Baranwal, S., Naydenov, N.G., Harris, G., Dugina, V., Morgan, K.G., Chaponnier, C., and Ivanov, A.I. (2012). Nonredundant roles of cytoplasmic beta- and gamma-actin isoforms in regulation of epithelial apical junctions. Mol Biol Cell *23*, 3542-3553.

Belin, B.J., Cimini, B.A., Blackburn, E.H., and Mullins, R.D. (2013). Visualization of actin filaments and monomers in somatic cell nuclei. Mol Biol Cell *24*, 982-994.

Belin, B.J., Goins, L.M., and Mullins, R.D. (2014). Comparative analysis of tools for live cell imaging of actin network architecture. Bioarchitecture *4*, 189-202.

Belyy, A., Merino, F., Sitsel, O., and Raunser, S. (2020). Structure of the Lifeact-F-actin complex. PLoS Biol *18*, e3000925.

Bendezu, F.O., Hale, C.A., Bernhardt, T.G., and de Boer, P.A. (2009). RodZ (YfgA) is required for proper assembly of the MreB actin cytoskeleton and cell shape in E. coli. EMBO J *28*, 193-204.

Bendezu, F.O., Vincenzetti, V., Vavylonis, D., Wyss, R., Vogel, H., and Martin, S.G. (2015). Spontaneous
Cdc42 polarization independent of GDI-mediated extraction and actin-based trafficking. PLoS Biol *13*,
e1002097.
Benninger, R.K., Onfelt, B., Neil, M.A., Davis, D.M., and French, P.M. (2005). Fluorescence imaging of two-
photon linear dichroism: cholesterol depletion disrupts molecular orientation in cell membranes. Biophys J
88, 609-622.
Blanchoin, L., Boujemaa-Paterski, R., Sykes, C., and Plastino, J. (2014). Actin dynamics, architecture, and
mechanics in cell motility. Physiol Rev *94*, 235-263.
Brasselet, S., and Alonso, M.A. (2023). Polarization microscopy: from ensemble structural imaging to
single-molecule 3D orientation and localization microscopy. Optica *10*, 1486-1510.
Brasselet, S., Ferrand, P., Kress, A., Wang, X., Ranchon, H., and Gasecka, A. (2013). Imaging Molecular
Order in Cell Membranes by Polarization-Resolved Fluorescence Microscopy. In Fluorescent Methods to
Study Biological Membranes, Y.M.a.G. Duportail, ed. (Springer-Verlag Berlin Heidelberg), pp. 311-338
Brault, V., Sauder, U., Reedy, M.C., Aebi, U., and Schoenenberger, C.A. (1999). Differential epitope tagging
of actin in transformed Drosophila produces distinct effects on myofibril assembly and function of the
indirect flight muscle. Mol Biol Cell *10*, 135-149.
Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.
Bubb, M.R., Senderowicz, A.M., Sausville, E.A., Duncan, K.L., and Korn, E.D. (1994). Jasplakinolide, a
cytotoxic natural product, induces actin polymerization and competitively inhibits the binding of phalloidin
to F-actin. J Biol Chem *269*, 14869-14871.
Burkel, B.M., von Dassow, G., and Bement, W.M. (2007). Versatile fluorescent probes for actin filaments
based on the actin-binding domain of utrophin. Cell Motil Cytoskeleton *64*, 822-832.
Cabantous, S., Terwilliger, T.C., and Waldo, G.S. (2005). Protein tagging and detection with engineered
self-assembling fragments of green fluorescent protein. Nat Biotechnol *23*, 102-107.
Carlson, H.J., Cotton, D.W., and Campbell, R.E. (2010). Circularly permuted monomeric red fluorescent
proteins with new termini in the beta-sheet. Protein Sci *19*, 1490-1499.
Chakraborty, S., Jasnin, M., and Baumeister, W. (2020). Three-dimensional organization of the
cytoskeleton: A cryo-electron tomography perspective. Protein Sci *29*, 1302-1320.
Chen, A., Arora, P.D., McCulloch, C.A., and Wilde, A. (2017). Cytokinesis requires localized beta-actin
filament production by an actin isoform specific nucleator. Nat Commun *8*, 1530.
Chen, L., Chen, X., Yang, X., He, C., Wang, M., Xi, P., and Gao, J. (2020). Advances of super-resolution
fluorescence polarization microscopy and its applications in life sciences. Comput Struct Biotechnol J *18*,
2209-2216.
Chen, Q., Nag, S., and Pollard, T.D. (2012). Formins filter modified actin subunits during processive
elongation. J Struct Biol *177*, 32-39.
Choidas, A., Jungbluth, A., Sechi, A., Murphy, J., Ullrich, A., and Marriott, G. (1998). The suitability and
application of a GFP-actin fusion protein for long-term imaging of the organization and dynamics of the
cytoskeleton in mammalian cells. Eur J Cell Biol *77*, 81-90.
Chugh, P., and Paluch, E.K. (2018). The actin cortex at a glance. J Cell Sci *131*.
Cortes, J.C., Sato, M., Munoz, J., Moreno, M.B., Clemente-Ramos, J.A., Ramos, M., Okada, H., Osumi,
M., Duran, A., and Ribas, J.C. (2012). Fission yeast Ags1 confers the essential septum strength needed
for safe gradual cell abscission. J Cell Biol *198*, 637-656.
Costa, J., Fu, C., Syrovatkina, V., and Tran, P.T. (2013). Imaging individual spindle microtubule dynamics
in fission yeast. Methods Cell Biol *115*, 385-394.
Costantini, L.M., Fossati, M., Francolini, M., and Snapp, E.L. (2012). Assessing the tendency of fluorescent
proteins to oligomerize under physiologic conditions. Traffic *13*, 643-649.
Courtemanche, N., Pollard, T.D., and Chen, Q. (2016). Avoiding artefacts when counting polymerized actin
in live cells with LifeAct fused to fluorescent proteins. Nat Cell Biol *18*, 676-683.
Cramer, L.P. (1999). Role of actin-filament disassembly in lamellipodium protrusion in motile cells revealed
using the drug jasplakinolide. Curr Biol *9*, 1095-1105.
Cranfill, P.J., Sell, B.R., Baird, M.A., Allen, J.R., Lavagnino, Z., de Gruiter, H.M., Kremers, G.J., Davidson,
M.W., Ustione, A., and Piston, D.W. (2016). Quantitative assessment of fluorescent proteins. Nat Methods
13, 557-562.
Dancker, P., Low, I., Hasselbach, W., and Wieland, T. (1975). Interaction of actin with phalloidin:
polymerization and stabilization of F-actin. Biochim Biophys Acta *400*, 407-414.
De Celis, J.F. (2003). Pattern formation in the Drosophila wing: The development of the veins. Bioessays
25, 443-451.
DeMay, B.S., Bai, X.B., Howard, L., Occhipinti, P., Meseroll, R.A., Spiliotis, E.T., Oldenbourg, R., and
Gladfelter, A.S. (2011a). Septin filaments exhibit a dynamic, paired organization that is conserved from
yeast to mammals. Journal of Cell Biology *193*, 1065-1081.
DeMay, B.S., Noda, N., Gladfelter, A.S., and Oldenbourg, R. (2011b). Rapid and quantitative imaging of
excitation polarized fluorescence reveals ordered septin dynamics in live yeast. Biophys J *101*, 985-994.

Dopf, J., and Horiagon, T.M. (1996). Deletion mapping of the Aequorea victoria green fluorescent protein.
Gene *173*, 39-44.
Doyle, T., and Botstein, D. (1996). Movement of yeast cortical actin cytoskeleton visualized in vivo. Proc
Natl Acad Sci U S A *93*, 3886-3891.
Dugina, V., Zwaenepoel, I., Gabbiani, G., Clement, S., and Chaponnier, C. (2009). Beta and gamma-
cytoplasmic actins display distinct distribution and functional diversity. J Cell Sci *122*, 2980-2988.
Edwards, K.A., Demsky, M., Montague, R.A., Weymouth, N., and Kiehart, D.P. (1997). GFP-moesin
illuminates actin cytoskeleton dynamics in living tissue and demonstrates cell shape changes during
morphogenesis in Drosophila. Dev Biol *191*, 103-117.
Eltzner, B., Hauke, L., Huckemann, S., Rehfeldt, F., and Wollnik, C. (2020). A Statistical and Biophysical
Toolbox to Elucidate Structure and Formation of Stress Fibers. Top Appl Phys *134*, 263-282.
Feng, S., Sekine, S., Pessino, V., Li, H., Leonetti, M.D., and Huang, B. (2017). Improved split fluorescent
proteins for endogenous protein labeling. Nat Commun *8*, 370.
Ferrand, P., Gasecka, P., Kress, A., Wang, X., Bioud, F.Z., Duboisset, J., and Brasselet, S. (2014). Ultimate
use of two-photon fluorescence microscopy to map orientational behavior of fluorophores. Biophys J *106*,
2330-2339.
Fletcher, D.A., and Mullins, R.D. (2010). Cell mechanics and the cytoskeleton. Nature *463*, 485-492.
Forkey, J.N., Quinlan, M.E., and Goldman, Y.E. (2000). Protein structural dynamics by single-molecule
fluorescence polarization. Prog Biophys Mol Biol *74*, 1-35.
Frokjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M., Olesen, S.P., Grunnet,
M., and Jorgensen, E.M. (2008). Single-copy insertion of transgenes in Caenorhabditis elegans. Nat Genet
40, 1375-1383.
Gally, C., Wissler, F., Zahreddine, H., Quintin, S., Landmann, F., and Labouesse, M. (2009). Myosin II
regulation during C. elegans embryonic elongation: LET-502/ROCK, MRCK-1 and PAK-1, three kinases
with different roles. Development *136*, 3109-3119.
Gasecka, A., Han, T.J., Favard, C., Cho, B.R., and Brasselet, S. (2009). Quantitative imaging of molecular
order in lipid membranes using two-photon fluorescence polarimetry. Biophys J *97*, 2854-2862.
Gayrard, C., and Borghi, N. (2016). FRET-based Molecular Tension Microscopy. Methods *94*, 33-42.
Griffin, B.A., Adams, S.R., and Tsien, R.Y. (1998). Specific covalent labeling of recombinant protein
molecules inside live cells. Science *281*, 269-272.
Hafi, N., Grunwald, M., van den Heuvel, L.S., Aspelmeier, T., Chen, J.H., Zagrebelsky, M., Schutte, O.M.,
Steinem, C., Korte, M., Munk, A.*, et al.* (2014). Fluorescence nanoscopy by polarization modulation and
polarization angle narrowing. Nat Methods *11*, 579-584.
Heisenberg, C.P., and Bellaiche, Y. (2013). Forces in tissue morphogenesis and patterning. Cell *153*, 948-
962.
Heuser, J.E., and Kirschner, M.W. (1980). Filament organization revealed in platinum replicas of freeze-
dried cytoskeletons. J Cell Biol *86*, 212-234.
Hochbaum, D., Ferguson, A.A., and Fisher, A.L. (2010). Generation of transgenic C. elegans by biolistic
transformation. J Vis Exp.
Huang, J.Q., Huang, Y.Y., Yu, H.C., Subramanian, D., Padmanabhan, A., Thadani, R., Tao, Y.Q., Tang,
X., Wedlich-Soldner, R., and Balasubramanian, M.K. (2012). Nonmedially assembled F-actin cables
incorporate into the actomyosin ring in fission yeast. Journal of Cell Biology *199*, 831-847.
Hunter, C., and Wieschaus, E. (2000). Regulated expression of nullo is required for the formation of distinct
apical and basal adherens junctions in the Drosophila blastoderm. J Cell Biol *150*, 391-401.
Jahnig, F. (1979). Structural order of lipids and proteins in membranes: evaluation of fluorescence
anisotropy data. Proc Natl Acad Sci U S A *76*, 6361-6365.
Johnson, H.W., and Schell, M.J. (2009). Neuronal IP3 3-kinase is an F-actin-bundling protein: role in
dendritic targeting and regulation of spine morphology. Mol Biol Cell *20*, 5166-5180.
Juanes, M.A., Isnardon, D., Badache, A., Brasselet, S., Mavrakis, M., and Goode, B.L. (2019). The role of
APC-mediated actin assembly in microtubule capture and focal adhesion turnover. J Cell Biol *218*, 3415-
3435.
Kampmann, M., Atkinson, C.E., Mattheyses, A.L., and Simon, S.M. (2011). Mapping the orientation of
nuclear pore proteins in living cells with polarized fluorescence microscopy. Nat Struct Mol Biol *18*, 643-
649.
Keeney, J.B., and Boeke, J.D. (1994). Efficient Targeted Integration at Leu1-32 and Ura4-294 in
Schizosaccharomyces-Pombe. Genetics *136*, 849-856.
Keep, N.H., Winder, S.J., Moores, C.A., Walke, S., Norwood, F.L., and Kendrick-Jones, J. (1999). Crystal
structure of the actin-binding region of utrophin reveals a head-to-tail dimer. Structure *7*, 1539-1546.
Kim, H.K., and Kaang, B.K. (1998). Truncated green fluorescent protein mutants and their expression in
neurons. Brain Res Bull *47*, 35-41.
Kokai, E., Beck, H., Weissbach, J., Arnold, F., Sinske, D., Sebert, U., Gaiselmann, G., Schmidt, V., Walther,
P., Munch, J.*, et al.* (2014). Analysis of nuclear actin by overexpression of wild-type and actin mutant
proteins. Histochem Cell Biol *141*, 123-135.

Kress, A., Ferrand, P., Rigneault, H., Trombik, T., He, H.T., Marguet, D., and Brasselet, S. (2011). Probing
orientational behavior of MHC class I protein and lipid probes in cell membranes by fluorescence
polarization-resolved imaging. Biophys J *101*, 468-476.
Kress, A., Wang, X., Ranchon, H., Savatier, J., Rigneault, H., Ferrand, P., and Brasselet, S. (2013).
Mapping the local organization of cell membranes using excitation-polarization-resolved confocal
fluorescence microscopy. Biophys J *105*, 127-136.
Kumari, A., Kesarwani, S., Javoor, M.G., Vinothkumar, K.R., and Sirajuddin, M. (2020). Structural insights
into actin filament recognition by commonly used cellular actin markers. EMBO J *39*, e104006.
Lamm, N., Read, M.N., Nobis, M., Van Ly, D., Page, S.G., Masamsetti, V.P., Timpson, P., Biro, M., and
Cesare, A.J. (2020). Nuclear F-actin counteracts nuclear deformation and promotes fork repair during
replication stress. Nat Cell Biol *22*, 1460-1470.
Lappalainen, P., Kotila, T., Jegou, A., and Romet-Lemonne, G. (2022). Biochemical and mechanical
regulation of actin dynamics. Nat Rev Mol Cell Biol *23*, 836-852.
Lawson, C.D., Peel, S., Jayo, A., Corrigan, A., Iyer, P., Baxter Dalrymple, M., Marsh, R.J., Cox, S., Van
Audenhove, I., Gettemans, J*., et al.* (2022). Nuclear fascin regulates cancer cell survival. Elife *11*.
Lazar, J., Bondar, A., Timr, S., and Firestein, S.J. (2011). Two-photon polarization microscopy reveals
protein structure and function. Nat Methods *8*, 684-690.
Li, X., Zhang, G., Ngo, N., Zhao, X., Kain, S.R., and Huang, C.C. (1997). Deletions of the Aequorea victoria
green fluorescent protein define the minimal domain required for fluorescence. J Biol Chem *272*, 28545-
28549.
Loison, O., Weitkunat, M., Kaya-Copur, A., Nascimento Alves, C., Matzat, T., Spletter, M.L., Luschnig, S.,
Brasselet, S., Lenne, P.F., and Schnorrer, F. (2018). Polarization-resolved microscopy reveals a muscle
myosin motor-independent mechanism of molecular actin ordering during sarcomere maturation. PLoS Biol
16, e2004718.
Lopata, A., Hughes, R., Tiede, C., Heissler, S.M., Sellers, J.R., Knight, P.J., Tomlinson, D., and Peckham,
M. (2018). Affimer proteins for F-actin: novel affinity reagents that label F-actin in live and fixed cells. Sci
Rep *8*, 6572.
Lukinavicius, G., Reymond, L., D'Este, E., Masharina, A., Gottfert, F., Ta, H., Guther, A., Fournier, M.,
Rizzo, S., Waldmann, H*., et al.* (2014). Fluorogenic probes for live-cell imaging of the cytoskeleton. Nat
Methods *11*, 731-733.
Madani, F., Lind, J., Damberg, P., Adams, S.R., Tsien, R.Y., and Graslund, A.O. (2009). Hairpin structure
of a biarsenical-tetracysteine motif determined by NMR spectroscopy. J Am Chem Soc *131*, 4613-4615.
Martin, B.R., Giepmans, B.N., Adams, S.R., and Tsien, R.Y. (2005). Mammalian cell-based optimization of
the biarsenical-binding tetracysteine motif for improved fluorescence and affinity. Nat Biotechnol *23*, 1308-
1314.
Mattheyses, A.L., Kampmann, M., Atkinson, C.E., and Simon, S.M. (2010). Fluorescence anisotropy
reveals order and disorder of protein domains in the nuclear pore complex. Biophys J *99*, 1706-1717.
Mavrakis, M. (2016). Visualizing septins in early Drosophila embryos. Methods Cell Biol *136*, 183-198.
Mavrakis, M., Azou-Gros, Y., Tsai, F.C., Alvarado, J., Bertin, A., Iv, F., Kress, A., Brasselet, S., Koenderink,
G.H., and Lecuit, T. (2014). Septins promote F-actin ring formation by crosslinking actin filaments into
curved bundles. Nat Cell Biol *16*, 322-334.
Mehta, S.B., McQuilken, M., La Riviere, P.J., Occhipinti, P., Verma, A., Oldenbourg, R., Gladfelter, A.S.,
and Tani, T. (2016). Dissection of molecular assembly dynamics by tracking orientation and position of
single molecules in live cells. Proc Natl Acad Sci U S A *113*, E6352-E6361.
Moores, C.A., and Kendrick-Jones, J. (2000). Biochemical characterisation of the actin-binding properties
of utrophin. Cell Motil Cytoskeleton *46*, 116-128.
Moreno, S., Klar, A., and Nurse, P. (1991). Molecular genetic analysis of fission yeast
Schizosaccharomyces pombe. Methods Enzymol *194*, 795-823.
Mullins, R.D., and Hansen, S.D. (2013). In vitro studies of actin filament and network dynamics. Curr Opin
Cell Biol *25*, 6-13.
Nakai, N., Sato, K., Tani, T., Saito, K., Sato, F., and Terada, S. (2019). Genetically encoded orientation
probes for F-actin for fluorescence polarization microscopy. Microscopy (Oxf) *68*, 359-368.
Nordenfelt, P., Moore, T.I., Mehta, S.B., Kalappurakkal, J.M., Swaminathan, V., Koga, N., Lambert, T.J.,
Baker, D., Waters, J.C., Oldenbourg, R*., et al.* (2017). Direction of actin flow dictates integrin LFA-1
orientation during leukocyte migration. Nat Commun *8*, 2047.
Ormo, M., Cubitt, A.B., Kallio, K., Gross, L.A., Tsien, R.Y., and Remington, S.J. (1996). Crystal structure of
the Aequorea victoria green fluorescent protein. Science *273*, 1392-1395.
Osawa, M., and Erickson, H.P. (2005). Probing the domain structure of FtsZ by random truncation and
insertion of GFP. Microbiology (Reading) *151*, 4033-4043.
Paluch, E., and Heisenberg, C.P. (2009). Biology and physics of cell shape changes in development. Curr
Biol *19*, R790-799.
Pedelacq, J.D., Cabantous, S., Tran, T., Terwilliger, T.C., and Waldo, G.S. (2006). Engineering and
characterization of a superfolder green fluorescent protein. Nat Biotechnol *24*, 79-88.

Pollard, T.D. (2016). Actin and Actin-Binding Proteins. Cold Spring Harb Perspect Biol *8*.

Pollard, T.D., and Cooper, J.A. (2009). Actin, a central player in cell shape and movement. Science *326*,
1208-1212.

Priess, J.R., and Hirsh, D.I. (1986). Caenorhabditis elegans morphogenesis: the role of the cytoskeleton in
elongation of the embryo. Dev Biol *117*, 156-173.

Raghunathan, G., Soundrarajan, N., Sokalingam, S., Yun, H., and Lee, S.G. (2012). Deletional protein
engineering based on stable fold. PLoS One *7*, e51510.

Redemann, S., Schloissnig, S., Ernst, S., Pozniakowsky, A., Ayloo, S., Hyman, A.A., and Bringmann, H.
(2011). Codon adaptation-based control of protein expression in C. elegans. Nat Methods *8*, 250-252.

Renshaw, M.J., Liu, J., Lavoie, B.D., and Wilde, A. (2014). Anillin-dependent organization of septin
filaments promotes intercellular bridge elongation and Chmp4B targeting to the abscission site. Open Biol
4, 130190.

Riedl, J., Crevenna, A.H., Kessenbrock, K., Yu, J.H., Neukirchen, D., Bista, M., Bradke, F., Jenne, D.,
Holak, T.A., Werb, Z.*, et al.* (2008). Lifeact: a versatile marker to visualize F-actin. Nat Methods *5*, 605-607.

Rimoli, C.V., Valades-Cruz, C.A., Curcio, V., Mavrakis, M., and Brasselet, S. (2022). 4polar-STORM
polarized super-resolution imaging of actin filament organization in cells. Nat Commun *13*, 301.

Royant, A., and Noirclerc-Savoye, M. (2011). Stabilizing role of glutamic acid 222 in the structure of
Enhanced Green Fluorescent Protein. J Struct Biol *174*, 385-390.

Schnorrer, F., Schonbauer, C., Langer, C.C., Dietzl, G., Novatchkova, M., Schernhuber, K., Fellner, M.,
Azaryan, A., Radolf, M., Stark, A.*, et al.* (2010). Systematic genetic analysis of muscle morphogenesis and
function in Drosophila. Nature *464*, 287-291.

Shaner, N.C., Lin, M.Z., McKeown, M.R., Steinbach, P.A., Hazelwood, K.L., Davidson, M.W., and Tsien,
R.Y. (2008). Improving the photostability of bright monomeric orange and red fluorescent proteins. Nat
Methods *5*, 545-551.

Shum, M.S., Pasquier, E., Po'uha, S.T., O'Neill, G.M., Chaponnier, C., Gunning, P.W., and Kavallaris, M.
(2011). gamma-Actin regulates cell migration and modulates the ROCK signaling pathway. FASEB J *25*,
4423-4433.

Slocum, J.D., and Webb, L.J. (2017). A Double Decarboxylation in Superfolder Green Fluorescent Protein
Leads to High Contrast Photoactivation. J Phys Chem Lett *8*, 2862-2868.

Spira, F., Cuylen-Haering, S., Mehta, S., Samwer, M., Reversat, A., Verma, A., Oldenbourg, R., Sixt, M.,
and Gerlich, D.W. (2017). Cytokinesis in vertebrate cells initiates by contraction of an equatorial actomyosin
network composed of randomly oriented filaments. Elife *6*.

Sugizaki, A., Sato, K., Chiba, K., Saito, K., Kawagishi, M., Tomabechi, Y., Mehta, S.B., Ishii, H., Sakai, N.,
Shirouzu, M.*, et al.* (2021). POLArIS, a versatile probe for molecular orientation, revealed actin filaments
associated with microtubule asters in early embryos. Proc Natl Acad Sci U S A *118*.

Sund, S.E., Swanson, J.A., and Axelrod, D. (1999). Cell membrane orientation visualized by polarized total
internal reflection fluorescence. Biophys J *77*, 2266-2283.

Svitkina, T.M., Verkhovsky, A.B., and Borisy, G.G. (1995). Improved procedures for electron microscopic
visualization of the cytoskeleton of cultured cells. J Struct Biol *115*, 290-303.

Swaminathan, V., Kalappurakkal, J.M., Mehta, S.B., Nordenfelt, P., Moore, T.I., Koga, N., Baker, D.A.,
Oldenbourg, R., Tani, T., Mayor, S.*, et al.* (2017). Actin retrograde flow actively aligns and orients ligand-
engaged integrins in focal adhesions. Proc Natl Acad Sci U S A *114*, 10648-10653.

Swulius, M.T., Nguyen, L.T., Ladinsky, M.S., Ortega, D.R., Aich, S., Mishra, M., and Jensen, G.J. (2018).
Structure of the fission yeast actomyosin ring during constriction. Proc Natl Acad Sci U S A *115*, E1455-
E1464.

Tiede, C., Tang, A.A., Deacon, S.E., Mandal, U., Nettleship, J.E., Owen, R.L., George, S.E., Harrison, D.J.,
Owens, R.J., Tomlinson, D.C.*, et al.* (2014). Adhiron: a stable and versatile peptide display scaffold for
molecular recognition applications. Protein Eng Des Sel *27*, 145-155.

Topell, S., Hennecke, J., and Glockshuber, R. (1999). Circularly permuted variants of the green fluorescent
protein. Febs Lett *457*, 283-289.

Tsien, R.Y. (1998). The green fluorescent protein. Annu Rev Biochem *67*, 509-544.

Valades Cruz, C.A., Shaban, H.A., Kress, A., Bertaux, N., Monneret, S., Mavrakis, M., Savatier, J., and
Brasselet, S. (2016). Quantitative nanoscale imaging of orientational order in biological filaments by
polarized superresolution microscopy. Proc Natl Acad Sci U S A *113*, E820-828.

van den Ent, F., Amos, L.A., and Lowe, J. (2001). Prokaryotic origin of the actin cytoskeleton. Nature *413*,
39-44.

Verkhusha, V.V., Tsukita, S., and Oda, H. (1999). Actin dynamics in lamellipodia of migrating border cells
in the Drosophila ovary revealed by a GFP-actin fusion protein. Febs Lett *445*, 395-401.

Vrabioiu, A.M., and Mitchison, T.J. (2006). Structural insights into yeast septin organization from polarized
fluorescence microscopy. Nature *443*, 466-469.

Vuong-Brender, T.T., Ben Amar, M., Pontabry, J., and Labouesse, M. (2017). The interplay of stiffness and
force anisotropies drives embryo elongation. Elife *6*.

Wang, X., Kress, A., Brasselet, S., and Ferrand, P. (2013). High frame-rate fluorescence confocal angle-
resolved linear dichroism microscopy. Rev Sci Instrum *84*, 053708.
Watanabe, N., and Mitchison, T.J. (2002). Single-molecule speckle analysis of actin filament turnover in
lamellipodia. Science *295*, 1083-1086.
Westphal, M., Jungbluth, A., Heidecker, M., Muhlbauer, B., Heizer, C., Schwartz, J.M., Marriott, G., and
Gerisch, G. (1997). Microfilament dynamics during cell movement and chemotaxis monitored using a GFP-
actin fusion protein. Curr Biol *7*, 176-183.
Wieland, T. (1977). Modification of actins by phallotoxins. Naturwissenschaften *64*, 303-309.
Wulf, E., Deboben, A., Bautz, F.A., Faulstich, H., and Wieland, T. (1979). Fluorescent phallotoxin, a tool for
the visualization of cellular actin. Proc Natl Acad Sci U S A *76*, 4498-4502.
Yang, F., Moss, L.G., and Phillips, G.N., Jr. (1996). The molecular structure of green fluorescent protein.
Nat Biotechnol *14*, 1246-1251.
Yi, J., Wu, X.S., Crites, T., and Hammer, J.A., 3rd (2012). Actin retrograde flow and actomyosin II arc
contraction drive receptor cluster dynamics at the immunological synapse in Jurkat T cells. Mol Biol Cell
23, 834-852.
Zacharias, D.A., Violin, J.D., Newton, A.C., and Tsien, R.Y. (2002). Partitioning of lipid-modified monomeric
GFPs into membrane microdomains of live cells. Science *296*, 913-916.
Zhang, G., Gurtu, V., and Kain, S.R. (1996). An enhanced green fluorescent protein allows sensitive
detection of gene transfer in mammalian cells. Biochem Biophys Res Commun *227*, 707-711.
Zhanghao, K., Chen, X., Liu, W., Li, M., Liu, Y., Wang, Y., Luo, S., Wang, X., Shan, C., Xie, H.*, et al.* (2019).
Super-resolution imaging of fluorescent dipoles via polarized structured illumination microscopy. Nat
Commun *10*, 4694.
1959

Fig2 Martins et al 2024

Fig3 Martins et al 2024

Fig4 Martins et al 2024

Fig5 Martins et al 2024

Fig6 Martins et al 2024

Fig7 Martins et al 2024

Figure legends

Figure 1. Measuring actin filament organization in cells with polarimetry

(A) Left, example of the polarization response of a sample at a given pixel of the image as obtained from a recorded polarimetry stack. The polarimetry stack is made of 18 polarized fluorescence images acquired using an incident linear polarization angle, α, varying from 0° to 170° with steps of 10°. Raw datapoints are shown as triangles and the theoretical fitting curve as a solid line. Right, schematic of a hypothetical organization of four fluorescently-labeled actin filaments in the confocal volume of the measured pixel, with the different orientations of the fluorophore dipoles shown by green double-headed arrows. The fluorophore dipoles are parallel to the actin filament axis in this example. The angle ρ corresponds to the mean orientation of all dipoles and thus the average orientation of actin filaments in the confocal volume. The ρ value is represented with a purple stick whose orientation and color depict the mean filament orientation in the pixel (see colorbar in (B)). The angle ψ corresponds to the angular aperture explored by all dipoles and is thus a readout of the average actin filament alignment in the confocal volume: the higher the filament alignment, the smaller the ψ angle.

(B) Representative examples of polarimetry measurements of actin filament organization in fixed U2OS cells labeled with AF488-phalloidin (top) or SiR-actin (bottom). The zoomed-out images on the left are summed intensity images of the respective polarimetry stacks. Insets on the right show zoom-ins of selected regions of interest (red outlined boxes) containing actin stress fibers (SFs) in different orientations, with the measured ρ and ψ angles per pixel. The angles ρ are represented as ρ stick maps ("orientation maps"), with a stick per pixel whose orientation and color depict the mean filament orientation in the pixel. The values of ρ, from 0° to 180°, are color-coded according to the colorbar. The angles ψ are represented as ψ stick maps ("organization maps"), with a stick per pixel whose orientation depicts the mean filament orientation (ρ) and whose color corresponds to the mean filament alignment (ψ) in the pixel. The values of ψ, from 40° to 180°, are color-coded according to the colorbar. Given that actin filaments are parallel to the long axis of SFs, the experimental measurements show that AF488 dipoles in the AF488-phalloidin conjugates are parallel to actin filaments (green double-headed arrows in the scheme on the far right), while SiR dipoles in the SiR-actin conjugates are oriented perpendicular to actin filaments (red double-headed arrows in the scheme on the far right).

(C) Box plots depicting the distribution of ψ angle measurements on SFs as shown in (B). The data points, color-coded according to the ψ colorbar, are plotted on top of the respective box plots. On each box, the central mark indicates the median, and the left and right edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers, and the outliers are plotted individually using the 'x' symbol. The number of measurements in each box plot is n = 258 and 45 for AF488-phalloidin and SiR-actin, respectively. The respective median values are 126° and 133°.

(D) Schematics showing the dependence of measured ψ angles on the underlying actin filament organization, the mobility of the fluorophore and the tilt angle, ξ, of the fluorophore with respect to the axis of the actin filament. ψ is color-coded as in (B). The mean filament orientation, ρ, is the same in all cases. Flexible fluorophores will lead to very high (>160°) ψ values and thus an overestimation of disorder even for highly aligned actin filaments (i). Constrained fluorophores allow us to detect changes in actin filament organization (ii and iii *vs.* iv).

(E) Representative ψ stick maps on SFs from measurements in live cells expressing widely used GFP fusions of actin-binding peptides or domains, or G-actin itself. The number shown in orange corresponds to the number of amino acid residues of the linker between the GFP and the actin-binding moiety. Mean ψ values are shown.

(F) Ribbon representation of the crystal structure of superfolder GFP (sfGFP) (PDB 2B3P). The N-terminal 3_{10} helix and beta-strands β1 and β11 are depicted in red, cyan and blue, respectively. Magenta arrowheads point to the beginning of β1 and the end of β11.

(G) Top, amino acid sequence of the N- and C-termini of monomeric EGFP (mEGFP) and monomeric sfGFP (msfGFP). The depicted secondary structure elements, color code and arrowheads are as in (F). Residue numbering is as shown. Bottom, screening of N- and C-terminal truncation mutants of mEGFP and msfGFP using fluorescence-activated cell sorting (FACS). Bars (mean + SD) depict the measured percentages of fluorescence-positive (FP+) cells for full-length (FL), N-terminally (ΔN) and C-terminally (ΔC) truncated proteins. The mean values are, from left to right: 100, 100, 74, 94, 53, 100, 42, 85, 4, 72, 9, 63, 0.4, 43, 0.1, 80, 65, 96, 72, 86, 69, 87, 0.2, and 80. Data are from three independent

experiments. Statistical significance was obtained using an unpaired t-test. The different constructs were compared to the respective FL; ns=not significant (P>0.05); * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

(H) Screening of the GFP constructs used in (G) with spinning disk fluorescence microscopy. Representative images of live cells expressing each construct are shown. For the sake of comparison, images are displayed with the same intensity range. In the case of weakly fluorescent cells, insets show contrast-enhanced images. No fluorescence was detectable for mEGFPΔN12ΔC11.

(I) Ribbon representation of the GFP dimerization interface from the crystal structure of GFP (PDB 1GFL). The hydrophobic residues A206 (V206 for sfGFP), L221 and F223 in the dimer interface are shown in brown. The respective monomerizing mutations used for impairing dimerization are shown.

(J-K) Designs used in this study to immobilize genetically-encoded fluorophore fusions to actin-binding domains (ABDs) (J) or G-actin itself (K). ABDs were fused to the N- or C-terminus of msfGFP (J, left) or placed in-between the N- and C-terminus of GFP using a circularly permuted GFP (cpGFP) (J, right). For G-actin terminal fusions, msfGFP or tetracysteine peptides were fused to the N-terminus of G-actin (K, left and right). For G-actin intramolecular fusions, msfGFP, the β11 strand alone, or tetracysteine peptides were placed intramolecularly within the G-actin structure (K, left, middle and right).

Figure 2. Engineering of Lifeact-based actin filament organization reporters for live-cell polarimetry

(A) List of Lifeact ("Life" for short)-GFP fusions tested for their usability in live polarimetry measurements in U2OS cells expressing each fusion. All constructs were expressed with a CMV$_{trunc}$ promoter except the ones with an asterisk (*) which used a CMV promoter. The left column lists the tested fusions as follows: a short name for each fusion (e.g., Life-msfGFPΔN6) followed by its designated number ("L5" in this case) and its primary sequence. Secondary structure elements of GFP are color-coded as in Fig.1F. To facilitate the tracking of modifications, only the region of the primary sequence that is modified is shown. Sequences that stay unchanged from one construct to the next one are shown with an outlined box for the first one (e.g. L1) and then represented as gray-filled boxes in subsequent constructs (in this case, L2-L12). Additional columns report whether the construct is fluorescent (column "fluo") and if it binds actin SFs (column "F-actin"). Constructs were classified as fluorescent (+), very weakly fluorescent (±), or nonfluorescent (-), and as binding to F-actin (+), very weakly binding to F-actin (±), or not binding to F-actin (-). For very weakly fluorescent or nonfluorescent constructs, F-actin binding was not determined (nd). Box plots on the far right depict the distribution of ψ angle measurements on SFs for the respective constructs. Box plots are depicted as in Fig.1C. Polarimetry measurements were not performed for constructs that were weakly fluorescent or weak F-actin binders and thus not usable for routine polarimetry. The number of measurements (see methods for details), from top to bottom, are n= 30, 40, 25, 25, 26, 21, 18, 22, 21, 15, 17, 15, 21, 22, 11, 19, 24, 23, 29, 11, 10, 20, 9, 17, 20, 10, 33, 15, 23, 20, 21, 26, 11, 18, 16, 21, 16, 8, 16, 3, 22, 7. The respective median ψ values are 162, 161, 160, 155, 152, 143, 151, 148, 141, 144, 137, 138, 163, 155, 161, 153, 132, 133, 130, 136, 147, 138, 138, 149, 147, 167, 160, 160, 154, 162, 142, 134, 142, 160, 163, 165, 141, 145, 155, 159, 158, 160°. Statistical significance (right-most column) was obtained using a non-parametric Kruskal-Wallis test followed by a Dunn's multiple comparisons test. The different constructs were compared to L1; ns=not significant (P>0.05); * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

(B-C) Representative ψ stick maps on SFs from measurements in live cells expressing the shown C-terminal (B) and N-terminal (C) GFP fusions, with mean ψ values indicated. The selected images correspond to median ψ values of the respective distributions. See also Figure S1.

Figure 3. Engineering of Utrophin, F-tractin, Affimer6- and red fluorescent protein-based actin filament organization reporters for live-cell polarimetry

(A) List of selected Utrophin calponin homology domain ("Utr" for short)-GFP fusions tested for their usability in live polarimetry measurements in U2OS cells expressing each fusion. The full screen is shown in Figure S2A. Fusion illustration and classification and box plots are as detailed in Figure 2A. Statistical significance (right-most column) was obtained using a non-parametric Kruskal-Wallis test followed by a Dunn's multiple comparisons test. The different constructs were compared to U1; ns=not significant (P>0.05); *** P<0.001, **** P<0.0001. The number of measurements and the respective median ψ values for each fusion are mentioned in Figure S2A.

(B-C) Representative ψ stick maps on SFs from measurements in live cells expressing the shown N-terminal (B) and C-terminal (C) GFP fusions, with mean ψ values indicated. The selected images correspond to median ψ values of the respective distributions. See also Figure S2.

(D) List of selected F-tractin ("Ftr" for short)-GFP fusions tested for their usability in live polarimetry measurements in U2OS cells expressing each fusion. The full screen is shown in Figure S3C. Fusion illustration and classification and box plots are as detailed in Figure 2A. Statistical significance (right-most column) was obtained using a non-parametric Kruskal-Wallis test followed by a Dunn's multiple comparisons test. The different constructs were compared to F1; ns=not significant (P>0.05); * P<0.05, ** P<0.01, **** P<0.0001. The number of measurements and the respective median ψ values for each fusion are mentioned in Figure S3C.

(E-F) Representative ψ stick maps on SFs from measurements in live cells expressing the shown C-terminal (E) and N-terminal (F) GFP fusions, with mean ψ values indicated. The selected images correspond to median ψ values of the respective distributions. See also Figure S3.

(G) List of selected Affimer6 ("Aff6" for short)-GFP fusions tested for their usability in live polarimetry measurements in U2OS cells expressing each fusion. The full screen is shown in Figure S3F. Fusion illustration and classification and box plots are as detailed in Figure 2A. Statistical significance (right-most column) was obtained using a non-parametric Kruskal-Wallis test followed by a Dunn's multiple comparisons test. The different constructs were compared to Af12; ns=not significant (P>0.05); **** P<0.0001. The number of measurements and the respective median ψ values for each fusion are mentioned in Figure S3F.

(H-I) Representative ψ stick maps on SFs from measurements in live cells expressing the shown C-terminal (H) and N-terminal (I) GFP fusions, with mean ψ values indicated. The selected images correspond to median ψ values of the respective distributions. See also Figure S3.

(J) List of Affimer6- and Lifeact-based sfCherry2 ("sfCh2" for short) fusions tested for their usability in live polarimetry measurements in U2OS cells expressing each fusion. sfCherry2 fusions were either C-terminal (Af26- Af30, L73-L77), or N-terminal (L78-L81). Fusion illustration and classification and box plots are as detailed in Figure 2A. The number of measurements (see methods for details), from top to bottom, are n= 23, 16, 23, 25, 18, 23, 1, 22, 22, 22, 20, 25. The respective median ψ values are 168, 161, 165, 155, 109, 170, 165, 169, 163, 140, 113, 111°. Statistical significance (right-most column) was obtained using a non-parametric Kruskal-Wallis test followed by a Dunn's multiple comparisons test. The different Affimer6 constructs were compared to Af12; the different Lifeact constructs were compared to L1; ns=not significant (P>0.05); * P<0.05, ** P<0.01, **** P<0.0001.

(K) Representative ψ stick maps on SFs from measurements in live cells expressing the shown Affimer6-based (Af) and Lifeact-based (L) sfCherry2 fusions, with mean ψ values indicated. The selected images correspond to median ψ values of the respective distributions.

Figure 4. Engineering of G-actin-based actin filament organization reporters for live-cell polarimetry

(A) List of human β-actin ("act" or "actin" for short) fusions tested for their usability in live polarimetry measurements in U2OS cells expressing each fusion. The fluorophores used were GFP, the β11 strand alone ("GFP11"), or tetracysteine peptides ("cys6" or "cys12") as shown in Fig.1K. For G-actin terminal fusions, GFP or tetracysteine peptides were fused to the N-terminus of G-actin (A1-A6). For G-actin intramolecular fusions, msfGFP (A7-A23), the β11 strand alone (A24-A37), or tetracysteine peptides (A38-A47) were placed intramolecularly within the G-actin structure (see (B)). Fusion illustration and classification and box plots are as detailed in Figure 2A. Fusions A6, A40 and A41 localized both to SFs and to nuclear F-actin, and box plots include measurements from both F-actin pools. Measurements for fusions A42 and A47 are from nuclear F-actin. The number of measurements (see methods for details), from top to bottom, are n= 23, 24, 26, 15, 18, 6, 12, 17, 17, 19, 18, 18, 26, 26, 17, 1, 1. The respective median ψ values are 161, 160, 158, 146, 148, 168, 166, 157, 163, 162, 150, 148, 158, 162, 164, 167, 165°. Statistical significance (right-most column) was obtained using a non-parametric Kruskal-Wallis test followed by a Dunn's multiple comparisons test. The different constructs were compared to A1; ns=not significant (P>0.05); *** P<0.001, **** P<0.0001.

(B-C) Ribbon representation of F-actin with three consecutive G-actin monomers colored in green, magenta and blue (PDB 5JLF) (B, left). Helix h7, used as an insertion site in intramolecular fusions, is shown in red. A close-up view of h7 (dashed box) shows residues in the loops (in cyan) flanking the helix, with arrowheads pointing to the insertion sites used in intramolecular fusions (B, right). (C)

WebLogo3 representation of the conservation of residues in h7 and the flanking residues. Forty-five actin sequences were used for this representation, including organisms as diverse as *Drosophila*, fungi, Dictyostelium, Arabidopsis, and sea animals. Negatively- and positively-charged residues are shown in red and blue, respectively.

(D-E) Representative ψ stick maps on SFs from measurements in live cells expressing the shown N-terminal (D) and intramolecular (E) GFP fusions, with mean ψ values indicated. The selected images correspond to median ψ values of the respective distributions. See also Figure S4.

Figure 5. Polarimetry measurements of actin filament organization in live dividing fission yeast expressing selected reporters

(A) Time-lapse maximum intensity projection images of a fission yeast cell (orange dash outline) expressing an acto-myosin ring marker (Rlc1-mCherry) and a spindle pole body marker (Sid4-mCherry) to monitor major cytokinetic events. The time interval between frames corresponds to 6 minutes. "A" represents the cytokinetic ring assembly stage, "M", the cytokinetic ring maturation stage and "C", the cytokinetic ring constriction stage. Scale bar, 2 μm.

(B) Quantification of the time taken for ring assembly completion, ring maturation, ring constriction, and the total time for cytokinesis completion in fission yeast strains expressing each actin reporter and the cytokinetic markers. As a control ("C"), a strain expressing only the cytokinetic markers was used. Scatter plots show means ± SD. The number of cells for each strain is, from left to right: 41, 31, 29, 31, 31, 33, 29, 36, 33. The mean measured times for each strain are, from left to right: 12, 13, 17, 13, 15, 15, 16, 13, 15 min for ring assembly; 10, 11, 12, 10, 11, 14, 11, 12, 16 min for ring maturation; 19, 18, 20, 17, 18, 20, 19, 22, 21 min for ring constriction; and 41, 42, 49, 41, 45, 49, 46, 48, 51 min for total cytokinesis. A t-test was applied to evaluate statistical differences between each strain and the control; ns=not significant, P>0.05; * 0.05>P>0.03; ** P<0.03.

(C) Serial dilution assay showing the sensitivity of the fission yeast strains expressing the corresponding actin reporter to the Arp2/3 complex inhibitor CK666, the G-actin sequestering drug latrunculin A (LatA), and DMSO (vehicle control). As controls, a previously published strain expressing Lifeact under the control of the actin promoter ('Lifeact') (Huang et al., 2012) and a wild-type strain ('wt') were included in the assay.

(D) Serial dilution assay showing the genetic interaction between the profilin mutant *cdc3-319* and the expression of the different actin reporters.

(E-G) Polarimetry measurements of actin filament organization in the cytokinetic ring of living dividing fission yeast cells expressing Affimer6-based (E) and Lifeact-based (F) reporters. Representative measurements are shown for fusions with unconstrained GFPs (Af1, L2) and constrained GFPs (Af7, L22). Left panels in (E), (F) show inverted grayscale summed intensity images of polarimetry stacks for the respective dividing yeast cells (orange dash outlines). ρ and ψ stick maps of actin organization in the cytokinetic ring (red dash box in left panels) are shown in the middle and right panels, respectively. The standard deviation of ρ values (SDρ), mean ρ and ψ values are shown for each map. Scatter plots in (G) show the quantification of ψ angle distributions for each reporter. Scatter plots show medians with interquartile range. The number of cells for each strain is, from left to right: 26, 20, 17, 9. The respective median ψ values are 151, 90, 155, 121°. Statistical significance was obtained using a non-parametric Mann-Whitney test; **** P<0.0001. See also Figure S5.

Figure 6. Polarimetry measurements of actin filament organization in live elongating *C. elegans* embryos expressing selected reporters

(A) Viability of *C. elegans* strains expressing different reporters assessed by the number of unhatched embryos 12-16 h after egg-laying. EL, embryonic lethality. n=number of scored embryos per genotype. See methods for details of the genotypes.

(B) Embryonic growth curves, showing fold-change of embryonic length until hatching based on differential interference contrast (DIC) filming of the indicated *C. elegans* strains. Curves show means ± SEM. n=10 embryos were measured for each genotype. See also Figure S6.

(C) Polarimetry measurements of actin filament organization in the circumferential bundles of dorsal and ventral epidermal cells (DV cells) in >2-fold stage embryos expressing the indicated reporters. Scatter plots show the quantification of ψ angle distributions for each reporter. Scatter plots show medians with interquartile range. The number of embryos for each strain is, from left to right: 7, 5, 8. The respective median ψ values are 130, 108, 81°.

(D-J) Polarimetry measurements of actin filament reorganization in the epidermis during embryonic elongation. (D) Scatter plots show the quantification of ψ angle distributions in DV cells and in seam cells in 1.5-fold, 1.5-2-fold and >2-fold stage embryos expressing Af7 as shown in (E), (G), (I). Scatter plots show medians with interquartile range. The number of embryos for each stage is, from left to right: 9, 4, 8, 9, 7, 4. The respective median ψ values are 115, 93, 81, 135, 126, 112°. (E), (G), (I) Representative ρ (left) and ψ (right) stick maps in DV and seam cells in 1.5-fold (E), 1.5-2-fold (G) and >2-fold (I) stage embryos expressing Af7. Insets show zoom-ins of selected ROIs (white outlined boxes) in the respective cell types. Mean ρ and ψ values are shown for each ROI. For all panels, anterior is to the left and dorsal is up. (F), (H), (J) Polar histograms of ρ value distributions in DV and seam cells in 1.5-fold (F), 1.5-2-fold (H) and >2-fold (J) stage embryos expressing Af7. ρ values are represented with respect to the DV/seam boundary (dotted line in (E)): considering that Af7 dipoles are parallel to actin filaments, the more perpendicular mean actin filament orientations are to the boundary, the closer the angle values are to 90° and the narrower the respective distributions. Means ± SD are shown. The number of embryos for each stage and type of cells is as in (D). See also Figure S6.

Figure 7. Polarimetry measurements of actin filament organization in live cellularizing *Drosophila* embryos and live flight muscle expressing selected reporters

(A-B) Polarimetry measurements of actin filament organization in cellularizing *Drosophila* embryos expressing Af7, in the actomyosin rings associated with the invaginating membrane front (A) and at the basal adherens junctions right apicolaterally to the former (B). The left images are summed intensity images of the respective polarimetry stacks. The middle and right images show the measured ρ and ψ angles per pixel, respectively. Insets show zoom-ins of selected ROIs (white outlined boxes)

(C) Flight tests were performed to compare the flight ability of strains expressing different actin-binding reporters (see methods for the assay and for details of the genotypes). The left scheme depicts the flight test assay: depending on their flight ability, flies were scored as "wild-type", "weak flier" or "flightless". The bar graph on the right quantifies the respective percentages for each strain. Reporters were expressed under the control of a muscle-specific driver, Mef2-GAL4, either throughout muscle development (*Mef2>*), or transiently after muscle development (*G80ts Mef2>*). Thirty flies were scored in total for each strain in three independent experiments. The mean percentages of "wild-type", "weak flier" or "flightless" flies are respectively, from left to right: 93,7,0; 90,10,0; 20,17,63; 90,3,7; 13,17,70; 100,0,0; 0,0,100; 97,3,0; 97,0,3; 100,0,0; 100,0,0; 100,0,0; 100,0,0.

(D-G) Polarimetry measurements of actin filament organization in live flight muscle expressing the indicated reporters. All reporters were expressed throughout flight muscle development except Af7 that was expressed transiently at the adult stage after flight muscle development. Scatter plots in (D) show the quantification of ψ angle distributions for each reporter measured as shown in (E)-(G). Scatter plots show medians with interquartile range. One datapoint represents one myofibril (see methods for details). The number of myofibrils for each strain is, from left to right: 70, 140, 190, 100. The respective median ψ values are 157, 114, 134, 74°. Statistical significance was obtained using a non-parametric Mann-Whitney test. The different constructs were compared to Lifeact; **** $P<0.0001$. (E)-(G) Left panels are summed intensity images of the respective polarimetry stacks. Middle and right panels are ρ and ψ stick maps in the flight muscle of the shown hemithorax, respectively. Ten myofibrils (red outlined boxes) were quantified in each field of view. Insets show zoom-ins of a selected sarcomere. The rightmost panels for each strain show examples of polar histograms of ρ value distributions and histograms of ψ value distributions for a single myofibril. Means ± SD are shown. See also Figure S7.

References

Huang, J.Q., Huang, Y.Y., Yu, H.C., Subramanian, D., Padmanabhan, A., Thadani, R., Tao, Y.Q., Tang, X., Wedlich-Soldner, R., and Balasubramanian, M.K. (2012). Nonmedially assembled F-actin cables incorporate into the actomyosin ring in fission yeast. Journal of Cell Biology *199*, 831-847.