Unraveling the Molecular Landscape of Uterine Fibroids, Insights into *HMGA2* and Stem Cell Involvement

Emmanuel N. Paul[1], Tyler J. Carpenter[1], Laura A. Pavliscak[1], Abigail Z. Bennett[1], Maria Ariadna Ochoa-Bernal[1], Asgerally T. Fazleabas[1], Jose M. Teixeira[1*]

[1]Department of Obstetrics, Gynecology and Reproductive Biology, College of Human Medicine, Michigan State University, Grand Rapids, MI 49503, USA

*Corresponding Author:
Jose M. Teixeira, PhD, Michigan State University, 400 Monroe Ave NW, Grand Rapids, MI 49503. Email: teixei15@msu.edu

Acknowledgments:
These studies have been funded by a grant from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NIH R01HD096259) to JMT and the SRI and Bayer Discovery/Innovation Grant to EP.

The authors have no conflicts to declare.

Abstract

Uterine fibroids are prevalent benign tumors in women that exhibit considerable

heterogeneity in clinical presentation and molecular characteristics, necessitating a

deeper understanding of their etiology and pathogenesis. *HMGA2* overexpression has

been associated with fibroid development, yet its precise role remains elusive. Mutations

in fibroids are mutually exclusive and largely clonal, suggesting that tumors originate from

a single mutant cell. We explored a possible role for *HMGA2* overexpression in

differentiated myometrial cells, hypothesizing its potential to induce a stem cell-like or

dedifferentiating phenotype and drive fibroid development. Myometrial cells were

immortalized and transduced with an *HMGA2* lentivirus to produce HMGA2hi cells. *In*

vitro stem cell assays were conducted and RNA from HMGA2hi and control cells and

fibroid-free myometrial and HMGA2 fibroid (HMGA2F) tissues were submitted for RNA-

sequencing. HMGA2hi cells have enhanced self-renewal capacity, decreased

proliferation, and have a greater ability to differentiate into other mesenchymal cell types.

HMGA2hi cells exhibit a stem cell-like signature and share transcriptomic similarities with

HMGA2F. Moreover, dysregulated extracellular matrix pathways are observed in both

HMGA2hi cells and HMGA2F. Our findings suggest that HMGA2 overexpression drives

myometrial cells to dedifferentiate into a more plastic phenotype and underscore a pivotal

role for HMGA2 in fibroid pathogenesis.

Introduction

Uterine fibroids, or leiomyomas, represent the most prevalent benign tumors in women of reproductive age, affecting approximately 70-80% of women during their lifetime (1). These tumors are mainly characterized by extensive extracellular matrix (ECM) deposition and remodeling (2). Despite their high incidence, the etiology and pathogenesis of uterine fibroids remain poorly understood.

Uterine fibroids exhibit considerable heterogeneity in terms of their clinical presentation, growth patterns, and response to therapeutic interventions (3), which suggests different phenotypic subtypes. Recent reports showing molecular characterization of fibroids have provided unprecedented insights into the complex interplay of genetic, epigenetic, and cellular factors contributing to the development and progression of uterine fibroids (4-6) and support the classification of uterine fibroids into distinct subtypes based on their combined molecular and histological characteristics. Mutations identified in various genes, including *MED12*, *FH*, and *COL4A5*, have been associated with distinct subtypes of fibroids, potentially contributing to the diverse clinical manifestations observed in affected individuals (7,8). Unraveling the molecular mechanisms driving the pathogenesis of the various fibroid subtypes is critical for developing targeted therapeutic strategies and personalized treatment approaches.

Another genetic factor associated with fibroid development is overexpression of the high-mobility group AT-hook 2 (*HMGA2*) gene (9-12). *HMGA2* is located on chromosome 12q15 and encodes a non-histone chromatin-binding protein that plays a crucial role in regulating gene expression especially during embryonic development, including stem cell pluripotency, lineage specification, and organogenesis (13). *HMGA2*

transcripts are detected in most embryonic tissues and progressively become absent in

adult tissues, except for the testis, the lung, and the kidney (14). Its dynamic expression

and regulatory functions are tightly orchestrated to ensure proper development, with

aberrations in *HMGA2* expression linked to developmental abnormalities and diseases

(13,15,16). Overexpression of *HMGA2* commonly resulting from the translocation

between chromosome bands 12q15 and 14q24 has been previously described as the

second major subtype of uterine fibroids (HMGA2F), constituting approximately 7% of

fibroid cases (17,18).

The discovery of a potential stem/progenitor cell population within the myometrium

has generated interest in uncovering the involvement of these cells in the initiation and

growth of fibroids (19,20). Indeed, uterine fibroids are thought to be a clonal disease

(21,22), and since most clonal diseases have a single cell origin, the prevailing hypothesis

has been that a dysregulated myometrium stem cell could be the cell of origin for uterine

fibroids (23). However, the molecular mechanisms governing the interplay between

myometrium stem cells and *HMGA2* in uterine fibroids remain largely unexplored. In this

study, we explored an alternative hypothesis regarding the cell origin for uterine fibroids

by postulating that a mutated differentiated myometrial cell could give rise to a fibroid. We

show that *HMGA2* overexpression in myometrium cells induces stem/progenitor cell

properties, suggesting a potential gain of function for the gene in fibroid development.

Such insights may pave the way for the development of novel therapeutic targets that can

effectively address uterine fibroids in a subtype-specific manner.

Methods

Myometrium cell line generation

HEK293T cells were transfected with hTERT plasmid, Addgene #85140 (24) and helper

plasmids (25) to prepare hTERT lentivirus. Myometrial cells were dissociated and isolated

from a fibroid-free myometrium patient (MyoN) as previously described (26) and

incubated at 37°C overnight in growth medium (DMEM/F12, 10% FBS) prior to

transduction with lentiviral preparations. The subsequent day, transduced MyoN cells

were selected with 50 μg/mL of hygromycin for 1 week. Following selection, MyoN-TERT

cells were maintained in medium with 5 μg/mL hygromycin.

HMGA2 **Overexpressing and Control MyoN-TERT Cell Lines**

MyoN-TERT cells were suspended in 1% bovine serum albumin blocking buffer,

incubated with a rabbit anti-human CRIP1 primary antibody (ThermoFisher, #PA5-24643).

Cells were then incubated with an Alexa-647 anti-rabbit secondary antibody and washed

with flow buffer, and resuspended in flow buffer with of 4′,6-diamidino-2-phenylindole

(DAPI). Cells were sorted by the flow cytometry core at Van Andel Research Institute

(VARI) using a FACSymphony S6 cytometer (BD Biosciences) and analyzed with FlowJo

Software (BD Biosciences, version 10.8.1). Lentiviral particles isolated from in HEK293T

cells transfected with either a custom *HMGA2* (VectorBuilder) or empty control plasmid

(VectorBuilder) were used to transduce CRIP1⁻ MyoN-TERT cells as described above.

Selection of successfully transduced MyoN-TERT live cells expressing GFP were sorted

by FACS. *HMGA2* overexpressing cells are called HMGA2hi cells.

Proliferation Assay

HMGA2hi and control cells were plated in triplicate a 96 well plate at 2,000 cells/well, and cell proliferation was recorded every 4 h using an Incucyte SX5 instrument (Sartorius), following the manufacturer's protocols, until cell confluence was reached. Cell proliferation was quantified using GFP and phase contrast images at each time point with the cell-by-cell software module of the instrument. The experiment was repeated three times using three different cell passages.

Colony Formation Assay

Matched passages of HMGA2hi and control cells were plated at 50 cells/cm^2 in triplicate overnight, then switched to MesenPro RS (Thermo Fisher, # 12746012) for 2 to 3 weeks. Cells were cultured in different wells of the same plate, and both cell types were assayed on the same day. Cultures were fixed in 4% paraformaldehyde (PFA) and stained with crystal violet for colony visualization. Images were taken using a Nikon SMZ18 microscope and Ds-Ri1 camera (Nikon Instruments Inc.). Colonies were counted and total surface area was estimated using ImageJ (version 1.53k), and the percent colony-forming units (CFUs) was calculated as (number of colonies/number of cells plated) × 100 and averaged for triplicates.

Differentiation Assays

For osteogenic and adipogenic differentiation, matched passage HMGA2hi and control cells were plated in duplicate at 50% to 80% confluency in growth media overnight and then cultured for 10 days in fresh StemPro Adipogenesis Differentiation (Thermo Fisher

Scientific, #A1007001) or StemPro Osteogenesis Differentiation (Thermo Fisher

Scientific, #A1007201) media according to the manufacturer's instructions. Cells were

cultured in regular growth media to serve as differentiation controls. To assay adipogenic

differentiation, cultures were fixed in 4% PFA and stained using Oil Red O (Sigma,

#01391) according to the manufacturer's instructions. To assay osteogenic differentiation,

cultures were stained for alkaline phosphatase activity using the Alkaline Phosphatase

(AP), Leukocyte kit (Sigma, #86R-1KT) according to the manufacturer's instructions.

3D Culture and Trichrome Staining

60,000 cells of HMGA2hi and control cells, as well as primary MyoN and fibroid cells,

were seeded in scaffold-free 96-well agarose micro-molds (MilliporeSigma, Z764043),

following the protocol outlined by Wiwatpanit et al. (27). Spheroids were cultured for 4

days in growth medium. Following culture, agarose micro-molds were sealed with

agarose and fixed in 4% PFA at 4°C overnight. Subsequently, they were briefly rinsed in

PBS and stored in 70% ethanol at 4°C until processed for standard paraffin embedding

and sectioning at 6 μm. Spheroid sections were stained with trichrome stain according to

the manufacturer's protocol (StatLab #KTTRBPT). Briefly, sections were deparaffinized

with two xylene baths and rehydrated with three absolute ethanol washes. After hydration,

spheroid sections were incubated in Bouin's fluid at room temperature (RT) overnight.

The next day, after rinsing in running tap water, slides were stained with Mayer's

hematoxylin and rinsed in running tap water. Spheroid sections were then immersed in

the one-step trichrome stain and quickly rinsed in running tap water. Slides were

dehydrated through three changes of absolute ethanol, cleared in three changes of

xylenes, and a coverslip was applied using Permount mounting medium (Fisher Scientific

#SP15). Collagen deposition was determined using the color deconvolution ImageJ

plugin (version 1.53k). Relative collagen area was calculated by the ratio of blue stain

(collagen) over the red stain (cytoplasm).

Tissue Collection and *HMGA2* fibroid mutation detection

The use of human tissue specimens was approved by the Spectrum Health Systems and

Michigan State University Institutional Review Boards (MSU IRB Study ID:

STUDY00003101, SR IRB #2017-198) as secondary use of biobank materials. Fibroids

samples (HMGA2F, n = 24) and fibroid-free myometrial (MyoN, n = 32) from pre-

menopausal women (aged 28-55) were obtained following total hysterectomy. All patients

who participated in the study gave consent to donate tissue through the Spectrum Health

Biorepository. Human samples were processed as previously described (4). Briefly,

samples were washed with phosphate-buffered saline, minced, and immersed in

RNAlater (Sigma, Saint Louis, MO, USA) for RNA-seq analyses. *HMGA2* fibroid

(HMGA2F) expression was determined by RT-qPCR using the iScript Advanced cDNA

Synthesis Kit (BioRad, #1725037) for the reverse transcription and *HMGA2* PrimePCR

primer (BioRad, #qHsaCED0021425) for the qPCR. RPL17 was use as a reference gene

with forward ACGAAAAGCCACGAAGTATCTG and reverse

GACCTTGTGTCCAGCCCCAT primers.

RNA Isolation, Library Preparation and Sequencing

Total RNA from HMGA2hi and control cells was isolated from 3 different passages using

an RNeasy mini kit (Qiagen) and stored at −80 °C in nuclease-free water. RNA integrity

values from cells and tissues were determined with an Agilent 2100 Bioanalyzer

(ThermoFisher), and values ≥7.5 were used for library preparation and paired-end (2 ×

100 bp) RNA-sequencing on an Illumina NextSeq 6000 instrument (Illumina). Libraries

were prepared using a Kapa RNA HyperPrep kit with ribosomal reduction, pooled, and

sequenced on flowcells to yield approximately 50–60 million reads/sample. Raw fastq

files were deposited in the NCBI Gene Expression Omnibus (GSEXXXX).

RNA Seq Analysis

RNA-seq reads were trimmed for quality and adapters using TrimGalore (version 0.6.4),

and quality trimmed reads were assessed with FastQC (version 0.11.7). Trimmed reads

were mapped to Homo sapiens genome assembly GRCh38 (hg38) using STAR (version

2.6.0c). Reads overlapping Ensembl annotations (version 99) were quantified with STAR

prior to model-based differential expression analysis using the edgeR-robust method.

Genes with low counts per million (CPM) were removed using the filterByExpr function

from edgeR (28). Scatterplots of two selected principal components were constructed

with the PCAtools R package (version 2.12.0) to verify sample separation prior to

statistical testing. Generalized linear models were used to determine if principal

components were significantly associated with cell type. Differential expression was

calculated by comparing HMGA2hi to control cells and HMGA2F to MyoN tissues and

differentially expressed genes (DEGs) were visualized with a volcano plot generated

using the EnhancedVolcano (version 1.18.0) package in R. Boxplots of the $\log_2$(CPM)

values were generated using the R package ggplot2 (version 3.4.4). Venn diagrams of

the overlapping DEG of the cells (HMGA2hi vs. control) and DEG of tissues (HMGA2F

vs. MyoN) were constructed using the eulerr package (version 7.0.0). Gene set

enrichment and overrepresentation analyses were completed with clusterProfiler

package (version 4.8.2) using MSigDB (29) and DOSE (30) reference databases.

Statistics and Reproducibility

Bioinformatic statistics were performed using the listed packages in R (version 4.3.2).

DEGs in the RNA-seq of HMGA2hi vs. control and the HMGA2F vs. MyoN were identified

as those having a Benjamini–Hochberg FDR corrected p <0.05 (31). Data with unequal

variances were log-transformed, and the homogeneity of variances was verified before

the completion of analyses. Hypergeometric testing was performed using the function

phyper from R. For the colony formation assays, comparison of two means was

performed with a two-sided student t test, and significance was determined at p <0.05

after confirming normal distribution using GraphPad Prism (version 10.2.2).

Results

HMGA2 **overexpression increases myometrium cell plasticity.**

Our initial objective was to examine whether *HMGA2* overexpression enhances stem

cell/progenitor-like properties in a differentiated myometrium cell line. Primary myometrial

cells were immortalized (MyoN-TERT) and subsequently depleted for stem/progenitor

cells via fluorescence-activated cell sorting (FACS) (Figure 1a and 1b), utilizing the cell

surface marker CRIP1, a putative myometrium stem/progenitor cell marker recently

identified in our laboratory (26). Among the MyoN-TERT cells, 73.7% of the MyoN-TERT

were CRIP1$^-$ (Figure 1b) and were sorted for downstream experiments. CRIP1-depleted

MyoN-TERT cells were then transduced with either *HMGA2* (HMGA2hi) or a control virus.

We performed proliferation, colony formation and differentiation assays to determine stem

cell-like properties induced by *HMGA2* overexpression. The proliferation assay revealed

reduced cell proliferation in HMGA2hi cells compared to the control cells (Figure 1c).

Quantitative analysis of the colony formation assay showed a significant 4.3-fold increase

in the number of cells with self-renewal capacity in the HMGA2hi cells (mean = 128)

compared to control cells (mean = 30) (Figure 1d). While both control and HMGA2hi cells

exhibited lipid droplets when cultured in adipogenic media (Figure 1e), indicating their

ability to differentiate into adipocytes, staining in the osteogenic differentiation assay

revealed alkaline phosphatase activity only in HMGA2hi cells and not in control cells when

cultured in osteogenic media (Figure 1f).

Stem cell-like transcriptomic signature is induced by *HMGA2* overexpression in

myometrial cells.

Principal component analysis (PCA) plot of HMGA2hi and control cells revealed sample

separation primarily along principal component (PC) 2, which accounted for 29% of the

variance, indicating that *HMGA2* overexpression induces significant transcriptomic

changes in the immortalized myometrial cells (Figure 2a). Notably, the most prominent

separation observed in PC1, representing 49% of the variance, was associated with cell

passage and was thus incorporated as a co-variable for subsequent analyses. A volcano

plot depicted a total of 709 DEGs with an FDR < 0.05 in HMGA2hi compared to control

cells (Figure 2b). Among these, 304 genes were downregulated, while 405, including

HMGA2 (Log_2FC = 2.2, FDR = 3.4 x 10^{-55}), were upregulated (Table S1). Additionally,

CSPG4, a gene associated with mesenchymal stem cells (26), was significantly

upregulated (Log_2FC = 0.4, FDR = 3.1 x 10^{-2}) in HMGA2hi cells (Figure 2c). However,

other putative myometrial stem/progenitor-related genes such as *PDGFRβ* (32), *SUSD2*

(32), *MCAM* (32), *CD44* (19), and *CD34* (33) did not exhibit significantly (FDR > 0.05)

higher expression in the HMGA2hi cells compared to control cells, except for *ITGA6* (also

known as *CD49f*) (33), which was downregulated (Figure S1). Gene set enrichment

analysis using the MSigDB (29) for cell type signature gene sets (C8) identified 62

significant pathways (FDR <0.05) (Table S2), including two upregulated stem cell gene

sets (Figure 2d).

HMGA2hi cells share a similar transcriptomic profile with HMGA2F.

Disease ontology analysis using DisGeNET database (34) of HMGA2hi compared to

control cells showed enrichment of multiple diseases associated with fibrosis, including

renal fibrosis (Unified Medical Language System: C0151650), chronic kidney disease

(C1561643), lung diseases (C0024115), and fatty liver disease (C4529962) (Figure 3a).

In addition, the uterine fibroids gene set (C0042133) was also enriched by the

overexpression of *HMGA2*. To determine if the transcriptomic profile of HMGA2hi cells is

similar to uterine fibroids, we sequenced RNA from *HMGA2* fibroids (HMGA2F) and

myometrium from non-fibroid patients (MyoN). The transcriptomes of the tissues

separated in PC1 with 41% of variance (Figure 3b) in the PCA plot. 12,929 DEG, 6,788

downregulated and 6,141 upregulated including *HMGA2* (Log$_2$FC = 11.4, FDR = 3.61 x

10^{-34}), were found in HMGA2F compared to MyoN samples (Table S3). A Venn diagram

of the DEGs from the HMGA2hi vs control cells and from the HMGA2F vs MyoN tissues

showed a significant overlap with 61% of the DEGs (n = 434) from the cells also found in

the tissue comparison HMGA2F vs. MyoN (Figure 3c).

***HMGA2*hi cells and HMGA2F tissue had dysregulated extracellular matrix.**

Overrepresentation analysis of gene ontology molecular functions (35) revealed co-

regulated significant pathways enriched in both cells (HMGA2hi compared to control) and

tissue (HMGA2F compared to MyoN). These co-regulated pathways included functions

related to ECM, cell adhesion, stem cell differentiation, and muscle development (Figure

4a). To assess ECM dysregulation in both cell lines and primary cells, 3D culture followed

by trichrome staining analyses were conducted. Spheroids derived from primary fibroid

cells exhibited well-defined round structures, and collagen quantification revealed

significantly higher deposition in fibroid spheroids compared to spheroids from MyoN cells

(Figure 4b and 4c). Similarly, spheroids derived from HMGA2hi cells displayed well-

defined structures and exhibited significantly elevated collagen deposition compared to

their respective controls.

Discussion

The clonal origin of fibroids from a mutated myometrial stem cell is the prevailing

hypothesis for the etiology of the disease (21,22). In this study, we explored an alternative

hypothesis, i.e., that fibroids could arise from a mutated differentiated myometrial cell.

Since elevated *HMGA2* expression in adult tissues is frequently encountered in neoplastic

processes, it is thought to be a key driver of tumorigenesis, but the mechanisms involved

are largely unknown. HMGA2 plays a crucial role in embryonic development and stem

cell maintenance (13). While its expression is generally low in most normal adult tissues

(14), it is notably elevated in some tumors (13,15,16), including HMGA2 fibroids, one of

the major fibroid subtypes (17,18). We exploited these HMGA2 characteristics to

investigate whether its overexpression was sufficient to induce a fibroid-like phenotype in

myometrial cell culture models. We found that overexpression of *HMGA2* induces a more

stem cell-like phenotype in HMGA2hi cells characterized by decreased cell proliferation,

enhanced colony formation, and enhanced differentiation into osteocytes and adipocytes

compared to control cells, all of which support a role for HMGA2 in inducing stem cell

properties in myometrial cells. Whether similar stem cell like properties can be induced in

myometrial cells with other fibroid subtype genetic alterations, for example with MED12

mutation or HMGA1 overexpression, remains to be determined.

Increased colony formation in myometrial cells overexpressing *HMGA2* compared

to control cells has been previously reported (12), but whether the parental cells used in

the study were first depleted of putative myometrial stem cells was not described.

Contradictory findings regarding cell proliferation were also noted with this previous report

and the current study. While our study indicated decreased proliferation, Li, *et al.* reported

a higher Ki67 proliferation index in *HMGA2*-overexpressing cells compared to the control

cells. This discrepancy may be attributed to differences in the methods used to assess

proliferation. In our study, we employed a continuous monitoring approach over a 10-day

period in a 2D model, whereas the study by Li et al. assessed Ki67 staining at a specific

time point in a 3D culture. The significance of this discrepancy is not clear and requires

further investigation.

Transcriptomic analysis of HMGA2hi cells further revealed increased stemness, as

evidenced by a significant increase in the expression of another mesenchymal marker,

CSPG4 (26), along with enrichment for two stem-related pathways. Several studies have

demonstrated the capability of myometrial and fibroid stem/progenitor cells to differentiate

into various cell types (26,32,36). However, to our knowledge, no previous reports have

investigated whether myometrium and fibroid cells depleted for stem/progenitor cells

retain the ability to differentiate into other cell types. Our findings suggest that the

myometrial cell line can differentiate into adipocytes, and notably, can differentiate into

osteocytes only when *HMGA2* is overexpressed, suggesting that *HMGA2*, a gene

expressed by stem cells, enhances cell plasticity in myometrium cells.

In addition to the acquisition of the stem-like properties induced by *HMGA2*

overexpression in the myometrial cell line, we have also uncovered significant

transcriptomic alterations associated with *HMGA2* overexpression, indicating a pivotal

role for HMGA2 in modulating key cellular processes in fibroid cells. HMGA2hi cells

showed a similar transcriptomic profile to HMGA2F based on DEGs. Disease ontology

analysis utilizing the sequencing data unveiled that HMGA2hi cells were enriched for the

uterine fibroids gene set and over 60% of the DEGs in the HMGA2hi cells overlapped

with those identified in HMGA2F tissues. This convergence in transcriptomic signatures

between HMGA2hi cells and HMGA2F tissues underscores the potential role of *HMGA2*

dysregulation in driving fibroid pathogenesis and further supports its involvement in

modulating gene expression patterns associated with fibroid development. Mas *et al*.

reported that overexpression of the truncated form of *HMGA2* in primary myometrial side

population and primary myometrium cells resulted in the formation of leiomyoma-like

tissue with similarities, including ECM content, to human fibroids in a xenotransplantation

in vivo model (11). Dysregulated ECM pathways were also found in the gene ontology

analysis in both HMGA2hi cells and HMGA2F tissues compared to their respective

controls. Additionally, collagen deposition was significantly increased with HMGA2hi cells

in a spheroid model, mimicking observations made using primary fibroid tissue cells

where collagen deposition is a key histological characteristic of uterine fibroids (37).

In conclusion, *HMGA2* emerges as a central player in the pathogenesis of uterine

fibroids, influencing cellular behavior and gene expression profiles. Our study sheds light

on the intricate interplay between HMGA2 and fibroid development, revealing its capacity

to induce a stem cell-like phenotype, alter transcriptomic profiles leading to significant

changes in ECM pathways, and mirror observations in fibroid tissues, all of which

underscores its role in driving fibroid pathogenesis. Further research is needed to

investigate if MED12mt cells exhibit similar characteristics, confirming the theory that the

mutation leads to a stem cell-like phenotype. Overall, our study highlights the significance

of HMGA2 in uterine fibroid biology and lays the groundwork for future studies aimed at

improving clinical management and treatment outcomes for individuals affected by this

condition.

Study approval

The use of human tissue specimens was approved by the Spectrum Health Systems and

Michigan State University Institutional Review Boards (MSU IRB Study ID:

STUDY00003101, SR IRB #2017-198) as secondary use of biobank materials.

Author contributions

Experimental design (ENP, JMT), collected data and performed experiments (ENP, TJC,

LP, AB, MAOB), analyzed data (ENP, JMT), wrote/reviewed manuscript (ENP, TJC, LP,

AB, MAOB, ATF, JMT).

Acknowledgments

We would like to thank the patients who consented for the study, the Spectrum Health

Systems Universal Biorepository staff, and the Van Andel Research Institute Genomics

Core for their help with the study. This work was supported in part through computational

resources and services provided by the Institute for Cyber-Enabled Research at Michigan

State University.

References

1. Stewart EA. Uterine fibroids. *Lancet*. 2001;357(9252):293-298.

2. Borahay MA, Al-Hendy A, Kilic GS, Boehning D. Signaling Pathways in
Leiomyoma: Understanding Pathobiology and Implications for Therapy. *Mol Med*.
2015;21(1):242-256.

3. Stewart EA, Laughlin-Tommaso SK, Catherino WH, Lalitkumar S, Gupta D,
Vollenhoven B. Uterine fibroids. *Nat Rev Dis Primers*. 2016;2:16043.

4. Paul EN, Burns GW, Carpenter TJ, Grey JA, Fazleabas AT, Teixeira JM.
Transcriptome Analyses of Myometrium from Fibroid Patients Reveals Phenotypic
Differences Compared to Non-Diseased Myometrium. *Int J Mol Sci*. 2021;22(7).

5. George JW, Fan H, Johnson B, Carpenter TJ, Foy KK, Chatterjee A, Patterson AL,
Koeman J, Adams M, Madaj ZB, Chesla D, Marsh EE, Triche TJ, Shen H, Teixeira
JM. Integrated Epigenome, Exome, and Transcriptome Analyses Reveal Molecular
Subtypes and Homeotic Transformation in Uterine Fibroids. *Cell Rep*.
2019;29(12):4069-4085 e4066.

6. Paul EN, Grey JA, Carpenter TJ, Madaj ZB, Lau KH, Givan SA, Burns GW,
Chandler RL, Wegienka GR, Shen H, Teixeira JM. Transcriptome and DNA
methylome analyses reveal underlying mechanisms for the racial disparity in
uterine fibroids. *JCI Insight*. 2022;7(20).

7. Mehine M, Kaasinen E, Heinonen HR, Makinen N, Kampjarvi K, Sarvilinna N,
Aavikko M, Vaharautio A, Pasanen A, Butzow R, Heikinheimo O, Sjoberg J,
Pitkanen E, Vahteristo P, Aaltonen LA. Integrated data analysis reveals uterine
leiomyoma subtypes with distinct driver pathways and biomarkers. *Proceedings of*
the National Academy of Sciences of the United States of America.
2016;113(5):1315-1320.

8. Makinen N, Mehine M, Tolvanen J, Kaasinen E, Li Y, Lehtonen HJ, Gentile M, Yan
J, Enge M, Taipale M, Aavikko M, Katainen R, Virolainen E, Bohling T, Koski TA,
Launonen V, Sjoberg J, Taipale J, Vahteristo P, Aaltonen LA. MED12, the mediator
complex subunit 12 gene, is mutated at high frequency in uterine leiomyomas.
Science. 2011;334(6053):252-255.

9. Nilbert M, Heim S, Mandahl N, Floderus UM, Willen H, Mitelman F. Characteristic
chromosome abnormalities, including rearrangements of 6p, del(7q), +12, and
t(12;14), in 44 uterine leiomyomas. *Hum Genet*. 1990;85(6):605-611.

10. Klemke M, Meyer A, Nezhad MH, Bartnitzke S, Drieschner N, Frantzen C, Schmidt
EH, Belge G, Bullerdiek J. Overexpression of HMGA2 in uterine leiomyomas points
to its general role for the pathogenesis of the disease. *Genes Chromosomes*
Cancer. 2009;48(2):171-178.

11. Mas A, Cervello I, Fernandez-Alvarez A, Faus A, Diaz A, Burgues O, Casado M,
Simon C. Overexpression of the truncated form of High Mobility Group A proteins
(HMGA2) in human myometrial cells induces leiomyoma-like tissue formation. *Mol*
Hum Reprod. 2015;21(4):330-338.

12. Li Y, Qiang W, Griffin BB, Gao T, Chakravarti D, Bulun S, Kim JJ, Wei JJ. HMGA2-
mediated tumorigenesis through angiogenesis in leiomyoma. *Fertil Steril*.
2020;114(5):1085-1096.

13. Parisi S, Piscitelli S, Passaro F, Russo T. HMGA Proteins in Stemness and
Differentiation of Embryonic and Adult Stem Cells. *Int J Mol Sci*. 2020;21(1).
14. Vignali R, Marracci S. HMGA Genes and Proteins in Development and Evolution.
Int J Mol Sci. 2020;21(2).
15. Mansoori B, Mohammadi A, Ditzel HJ, Duijf PHG, Khaze V, Gjerstorff MF,
Baradaran B. HMGA2 as a Critical Regulator in Cancer Development. *Genes*
(Basel). 2021;12(2).
16. Li Y, Zhao Z, Xu C, Zhou Z, Zhu Z, You T. HMGA2 induces transcription factor Slug
expression to promote epithelial-to-mesenchymal transition and contributes to
colon cancer progression. *Cancer Lett*. 2014;355(1):130-140.
17. Nibert M, Heim S. Uterine leiomyoma cytogenetics. *Genes Chromosomes Cancer*.
1990;2(1):3-13.
18. Berta DG, Kuisma H, Valimaki N, Raisanen M, Jantti M, Pasanen A, Karhu A,
Kaukomaa J, Taira A, Cajuso T, Nieminen S, Penttinen RM, Ahonen S, Lehtonen
R, Mehine M, Vahteristo P, Jalkanen J, Sahu B, Ravantti J, Makinen N, Rajamaki
K, Palin K, Taipale J, Heikinheimo O, Butzow R, Kaasinen E, Aaltonen LA.
Deficient H2A.Z deposition is associated with genesis of uterine leiomyoma.
Nature. 2021;596(7872):398-403.
19. Mas A, Nair S, Laknaur A, Simon C, Diamond MP, Al-Hendy A. Stro-1/CD44 as
putative human myometrial and fibroid stem cell markers. *Fertil Steril*.
2015;104(1):225-234 e223.
20. Yin P, Ono M, Moravek MB, Coon JSt, Navarro A, Monsivais D, Dyson MT,
Druschitz SA, Malpani SS, Serna VA, Qiang W, Chakravarti D, Kim JJ, Bulun SE.
Human uterine leiomyoma stem/progenitor cells expressing CD34 and CD49b
initiate tumors in vivo. *J Clin Endocrinol Metab*. 2015;100(4):E601-606.
21. Canevari RA, Pontes A, Rosa FE, Rainho CA, Rogatto SR. Independent clonal
origin of multiple uterine leiomyomas that was determined by X chromosome
inactivation and microsatellite analysis. *Am J Obstet Gynecol*. 2005;193(4):1395-
1403.
22. Zhang P, Zhang C, Hao J, Sung CJ, Quddus MR, Steinhoff MM, Lawrence WD.
Use of X-chromosome inactivation pattern to determine the clonal origins of uterine
leiomyoma and leiomyosarcoma. *Hum Pathol*. 2006;37(10):1350-1356.
23. Bulun SE. Uterine fibroids. *N Engl J Med*. 2013;369(14):1344-1355.
24. Hayer A, Shao L, Chung M, Joubert LM, Yang HW, Tsai FC, Bisaria A, Betzig E,
Meyer T. Engulfed cadherin fingers are polarized junctional structures between
collectively migrating endothelial cells. *Nat Cell Biol*. 2016;18(12):1311-1323.
25. Dull T, Zufferey R, Kelly M, Mandel RJ, Nguyen M, Trono D, Naldini L. A third-
generation lentivirus vector with a conditional packaging system. *J Virol*.
1998;72(11):8463-8471.
26. Paul EN, Carpenter TJ, Fitch S, Sheridan R, Lau KH, Arora R, Teixeira JM.
Cysteine-rich intestinal protein 1 is a novel surface marker for human myometrial
stem/progenitor cells. *Commun Biol*. 2023;6(1):686.
27. Wiwatpanit T, Murphy AR, Lu Z, Urbanek M, Burdette JE, Woodruff TK, Kim JJ.
Scaffold-Free Endometrial Organoids Respond to Excess Androgens Associated
With Polycystic Ovarian Syndrome. *J Clin Endocrinol Metab*. 2020;105(3):769-
780.

28. Zhou X, Lindsay H, Robinson MD. Robustly detecting differential expression in RNA sequencing data using observation weights. *Nucleic Acids Res*. 2014;42(11):e91.

29. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;102(43):15545-15550.

471 30. Yu G, Wang LG, Yan GR, He QY. DOSE: an R/Bioconductor package for disease ontology semantic and enrichment analysis. *Bioinformatics*. 2015;31(4):608-609.

473 31. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Statistical Methodology*. 1995;57(1):289-300.

476 32. Patterson AL, George JW, Chatterjee A, Carpenter TJ, Wolfrum E, Chesla DW, Teixeira JM. Putative human myometrial and fibroid stem-like cells have mesenchymal stem cell and endometrial stromal cell properties. *Human reproduction*. 2020;35(1):44-57.

480 33. Ono M, Kajitani T, Uchida H, Arase T, Oda H, Uchida S, Ota K, Nagashima T, Masuda H, Miyazaki K, Asada H, Hida N, Mabuchi Y, Morikawa S, Ito M, Bulun SE, Okano H, Matsuzaki Y, Yoshimura Y, Maruyama T. CD34 and CD49f Double-Positive and Lineage Marker-Negative Cells Isolated from Human Myometrium Exhibit Stem Cell-Like Properties Involved in Pregnancy-Induced Uterine Remodeling. *Biology of reproduction*. 2015;93(2):37.

486 34. Pinero J, Sauch J, Sanz F, Furlong LI. The DisGeNET cytoscape app: Exploring and visualizing disease genomics data. *Comput Struct Biotechnol J*. 2021;19:2960-2967.

489 35. Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou LP, Mi H. PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci*. 2022;31(1):8-22.

492 36. Banerjee S, Xu W, Chowdhury I, Driss A, Ali M, Yang Q, Al-Hendy A, Thompson WE. Human Myometrial and Uterine Fibroid Stem Cell-Derived Organoids for Intervening the Pathophysiology of Uterine Fibroid. *Reprod Sci*. 2022;29(9):2607-2619.

496 37. Leppert PC, Jayes FL, Segars JH. The extracellular matrix contributes to mechanotransduction in uterine fibroids. *Obstet Gynecol Int*. 2014;2014:783289.

Figures

Figure 1. *HMGA2* overexpression increases myometrial cell plasticity. Scatter plot

of the gating strategy used for cell sorting of depleted stem/progenitor MyoN-TERT, using

the unstained sample as control **(a)** and the full stain sample to isolate the depleted cell

population **(b)**. Quantification of the cell proliferation from the cell-by-cell function of the

Incucyte, measuring the area occupied by cells (cm^2/image) for each image scan

normalized to time = 0 h (n = 3) (**c**). The yellow curve represents the control MyoN-TERT

cells, and the blue curve the HMGA2hi cells, error bars represent standard error of the

mean. Representative images of colonies formed by the control and HMGA2hi cells and

plot of colony forming efficiency represented as %CFUs (#CFU/cells seeded × 100) (n

= 3) **(d)**. Representative images of control and HMGA2hi cells grown in control growth

511   media and adipogenic **(e)** or osteogenic **(f)** differentiation media (n = 3). Adipogenic and

512   control cultures were stained with Oil Red O (red color, black arrows), and osteogenic

513   and control cultures were stained for alkaline phosphatase activity (purple color). Scale

514   bar for the adipogenic and osteogenic assays are 500 μm. **p<0.01; by student t-test.

Figure 2. HMGA2hi transcriptome is enriched for stem cell markers and pathways.

Principal component (PC) plot analyses for PC1 vs PC2 shows samples for control and

HMGA2hi cells (n = 3/group) in two-dimensional space. Each dot represents one sample

(a). Volcano plots showing the DEGs from the comparison of HMGA2hi to control cells

(b). Those DEGs with an FDR < 0.05 are depicted as red dots, and the top 10 down- and

up-regulated genes are annotated. Boxplot of two stem cells markers, *HMGA2* (**c**) and

CSPG4 (**d**), that are significantly induced (FDR<0.05) in the HMGA2hi cells compared to

control cells (n = 3 each). Gene expression is shown as log$_2$CPM. Gene set enrichment

analysis (GSEA) using the cell type signature gene sets showing two upregulated stem

526   cell-related pathways in HMGA2hi cells compared to control cells (**e**). Gene count and

527   significance level are shown by the size and color of each circle.

Figure 3. *HMGA2*hi cells have a similar transcriptomic profile with HMGA2 fibroid tissues. Disease-associated gene analysis using the DisGeNET database showing the top 30 significantly enriched pathways (FDR < 0.05) (**a**). Gene count and significance level are shown by the size and color of each circle. Principal component (PC) plot analyses for PC1 vs PC2 show samples from MyoN (n = 32) and HMGA2F tissues (n = 24) in two-dimensional space. Each dot represents one tissue sample (**b**). Venn

536   diagram illustrate the significant (hypergeometric test; $p = 2.2 \times 10^{-22}$) overlap of DEGs

537   between the HMGA2hi vs. control cells and the HMGA2F vs. MyoN tissue samples (**c**).

Figure 4. *HMGA2* **overexpression in MyoN cells modulates extracellular matrix-associated pathways similarly to HMGA2F tissue.** Gene ontology (GO) analysis using the biological processes (BP) gene sets showing the top 25 down- and up-regulated pathways in both cells (HMGA2hi vs. control) and tissues (HMGA2F vs. MyoN) (**a**). Gene ratio and significance level are shown by the size and color of each circle. Representative brightfield and trichrome-stained cross-section images of spheroids from primary cells and cell lines (**b**). The scale bar for the trichrome staining at higher magnitude is 10 µm. Collagen is stained in blue and cytoplasm in red. Relative collagen area, ratio of the

548 blue/red stain, was calculated using the trichrome-stained cross sections from primary

549 cells and cell lines (**c**). *p<0.05, **p<0.01; by student t test.