Structural and Dynamic Impacts of Single-atom Disruptions to Guide RNA Interactions within the Recognition Lobe of *Geobacillus stearothermophilus* Cas9

Helen B. Belato[1,#], Alexa L. Knight[1,#], Alexandra M. D'Ordine[1], Chinmai Pindi,[2] Zhiqiang Fan[2],
Jinping Luo[2], Giulia Palermo,[2] Gerwald Jogl[1], George P. Lisi[1,4]*****

[1]Department of Molecular Biology, Cell Biology & Biochemistry, Brown University,
Providence, RI USA

[2]Departments of Bioengineering and Chemistry, University of California Riverside, Riverside,
CA USA

[3]Brown University Transgenic Mouse & Gene Targeting Facility, Providence, RI USA

[4]Brown University RNA Center, Providence, RI USA

[#]These authors contributed equally to this work
10
11 *** Correspondence:**
12 george_lisi@brown.edu
13
14 **Keywords: CRISPR-Cas9, guide RNA, protein dynamics, NMR, thermophile**
15

Abstract

The intuitive manipulation of specific amino acids to alter the activity or specificity of CRISPR-

Cas9 has been a topic of great interest. As a large multi-domain RNA-guided endonuclease, the

intricate molecular crosstalk within the Cas9 protein hinges on its conformational dynamics, but a

comprehensive understanding of the extent and timescale of the motions that drive its allosteric

function and association with nucleic acids remains elusive. Here, we investigated the structure

and multi-timescale molecular motions of the recognition (Rec) lobe of *Geo*Cas9, a thermophilic

Cas9 from *Geobacillus stearothermophilus.* Our results provide new atomic details about the

*Geo*Rec subdomains (*Geo*Rec1, *Geo*Rec2) and the full-length domain in solution. Two rationally

designed mutants, K267E and R332A, enhanced and redistributed micro-millisecond flexibility

throughout *Geo*Rec, and NMR studies of the interaction between *Geo*Rec and its guide RNA

showed that mutations reduced this affinity and the stability of the ribonucleoprotein complex.

Despite measured biophysical differences due to the mutations, DNA cleavage assays reveal no

functional differences in on-target activity, and similar specificity. These data suggest that guide

RNA interactions can be tuned at the biophysical level in the absence of major functional losses,

but also raise questions about the underlying mechanism of *Geo*Cas9, since analogous single-point

mutations have significantly impacted on- and off-target DNA editing in mesophilic *S. pyogenes*

Cas9. A K267E/R332A double mutant did also did not enhance *Geo*Cas9 specificity, highlighting

the robust tolerance of mutations to the Rec lobe of *Geo*Cas9 and species-dependent complexity

of Rec across Cas9 paralogs. Ultimately, this work provides an avenue by which to modulate the

structure, motion, and guide RNA interactions at the level of the Rec lobe of *Geo*Cas9, setting the

stage for future studies of *Geo*Cas9 variants and their effect on its allosteric mechanism.

Introduction

The vast majority of Cas systems explored as genome editors originate from mesophilic

hosts. The emergence of the thermophilic *Geo*Cas9, with DNA cleavage function up to 85°C, can

expand CRISPR technology to higher temperature regimes and stabilities,[1,2] but its regulatory

mechanism relative to canonical Cas9s must be established. The *Sp*Cas9, which originates from

the mesophilic *Streptococcus pyogenes*, as well as *Geo*Cas9, are both effectors of Type-II CRISPR

systems. Interestingly, the Type II-A *Sp*Cas9 has been by far the most used Cas enzyme, including

in ongoing clinical trials.[3,4] But Cas9 homologs of the Type II-C class, such as *Neisseria meningitis*

(*Nme*Cas9) and *Campylobacter jejuni* (*Cje*Cas9), to which *Geo*Cas9 belongs, have been validated

for mammalian genome editing,[2,5,6] reinforcing the need to better understand this CRISPR class.

The similar domain arrangements of *Geo*Cas9 and *Sp*Cas9 led us to initially speculate that

these could share atomic level mechanistic similarities.[7] *Geo*Cas9 utilizes a guide RNA (gRNA)

to localize and unwind a double-stranded DNA (dsDNA) target after recognition of its

5'NNNNCRAA-3' protospacer adjacent motif (PAM).[2,8] Upon recognition of the PAM sequence

by the PAM-Interacting domain (PI), Cas9-bound guide (gRNA) forms an RNA:DNA hybrid with

the target DNA strand. Initially thought to be part of the PI domain[2], the wedge (WED) domain

recognizes the repeat:anti-repeat region of the gRNA and the dsDNA upstream of the target

region.[9] The Rec lobe of Cas9 is responsible for orienting the RNA:DNA hybrid, as well as the

adjacent nuclease domains, into their active conformations.[10-13] Coordinated cleavage of the target

and non-target DNA strand then occurs via the HNH and RuvC nucleases, respectively.   The

*Geo*Cas9 nuclease active sites within HNH and RuvC are spatially distinct from the PAM

recognition site in the PI domain, necessitating structural and dynamic changes that allosterically

couple dsDNA binding to cleavage. Biochemical[11,14,15] and structural[16,17] experiments using the

extensively studied *Sp*Cas9 have revealed that its function is governed by a sophisticated allosteric

mechanism that transfers gRNA and dsDNA binding information from the Rec lobe to the distal

catalytic sites. A dynamically driven allosteric signal spans the HNH domain of *Sp*Cas9, enabled

by the plasticity of the Rec lobe, which orchestrates the conformational activation required for

DNA cleavage.[14,18] Our prior work revealed a divergence in the timescales of allosteric motions in

the *Sp*Cas9 and *Geo*Cas9 HNH domains[7,17] suggesting an unusually flexible HNH and unique

allosteric mode of regulation for *Geo*Cas9. It is therefore also possible that docking of the gRNA

with *Geo*Cas9, and thus its interaction with the RNA:DNA hybrid, may differ from the *Sp*Cas9

system, as *Geo*Cas9 contains a truncated Rec lobe with only two of the three canonical

subdomains.

71   The high thermal stability and more compact size of *Geo*Cas9 (it is 281 residues shorter

than *Sp*Cas9) can be especially important for *in vivo* delivery applications, since promising viral

vectors (*i.e.* adeno-associated virus, AAV) have cargo capacities of ~4.7kb,[19] which prevents

*Sp*Cas9-gRNA packaging into a single AAV vector but permits "all-in-one" delivery of *Geo*Cas9-

sgRNA.[12] Until the very recent cryo-EM structures of *Geo*Cas9,[20,21] little was known about

specific residues that influence its structure, gRNA binding, or function. Our recent NMR work

with *Sp*Cas9 uncovered pathways of micro-millisecond timescale motions that propagate chemical

information related to allostery and specificity through *Sp*Rec and its RNA:DNA hybrid,[16,17]

prompting us to investigate this phenomenon in *Geo*Rec.

An atomic-level structural understanding of specificity in large multi-domain protein-

nucleic acid complexes like Cas9 is often difficult to address by NMR spectroscopy. Although

dynamic ensembles in DNA repair enzymes have provided some insight,[22] many efforts to improve

Cas9 specificity and reduce off-target activity have relied on large mutational screens[23] or error-

prone PCR[24], which are less intuitive. Inter-subunit allosteric communication between the catalytic

HNH domain and the Rec lobe is critical to Cas9 specificity, as the binding of off-target DNA

sequences at Rec alter HNH dynamics to affect DNA cleavage.[8,25,26] To further probe the

fundamental role of protein motions in the function and specificity of *Geo*Cas9, as well as the

effect of protein-nucleic acid interactions on its structural signatures, we engineered two mutations

in *Geo*Rec (K267E and R332A, housed within *Geo*Rec2). We hypothesized that these variants

could enhance *Geo*Cas9 specificity (*i.e.* limit its off-target cleavage) for two reasons. First, the

chosen mutation sites are homologous to those of specificity-enhancing variants of *Sp*Cas9.[24,27]

Second, altered Cas9-gRNA interactions have been shown to be a consequence of specificity-

enhancement and these charged residues appear to directly interact with the gRNA.[10,11,15,28]

Balancing these two points is the fact that Type-II Cas systems generally have conserved nuclease

domains, but are delineated by highly varied Rec lobes.[12] This implies that the structural and

dynamic properties of Rec may play an outsized role in differentiating the functions of *Sp*Cas9 and

*Geo*Cas9, which may not be identical. Nevertheless, our work provides new insight into the

biophysical, biochemical, and functional role of the *Geo*Rec lobe and how mutations modulate the

domain itself and its interaction with gRNA in full-length *Geo*Cas9.

Results

The structural similarity of GeoRec1, GeoRec2, and GeoRec facilitates NMR analysis of protein

dynamics and RNA affinity

*Geo*Cas9 is a 1087 amino acid polypeptide, thus we employed a "divide and concur"

approach for NMR studies, which we previously showed to be useful for quantifying allosteric

structure and motion in *Sp*Cas9.[16,17,29,30] The *Geo*Rec lobe is comprised of subdomains *Geo*Rec1

and *Geo*Rec2, which likely work together to recognize nucleic acids. We engineered constructs of

the *Geo*Rec1 (136 residues, 16 kDa) and *Geo*Rec2 (212 residues, 25 kDa) subdomains and solved

the X-ray crystal structure of *Geo*Rec2 at 1.49 Å, which aligns remarkably well with the structure

of the *Geo*Rec2 domain within the AlphaFold model (RMSD 1.03 Å) and new cryo-EM structure

of *Geo*Cas9 (RMSD 1.10 Å, **Figure 1A**). We were neither able to crystallize *Geo*Rec1 nor full-

length *Geo*Cas9 in the apo state, but our *Geo*Rec2 crystal structure represents the structure of the

subdomain within the full-length *Geo*Cas9 protein quite well. Our previous studies of *Geo*HNH

also show identical superpositions of X-ray crystal structures with full-length Cas complexes.[7] In

addition to the individual subdomains, we also generated an NMR construct of the intact *Geo*Rec

(370 residues, 43 kDa).

Despite only 22% sequence identity, the structure of *Sp*Rec3 and *Geo*Rec2 are highly

similar (RMSD 2.00 Å, **Figure S1**). The structure of *Geo*Rec1, in contrast, does not align perfectly

with *Sp*Rec1, instead, it partially aligns with both *Sp*Rec1 and *Sp*Rec2 **(Figure S1)**. Thus, the

nearly identical *Sp*Rec3 and *Geo*Rec2 architectures and their intrinsic dynamics may be a common

thread among Type II Cas9s of different size and PAM preference. To capture atomic-level

signatures of *Geo*Rec, we obtained well-resolved ^{1}H-^{15}N NMR fingerprint spectra for all three

protein constructs and assigned the amide backbones (**Figure S2**). ^{1}H-^{15}N amide and ^{1}H-^{13}CH$_3$ Ile,

Leu, and Val (ILV)-methyl NMR spectra (**Figure 1B, S3**) of *Geo*Rec overlay very well with those

of its individual subdomains, suggesting that the linkage of subdomains within the full-length

Figure 1. (A) Arrangement of *Geo*Cas9 domains across the primary sequence. The cryo-EM structure of *Geo*Cas9 in complex with gRNA (PDB: 8JTR) shows poor resolution of HNH. The *Geo*Rec2 domain from PDB: 8JTR (gray) is overlaid with our X-ray structure of *Geo*Rec2 (red, PDB: 9B72). **(B)** ^{1}H^{15}N TROSY HSQC NMR spectrum of *Geo*Rec collected at 850 MHz. Overlays of this spectrum with resonances from spectra of *Geo*Rec1 (black) and *Geo*Rec2 (blue) demonstrate a structural similarity between the isolated subdomains and intact *Geo*Rec.

*Geo*Rec polypeptide does not alter their individual folds. Consistent with this observation, circular

dichroism (CD) thermal unfolding profiles of *Geo*Rec1 ($T_m \sim 34$ °C) and *Geo*Rec2 ($T_m = 61.50$

°C) are distinct and occur as separate events in the unfolding profile of *Geo*Rec (**Figure S4**). The

dumbbell shape of *Geo*Rec, with its two globular subdomains connected by a short flexible linker,

is a likely contributor to these biophysical properties.

Rationally designed GeoRec2 mutants do not substantially impact the GeoRec structure

To understand how the structure and gRNA interactions of *Geo*Cas9 can be modulated at

the level of *Geo*Rec, we engineered two charge-altering point mutants in the *Geo*Rec2 subdomain,

K267E and R332A. Based on the AlphaFold2 model of *Geo*Cas9, both of these residues are < 5Å

from the bound RNA:DNA hybrid and were predicted to interface with the nucleic acids directly

(**Figure 2A/B**). A new experimental cryo-EM structure of *Geo*Cas9 confirmed the interaction

between K267 and the gRNA, but does not report a < 5Å interaction of R332 with the gRNA.[31]

The rationale for our designed mutations was also that removal of positive charge would weaken

the interactions between *Geo*Cas9 and the gRNA, affecting K_d via the electrostatics or dynamics

of the *Geo*Rec lobe. Studies of *Sp*Cas9 revealed that interaction of *Sp*Rec3 (analogous to *Geo*Rec2)

with its RNA:DNA hybrid triggers conformational rearrangements that allow the catalytic HNH

domain to sample its active conformation.[11] Thus, *Sp*Rec3 acts as an allosteric effector that

recognizes the RNA:DNA hybrid to activate HNH. Mismatches (*i.e.* off-target DNA sequences)

in the target DNA generally prevent *Sp*Rec3 from undergoing the full extent of its required

conformational rearrangements, leaving HNH in a "proofreading" state with its catalytic residues

too far from the DNA cleavage site. Off-target DNA cleavage by Cas9 remains an area of intense

study and substantial effort from various groups has gone into mitigating such effects.[15,23,28,32,33]

Indeed, many high-specificity *Sp*Cas9 variants contain mutations within *Sp*Rec3 that increase the

threshold for its conformational activation, reducing the propensity for HNH to sample its active

state in the presence of off-target DNA sequences.[11,15,23] Studies of flexibility within Rec itself, as

well as its gRNA interactions in the presence of mutations, are therefore essential to connecting

biophysical properties to function and specificity in related Cas9s.

The K267E *Geo*Rec2 variant is sequentially and structurally similar to a specificity enhancing

Figure 2. (A, B) Sites of selected mutations within *Geo*Rec2, K267 and R332, are highlighted as purple sticks directly facing the RNA and DNA modeled from *Nme*Cas9 (PDB ID: 6JDV), allowing for prediction of the binding orientation within *Geo*Cas9. NMR chemical shift perturbations caused by the K267E **(C)** or R332A **(D)** mutations are plotted for each residue of *Geo*Rec. Gray bars denote sites of line broadening, the blue bar denotes an unassigned region of *Geo*Rec corresponding to the native Rec1-Rec2 linker, and the red bar indicates the mutation site. The red dashed line indicates 1.5σ above the 10% trimmed mean of the data. Chemical shift perturbations 1.5σ above the 10% trimmed mean are mapped onto K267E **(E)** and R332A **(F)** *Geo*Rec (red spheres). Resonances that have broadened beyond detection are mapped as yellow spheres and the mutation sites are indicated by a black sphere and green arrow.

site in *Sp*Cas9 (K526E), within the evoCas9 system.[27] The *Sp*Cas9 K526E mutation substantially

reduced off-target activity alone, but was even more effective in conjunction with three other

single-point mutations in *Sp*Rec3.[27] The R332A *Geo*Rec2 variant also resembles one mutation

within a high-specificity *Sp*Cas9 variant, an early iteration of HiFi *Sp*Cas9 called HiFi Cas9-

R691A.[24] We assessed mutation-induced changes to local structure in *Geo*Rec via NMR chemical

shift perturbations in ¹H-¹⁵N HSQC backbone amide spectra. Consistent with experiments using

*Geo*Rec2 alone, chemical shift perturbations and line broadening are highly localized to the

mutation sites. (**Figure 2C-F**). Perturbation profiles of the *Geo*Rec2 subdomain and intact *Geo*Rec

also implicate the same residues as sensitive to the mutations (**Figure S5**).

CD spectroscopy revealed that wild-type (WT), K267E, and R332A GeoRec2 maintained

similar alpha-helical secondary structure, though the thermostability of both variants was slightly

reduced from that of WT GeoRec2 (**Figure S6)**. The T_m of WT GeoRec2 is ~62 °C, consistent

with the T_m of the full-length GeoCas9, while that of K267E GeoRec2 was decreased to ~55 °C.

Though the R332A GeoRec2 T_m remains ~62 °C, this variant underwent a smaller unfolding event

near 40 °C before completely unfolding. These data suggest that despite small structural

perturbations, both mutations are destabilizing to GeoRec2, which led us to expect a change in

NMR-detectable protein dynamics.

Mutations enhance and redistribute molecular motions within GeoRec2

Due to the high molecular weight of the intact GeoRec lobe, decays in NMR signal

associated with spin relaxation experiments were significant and hampered data quality. Thus, we

focused on quantifying the molecular motions of the GeoRec2 subdomain, where the K267E and

R332A mutations reside, and the chemical shift perturbations are most apparent. To obtain high-

quality per-residue information representative of GeoRec, we measured longitudinal (R_1) and

transverse (R_2) relaxation rates and heteronuclear ^{1}H-[^{15}N] NOEs (**Figure S7**), then used these data

in a Model-free analysis of per-residue order parameters (S^2). Previous measurements of S^2 across

the adjacent GeoHNH nuclease revealed substantial ps-ns timescale flexibility,[7] leading us to

wonder whether a similar observation would be made for GeoRec2, which abuts GeoHNH. Such

a finding could suggest that HNH-Rec2 crosstalk in GeoCas9 is driven primarily by rapid bond

vector fluctuations. However, unlike GeoHNH, S^2 values for GeoRec2 are globally elevated,

suggesting that the ps-ns motions of this subdomain arise primarily from global tumbling of the

protein in solution. We therefore carried out Carr-Purcell-Meiboom-Gill (CPMG) relaxation

dispersion NMR experiments to assess the flexibility of GeoRec2 on slower timescales, which has

been linked to chemical information transfer in the well-studied *Sp*Cas9.[10,16,17,29] Evidence of μs-

ms motions (*i.e.* curved relaxation dispersion profiles) is observed in 17 residues within the

*Geo*Rec2 core, spanning its interfaces to Rec1 and HNH **(Figure 3B)**. Such motions are

completely absent from *Geo*HNH, thus two neighboring domains, *Geo*Rec2 and *Geo*HNH, diverge

in their intrinsic flexibility (at least in isolation), raising questions about the functional implications

of these motions in *Geo*Rec2. We previously showed that heightened flexibility of *Sp*Rec3 via

specificity-enhancing mutations concomitantly narrowed the conformational space sampled by

*Sp*HNH, highlighting a "motional trade-off" between the domains. Manipulation of the flexibility

Figure 3. (A) CPMG relaxation dispersion profiles of all residues with evidence of μs-ms motion, fit to a global k_{ex} of 147 ± 41 s^{-1} (WT *Geo*Rec2, left), 376 ± 89 s^{-1} (K267E *Geo*Rec2, center), and 142 ± 28 s^{-1} (R332A *Geo*Rec2, right) collected at 25 °C and 600 MHz. Residues are colored in accordance with Table S1. Relaxation dispersion profiles for individual resonances are shown in Figures S8-S10. **(B)** Sites exhibiting CPMG relaxation dispersion in **(A)** are mapped to *Geo*Rec as blue spheres. Adjacent domains within the cryo-EM structure of *Geo*Cas9 are also shown. **(C)** Per-residue NMR order parameters of WT (black), K267E, and R332A (red, separate plots) *Geo*Rec.

of *Sp*Cas9 and *Geo*Cas9 domains by mutagenesis also impacts aspects of nucleic acid binding and

cleavage,[1,10,14,15,29] which led us to investigate similar perturbations in *Geo*Rec2.

Since *Sp*Rec3 and *Geo*Rec2 have similar structures and μs-ms flexibility, we speculated

that charge-altering mutations would modulate the biophysical properties of *Geo*Rec and the

function of *Geo*Cas9, as observed for *Sp*Cas9. We investigated K267E and R332A *Geo*Rec2 with

NMR spin relaxation, as described for WT *Geo*Rec2 (*vide supra*). An analysis of chemical

exchange rates, k_{ex}, derived from dual-field CPMG relaxation dispersion show a global k_{ex} for WT

*Geo*Rec2 of 147 ± 41 s^{-1}. The K267E mutation, which directly contacts the nucleic acids, shifts the

globally fitted k_{ex} to 376 ± 89 s^{-1}, while the R332A variant maintains a global k_{ex} similar to that of

WT *Geo*Rec2 (142 ± 28 s^{-1}) and consistent with its similar thermal stability. The global fit of the

K267E variant is based on CPMG profiles of 33 residues, while that of R332A is derived from 18

residues (**Table S1**). Interestingly, the residues participating in the global motions of both variants

are distinct from those of WT *Geo*Rec2, demonstrating that residue-specific flexibility is

redistributed throughout *Geo*Rec2, which suggests an altered intradomain molecular crosstalk

within the larger *Geo*Rec. Indeed, perturbation to NMR-detectable motions in *Sp*Cas9 rewired its

allosteric signaling and enzymatic function.[16,29] A similar dynamic modulation of *Geo*Cas9 may

fine-tune its DNA cleavage, which has been demonstrated within the *Geo*HNH nuclease[1] and

wedge (WED) domains.[32] We also assessed the ps-ns fluctuations of *Geo*Rec2 variants (a

negligible contribution to the WT *Geo*Rec2 dynamic profile) and calculated order parameters from

R_1, R_2 and ^{1}H-[^{15}N] NOE relaxation measurements (**Figure 3C**). Bond vector fluctuations on the

ps-ns timescale are only locally altered, thus the mutation-induced reshuffling of these motions is

negligible ($<\Delta S^2> \leq 0.1$) and suggests that, like WT *Geo*Rec2, ps-ns motion arises primarily from

global tumbling in solution.

Mutations within GeoRec alter its affinity for guide RNA

The role of the Rec lobe in orienting the RNA:DNA hybrid within Cas9 is crucial to its

function.[10-13] Thus, the structure, motions, and nucleic acid interactions of Rec represent a critical

piece of the Cas9 signaling machinery. Previous studies of *Sp*Cas9 revealed that gRNA binding to

the Rec lobe induces a global structural rearrangement of the protein that positions the adjacent

HNH into its "proofreading" state,[15] after which target DNA binding positions the nucleases into

active conformations for cleavage.[15,34] We wondered if the atomistic details of the apo *Geo*Cas9-

to-RNP transition could be captured by NMR using the *Geo*Rec construct. In our previous studies,

we used an *in vitro* DNA cleavage assay with *Geo*Cas9 and a 141nt gRNA containing a 21nt spacer

targeting the mouse *Tnnt2* gene locus.[1] Since this assay was already established, we utilized the

same gRNA sequence. However, truncating this gRNA was necessary to optimize binding studies

for NMR analysis. We focused on the 5' end of the gRNA, which includes the spacer sequence,

based on the *Geo*Cas9 AlphaFold2 model and structural data from *Nme*Cas9 and *Sp*Cas9 showing

interactions between the Rec lobe and this region of the gRNA. The subsequent cryo-EM structure

of *Geo*Cas9 corroborated this interaction.[20] Initial attempts using a truncated 101nt gRNA resulted

in poor NMR spectra. An overlay of the ^{1}H-^{15}N HSQC NMR spectra of apo *Geo*Rec and *Geo*Rec-

RNP at a 1:1 molar ratio showed extensive line broadening (**Figure S11**), likely due to the large

size of the complex (75.5 kDa). To mitigate this issue, a 39nt gRNA containing the 21bp spacer

sequence was selected for its ability to maintain the NMR signal while being long enough to

interact fully with the Rec lobe, as suggested by prior structures. When bound to *Geo*Rec, this

complex is 55.6 kDa and a ^{1}H-^{15}N NMR spectral overlay of apo *Geo*Rec and the domain bound to

39nt gRNA shows clear, resolved resonances with significant chemical shift perturbations and line

broadening (**Figure 4A/B, S11**). The strongest chemical shift perturbations are localized to the

*Geo*Rec2 subdomain that interfaces with the RNA:DNA hybrid at the PAM distal end, where

previous studies of specificity-enhancing variants of *Sp*Cas9 have identified alterations in nucleic

acid binding to *Sp*Rec3.[16] It is not known whether specific residues at the PAM distal binding

interface of *Geo*Rec2 play a similar role. Line broadening is evident in both *Geo*Rec1 and

*Geo*Rec2, primarily localized to the RNA:DNA hybrid interface revealed in recent *Geo*Cas9

structures. Microscale thermophoresis (MST) experiments quantified the affinity of *Geo*Rec for

gRNA, producing a $K_d = 3.3 \pm 1.5$ µM that is consistent with the concentration-dependent NMR

chemical shift perturbations (**Figure 5A**).

247   To understand how the K267E and R332A mutants impact gRNA binding to *Geo*Rec, we

248   conducted gRNA titration experiments via NMR and observed that chemical shift perturbations

Figure 4. (A) NMR chemical shift perturbations caused by gRNA binding to WT, K267E, and R332A *Geo*Rec. Gray bars denote sites of line broadening, and the blue bar denotes an unassigned region of *Geo*Rec corresponding to the flexible Rec1-Rec2 linker. The red dashed line indicates 1.5σ above the 10% trimmed mean of the data. **(B)** Representative NMR resonance shifts caused by titration of 39nt gRNA into WT *Geo*Rec. **(C)** NMR titration of 39nt gRNA into K267E (top) and R332A (bottom) *Geo*Rec. The left panel of each pair demonstrates that minimal change in NMR chemical shift or resonance intensity is apparent at gRNA concentrations mimicking the WT titration. The right panel of each pair depicts the titration over a three-fold wider concentration range of gRNA, where shifts and line broadening are visible. Representative resonances are colored by increasing gRNA concentration in the legend.

were attenuated in both variants, relative to WT *Geo*Rec. Despite this muted structural effect, the

impact from gRNA-induced line broadening remains substantial in the *Geo*Rec1 subdomain. Our

NMR data revealed that a three-fold greater concentration of gRNA was required to induce the

maximal structural and dynamic effects in the variants than is required for WT *Geo*Rec (**Figure**

4A/C), suggesting that the variants have a reduced gRNA affinity. MST experiments showed

statistically significant reductions in gRNA affinity for the K267E and R332A constructs, relative

to WT *Geo*Rec, where K267E *Geo*Rec produced a $K_d = 7.2 \pm 1.0$ µM and R332A *Geo*Rec

produced a $K_d = 7.2 \pm 1.5$ µM (**Figure 5B**). The ~3-fold decrease in K_d may also be due, in part,

to a change in the binding mode of the gRNA, such as a faster k_{off}. Collectively, these data reveal

that mutations within *Geo*Rec primarily alter its structure around the mutation site with weaker

distal effects, but more significantly impact protein dynamics and in turn, the gRNA interaction.

NMR experiments also demonstrate that the presence of gRNA impacts both subdomains of

*Geo*Rec, providing a significant structural interface for additional molecular tuning of nucleic acid

binding.

To investigate the impact of gRNA binding on *Geo*Rec2 in greater detail, we conducted

Figure 5. Representative MST-derived profiles of WT **(A)**, K267E, and R332A **(B)** *Geo*Rec binding to a Cy5-labeled 39nt gRNA, yielding $K_d = 3.3 \pm 1.5$ µM, $K_d = 7.2 \pm 1.0$ µM and $K_d = 7.2 \pm 1.5$ µM, respectively. Bar graphs comparing K_d values across n ≥ 3 replicate samples are shown for Tnnt2 gRNA **(C)** and 8UZA gRNA from a recent cryo-EM structure **(D)**. *$p < 0.05$, **$p < 0.004$

NMR titration experiments using the isolated domain, which yielded even clearer NMR spectra.

Figure S12 shows NMR spectra of WT, K267E, and R332A *Geo*Rec2 overlaid with their

corresponding gRNA-bound spectra (39nt Tnnt2 RNA). At protein:RNA molar ratios used for full-

length *Geo*Rec studies, the WT *Geo*Rec2 spectrum exhibited significant line broadening across

the *Geo*Rec2 sequence. Plots of NMR peak intensities (I_{bound}/I_{free}) show substantial resonance

intensity losses (**Figure S12**), with many residues likely in the intermediate exchange regime, in

addition to the assumed changes in rotational correlation of the domain. In comparison, spectra of

the gRNA-bound K267E and R332A *Geo*Rec2 variants showed less pronounced signal decay at

the same levels of titrant, retaining nearly double the I_{bound}/I_{free} ratio across these spectra (**Figure**

S12). These data are consistent with the results of NMR experiments with the 43 kDa *Geo*Rec,

supporting the premise that *Geo*Rec2 mutations weaken its interaction with gRNA. The less

crowded NMR spectrum of isolated *Geo*Rec2 facilitated the resolution of distinct structural

features that explain the impact of the mutations on gRNA binding (**Figure S9**). For example,

residue I53 adopts a similar conformation in gRNA-bound WT and K267E *Geo*Rec2 but assumes

a different structural state in R332A. Conversely, residue R25 populates a WT-like structure in

gRNA-bound R332A *Geo*Rec2, unlike K267E. Additionally, two resonances are observed for

residue K71 in the gRNA-bound R332A NMR spectrum, indicating real-time equilibration

between two structural states. This effect is unique to the R332A variant and underscores subtle

structural and dynamic changes to *Geo*Rec during gRNA binding.

We further examined the NMR data to attempt to identify residues most critical for gRNA

binding to *Geo*Rec. In an overlay of the WT and mutant gRNA-induced chemical shift

perturbations ($\Delta\delta$, **Figure S13**), it became clear that the effect of gRNA binding to *Geo*Rec variants

was muted, where even at saturating concentrations, the chemical shift perturbations across the

K267E and R332A *Geo*Rec2 sequences were weaker than those of same residues in WT *Geo*Rec.

The residual $\Delta\delta$ (WT - mutant) was plotted (**Figure S13**), where positive values indicate that

residues in a *Geo*Rec variant are weakly affected by gRNA, relative to WT. Negative residual $\Delta\delta$

denote sites where *Geo*Rec variants experience a greater structural impact from gRNA than

corresponding sites in WT. Of particular interest are the positive residuals that hint at the sites in

*Geo*Rec most critical for tight gRNA binding. These residues were mapped onto the *Geo*Rec

structure (**Figure S13**) and termed allosteric hotspots, as many are not at the RNA interface.

Mutations of these hotspots in future studies offers a potential means of precisely tuning the

affinity of *Geo*Rec to gRNA. Notably, residues with positive residual Δδ (suggested as critical for

tight RNA binding) largely overlap in the analysis of both variants. Specifically, residues F170,

R192, H264, R269, L270, L279, H300, D301, E368, D376, D403, E405, E408, and I429 appear

as allosteric hotspots (with CPMG relaxation dispersion) critical to WT-like gRNA interaction.

299   Having observed a reduced affinity of *Geo*Rec variants for gRNA by NMR and MST, we

next quantified the impact of the K267E and R332A mutations on RNP formation and stability in

full-length *Geo*Cas9. The thermal unfolding midpoint of full-length WT *Geo*Cas9 determined by

CD is ~60 °C and the K267E and R332A mutations do not change the T_m of the apo protein (**Figure**

S14). Upon formation of an RNP, the T_m of WT *Geo*Cas9 increases to 73°C. K267E *Geo*Cas9

retains a similar T_m increase to 70 °C, while R332A *Geo*Cas9 forms a less stable RNP with T_m of

61 °C. The trend of these data is consistent with NMR and MST, which highlight that although

K267E and R332A mutations within *Geo*Rec have somewhat muted structural effects, these

changes alter protein dynamics and the interaction with gRNA.

Mutations in full-length GeoCas9 alter its structural dynamics and interaction with gRNA

309   To further investigate the effects of mutations on protein dynamics, we performed

molecular dynamics (MD) simulations based on the cryo-EM structure of full-length *Geo*Cas9

(PDB: 8UZA) in complex with gRNA and target DNA. We simulated the full-length WT *Geo*Cas9

and its K267E and R332A mutants as well as a double mutant combining K267E and R332A

(**Figure 6A**), in three replicates of approximately 2 μs each. Multi-microsecond simulations

revealed substantial changes in the dynamics of the GeoCas9 mutants compared to the WT (**Figure**

S15). Specifically, we observed that mutations in the REC domain significantly altered the

Figure 6: Effects of mutations in full-length *Geo*Cas9 revealed by MD simulations. **(A)** The structure of *Geo*Cas9 (PDB: 8UZA, protein in gray) bound to gRNA (orange) and DNA (magenta) is shown. Mutations studied include K267E (red), R332A (blue), and the 10 mutations of i*Geo*Cas9 (lime green), all highlighted in surface representation. **(B)** Differential root-mean-square fluctuations (ΔRMSF) of protein residues computed between WT *Geo*Cas9 and the K267E (red), R332A (blue), and double mutant (pink). **(C)** Distribution of protein-RNA contacts for WT and *Geo*Cas9 variants computed over the 6 μs simulation ensemble. **(D)** Comparison of gRNA binding free energy to the Rec domain in WT *Geo*Cas9 and variants. **(E)** Representative snapshots from MD simulations illustrating structural changes in Rec-gRNA association in WT *Geo*Cas9 (left) and variants (right).

dynamics of both the Rec and the adjacent HNH (**Figure S15**). Differential root-mean-square

fluctuations (ΔRMSF) analysis of protein residues between the WT and variants further

highlighted these alterations, showing increased dynamics in the HNH and Rec domains induced

by the mutations (**Figure 6B**). To quantify the impact of these mutations, we analyzed protein-

RNA interactions by calculating the number of contacts between *Geo*Cas9 and gRNA in the WT

and mutant systems. A contact was defined as a distance between two atoms of ≤ 4.5 Å. The

number of contacts was significantly reduced in all variants compared to the WT, with the most

pronounced reduction observed in the K267E variant (**Figure 6C**). We next quantified gRNA-Rec

domain binding by calculating the binding free energy using the MM-GBSA method over ~200 ns

of stable simulation trajectories (details in Materials and Methods). Consistent with a reduction in

gRNA contacts, variants with the K267E mutation exhibited a substantial reduction in binding free

energy (>60 kcal mol^{-1}) relative to the WT, whereas the R332A variant displayed a smaller

reduction (<20 kcal mol^{-1}, **Figure 6D**). Notably, protein-DNA interactions remained largely

unaffected, suggesting that these mutations do not impair *Geo*Cas9 DNA cleavage ability.

Additionally, we simulated a novel variant, i*Geo*Cas9 (PDB: 8UZB), containing mutations

in the Rec1 and WED domains (**Figure 6A**, mutations highlighted in lime green). This variant was

recently demonstrated to have enhanced specificity in genome-editing.[21] Intriguingly, i*Geo*Cas9

exhibited increased dynamics in the HNH and Rec domains, along with a reduction in gRNA

binding free energy similar to the K267E and R332A mutants. These results suggest a distinct

allosteric pathway involving additional residues that enables i*Geo*Cas9 to maintain improved DNA

cleavage activity despite reduced gRNA binding affinity. In fact, i*Geo*Cas9 samples the greatest

conformational space of any variant tested (**Figure S15**), suggesting a high level of flexibility is

critical to enhanced specificity in *Geo*Cas9. Collectively, our simulations reveal that mutations

K267E and R332A destabilize the *Geo*Cas9 interaction with gRNA, consistent with NMR

observations. Furthermore, the enhanced dynamics and altered binding affinities observed in

i*Geo*Cas9 indicate potential allosteric mechanisms that optimize its genome-editing functionality,

where K267E and R332A evoke a similar, but lesser degree of biophysical change.

DNA cleavage assays suggest the highly stable GeoCas9 is resistant to functional changes by

K267E or R332A mutations

The dynamic impact of the *Geo*Rec mutations and their altered gRNA interactions at the

biophysical level led us to speculate that either mutation incorporated into full-length *Geo*Cas9

would also alter its DNA cleavage function, especially at elevated temperatures where WT

*Geo*Cas9 is most active. Temperature-dependent functional alterations were previously observed

for single-point mutations within *Geo*HNH.[1] Although the K267E and R332A mutations slightly

diminished on-target DNA cleavage by *Geo*Cas9, the effect was very subtle and these overall

cleavage activities followed the temperature dependence of WT *Geo*Cas9 quite closely (**Figure**

S16).

To assess the impact of the K267E and R332A mutations on *Geo*Cas9 specificity, we

assayed the propensity for off-target cleavage using DNA substrates with mismatches 5-6 or 19-

20 base pairs from the PAM site (**Figure S17, Table S2**). As a control for on- and off-target

activity, we assayed WT *Sp*Cas9 alongside the widely used high-specificity HiFi-*Sp*Cas9 variant[24]

(**Figure S17, Table S3**) and found a lower percent of digested off-target (mismatched) DNA

sequences when compared to WT *Sp*Cas9. As expected, WT *Geo*Cas9 was increasingly sensitive

to mismatched target sequences closer to the seed site, which has been demonstrated with *Sp*Cas9

and other Cas systems.[2,6,8,35] No significant differences in activity were observed with digestion

durations ranging from 1-60 minutes,[2] implying that a 1-minute digestion is sufficient for *in vitro*

activity of *Geo*Cas9 with the target DNA template. While these findings generally align with prior

investigations of off-target DNA cleavage,[2,6,8] there are nuanced differences. Specifically, a

previous study reported ~10% cleavage of off-target DNA with a mismatch 5-6 base pairs from

the PAM by WT *Geo*Cas9.[2] Our results showed nearly 50% cleavage for the same off-target

mismatch, but still a significant decrease in cleavage from on-target or 19-20 base pair distal

mismatches. This could be due to the relatively high RNP concentrations (600-900 nM) in our

assay (for clear visibility on the gel), compared to prior studies with RNP concentrations $\leq$ 500

nM.[2] Our results corresponded closely to those of prior studies with a 19-20 base pair mismatch,

where off-target cleavage is tolerated by WT *Geo*Cas9.[2] Single-point mutants K267E and R332A

*Geo*Cas9 have negligible impact on *Geo*Cas9 specificity (both variants follow the trend of WT

*Geo*Cas9, **Figure S17**), which contrasts prior work with *Sp*Cas9 that demonstrated robust

specificity enhancement with single-point mutations in Rec.[24,27] Additionally, a *Geo*Cas9 double

mutant K267E/R332A exhibits decreased on-target cleavage efficiency, which has been noted in

high-specificity Cas systems.[15,23,24,27,36-38] However, the additive effect of the K267E/R332A

double mutant still does not enhance *Geo*Cas9 specificity in our assay. MST-derived binding

affinities using full-length Tnnt2 gRNA and full-length WT, K267E, or R332A *Geo*Cas9 indicate

that all three proteins have similar affinities for the gRNA used in the functional assays (**Figure**

7). Thus, mutations do not substantially alter full-length *Geo*Cas9 binding to Tnnt2 gRNA,

supporting similar cleavage activities for these proteins. We repeated the MST experiments using

a different gRNA sequence derived from the new cryo-EM structure of *Geo*Cas9 (PDB:8UZA),

which has a different spacer sequence. We observed a similar trend, as all *Geo*Cas9 variants

exhibited comparable affinities for this gRNA. Our functional studies illustrate an apparent

resilience of *Geo*Cas9 to major functional changes at the level of Rec, despite comparable

mutations having profound functional impacts in mesophilic Cas9s.

Figure 7. (A) Representative MST-derived profiles of WT, K267E, and R332A *Geo*Cas9 binding to a Cy5-labeled full-length Tnnt2 gRNA. **(B)** Bar graph comparing K_d values across n ≥ 3 replicate samples are shown for Tnnt2 and 8UZA gRNA from a recent cryo-EM structure of *Geo*Cas9. *$p < 0.01$

Discussion

CRISPR-Cas9 is a powerful tool for targeted genome editing with high efficiency and

modular specificity.[2,15,24,27] Allosteric signals propagate DNA binding information to the HNH and

RuvC nuclease domains, facilitating their concerted cleavage of double-stranded DNA.[8,10,14,15] The

intrinsic flexibility of the nucleic acid recognition lobe plays a critical role in this information

transfer, exerting a measure of conformational control over catalysis.[10] This study provides new

insights into the structural, dynamic, and functional role of the thermophilic *Geo*Cas9 recognition

lobe. Novel constructs of subdomains *Geo*Rec1 and *Geo*Rec2, as well as intact *Geo*Rec show a

high structural similarity to the domains in full-length *Geo*Cas9, facilitating solution NMR

experiments that captured the intrinsic allosteric motions across *Geo*Rec2. These studies revealed

the existence of μs-ms timescale motions that are classically associated with allosteric signaling

and enzyme function, which span the entire *Geo*Rec2 domain to its interfaces with *Geo*Rec1 and

the adjacent *Geo*HNH domain.

Based on homology to specificity-enhancing variants of the better studied *Sp*Cas9, the

biophysical and biochemical consequences of two mutations were tested in *Geo*Rec2, the larger

*Geo*Rec lobe, and full-length *Geo*Cas9. We speculated that removing positively charged residues

with potential to interact with negatively charged nucleic acids could disrupt *Geo*Cas9-gRNA

complex formation, stability, and subsequent function by altering the protein or nucleic acid

motions. Indeed, CPMG relaxation dispersion experiments revealed that mutations enhanced and

reorganized the μs-ms flexibility of *Geo*Rec2. Further, NMR titrations showed the affinity of

K267E and R332A *Geo*Rec for gRNA to be weaker than that of WT *Geo*Rec, consistent with

MST-derived K_d values using the isolated domain. The mutations also diminished the stability of

the full-length *Geo*Cas9 RNP complex.

The collective changes to protein dynamics, gRNA binding, and RNP thermostability

suggested that mutations could modulate *Geo*Cas9 function, as observed in similar studies of

*Sp*Cas9 reporting that gRNA dynamics, affecting the potential for the RNA:DNA hybrid to

dissociate, have affected function.[11,15,39] Yet, the functional impact of single-point and double

mutations in this work were negligible, despite homologous K-to-E and R-to-A single point

mutations enhancing specificity of the mesophilic *Sp*Cas9. The biophysical impact of mutations

within *Geo*Rec2 and *Geo*Rec may be tempered by its evolutionary resilience and the highly stable

neighboring domains in the context of full-length *Geo*Cas9, reflected in an unchanged affinity for

target DNA once the RNP was formed.[40,41] Thus, a greater number of additive (or synergistic)

mutations within *Geo*Rec would be required to fine-tune activity or specificity to a large degree.

It should be noted that the effects of these and other *Geo*Rec mutations may vary *in vivo*

or with alternative target cleavage sites and cell types. Such studies will be the subject of future

work, as will biochemical assays of homologous mutations across diverse Cas9s, which have

contributed to the wide use of CRISPR technology.[42] We also note that despite the homology

between *Geo*Rec2 and *Sp*Rec3 and the latter's role in evo- and HiFi-*Sp*Cas9 variants that inspired

the K267E and R332A mutations, the maximally enhanced *Sp*Cas9 variants contain four mutations

each. Presumably each individual substitution plays a small role modulating specificity. However,

there is no consistent pattern that discerns whether multiple mutations will have additive or

synergistic impacts on Cas9 function. NMR and MD studies of high-specificity *Sp*Cas9 variants

(HF-1, Hypa, and Evo, each with distinct mutations in the *Sp*Rec3 domain) reveal universal

structural and dynamic variations in regions of *Sp*Rec3 that interface with the RNA;DNA hybrid.[16]

Notably, a recently published variant, i*Geo*Cas9,[32] demonstrated enhanced genome-editing

capabilities in HEK293T cells with eight mutations, though none in the Rec2 subdomain. This

study highlighted the functional adaptability of i*Geo*Cas9 under low magnesium conditions, a trait

beneficial in mammalian cells, distinguishing it from WT *Geo*Cas9. These very recently published

data, as well as the findings reported here still advance our molecular understanding of the

functional handles in *Geo*Cas9, relevant to the design of new enhanced variants.

This study marks the first phase of mapping allosteric motions and pathways of information

flow in the *Geo*Rec lobe with solution NMR experiments. Such information transfer is critical to

the crosstalk between Rec and HNH in several Cas9s. Despite NMR advancements in

perdeuteration,[43] transverse relaxation-optimized spectroscopy (TROSY),[44] and sparse isotopic

labeling,[45] per-residue dynamics underlying allosteric signaling in large multi-domain proteins

such as *Geo*Cas9 (~126 kDa) have remained challenging to characterize. Novel cryo-EM

structures of *Geo*Cas9[32] will facilitate the merging of future NMR and MD simulation studies to

report on RNP dynamics and atomic level networks of communication. The identification of

additional (or synergistic) allosteric hotspots within *Geo*Rec using an integrated workflow will

help to further resolve the balance between structural flexibility and the unusually high stability of

*Geo*Cas9, leading to new insight into targeted manipulation of RNA affinity and enhanced

variants.

Here, we set out to biophysically characterize the Rec lobe of *Geo*Cas9 to obtain new

understanding of its function (in the context of well-studied mesophilic Cas9s). Using an

AlphaFold2 model, and later a cryo-EM structure of *Geo*Cas9, we introduced mutations based on

proximal gRNA interactions and homology to specificity-enhancing sites in *Sp*Cas9. However, the

mutations did not affect *Geo*Cas9 function as expected, highlighting the complicated interplay

between the biophysics of mesophilic and thermophilic Cas enzymes and the difficulty of applying

universal functional predictions to Cas9. The very recent report of the i*Geo*Cas9 variant further

reinforces this point.[21] While high-specificity *Sp*Cas9 variants are heavily mutated in Rec3

(analogous to *Geo*Rec2), i*Geo*Cas9 lacks mutations in Rec2 entirely, raising new questions about

the functional role of *Geo*Rec. MD simulations of WT *Geo*Cas9, i*Geo*Cas9, and the Rec variants

revealed that while K267E and R332A induce dynamic effects on a similar trajectory to i*Geo*Cas9,

a true high-specificity variant samples a very wide conformational space with displacements of

both Rec and HNH (**Figures 6** and **S15**). Through further study of the fundamental mechanism of

*Geo*Cas9, it remains possible that engineering of *Geo*Rec may produce high-specificity variants.

Materials and Methods

Expression and purification of *Geo*Rec1, *Geo*Rec2, *Geo*Rec, and *Geo*Cas9

The *Geo*Rec1 (residues 90-225) and *Geo*Rec2 (residues 245-456) subdomains, as well as

the entire *Geo*Rec lobe (residues 90-456) of *G. stearothermophilus* Cas9 were engineered into a

pET28a vector with a N-terminal His$_6$-tag and a TEV protease cleavage site. The K267E and

R332A mutations were separately introduced into the *Geo*Rec2 plasmid. Plasmids were

transformed into BL21 (DE3) cells (New England Biolabs). Protein samples for CD spectroscopy,

MST, and functional assays were grown in Lysogeny Broth (LB, Fisher), while isotopically

labeled samples for NMR were grown in M9 minimal media (deuterated for *Geo*Rec2 and

*Geo*Rec) containing $CaCl_2$, $MgSO_4$, MEM vitamins, and 1.0 g/L ^{15}N ammonium chloride

and 2.0 g/L ^{13}C glucose (Cambridge Isotope Laboratories), as the sole nitrogen and carbon sources,

respectively. Cells were induced with 1 mM IPTG after reaching an OD_{600} of 0.8−1.0 and grown

for 4 hours at 37 °C post induction. The cells were harvested by centrifugation, resuspended in a

buffer of 50 mM Tris-HCl, 250 mM NaCl, 5 mM imidazole, and 1 mM PMSF at pH 7.4, lysed by

ultrasonication, and purified by Ni−NTA affinity chromatography. Following TEV proteolysis of

the terminal His-tag, the samples were further purified on a Superdex75 size exclusion column.

NMR samples were dialyzed into a buffer containing 20 mM NaPi, 80mM KCl, 1mM DTT, and

1mM EDTA at pH 7.4.

The full-length *Geo*Cas9 plasmid was acquired from Addgene (#87700), expressed in TB

media and was expressed and purified as previously described.[2] The K267E, R332A, and

K267E/R332A variants were introduced into full-length *Geo*Cas9 by modifying the original

plasmid acquired from Addgene.

NMR spectroscopy

Backbone resonance assignments of *Geo*Rec1 and *Geo*Rec2 were carried out on a Bruker

Avance NEO 600 MHz spectrometer at 25 °C. The following triple resonance experiments were

collected for each sample: ^{1}H-^{15}N TROSY-HSQC, HNCA, HN(CO)CA, HN(CA)CB,

HN(COCA)CB, HN(CA)CO and HNCO. All spectra were processed in NMRPipe[46] and analyzed

in Sparky[47]. Three-dimensional correlations and assignments were made in CARA[48] and *Geo*Rec1

and *Geo*Rec2 backbone assignments were deposited in the BMRB under accession numbers 52363

and 51197, respectively. Backbone resonance assignments of *Geo*Rec were completed by

transferring assignments from the individually assigned spectra of *Geo*Rec1 and *Geo*Rec2, as done

previously for other large Cas9 fragments.[49,50]

NMR spin relaxation experiments were carried out in a temperature-compensated manner

at 600 and 850 MHz on Bruker Avance NEO and Avance III HD spectrometers, respectively.

CPMG experiments were adapted from the report of Palmer and coworkers[51] with a constant

relaxation period of 20 ms and ν_{CPMG} values of 0, 25, 50, 75, 100, 150, 250, 500, 750, 800, 900,

and 1000 Hz. Exchange parameters were obtained from global fits of the data carried out with

RELAX[52] using the R2eff, NoRex, and CR72 models, as well as in-house fitting in GraphPad

Prism with the following models:

Model 1: No exchange

502
$$R_2^{eff} = R_2^0 \tag{1}$$

503
504   Model 2: Two-state, fast exchange (Meiboom equation [53])
505
$$R_2^{eff} = R_2^0 + \frac{\Phi}{k_{ex}}\left[1 - \frac{4\nu_{CPMG}}{k_{ex}}\tan h\left(\frac{k_{ex}}{4\nu_{CPMG}}\right)\right] \tag{2}$$

Global fitting of CPMG profiles was determined to be superior to individual fits based on the

Akaike Information Criterion.[54] Uncertainties in these rates were determined from replicate spectra

with duplicate relaxation delays of 0, 25, 50 (×2), 75, 100, 150, 250, 500 (×2), 750, 800 (×2), 900,

and 1000 Hz.

Longitudinal and transverse relaxation rates were measured with randomized T_1 delays of

0, 20, 60, 100, 200, 600, 800, and 1200 ms and T_2 delays of 0, 16.9, 33.9, 50.9, 67.8, 84.8, and

101.8 ms. Peak intensities were quantified in Sparky and the resulting decay profiles were analyzed

in Sparky with errors determined from the fitted parameters. Uncertainties in these rates were

determined from replicate spectra with duplicate relaxation delays of 20 (x2), 60 (x2), 100, 200,

600 (x2), 800, and 1200 ms for T_1 and 16.9, 33.9 (x2), 50.9 (x2), 67.8 (x2), 84.8, 101.8 (x2) ms

for T_2. Steady-state ^{1}H-[^{15}N] NOE were measured with a 6 second relaxation delay followed by a

3 second saturation (delay) for the saturated (unsaturated) experiments and calculated by I_{sat}/I_{ref}.

All relaxation experiments were carried out in a temperature-compensated interleaved manner.

Model-free analysis was carried out by fitting relaxation rates to five different forms of the

spectral density function with local τ_m, spherical, prolate spheroid, oblate spheroid, or ellipsoid

diffusion tensors.[55-60] The criteria for inclusion of resonances in the diffusion tensor estimate was

based on the method of Bax and coworkers.[61] N-H bond lengths were assumed to be 1.02 Å and

the ^{15}N chemical shift anisotropy tensor was -160 ppm. Diffusion tensor parameters were

optimized simultaneously in RELAX under the full automated protocol.[52] Model selection was

iterated until tensor and order parameters did not deviate from the prior iteration.

NMR titrations were performed on a Bruker Avance NEO 600 MHz spectrometer at 25 °C

by collecting a series of ^{1}H-^{15}N TROSY HSQC spectra with increasing ligand (*i.e.* gRNA)

concentration. The ^{1}H and ^{15}N carrier frequencies were set to the water resonance and 120 ppm,

respectively. Samples of WT, K267E, and R332A *Geo*Rec were titrated with gRNA until no

further spectral perturbations were detected. NMR chemical shift perturbations were calculated as:

532
$$\Delta\delta = \sqrt{(\Delta\delta_{HN}^2 + \Delta\delta_{NH}^2/25)/2}$$

Microscale thermophoresis (MST)

MST experiments were performed on a Monolith X instrument (NanoTemper

Technologies), quantifying WT, K267E, R332A, and K267E/R332A *Geo*Rec binding to a 39-nt

Cy5-labeled gRNA at a concentration of 20 nM in a buffer containing 20 mM sodium phosphate,

150 mM KCl, 5 mM $MgCl_2$, and 0.1% Triton X-100 at pH 7.6. The *Geo*Rec proteins were serially

diluted from a 200 μM stock into 16 microcentrifuge tubes and combined in a 1:1 molar ratio with

serially diluted gRNA from a 40 nM stock. After incubation for 5 minutes at 37 °C in the dark,

each sample was loaded into a capillary for measurement. K_d values for the various complexes

were calculated using the MO Control software (NanoTemper Technologies). Statistical

significance was calculated using a two-tailed T-test.

Circular dichroism (CD) spectroscopy

All *Geo*Cas9 and *Geo*Rec proteins were buffer exchanged into a 20 mM sodium phosphate buffer at pH 7.5, diluted to 1 µM, and loaded into a 2 mm quartz cuvette (JASCO instruments). A CD spectrum was first measured between 200 - 250 nm, after which the sample was progressively heated from 20 – 90 °C in 1.0 °C increments while ellipticity was monitored at 222 and 208 nm. Phosphate buffer baseline spectra were subtracted from the sample measurements. Prior to CD measurements, *Geo*Cas9-RNP was formed by incubating 3 µM *Geo*Cas9 with gRNA at a 1:1.5 molar ratio at 37 °C for 10 minutes. The unfolding CD data was fit in GraphPad Prism to:

$$Ellipticity\ (T) = \frac{[(m_f T + b_f) + (m_u T + b_u)] exp\left[\left(-\frac{\Delta H_{D,vH}}{R}\right)\left(\frac{1}{T} - \frac{1}{T_m}\right)\right]}{1 + exp\left[\left(-\frac{\Delta H_{D,vH}}{R}\right)\left(\frac{1}{T} - \frac{1}{T_m}\right)\right]}$$

X-ray crystallography

*Geo*Rec protein purified as described above was crystallized by sitting drop vapor diffusion at room temperature by mixing 1.0 µL of 15 mg/mL *Geo*Rec in a buffer of 20 mM HEPES and 100 mM KCl at pH 7.5 with 2.0 µL of crystallizing condition: 0.15 M calcium chloride, 15 % polyethylene glycol 6000, 0.1 M HEPES at pH 7.0. Crystals were cryoprotected in crystallizing condition supplemented with 30% ethylene glycol. Diffraction images were collected at the NSLS-II AMX beamline at Brookhaven National Laboratory under cryogenic conditions. Images were processed using XDS[62] and Aimless in CCP4.[63] Chain A of the *N. meningitidis* Cas9 X-ray structure (residues 249-445 only, PDB ID: 6JDQ) was used for molecular replacement with Phaser followed by AutoBuild in Phenix.[64] Electron density was only observed for the *Geo*Rec2 subdomain. The *Geo*Rec2 structure was finalized through manual building in Coot[65] and refinement in Phenix.

Molecular dynamics (MD) simulations

Molecular Dynamics (MD) simulations were based on the cryo-EM structure of full-length

*Geo*Cas9 (PDB: 8UZA, resolution 3.17 Å) in complex with gRNA and target DNA with two

mutations in *Geo*Cas9 (at residues 8 and 582). Four systems were considered for the MD studies:

WT, K267E, R332A, K267E/R332A and i*Geo*Cas9. We generated the WT *Geo*Cas9 by back-

mutating A8D and A582H from the cryo-EM structure (PDB: 8UZA), followed by introducing the

mutations K267E, R332A, or a double mutation (with both K267E and R332A) for the variant

systems. Subsequently, we performed MD simulation of i*Geo*Cas9 (PDB: 8UZB, resolution 2.63

Å) consisting of 10 mutations (D8A, E149G, T182I, N206D, P466Q, H582A, Q817R, E843K,

E854G, K908R). All systems were solvated with explicit water in a periodic box of ~ 134 Å x ~

154 Å x ~ 151 Å resulting in ~ 276,000 atoms. Counter ions were added to neutralize the systems.

MD simulations were performed using a protocol tailored for protein-nucleic acid complexes,[66]

previously applied in studies of CRISPR-Cas systems.[67-69] All the simulations were performed by

using Amber ff19SB force field for protein,[70] ff99bsc1 corrections and χOL3 corrections for DNA

and RNA, respectively.[71,72] Water molecules were described by TIP3P model.[73] All bonds

involving hydrogens were constrained using the LINCS algorithm. A particle mesh Ewald method

(PME) with a 10 Å cutoff was used to calculate electrostatics. Energy minimization was performed

to relax the water molecules and counterions, keeping the protein-nucleic acid complex fixed with

harmonic potential restraints of 100 kcal/mol Å^2. Equilibration was performed by gradually

increasing the temperature from 0 to 100 K and then to 200 K in canonical NVT ensemble and

isothermal-isobaric NPT ensemble. A final temperature of 300 K was maintained via Langevin

dynamics with a collision frequency γ = 1/ps and a reference pressure of 1 atm was achieved

through Berendsen barostat. Production runs were carried out in NVT ensemble for 2 μs for each

system in three replicates, resulting in 6 μs per system (totaling 30 μs for all systems). The

equations of motion were integrated with the leapfrog Verlet algorithm with a time step of 2 fs.

All simulations were conducted using the GPU-empowered version of AMBER 22.[74] Analysis was

performed on the aggregate ensemble (i.e., ~6 µs per system).

To characterize the protein-nucleic acid interactions in all the systems under investigation,

we performed contact analysis. A contact was considered to form between two atoms within a

cutoff distance of ≤4.5 Å. The binding free energy of gRNA and DNA with *Geo*Cas9 was

calculated using the Molecular Mechanics Generalized Born Surface Area (MM-GBSA)

method.[75-77] This approach was used to compare the Rec-gRNA binding affinity of WT *Geo*Cas9

with its mutants. For each system, the binding energies were calculated over the ~200 ns ensemble

of the stable trajectories at an interval of ~20 ns.

DNA cleavage assays

*Geo*Cas9 gRNA templates containing 21-nt spacers targeting the mouse *Tnnt2* gene locus

were introduced into EcoRI and BamHI sites in pUC57 (Genscript). The plasmid was transformed

into BL21(DE3) cells (New England BioLabs) and subsequent restriction digest of the plasmid

DNA was carried out using the BamHI restriction enzyme (New England BioLabs) according to

the manufacturer's instructions. Linearized plasmid DNA was immediately purified using the

DNA Clean and Concentrator-5 kit (Zymo Research) according to the manufacturer's instructions.

RNA transcription was performed *in vitro* with the HiScribe T7 High Yield RNA Synthesis Kit

(New England BioLabs). DNA substrates containing the target cleavage site (479 base pairs,

Figure S13) were produced by polymerase chain reaction (PCR) using mouse genomic DNA as a

template   and   primer   pairs   5'CAAAGAGCTCCTCGTCCAGT3'   and   5'

ATGGACTCCAGGACCCAAGA3' followed by a column purification using the NucleoSpin®

Gel and PCR Clean-up Kit (Macherey-Nagel). For the *in vitro* activity assay, RNP formation was

achieved by incubating 3 µM *Geo*Cas9 (WT, K267E, R332A, or K267E/R332A mutant) and 3

µM gRNA at either 37 °C, 60 °C, 75 °C, or 85 °C for 30 minutes in a reaction buffer of 20 mM

Tris, 100 mM KCl, 5 mM $MgCl_2$, 1 mM DTT, and 5% glycerol at pH 7.5. The 10 µL cleavage

reactions were set up by mixing RNP at varying concentrations with 149 nanograms of PCR

products on ice followed by incubation at 37 °C for 30 minutes. The reaction was quenched with

1 µL of proteinase K (20 mg/mL) and subsequent incubation at 56 °C for 10 minutes. 6x DNA

loading buffer was added to each reaction and 10 µL of reaction mixture per lane was loaded onto

an agarose gel. DNA band intensity measurements were carried out with ImageJ.

For *in vitro* off-target activity assays, RNP formation was achieved by incubating 10 µM

*Geo*Cas9 (WT, K267E, R332A, or K267E/R332A mutant) and 10 µM gRNA at 37 °C for 30

minutes in the reaction buffer described above. The 10 µL cleavage reactions were set up by

mixing 1 µM RNP with 150 nanograms of PCR products (off-target DNA sequences listed in

Table S2) on ice followed by incubation at 37 °C for varying time points. The reaction was

quenched with 1 µL of proteinase K (20 mg/mL) and subsequent incubation at 56 °C for 10

minutes. 6x DNA loading buffer was added to each reaction and 10 µL of reaction mixture per

lane was loaded onto an agarose gel. DNA band intensity measurements were carried out with

ImageJ. WT and HiFi *Sp*Cas9 control proteins were purchased from Integrated DNA Technologies

(IDT, cat. No. 108158 and No. 108160, respectively), as was the associated *Sp*Cas9 gRNA, Alt-

R™ CRISPR-Cas9 gRNA, with an RNA spacer sequence complementing 5'-

TGGACAGAGCCTTCTTCTTC-3'. The on-target and off-target DNA sequences used for the

*Sp*Cas9 *in vitro* cleavage assay can be found in **Table S3**.

Author Contributions

HBB and **ALK** produced *Geo*Rec2, *Geo*Rec1, *Geo*RecFL, and *Geo*Cas9 proteins and

sgRNA, conducted the NMR and biophysical experiments, analyzed the data, and wrote the

original draft of the manuscript. **AMD** solved the X-ray crystal structure of *Geo*Rec2. **CP** carried

out MD simulations and analyzed the data. **ZF** and **JL** conducted *Geo*Cas9 functional assays and

analyzed the data. **GP** supervised the MD studies and obtained funding. **GJ** supervised collection

of X-ray crystallographic data. **GPL** conceived the study, supervised collection of NMR

spectroscopic data, obtained funding, and wrote the original draft. The final manuscript was

written and edited with contributions from all authors.

Acknowledgments

This work was supported by NIH grant R01 GM 136815 (to GP and GPL) and NSF grant

MCB 2143760 (to GPL). GP acknowledges support from the NIH (Grant No. R01GM141329) and

the NSF (CHE- 2144823), as well as from the Sloan Foundation (FG-2023-20431) and the Camille

and Henry Dreyfus Foundation (TC-24-063). This research used the AMX beamline of the

National Synchrotron Light Source II, a U.S. Department of Energy (DOE) Office of Science User

Facility operated for the DOE Office of Science by Brookhaven National Laboratory under

Contract No. DE-SC0012704. The Center for BioMolecular Structure (CBMS) is primarily

supported by NIGMS through a Center Core P30 Grant (P30 GM133893), and by the DOE Office

of Biological and Environmental Research (KP1607011). Computational studies were carried out

using Expanse at the San Diego Supercomputing Center through allocation MCB160059 and

Bridges2 at the Pittsburgh Supercomputer Center through allocation BIO230007 from the

Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program,

which is supported by NSF grants #2138259, #2138286, #2138307, #2137603, and #2138296.

References

Belato, H. B. *et al.* Disruption of electrostatic contacts in the HNH nuclease from a thermophilic Cas9 rewires allosteric motions and enhances high-temperature DNA cleavage. *The Journal of Chemical Physics* **157**, 225103, doi:10.1063/5.0128815 (2022).

Harrington, L. B. *et al.* A thermostable Cas9 with increased lifetime in human plasma. *Nat Commun* **8**, 1424, doi:10.1038/s41467-017-01408-4 (2017).

Zhang, S. *et al.* Current Trends of Clinical Trials Involving CRISPR/Cas Systems. *Front. Med.* **10**, DOI: 10.3389/fmed.2023.1292452 (2023).

Li, T. *et al.* CRISPR/Cas Therapeutics: Progress and Prospects. *Signal Trans. Targ. Ther.* **8**, DOI: 10.1038/s41392-41023-01309-41397 (2023).

Kim, E. *et al.* In vivo genome editing with a small Cas9 orthologue derived from Campylobacter jejuni. *Nat Commun* **8**, 14500, doi:10.1038/ncomms14500 (2017).

Lee, C. M., Cradick, T. J. & Bao, G. The Neisseria meningitidis CRISPR-Cas9 System Enables Specific Genome Editing in Mammalian Cells. *Mol Ther* **24**, 645-654, doi:10.1038/mt.2016.8 (2016).

Belato, H. B. *et al.* Structural and dynamic insights into the HNH nuclease of divergent Cas9 species. *J Struct Biol* **214**, 107814, doi:10.1016/j.jsb.2021.107814 (2021).

Jinek, M. *et al.* A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816-821, doi:10.1126/science.1225829 (2012).

Eggers, A. R. *et al.* Rapid DNA unwinding accelerates genome editing by engineered CRISPR-Cas9. *Cell* **187**, 3249-3261.e3214, doi:10.1016/j.cell.2024.04.031 (2024).

Palermo, G. *et al.* Key role of the REC lobe during CRISPR-Cas9 activation by 'sensing', 'regulating', and 'locking' the catalytic HNH domain. *Q Rev Biophys* **51**, doi:10.1017/s0033583518000070 (2018).

Dagdas, Y. S., Chen, J. S., Sternberg, S. H., Doudna, J. A. & Yildiz, A. A conformational checkpoint between DNA binding and cleavage by CRISPR-Cas9. *Sci Adv* **3**, eaao0027, doi:10.1126/sciadv.aao0027 (2017).

Mir, A., Edraki, A., Lee, J. & Sontheimer, E. J. Type II-C CRISPR-Cas9 Biology, Mechanism, and Application. *ACS Chem Biol* **13**, 357-365, doi:10.1021/acschembio.7b00855 (2018).

Jiang, F., Zhou, K., Ma, L., Gressel, S. & Doudna, J. A. STRUCTURAL BIOLOGY. A Cas9-guide RNA complex preorganized for target DNA recognition. *Science* **348**, 1477-1481, doi:10.1126/science.aab1452 (2015).

Sternberg, S. H., LaFrance, B., Kaplan, M. & Doudna, J. A. Conformational control of DNA target cleavage by CRISPR-Cas9. *Nature* **527**, 110-113, doi:10.1038/nature15544 (2015).

Chen, J. S. *et al.* Enhanced proofreading governs CRISPR-Cas9 targeting accuracy. *Nature* **550**, 407-410, doi:10.1038/nature24268 (2017).

Skeens, E. *et al.* High-fidelity, hyper-accurate, and evolved mutants rewire atomic-level communication in CRISPR-Cas9. *Sci Adv* **10**, eadl1045, doi:10.1126/sciadv.adl1045 (2024).

East, K. W. *et al.* Allosteric Motions of the CRISPR-Cas9 HNH Nuclease Probed by NMR and Molecular Dynamics. *Journal of the American Chemical Society*, doi:10.1021/jacs.9b10521 (2020).

18   Palermo, G., Miao, Y., Walker, R. C., Jinek, M. & McCammon, J. A. Striking Plasticity
of CRISPR-Cas9 and Key Role of Non-target DNA, as Revealed by Molecular
Simulations. *ACS Cent Sci* **2**, 756-763, doi:10.1021/acscentsci.6b00218 (2016).
19   Wu, Z., Yang, H. & Colosi, P. Effect of genome size on AAV vector packaging. *Mol
Ther* **18**, 80-86, doi:10.1038/mt.2009.255 (2010).
20   Shen, P. *et al.* Structure of *Geobacillus stearothermophilus* Cas9: Insights into the
Catalytic Process and Thermostability of CRISPR-Cas9. *ACS Catalysis* **14**, 13227-13235
(2024).
21   Eggers, A. R. *et al.* Rapid DNA Unwinding Accelerates Genome Editing by Engineered
CRISPR-Cas9. *Cell* **187**, 3249-3261 (2024).
22   Lisi, G. P., East, K. W., Batista, V. S. & Loria, J. P. Altering the allosteric pathway in
IGPS suppresses millisecond motions and catalytic activity. *Proc Natl Acad Sci U S A*
114, E3414-e3423, doi:10.1073/pnas.1700448114 (2017).
23   Slaymaker, I. M. *et al.* Rationally engineered Cas9 nucleases with improved specificity.
Science **351**, 84-88, doi:10.1126/science.aad5227 (2016).
24   Vakulskas, C. A. *et al.* A high-fidelity Cas9 mutant delivered as a ribonucleoprotein
complex enables efficient gene editing in human hematopoietic stem and progenitor cells.
Nat Med **24**, 1216-1224, doi:10.1038/s41591-018-0137-0 (2018).
25   Guzman, I., Ghaemi, Z., Baranger, A., Luthey-Schulten, Z. & Gruebele, M. Native
Conformational Dynamics of the Spliceosomal U1A Protein. *The Journal of Physical
Chemistry B* **119**, 3651-3661, doi:10.1021/jp511760m (2015).
26   Doudna, J. A. & Charpentier, E. Genome editing. The new frontier of genome
engineering with CRISPR-Cas9. *Science* **346**, 1258096, doi:10.1126/science.1258096
(2014).
27   Casini, A. *et al.* A highly specific SpCas9 variant is identified by in vivo screening in
yeast. *Nat Biotechnol* **36**, 265-271, doi:10.1038/nbt.4066 (2018).
28   Ricci, C. G. *et al.* Deciphering Off-Target Effects in CRISPR-Cas9 through Accelerated
Molecular Dynamics. *ACS Cent Sci* **5**, 651-662, doi:10.1021/acscentsci.9b00020 (2019).
29   Nierzwicki, L. *et al.* Enhanced specificity mutations perturb allosteric signaling in
CRISPR-Cas9. *Elife* **10**, doi:10.7554/eLife.73601 (2021).
30   Nierzwicki, Ł. *et al.* Principles of target DNA cleavage and the role of Mg2+ in the
catalysis of CRISPR–Cas9. *Nature Catalysis* **5**, 912-922, doi:10.1038/s41929-022-
00848-6 (2022).
31   Shen, P. *et al.* Structure of Geobacillus stearothermophilus Cas9: Insights into the
Catalytic Process and Thermostability of CRISPR-Cas9. *ACS Catalysis* **14**, 13227-13235,
doi:10.1021/acscatal.4c03278 (2024).
32   Eggers, A. R. *et al.* Rapid DNA unwinding accelerates genome editing by engineered
CRISPR-Cas9. *bioRxiv*, 2023.2012.2014.571777, doi:10.1101/2023.12.14.571777
(2023).
33   Doudna, J. A. The promise and challenge of therapeutic genome editing. *Nature* **578**,
229-236, doi:10.1038/s41586-020-1978-5 (2020).
34   Palermo, G. *et al.* Protospacer Adjacent Motif-Induced Allostery Activates CRISPR-
Cas9. *J Am Chem Soc* **139**, 16028-16031, doi:10.1021/jacs.7b05313 (2017).
35   Hou, Z. *et al.* Efficient genome engineering in human pluripotent stem cells using Cas9
from Neisseria meningitidis. *Proceedings of the National Academy of Sciences of the
United States of America* **110**, 15644-15649, doi:10.1073/pnas.1313587110 (2013).

36   Kleinstiver, B. P. *et al.* High-fidelity CRISPR-Cas9 nucleases with no detectable
749   genome-wide off-target effects. *Nature* **529**, 490-495, doi:10.1038/nature16526 (2016).
750   37   Hu, J. H. *et al.* Evolved Cas9 variants with broad PAM compatibility and high DNA
751   specificity. *Nature* **556**, 57-63, doi:10.1038/nature26155 (2018).
752   38   Lee, J. K. *et al.* Directed evolution of CRISPR-Cas9 to increase its specificity. *Nat*
753   *Commun* **9**, 3048, doi:10.1038/s41467-018-05477-x (2018).
754   39   Ma, E., Harrington, L. B., O'Connell, M. R., Zhou, K. & Doudna, J. A. Single-Stranded
755   DNA Cleavage by Divergent CRISPR-Cas9 Enzymes. *Mol Cell* **60**, 398-407,
756   doi:10.1016/j.molcel.2015.10.030 (2015).
757   40   Nguyen, V. *et al.* Evolutionary drivers of thermoadaptation in enzyme catalysis. *Science*
758   **355**, 289-294, doi:10.1126/science.aah3717 (2017).
759   41   Saavedra, H. G., Wrabl, J. O., Anderson, J. A., Li, J. & Hilser, V. J. Dynamic allostery
760   can drive cold adaptation in enzymes. *Nature* **558**, 324-328, doi:10.1038/s41586-018-
761   0183-2 (2018).
762   42   Shmakov, S. *et al.* Diversity and evolution of class 2 CRISPR-Cas systems. *Nat Rev*
763   *Microbiol* **15**, 169-182, doi:10.1038/nrmicro.2016.184 (2017).
764   43   Venters, R. A., Farmer, B. T., 2nd, Fierke, C. A. & Spicer, L. D. Characterizing the use
765   of perdeuteration in NMR studies of large proteins: 13C, 15N and 1H assignments of
766   human carbonic anhydrase II. *Journal of molecular biology* **264**, 1101-1116,
767   doi:10.1006/jmbi.1996.0699 (1996).
768   44   Pervushin, K., Riek, R., Wider, G. & Wüthrich, K. Attenuated T2 relaxation by mutual
769   cancellation of dipole-dipole coupling and chemical shift anisotropy indicates an avenue
770   to NMR structures of very large biological macromolecules in solution. *Proceedings of*
771   *the National Academy of Sciences of the United States of America* **94**, 12366-12371,
772   doi:10.1073/pnas.94.23.12366 (1997).
773   45   Tugarinov, V., Kanelis, V. & Kay, L. E. Isotope labeling strategies for the study of high-
774   molecular-weight proteins by solution NMR spectroscopy. *Nat Protoc* **1**, 749-754,
775   doi:10.1038/nprot.2006.101 (2006).
776   46   Delaglio, F. *et al.* NMRPipe: a multidimensional spectral processing system based on
777   UNIX pipes. *J Biomol NMR* **6**, 277-293, doi:10.1007/bf00197809 (1995).
778   47   Lee, W., Tonelli, M. & Markley, J. L. NMRFAM-SPARKY: enhanced software for
779   biomolecular NMR spectroscopy. *Bioinformatics* **31**, 1325-1327,
780   doi:10.1093/bioinformatics/btu830 (2015).
781   48   Keller, R. L. J. *Optimizing the process of nuclear magnetic resonance spectrum analysis*
782   *and computer aided resonance assignment* Doctoral Thesis thesis, ETH Zürich, (2005).
783   49   Nerli, S., De Paula, V. S., McShan, A. C. & Sgourakis, N. G. Backbone-independent
784   NMR Resonance Assignments of Methyl Probes in Large Proteins. *Nat. Commun.* **12**,
785   691 (2021).
786   50   De Paula, V. S., Dubey, A., Arthanari, H. & Sgourakis, N. G. Dynamic Sampling of a
787   Surveillance State Enables DNA Proofreading by Cas9. *Cell Chem. Biol.*
788   **doi.org/10.1016/j.chembiol.2024.10.001** (2024).
789   51   Loria, J. P., Rance, M. & Palmer, A. G. A Relaxation-Compensated
790   Carr−Purcell−Meiboom−Gill Sequence for Characterizing Chemical Exchange by NMR
791   Spectroscopy. *Journal of the American Chemical Society* **121**, 2331-2332,
792   doi:10.1021/ja983961a (1999).

793   52   Bieri, M., d'Auvergne, E. J. & Gooley, P. R. relaxGUI: a new software for fast and
794   simple NMR relaxation data analysis and calculation of ps-ns and μs motion of proteins.
795   *J Biomol NMR* **50**, 147-155, doi:10.1007/s10858-011-9509-1 (2011).

796   53   Luz, Z. & Meiboom, S. Nuclear Magnetic Resonance Study of the Protolysis of
797   Trimethylammonium Ion in Aqueous Solution—Order of the Reaction with Respect to
798   Solvent. *The Journal of Chemical Physics* **39**, 366-370, doi:10.1063/1.1734254 (1963).

799   54   Cavanaugh, J. E. & Neath, A. A. The Akaike Information Criterion: Background,
800   Derivation, Properties, Application, Interpretation, and Refinements. *WIREs Comp. Mol.*
801   *Sci.* **11**, e1460 (2019).

802   55   Brüschweiler, R., Liao, X. & Wright, P. E. Long-range Motional Restrictions in a
803   Multidomain Zinc-finger Protein from Anisotropic Tumbling. *Science* **268**, 886-889
804   (1995).

805   56   Mandel, A. M., Akke, M. & Palmer, A. G. Backbone Dynamics of Escherichia coli
806   Ribonuclease HI: Correlations with Structure and Function in an Active Enzyme. *J. Mol.*
807   *Biol.* **246**, 144-163 (1995).

808   57   Fushman, D., Cahill, S. & Cowburn, D. The Main-chain Dynamics of the Dynamin
809   Pleckstrin Homology (PH) Domain in Solution: Analysis of 15N Relaxation with
810   Monomer/Dimer Equilibration. *J. Mol. Biol.* **266**, 173-194 (1997).

811   58   Orekhov, V. Y., Korzhnev, D. M., Diercks, T., Kessler, H. & Arseniev, A. S. 1H-15N
812   NMR Dynamic Study of an Isolated α-helical Peptide (1-36)bacteriorhodopsin Reveals
813   the Equilibrium Helix-coil Transitions. *J. Biomol. NMR* **14**, 345-356 (1999).

814   59   Korzhnev, D. M., Billeter, M., Arseniev, A. S. & Orekhov, V. Y. NMR Studies of
815   Brownian Tumbling and Internal Motions in Proteins. *Prog. NMR Spectros.* **38**, 197-266
816   (2001).

817   60   Zhuravleva, A. V. *et al.* Gated Electron Transfers and Electron Pathways in Azurin: A
818   NMR Dynamic Study at Multiple Fields and Temperatures. *J. Mol. Biol.* **342**, 1599-1611
819   (2004).

820   61   Tjandra, N., Feller, S. E., Pastor, R. W. & Bax, A. Rotational Diffusion Anisotropy of
821   Human Ubiquitin from N-15 NMR Relaxation. *J. Am. Chem. Soc.* **117**, 12562-12566
822   (1995).

823   62   Kabsch, W. XDS. *Acta Crystallogr D Biol Crystallogr* **66**, 125-132,
824   doi:10.1107/s0907444909047337 (2010).

825   63   Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta*
826   *Crystallogr D Biol Crystallogr* **67**, 235-242, doi:10.1107/s0907444910045749 (2011).

827   64   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
828   electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-877,
829   doi:10.1107/s2059798319011471 (2019).

830   65   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
831   *Acta Crystallogr D Biol Crystallogr* **66**, 486-501, doi:10.1107/s0907444910007493
832   (2010).

833   66   Sinha, S., Pindi, C., Ahsan, M., Arantes, P. R. & Palermo, G. Machines on Genes through
834   the Computational Microscope. *J. Chem. Theor. Comput.* **19**, 1945-1964 (2023).

835   67   Saha, A. *et al.* An Alpha-helical Lid Guides the Target DNA Toward Catalysis in
836   CRISPR-Cas12a. *Nat. Commun.* **15**, DOI: 10.1038/s41467-41024-45762-41466 (2024).

68   Arantes, P. R. *et al.* Dimerization of the Deaminase Domain and Locking Interactions
with Cas9 Boost Base Editing Efficiency in ABE8e. *Nucl. Acids Res.* **52**, 13931-13944
(2023).
69   Sinha, S. *et al.* Unveiling the RNA-mediated Allosteric Activation Discloses Functional
Hotspots in CRISPR-Cas13a. *Nucl. Acids Res.* **52**, 906-920 (2024).
70   Tian, C. *et al.* f19SB: Amino-Acid-Specific Protein Backbone Parameters Trained
against Quantum Mechanics Energy Surfaces in Solution. *J. Chem. Theor. Comput.* **16**,
528-552 (2019).
71   Galindo-Murillo, R. *et al.* Assessing the Current State of Amber Force Field
Modifications for DNA. *J. Chem. Theor. Comput.* **12** (2016).
72   Zgarbova, M. *et al.* Refinement of the Cornell et al. Nucleic Acids Force Field Based on
Reference Quantum Chemical Calculations of Glycosidic Torsion Profiles. *J. Chem.*
Theor. Comput. **7**, 2886-2902 (2011).
73   Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L.
Comparison of Simple Potential Functions for Simulating Liquid Water. *J. Chem. Phys.*
79, 926-935 (1983).
74   Case, D. A. *et al.* The Amber Biomolecular Simulation Programs. *J. Comput. Chem.* **26**,
1668-1688, doi:10.1002/jcc.20290 (2005).
75   Mongan, J., Simmerling, C., McCammon, J. A., Case, D. A. & Onufriev, A. Generalized
Born Model with a Simple, Robust Molecular Volume Correction. *J. Chem. Theor.*
Comput. **3**, 156-169 (2007).
76   Nguyen, H., Pérez, A., Bermeo, S. & Simmerling, C. Refinement of Generalized Born
Implicit Solvation Parameters for Nucleic Acids and their Complexes with Proteins. *J.*
Chem. Theor. Comput. **11**, 3714-3728 (2015).
77   Nguyen, H., Roe, D. R. & Simmerling, C. Improved Generalized Born Solvent Model
Parameters for Protein Simulations. *J. Chem. Theor. Comput.* **9**, 2020-2034 (2013).
863