1 **The forgotten wine: understanding the fermentation and composition of palm wine**
2

3 I Nyoman Sumerta[1,2], Xinwei Ruan[1] and Kate Howell[1]*
4
5 [1]School of Agriculture, Food and Ecosystem Sciences, Faculty of Science, The University of
6 Melbourne, Victoria 3010, Australia
7 [2]National Research and Innovation Agency (BRIN), Jakarta 10340, Indonesia
8
9 *Corresponding author: Kate Howell, khowell@unimelb.edu.au
10

Abstract

Palm wine is an alcoholic beverage that has existed for centuries and has important
economic and socio-culture values in many tropical and sub-tropical countries. Palm wine is
made by spontaneous fermentation of palm sap by naturally occurring microbial
communities, which determine product quality. Defining the core microbial communities
driving the fermentation process is pivotal for the distinctive of palm wine characteristics;
however, it is also challenging due to micro and macro-ecological factors, which contribute to
geographic specificity. A dynamic consortium of microbial species occupies every stage of
palm wine fermentation and consequently releasing metabolites to trigger growth of other
associated species. Studies have been conducted to investigate palm wine composition and
microbial diversity but these studies and have not been extrapolated to understand the
function of microbial communities and metabolite diversity. In this review, we gather
information of microbial communities and metabolite profiles from 47 published studies,
covering a wide range of methodologies and regions to better understand the causal links
between the principal microbial species and the metabolites of palm wine. We address palm
wine quality across production regions to provide specific regional information to elevate our
knowledge of this fermented beverage. This understanding will help to manage the causal
effects of palm wine fermentation and define the regional signature of palm wine to help with
the challenge of ensuring product quality in this unique, yet overlooked, fermented beverage.
30
31 **Keywords:** fermentation, metabolite signature, microbial community, palm wine, wine
32 specificity
33

1. Introduction

Microbial communities are the main factor to reveal fermented beverage characteristics,
attributed to chemical release as a results of ecological interactions (1–3). Other contributing
factors to the determination of beverage sensory characteristics are regional variations of
plant species and cultivar, genetic diversity of plant and microbe, which are expressed by the
metabolic profile of the end product (4,5). We know that there are dynamic complex
interactions, which impact upon the product quality and have been a major challenge for
product development (6). Setting a biogeographical approach to understand and define the
distinctiveness of fermented beverages will preserve the sensory diversity and increase the
product value (4). This is an important concern to protect designations of origins (PDOs) and
geographical indication (GIs) of fermented beverages, that have a significant impact in
economic aspect of the product (7,8).
46
Palm wine is less acknowledged compared to the better-known fermented beverages made
from grapes and rice. Palm wine is extremely important in many tropical countries as a
traditional alcoholic beverage, contributing to local economy and with distinctive impacts
upon social culture (9–12). Palm wine production is performed as a traditional practice and is
treated differently based on its origin, which leads to distinctive regional variation. The
production of palm wine relies on native microbial communities, which are likely have

adapted to local biogeographical constraints to conduct the fermentation (13). Some microbial species (including yeasts) have evolved to occupy a niche to engage with a wider range of ecological interactions (14). However, biotic and abiotic factors can specifically modify microbial community structure (15), and drive the differentiation of evolutionary traits and chemical release of the principal fermentative species, *Saccharomyces cerevisiae* (16). Understanding biogeographical ranges of microbial communities will contribute to the comparison of global populations, especially on domestication trajectory of palm wine's principal species (17–19) and gain cultural traction by being associated with the regionality of palm wine from topical countries.

Spontaneous fermentation of palm wine demonstrates dynamic nature and changing occurrence of fermentative and non-fermentative microbes, which can lead to flavour formation. Studying microbial ecology is important to conceptualize a fermentation model for developing a consistent beverage product (20). However, reports of palm wine ecology, especially on comprehensive microbial and metabolite profiles covering biogeographical perspective are limited. Compared to wines made from grapes and rice, the microbial ecology of palm wine study has not been intensively conducted to examine regional typicity. This scant information results in gaps for data comparison and general strategies for palm wine fermentation linked to the natural environment. Gathering and compiling recent studies can construct a general pattern of palm wine fermentation. For example, comprehensive studies of microbial communities and regional characteristics of grape wine have helped understanding of how the fermentation technique and ecological factors influencing product specificity (4,16). Examining palm wine microbial ecology across tropical areas will give new insights into the range, activity and formation of yeast and bacterial communities in sap from palm trees.

The important links of compromised microbial composition and metabolic properties to develop and preserve traditional fermented product are essential to better understand palm wine. This review focuses on the principal microbial communities and chemical compounds, that represent the characteristics of palm wine linked to ecology and geography, that influence the specific profiles for wine. We begin by discussing the distinctive composition of palm wine and the biogeographical areas of palm wine production. We explain the contribution of ecological factors on the fermentation process and how they affect palm wine sensory by collating the information present in published studies and examining them together. We conclude by examining the influence of local cultural practices on palm wine, followed by suggestions for directions in product development.

2. The distinctive composition and flavour of palm wine

Grapes and rice wines are the most known types of fermented beverages across the world and play an important part in social and economic development of many countries (4,21). Meanwhile, palm wine from topical countries is largely unknown. Palm wine is a traditional alcoholic beverage in Africa, south Asia, southeast Asia, and central America (11). Palm wine is derived from palm sap, which is tapped form the blossom or apical structure of palm tree (10,22). The fresh sap drips from the cutting part with slightly white turbid colour and moderate sweetness (23), containing sugars (predominantly sucrose) (24,25). In grapes, fructose and glucose predominate (26) and a crushing step is needed to release these sugars from the grapes to initiate the fermentation process. Rice requires extensive processing before starch conversion is ready to ferment (27). In contrast, palm sap can be directly fermented after tapping without any pre-processing steps.

Palm sap is a versatile commodity from palm trees, that can be transformed into further derivative products (**Fig. 1**). The fresh sap can be directly consumed and has various beneficial metabolites, including organic acids (24,28–30), vitamins (31,32), proteins (30),

phenolics (33), and minerals (31). A shift from sugar fermentation to acidic fermentation generates a variety of derivative products, such as palm sugar, alcoholic beverages, and palm vinegar (25,32,34,35). The fresh sap can be refined to obtain palm sugar and left naturally at room temperature to produce palm wine. Palm wine typically has shelf life only for 2-3 days of fermentation with alcohol content similar to beer (up to 6% v/v ethanol). Before a subsequent acidic fermentation, palm wine can be distilled to produce spirits, which has more than 40% ethanol (v/v) (36). A longer fermentation generates organic acids as a source to produce a palm vinegar. In all cases, the typical fermentation is conducted by palm's natural microbial community without addition of cultured microbes and in general swift and dynamic fermentation process occur (9,10).

3. Biogeographical perspectives of palm wine

Palm tree distribution across the sub and tropical countries is related to palm wine making practice. The utilization of specific palm species for palm wine making and the ecological factors exposure can mediate microbial ecology, which determines the distinctiveness of palm wine sensory attributes. Limited studies have been performed to establish and define palm wine regional typicity and so we have compiled recent studies to construct a clear picture of palm wine practices. Here, we have collated data from 47 studies, specifically focused on microbial and biochemical diversity topics within various approaches (**Supplementary Fig. S1**). Based on our compiled studies, palm wine has been studied across three continents over 14 tropical and subtropical countries (**Fig. 2A**). More than half of the studies was conducted in Africa, with the most studies in India. Each continent has specific palm species, utilized for palm wine making; *Nypha* sp. in Asia, *Acrocomia* sp. in Central America, *Elaesis* spp. and *Raphia* spp. in Africa. Of the studies collated, 17 focussed on description of the microbes present, 7 on the small molecules, which are microbially derived and likely to be sensorially active and 23 studies combined these approaches (**Fig.2B**).

Palm species can contribute to the regionality of palm wine characters, which are also attributed into a traditional name. In Burkina Faso and Cote d'Ivoire, palm wine is widely known as *Bandji* (37,38); *Mimbo* in Cameroon (20,39); *Akpeteshie* in Ghana (40); *Toddy* in India (41); *Tuak* in Indonesia (12,42); *Taberna* or *Tavern* in Mexico (11,43); *Ogogoro* in Nigeria (44); *Lambanog* in Philippines (45); *Ra* in Sri lanka (20); *Songkhla* in Thailand (24); *Lagmi* in Tunisia (46).

3.1 Macro-ecology affects palm wine signatures

Environmental variables have a significant role on palm sap production and characteristics (47). Although it may be less appreciated, environmental factors contribute to the palm physiology, which can directly affect the sap yield and physicochemical characters, inducing the microbial communities, and alter the fermentation process. Astudillo-Melgar et al. (13). believed that microbial diversity is correlated to biogeographical factors from where palm wine is produced and adapted with specific ecological variables. Factors such as climate, weather, palm species, microbial vector, and tapping process may not work independently to alter the microbial composition, but often exert an integrated effect. Therefore, impacts will be discussed as an important aspect to taking further action on expected mitigation for product development.

3.1.1 Biogeographical locality
Although palm is mostly observed in coastal area of tropical and sub-tropical countries (48), the variety of local factors could also impact the diversity of microbial community. Barata et al. (49) emphasised that microbial diversity is relied on climatic condition, temperature, sunlight exposure, wind, and rainfall where the plant origin, indicating the sign of terroir.

Environmental factors have been observed on their influence for plant and soil-microbes
association (15); however, there is a lack information on arboreal microbes, correlated to
palm sap tapping. A study by Ziadi et al. (47) reported that sap production and
physicochemical characters are significantly influenced by tapping season. Higher sugar
content and less acidic are characterized from palm sap in warmer seasons, which have
more sunlight exposure to increase palm photosynthesis and thus sugar content. Microbial
counts are slightly higher, which is less affected by the warmer temperature. The fluctuation
of microbial occurrence in a specific region is altered by rainfall although a consistent pattern
are not shown (49). The variety of regional factors and microbial characteristics determine
how a microbial species might adapt to those specific niches. In conjunction to the local
climate and specific biogeographical factors, there are very limited studies which might help
to determine the regional typicity of palm wine. It is likely that these factors are important as
the interplay among environment, plant species, and winemaking practice impacts the
microbial populations and their metabolism and contributes to formation of the final product
characteristics, reflecting a fermented beverage product's origin and terroir (2,4,16).
173
3.1.2 Palm diversity
Palm species are a critical aspect that contribute to the differences of palm sap
characteristics and thus flavour. There are more than 10 species of palm trees, that have
utilized for making palm wine (22). As showed in **Fig. 2C**, palm species are established and
well-adapted to local biogeographical characteristics. For example, *Elaensis* and *Raphia*
palm species are used in Africa but unrecorded in both Asia and America. Palm tree
specificity can be considered using chemical analysis of the sap where each palm species
likely has different composition and quantity of macro and micronutrient, contributing to the
growth of microbial communities (31,50). This concords to a unique microbial composition in
each palm species (10,25,29), which depends upon geographical location (9,10,19,25).
Although a general pattern might be showed by each palm species, that generally the
fermentative species, *S. cerevisiae*, conquers to all palm sap niches. The diversity
differentiation likely occurs in non-*Saccharomyces*, determining the flavour complexity. This
variation indicates that palm species genetic diversity and the influence of geographical
locality are the significant factors to determine palm wine specificity. These processes and
the collated data from our survey are explained in the following sections.

3.2 Introduction of microbes into palm sap

Palm sap is believed in a sterile condition upon collection and microbial communities are
inoculated through insect visitation and the tapping process (24,51,10). Insects and
microbes have an intimate relationship with plants, signalling symbiotic associations to help
them detecting nutrient, protecting from pathogens, and as parasites (52,53). Insects find
sugar source through the detection of released volatile compounds by plants (52,54). Palm
sap is rich in sugars, which attract bees and other insects to the collected sap and can be
observed floating in the liquid and visiting the collection vessels. Honeybees seek sugary
elements of plants, and then acts as pollinator while vectoring microbes (55). Wasps and
vinegar flies are often observed in sugar-rich environments in other fermentations (56,57).
The close relationships of palm sap and insects reflect the vital role of microbial vectors to
contribute to the palm wine flavour characteristics. On the other hands, artisanal processes
can also contribute to the introduction of microbial communities. Since the tapping process is
carried out with a traditional way, and without deliberates sanitation, the possibility of
microbe introduction is inevitable and may be from the collection container, human hands,
and cutting equipment such as knives. Microbial loads in palm sap can be influenced by a
perpetual tapping practice, which can reduce up to 25% (47). Factor of slicing stress during
the tapping process is likely related to the decrease of palm sap nutritive value after 10
weeks consecutive tapping and impacts to microbial load. The decrease of microbial loads
directly impacts to the fermentation process, affecting the quality and stability of final

product. It is likely that the tapping process is a crucial management for product hygiene and
sap production sustainability.

4   How does the nature of palm wine fermentation contribute to the construction of wine flavour?

The nature of palm wine fermentation is extremely fast and complex, involving sugars
accessibility (10,58). Spontaneous fermentations allow all natural microorganisms to
compete organic resources without deliberate addition of selected yeast strains. The
limitation of the fermentation environment is one of significant factor to alter microbial
interactions and ecological dynamics (59,60). As a result, a change in microbial structure is
often observed along the fermentation process. Here, to see microbial diversity and dynamic
along the fermentation process, we reanalysed next generation sequencing (NGS) data of 7
abstracted studies with further data from Sumerta et al. (61); (**Supplementary Table S4**).
We subsequently selected several studies by inclusion criteria, representing a vast diversity
change in 24 hours fermentation (**Fig. 3A & 3B)**.

Comparison within country or region, type of sample, and some time frames are less
informative because of the limitation in comparable parameters, such as research focus,
methodology, and data distribution. Since palm wine fermentation is swift, the species loss of
fermenting microbial communities is seen within 24 hours (**Fig. 3A)**; demonstrated by the
decrease in Shannon index during fermentation (0, 12, and 24 hours of fermentation). Higher
diversity is observed in early stage of the fermentation and decrease to the following process
(62). Nwachukwu et al. (63) used isolation methods and found a diversity loss because of
succession of some microbial species. Djeni et al. (50) reiterated that microbial diversity loss
was due to a unique inhabitant species in every stage of palm wine fermentation, that
dominant growth.

The relative abundance of bacterial and fungal amplicons from next generation sequencing
and collated demonstrate that while the initial microbial populations may be maintained for
12 hours, the dominance of *Lactococcus* and *Saccharomyces* is evident (**Fig. 3B**). There are
10 studies that describe a collapse of one microbial group in every stage of palm wine
fermentation (**Supplementary Table S6**). Microbial species fluctuate over 24 hours
fermentation of palm wine (**Fig. 3B**). The decrease of *Leuconostoc* and the growth of
Lactococcus present in early stage, with a concurrent drop in *Hanseniaspora* species
occurrence and the increase of *Saccharomyces* species. A shift between *Saccharomyces*
and non-*Saccharomyces* is inevitable in every stage of the fermentation (50,58,63,64). This
succession of microbial composition over the fermentation process is identified as a key
feature to determining palm wine characteristics (50).

4.1 *Saccharomyces cerevisiae* is prominent in palm wine fermentations

In a natural fermentation, the complex microbial interactions may result in the dominance of
certain species. We sought to find the microbial species most regularly associated with palm
fermentation Through the comparison of a conventional microbial isolation and species
inferred from amplicon barcodes, we selected data of top 10 abundance of culturable
species (CD) (**Supplementary Table S1)** and NGS profiling (CI) (**Supplementary Table
S4)**. We found the general pattern of microbial composition in palm wine with the presence
of both principal yeast species and principal bacterial genera, which are *S. cerevisiae* and
Leuconostoc, respectively (**Fig.3C**). Both of methods confirm that the role of *S. cerevisiae*
and *Leuconostoc* and this contributes to more than 40% of the total microbial abundance.
Some species that do not present in one of those methods may relate to the limitation of
each method. Meanwhile, as described in previous studies, *S. cerevisiae* is dominant in all
fermentation stages while *Leuconostoc* fluctuates during the final stage (9,50). The principal
species may dominant for the primary product, but non-fermentative or non-*Saccharomyces*
species enriches the sensory outputs for fermented beverages more broadly (65,66).

262
263
4.1.1 What circumstances allow for dominance of microbes?
The role of the key yeast species on producing ethanol as the primary product is pivotal in palm wine fermentation. Some bacterial species are reported to produce ethanol, such as *Zymomonas* spp. (62), but production is less effective than *S. cerevisiae.* The fermentation process is driven by *S. cerevisiae* in the production of many beverages, as this yeast is able to readily transport sugars, other carbohydrates are hydrolysed and various compound are produced and released. Some of the produced compounds can inhibit other species and the others yet may display e synergistic actions to adapt with the change of ecological factors during the fermentation process (67,68). This often leads to the dominance of certain species in a complex spontaneous fermentation of alcoholic beverages of which *S. cerevisiae* in the prime example (66,69,70).

S. cerevisiae is widely regarded as the most tolerance species to ethanol, acidity, and physical interactions (71–74). Perrone et al. (69) suggested that its dominancy is triggered by the occurrence of other species, driving various mechanisms to suppress non-*Saccharomyces*. A regulatory mechanism of its alcohol dehydrogenases (*ADH*) genes controls the ethanol production and at the same time can recover energy when the ethanol excess (75). *ADH1* gene regulates the fermentation process to produce ethanol, while *ADH2* gene is responsible to transferring energy from ethanol conversion. Non-*Saccharomyces* species are relatively vulnerable and non-competitive to the alcohol and other environmental factors (74). *S. cerevisiae* may secrete compounds, such fungicide like mycosin and perform cell-contact abundance and triggering to the death of other species in mixed culture fermentation (76,77). Albergaria et al. (78) found that the 2-10 kDa size of peptides secreted by *S. cerevisiae* can inhibit the growth of *Kluyveromyces marxianus, Kluyveromyces thermotolerans, Torulaspora delbrueckii* and *Hanseniaspora guilliermondii*. The type of cell-to-cell contact and antibiotic production mechanism can also result in the death of certain species (76). Meanwhile, cell-to-cell contact induces the metabolites properties of the species and affect cell viability in late stage of the fermentation (66,77). However, physical contact is not the major factor to decrease the presence of non-*Saccharomyces* and confirmed that the metabolites produced by *S. cerevisiae* mostly sharps the non-*Saccharomyces* population (77). With the efficiency of energy use, less osmotic stress, high ethanol production, tolerant to lower pH, and enable to inhibit other species, *S. cerevisiae* often survives until the final stage of fermentation (19). It is likely that many of these characteristics are also true of *S. cerevisiae* found in palm wine environments, but they have not been directly shown.

4.1.2 Diversity of *Saccharomyces* and the implications for palm wine fermentations
S. cerevisiae is a model species for genomics and plays a pivotal role in many food systems, having been selected, adapted, and well domesticated across human history (79–82). Peter et al. (83) described that the highest variation in *S. cerevisiae* genomes is correlated to the wider differences of phenotypic characters, and it is these characters that are responsible for the signature and originality of fermentation products. The evolutionary process of *S. cerevisiae* at genome level can be modified by human interferences, and the yeasts adapt to various growth conditions, alter phenotypes, and change reproductive mode (84,85). In natural conditions, the population structure of *S. cerevisiae* is primarily shaped by ecology (86). *S. cerevisiae* evolution can be triggered by its respond to ecological interactions in harsh fermentation environments (1).

Spontaneous fermentation on fermented beverage making has been reproducibly conducted and if it affects on microbial diversity loss, this potentially leads to a species domestication (87). Interaction with the natural environment is often possible, where new strains may contribute, and experience variable abiotic conditions (80,84). Legras et al. (88) stated that a rich environmental condition, geographical condition, and repetitive use in the same

condition results in gradual genotypes evolution, species distinction, and suggest local
domestication pattern. However, spontaneous fermentation of palm wine is still argued to
generate domestication and leaved as an open question (89). Some reports released that *S.
cerevisiae* population of African palm wine is admixed compared to global references,
showing a unique signature relatively to different continents, niches, and favour strain
migration through human interference (18,19). Tra Bi et al. (19) showed that the genetic
relationship of *S. cerevisiae* in palm wine is significantly related to ecological niches and
sharing the similar geographical origin generates less genetic diversity. A local
domestication may be attributed to *S. cerevisiae* from palm wine strains, which are observed
as homozygous strains (18,90).

327 4.2 Interactions amongst microbes in fermentation

We conducted a network analysis of the co-occurrence of species groups from compiled
studies which used culture-dependent methods to assess microbial diversity (**Fig. 4**). The
contribution of *Saccharomyces* is dominant and acts as a central hub to other species group
in palm wine fermentation. It also has a strong relationship to other yeast genera
(*Hanseniaspora, Candida*, and *Pichia*) and the core bacteria in organic acids production,
such as *Lactobacillus*, *Leuconostoc*, and acetic acid bacteria (AAB). These show a vital
connection pattern of those microbial groups on constructing palm wine characters and
suggest a fermentation model retrieved from multi-biogeographical studies. This pattern
reveals a network of microbial ecology, and reveals potential microbial interactions which
may be able to predict the occurrence of unwanted species (91).
338
In spontaneous fermentation, interactions have a significant role on driving the fermentation
process, involving various microbial groups. Interactions among microbial species may
deliver positive and negative impacts, which determine the product formation (92–95). In
spontaneous fermentations, synergistic interactions among different species mediate
tolerance to the robustness of dynamic condition in particular level. Mutualism and
commensalism interaction are attributed to the synergistic interaction with cross-feeding
mechanism or supporting certain metabolite producer (3). For instance, yeasts ferment the
sugar into ethanol while acetic acid bacteria convert the ethanol into acetic acid which is not
affect the yeast (96). Lactic acid bacteria and yeast sometimes support each other by
exchanging some metabolites for their metabolism (3). Each group of microbial species has
different way to develop its functionality by interacting with other species. Fleet (68)
highlighted the interaction among yeast species contributes to the fermentation behaviour
while yeast-bacteria interactions determine the ecological dynamic. This interesting feature
of complex microbial communities interaction can be proposed as new model of the
fermentation to create flavourful product (67). Specific interactions between microbes have
not been described in a palm wine environment, but they are likely to occur.

355 4.3 The implications of metabolite release on fermentation dynamics

The associations of microbial communities and chemical release have a major contribution
on constructing palm wine characteristics. A wide range of microbial groups occupies in
every stage of the fermentation process but following the depletion of sugar in the
fermentation process, microbial diversity decreases because of biochemicals alteration and
ecological interactions. Here, we abstracted 10 studies (**Supplementary Table S6**),
exploring both in microbial communities and chemical composition along the fermentation
stages, representing palm wine fermentation dynamics and summarised in **Fig. 5**. The
abundance of ethanol is likely attributed to the presence of *Saccharomyces* yeasts during
fermentation. Lactic acid bacteria (LAB) and acetic acid bacteria (AAB) grow on palm wine
fermentation related to the significant number of organic acids in final stages of fermentation
(38,97–99). Total phenolics and flavonoids impact shifts in microbial community composition
through the fermentation process (58). We presented the examples of chemical yield in the
table, which represents the gradual change in line graph. It is important to note that chemical

content of palm wine ultimately depends on, geographical location, palm species, and the way analytical process and research design conducted, so the results vary among studies. By summarising related studies, the general pattern was demonstrated to provide a better understanding of palm wine fermentation.

We simulated the co-occurrence of microbial groups and compounds based on the correlation analysis of each microbial group follows the biochemical pattern (**Fig. 6**). Data was summarised from the frequency of microbial groups and metabolite components presented in each study. Lactic and acetic acid production in the final stage of palm wine fermentation positively corelate to the frequency of lactic acid and acetic acid bacteria group while *Saccharomyces* and *Candida* group may also have similar contribution. These compounds reduce the pH and alter the microbial diversity. Since pH decreasing, the evaporation of volatile fatty acids is possibly high (100). Meanwhile in alcohol production, non-*Saccharomyces* yeasts have a negative correlation compared to bacterial groups and *Saccharomyces*. This simulation provides the influence of metabolite components on the composition of microbial diversity, which can demonstrate the fermentation dynamics.

5 The intersection of culture on palm wine production

The process of palm wine making is still performed in traditional ways from tapping process to the fermentation. Depend on palm species, tapping process can be conducted through destructive and non-destructive methods (10). In West Africa, destructive tapping process is mostly observed by notching or perforating the apical meristem, often resulting in the death of the tree. The widely used method is non-destructive by cutting the inflorescence and the sap drips from cutting tip to the collection container (25). The collected palm sap is then processed into various products for daily consumption or sold. For some areas, this activity is the main job of local communities and maintains economic stability in many villages.

The use of palm sap and/or wine has an important role as part of culture and tradition (12). Palm spirits can be used as part of wedding ceremonies, event celebration, welcoming rituals, and religious rituals. In some tropical countries, palm sap products are also responsible for ethnomedicines. Anthropologists have acknowledged that palm wine is important as stimulant, depressant, and part of traditional medication, well-written in traditional transcripts (101). Recent scientific reports also suggested that palm sap/wine compounds have health benefits, related to the empirical thoughts of local people, such as decrease diabetes risk, antioxidant, antibiotics, anaemia treatment, and immunomodulators (22,31,102–105). Palm wine contains phenolic compounds that may contribute to antioxidant activity in the plasma (22,33) and modulate the growth of gut microbiota (106–108). A probiotic-type microbiome that benefits for digestive system is also found in fresh palm sap (64) and fermented sap showed bioactive chemical activities (109). Despite the significance of pharmacological and ethnomedicinal utilization, basic information on microbial community structure and dynamic of metabolite profile benefits to the traditional thoughts and practices that may evolve driving regional identity.

6 The future of palm wine and derivatives in modern food systems

Understanding microbial ecology and controlling microbial composition are a pivotal aspect to the beverage quality and a major challenge for future product development (6). A rough and inconsistent characteristic of beverages results from spontaneous fermentations and it is difficult to control the product quality and consistency. By the application of defined starter culture, more consistency, efficiency, and desire characters of the alcohol products are achieved; however, the product might lack complexity of organoleptic characteristics (82). The presence of spoilage bacteria is inevitable amid a spontaneous fermentation. Spoilage bacteria, for example lactic acid and acetic acid bacteria, can compete for nutrients and produce lactic and acetic acids which limit the ethanol production during the fermentation

process (110). Intervention at harvesting can obtain higher ethanol production before the
spoilage bacteria rapidly conducts the acidic fermentation.

As palm wine is product of spontaneous fermentation and has demand in local communities,
preserving the product and maintaining the product quality and stability are the major
challenge to bring this beverage to a global market. Palm wine has a swift fermentation for
alcohol and organic acids that the selection of fermentation time can help to obtain the best
yield. However, many ecological factors in palm productivity, season, tapping process, and
microbial composition may create unstable product quality and quantity. Defining processing
solution and while maintaining palm wine regional distinction will stimulate local economies
and support maintaining traditional practices.

Mimicking the natural fermentation in the laboratory from raw materials to environmental
conditioning can support product quality and stability at an industrial scale. Indeed, alcoholic
fermentation development begun from spontaneous fermentation, evolving to current
commercial practices with selected yeast strains (82,111). Procedures could adapt by
shifting the fermentation practices from natural fermentation to the use of defined starter to
maintain the sensory characters by mixing microbes together in a starter culture. The mixed
cultures can provide a broader profile of aroma and flavour than using single culture starter,
at least in grape juice fermentations (112,113). Using non-*Saccharomyces* associated with
S. cerevisiae is a growing trend in the fermented beverage industry (67). Non-
Saccharomyces play a significant role on constructing the product value, acting as a co-
culture to support the fermentation process (65) and their metabolic releases might be
adjusted through the mixed fermentation (67). We recognise that there may be limitations
have to further action as there is a need to retain and foster traditional food products, which
may at odds with the use of commercial starter cultures.

7 Conclusions

Palm wine fermentation is challenged by many factors, which contribute to differences of
product composition and distinction. The main actor is the microbial communities, which
affect palm wine composition, metabolic release, and genetic diversity to drive the
fermentation process and determine product characteristics. Environmental factors are an
additional aspect of typical raw material, collection and tapping strategy, then
biogeographical location leads to the product specificity. The interplay of those factors
contributes to the construction of palm wine regional signature, fostering local product quality
into a global economic quality. We support further actions to strengthen the fundamental
understanding of palm wine production and here we have provided an overview of the
challenges for sustainable product development and appreciating a local knowledges and
traditions.

Supplementary materials

Supplementary Fig. S1 Literature searching and data variability of retrieved studies
Supplementary Table S1 Data of included studies in microbial diversity
Supplementary Table S2 Data of included studies in metabolite profile
Supplementary Table S3 Data of studies both in microbial diversity and metabolite profile
Supplementary Table S4 Abstracted studies conducting NGS method
Supplementary Table S5 Abstracted data of studies in frequency and occurrence of
microbial groups
Supplementary Table S6 Abstracted studies representing fermentation stages

Conflict of Interest

468 Authors declare no conflict of interest

Funding

This study is supported by Australia Awards of the Government of Australia, which the views and opinions expressed in this review are from authors and do not represent the Australian Government.

Acknowledgements

Authors would like to pass our gratitude to University of Melbourne for access to the high-performance computing for faster analysis. Theodore N. Djeni and Kumaraswamy Jeyaram who provided the sequencing data available publicly for compiled analysis. SSEAC (Sydney Southeast Asia Centre) for the intensive writing program, supporting on the writing process.

Author contributions

INS: research design, data compilation and analysis, and writing the manuscript
XR: data analysis supervision and review the manuscript
KH: research design supervision, writing, and review the manuscript

References

1. Conacher C, Luyt N, Naidoo-Blassoples R, Rossouw D, Setati M, Bauer F. 2021.The ecology of wine fermentation: a model for the study of complex microbial ecosystems. *Appl Microbiol Biotechnol* 105(8):3027–43.

2. Liu D, Zhang P, Chen D, Howell K. 2019. From the vineyard to the winery: how microbial ecology drives regional distinctiveness of wine. *Front Microbiol* 10:2679.

3. Smid EJ, Lacroix C. 2013. Microbe–microbe interactions in mixed culture food fermentations. *Curr Opin Biotechnol* 24(2):148–54.

4. Bokulich NA, Collins TS, Masarweh C, Allen G, Heymann H, Ebeler SE, et al. 2016. Associations among wine grape microbiome, metabolome, and fermentation behavior suggest microbial contribution to regional wine characteristics. *mBio* 7(3):e00631-16.

5. Liu D, Chen Q, Zhang P, Chen D, Howell KS. 2020. The fungal microbiome is an important component of vineyard ecosystems and correlates with regional distinctiveness of wine. *mSphere* 5(4):e00534-20.

6. Gobert A, Evers MS, Morge C, Sparrow C, Delafont V. 2022. Comparison of DNA purification methods for high-throughput sequencing of fungal communities from wine fermentation. *MicrobiologyOpen* 11(5):e1321.

7. Candiago S, Tscholl S, Bassani L, Fraga H, Egarter Vigl L. 2022. A geospatial inventory of regulatory information for wine protected designations of origin in Europe. *Sci Data* 9(1):394.

8. Deselnicu OC, Costanigro M, Souza-Monteiro DM, McFadden DT. 2013. A meta-analysis of geographical indication food valuation studies: what drives the premium for origin-based labels? *J Agric Resour Econ* 38(2):204–19.

9. Das S, Tamang JP. 2021. Changes in microbial communities and their predictive functionalities during fermentation of toddy, an alcoholic beverage of India. *Microbiol Res* 248:126769.

10. Djeni TN, Kouame KH, Ake FDM, Amoikon LST, Dje MK, Jeyaram K. 2020. Microbial diversity and metabolite profiles of palm wine produced from three different palm tree species in Côte d'Ivoire. *Sci Rep* 10(1):1715.

11. Santiago-Urbina JA, Ruíz-Terán F. 2014. Microbiology and biochemistry of traditional palm wine produced around the world. *Int Food Res J* 21(4):1261–9.

12. Wijaya L, Sumerta IN, Napitupulu TP, Kanti A, Keim AP, Howell K, et al. 2024. Cultural, nutritional and microbial perspectives of tuak, a traditional Balinese beverage *J Ethn Foods* 11(1):4.

13. Astudillo-Melgar F, Ochoa-Leyva A, Utrilla J, Huerta-Beristain G. 2019. Bacterial diversity and population dynamics during the fermentation of palm wine from Guerrero Mexico. *Front Microbiol* 10:531.

14. Gray ND, Head IM. 2008. Microbial ecology. *In*: Jørgensen SE, Fath BD (ed). Encyclopedia of ecology, Internet. Academic Press, Oxford.

15. Zhou J, Cavagnaro TR, De Bei R, Nelson TM, Stephen JR, Metcalfe A, et al. 2021. Wine Terroir

521   and the Soil Bacteria: An amplicon sequencing–based assessment of the barossa valley and its
522   sub-regions. *Front Microbiol* 11:597944.

523   16.   Knight S, Klaere S, Fedrizzi B, Goddard MR. 2015. Regional microbial signatures positively
524   correlate with differential wine phenotypes: evidence for a microbial aspect to terroir. *Sci Rep*
525   5(1):14233.

526   17.   De Guidi I, Legras JL, Galeote V, Sicard D. 2023. Yeast domestication in fermented food and
527   beverages: past research and new avenues. *Curr Opin Food Sci* 101032.

528   18.   Ezeronye O u., Legras JL. 2009. Genetic analysis of *Saccharomyces cerevisiae* strains isolated
529   from palm wine in eastern Nigeria: Comparison with other African strains. *J Appl Microbiol*
530   106(5):1569–78.

531   19.   Tra Bi CY, Amoikon TLS, Kouakou CA, Noemie J, Lucas M, Grondin C, et al. 2019. Genetic
532   diversity and population structure of *Saccharomyces cerevisiae* strains isolated from traditional
533   alcoholic beverages of Côte d'Ivoire. Int J Food Microbiol 297:1–10.

534   20.   Ciani M, Stringini M, Comitini F. 2012. Palm Wine. Handbook of Plant-Based Fermented Food
535   and Beverage Technology, Internet. CRC Press, Boca Raton.

536   21.   Anderson K, Wittwer G. 2015. Asia's evolving role in global wine markets. *China Econ Rev* 35:1–
537   14.

538   22.   Hebbar KB, Pandiselvam R, Manikantan MR, Arivalagan M, Beegum S, Chowdappa P. 2018.
539   Palm sap—quality profiles, fermentation chemistry, and preservation methods. *Sugar Tech*
540   20(6):621–34.

541   23.   Chandrasekhar K, Sreevani S, Seshapani P, Pramodhakumari J. 2012. A Review on palm wine.
542   *Int J Res Bio Sci* 2(1):33–8.

543   24.   Naknean P, Meenune M, Roudaut G. 2010. Characterization of palm sap harvested in Songkhla
544   province, Southern Thailand. *Int Food Res J* 17:977–86.

545   25.   Limtong S, Am-In S, Kaewwichian R, Kaewkrajay C, Jindamorakot S. 2020. Exploration of yeast
546   communities in fresh coconut, palmyra, and nipa palm saps and ethanol-fermenting ability of
547   isolated yeasts. *Antonie Van Leeuwenhoek* 113(12):2077–95.

548   26.   Corbin KR, Hsieh YSY, Betts NS, Byrt CS, Henderson M, Stork J, et al. 2015. Grape marc as a
549   source of carbohydrates for bioethanol: Chemical composition, pre-treatment and
550   saccharification. *Bioresour Technol* 193:76–83.

551   27.   Lee TJ, Liu Y, Liu WA, Lin YF, Lee HH, Ke HM, et al. 2022. Extensive sampling of *Saccharomyces*
552   *cerevisiae* in Taiwan reveals ecology and evolution of predomesticated lineages. *Genome Res*
553   gr.276286.121.

554   28.   Doddipalla R, Rendedula D, Ganneru S, Kaliyaperumal M, Mudiam MKR. 2022. Understanding
555   metabolic perturbations in palm wine during storage using multi-platform metabolomics. *LWT*
556   155:112889.

557   29.   Nwaiwu O, Ibekwe VI, Amadi ES, Udebuani AC, Nwanebu FC, Oguoma OI, et al. 2016.
558   Evaluation of fermentation products of palm wine yeasts and role of *Sacoglottis gabon*ensis
559   supplement on products abundance. *Beverages* 2(2):9.

560   30.   Pammi N, Bhukya KK, Lunavath RK, Bhukya B. 2021. Bioprospecting of Palmyra Palm (*Borassus*
561   *flabellifer*) Nectar: Unveiling the probiotic and therapeutic potential of the traditional rural
562   drink. *front Microbiol* 12:683996.

563   31.   Barh D, Mazumdar BC. 2008. Comparative nutritive values of palm saps before and after their
564   partial fermentation and effective use of wild date (*Phoenix sylvestris* Roxb.) sap in treatment
565   of anemia. Med Sci 3(2):173–6.

566   32.   Zongo O, Tapsoba F, Leray F, Bideaux C, Guillouet S, Traoré Y, et al. 2020. Nutritional,
567   biochemical and microbiological composition of *Borassus aethiopum* Mart. sap in Burkina Faso.
568   *J Food Sci Technol* 57(2):495–504.

569   33.   Xia Q, Li R, Zhao S, Chen W, Chen H, Xin B, et al. 2011. Chemical composition changes of post-
570   harvest coconut inflorescence sap during natural fermentation. *Afr J Biotechnol* 10(66):14999–
571   5005.

572   34.   González Á, De Vuyst L. 2009. Vinegars from Tropical Africa. *In*: Solieri L, Giudici P (ed).
573   Vinegars of the World, Internet. Springer, Milan.

574   35.   Somawiharja Y, Wonohadidjojo DM, Kartikawati M, Suniati FRT, Purnomo H. 2018. Indigenous
575   technology of tapping, collecting and processing of coconut (*Cocos Nucifera*) sap and its quality
576   in Blitar Regency, East Java, Indonesia. *Food Res* 2(4):398–403.

577   36.   Pratiknjo MH, Mambo R. 2019. The cultural value of the minahasa people about liquor "Cap
578   Tikus." *J Drug Alcohol Res* 4.

579   37.   Karamoko D, Djeni NT, N'guessan KF, Bouatenin KMJP, Dje KM. 2012. The biochemical and
580   microbiological quality of palm wine samples produced at different periods during tapping and
581   changes which occured during their storage. *Food Control* 26(2):504–11.

582   38.   Ouoba L i. i., Kando C, Parkouda C, Sawadogo-Lingani H, Diawara B, Sutherland J p. 2012. The
583   microbiology of Bandji, palm wine of *Borassus akeassii* from Burkina Faso: identification and
584   genotypic diversity of yeasts, lactic acid and acetic acid bacteria. *J Appl Microbiol* 113(6):1428–
585   41.

586   39.   Jespersen L. 2003. Occurrence and taxonomic characteristics of strains of Saccharomyces
587   cerevisiae predominant in African indigenous fermented foods and beverages. *Fems Yeast Res*
588   3(2):191–200.

589   40.   Amoa-Awua WK, Sampson E, Tano-Debrah K. 2007. Growth of yeasts, lactic and acetic acid
590   bacteria in palm wine during tapping and fermentation from felled oil palm (*Elaeis guineensis*)
591   in Ghana. *J Appl Microbiol* 102(2):599–606.

592   41.   Tamang JP. 2022. "Ethno-microbiology" of ethnic Indian fermented foods and alcoholic
593   beverages. *J Appl Microbiol*. 133(1):145–61.

594   42.   Hermansyah H, Novia N, Minetaka S, Satoshi H. 2015. *Candida tropicalis* Isolated from tuak, a
595   north sumatera- indonesian traditional beverage, for bioethanol production. *Microbiol*
596   *Biotechnol Lett* 43(3):241–8.

597   43.   Coutiño B, Flores AC, Vela-Gutiérrez G, Sepúlveda L, Aguilar CN, Chavez-Gonzalez M, et al.
598   2020. Tavern or coyol wine: a beverage from palm sap with biotechnological potential. *In*:
599   Grumezescu AM, Holban AM (ed). Biotechnological progress and beverage consumption,

600   Internet. Academic Press, UK.

601   44.   Ikeh IM, Anele BC, Ukanwa CC, Njoku SO. 2021. Analysis of the microbial quality of locally
602   consumed palm wine sold in elele community of Rivers State Nigeria. *Eur J Nutr Food Saf* 62–9.

603   45.   Culla S, Perey CL, Tagasa J, Uy JK. 2019. Quantification of methanol content in homemade palm
604   wine (Lambanog) produced in Barangay Balungay, Panukulan, Quezon Province. *Salle Med*
605   *Helath Sci Inst,* Available from: https://greenprints.dlshsi.edu.ph/grade_12/3

606   46.   Malek G, Moktar H, Luciano B, Vittorio C, Salvatore M. 2017. Identification of acetic acid
607   bacteria isolated from Tunisian palm sap. *Afr J Microbiol Res* 11(15):596–602.

608   47.   Ziadi M, Gaabeb N, Mrabet A, Ferchichi A. 2014. Variation in physicochemical and
609   microbiological characteristics of date palm sap (*Phoenix dactylifera*) during the tapping period
610   in oasian ecosystem of Southern Tunisia. *Int Food Res J* 21(2).

611   48.   Moore HE, Uhl NW. 2019. Palm. Available from: https://www.britannica.com/plant/palm-tree

612   49.   Barata A, Malfeito-Ferreira M, Loureiro V. 2012. The microbial ecology of wine grape berries.
613   *Int J Food Microbiol* 153(3):243–59.

614   50.   Djeni TN, Keisam S, Kouame KH, Assohoun-Djeni CN, Ake FDM, Amoikon LST, et al. 2022.
615   Dynamics of microbial populations and metabolites of fermenting saps throughout tapping
616   process of ron and oil palm trees in Côte d'Ivoire. *Front Microbiol* 13.

617   51.   Santiago-Urbina JA, Arias-García JA, Ruiz-Terán F. 2015. Yeast species associated with
618   spontaneous fermentation of taberna, a traditional palm wine from the southeast of Mexico.
619   *Ann Microbiol* 65(1):287–96.

620   52.   Becher PG, Hagman A, Verschut V, Chakraborty A, Rozpędowska E, Lebreton S, et al. 2018.
621   Chemical signaling and insect attraction is a conserved trait in yeasts. *Ecol Evol* 8(5):2962–74.

622   53.   Douglas AE. 2011. Lessons from Studying Insect Symbioses. *Cell Host Microbe* 10(4):359–67.

623   54.   Madden AA, Epps MJ, Fukami T, Irwin RE, Sheppard J, Sorger DM, et al. 2018. The ecology of
624   insect–yeast relationships and its relevance to human industry. *Proc R Soc B Biol Sci*
625   285(1875):20172733.

626   55.   Evans JD, Schwarz RS. 2011. Bees brought to their knees: microbes affecting honey bee health.
627   *Trends Microbiol* 19(12):614–20.

628   56.   Lam SSTH, Howell KS. 2015. Drosophila-associated yeast species in vineyard ecosystems. FEMS
629   *Microbiol Lett* 362(20).

630   57.   Stefanini I, Dapporto L, Legras JL, Calabretta A, Di Paola M, De Filippo C, et al. 2012. Role of
631   social wasps in Saccharomyces cerevisiae ecology and evolution. *Proc Natl Acad Sci*
632   109(33):13398–403.

633   58.   Das S, Tamang JP. 2023. Fermentation Dynamics of Naturally Fermented Palm Beverages of
634   West Bengal and Jharkhand in India. *Fermentation* 9(3):301.

635   59.   Haloin JR, Strauss SY. 2008. Interplay between ecological communities and evolution: review of
636   feedbacks from microevolutionary to macroevolutionary scales. *Ann N Y Acad Sci* 1133:87–125.

60. Kowallik V. 2015. The natural ecology of *Saccharomyces* yeasts [PhD Thesis]. the Christian
Albrechts University of Kiel, Germany.

61. Sumerta IN, Howell K, Sudiana IM, Kanti A. 2024. Co-occurence of non Saccharomyces yeast in
the natural fermentation of palm wine in Indonesia. doi: 10.26188/25309117.v1

62. Alcántara-Hernández RJ, Rodríguez-Álvarez JA, Valenzuela-Encinas C, Gutiérrez-Miceli FA,
Castañón-González H, Marsch R, et al. 2010. The bacterial community in 'taberna' a traditional
beverage of Southern Mexico. *Lett Appl Microbiol* 51(5):558–63.

63. Nwachukwu I, Ibekwe VI, Anyanwu BN. 2017. Investigation of physicochemical and microbial
succession parameters of palm wine. *J Food Tech* 4(4).

64. Gopal M, Shil S, Gupta A, Hebbar KB, Arivalagan M. 2021. Metagenomic investigation uncovers
presence of probiotic-type microbiome in Kalparasa® (Fresh Unfermented Coconut
Inflorescence Sap). *Front Microbiol* 12:662783.

65. Li X, Liu SQ. 2016. Antagonism between *Saccharomyces cerevisiae* and *Williopsis mrakii* and
significance for mixed yeast alcoholic beverage fermentations. *Int J Food Sci Technol*
51(3):656–63.

66. Pietrafesa A, Capece A, Pietrafesa R, Bely M, Romano P. 2020. *Saccharomyces cerevisiae* and
Hanseniaspora uvarum mixed starter cultures: Influence of microbial/physical interactions on
wine characteristics. *Yeast* 37(11):609–21.

67. Ciani M, Comitini F, Mannazzu I, Domizio P. 2010. Controlled mixed culture fermentation: a
new perspective on the use of non-*Saccharomyces* yeasts in winemaking. *FEMS Yeast Res*
10(2):123–33.

68. Fleet GH. 2003. Yeast interactions and wine flavour. *Int J Food Microbiol* 86(1):11–22.

69. Perrone B, Giacosa S, Rolle L, Cocolin L, Rantsiou K. 2013. Investigation of the dominance
behavior of *Saccharomyces cerevisiae* strains during wine fermentation. *Int J Food Microbiol*
165(2):156-62.

70. Wang C, Mas A, Esteve-Zarzoso B. 2015. Interaction between *Hanseniaspora uvarum* and
Saccharomyces cerevisiae during alcoholic fermentation. *Int J Food Microbiol* 206:67–74.

71. Ma M, Liu ZL. 2010. Mechanisms of ethanol tolerance in *Saccharomyces cerevisiae*. *Appl
Microbiol Biotechnol* 87(3):829–45.

72. Henderson CM, Block DE. 2014. Examining the role of membrane lipid composition in
determining the ethanol tolerance of *Saccharomyces cerevisiae*. *Appl Environ Microbiol*
80(10):2966–72.

73. Albergaria H, Arneborg N. 2016. Dominance of *Saccharomyces cerevisiae* in alcoholic
fermentation processes: role of physiological fitness and microbial interactions. *Appl Microbiol
Biotechnol* 100(5):2035–46.

74. Pérez-Torrado R, Rantsiou K, Perrone B, Navarro-Tapia E, Querol A, Cocolin L. 2017. Ecological
interactions among *Saccharomyces cerevisiae* strains: insight into the dominance
phenomenon. *Sci Rep* 7(1):43603.

75. Goddard MR, Greig D. 2015. *Saccharomyces cerevisiae*: a nomadic yeast with no niche? FEMS
Yeast Res 15(3)

76. Kemsawasd V, Branco P, Almeida MG, Caldeira J, Albergaria H, Arneborg N. 2015. Cell-to-cell
contact and antimicrobial peptides play a combined role in the death of *Lachanchea*
thermotolerans during mixed-culture alcoholic fermentation with *Saccharomyces cerevisiae*.
FEMS Microbiol Lett 362.

77. Wang C, Mas A, Esteve-Zarzoso B. 2015. Interaction between *Hanseniaspora uvarum* and
Saccharomyces cerevisiae during alcoholic fermentation. *Int J Food Microbiol* 206:67–74.

78. Albergaria H, Francisco D, Gori K, Arneborg N, Gírio F. 2010. *Saccharomyces cerevisiae* CCMI
885 secretes peptides that inhibit the growth of some non-*Saccharomyces* wine-related
strains. *Appl Microbiol Biotechnol* 86(3):965–72.

79. Aa E, Townsend JP, Adams RI, Nielsen KM, Taylor JW. 2006. Population structure and gene
evolution in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 6(5):702–15.

80. Gibbons JG, Rinker DC. 2015. The genomics of microbial domestication in the fermented food
environment. *Curr Opin Genet Dev* 35:1–8.

81. Steensels J, Gallone B, Voordeckers K, Verstrepen KJ. 2019. Domestication of industrial
microbes. *Curr Biol* 29(10):R381–93.

82. Steensels J, Verstrepen KJ. 2014. Taming wild yeast: potential of conventional and
nonconventional yeasts in industrial fermentations. *Annu Rev Microbiol* 68(1):61–80.

83. Peter J, De Chiara M, Friedrich A, Yue JX, Pflieger D, Bergström A, et al. 2018. Genome
evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature* 556(7701):339–44.

84. Gallone B, Steensels J, Prahl T, Soriaga L, Saels V, Herrera-Malaver B, et al. 2016. Domestication
and divergence of *Saccharomyces cerevisiae* beer yeasts. *Cell* 166(6):1397-1410.e16.

85. Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, et al. 2009. Population genomics
of domestic and wild yeasts. *Nature* 458(7236):337–41.

86. Han DY, Han PJ, Rumbold K, Koricha AD, Duan SF, Song L, et al. 2021. Adaptive gene content
and allele distribution variations in the wild and domesticated populations of *Saccharomyces*
cerevisiae. *Front Microbiol* 12:631250.

87. Friedrich A, Gounot JS, Tsouris A, Bleykasten C, Freel K, Caradec C, et al. 2023. Contrasting
genomic evolution between domesticated and wild *Kluyveromyces lactis* yeast populations.
Genome Biol Evol 15(2).

88. Legras JL, Merdinoglu D, Cornuet JM, Karst F. 2007. Bread, beer and wine: *Saccharomyces*
cerevisiae diversity reflects human history. *Mol Ecol* 16(10):2091–102.

89. Fay JC, Benavides JA. 2005. Evidence for domesticated and wild populations of *Saccharomyces*
cerevisiae. *PLoS Genet* 1(1):66–71.

90. Tapsoba F, Legras JL, Savadogo A, Dequin S, Traore AS. 2015. Diversity of *Saccharomyces*
cerevisiae strains isolated from *Borassus akeassii* palm wines from Burkina Faso in comparison
to other African beverages. *Int J Food Microbiol* 211:128–33.

91.   Fuhrman JA. 2009. Microbial community structure and its functional implications. *Nature*
459(7244):193–9.

92.   Ciani M, Comitini F. 2015. Yeast interactions in multi-starter wine fermentation. *Curr Opin Food*
Sci 1:1–6.

93.   Cordero OX, Datta MS. 2016. Microbial interactions and community assembly at microscales.
Curr Opin Microbiol 31:227–34.

94.   Viljoen BC. 2001. The interaction between yeasts and bacteria in dairy environments. *Int J Food*
Microbiol 69(1):37–44.

95.   Zuñiga C, Zaramela L, Zengler K. 2017. Elucidation of complexity and prediction of interactions
in microbial communities. *Microb Biotechnol* 10(6):1500–22.

96.   De Vuyst L, Leroy F. 2020. Functional role of yeasts, lactic acid bacteria and acetic acid bacteria
in cocoa fermentation processes. *FEMS Microbiol Rev* 44(4):432–53.

97.   Adamu-Governor OL, Shittu TA, Afolabi OR, Uzochukwu SVA. 2018. Screening for gum-
producing Lactic acid bacteria in oil palm (*Elaeis guineensis*) and raphia palm (*Raphia regalis*)
sap from South-West Nigeria. *Food Sci Nutr* 6(8):2047–55.

98.   Kouame HK, Ake MDF, Assohoun NMC, Dje MK, Djeni NT. 2020. Dynamics and species diversity
of lactic acid bacteria involved in the spontaneous fermentation of various palm tree saps
during palm wine tapping in Cote d'Ivoire. *World J Microbiol Biotechnol* 36(5):64.

99.   Somashekaraiah R, Shruthi B, Deepthi B, Sreenivasa MY. 2019. Probiotic Properties of Lactic
Acid Bacteria Isolated From Neera: A Naturally Fermenting Coconut Palm Nectar. *Front*
Microbiol 10:1382.

100. Derikx PJL, Willers HC, ten Have PJW. 1994. Effect of pH on the behaviour of volatile
compounds in organic manures during dry-matter determination. *Bioresour Technol* 49(1):41–
5.

101. Jakl J. 2021. Alcohol in early Java: its social and cultural significance. Boston: Brill, Leiden.

102. Laklaeng S ngob, Kwanhian W. Immunomodulation 2020. Effect of *Nypa fruticans* Palm vinegar.
Walailak J Sci Technol WJST 17(11):1200–10.

103. Leida IM, Thaha RM, Yusnitasari AS, Afsahyana. 2020. Effect of sap palm (*Borassus flabellifer*)
on blood glucose level in pre-diabetic patients. *Int J Curr Res Rev* 12(24):96–100.

104. Mia MAT, Mosaib MG, Khalil MI, Islam MA, Gan SH. 2020. Potentials and safety of date palm
fruit against diabetes: a critical review. *Foods* 9(11):1557.

105. Senghoi W, Klangbud WK. 2021. Antioxidants, inhibits the growth of foodborne pathogens and
reduces nitric oxide activity in LPS-stimulated RAW 264.7 cells of nipa palm vinegar. *PeerJ*
9:e12151.

106. Cardona F, Andrés-Lacueva C, Tulipani S, Tinahones FJ, Queipo-Ortuño MI. 2013. Benefits of
polyphenols on gut microbiota and implications in human health. *J Nutr Biochem* 24(8):1415–
22.

107. Nash V, Ranadheera CS, Georgousopoulou EN, Mellor DD, Panagiotakos DB, McKune AJ, et al.
2018. The effects of grape and red wine polyphenols on gut microbiota – A systematic review.
Food Res Int 113:277–87.

108. Queipo-Ortuño MI, Boto-Ordóñez M, Murri M, Gomez-Zumaquero JM, Clemente-Postigo M,
Estruch R, et al. 2012. Influence of red wine polyphenols and ethanol on the gut microbiota
ecology and biochemical biomarkers. *Am J Clin Nutr* 95(6):1323–34.

109. Kurniawan T, Jayanudin J, Kustiningsih I, Adha Firdaus M. 2018. Palm sap sources,
characteristics, and utilization in Indonesia. *J Food Nutr Res* 6(9):590–6.

110. Beckner M, Ivey ML, Phister TG. 2011. Microbial contamination of fuel ethanol fermentations:
Bioethanol contamination. *Lett Appl Microbiol* 53(4):387–94.

111. Cavalieri D, McGovern PE, Hartl DL, Mortimer R, Polsinelli M. 2003. Evidence for *S. cerevisiae*
fermentation in ancient wine. *J Mol Evol* 1:S226-232.

112. Medina K, Boido E, Fariña L, Gioia O, Gomez ME, Barquet M, et al. 2013. Increased flavour
diversity of Chardonnay wines by spontaneous fermentation and co-fermentation with
Hanseniaspora vineae. *Food Chem* 141(3):2513–21.

113. Rodríguez-Lerma G k., Gutiérrez-Moreno K, Cárdenas-Manríquez M, Botello-Álvarez E,
Jiménez-Islas H, Rico-Martínez R, et al. 2011. Microbial ecology studies of spontaneous
fermentation: starter culture selection for prickly pear wine production. *J Food Sci*
76(6):M346–52.

114. Schweinberger M. 2024. Network Analysis using R. Available from:
https://ladal.edu.au/net.html.

Figure legends

Fig. 1 Schematic to describe the processing of palm sap with its derivative products. Created with BioRender.com

Fig.2 A survey of palm wine studies which include measures of microbial ecology and metabolite profiling (n= 47), details are available on **Supplementary Table S1 & S2**. A) Country distribution of scientific studies in both microbes and metabolites across continents; B) The Venn diagram shows the focus of compiled scientific studies; C) Proportion of the studies of palm species for all compiled studies.

Fig 3. Microbial dynamics and community proportion of palm wine fermentation. Next generation sequencing (NGS) data of 7 abstracted studies with data from Sumerta et al. (2024); **Supplementary Table S4**) were analysed together. A) Microbial diversity loss (measured as Shannon index) over 24 hours palm wine fermentation (Wilcoxon test; p-value <0.05); B) Change in top 10 abundance of microbial species in 24 hours palm wine fermentation. C) The proportion of top 10 microbial species, obtained by CD (inner circle; n =33) compared to CI (outer circle; bacteria n=7-1, excluded due to technical limitation; fungi n = 5).

Fig 4. Network analysis of the relationship of microbial group in palm wine fermentation. The contribution of each group is represented as an edge. Different colour showing different microbial group (blue= yeasts, red= bacteria), then different size of font represents the abundance value. The data set used for this study consists of 23 studies based on the abundance and frequency each species group in **Supplementary Table S3**, omitting 7 studies of NGS, where data was abstracted in **Supplementary Table S5** and used R studio adopting the codes by Schweinberger (2024).

Fig. 5 Schematic of the process of palm wine fermentation which summarises the microbial and chemical composition, dynamics of microbial communities, raw materials, and metabolites over time. A gradual pattern of palm wine fermentation was reconstructed from compiled studies (n= 10, **Supplementary Table S6**) and table shows the examples of chemical yield during the fermentation process, created with BioRender.com.

Fig. 6 Correlation heatmap between microbial and metabolite groups (n= 23-3, studies which used amplicon sequencing were disregarded). The frequency of microbial and metabolite groups were abstracted from **Supplementary Table S3** and analysed with Spearman correlations (p value > 0.05).

Figure 1

814
815   Figure 2
816

817
818   Figure 3
819

820
821   Figure 4
822

Criteria	Early	Middle	Final	Reference
Total Sugars	11%	5%	3%	Amoa-Awua et al., 2007
Ethanol	34 g/L	35 g/L	30 g/L	Tapsoba et al., 2016
Acetic acid	0.50%	0.60%	3%	Amoa-Awua et al., 2007
Lactic acid	0.20%	0.30%	2.80%	Amoa-Awua et al., 2007
pH	4.38	3.54	3.53	Alcantara-Hernandez et al. 2020
Total Phenolics	70 mg/L	100 mg/L	150 mg/L	Das & Tamang 2023
Total Flavonoids	40 mg/L	90 mg/L	150 mg/L	Das & Tamang 2023

823
824   Figure 5
825

826