Novel sulfatase cancer therapeutics negatively impact Bacteroidota of the colonic microbiota in a non-sulfatase dependent manner

Conor J. Crawford[1], Christian Gunawan[2], Zongjia Chen[2], Dominic P Byrne[3], Cosette Darby[4], Ana S Luis[5], Stefano Elli[6], Edwin A Yates[3], David N Bolam[4], Sjoerd van der Post[5], Spencer J Williams[2]*, and Alan Cartmell[7,8,9]*

[1]Max Planck Institute of Colloids and Interfaces, Am Mühlenberg 1, 14476, Potsdam, Germany
[2]School of Chemistry and Bio21, Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria3010, Australia
[3]Department of Biochemistry, Cell and Systems biology, Institute of Systems, Molecular and Integrative biology, University of Liverpool, Liverpool L69 7ZB, U.K.
[4]Biosciences Institute, Faculty of medical sciences, Newcastle university, Medical school, Framlington place, Newcastle upon tyne, NE2 4HH, U.K.
[5]Department of Medical Biochemistry and Cell Biology, University of Gothenburg, Gothenburg, Sweden
[6]Istituto di Ricerche Chimiche e Biochimiche G. Ronzoni, Milan,20133, Italy
[7]Department of Biology, University of York, Wentworth Way, York, YO10 5DD, U.K.
[8]York Structural Biology Laboratory (YSBL), University of York, Wentworth Way, York, YO10 5DD, U.K.
[9]York Biomedical Research Institute (YBRI), University of York, Wentworth Way, York, YO10 5DD, U.K.

*To whom correspondence should be addressed:
alan.cartmell@york.ac.uk
sjwill@unimelb.edu.au

Abstract

The S1 family of sulfatases is the sole family of enzymes that desulfate carbohydrates, linking them to inflammatory bowel diseases and various cancers. There is an unmet need for therapeutics against carbohydrate sulfatases, and while effective, orally available inhibitors of S1 steroid sulfatases exist, their efficacy against S1 carbohydrate sulfatases remains largely unexplored. In this study we assess a library of aryl- and carbohydrate sulfamates/sulfonates for their ability to inhibit carbohydrate sulfatases from human colonic bacteria. Surprisingly, these compounds prove ineffective. We show that arylsulfamate inhibitors inhibit the growth of anti-cancer human gut microbiota species independent of sulfatase activity. Leveraging thermal proteome profiling, we identify the non-sulfatase targets responsible for these effects. This work highlights the imperative for developing specific inhibitors targeting carbohydrate sulfatases and unveils the adverse effects of arylsulfamates, a class of drugs designed for hormone cancer treatment, on the human gut microbiota, potentially influencing their therapeutic efficacy.

Introduction

Sulfatases are enzymes that perform the enzymatic desulfation of various biomolecules.

These biomolecules include small molecules, such as hormones, and larger

macromolecules, such as proteins and complex carbohydrates. The SulfAtlas database

classifies sulfatases into four families, S1, S2, S3, and S4[1]. Of the over 160,000 sequences

catalogued in the SulfAtlas database, the majority (over 90%, or >145,000 sequences)

belong to the S1 family[2]. The S1 family occurs across all domains of life and members

desulfate all known classes of sulfated biomolecules[1,3,4]. The remaining approximately 10%

of sulfatase sequences are distributed across the S2, S3, and S4 families. These are

exclusive to bacteria and are active on alkyl- and arylsulfates[1].

The significance of the S1 sulfatase family lies in its size and distribution, with 110

subfamilies based on sequence homology. The dysfunction of S1 sulfatases is implicated in

a range of human diseases. Human steroid sulfatase (STS) plays a critical role in catalysing

the desulfation of the hormones dehydroepiandrosterone (DHEA) sulfate and esterone (E1)

sulfate[5], as well as other oestrogen-based hormones, converting them into their unsulfated

active forms[6]. Elevated levels of STS are correlated with increased hormone levels in breast

cancer tumours, primarily in post-menopausal women, leading to poorer patient prognosis[5].

Furthermore, STS activity has also been linked to prostate cancer in men, influencing the

production of testosterone and androstenediol from DHEA sulfate[7].

The S1 human carbohydrate sulfatases Sulf1 and Sulf2, catalyse desulfation of the

glycosaminoglycan (GAG) heparan sulfate at the *O6* position of *N*-acetyl-D-glucosamine

(GlcNAc) within the extracellular matrix (ECM)[8,9]. Both Sulf1 and Sulf2 are overexpressed in

most cancers, and their heightened expression is essential for metastasis in small cell lung

carcinomas[10,11]. Bacterial S1 carbohydrate sulfatases are essential for some symbiotic

colonic bacteria to colonise and utilise host colonic mucin and have been implicated in

ulcerative colitis (UC)[12,13], and colon cancer (CC)[14,15]. Excessive activity of these bacterial

sulfatases leads to over-degradation and alteration of the protective colonic mucin layer,

exposing the gut epithelium to the normally symbiotic human gut microbiota (HGM). This

exposure leads to gut inflammation, microbial dysbiosis, and the development of disease

states such as UC[12,13,16] and CC[14,17]. Inhibitors targeting S1 sulfatases may prove effective in

treating these diseases. Some promising inhibitors of the S1 family have already been

developed[18,19], offering potential approaches to treat conditions like UC and CC.

The S1 family of sulfatases has a conserved alkaline phosphatase-like fold

characterized by a larger N-terminal domain housing a central mixed beta sheet flanked by

alpha helices. This domain is abutted by a smaller C-terminal subdomain comprising a 4

stranded antiparallel beta sheet, and a single alpha helix[20] **(Figure 1a)**. Within this structural

framework the S1 family employs a non-canonical amino acid as the catalytic nucleophile
known as formylglycine (FGly). FGly is generated co-translationally from a Cys or Ser within
the consensus sequence C/S-X-P/A-X-R, and is situated within an invariant sulfate (S)
binding site[3] **(Figure 1a)**. The arylsulfamate class of inhibitors have been developed to
target the distinctive FGly residue of the S1 family.[21]

Arylsulfamate inhibitors have the general formula $Ar-O-SO_2NH_2$. These inhibitors,
developed by replacing the sulfate group with a sulfamate on aryl scaffolds, prove to be
effective, irreversible, active site directed inhibitors of S1 steroid sulfatases. Despite their
efficacy, the precise nature of the dead-end complex formed is unknown[21]. Arylsulfamates
demonstrate broad effectiveness against S1 steroid sulfatases of both bacterial and
eukaryotic origin[18]. The initial development of arylsulfamates targeted human STS for breast
cancer treatment. Arylsulfamates mimic sulfated steroids, the natural substrates of STS.
Notable arylsulfamate inhibitors of STS are Irosustat[22], estrone 3-O-sulfamate (EMATE)[23],
and estradiol sulfamate (E2MATE)[24] **(Figure 1b)**. Irosustat, a first-in-class STS inhibitor has
progressed into phase II clinical trials[25]. It has shown positive outcomes in treating oestrogen
receptor-positive breast cancer in post-menopausal women, both in early[26] and late-stage
cancers[25], and oestrogen receptor-positive endometrial cancer[27]. EMATE, the first oestrogen
sulfamate tested against cancer cells, exhibits almost complete reduction in STS activity and
a reduction in mammary tumour growth in rats upon both oral and subcutaneous
administration[28]. E2MATE, originally developed as an estradiol prodrug, reached phase II
trials for hormone replacement therapy, demonstrating effectiveness in a clinical trial[28].
Additionally, it was explored as an STS inhibitor for treatment of endometriosis, and showed
a 91% reduction in endometrial STS in a phase I clinical trial of 24 healthy women of
reproductive age[28]. All three inhibitors exhibit good bioavailability, are well tolerated, and
have nanomolar activities against STS[22,25,27,28].

Given the invariant nature of the sulfate binding site, it was initially assumed that
sulfamate based inhibitors would be universally effective across all sulfatases within the S1
family. However, limited tests have suggested arylsulfamates are ineffective inhibitors of S1
carbohydrate sulfatases[29,30], but this has yet to be systematically investigated. The
specificity and a significant portion of the binding energy of carbohydrate sulfatases are
dictated by their glycan binding regions, which target hydrophilic carbohydrates rather than
hydrophobic steroids[14,20], and is line with low activity on arylsulfates. Consequently, efforts
have been made to incorporate sulfamate groups into carbohydrate substrates of S1
carbohydrate sulfatases, aiming to achieve results similar to those observed for S1 steroid
sulfatases.

While some success has been reported against the human carbohydrate sulfatase
Sulf1 using sulfamate-modified disaccharide, trisaccharide, and tetrasaccharide substrate

analogues, these compounds functioned as reversible competitive inhibitors[31,32]. This
suggests a fundamental change in the reactivity of the sulfamate group when transitioning
from an aryl to a carbohydrate scaffold. There has not been systematic analysis of the
effectiveness of sulfamate inhibitors, nor the newer sulfur-based covalent modifiers such as
vinyl sulfones and fluorosulfates, against S1 carbohydrate sulfatases, nor has there been an
assessment of the impact of these inhibitors on the HGM. *Bacteroides* species within the
HGM encode a large number of S1 sulfatases[33,34] and may be susceptible to arylsulfamates
themselves, making this point relevant for two reasons: 1) *Bacteroides* species are crucial
for positive outcomes for patients undergoing chemotherapy for pancreatic ductal
adenocarcinoma (PDAC)[35], which may be applicable to a broader range of cancers; 2)
concentrations of orally ingested drugs are elevated in the colon compared to plasma, and
recent data suggests this significantly influences the HGM[36,37].

Here, we examine the ability of a library of aryl- and carbohydrate
sulfamates/sulfonates, as well as vinyl sulfones and fluorosulfates, **(Figure 1c,d)** to inhibit
carbohydrate sulfatases from human colonic bacteria, demonstrating a lack of effectiveness.
Furthermore, we show that arylsulfamate inhibitors, which are delivered orally to treat
cancer, inhibit the growth of anti-cancer HGM Bacteroidota species in a non-sulfatase
dependent manner, but these effects can be overcome through growth on select complex
glycans. Finally, utilising thermal proteome profiling we identify the non-sulfatase targets
through which these effects are mediated.

Results
Arylsulfamates do not inhibit S1 HGM carbohydrate sulfatases
In our initial experiments, we evaluated a panel of arylsulfamates to assess their inhibitory
potential on ten recombinantly expressed, S1 carbohydrate sulfatases from the HGM
members *Akkermansia muciniphila ATCC BAA-835 (A. muc*) and *Bacteroides*
thetaiotaomicron VPI-5482 (*B. theta*). As a positive control we conducted assays on our
arylsulfamate library, which included three negative control methylated sulfamates. We
included two S1 steroid sulfatases: PaAstA, belonging to S1_4 and from *Pseudomonas*
aeruginosa, and the commercially available snail sulfatase from *Helix pomatia* (HpSulf), from
S1_2. The efficacy of each compound was determined by measuring the sulfatase activity of
either enzyme incubated in the presence of inhibitor, or following ~24h pre-incubation and
jump-dilution 'washout'[38]. At a concentration of 1 mM, all the arylsulfamates were broadly
equal in their ability to completely inhibit both enzymes with only minor differences observed
between the two **(Figure 2, Figure S1-S4, Table S1 and S2).** Against PaAsta, partial
sulfatase activity could be measured for 17 and 18 before complete inactivation **(Figure 2a,**
left panel and Figure S1), and this residual activity could be eliminated by pre-incubation

with the two compounds **(Figure 2a, right panel)**. Similarly, for HpSulf, initial rates that
could be measured for 8 and 14 were blocked in the jump dilution experiment **(Figure 2a**
and Figure S3). These findings suggest different on rates between the enzymes and are
consistent with the irreversible, covalent, nature of the inhibitor compounds. Importantly,
near wildtype activity was observed for the *N* methyl and *N, N* dimethyl sulfamate control
compounds 2, 5, and 6 **(Figure 2a, right panel)**.

Having confirmed the effectiveness of the arylsulfamates inhibitors against two S1
steroid sulfatases we next assayed their effectiveness against 10 bacterial carbohydrate
sulfatases that target common linkages found in host glycans: four from S1_11 (BT3177[6S-
GlcNAc], BT4656[6S-GlcNAc], Amuc1033[6S-GlcNAc], and Amuc1074[6S-GlcNAc]) targeting *O*6 sulfated *N*-
acetyl-D-glucosamine (6S-GlcNAc), four from S1_20 (BT1636[3S-Gal], BT1622[3S-Gal/GalNAc],
Amuc0451[3S-Gal], and Amuc0491[3S-Gal/GalNAc]) targeting *O*3 sulfated D-galactose/*N*-acetyl-D-
galactosamine (3S-Gal/GalNAc), and two from S1_16 (BT3796[4S-Gal/GalNAc] and Amuc1755[4S-
Gal]) targeting *O*4 sulfated D-galactose/*N*-acetyl-D-galactosamine (4S-Gal/GalNAc). No
significant inhibition was observed for any of the enzymes when assaying their activity with 1
mM compound, even after pre-incubation **(Figure 2b, Supplementary Figure S5-S24,**
Table S1 and S2). This indicates that the arylsulfamate compounds do not inhibit S1 HGM
carbohydrate sulfatases.

Arylsulfamates affect the growth of Bacteroides thetaiotaomicron
Given that arylsulfamate inhibitors of STS are administered orally[25,27], exposing members of
the HGM such as *B. theta* in any treatment regimen, we aimed to explore their impact on the
growth of HGM species. We cultivated *B. theta* in brain heart infusion (BHI) media, and in
minimal media supplemented with various carbon sources including D-glucose (Glc),
chondroitin sulfate C (CSC), heparin (Hep), larch arabinogalactan (LAG), and potato
galactan (PG), in the presence of each arylsulfamate compound at 1 mM. When DMSO was
used at a final concentration of 5 % DMSO (as used in the *in vitro* assay) *B. theta* failed to
grow in minimal medium supplemented with Hep or LAG, and grew poorly with CSC and
Glc. Instead, a final DMSO concentration of 1 % was used, which did not affect growth of *B.*
theta under any minimal media conditions tested **(Figure S25)**.

Interestingly, we observed variable bacterial growth rates with different compound
and carbon source combinations **(Figure 3 and Figure S26-S31)**. Consistently however,
arylsulfamates based on a napththyl scaffold (9, 10, 11, 12, 13, 14, and 16) effectively
inhibited growth in all conditions. In contrast, the effect of the phenyl-based compounds (1,
2, 4, 17 and 18) varied substantially. Whilst we observed little effect of phenyl-based
compounds for bacteria grown on BHI and Glc, growth on the sulfated host glycan CSC was
strongly perturbed by 1 and 5. Moreover, there was an increase in lag time with 4 and 17,

and total cessation of growth for 18, on the sulfated host glycan Hep **(Figure 3)**. Similarly to
CSC, large growth defects were caused for 1 and 5 when using the unsulfated plant glycan
PG as a carbon source. For the unsulfated plant glycan LAG, 1 and 17 caused elongation of
the lag phase whilst 4 completely inhibited growth. Interestingly, compound 3, which is a 4-
phenylphenyl sulfamate, inhibited growth under all conditions tested. When *B. theta* was
grown on BHI and Glc, the coumate scaffolds (5, 6, and 7) displayed little to no effect. These
compounds, however, increased the lag phase and reduced growth when CSC was the
carbon source but completely inhibited growth with Hep. On the plant glycan LAG 5 and 7
increased the lag phase and reduced growth whilst 6 completely inhibited growth. By
contrast, when PG was the carbon source the effects of 5, 6, and 7 were relatively mild.

The variable impact of arylsulfamates on the growth of *B. theta*, observed across
both sulfated and unsulfated glycan substrates, aligns with the *in vitro* biochemical data
showing that these compounds do not target S1 carbohydrate sulfatases. Additionally, the
fact that the 3 negative control methyl sulfamate compounds (2, 5, and 6) also influenced the
growth *B. theta* provides further support that the observed growth defects are not mediated
through an S1 sulfatase dependent mechanism.

Inhibition assays using cell lysates show carbohydrate sulfatase activity
To confirm the hypothesis that the growth defects observed on the sulfated host glycans
were not due to S1 carbohydrate sulfatase inhibition, and that the lack of inhibition on
recombinantly produced enzymes was not due to any improper FGly formation, we next
grew *B. theta* to mid-exponential growth phase on chondroitin sulfate A (CSA) and Hep to
stimulate upregulation of native GAG S1 carbohydrate sulfatases. Cells were then harvested
by centrifugation, washed, and resuspended in PBS, before being lysed and incubated with
two arylsulfamates inhibitors for 1 hour before exposure to sulfated mono- or disaccharide
substrates. Compounds 3 and 12, for which kill curves were plotted **(Extended data 1a and**
Figure S32), and were effective at inhibiting growth on all carbon sources tested (**Figure 3**),
did not impact the production of the final desulfated products (GlcNAc for Hep substrates
and GalNAc for CSA substrates, as predicted from the combined action of endogenous S1
carbohydrate sulfatases and glycoside hydrolases), as judged by TLC and HPAEC
(Extended data 1b,c). This confirms that compounds 3 and 12 are not effective inhibitors of
the GAG S1 carbohydrate sulfatases produced *in vivo*, and that observed effects on
bacterial growth were likely due to off target interactions.

Carbohydrate sulfamates failed to inhibit the S1_20 sulfatase BT1636[3S-Gal]
We hypothesised that the refractory nature of S1 carbohydrate sulfatases to arylsulfamate
inhibition may be related to an inability of the compound to engage with the glycan binding

region of the enzymes. This is consistent with the inactivity of S1 carbohydrate sulfatases
towards 4-nitrophenol-sulfate (4NP-SO$_3$), which in contrast is rapidly hydrolysed by the
steroid sulfatase PaAsta, and emphasises the divergent substrate specificities (and by
extension inhibitor profiles) of the two enzyme classes **(Extended data figure 2)**. We next
selected the S1_20 sulfatase BT1636[3S-Gal] to test if carbohydrate sulfamates would act as
effective irreversible, covalent, active site inhibitors. BT1636[3S-Gal] desulfates 3-*O* sulfated D-
galactose (3S-Gal) and is essential for *B. theta* to utilise colonic mucin *O*-glycans, and
competitively colonise the colon in mice[14].

The substrate analogues, 3-*O* sulfamate D-galactose (23) and 3-*O* vinylsulfonyl D-
galactose (25) were synthesised, as well the control compound 2-*O* sulfamate D-galactose
(24), and screened alongside aryl vinyl sulfone analogues 19 and 22, aryl fluorosulfate (20),
and aryl sulfamate (21) for inhibitory activity towards BT1636[3S-Gal] **(Figure 4a)**. For the aryl
compounds, 20 and 22 showed moderate inhibition after pre-incubation with BT1636[3S-Gal]
(Figure 4b, Figure S33, S34, and Table S3). For the carbohydrate scaffold 24, which
possesses a 2-*O* sulfamate, as expected, showed no inhibition but surprisingly neither did
the 3-*O* sulfamate 23. Despite the moderate inhibition activity by 22 (aryl vinyl sulfone), the
D-galactose variant (25), showed no activity **(Figure 4b, Figure S33, S34, and Table S3)**.
Synthesis of fluorosulfate armed D-galactose could not be achieved and therefore could not
be tested. Thermal shift assays (TSA) performed using wildtype Ser77 form of BT1636[3S-]
[Gal], which is inactive when expressed in *E. coli*, (as *E. coli* can only covert Cys to FGly),
showed no change in melting temperature (indicative of compound-binding) for 23, 24, or 25
at 0.01, 0.1, or 1.0 mM compound concentrations, whilst both 21 and 22 destabilised the
protein (**Figure 4c, Figure S35, and Table S4**). In contrast, the canonical substrate of
BT1636[3S-Gal], 3S-Gal, potently stabilised the protein (ΔTm = 0.10, 0.98, and 3.02 ºC) at
equivalent concentrations **(Figure 4c, Figure S35 and Table S4)**. Collectively, these data
demonstrate that the carbohydrate sulfamate 23, and its vinylsulfonyl equivalent 25, are not
effective inhibitors of BT1636[3S-Gal] and appear unable to bind to the enzyme.

The effects of arylsulfamates apply across multiple HGM Bacteroidota species
We next selected 10 additional HGM Bacteroidota species to see if the growth effects
observed for *B. theta* in the presence of arylsulfamates applied more broadly. We conducted
growth experiments using the arylsulfamates 3 and 12, as they inhibited the growth of *B.*
theta across all conditions tested (**Figure 3**), as well the phase II arylsulfamate drugs
Irosustat and estradiol sulfamate. No growth was observed for any species in the presence
of arylsulfamate 3, except *Phocaeicola plebius* (*P. plebius*) which had highly variable growth
(Figure 5 and Figure S36). The inhibitory effect of arylsulfamate 12 treatment was milder,
but all species (and in particular *B. cellulosilyticus* for which no growth could be detected)

still exhibited growth defects. Irosustat potently inhibited growth, with all but one species (*B.*
fragilis) showing limited or no growth **(Figure 5 and Figure S37)**. Surprisingly, only mild
growth defects could be detected with the drug estradiol sulfamate **(Figure 5 and Figure**
S37).

It is clear that the effects of arylsulfamates apply broadly across the *Bacteroides* and
Phocaeicola genera, but the response varies significantly by arysulfamate structure, and,
although less widely, by species. Estradiol sulfamate (EMATE2) is a structural mimetic of the
endogenous human steroid hormone, estradiol sulfate, and this chemical resemblance to a
common host product may partly explain the reduced impact on the HGM species tested
here.

279
Thermal proteome profiling to identifies off target interactions
We next took the arylsulfamate compounds 3 and 12, and determined the effect on thermal
protein stability across the proteome using thermal proteome profiling (TPP) in *B. theta* cell
lysates grown on CSA as a substrate. TPP is a non-targeted mass spectrometry technique
which can identify protein-compound interactions based on acquired changes in a protein's
thermal stability. This is quantified as an increase in the relative abundance of peptides
originating from the soluble protein fraction over a temperature gradient following incubation
with a ligand[39]. Protein lysates were heat treated at 10 different temperatures ranging from
45 – 72 °C in the presence of inhibitor 3, 12 or DMSO solvent control.

The *B. theta* lysate proteome showed a narrow melt temperature ranging between 50
and 58 °C (90% range), much narrower than that reported for *E. coli* from whole cell TPP
experiments **(Extended data 3a)**. A total of 2550 proteins were identified in the combined
analysis, from which 11 S1 sulfatases, including the three upregulated in response to CSA
(BT1596, BT3333, and BT3349)[40] **(Extended data figure 3b,c)** were identified. The melting
temperature (Tm) was established for 2112 proteins **(Extended data figure 3d)** based on
the criteria of detecting at least two unique peptides in a minimum of two replicate analysis
(see methods). This allowed the determination of the melting curves for 8 out of the 11
identified S1 sulfatases **(Extended data figure 4)**. The protein stability of none of the S1
sulfatases was significantly affected upon treatment with compound 3 or 12 suggesting no
direct interaction with the compounds. This is in line with the hypothesis that arylsulfamate
inhibitors do not exert their effects on *B. theta* via S1 sulfatases.

A total of 7 proteins for arylsulfamate 3 and 4 proteins for arylsulfamate 12 were
identified as having significant changes in Tm indicating a potential interaction **(Figure 6a, b,**
c). Some of these hits can be discounted as causing the defective growth phenotype:
BT0600, BT2504, and BT3464 were recently disrupted through transposon mutagenesis
and no phenotype was reported for these genes when grown on a supplemented BHI

media[41]. BT0133 is a homologue of the *E. coli* protein murQ (having 45 % identity), a protein

which produces 6-*O* phosphorylated GlcNAc from 6-*O* phosphorylated *N*-acetyl-D-muramic

acid (MurNAc). Loss of MurQ in *E. coli* does not cause a no growth phenotype[42] and no

MurNAc was present in our experiments. This leaves a total 7 unique proteins (BT0218,

BT1443, BT1889, BT4322, BT4335, BT4346, and BT4480) through which an inhibitory

arylsulfamate interaction could drive the observed growth defects, and of which only one

was targeted by both arylsulfamate molecules **(Figure 6b,c)**; the uncharacterised protein

BT4322.

BT4322 is a distant homologue of the cytoplasmic diacylglycerol kinases (DGK)

YerQ and DgkB, from the gram-positive organisms *Bacillus subtilis*[43] and *Staphylococcus*

aureus[44], respectively, and the human sphingosine kinase SphK1[45]. YerQ is essential for the

production of the cell wall component, lipoteichoic acid (LTA), and growth of *B. subtilis*[43].

BT4322 is predicted to have the same fold and catalytic apparatus as both DgkB

(PDB:2QV7) and SphK1 (PDB:3VZB), as well as similar ATP binding site motifs and lipid

binding pockets **(Extended data figure 5)**. BT4322 is therefore highly likely to be a *B. theta*

lipid kinase, but the nature of the lipid substrate is yet to be determined. Although LTA is

specific to gram positive bacteria, BT4322 may have an analogous role to YerQ in *B. theta*

membrane lipid metabolism; this would be consistent with the lack of growth phenotype in

the presence of compounds 3 and 12. Furthermore, a blastp of BT4322 against the 10 other

compound-sensitive bacteria (**Figure 4**) returned a single orthologue from each species,

with a 100 % query coverage, and a minimum of 84 % identity for the *Bacteroides* species

and 73 % for *P. plebius*, showing a high degree of conservation and lack of redundancy

between these species **(Extended data figure 6 and Table S5)**, and suggesting an

important role for these proteins.

Diacylglycerol kinase inhibitors significantly inhibit the growth of Bacteroides species

With BT4322 being a lipid kinase we next sought to see if we could reproduce the

arylsulfamate induced growth defects using commercially available sphingosine and

diacylglycerol kinase inhibitors. Sphingosine kinase inhibitor (SKI) had a maximum solubility

of 0.1 mM and no notable growth defect could be observed **(Figure 6d and Extended data**

figure 7). Sphingosine kinase 1 inhibitor (SK1I)-II dissolved to 0.5 mM and caused mild

phenotypes in four species, manifesting as an elongated lag phase, except for *B.*

xylanisolvens which showed severe defects, especially in its lag phase **(Figure 6d and**

Extended data figure 7). The solubility of diacylglcerol kinase inhibitor (DAGKI)-I and II was

even lower at 125 μM and 62.5 μM, respectively. DAGKI-I was effective at inhibiting the

growth of all species with only *B. cellulosilyticus*, *B. intestinalis*, and *B. oleicipnenius* showing

significant growth, albeit highly variable with very extended lag phases **(Figure 6d and**

Extended data figure 8). DAGKI-II showed a milder phenotype but still inhibited the growth
of all species, lowering final OD and extending the lag phase, except for *B. oleicipnenius*
which reached a similar OD to non-inhibitor growths **(Figure 6d and Extended data figure**
8). These data support the view that the arylsulfamate observed growth defects are
mediated through inhibition of a diacylglycerol kinase, not a sphingosine kinase, of which
BT4322 is the likely target.

Ligand docking of arylsulfamates into off-target proteins
BT4322 emerged from the TPP analyses as the top protein target through which the
arylsulfamates mediated their off-target effects, and as such we performed blind ligand
docking (no pocket specified) with arylsulfamates 3, 12 and Irosustat using the AF2
predicted model of BT4322. The docking results for Irosustat with BT4322 predicted binding
almost exclusively to the lipid pocket, with some minor populations elsewhere. The situation
was similar for arylsulfamate 3 and 12, but some binding in the ATP site was also predicted
(Extended data figure 9). Interestingly, when the same experiment was performed for the
human sphingosine kinase, SphK1, no specific site was favoured **(Extended data 6)**. These
results suggest that *in vivo* arylsulfamates bind to human S1 steroid sulfatase STS but not
SphK1, but reciprocally in *Bacteroides* species they bind lipid kinases (distant SphK1
orthologues), but not carbohydrate S1 sulfatases. This would be in line with the
arylsulfamates, such as Irosustat, being well tolerated in humans with relatively mild side
effects.

Discussion
Arylsulfamates are effective irreversible, active site targeting, inhibitors of S1 steroid
sulfatases[21] and display relatively mild side effects when administered to humans to treat
hormone dependent cancers in phase II clinical trials[25,26]. S1 carbohydrate sulfatases have
been implicated in driving several disease states and it was presumed that arylsulfamates
would be broadly effective across the S1 family, but it is clear from the work presented here
that bacterial S1 carbohydrate sulfatases of the HGM, despite sharing the invariant sulfate
binding subsite (S subsite), are unaffected by arylsulfamates both *in vitro* and *in vivo*.
Furthermore, the generation of an equivalent carbohydrate sulfamate not only failed to inhibit
the target S1 carbohydrate sulfatase, but showed no evidence of binding **(Figure 4b,c)**.
Collectively, these data demonstrate that sulfamate based inhibitors are not effective against
the HGM S1 carbohydrate sulfatases tested here. The addition of an amino group to the
sulfur atom must alter both its binding and/or reactivity in the S subsite. Similarly, the vinyl
sulfone and fluorosulfate modifications also do not appear to be well tolerated. It is clear S1
steroid sulfatases can accommodate the changes required for arylsulfamate reactivity, but

the S1 carbohydrate sulfatases tested here cannot. With the S subsite being invariant, the
apparent differing susceptibilities of these enzyme classes must be due to the distinctive
chemical environments and/or structural geometries of glycan versus steroid binding
subsites. Steroid binding sites have evolved to provide a hydrophobic, 'greasy', patch that
can accommodate lipophilic molecules and may allow the aryl groups of arylsulfamates to
adopt an alternative binding configuration relative to the natural steroid substrate, and thus
facilitate binding of a sulfamate group in the S subsite. In contrast, carbohydrates are
hydrophilic, and S1 carbohydrate sulfatases have polar substrate binding regions which form
extensive hydrogen bonding networks[14,20,46,47]. The polar, hydrogen bond-driven nature of
the glycan binding site may impose a level of rigidity on substrate interactions which may be
incompatible with alternate binding modes that may be necessary to accommodate
sulfamate groups.

Carbohydrate sulfamate inhibitors have been developed against the human S1
carbohydrate sulfatase Sulf1[30,31]. The authors replaced the labile sulfate in HS di-, tri, and
tetrasaccharides with a sulfamate. These inhibitors have a competitive mode of action with
IC_{50} values of ~700, ~30, and ~0.5 μM for the di-, tetra-, and triasaccharide sulfamates,
respectively. The trisaccharide substrate has a K_M of ~26 μM, ~50 fold higher than the IC_{50}
value for the trisaccharide sulfamate. However, it should be noted that the data from which
the K_M was derived contain only 5 data points, 4 of which are close to, or at, V_{max} and 1 at
the K_M. It is therefore likely that this K_M calculation is not accurate, and the true K_M will be
substantially lower[31]. Taking this into account, and the fact HS has a disaccharide repeat, it
is likely that the affinity of the trisaccharide sulfamate is largely driven by glycan binding, not
the presence of the sulfamate groups, as only the trisaccharide could occupy 3 subsites
spanning -1, 0, +1. The di- and tetrasaccharide sulfamates could only occupy either -1, 0 or
0, +1 subsites depending on the orientation of the bound sugar chain. Despite this, the
carbohydrate sulfamate analogues did bind to the *endo*-acting Sulf1 which is in stark
contrast to the *exo*-acting BT1636[3S-Gal] assayed here[31,32], suggesting the *endo*-acting S1
carbohydrate sulfatases may have increased tolerance for the binding of sulfamate groups in
their S subsite.

Despite a lack of activity towards carbohydrate sulfatases, the panel of
arylsulfamates tested here profoundly impacted the growth of 10 HGM *Bacteroides*, and 1
Phocaeicola species, both from the Bacteroidota (formerly Bacteroidetes) phylum. When
considering arylsulfamates as part of hormone dependent cancer regimens, this could have
a significant impact on their long-term efficacy. It was recently shown that in PDAC, the
species *B. theta and B. fragilis* were required for effective response to the chemotherapy
regimen FOLFIRINOX[35]. This was mediated through the production of the HGM metabolite,

3-IAA, which is found systemically, increases ROS production, and reduces autophagy in

cancer cells; compromising their metabolic fitness and proliferation. The arylsulfamate drug,

Irosustat, has had positive outcomes for both early and late breast cancer treatment in

phase II clinical trials[25,26] but its effect on the HGM, and how this affects Irosustat's efficacy,

has not been considered. Our data show that Irosustat, and arylsulfamates of varying

structure, negatively impact HGM Bacteroidota species, including those identified as helpful

in PDAC. Our data, also show that some structures from our library are less detrimental than

others, and some of these effects can be mitigated through carbon source selection.

Importantly, the phase II arylsulfamate drug estradiol sulfamate (EMATE2), tested for

hormone replacement therapy and endometriosis[21,28], had very little effect on the growth of

HGM Bacteroidota species. It is possible that arylsulfamate based cancer treatment could be

improved by ameliorating their negative impacts on the HGM. This could be achieved

through the generation of compounds with altered chemical structures and/or by co-

application of selected complex glycan prebiotics.

The most likely target through which the arylsulfamates mediate their growth

inhibiting effects on the HGM species was BT4322, a lipid kinase. A single homologue of

BT4322, with high sequence identity, is found in the 10 other species examined here,

suggesting lack of functional redundancy. Furthermore, a homologue of BT4322 has been

genetically knocked out in *Bacillus subtilis*, which displayed a no growth phenotype, and is

implicated in peptidoglycan metabolism and cell membrane integrity[43]. If BT4322 has a role

in phospholipid biosynthesis, it may be a critical regulator of membrane integrity. Moreover,

accumulation of DAG in *E. coli* membranes prevents growth under stress conditions[48], and

the loss of YerQ in *B. substilis* also causes DAG accumulation in cell membranes,

confirming important roles for two distantly related BT4322 homologues[43]. Unusually for

bacteria, *Bacteroides spp.* also contain sphingosine lipids in their membranes[49], which may

implicate BT4322 as a sphingosine kinase (given its modest homology to mammalian

SphK1). However, the fact that the diacylglycerol kinase inhibitors, and not the sphingosine

kinase inhibitors, caused growth defects like that observed for arylsulfamates suggests that

BT4322 is a DAGK, and that accumulation of DAG in the lipid membrane is toxic for the

Bacteroides species examined here. Furthermore, inhibition of sphingosine synthesis does

not arrest growth in *B. fragilis,* but does reduce cell survival under stress conditions, and at

stationary phase[49].

The work presented here reveals three significant findings: first is that aryl- and

carbohydrate sulfamates and sulfonates are not effective inhibitors of *exo*-acting S1 HGM

carbohydrate sulfatases. Currently, no inhibitors of these enzymes have been developed

and is an area where further research is needed. Secondly, it is clear that the arylsulfamates

have unintended off-target effects which negatively impact the HGM and this could have

implications for the evolution of arylsulfamates as effective pharmacological agents for the
treatment of hormone dependent cancers/conditions. Thirdly, select complex carbohydrates
can protect the HGM *Bacteroides* species from the harmful effects of arylsulfamates opening
up future paired drug-glycan co-treatment strategies.

Acknowledgements
CJC was funded by MSCA grant MARINEGLYCAN (101029842). PHS acknowledges
funding from the Max Planck Society. SVDP was supported by the Swedish Research
Council (2020-02536) This project has received funding from the Academy of Medical
Sciences/Wellcome Trust through the Springboard Grant (SBF005\1065 163470), the Royal
Society through their research grant (RGS\R2\212050), and Wellcome through the career
development award (225897/Z/22/Z) awarded to AC.

465
Author contributions
AC wrote the draft manuscript. AC, DPB, carried out enzymology experiments. AC, and DPB
analysed enzymological data. CJC, EAY, and SJW synthesised compounds. SVDP carried
out and analysed proteomic experiments and data. AC carried out bacterial growth
experiments. AC supervised and provided funding for the project. All authors read and
approved the manuscript.

472
473 **Data availability statement**
474 Source Data for all experiments, along with corresponding statistical test values, where
475 appropriate, are provided within the paper and in Supplementary information. The mass
476 spectrometry data has been deposited to the ProteomeXchange consortium via PRIDE with
477 the dataset identified PXDXXXXXX. All data is available on request from the authors.

478
479 **Code availability statement**
480 No new codes were developed or compiled in this study

481
482 **Competing interest statement**
483 The authors declare no competing interests.

484
485 **Methods**

486
Recombinant protein production.
Catalytically active recombinant S1 sulfatases (in which Ser, within the C/S-X/P/A-X-R
consensus sequence, for *B. thetaiotaomicron* sulfatases was mutated to Cys, to permit
conversion of the thiol to the formylglycine (FGly) nucleophile by the *E.coli* machinery[3]), or
catalytically-inactive sulfatases (in which Cys in the consensus sequence is mutated to Ser,
thus preventing formation of formylglycine) were expressed in the *E. coli* strain TUNER
(Novagen), and cultured to mid-exponential phase in LB supplemented with 50 μg/mL
kanamycin at 37 °C, in an orbital shaker set to 180 rpm. Cells were then cooled to 16 °C,
and recombinant gene expression was induced by the addition of 0.1 mM isopropyl β-D-1-
thiogalactopyranoside for 16 h at 16 °C, and 180 rpm. The cells were then centrifuged at
5,000 × g and pellets resuspended in 20 mM HEPES, pH 7.4, with 500 mM NaCl before
being sonicated on ice. Recombinant protein was then purified by immobilized metal ion

affinity chromatography using a cobalt-based matrix (Talon, Clontech) and, after a wash with
resuspension buffer, eluted with a step gradient of 10, 50, and 100 mM imidazole in
resuspension buffer. Proteins were then analysed by SDS-PAGE and appropriately pure
fractions were pooled and dialysed into 10 mM HEPES pH 7.0 with 150 mM NaCl. Protein
concentrations were determined by measuring absorbance at 280 nm using the molar
extinction   coefficient   calculated   by   ProtParam   on   the   ExPasy   server
(web.expasy.org/protparam/).

Spectrophotometric based sulfatase assays

508   Production of *para*-nitrophenol from *para*-nitrophenol sulfate by steroid sulfatases was
509   monitored at A_{400nm} using a HIDEX sense or spectra max plus 384 (molecular devices) plate
510   reader in a 96 well plate format. Reactions were monitored at ambient tempertature (20-25
511   °C) and were 100 µl in volume in the presence of 1 mM substrate with 100 mM Bis-Tris-
512   Propane, pH 7.0 with 5 % (v/v) DMSO, 150 mM NaCl, and 5 mM CaCl$_2$.

Microfluidic-based de-sulfation assay

Reducing end BODIPY (maximal excitation/emission coefficient of ~503/511 nm) labelled
sulfated substrates were detected using the EZ Reader II platform (Ret biochem) *via* LED-
induced fluorescence, as described previously[29]. Non-radioactive microfluidic mobility shift
carbohydrate sulfation assays were optimised in solution with a 12-sipper chip coated with
CR8 reagent and performed using a PerkinElmer EZ Reader II system employing EDTA-
based separation buffer and real-time kinetic evaluation of substrate desulfation. Pressure
and voltage settings were adjusted manually (1.8 psi, upstream voltage: 2250 V,
downstream voltage: 500 V) to afford optimal separation in the reaction mixture of the
sulfated substrate and unsulfated glycan product, with a sample (sip) time of 0.2 s, and total
assay times appropriate for the experiment. Individual de-sulfation assays were carried out
at 28°C and were pre-assembled in a 384-well plate in a volume of 80 µl in the presence of
substrate concentrations of 1 µM with 100 mM Bis-Tris-Propane, MES, or Tris, dependent
on the pH optimum of the sulfatase being assayed, and 5 % DMSO, 150 mM NaCl, 0.02%
(v/v) Brij-35 and 5 mM CaCl$_2$. The amount of de-sulfation was directly calculated in real-time
using EZ Reader II software by measuring the sulfated carbohydrate☐:☐unsulfated
carbohydrate ratio at each time-point during the assay. The activity of sulfatase enzymes
was quantified in 'kinetic mode' by monitoring the amount of unsulfated glycan generated
over the assay time, relative to control assay with no enzyme; with sulfate elimination from
the   substrate   limited   to   ~20%   to   prevent   loss   assay   linearity   via   substrate

depletion. k_{cat}/K_M values, using the equation $V_0=(V_{max}/K_M)/S$, were determined by linear
regression analysis with GraphPad Prism software.

Differential scanning fluorimetry

Thermal shift/stability assays (TSAs) were performed using a StepOnePlus Real-Time PCR
machine (LifeTechnologies) and SYPRO-Orange dye, at a 1:1000 dilution, (excitation 470
nm, emission maximum 570 nm, Invitrogen) with thermal ramping between 20 and 95°C in
0.3°C step intervals per data point to induce denaturation of purified, folded, inactive, wild-
type (Ser77) BT1636[S1–20] in the presence or absence of substrate or putative inhibitors. The
melting temperature (Tm) corresponding to the midpoint for the protein unfolding transition
was calculated by fitting the sigmoidal melt curve using the Boltzmann equation in GraphPad
Prism, with R^2 values of ≥0.99, as described in[29]. Data points after the fluorescence intensity
maximum were excluded from the fitting. Changes in the unfolding transition temperature
compared with the control curve (ΔT_m) were calculated for each ligand. A positive ΔT_m value
indicates that the ligand stabilises the protein from thermal denaturation, and confirms
binding to the protein. All TSA experiments were conducted using a final protein
concentration of 5 µM in 100 mM Bis-Tris-Propane (BTP), pH 7.0, and 150 mM NaCl,
supplemented with the appropriate ligand concentration. Three independent assays were
performed for each protein and protein ligand combination.

Bacterial growth experiments

Growth of various *Bacteroides* species in brain heart infusion media (BHI [sigma 53286-
500G]) was carried out by dissolving BHI in 18.2Ω deionised water (37g per litre), and then
sterilised by autoclaving. The media was then allowed to cool prior to adding haematin (1.2
mg/ml in 0.2 His-HCl pH8.0) in a 1:1000 dilution. 5 mls of BHI media was then inoculated
with *Bacteroides* species and grown overnight, and diluted 1:20 into 96 well plate growths
(10 µl in 200) to monitor growth. Growth of the *B. thetaiotaomicron* VPI-5482 on specified
glycan sources was achieved by mixing 1:1 the 0.22 micron filter-sterilised polysaccharides
(10.2 mg/ml in H_2O) with 0.22 micron filter-sterilised 2 x minimal media (per 50 ml was: 0.1g
of ammonium sulfate and sodium carbonate, 0.05g cysteine, 10 ml 1 M potassium
phosphate pH 7.2, 0.1 ml 1 mg.ml^{-1} vitamin K, 1 ml of 0.4 mg.ml^{-1} iron sulfate, 0.4 ml of 0.25
mg.ml resazurin , 0.05 ml of 0.01 mg.ml^{-1} vitamin$_{B12}$, 5 ml of mineral salts for defined media,
and 100 µl of 1.2 mg/ml haematin in 0.2 M His-HCl pH8.0); this gives a final glycan
concentration of 5 mg/ml in a 1 x minimal media. All growth experiments were monitored
continuously in 96-well plates using a cerillo stratus plate reader within a Don whitely VA-
500, A85, or A35 cabinet configured for anaerobic conditions (80 % N_2, 10% CO_2, 10 % H_2).

Where appropriate, either 1 % DMSO (control growths) or 1 % DMSO and 1 mM
arylsulfamates was added to growth conditions. For sphingosine kinase inhibitor (SKI,
567731) and sphingosine kinase 1 inhibitor II (SK1-II, 567741) a maximum of 0.1 and 0.5
mM, respectively, could be dissolved in BHI with 1 % DMSO. For diacylglycerol kinase
inhibitors i (D5794-5MG) and ii (D5919-5MG) a maximum concentration of 125 μM and 62.5
μM, respectively, could be dissolved in BHI with 1 % DMSO but had were hazy; this did not
affect measurement readings at OD_{600nm}. Overnight growth cultures were diluted 1:20 into 96
well plate growths (10 μl in 200 μl). Growth curves presented are averages of three technical
replicates.
579
Thin layer chromatography (TLC)
B. thetaiotaomicron VPI-5482 was grown in 10 ml minimal media with 10 mg/ml (w/v)
appropriate GAG (either CSA or Heparin) as the sole carbon source to mid-exponential
phase in glass test tubes. Cells were harvested by centrifugation at 5,000 × g for 10 min at
room temperature and washed 2 x with 5 mL PBS (pH 7.2) before being resuspended in 1
mL PBS. Cells washed in PBS were lysed and the lysates assayed against 5 mM of the
appropriate substrate at 37 °C for up to 24 h. Assays were analysed by TLC, and 2 μL each
sample was spotted onto silica plates and resolved in butanol:acetic acid:water (2:1:1)
running buffer. The plates were dried, and the sugars were visualized using diphenylamine
stain (DPA: 1 ml of 37.5% HCl, 2 ml of aniline, 10 ml of 85% H_3PO_3, 100 ml of ethyl acetate,
and 2 g of diphenylamine) by heating using a heat gun set to 450°C on medium flow.
Arylsulfates and their pNP products were visualised using 1 M NaOH and then heated as
described for DPA stains.
593
High performance anion exchange chromatography (HPAEC)
Sonicated bacterial cell lysates grown on the glycosaminoglycans CSA and heparin were
treated with 1 mM of arylsulfamates 3 and 12 for 1 hour then challenged with CSA or
heparin mono- and disaccharides by mixing 1:1, to a final concentration of 5 mM, to see if
sulfatase activity was still present. Analysis of the sugar products was performed by HPAEC
using an ICS-6000 with pulsed amperometic detection using a gold work electrode and pDH
reference electrode with a standard quad carbohydrate waveform. Separation was done
using a Dionex PA-200 analytical column preceded by a PA-200 guard column. Samples
were resolved in an isocratically in 100 mM NaOH for 20 mins, then clean with 500 mM
NaOH for 10 mins, finally ran back into 100 mM NaOH for 5 mins, with a flow rate of 0.5
ml/min.
605
606   **Synthesis of chemical compunds**

607 Full detail of the synthetic procedures can be found in the supporting information.
608

Thermal proteome profiling

609 A 300 ml culture of minimal media (see 'Bacterial growth experiments' section)
610

supplemented with 5 mg/ml chondroitin sulfate A (CSA), stored in the anaerobic overnight,
and then inoculated with 10 mls of *B. thetaiotaomicron* VPI-5482 was grown overnight in BHI
media. This culture was then grown to a target OD_{600nm} of ~0.5-0.6 at which point cells were
centrifuged at 5000 x g and washed with PBS, this process was repeated two more times.
Cells were then treated as in *Mateus et al*[39]; briefly, cells were then resuspended in lysis
buffer (50µg/ml lysozyme, 1×protease inhibitor(Roche), 250 U/ml benzonase, and 1 mM
$MgCl_2$ in PBS) to give an OD_{600nm} of 50. Samples were then sonicated on ice for 15 seconds
three times. The lysate was then split into three: One sample was a control (1 % (v/v)
DMSO) and experimental samples contained 1 mM arylsulfamate 3 or 1 mM arylsulfamate
12 (1 % (v/v) DMSO). Next 20 µl of each condition were aliquoted into a 96 well PCR plate
across 10 wells in quadruplicate. The plates were then subjected to a temperature gradient
from 45 – 72 °C (45, 46.8, 49.1, 52.1, 55.9, 60.3, 64.4, 67.4, 69.9, 72°C) for 3 min, followed
by 3 min at room temperature. NP-40 was then added to samples to a final concentration of
0.8 % (v/v). Samples were then transferred to a 0.22 µm 96 well filter plate (Millipore:
MSGVS2210) and centrifuged at 500 x g for 5 min to remove protein aggregates. Following
filtration, the flowthrough was combined 1:1 with 400 mM Tris pH 8.0 and 4 % SDS (w/v).
The protein concentration for each sample was determined by Bradford assay, and a volume
corresponding to 10 µg for the lowest-temperature condition was used as input for all other
samples in a set for analysis. Protein digestion was performed using a modified solid-phase-
enhanced sample preparation (SP3) based method adapted to a 96 well format in 0.45µM
filter plates (Millipore, MSRPN04)[50,51]. Premixed magnetic SpeedBeads (Cytiva, (1:1,
GE45152105050250 and GE 65152105050250)) were added to the samples, combined with
two volumes of ethanol and incubated for 10 min while mixing at 150 RPM. Samples were
transferred to the filter plates and washed four times with 200 µL of 70% ethanol (v/v) by
centrifugation for 2 min. at 2000 x g. Beads were resuspended in 40 µL 0.1 mM HEPES pH
8.5 containing trypsin/Lys-c (Thermo Scientific), 1.25 mM TCEP and 5 mM chloroacetamide
(Sigma-Aldrich) and incubated at room temperature overnight while mixing 500
RPM. Peptides were collected by centrifugation, followed by a second elution with 2%
DMSO. TMT 10-plex 17 labelling (Thermo Scientific) was performed at a 2:1 ratio for 1 hr
while mixing at 500 RPM. The 17 labelling reaction was quenched with 0.4% hydroxylamine
(v/v) for 15 min before each sample was combined and dried under vacuum. Peptides were
resolved in 0.1% trifluoroacetic acid (TFA) and fractionated using 0.1% triethylamine with
increasing concentrations of acetonitrile by high pH reverse phase into 8 fractions using C18

desalting columns (Pierce, 89852) pooled into 4 fractions and dried under a vacuum.
Lyophilized samples are resolved in 0.1% TFA and analyzed by LC-MS/MS using a nano
HPLC system (EASY-nLC 1200, Thermo Scientific) coupled to a Q-Exactive HF-X mass
spectrometer (Thermo Scientific). Peptides were separated using in-house packed columns
(150 x 0.075 mm) packed with Reprosil-Pur C18-AQ 3 µm particles (Dr. Maisch). Elution
was performed with a 5 to 45% gradient (A: 0.1% formic acid, B :0.1% formic acid ,80%
Acetontrile) in 135 minutes at 250 nl. Full mass spectra were acquired over a mass range of
minimum 400 m/z and maximum 1600 m/z, with a resolution of 60,000 at 200 m/z. The
isolation window for fragmentation spectra collection was set to 1Da and a fixed first mass of
100 m/z with a resolution of 30,000 at 200 m/z. The top 15 most intense peaks with a charge
state ≥2 – 5 were selected for fragmentation by HCD with a normalized collision energy of
32% and subsequent excluded for selection for 30 seconds. Thermo .RAW files were
converted into MZml using msConvert (proteowizard) and FragPipe (v20.0) configured with
Msfragger (v3.8) and Philosopher (v5.0)[52,53] was used for protein identification and TMT
quantification. Database searches were performed against the *Bacteroides thetaiotaomicron*
reference proteome (4782 entries, UP000001414, 2023_10) with the following settings,
parent ion mass and fragment ion mass accuracy after recalibration was set to 20 ppm.
Enzyme specificity was set to trypsin, fixed modifications were set for cysteine
carbamidomethylation and TMT modified lysine, variable modifications considered were
methionine oxidation, protein n-terminal acetylation and TMT modification on peptide n-
terminal or serine. The identified proteins and peptides were both filtered at a 1% and only
identifications based on ≥2 unique peptides with a minimum ion purity of 0.75 were
considered for quantification and thermal proteome profiling. Further data analysis was
performed in R environment to combine the search outcome, normalize reporter ion
channels. Meltome curve fitting analysis and statistical analysis were executed using the
TPP-TR workflow in the TPP package[54].
670
Ligand docking experiments
All docking experiments were conducted using SwissDock[55] on the Expasy server using
default parameter; no binding site was specified. The alphafold model of BT4322 (Q89ZQ4)
and the crystal structure of SphK1 (3VZB) were downloaded, and where appropriate
heteroatoms removed, and uploaded to the Swiss dock as PDB files. Small molecules to be
docked where built in Jligand[56], the generated PDB or CIF files converted to MOL2 files
using the openbabel server and uploaded to SwissDock as MOL2 files.
678
679
**Figure 1 General example of the conserved S1 sulfatase fold and consensus active
site sequence.**

a. The 3-D structure of the human lysosomal sulfatase GALNS (PDB:4FDJ), the N-terminal
domain is coloured red and yellow for alpha helices and beta sheets, with the C-terminal
subdomain coloured cyan and magenta for alpha helices and beta sheets; a zoom in of the
consensus sequence of GALNS with a converted formylglycine, and with the general
consensus pictured to the right; **b.** Structure of phase I and II arylsulfamate drugs developed
to inhibit STS **c.** Chemical structures of aryl- and carbohydrate sulfamate inhibitors used in
the study. **d.** Chemical structures of phenylvinyl sulfone, phenylsulfonyl fluoride, phenylvinyl
sulfonate, and carbohydrate vinyl sulfone.

Figure 2 The effectiveness of arylsulfamate inhibitors against S1 steroid and
carbohydrate sulfatases.
a. The enzymes PaAsta and HpSulf, belonging to S1_4 (green bar) and S1_2 (yellow bar),
respectively, assayed against the panel of arylsulfamates (1-19 listed in Figure 2).; **b.** The
S1_11 sulfatases (BT3177$^{6S-GlcNAc}$, BT4656$^{6S-GlcNAc}$, Amuc_1033$^{6S-GlcNAc}$, Amuc_1074$^{6S-GlcNAc}$;
blue bar above), the S1_20 sulfatases (BT1622$^{3S-Gal/GalNAc}$, BT1636^{3S-Gal}, Amuc_0451^{3S-Gal},
Amuc_0491^{3S-Gal}; pink bar above), and the S1_16 enzymes (BT3796$^{4S-Gal/GalNAc}$ and
BT1755^{4S-Gal}; orange bar above) against the panel of arylsulfamates (1-19 listed in **Figure**
2). For both a and b the left two panels display relative activity to wildtype protein with the
arylsulfamate compound in the assay, whilst the left two panels are overnight preincubation
of the enzyme with selected arylsulfamates then dilution of this preincubation mix into the
reaction. The numbers indicate which arylsulfamates from **Figure 1** were used and in all
instances 1 mM of compound was deployed. Wt indicates reaction with enzyme and no
arylsulfamate. Kinetic curves from which the data are derived are shown in supplemental
figures S1-S24.

Figure 3 Growth of *Bacteroides thetaiotaomicron* on varied carbon sources in the
presence of arylsulfamate inhibitors
Bacteroides thetaiotaomicron (*B.theta*) grown on varied carbon sources; two rich carbon
sources, brain heart infusion media (BHI) and glucose (Glc); two sulfated host glycan carbon
sources, chondroitin sulfate c (CSC) and Heparin (Hep), and two plant glycan carbon
sources potato galactan (PG) and Larch arabinogalactan (LAG), in the presence of
arylsulfamates inhibitors, labelled 1-19 (see **Figure 2** for structures). Complex glycans and
arylsulfamates inhibitors were used at a concentration of 5 mg/ml and 1 mM, respectively.
All carbon source were utilised in a prepared minimal media except BHI which was
purchased from sigma and simply dissolved in water. The traces are the mean of 3
independent growths but errors bars have been omitted for clarity; individual graphs with
error bars are presented in supplemental figures S25-S30.

Figure 4. The effectiveness of carbohydrate sulfamate substrate mimics against the
S1 carbohydrate sulfatase BT1636^{3S-Gal}.
a. Structure of the carbohydrate sulfonate/sulfamates and their aryl equivalents**. b.** The
graphs show the effectiveness of carbohydrate sulfamates, carbohydrate sulfovinyl, and
arylsulfonates against BT1636^{3S-Gal}; the left graph shows relative activity to wildtype with the
compounds included in the assay, whilst the right graph show the relative activity after the
enzyme had been preincubated with the compounds overnight then diluted into the reaction
mixture; **c.** Thermostability shift assays (TSA) using inactive BT1636^{3S-Gal} against the
substrate 3S-Gal, and the putative inhibitor structures carbohydrate sulfamates,
carbohydrate sulfovinyl, and arylsulfonates. All compounds in **b** were used at a
concentration of 1 mM.

Figure 5. Arylsulfamates broadly affect *Bacteroides* spp. of the HGM
Bacteroides species were grown in BHI media in the presence, or absence, of 1 mM of the
listed arylsulfamate. The data plotted are the maximal OD$_{600nm}$ observed for arylsulfamate
treated growths relative to the untreated controls which were both ran in triplicate and

randomly paired then averages derived. For all the estradiol growths a single experiment
was conducted except for *B. theta* which was ran in triplicate; this was due to limited
material.
740
Figure 6. Global protein thermal stability reveals off target interactions of
arysulfamate inhibitors 3 and 12.
a. Table of proteins which displayed an altered Tm in the presence of arylsulfamates 3 and
12. **b.** Proteins identified with a significance change in melting point upon inhibitor 3
treatment (n=2). **c.** Proteins identified with a significance change in melting point upon
inhibitor 12 treatment (n=2). A dashed box indicates BT4322 is the only protein common
across both groups. **d.** *Bacteroides* species were grown in BHI media in the presence, or
absence, of 0.125 mM or 0.0625 mM diacylglycerol kinase inhibitor I and II, respectively.
The data plotted are the maximal OD_{600nm} observed for inhibitor treated growths relative to
the untreated controls which were both ran in triplicate and randomly paired then averages
derived.
752
Extended data figure 1. Cell lysates of *Bacteroides thetaiotaomicron* incubated with
two arylsulfamates inhibitors
a. The effects of varied concentrations of arylsulfamate 3 and 12 on the growth of *B. theta* in
BHI media. **b.** Thin layer chromatography (TLC) analysis of cell lysates from *B. theta* cells
grown to mid-exponential phase on CSA as a carbon source, and incubated with
arylsulfamates 4 and 13 **(Figure 1)**, mixed 1:1 with the sulfated substrates: *O*2 sulfated
uronic acid β1,3 linked to *O*4 sulfated *N*-acetylgalactosmaine (UA2S-GalNAc4S), uronic acid
β1,3 linked to *O*4 sulfated *N*-acetylgalactosmaine (UA-GalNAc4S), and *O*6 sulfated *N*-
acetylgalactosmaine (GalNAc6S). Below are high performance anion exchange
chromatography (HPAEC) traces showing the production of GalNAc in all instances except
the negative control; **c.** TLC analysis of cell lysates from *B. theta* cells grown to mid-
exponential phase on Heparin as a carbon source, and incubated with arylsulfamates 4 and
13 **(Figure 1)**, mixed 1:1 with the sulfated substrates: *O*2 sulfated uronic acid β1,4 linked to
*O*4 sulfated *N*-acetylgalactosmaine (UA2S-GalNAc4S), *O*6 sulfated *N*-acetylglucosmaine
(GlcNAc6S), and *N*-sulfated glucosamine (GlcNS). Below are high performance anion
exchange chromatography (HPAEC) traces showing the production of GlcNAc in all
instances except the negative control. All reactions were in PBS, with 5 % DMSO, and
incubated at 37°C overnight.
771
Extended data figure 2. Activity of carbohydrate sulfatases on *para*-nitrophenol
sulfate.
Sulfatases were incubated with 5 mM *para*-nitrophenol sulfate overnight in 100 mM BTP pH
7.0 with 150 mM NaCl and 5 mM $CaCl_2$. Reactions were then ran on silica based thin layer
chromatography using a resolving buffer of butanol:acetic acid:water (2:1:1) and develop by
soaking the plates with 1 M NaOH and heating with a heat gun. It can be seen only the
steroid sulfatase *Pa*Asta shows the production of a *para*-nitrolphenol product.
779
Extended data figure 3. Global features of the thermal proteome profiling data
a. Distribution of the melting temperature (Tm) of *Bacteroides thetaiotaomicron* VPI-5482 (*B.*
theta) lysate proteome and the whole cell *E.coli* proteome. **b.** Ranked abundance plot of all
identified proteins in the 1% (v/v) DMSO treated controls, S1 sulfatase are highlighted. **c.**
Relative abundance of 11 S1 sulfatases identified by mass spectrometry for *B. theta* cultured
in the presence of chondroitin sulfate A (n=6). **d.** Melting curve profiles for all identified *B.*
theta proteins in control (1% (v/v) DMSO) treated lysate. The thermal stability profiles for
each individual S1 sulfatases are highlighted in blue.
788
789 **Extended data figure 4. Global protein thermal stability analysis confirms that**
790 **arysulfamate inhibitors 3 and 12 do not directly interact with S1 sulfatases.**

Individual melt curve for selected S1 sulfatase for control and arysulfamate inhibitors 3 and 12 treated lysates (n=2 per treatment group).

Extended data figure 5. Structure of BT4322, SaDgKB, and Sphk1 and comparison of the ATP helix
a. Tertiary structures of BT4322, SaDgKB, and SphK1 shown as with α helices, β sheets, and loops coloured cyan, magenta, and pink. The 'ATP helix' is coloured in green. **b.** An expanded view of the ATP helix and its sequence.

Extended data figure 6. Sequence alignment of putative DAGKs from select Bacteroides species.
Sequences were aligned using the MAFFT online server and visualised using Jalview.

Extended data figure 7. The effects of sphigosine kinase inhibitors on the growth of HGM Bacteroidota species
Bacteroidota species were grown in BHI, 1 % DMSO, with 0.1 mM of sphingosine kinase inhibitor (SKI) and 0.5 mM sphingosine kinase 1 inhibitor II and the effects on growth observed. Data are technical triplicates with the standard error of the mean.

Extended data figure 8. The effects of diacylglycerol kinase inhibitors on the growth of HGM Bacteroidota species
Bacteroidota species were grown in BHI, 1 % DMSO, with 0.125 mM of diacylglycerol kinase inhibitor I (DAGKI-i) and 0.0625 mM diacylglycerol kinase inhibitor II (DAGKI-ii) and the effects on growth observed. Data are technical triplicates with the standard error of the mean.

Extended data figure 9. Blind docking of arylsulfamates onto BT4322 and SphK1.
The results of blind docking (no site specified) arylsulfamate inhibitors into the alphafold model of BT4322 (Q89ZQ4) and the crystal structure of Sphk1 (3VZB). It can be observed all arylsulfamates preferentially cluster in, or near, the lipid binding pocket. For SpKH1 there appears to be no single preferred clustering site.

Supplementary figure 1. Kinetic curves of *Pa*Asta assayed against *para*-nitrophenol sulfate with, and without, arylsulfamate inhibitors
*Pa*Asta, at a concentration of 25 nM, was assayed against 1 mM *para*-nitrophenol with and without 1 mM of various arylsulfamate inhibitors included in the assay. The assay was performed in 100 mM of Bis-Tris-propane pH 7.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.

Supplementary figure 2. Kinetic curves of *Pa*Asta assayed against *para*-nitrophenol sulfate after pre-incubation of the enzyme with arylsulfamate inhibitors
*Pa*Asta, which had been incubated for ~24h with 1 mM of the appropriate arylsulfamate inhibitor, was assayed using a concentration of 25 nM against 1 mM para-nitrophenol. The assay was performed in 100 mM of Bis-Tris-propane pH 7.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.

Supplementary figure 3. Kinetic curves of *Hp*Sulf assayed against para-nitrophenol sulfate with, and without, arylsulfamate inhibitors
*Hp*Sulf, at a concentration of 5 μg/ml, was assayed against 1 mM para-nitrophenol with and without 1 mM of various arylsulfamate inhibitors included in the assay. The assay was performed in 100 mM of Bis-Tris-propane pH 7.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.

846
Supplementary figure 4. Kinetic curves of *Hp*Sulf assayed against para-nitrophenol
sulfate after pre-incubation of the enzyme with arylsulfamate inhibitors
*Hp*Sulf, which had been incubated for ~24h with 1 mM of the appropriate arylsulfamate
inhibitor, was assayed using a concentration of 50 μg/ml against 1 mM para-nitrophenol. The
assay was performed in 100 mM of Bis-Tris-propane pH 7.0 with 5 % DMSO, 150 mM NaCl,
0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate
arylsulfamate compound as in Figure 1.
854
Supplementary figure 5. Kinetic curves of BT3177[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine with, and without, arylsulfamate inhibitors
BT3177[6S-GlcNAc], at a concentration of 446.5 nM, was assayed against 1 μM BODIPY labelled
6S-*N*-acetylglucosamine with and without 1 mM of various arysulfamate inhibitors included
in the assay. The assay was performed in 100 mM of Bis-Tris-propane pH 8.0 with 5 %
DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
862
Supplementary figure 6. Kinetic curves of BT3177[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine after pre-incubation of the enzyme with
arylsulfamate inhibitors
BT3177[6S-GlcNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 470 nM against 1 μM BODIPY
labelled 6S-*N*-acetylglucosamine. The assay was performed in 100 mM of Bis-Tris-propane
pH 8.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were
performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
871
Supplementary figure 7. Kinetic curves of BT4656[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine with, and without, arylsulfamate inhibitors
BT4656[6S-GlcNAc], at a concentration of 200 nM, was assayed against 1 μM BODIPY labelled
6S-*N*-acetylglucosamine with and without 1 mM of various arysulfamate inhibitors included
in the assay. The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM
NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers
indicate arylsulfamate compound as in Figure 1.
879
Supplementary figure 8. Kinetic curves of BT4656[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine after pre-incubation of the enzyme with
arylsulfamate inhibitors
BT4656[6S-GlcNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 200 nM against 1 μM BODIPY
labelled 6S-*N*-acetylglucosamine. The assay was performed in MES pH 6.0 with 5 % DMSO,
150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate.
Numbers indicate arylsulfamate compound as in Figure 1.
888
Supplementary figure 9. Kinetic curves of Amuc1074[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine after pre-incubation of the enzyme with
arylsulfamate inhibitors
Amuc1074[6S-GlcNAc], at a concentration of 6.4 μM, was assayed against 1 μM BODIPY
labelled 6S-*N*-acetylglucosamine with and without 1 mM of various arysulfamate inhibitors
included in the assay. The assay was performed in 100 mM of MES pH 6.0 with 5 %
DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
897
Supplementary figure 10. Kinetic curves of Amuc1074[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine with, and without, arylsulfamate inhibitors

Amuc1074[6S-GlcNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 6.08 µM against 1 µM BODIPY
labelled 6S-*N*-acetylglucosamine. The assay was performed in 100 mM of MES pH 6.0 with
5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
905
Supplementary figure 11. Kinetic curves of Amuc1033[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine with, and without, arylsulfamate inhibitors
Amuc1033[6S-GlcNAc], at a concentration of 4.6 µM, was assayed against 1 µM BODIPY
labelled 6S-*N*-acetylglucosamine with and without 1 mM of various arysulfamate inhibitors
included in the assay. The assay was performed in 100 mM of MES pH 6.0 with 5 %
DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
913
Supplementary figure 12. Kinetic curves of Amuc1033[6S-GlcNAc] assayed against BODIPY
labelled 6S-*N*-acetylglucosamine after pre-incubation of the enzyme with
arylsulfamate inhibitors
Amuc1033[6S-GlcNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 4.37 µM against 1 µM BODIPY
labelled 6S-*N* acetylglucosamine. The assay was performed in 100 mM of MES pH 6.0 with
5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
922
Supplementary figure 13. Kinetic curves of BT1636[3S-Gal] assayed against BODIPY
labelled 3S-galactose with, and without, arylsulfamate inhibitors
BT1636[3S-Gal], at a concentration of 400 nM, was assayed against 1 µM BODIPY labelled 3S-
galactose with and without 1 mM of various arysulfamate inhibitors included in the assay.
The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM NaCl, 0.02%
(v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate
arylsulfamate compound as in Figure 1.
930
Supplementary figure 14. Kinetic curves of BT1636[3S-Gal] assayed against BODIPY
labelled 3S-galactose sulfate after pre-incubation of the enzyme with arylsulfamate
inhibitors
BT1636[3S-Gal], which had been incubated for ~24h with 1 mM of the appropriate arylsulfamate
inhibitor, was assayed using a concentration of 200 nM against 1 µM BODIPY labelled 3S-
galactose. The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM
NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers
indicate arylsulfamate compound as in Figure 1.
939
Supplementary figure 15. Kinetic curves of BT1622[3S-Gal/GalNAc] assayed BODIPY labelled
3S-*N*-acetylgalactosamine with, and without, arylsulfamate inhibitors
BT1622[3S-Gal/GalNAc], at a concentration of 6.5 µM, was assayed against 1 µM BODIPY labelled
3S-*N*-acetylgalactosamine with and without 1 mM of various arysulfamate inhibitors included
in the assay. The assay was performed in 100 mM of BTP pH 8.5 with 5 % DMSO, 150 mM
NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers
indicate arylsulfamate compound as in Figure 1.
947
Supplementary figure 16. Kinetic curves of BT1622[3S-Gal/GalNAc] assayed against BODIPY
labelled 3S-*N*-acetylgalactosamine after pre-incubation of the enzyme with
arylsulfamate inhibitors
BT1622[3S-Gal/GalNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 2 µM against 1 µM BODIPY
labelled 3S-*N*-acetylgalactosamine. The assay was performed in 100 mM of BTP pH 8.5

with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were
performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
956
Supplementary figure 17. Kinetic curves of Amuc0451[3S-Gal] assayed against BODIPY
labelled 3S-galactose with, and without, arylsulfamate inhibitors
Amuc0451[3S-Gal], at a concentration of 15.6 μM, was assayed against 1 μM BODIPY labelled
3S-galactose with and without 1 mM of various arysulfamate inhibitors included in the assay.
The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM NaCl, 0.02%
(v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate
arylsulfamate compound as in Figure 1.
964
Supplementary figure 18. Kinetic curves of Amuc0451[3S-Gal] assayed against BODIPY
labelled 3S-galactose after pre-incubation of the enzyme with arylsulfamate inhibitors
Amuc0451[3S-Gal], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 15.6 μM against 1 μM BODIPY
labelled 3S-galactose. The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO,
150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate.
Numbers indicate arylsulfamate compound as in Figure 1.
972
Supplementary figure 19. Kinetic curves of Amuc0491[3S-Gal/GalNAc] assayed against
BODIPY labelled 3S-*N*-acetylgalactosamine with, and without, arylsulfamate inhibitors
Amuc0491[3S-Gal/GalNAc], at a concentration of 3.4 μM, was assayed against 1 μM BODIPY
labelled 3S-*N*-acetylgalactosamine with and without 1 mM of various arysulfamate inhibitors
included in the assay. The assay was performed in 100 mM of MES pH 6.0   with 5 %
DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
980
Supplementary figure 20. Kinetic curves of Amuc0491[3S-Gal/GalNAc] assayed against
BODIPY labelled 3S-*N*-acetylgalactosamine after pre-incubation of the enzyme with
arylsulfamate inhibitors
Amuc0491[3S-Gal/GalNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 3.2 μM against 1 μM BODIPY
labelled 3S-*N*-acetylgalactosamine. The assay was performed in 100 mM of MES pH 6.0
with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were
performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
989
Supplementary figure 21. Kinetic curves of BT3796[4S-Gal/GalNAc] assayed against BODIPY
labelled 4S-galactose with, and without, arylsulfamate inhibitors
BT3796[4S-Gal/GalNAc], at a concentration of 630 nM, was assayed against 1 μM BODIPY
labelled 4S-galactose with and without 1 mM of various arylsulfamate inhibitors included in
the assay. The assay was performed in 100 mM of BTP pH 8.5   with 5 % DMSO, 150 mM
NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers
indicate arylsulfamate compound as in Figure 1.
997
Supplementary figure 22. Kinetic curves of BT3796[4S-Gal/GalNAc] assayed against BODIPY
labelled 4S-galactose after pre-incubation of the enzyme with arylsulfamate inhibitors
BT3796[4S-Gal/GalNAc], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 630 nM against 1 μM BODIPY
labelled 4S-galactose. The assay was performed in 100 mM of BTP pH 8.5 with 5 % DMSO,
150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate.
Numbers indicate arylsulfamate compound as in Figure 1.
1005
Supplementary figure 23. Kinetic curves of Amuc1755[4S-Gal] assayed against BODIPY
labelled 4S-galactose with, and without, arylsulfamate inhibitors

Amuc1755[4S-Gal], at a concentration of 2.3 µM, was assayed against 1 µM BODIPY labelled
4S-galactose with and without 1 mM of various arysulfamate inhibitors included in the assay.
The assay was performed in 100 mM of MES pH 6.0   with 5 % DMSO, 150 mM NaCl,
0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in triplicate. Numbers indicate
arylsulfamate compound as in Figure 1.
1013
Supplementary figure 24. Kinetic curves of Amuc1755[4S-Gal] assayed against BODIPY
labelled 4S-galactose after pre-incubation of the enzyme with arylsulfamate inhibitors
Amuc1755[4S-Gal], which had been incubated for ~24h with 1 mM of the appropriate
arylsulfamate inhibitor, was assayed using a concentration of 2.185 µM against 1 µM
BODIPY labelled 4S-galactose. The assay was performed in 100 mM of MES pH 6.0 with 5
% DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. Assays were performed in
triplicate. Numbers indicate arylsulfamate compound as in Figure 1.
1021
Supplementary figure 25. The effects of DMSO on the growth of *Bacteroides*
***thetaiotaomicron* VPI-5482**
Bacteroides thetaiotaomicron VPI-5482 was grown in rich media (BHI), or minimal media
supplemented 5 mg/ml of an appropriate polysaccharide with either 5 % or 1 % DMSO. Data
are technical triplicates with the standard error of the mean. Numbers indicate arylsulfamate
compound as in Figure 1.
1028
1029   **Supplementary figure 26. The effects of arylsulfamates on the growth of *Bacteroides***
1030   ***thetaiotaomicron* VPI-5482 grown in BHI media**
1031   *Bacteroides thetaiotaomicron* VPI-5482 was grown in rich media (BHI), with 1 % DMSO, and
1032   with or without 1 mM of the appropriate arylsulfamate. Data are technical triplicates with the
1033   standard error of the mean. Numbers indicate arylsulfamate compound as in Figure 1.
1034
Supplementary figure 27. The effects of arylsulfamates on the growth of *Bacteroides*
***thetaiotaomicron* VPI-5482 grown in minimal media with D-glucose**
Bacteroides thetaiotaomicron VPI-5482 was grown in minimal media supplemented with 5
mg/ml D-glucose (Glc), 1 % DMSO, and with or without 1 mM of the appropriate
arylsulfamate. Data are technical triplicates with the standard error of the mean. Numbers
indicate arylsulfamate compound as in Figure 1.
1041
Supplementary figure 28. The effects of arylsulfamates on the growth of *Bacteroides*
***thetaiotaomicron* VPI-5482 grown in minimal media with Chondroitin sulfate C (CSC)**
Bacteroides thetaiotaomicron VPI-5482 was grown in minimal media supplemented with 5
mg/ml Chondroitin sulfate C (CSC), 1 % DMSO, and with or without 1 mM of the appropriate
arylsulfamate. Data are technical triplicates with the standard error of the mean. Numbers
indicate arylsulfamate compound as in Figure 1.
1048
Supplementary figure 29. The effects of arylsulfamates on the growth of *Bacteroides*
***thetaiotaomicron* VPI-5482 grown in minimal media with Heparin**
Bacteroides thetaiotaomicron VPI-5482 was grown in minimal media supplemented with 5
mg/ml Heparin (Hep), 1 % DMSO, and with or without 1 mM of the appropriate
arylsulfamate. Data are technical triplicates with the standard error of the mean. Numbers
indicate arylsulfamate compound as in Figure 1.
1055
Supplementary figure 30. The effects of arylsulfamates on the growth of *Bacteroides*
***thetaiotaomicron* VPI-5482 grown in minimal media with Potato galactan**
Bacteroides thetaiotaomicron VPI-5482 was grown in minimal media supplemented with 5
mg/ml Potato galactan (PG), 1 % DMSO, and with or without 1 mM of the appropriate
arylsulfamate. Data are technical triplicates with the standard error of the mean. Numbers
indicate arylsulfamate compound as in Figure 1.
1062

Supplementary figure 31. The effects of arylsulfamates on the growth of *Bacteroides thetaiotaomicron* VPI-5482 grown in minimal media with Larch arabinogalactan

Bacteroides thetaiotaomicron VPI-5482 was grown in minimal media supplemented with 5 mg/ml Larch arabinogalactan (LAG), 1 % DMSO, and with or without 1 mM of the appropriate arylsulfamate. Data are technical triplicates with the standard error of the mean. Numbers indicate arylsulfamate compound as in Figure 1.

Supplementary figure 32. The effects of varied concentration arylsulfamates 3 and 12 on the growth of *Bacteroides thetaiotaomicron* VPI-5482 grown in BHI media

Bacteroides thetaiotaomicron VPI-5482 was grown in BHI, 1 % DMSO, with varied concentrations of arylsulfamate 3 and 12. Data are technical triplicates with the standard error of the mean. Numbers indicate arylsulfamate compound as in Figure 1.

Supplementary figure 33. Kinetic curves of BT1636$^{3S\text{-}Gal}$ assayed against BODIPY labelled 3S-galactose with, and without, carbohydrate and aryl- sulfonates/sulfamates

BT1636$^{3S\text{-}Gal}$, at a concentration of 270 nM, was assayed against 1 μM BODIPY labelled 3S-galactose with and without 1 mM of various arysulfamate inhibitors included in the assay. The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM CaCl$_2$. Assays were performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.

Supplementary figure 34. Kinetic curves of BT1636$^{3S\text{-}Gal}$ assayed against BODIPY labelled 3S-galactose sulfate after pre-incubation of the enzyme with carbohydrate and aryl- sulfonates/sulfamates

BT1636$^{3S\text{-}Gal}$, which had been incubated for ~24h with 1 mM of the appropriate arylsulfamate inhibitor, was assayed using a concentration of 270 nM against 1 μM BODIPY labelled 3S-galactose. The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM CaCl$_2$. Assays were performed in triplicate. Numbers indicate arylsulfamate compound as in Figure 1.

Supplementary figure 35. Thermal melt curves of BT1636$^{3S\text{-}Gal}$ incubated substrate and various carbohydrate and aryl- sulfonates/sulfamates

BT1636$^{3S\text{-}Gal}$, at a concentration of 5 μM, was incubated with no compound, and varying concentrations and the effect on its melting temperature monitored, a shift in melting temperature is indicative of an interaction. The assays were performed in 100 mM of BTP pH 7.0 with 5 % DMSO, 150 mM NaCl. Assays were performed in triplicate.

Supplementary figure 36. The effects of arylsulfamates 3 and 12 on the growth of HGM Bacteroidota species

Bacteroidota species were grown in BHI, 1 % DMSO, with 1 mM of arylsulfamate 3 and 12 and the effects on growth observed. Data are technical triplicates with the standard error of the mean.

Supplementary figure 37. The effects of the phase I/II arylsulfamate drugs Irosustat and estradiol sulfamate on the growth of Bacteroidota species

Bacteroidota species were grown in BHI, 1 % DMSO, with 1 mM of Irosustat and estradiol sulfamate and the effects on growth observed. Growths with Irosustat are technical triplicates with the standard error of the mean. Growths with estradiol sulfamate represent single growth experiments with the exception of *Bacteroides thetaiotaomicron* VPI-5482 where the data is from triplicate growths with standard error of the mean.

Supplementary table 1. Kinetic rates of sulfatase when arylsulfamates were included in the assay conditions.

The concentration of arylsulfamate was always 1 mM. For the steroid sulfatases the substrate *para*-nitrophenol sulfate was at a concentration of 1 mM, whilst for carbohydrate sulfatases substrate concentration was 1 µM. NA means no activity when compared to the control. NT means not tested. Numbers in the left most column refer to the arylsulfamate in Figure 1c added to the reaction; control represents substrate only whilst WT is substrate plus enzyme without an arylsulfamate inhibitor. Rates are µM product/min.

Supplementary table 2. Kinetic rates of sulfatase when sulfatases were pre-incubated with arylsulfamates then assayed.

The concentration of arylsulfamate used for preincubation was always 1 mM and the enzyme concentration 25 - 50 µM. For the steroid sulfatases the substrate *para*-nitrophenol sulfate was at a concentration of 1 mM, whilst for carbohydrate sulfatases substrate concentration was 1 µM. NA means no activity when compared to the control. NT means not tested. Numbers in the left most column refer to the arylsulfamate in Figure 1c added to the reaction; control represents substrate only whilst WT is substrate plus enzyme without an arylsulfamate inhibitor. Rates are µM product/min.

Supplementary table 3. Kinetic rate of BT1636$^{3S\text{-}Gal}$ assayed against BODIPY labelled 3S-galactose sulfate with various carbohydrate and aryl- sulfonates/sulfamates.

BT1636$^{3S\text{-}Gal}$ was assayed against 1 µM BODIPY labelled 3S-galactose with 1 mM of **with** various carbohydrate and aryl- sulfonates/sulfamates the assay, or after BT1636$^{3S\text{-}Gal}$ had been incubated with 1 mM of the same compounds; concentrations of 300 nM and 285 nM of BT1636$^{3S\text{-}Gal}$ were used, respectively. The assay was performed in 100 mM of MES pH 6.0 with 5 % DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM CaCl$_2$. Assays were performed in triplicate. Rates are fluorescence of product/min. The values in green indicate 5 % was not present.

Supplementary table 4. Thermal melt values of BT1636$^{3S\text{-}Gal}$ in the absence and presence of various carbohydrate and aryl- sulfonates/sulfamates.

BT1636$^{3S\text{-}Gal}$, at a concentration of 5 µM, was incubated with no compound, and varying concentrations and the effect on its melting temperature monitored, a shift in melting temperature is indicative of an interaction. The assays were performed in 100 mM of BTP pH 7.0 with 5 % DMSO, 150 mM NaCl. Assays were performed in triplicate.

Supplementary table 5. Conservation of BT4322 in select Bacteroidota species of the human gut microbiota.

BT4322 was used as the query against the genomes of the 10 species from Figure 5 against which arylsulfamates 3 and 12 were tested. The BLASTp search was done using default settings with the blastp algorithm accessed through the NCBI interface.

References:

1   Barbeyron, T. *et al.* Matching the Diversity of Sulfated Biomolecules: Creation of a Classification Database for Sulfatases Reflecting Their Substrate Specificity. *PloS one* **11**, e0164846 (2016). https://doi.org:10.1371/journal.pone.0164846

2   Stam, M. *et al.* SulfAtlas, the sulfatase database: state of the art and new developments. *Nucleic Acids Res* **51**, D647-D653 (2023). https://doi.org:10.1093/nar/gkac977

3   Hanson, S. R., Best, M. D. & Wong, C. H. Sulfatases: structure, mechanism, biological activity, inhibition, and synthetic utility. *Angewandte Chemie* **43**, 5736-5763 (2004). https://doi.org:10.1002/anie.200300632

1167   4   Luis, A. S., Yates, E. A. & Cartmell, A. Functions and specificity of bacterial
1168   carbohydrate sulfatases targeting host glycans. *Essays Biochem* **67**, 429-442 (2023).
1169   https://doi.org:10.1042/EBC20220120

1170   5   Stanway, S. J. *et al.* Steroid sulfatase: a new target for the endocrine therapy of
1171   breast cancer. *Oncologist* **12**, 370-374 (2007).
1172   https://doi.org:10.1634/theoncologist.12-4-370

1173   6   Reed, M. J., Purohit, A., Woo, L. W., Newman, S. P. & Potter, B. V. Steroid sulfatase:
1174   molecular biology, regulation, and inhibition. *Endocr Rev* **26**, 171-202 (2005).
1175   https://doi.org:10.1210/er.2004-0003

1176   7   Armstrong, C. M. *et al.* Steroid Sulfatase Stimulates Intracrine Androgen Synthesis
1177   and is a Therapeutic Target for Advanced Prostate Cancer. *Clin Cancer Res* **26**, 6064-
1178   6074 (2020). https://doi.org:10.1158/1078-0432.CCR-20-1682

1179   8   Ai, X. *et al.* Substrate specificity and domain functions of extracellular heparan
1180   sulfate 6-O-endosulfatases, QSulf1 and QSulf2. *The Journal of biological chemistry*
1181   **281**, 4969-4976 (2006). https://doi.org:10.1074/jbc.M511902200

1182   9   Lamanna, W. C. *et al.* Heparan sulfate 6-O-endosulfatases: discrete in vivo activities
1183   and functional co-operativity. *The Biochemical journal* **400**, 63-73 (2006).
1184   https://doi.org:10.1042/BJ20060848

1185   10   Hur, K. *et al.* Up-regulated expression of sulfatases (SULF1 and SULF2) as prognostic
1186   and metastasis predictive markers in human gastric cancer. *The Journal of pathology*
1187   **228**, 88-98 (2012). https://doi.org:10.1002/path.4055

1188   11   Vicente, C. M., Lima, M. A., Yates, E. A., Nader, H. B. & Toma, L. Enhanced
1189   tumorigenic potential of colorectal cancer cells by extracellular sulfatases. *Molecular*
1190   *cancer research : MCR* **13**, 510-523 (2015). https://doi.org:10.1158/1541-7786.MCR-
1191   14-0372

1192   12   Van der Sluis, M. *et al.* Muc2-deficient mice spontaneously develop colitis, indicating
1193   that MUC2 is critical for colonic protection. *Gastroenterology* **131**, 117-129 (2006).
1194   https://doi.org:10.1053/j.gastro.2006.04.020

1195   13   Tsai, H. H., Dwarakanath, A. D., Hart, C. A., Milton, J. D. & Rhodes, J. M. Increased
1196   faecal mucin sulphatase activity in ulcerative colitis: a potential target for treatment.
1197   *Gut* **36**, 570-576 (1995). https://doi.org:10.1136/gut.36.4.570

1198   14   Luis, A. S. *et al.* A single sulfatase is required to access colonic mucin by a gut
1199   bacterium. *Nature* **598**, 332-337 (2021). https://doi.org:10.1038/s41586-021-03967-
1200   5

1201   15   Velcich, A. *et al.* Colorectal cancer in mice genetically deficient in the mucin Muc2.
1202   *Science* **295**, 1726-1729 (2002). https://doi.org:10.1126/science.1069094

1203   16   Bloom, S. M. *et al.* Commensal Bacteroides species induce colitis in host-genotype-
1204   specific fashion in a mouse model of inflammatory bowel disease. *Cell host &*
1205   *microbe* **9**, 390-403 (2011). https://doi.org:10.1016/j.chom.2011.04.009

1206   17   Hickey, C. A. *et al.* Colitogenic Bacteroides thetaiotaomicron Antigens Access Host
1207   Immune Cells in a Sulfatase-Dependent Manner via Outer Membrane Vesicles. *Cell*
1208   *host & microbe* **17**, 672-680 (2015). https://doi.org:10.1016/j.chom.2015.04.002

1209   18   Bojarova, P. & Williams, S. J. Aryl sulfamates are broad spectrum inactivators of
1210   sulfatases: effects on sulfatases from various sources. *Bioorg Med Chem Lett* **19**,
1211   477-480 (2009). https://doi.org:10.1016/j.bmcl.2008.11.059

1212   19   Thomas, M. P. & Potter, B. V. Discovery and Development of the Aryl O-Sulfamate
1213   Pharmacophore for Oncology and Women's Health. *Journal of medicinal chemistry*
1214   **58**, 7634-7658 (2015). https://doi.org:10.1021/acs.jmedchem.5b00386

1215   20   Luis, A. S. *et al.* Sulfated glycan recognition by carbohydrate sulfatases of the human
1216   gut microbiota. *Nature chemical biology* **18**, 841-849 (2022).
1217   https://doi.org:10.1038/s41589-022-01039-x

1218   21   Potter, B. V. L. SULFATION PATHWAYS: Steroid sulphatase inhibition via aryl
1219   sulphamates: clinical progress, mechanism and future prospects. *J Mol Endocrinol*
1220   **61**, T233-T252 (2018). https://doi.org:10.1530/JME-18-0045

1221   22   Palmieri, C., Januszewski, A., Stanway, S. & Coombes, R. C. Irosustat: a first-
1222   generation steroid sulfatase inhibitor in breast cancer. *Expert Rev Anticancer Ther*
1223   **11**, 179-183 (2011). https://doi.org:10.1586/era.10.201

1224   23   Elger, W. *et al.* Estrogen sulfamates: a new approach to oral estrogen therapy.
1225   *Reprod Fertil Dev* **13**, 297-305 (2001). https://doi.org:10.1071/rd01029

1226   24   Woo, L. W. *et al.* Steroidal and nonsteroidal sulfamates as potent inhibitors of
1227   steroid sulfatase. *Journal of medicinal chemistry* **41**, 1068-1083 (1998).
1228   https://doi.org:10.1021/jm970527v

1229   25   Palmieri, C. *et al.* IRIS study: a phase II study of the steroid sulfatase inhibitor
1230   Irosustat when added to an aromatase inhibitor in ER-positive breast cancer
1231   patients. *Breast Cancer Res Treat* **165**, 343-353 (2017).
1232   https://doi.org:10.1007/s10549-017-4328-z

1233   26   Palmieri, C. *et al.* IPET study: an FLT-PET window study to assess the activity of the
1234   steroid sulfatase inhibitor irosustat in early breast cancer. *Breast Cancer Res Treat*
1235   **166**, 527-539 (2017). https://doi.org:10.1007/s10549-017-4427-x

1236   27   Pautier, P. *et al.* A Phase 2, Randomized, Open-Label Study of Irosustat Versus
1237   Megestrol Acetate in Advanced Endometrial Cancer. *Int J Gynecol Cancer* **27**, 258-
1238   266 (2017). https://doi.org:10.1097/IGC.0000000000000862

1239   28   Thomas, M. P. & Potter, B. V. Estrogen O-sulfamates and their analogues: Clinical
1240   steroid sulfatase inhibitors with broad potential. *J Steroid Biochem Mol Biol* **153**,
1241   160-169 (2015). https://doi.org:10.1016/j.jsbmb.2015.03.012

1242   29   Byrne, D. P., London, J. A., Eyers, P. A., Yates, E. A. & Cartmell, A. Mobility shift-based
1243   electrophoresis coupled with fluorescent detection enables real-time enzyme
1244   analysis of carbohydrate sulfatase activity. *The Biochemical journal* **478**, 735-748
1245   (2021). https://doi.org:10.1042/BCJ20200952

1246   30   Schelwies, M. *et al.* Glucosamine-6-sulfamate analogues of heparan sulfate as
1247   inhibitors of endosulfatases. *Chembiochem : a European journal of chemical biology*
1248   **11**, 2393-2397 (2010). https://doi.org:10.1002/cbic.201000401

1249   31   Chiu, L. T. *et al.* Trisaccharide Sulfate and Its Sulfonamide as an Effective Substrate
1250   and Inhibitor of Human Endo-O-sulfatase-1. *J Am Chem Soc* **142**, 5282-5292 (2020).
1251   https://doi.org:10.1021/jacs.0c00005

1252   32   Tseng, K. Y., Tzeng, Z. H., Cheng, T. R., Liang, P. H. & Hung, S. C. Design and Synthesis
1253   of 1-O- and 6'-C-Modified Heparan Sulfate Trisaccharides as Human Endo-6-O-
1254   Sulfatase 1 Inhibitors. *Front Chem* **10**, 947475 (2022).
1255   https://doi.org:10.3389/fchem.2022.947475

1256   33   Berteau, O., Guillot, A., Benjdia, A. & Rabot, S. A new type of bacterial sulfatase
1257   reveals a novel maturation pathway in prokaryotes. *The Journal of biological*
1258   *chemistry* **281**, 22464-22470 (2006). https://doi.org:10.1074/jbc.M602504200

34 Benjdia, A., Martens, E. C., Gordon, J. I. & Berteau, O. Sulfatases and a radical S-
adenosyl-L-methionine (AdoMet) enzyme are key for mucosal foraging and fitness of
the prominent human gut symbiont, Bacteroides thetaiotaomicron. *The Journal of*
biological chemistry **286**, 25973-25982 (2011).
https://doi.org:10.1074/jbc.M111.228841
35 Tintelnot, J. *et al.* Microbiota-derived 3-IAA influences chemotherapy efficacy in
pancreatic cancer. *Nature* **615**, 168-174 (2023). https://doi.org:10.1038/s41586-023-
05728-y
36 Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria.
Nature **555**, 623-628 (2018). https://doi.org:10.1038/nature25979
37 Ricaurte, D. *et al.* High-throughput transcriptomics of 409 bacteria-drug pairs reveals
drivers of gut microbiota perturbation. *Nature microbiology* (2024).
https://doi.org:10.1038/s41564-023-01581-x
38 Copeland, R. A. Evaluation of enzyme inhibitors in drug discovery. A guide for
medicinal chemists and pharmacologists. *Methods Biochem Anal* **46**, 1-265 (2005).
39 Mateus, A. *et al.* Thermal proteome profiling in bacteria: probing protein state in
vivo. *Mol Syst Biol* **14**, e8242 (2018). https://doi.org:10.15252/msb.20188242
40 Martens, E. C. *et al.* Recognition and degradation of plant cell wall polysaccharides
by two human gut symbionts. *PLoS biology* **9**, e1001221 (2011).
https://doi.org:10.1371/journal.pbio.1001221
41 Arjes, H. A. *et al.* Construction and characterization of a genome-scale ordered
mutant collection of Bacteroides thetaiotaomicron. *BMC Biol* **20**, 285 (2022).
https://doi.org:10.1186/s12915-022-01481-2
42 Jaeger, T., Arsic, M. & Mayer, C. Scission of the lactyl ether bond of N-acetylmuramic
acid by Escherichia coli "etherase". *The Journal of biological chemistry* **280**, 30100-
30106 (2005). https://doi.org:10.1074/jbc.M502208200
43 Jerga, A., Lu, Y. J., Schujman, G. E., de Mendoza, D. & Rock, C. O. Identification of a
soluble diacylglycerol kinase required for lipoteichoic acid production in Bacillus
subtilis. *The Journal of biological chemistry* **282**, 21738-21745 (2007).
https://doi.org:10.1074/jbc.M703536200
44 Miller, D. J., Jerga, A., Rock, C. O. & White, S. W. Analysis of the Staphylococcus
aureus DgkB structure reveals a common catalytic mechanism for the soluble
diacylglycerol kinases. *Structure* **16**, 1036-1046 (2008).
https://doi.org:10.1016/j.str.2008.03.019
45 Wang, Z. *et al.* Molecular basis of sphingosine kinase 1 substrate recognition and
catalysis. *Structure* **21**, 798-809 (2013). https://doi.org:10.1016/j.str.2013.02.025
46 Cartmell, A. *et al.* How members of the human gut microbiota overcome the
sulfation problem posed by glycosaminoglycans. *Proceedings of the National*
Academy of Sciences of the United States of America **114**, 7037-7042 (2017).
https://doi.org:10.1073/pnas.1704367114
47 Ndeh, D. *et al.* Metabolism of multiple glycosaminoglycans by Bacteroides
thetaiotaomicron is orchestrated by a versatile core genetic locus. *Nature*
communications **11**, 646 (2020). https://doi.org:10.1038/s41467-020-14509-4
48 Badola, P. & Sanders, C. R., 2nd. Escherichia coli diacylglycerol kinase is an
evolutionarily optimized membrane enzyme and catalyzes direct phosphoryl
transfer. *The Journal of biological chemistry* **272**, 24176-24182 (1997).
https://doi.org:10.1074/jbc.272.39.24176

49   An, D., Na, C., Bielawski, J., Hannun, Y. A. & Kasper, D. L. Membrane sphingolipids as
essential molecular signals for Bacteroides survival in the intestine. *Proceedings of*
the National Academy of Sciences of the United States of America **108 Suppl 1**, 4666-
4671 (2011). https://doi.org:10.1073/pnas.1001501107

50   Moggridge, S., Sorensen, P. H., Morin, G. B. & Hughes, C. S. Extending the
Compatibility of the SP3 Paramagnetic Bead Processing Approach for Proteomics. *J*
Proteome Res **17**, 1730-1740 (2018).
https://doi.org:10.1021/acs.jproteome.7b00913

51   Smits, A. H. *et al.* Biological plasticity rescues target activity in CRISPR knock outs.
Nat Methods **16**, 1087-1093 (2019). https://doi.org:10.1038/s41592-019-0614-5

52   Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I.
MSFragger: ultrafast and comprehensive peptide identification in mass
spectrometry-based proteomics. *Nat Methods* **14**, 513-520 (2017).
https://doi.org:10.1038/nmeth.4256

53   da Veiga Leprevost, F. *et al.* Philosopher: a versatile toolkit for shotgun proteomics
data analysis. *Nat Methods* **17**, 869-870 (2020). https://doi.org:10.1038/s41592-020-
0912-y

54   Franken, H. *et al.* Thermal proteome profiling for unbiased identification of direct
and indirect drug targets using multiplexed quantitative mass spectrometry. *Nat*
Protoc **10**, 1567-1593 (2015). https://doi.org:10.1038/nprot.2015.101

55   Grosdidier, A., Zoete, V. & Michielin, O. Fast docking using the CHARMM force field
with EADock DSS. *J Comput Chem* **32**, 2149-2159 (2011).
https://doi.org:10.1002/jcc.21797

56   Lebedev, A. A. *et al.* JLigand: a graphical tool for the CCP4 template-restraint library.
Acta crystallographica. Section D, Biological crystallography **68**, 431-440 (2012).
https://doi.org:10.1107/S090744491200251X

1332

Figure 1 General example of the conserved S1 sulfatase fold and consensus active site sequence.
a. The 3-D structure of the human lysosomal sulfatase GALNS (PDB:4FDJ), the N-terminal domain is coloured red and yellow for alpha helices and beta sheets, with the C-terminal subdomain coloured cyan and magenta for alpha helices and beta sheets; a zoom in of the consensus sequence of GALNS with a converted formylglycine, and with the general consensus pictured to the right; **b.** Structure of phase I and II arylsulfamate drugs developed to inhibit STS **c.** Chemical structures of aryl- and carbohydrate sulfamate inhibitors used in the study. **d.** Chemical structures of phenylvinyl sulfone, phenylsulfonyl fluoride, phenylvinyl sulfonate, and carbohydrate vinyl sulfone.

Figure 2 The effectiveness of arylsulfamate inhibitors against S1 steroid and carbohydrate sulfatases.
a. The enzymes PaAsta and HpSulf, belonging to S1_4 (green bar) and S1_2 (yellow bar), respectively, assayed against the panel of arylsulfamates (1-19 listed in Figure 2). **b**. The S1_11 sulfatases (BT3177 6S-GlcNAc, BT4656 6S-GlcNAc, Amuc_1033 6S-GlcNAc, Amuc_1074 6S-GlcNAc; blue bar above), the S1_20 sulfatases (BT1622 3S-Gal/GalNAc, BT1636 3S-Gal, Amuc_0451 3S-Gal, Amuc_0491 3S-Gal; pink bar above), and the S1_16 enzymes (BT3796 4S-Gal/GalNAc and BT1755 4S-Gal; orange bar above) against the panel of arylsulfamates (1-19 listed in Figure 2). For both a and b the left two panels display relative activity to wildtype protein with the arylsulfamate compound in the assay, whilst the left two panels are overnight preincubation of the enzyme with selected arylsulfamates then dilution of this preincubation mix into the reaction. The numbers indicate which arylsulfamates from Figure 2 were used and in all instances 1 mM of compound was deployed. Wt indicates reaction with enzyme and no arylsulfamate. Kinetic curves from which the data are derived are shown in supplemental figures S1-S24.

Figure 3 Growth of Bacteroides thetaiotaomicron on varied carbon sources in the presence of arylsulfamate inhibitors

Bacteroides thetaiotaomicron (B.theta) grown on varied carbon sources; two rich carbon sources, brain heart infusion media (BHI) and glucose (Glc); two sulfated host glycan carbon sources, chondroitin sulfate c (CSC) and Heparin (Hep), and two plant glycan carbon sources potato galactan (PG) and Larch arabinogalactan (LAG), in the presence of arylsulfamates inhibitors, labelled 1-19 (see Figure 2 for structures). Complex glycans and arylsulfamates inhibitors were used at a concentration of 5 mg/ml and 1 mM, respectively. All carbon source were utilised in a prepared minimal media except BHI which was purchased from sigma and simply dissolved in water. The traces are the mean of 3 independent growths but errors bars have been omitted for clarity; individual graphs with error bars are presented in supplemental figures S25-S30.

Figure 4. The effectiveness of carbohydrate sulfamate substrate mimics against the S1 carbohydrate sulfatase BT1636[3S-Gal].

a. Structure of the carbohydrate sulfonate/sulfamates and their aryl equivalents. **b.** The graphs show the effectiveness of carbohydrate sulfamates, carbohydrate sulfovinyl, and arylsulfonates against BT16363S-Gal; the left graph shows relative activity to wildtype with the compounds included in the assay, whilst the right graph show the relative activity after the enzyme had been preincubated with the compounds overnight then diluted into the reaction mixture; **c.** Thermostability shift assays (TSA) using inactive BT1636[3S-Gal] against the substrate 3S-Gal, and the putative inhibitor structures carbohydrate sulfamates, carbohydrate sulfovinyl, and arylsulfonates. All compounds in b were used at a concentration of 1 mM.

Figure 5. Arylsulfamates broadly affect Bacteroides spp. of the HGM
Bacteroidota species were grown in BHI media in the presence, or absence, of 1 mM of the listed arylsulfamate. The data plotted are the maximal OD_{600nm} observed for arylsulfamate treated growths relative to the untreated controls which were both ran in triplicate and randomly paired then averages derived. For all the estradiol growths a single experiment was conducted except for *B. theta* which was ran in triplicate; this was due to limited material.

Figure 6. Global protein thermal stability reveals off target interations of arysulfamate inhibitors 3 and 12.
a. Table of proteins which displayed an altered Tm in the presence of arylsulfamates 3 and 12. **b.** Proteins identified with a significance change in melting point upon inhibitor 3 treatment (n=2). **c.** Proteins identified with a significance change in melting point upon inhibitor 12 treatment (n=2). A dashed box indicates BT4322 is the only protein common across both groups. **d.** Bacteroides species were grown in BHI media in the presence, or absence, of 0.125 mM or 0.0625 mM diacylglycerol kinase inhibitor I and II, respectively. The data plotted are the maximal OD_{600nm} observed for inhibitor treated growths relative to the untreated controls which were both ran in triplicate and randomly paired then averages derived.

Extended data 1 Cell lysates of *Bacteroides thetaiotaomicron* incubated with two arylsulfamates inhibitors

a.   The effects of varied concentrations of arylsulfamate 3 and 12 on the growth of *B. theta* in BHI media.

b. Thin layer chromatography (TLC) analysis of cell lysates from *B.theta* cells grown to mid-exponential phase on CSA as a carbon source, and incubated with arylsulfamates 4 and 13 (Figure 2), mixed 1:1 with the sulfated substrates: O2 sulfated uronic acid β1,3 linked to O4 sulfated N-acetylgalactosmaine (UA2S-GalNAc4S), uronic acid β1,3 linked to O4 sulfated N-acetylgalactosmaine (UA-GalNAc4S), and O6 sulfated N-acetylgalactosmaine (GalNAc6S). Below are high performance anion exchange chromatography (HPAEC) traces showing the production of GalNAc in all instances except the negative control; **c**. Thin layer chromatography (TLC) analysis of cell lysates from *B.theta* cells grown to mid-exponential phase on Heparin as a carbon source, and incubated with arylsulfamates 4 and 13 (Figure 2), mixed 1:1 with the sulfated substrates: O2 sulfated uronic acid α1,4 linked to O4 sulfated N-acetylgalactosmaine (UA2S-GalNAc4S), O6 sulfated N-acetylglucosmaine (GlcNAc6S), and N-sulfated glucosamine (GlcNS). Below are high performance anion exchange chromatography (HPAEC) traces showing the production of GlcNAc in all instances except the negative control. All reactions were in PBS, with 5 % DMSO, and incubated at 37ºC overnight.

Extended data 2 Activity of carbohydrate sulfatases on para-nitrophenol sulfate.
Sulfatases were incubated with 5 mM para-nitrophenol sulfate over night in 100 mM BTP pH 7.0 with 150 mM NaCl and 5 mM CaCl2. Reactions were then ran on silica based thin layer chromatography using a resolving buffer of butanol:acetic acid:water (2:1:1) and develop by soaking the plates with 1 M NaOH and heating with a heat gun. It can be seen only the steroid sulfatase PaAsta shows the production of a para-nitrolphenol product.

Extended data 3 Global features of the thermal proteome profiling data
a. Distribution of the melting temperature (Tm) of *Bacteroides thetaiotaomicron* VPI-5482 (*B. theta*)
lysate proteome and the whole cell *E.coli* proteome. **b.** Ranked protein abundance plot of all identified
proteins in the 1%DMSO treated controls, S1 sulfatase are highlighted. **c.** Relative abundance of 11
S1 sulfatases identified by mass spectrometry for *B. theta* cultured in the presence of chondroitin sulfate A
(n=6). (b) Melting curve profiles for all identified *B. theta* proteins in control (1%DMSO) treated lysate.
The thermal stability profiles for each individual S1 sulfatases are highlighted in blue. **d.** Melting curve
profiles for all identified *B. theta* proteins in control (1%DMSO) treated lysate. The thermal stability profiles
for each individual S1 sulfatases are highlighted in blue.

Extended data 4. Global protein thermal stability analysis confirms that arysulfamate inhibitors 3 and 12 do not directly interact with S1 sulfatases.

Individual melt curve for selected S1 sulfatase for control and arysulfamate inhibitors 3 and 12 treated lysates (n=2 per treatment group).

Extended data 5 Structure of BT4322, SaDgKB, and SphK1 and comparison of the ATP helix
a. Tertiary structures of BT4322, SaDgKB, and SphK1 shown as with a helices, b sheets, and loops coloured cyan, magenta, and pink. The 'ATP helix' is coloured in green. **b.** An expanded view of the ATP helix and its sequence.

Extended data 6 Sequence alignment of putative DAGKs from select Bacteroides species.
Sequences were aligned using the MAFFT online server and visualised using Jalview.

Extended data 7. The effects of sphigosine kinase inhibitors on the growth of HGM Bacteroidota species
Bacteroidota species were grown in BHI, 1 % DMSO, with 0.1 mM of sphingosine kinase inhibitor (SKI) and 0.5 mM sphingosine kinase 1 inhibitor II and the effects on growth observed. Data are technical triplicates with the standard error of the mean.

Extended data 8. The effects of diacylglycerol kinase inhibitors on the growth of HGM Bacteroidota species
Bacteroidota species were grown in BHI, 1 % DMSO, with 0.125 mM of diacylglycerol kinase inhibitor I (DAGKI-i) and 0.0625 mM diacylglycerol kinase inhibitor II (DAGKI-ii) and the effects on growth observed. Data are technical triplicates with the standard error of the mean.

Extended data 9 Blind docking of arylsulfamates onto BT4322 and SpkH1.
The results of blind docking (no site specified) arylsulfamate inhibitors into the alphafold model of BT4322 (Q89ZQ4) and the crystal structure of SpKH1 (3VZB).
It can be observed all arylsulfamates preferentially cluster in, or near, the lipid binding pocket. For SpKH1 there appears to be no single preferred clustering site.