Arylsulfamates inhibit colonic *Bacteroides* growth through lipid kinases and not sulfatases

Conor J Crawford[1+], Charles Tomlinson[2,3,4+], Christian Gunawan[5], Zongjia Chen[5], Dominic P Byrne[6], Cosette Darby[7], Martina L. G. Conti[2], Tony Larson[2], Ana S Luis[8,9], Stefano Elli[10], Edwin A Yates[6], David N Bolam[7], Sjoerd van der Post[8], Spencer J Williams[5]*, and Alan Cartmell[2,3,4]*

[1]Max Planck Institute of Colloids and Interfaces, Am Mühlenberg 1, 14476, Potsdam, Germany
[2]Department of Biology, University of York, Wentworth Way, York, YO10 5DD, U.K. w
[3]York Structural Biology Laboratory (YSBL), University of York, Wentworth Way, York, YO10 5DD, U.K.
[4]York Biomedical Research Institute (YBRI), University of York, Wentworth Way, York, YO10 5DD, U.K.
[5]School of Chemistry and Bio21, Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria3010, Australia
[6]Department of Biochemistry, Cell and Systems Biology, Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool L69 7ZB, U.K.
[7]Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Medical School, Framlington Place, Newcastle upon Tyne, NE2 4HH, U.K.
[8]Department of Medical Biochemistry and Cell Biology, University of Gothenburg, Gothenburg, Sweden
[9]SciLifeLab, University of Gothenburg, 41390 Gothenburg, Sweden
[10]Istituto di Ricerche Chimiche e Biochimiche G. Ronzoni, Milano,20133, Italy

+Indicates these authors contributed equally

*To whom correspondence should be addressed:
alan.cartmell@york.ac.uk
sjwill@unimelb.edu.au

Abstract

Excessive degradation of the colonic mucin layer by *Bacteroides* within the human gut microbiota drives inflammatory bowel disease in mice. Bacterial carbohydrate sulfatases are key enzymes in gut colonization, as they are elevated in human inflammatory bowel disease and correlate with disease severity. Selective inhibitors of carbohydrate sulfatases could function as sulfatase-selective drugs, allowing precise control of sulfatase activity while preserving these otherwise beneficial bacteria. Arylsulfamates are covalent inhibitors that target a catalytic formylglycine residue of steroid sulfatases, a residue that is also conserved in carbohydrate sulfatases. Here, we find that a library of aryl- and carbohydrate sulfamates is ineffective against *Bacteroides* carbohydrate sulfatases, yet can inhibit human gut microbiota species grown on sulfated glycans. Leveraging thermal proteome profiling, we identify a lipid kinase as the target responsible for these effects. This work highlights the imperative for developing specific inhibitors targeting carbohydrate sulfatases and reveals the adverse effects that arylsulfamates have on *Bacteroides* species of the human gut microbiota.

49
50

Significance statement

Arylsulfamates are currently the only effective class of sulfatase inhibitors available and offer a potential strategy to treat inflammatory bowel disease driven by gut microbiota carbohydrate sulfatases. Although arylsulfamates inhibit the growth of microbiota *Bacteroides* species on sulfated glycans, this is not mediated through carbohydrate sulfatases but, via a conserved lipid kinase. Carbohydrate sulfatases are resistant to arylsulfamates whilst steroid sulfatases are susceptible despite a conserved active site. Finally, selected complex plant glycans confer a resistant/protective phenotype against the harmful effects of arylsulfamates. These data guide the future development of targeted carbohydrate sulfatase inhibitors and potential drug-prebiotic pairings.

Introduction

The human gut microbiota (HGM) is a microbial community found throughout the gastrointestinal tract but is densest in the distal colon where it is composed of trillions of bacteria. This community is critical to human health, providing essential vitamins[1,2], such as soluble B vitamins, calories through short chain fatty acid production[3], immune system regulation[4,5], and production of metabolites that influence the gut-brain axis[6,7]. Fermentation of complex carbohydrates underpins all these processes; the complex carbohydrates being derived either from dietary fibre (plant glycans) or the host. In the latter case this involves mainly sulfated glycans from colonic mucin and glycosaminoglycans.

The colonic mucin layer is the most abundant host glycan in the colon. It is composed of a gel forming glycoprotein, MUC2, is 80% glycan by mass and is heavily sulfated[8]. This layer has many biological functions, and provides a colonisable niche and food source for colonic bacteria, whilst simultaneously acting as largely impenetrable barrier protecting the colonic epithelium[9,10]. The Bacteroidota phylum are the major glycan degraders present in the HGM and are enriched in carbohydrate sulfatases, which are essential enzymes for utilisation of sulfated host glycans. In host symbiosis, the model Bacteroidota, *Bacteroides thetaiotaomicron* VPI-5482 (*B. theta*), grazes on colonic mucin *O*-glycans in a carbohydrate sulfatase dependent process[11]. However, in dysbiosis, excessive degradation of the sulfated colonic mucin drives inflammatory bowel disease (IBD) and carbohydrate sulfatases are the enzymatic drivers of this effect in a 'friend turned foe' scenario[12-14]. Excessive inflammation caused by loss of the mucin barrier is also a risk factor for colon cancer[15], the second leading cause of cancer deaths. Thus, effective inhibitors of HGM carbohydrate sulfatases are of high medical relevance.

Sulfatases are classified into four families, S1, S2, S3, and S4[16]. Of the over 160,000 sequences catalogued in the SulfAtlas database, the majority (over 90%, or >145,000 sequences) belong to the S1 family[17]. The S1 family occurs across all domains of life and is

the only family that contains sulfatases able to desulfate carbohydrates[16,18,19]. Members of

the S1 family of sulfatases have a conserved alkaline phosphatase-like fold characterised by

a larger N-terminal domain housing a central mixed β sheet flanked by α helices. This

domain abuts a smaller C-terminal subdomain composed of a 4-stranded antiparallel β sheet

and a single α helix[20] **(Figure 1a)**. Within this structural framework, the S1 family employs a

catalytic formylglycine (FGly) residue, which is generated co-translationally from a Cys or

Ser within the consensus sequence **C/S**-X-P/A-X-R, and is situated within an invariant

sulfate binding (S) site[18] **(Figure 1a)**.

The major class of sulfatase inhibitors are arylsulfamates, with the general formula

Ar-O-SO$_2$NH$_2$. These inhibitors, developed by replacing the sulfate group of assorted aryl

sulfate substrates with a sulfamate group, are irreversible, active-site directed inhibitors that

are broadly effective against steroid sulfatases of both bacterial and eukaryotic origin[21].

Clinical development of arylsulfamates targeting human steroid sulfatase (STS) for breast

cancer treatment has advanced to phase II clinical trials, with positive outcomes[22-24].

Arylsulfamates work by targeting the unusual catalytic FGly found in the invariant S site of

S1 family of sulfatases, and are believed to be pan-inhibitors of the family[25] but this not been

systematically tested.

Here, we investigate the ability of a panel of aryl- and carbohydrate

sulfamates/sulfonates, as well as vinyl sulfones and fluorosulfates **(Figure 1b,c)**, to inhibit

the growth of S1 sulfatase-containing Bacteroidota bacteria on sulfated glycans, and to

directly inhibit recombinant and lysate derived carbohydrate sulfatases from these species,

with the goal of developing novel, targeted strategies to treat IBD. Our data revealed that

sulfamate based inhibitors inhibit the growth of *Bacteroides* species on sulfated glycans but,

unexpectedly, do not inhibit any of the carbohydrate sulfatases from these organisms.

Leveraging thermal proteome profiling we identify lipid kinases, and not sulfatases, as the

targets through which arylsulfamates mediate their effects. We also show that arylsulfamate

inhibition of growth can be overcome through the utilisation of select, non-sulfated, plant

complex glycans. Our data suggest that arylsulfamates may have potential for the treatment

of IBD but at the expense of *Bacteroides* species, whilst also uncovering potential prebiotic

strategies to protect *Bacteroides* species of the HGM from arylsulfamate drugs when they

are delivered orally to treat cancer, or IBD.

Results

Arylsulfamates inhibit growth of Bacteroides thetaiotaomicron

Bacteroides species of the HGM encode large numbers of S1 sulfatases, which are induced

during growth on sulfated glycans. The model organism *Bacteroides thetaiotaomicron* VPI-

5482 (*B. theta*) encodes 28 S1 sulfatases, 16 of which have been characterised as
carbohydrate sulfatases. We hypothesized that growth of *B. theta* would be affected by the
action of arylsulfamates when grown on sulfated glycans, but not on unsulfated glycans.
Therefore, we cultivated *B. theta* in rich brain heart infusion media, and in minimal media
supplemented with sulfated (chondroitin sulfate C and heparin and unsulfated (D-glucose,
larch arabinogalactan, and potato galactan) carbon sources, each in the presence of a panel
of arylsulfamate compounds at 1 mM **(Figure 2)**. This panel includes 3 control compounds:
N-methyl and *N,N*-dimethyl sulfamates **6-8**, which should not inhibit S1 sulfatases. DMSO
was used at a final concentration of 1%, which did not affect growth of *B. theta* **(Figure S1)**.

We observed variable growth rates for *B. theta* with different arylsulfamate and
carbon source combinations **(Figure 2 and Figure S2-S7)**. Arylsulfamates based on a
naphthyl scaffold (compounds 11-17) caused severe growth defects on all substrates. In
contrast, the effect of arylsulfamates based on a substituted phenyl group (**1**, **3-6**) varied
substantially. *B. theta* displayed mild growth defects in the presence of these compounds
when grown on brain heart infusion media and glucose, while growth on the sulfated host
glycan chondroitin sulfate C, and the unsulfated plant glycan potato galactan, was strongly
perturbed by **1** and **5**. Compound **3** induced an increase in lag phase on chondroitin sulfate
C, and compound **4** caused complete inhibition of growth on the sulfated host glycan heparin
(Figure 2). On the unsulfated plant glycan larch arabinogalactan, **1** and **3** caused extension
of *B. theta* lag phase whilst **5** completely inhibited growth. Compound **2**, a 4-phenylphenyl
sulfamate, inhibited growth on all substrates. For the coumate scaffold, compounds **7-9** had
only mild effects when *B. theta* was grown on brain heart infusion media and glucose.
However, they increased the lag phase and reduced on chondroitin sulfate C , and
completely inhibited growth on heparin. Compounds **3** and **7** increased lag phase and
reduced growth on the plant glycan larch arabinogalactan, whilst compound **8** completely
inhibited growth. When potato galactan was used as the carbon source **3** caused a mild
growth defect whilst **7** reduced the lag phase; **8** caused a significant defect but still allowed
growth.

The variable impact of arylsulfamates on the growth of *B. theta*, observed for both
sulfated and unsulfated glycan substrates, suggests that these compounds do not target S1
carbohydrate sulfatases. Additionally, the fact that the 3 negative control *N*-
methylsulfamates (compounds **6-8**) also influenced growth of *B. theta* provides further
support that the observed growth defects are not mediated through an S1 sulfatase
dependent mechanism. A significant finding is that the same arylsulfamate structure has a
different effect depending on the carbon source utilised. Brain heart infusion media, and the
non-sulfated substrates potato galactan and glucose in a minimal media context, appear to

switch *B. theta* to a more resistant/protective metabolic state evinced by milder phenotypes

and fewer severe effects.

Arylsulfamates do not inhibit HGM B. theta S1 carbohydrate sulfatases

With the observation that the arylsulfamates, including the negative control *N*-

methylsulfamates, affected growth of *B. theta* on unsulfated and sulfated polysaccharides

we next wanted to investigate the effect of our panel of arylsulfamates on purified S1

sulfatases. We first confirmed the potency of our panel of arylsulfamates against two S1

steroid sulfatases: *Pa*AstA from *Pseudomonas aeruginosa*, belonging to S1 subfamily 4

(S1_4), and commercially available snail sulfatase *Hp*Sulf from *Helix pomatia*, belonging to

S1_2. The inhibitory activity of each compound was determined by measuring the sulfatase

activity of each enzyme incubated in the presence of inhibitor, or following ~24 h pre-

incubation and jump-dilution 'washout'[26]. At a concentration of 1 mM, all arylsulfamates were

broadly equivalent in their ability to completely inhibit both enzymes; only minor differences

being observed **(Figure 3, Figure S8-S11, Table S3 and S4).** Against *Pa*Asta, partial

inhibition was observed for **3** and **4** before complete inactivation **(Figure 3a, left panel and**

Figure S8), whilst pre-incubation with the two compounds resulted in complete inhibition

(Figure 3a, right panel). Competitive inhibition of *Pa*Asta was observed with **7** in the in

assay experiments but not in the enzyme incubation and jump dilution assays **(Figure 3a, b**

). Similarly, for *Hp*Sulf, residual activity observed when treated with **10** and **12** was

completely lost in the jump dilution experiment **(Figure 3a and Figure S11)**. These findings

suggest different inactivation rates between the enzymes and are consistent with the

irreversible, covalent, nature of the inactivation process. Importantly, near control levels of

activity were observed upon treatment with the *N*-methyl and *N,N*-dimethyl sulfamates **6-8**

(Figure 3a, right panel).

185   Having confirmed the activities of the arylsulfamates against two S1 steroid

sulfatases, we next examined their activity towards 10 bacterial S1 carbohydrate sulfatases

from the HGM members *B. theta* and *Akkermansia muciniphila ATCC BAA-835 (A. muc)*.

These enzymes target common linkages found in host glycans: four from S1_11 (BT3177[6S-

GlcNAc], BT4656[6S-GlcNAc], Amuc1033[6S-GlcNAc], and Amuc1074[6S-GlcNAc]) target *O*6 sulfated *N*-

acetyl-D-glucosamine (6S-GlcNAc), four from S1_20 (BT1636[3S-Gal], BT1622[3S-Gal/GalNAc],

Amuc0451[3S-Gal], and Amuc0491[3S-Gal/GalNAc]) target *O*3 sulfated D-galactose/*N*-acetyl-D-

galactosamine (3S-Gal/GalNAc), and two from S1_16 (BT3796[4S-Gal/GalNAc] and Amuc1755[4S-

Gal]) target *O*4 sulfated D-galactose/*N*-acetyl-D-galactosamine (4S-Gal/GalNAc). No

significant inhibition of any of the enzymes was observed when assaying their activity in the

presence of 1 mM arylsulfamate, even after pre-incubation **(Figure 3b, Supplementary**

Figure S12-S31, Table S3 and S4). Thus, while the panel of arylsulfamates inhibit bacterial
and eukaryotic S1 steroid sulfatases, they do not inhibit S1 HGM carbohydrate sulfatases.

We also explored the ability of two potent growth inhibitory sulfamates, **2** and **17**
(Supplementary Figure S32 and Figure S33), to inhibit S1 carbohydrate sulfatases in cell
lysates. Cell lysates derived from *B. theta* grown to mid-exponential growth phase on
chondroitin sulfate A (CSA) or Hep (to stimulate production of native S1 GAG carbohydrate
sulfatases) were incubated with arylsulfamates **2** and **17** for 1 hour, then sulfated mono- or
disaccharide substrates were added. Compounds **2** and **17** did not impact the production of
the final desulfated products (GlcNAc for Hep substrates and GalNAc for CSA substrates) as
judged by TLC and HPAEC **(Supplementary Figure S32b,c)**. Thus, compounds **2** and **17**
are not effective inhibitors of natively produced S1 GAG carbohydrate sulfatases, and the
effects observed on bacterial growth are likely to be due to effects on a non-sulfatase target.

Carbohydrate sulfamates and vinylsulfonates do not inhibit the S1_20 sulfatase BT1636^{3S-Gal}
The refractory nature of S1 carbohydrate sulfatases to arylsulfamate inhibition may arise
from the inability to bind at the active site of these enzymes, which has evolved to bind
sulfated glycans. Notably, S1 carbohydrate sulfatases exhibit low activity towards 4-
nitrophenyl sulfate (4NP-SO$_3$), which in contrast is rapidly hydrolysed by the family S1
steroid sulfatase *Pa*Asta **(Supplementary Figure S34)**. We selected the well-studied S1_20
sulfatase BT1636^{3S-Gal}, which desulfates 3*O* sulfated D-galactose (3S-Gal) and is essential
for *B. theta* to utilise colonic mucin *O*-glycans, and to competitively colonise the colon of
mice[11]. We prepared two substrate analogues, D-galactose-3-*O*-sulfamate (**20**) and 3-*O*-
vinylsulfonyl D-galactose (**25**), the isomeric compound D-galactose-2-*O*-sulfamate (**21**), aryl
vinyl sulfone analogues **22** and **24**, aryl fluorosulfate (**23**), and aryl sulfamate (**19**) **(Figure**
4a). Of these compounds, the only activity towards BT1636^{3S-Gal} was for compounds **22**, **23**
and **24 but only** after pre-incubation; **22** showed a ~3 fold increase in activity while **23** and
24 showed just over 50 % inhibition **(Figure 4b, Supplementary Figure S35, S36, and**
Table S5). We assessed whether any of these compounds could bind by studying the
inactive mutant Cys77Ser BT1636^{3S-Gal} using a thermal shift assay (TSA).). No change in
melting temperature was observed for compounds **20**, **21**, **23**, or **25** at concentrations up to
1.0 mM, whilst compounds **19, 22,** and **24** reduced the melting temperature (Tm),
suggesting binding causes protein destabilisation **(Figure 4c, Supplementary Figure**
Figure S37, and Table S6). In contrast, a substrate of BT1636^{3S-Gal}, 3S-Gal, stabilised the
protein (ΔTm = 3.0 ºC) at a concentration of 1 mM **(Figure 4c, Supplementary Figure S37**
and Table S6). Collectively, these data demonstrate that the carbohydrate sulfamate **20**,
and its vinylsulfonyl analogue **25**, are not inhibitors of BT1636^{3S-Gal} and do not bind the
enzyme.

Growth inhibition by arylsulfamates extends to multiple HGM Bacteroidota species

We selected nine additional HGM *Bacteroides* species to see if the growth effects observed for *B. theta* in the presence of arylsulfamate compounds **2** and **17 (Figure 2)** applied more broadly. We also included the arylsulfamate anticancer drugs Irosustat and estradiol sulfamate. No growth was observed for any species in the presence of 1 mM of arylsulfamate **2 (Figure 5 and Supplementary Figure S38)**. Arylsulfamate **17** exhibited growth defects across all species; in particular, *B. cellulosilyticus* for which no growth could be detected. Irosustat potently inhibited growth with all but one species, *B. fragilis*, showing limited or no growth **(Figure 5 and Supplementary Figure S39)**. Only very mild growth defects were observed for estradiol sulfamate **(Figure 5 and Supplementary Figure S39)**.

Thermal proteome profiling identifies a lipid kinase target for arylsulfamates

We next applied thermal proteome profiling (TPP) to determine the targets of arylsulfamate compounds **2** and **17**. TPP is a non-targeted mass spectrometry technique that can identify protein-compound interactions based on changes in a protein's thermal stability. Increased, or decreased, thermal stability is quantified as an increase, or decrease, in the relative abundance of peptides originating from the soluble protein fraction over a temperature gradient following incubation with a ligand[27].

Cell lysates were derived from *B. theta* grown on chondroitin sulfate A. *B. theta* lysates were treated with inhibitors **2** or **17**, or DMSO control and then heated at 10 different temperatures between 45–72 °C, and the soluble proteome subjected to trypsinisation, and analysed by liquid chromatography-mass spectrometry. The *B. theta* lysate proteome showed a relatively narrow melting temperature ranging between 50 and 58 °C (90% range) **(Supplementary Figure S40a)**. A total of 2550 proteins were identified in the combined analysis, from which 11 S1 sulfatases were identified, including the three upregulated in response to chondroitin sulfate A (BT1596, BT3333, and BT3349)[28] **(Supplementary Figure S40b,c)**. The melting temperature (Tm) was established for 2112 proteins **(Supplementary Figure S40d)** based on the detection of at least two unique peptides in a minimum of two replicate analyses (see Methods). Melting curves were determined for 8 out of the 11 identified S1 sulfatases **(Supplementary Figure S41)**, none of which was significantly affected by treatment with compounds **2** or **17**, indicating no direct interaction.

TPP identified 7 proteins for which arylsulfamate **2** induced significant changes in Tm, and 4 proteins for arylsulfamate **17**, indicating a potential arylsulfamate-protein interaction **(Figure 6a, b, c)**. Some of these hits can be discounted as causing the defective growth phenotype: disruption of BT0600, BT2504, and BT3464 through transposon mutagenesis results in no phenotype when grown on supplemented BHI media[29]. BT0133 is

a homologue of the *E. coli* protein MurQ (having 45% identity), a protein that produces
GlcNAc-6-phosphate from 6*O* phosphorylated *N*-acetyl-D-muramic acid (MurNAc). Loss of
MurQ in *E. coli* does not cause a no growth phenotype[30] and no MurNAc was detected in our
experiments. This leaves a total 7 candidate proteins (BT0218, BT1443, BT1889, BT4322,
BT4335, BT4346, and BT4480), of which only one was identified in TPP experiments with
arylsulfamates **2** and **17**, the uncharacterised protein BT4322 **(Figure 6b,c)**.

BT4322 is homologous to the cytoplasmic diacylglycerol kinases (DGK) YerQ and
DgkB, from the gram-positive bacteria *Bacillus subtilis*[31] and *Staphylococcus aureus*[32],
respectively, and the human sphingosine kinase SphK1[33]. YerQ is essential for growth of *B.*
subtilis and is involved in production of   lipoteichoic acid (LTA), a cell wall
glycophospholipid[31]. BT4322 is predicted to have the same fold and catalytic apparatus as
both DgkB (PDB:2QV7) and SphK1 (PDB:3VZB), and similar ATP binding site motifs and
lipid binding pockets **(Supplementary Figure S42)**. BT4322 is therefore likely to be a *B.*
theta lipid kinase, but the nature of the lipid substrate is yet to be determined. LTA is specific
to gram positive bacteria, and so BT4322 may act on another membrane lipid in *B. theta*, or
another as yet unknown target. BLASTp using BT4322 against the nine other arylsulfamate-
sensitive *Bacteroides* bacteria returned a single orthologue from each species, with 100%
query coverage, and a minimum of 84% identity suggesting a conserved role for these
proteins across these organisms **(Supplementary Figure S43 and Table S7)**.

289
Diacylglycerol kinase inhibitors significantly inhibit the growth of Bacteroides species
We next investigated whether we could recapitulate the arylsulfamate induced growth
defects assigned to inhibition of the putative lipid kinase BT4322 using commercially
available sphingosine and diacylglycerol kinase inhibitors. Diacylglcerol kinase inhibitor
(DAGKI)-I was effective at inhibiting the growth of all species with only *B. cellulosilyticus*, *B.*
intestinalis, and *B. oleicipnenius* showing limited, albeit highly variable, growth with extended
lag phases **(Figure 6d and Supplementary Figure S44)**. Bacteria treated with DAGKI-II
exhibited a milder phenotype, but this compound still inhibited the growth of all species
(except for *B. oleicipnenius*), lowering the final OD and extending the lag phase **(Figure 6d**
and Supplementary Figure S44). No growth defect could was observed with sphingosine
kinase inhibitor (SKI), whilst sphingosine kinase 1 inhibitor (SK1I)-II caused mild phenotypes
in four species, manifesting as an elongated lag phase, except for *B. xylanisolvens* which
showed severe defects **(Figure 6d and Supplementary Figure S45)**. These data support
the view that the observed arylsulfamate growth defects are mediated through inhibition of
BT4322, which functions as a diacylglycerol kinase, and not a sphingosine kinase.

305
Destabilising inhibitor interactions with BT4322 correlate with severe phenotypes

BT4322 was expressed recombinantly in *E. coli* and its interaction with inhibitors and ATP
assessed by DSF. Arylsulfamates **2** and **17** show a concentration dependent destabilisation
up to 1 mM **(Figure 7)** which is consistent with the TPP data **(Figure 6b,c)**. The DAGK-I
inhibitor also caused concentration-dependent destabilisation, similar to arylsulfamates **2**
and **17**, and correlates with a no growth phenotype for *B. theta* grown in the presence of
DAGK-I **(Figure 6d)**. By contrast DAGK-II caused a concentration dependent stabilisation
(Figure 7) and correlated with only a mild growth defect **(Figure 6d)**. The substrate ATP
also stabilised BT4322, supporting its activity as a lipid kinase.
315
Ligand docking of arylsulfamates into putative lipid kinase BT4322
To understand how arylsulfamates may interact with BT4322 we performed blind ligand
docking (no pocket specified) with arylsulfamates **2**, **17** and Irosustat using the AlphaFold 2
predicted model of BT4322. The docking results for Irosustat with BT4322 predicted binding
almost exclusively to the putative lipid binding pocket, with some minor solutions elsewhere.
Similar results were obtained for arylsulfamate **2** and **17**, but with some models also
predicting binding in the ATP site **(Supplementary Figure S46)**. This binding of the
arylsulfamates in the lipid pocket is consistent with what has been observed for other lipid
kinase inhibitors[32,34].
325
In vitro and in vivo effects of arylsulfamates on colonic bacteria
Growth of *B. theta* in the presence of sublethal doses of arylsulfamate **2**, and subsequent
lipidomic analysis, showed a drastically altered membrane lipid composition whilst only
causing a 15% reduction in total cell mass. These observations indicate arylsulfamate **2**
affects lipid metabolism, supporting the TPP target identification of a lipid kinase as
mediating the effects of arylsulfamates **(Figure 8a)**. The largest decrease ($\log^2$ FC ~-5) was
in the phosphorylated ceramides, suggesting ceramides as the potential lipid substrate for
BT4322, whilst large increase ($\log^2$ FC >8) was in membrane unphosphorylated triglycerides
were also observed **(Figure 8a)**.
In *in vivo* animal studies, where mice were gavaged with Irosustat, showed no
significant alteration of Bacteroidota and Bacillota levels, or at the genus level for
Bacteroides **(Figure 8b)**. However, these data lack the resolution to resolve at the species
level where it is possible changes may have occurred. Furthermore, any changes in
bacterial lipid composition *in vivo* were not assessed. Additionally, Irosustat may be
absorbed in the small intestine and not reach the colon to exert an effect on colonic bacterial
species.
342
Discussion

Arylsulfamates are effective irreversible, active site directed, inhibitors of S1 steroid
sulfatases[25] and display relatively mild side effects when administered to humans to treat
hormone dependent cancers in phase II clinical trials[24,35]. S1 carbohydrate sulfatases have
been implicated in driving several disease states and it was presumed that arylsulfamates
would be broadly effective across the S1 family. However, the work presented here shows
that bacterial S1 carbohydrate sulfatases of the HGM, despite sharing the invariant sulfate
binding subsite (S subsite) and a FGly nucleophile, are not inhibited by arylsulfamates *in*
vitro and *in vivo*. Furthermore, substrate mimicking carbohydrate sulfamate not only failed to
inhibit the target S1 carbohydrate sulfatase, but showed no evidence of binding **(Figure**
4b,c). Collectively, these data demonstrate that aryl or carbohydrate sulfamates are not
effective inhibitors of the HGM S1 carbohydrate sulfatases tested here. Similarly, the vinyl
sulfone and fluorosulfate modifications are ineffective inhibitors. With the S subsite being
invariant, the apparent differing susceptibilities of these enzyme classes must be due to the
distinctive chemical environments and/or structural geometries of glycan versus steroid
binding subsites. Steroid binding sites include a hydrophobic patch that can accommodate
lipophilic molecules and may allow the sulfamate moiety to adopt an alternative binding
conformation relative to when arylsulfamates bind to carbohydrate sulfatases. In contrast,
carbohydrates are hydrophilic, and S1 carbohydrate sulfatases have polar substrate binding
regions, which form extensive hydrogen bonding networks[11,20,36,37]. The polar, hydrogen
bond-driven nature of the glycan binding site may impose a level of rigidity on substrate
interactions that may be incompatible with alternative binding modes that may be necessary
to accommodate sulfamate groups. Moreover, the aryloxy group in arylsulfamates is a good
leaving group ($pK_a \leq 10$, depending on structure), whereas a carbohydrate alcohol is a much
poorer leaving group (pK_a value approximately 16).

Carbohydrate sulfamate inhibitors have been developed against the *endo*-acting
human S1 carbohydrate sulfatase Sulf1 by replacing the sulfate in HS di-, tri, and
tetrasaccharides with a sulfamate[38,39]. The most potent compound was the trisaccharide
sulfamate, with an IC_{50} value of 0.5 μM (versus 70 μM for the disaccharide), but this
compound operated through a competitive mechanism[39,38] distinct from the time-dependent
inactivation seen with arylsulfamates, with potency largely driven by the number of sugar
residues. In contrast, TSA gave no evidence of the monosaccharide sulfamate **20** binding to
the *exo*-acting BT1636[3S-Gal], suggesting that the sulfamate cannot interact with this sulfatase.

TPP analysis highlighted BT4322 as the most likely target through which the
arylsulfamates mediate their growth inhibiting effects on *B. theta* and lipidomic analyses of
B. theta, grown in the presence of arylsulfamate **2**, showed large changes in membrane lipid
composition supporting BT4322 as the target. Studies using lipid kinase inhibitors further
support BT4322 as a lipid kinase. A single homologue of BT4322, with high sequence

identity, is found in *B. theta* and the nine other species examined here, suggesting an
important role for BT4322 in phospholipid biosynthesis and a lack of functional redundancy
in *Bacteroides*. A homologue of BT4322 in *Bacillus subtilis* is implicated in peptidoglycan
metabolism and cell membrane integrity, and the corresponding genetic knock out mutant
displayed a no-growth phenotype[31]. *In vivo* mouse data upon oral dosing with Irosustat did
not identify any significant effects on the colonic microbiota at the phylum or *Bacteroides*
genus level. This may be as a result of rapid uptake of the drug in the stomach or small
intestine. It will be important to identify sulfamates that can transit to the distal gut and study
mice inoculated with HGM bacteria, which will allow studies of species level changes and
alteration of membrane lipid composition. Interestingly, it has recently been shown that
altered *B. theta* membrane lipid composition can lead to the transfer of bacterial lipids to the
host cell membrane[40].

This work reveals three findings with corresponding implications: (1) aryl and
carbohydrate sulfamates and sulfonates are not effective inhibitors of *exo*-acting HGM S1
carbohydrate sulfatases. There are no known inhibitors of these enzymes, and this is an
area where further research is needed. (2) Arylsulfamates have unexpected off-target effects
with negative impacts on HGM *Bacteroides* species. Potentially, the removal of these mucin
degrading bacteria could help treat IBD but the effects on the HGM may also lead to
undesirable side-effects on the host. (3) Selected non-sulfated complex carbohydrates
confer a resistant/protective phenotype on HGM *Bacteroides* species that can protect from
the harmful effects of arylsulfamates. This could help to mitigate the effect of arylsulfamates
(and potentially other drugs that transit it to the colon), opening the potential for paired drug-
glycan co-treatment strategies.

Acknowledgements
CJC was funded by MSCA grant MARINEGLYCAN (101029842) and acknowledges funding
from the Max Planck Society. SVDP was supported by the Swedish Research Council
(2020-02536). ASL was supported by Swedish Research Council grant (2021-01409)
Swedish Society for Medical Research (Svenska Sällskapet för Medicinsk Forskning, grant
S21-0026), Sahlgrenska Academy International Starting Grant (GU2021/1070), and
Jeanssons Foundation grant. This project has received funding from the Academy of
Medical Sciences/Wellcome Trust through the Springboard Grant (SBF005\1065 163470
awarded to AC), the Australian Research Council (DP210100235, DP240100126,
DP250100819), the Royal Society (RGS\R2\212050), and the Wellcome Trust (career
development award to AC, 225897/Z/22/Z).

Author contributions
AC conceived the project. AC, CJC and SW wrote the manuscript. AC, CT, DPB, and ASL
carried out enzymology experiments. AC, CT, and DPB analysed enzymological data. CJC,
EAY, and CG and ZC synthesised compounds. SE carried out computational analyses.
SVDP carried out and analysed proteomic experiments and data. AC, CT, CD, and DN
carried out bacterial growth experiments. MC and TL carried out lipidomic analysis. All
authors read and approved the manuscript.

Data availability statement
Source Data for all experiments, along with corresponding statistical test values, where appropriate, are provided within the paper and in Supplementary information. The mass spectrometry data has been deposited to the ProteomeXchange consortium via PRIDE with the dataset identified PXDXXXXXX.

Code availability statement
No new codes were developed or compiled in this study

Competing interest statement
The authors declare no competing interests.

Methods

Recombinant protein production.

Catalytically-active recombinant S1 sulfatases (in which Ser, within the C/S-X/P/A-X-R consensus sequence, for *B. thetaiotaomicron* sulfatases was mutated to Cys, to permit conversion to the formylglycine (FGly) residue by *E. coli*[18]), or catalytically-inactive sulfatases (in which Cys in the consensus sequence is mutated to Ser, thus preventing formation of FGly) were expressed in the *E. coli* strain TUNER (Novagen). Cultures were grown to mid-exponential phase in LB media supplemented with 50 μg/mL kanamycin at 37 °C, in an orbital shaker set to 180 rpm. Cultures were then cooled to 16 °C, and recombinant gene expression was induced by the addition of 0.1 mM isopropyl β-D-1-thiogalactopyranoside for 16 h at 16 °C, and 180 rpm. Cells were collected by centrifugation at 5,000 × g and pellets were resuspended in 20 mM HEPES, pH 7.4, with 500 mM NaCl, then were sonicated on ice. Recombinant protein was then purified by immobilized metal ion affinity chromatography using a cobalt-based matrix (Talon, Clontech) and, after a wash with resuspension buffer, eluted with a step gradient of 10, 50, and then 100 mM imidazole in resuspension buffer. Proteins were then analysed by SDS-PAGE and appropriately pure fractions were pooled and dialysed into 10 mM HEPES pH 7.0 with 150 mM NaCl. Protein concentrations were determined by measuring absorbance at 280 nm using the molar extinction coefficient calculated by ProtParam on the ExPasy server (web.expasy.org/protparam/).

Spectrophotometric based sulfatase assays

Production of *para*-nitrophenolate from *para*-nitrophenol sulfate by steroid sulfatases was monitored at A_{400nm} using a HIDEX sense or spectra max plus 384 (molecular devices) plate reader in a 96 well plate format. Reaction mixtures (100 μl) were monitored at ambient temperature (20-25 °C) and contained 1 mM substrate with 100 mM Bis-Tris-Propane, pH 7.0 with 5% (v/v) DMSO, 150 mM NaCl, and 5 mM $CaCl_2$.

Microfluidic-based de-sulfation assay

Reducing end BODIPY (maximal excitation/emission coefficient of ~503/511 nm) labelled sulfated substrates were detected using the EZ Reader II platform (Ret biochem) *via* LED-induced fluorescence, as described previously[41]. Real-time kinetic evaluation of substrate desulfation was achieved using a non-radioactive microfluidic mobility shift carbohydrate sulfation assays were optimised in solution with a 12-sipper chip coated with CR8 reagent and performed using a PerkinElmer EZ Reader II system employing EDTA-based separation buffer. Pressure and voltage settings were adjusted manually (1.8 psi, upstream voltage: 2250 V, downstream voltage: 500 V) to afford optimal separation in the reaction mixture of the sulfated substrate and unsulfated glycan product, with a sample (sip) time of 0.2 s, and total assay times appropriate for the experiment. Individual de-sulfation assays were carried out at 28°C and were pre-assembled in a 384-well plate in a volume of 80 µl in the presence of substrate concentrations of 1 µM with 100 mM Bis-Tris-Propane, MES, or Tris, dependent on the pH optimum of the sulfatase being assayed, and 5% DMSO, 150 mM NaCl, 0.02% (v/v) Brij-35 and 5 mM $CaCl_2$. The amount of de-sulfation was directly calculated in real-time using EZ Reader II software by measuring the sulfated carbohydrate : unsulfated carbohydrate ratio at each time-point during the assay. The activity of sulfatase enzymes was quantified in 'kinetic mode' by monitoring the amount of unsulfated glycan generated over the assay time, relative to control assay with no enzyme; with desulfation of the substrate limited to ~20% to prevent loss assay linearity via substrate depletion. k_{cat}/K_M values, using the equation $V_0 = (k_{cat}/K_M)/[E][S]$, were determined by linear regression analysis with GraphPad Prism software.

Differential scanning fluorimetry

Thermal shift/stability assays (TSAs) were performed using a StepOnePlus Real-Time PCR system (LifeTechnologies) and SYPRO-Orange dye, at a 1:1000 dilution, (excitation 470 nm, emission maximum 570 nm, Invitrogen) with thermal ramping between 20 and 95°C in 0.3°C step intervals per data point to induce denaturation of purified, folded, inactive, wild-type (Ser77) BT1636[S1–20] in the presence or absence of substrate or potential inhibitors. The melting temperature (Tm) corresponding to the midpoint for the protein unfolding transition was calculated by fitting the sigmoidal melt curve using the Boltzmann equation in GraphPad Prism, with R^2 values of ≥0.99, as described in[41]. Data points after the fluorescence intensity maximum were excluded from the fitting. Changes in the unfolding transition temperature compared with the control curve (ΔT_m) were calculated for each ligand. A positive ΔT_m value indicates that the ligand stabilises the protein from thermal denaturation, and confirms binding to the protein. All TSA experiments were conducted using a final protein

concentration of 5 µM in 100 mM Bis-Tris-Propane (BTP), pH 7.0, and 150 mM NaCl,

supplemented with the appropriate ligand concentration and 5 % DMSO. Three independent

assays were performed for each protein and protein ligand combination.

Bacterial growth experiments

Growth of various *Bacteroides* species was conducted in brain heart infusion media. BHI

[Sigma 53286-500G], 37g per litre) was dissolved in 18.2Ω deionised water, and then

sterilised by autoclaving. The media was then allowed to cool then haematin (1.2 mg/ml in

0.2 His-HCl pH 8.0) was added in a 1:1000 dilution. Five ml of the resulting BHI media was

then inoculated with *Bacteroides* species and grown overnight, and diluted 1:20 into 96 well

plate growths (10 µl in 200) to monitor growth. Growth of the *B. thetaiotaomicron* VPI-5482

on specified glycan sources was achieved by mixing 1:1 the 0.22 micron filter-sterilised

polysaccharides (10.2 mg/ml in H_2O) with 0.22 micron filter-sterilised 2 x minimal media (per

50 ml was: 0.1g of ammonium sulfate and sodium carbonate, 0.05g cysteine, 10 ml 1 M

potassium phosphate pH 7.2, 0.1 ml 1 mg.ml^{-1} vitamin K, 1 ml of 0.4 mg.ml^{-1} iron sulfate, 0.4

ml of 0.25 mg.ml resazurin, 0.05 ml of 0.01 mg.ml^{-1} vitamin B12, 5 ml of mineral salts for

defined media, and 100 µl of 1.2 mg/ml haematin in 0.2 M His-HCl pH8.0); this gives a final

glycan concentration of 5 mg/ml in a 1 x minimal media. All growth experiments were

monitored continuously in 96-well plates using a cerillo stratus plate reader within a Don

whitely VA-500, A85, or A35 cabinet configured for anaerobic conditions (80% N_2, 10% CO_2,

10% H_2). Where appropriate, either 1% DMSO (control growths) or 1% DMSO and 1 mM

arylsulfamates was added to growth conditions. For sphingosine kinase inhibitor (SKI,

567731) and sphingosine kinase 1 inhibitor II (SK1-II, 567741) a maximum of 0.1 and 0.5

mM, respectively, could be dissolved in BHI with 1% DMSO. For diacylglycerol kinase

inhibitors i (D5794-5MG) and ii (D5919-5MG) a maximum concentration of 125 µM and 62.5

µM, respectively, could be dissolved in BHI with 1% DMSO but were hazy; this did not affect

measurement readings at OD_{600nm}. Overnight growth cultures were diluted 1:20 into 96 well

plate growths (10 µl in 200 µl). Growth curves presented are averages of three technical

replicates.

Thin layer chromatography (TLC)

B. thetaiotaomicron VPI-5482 was grown in 10 ml minimal media with 10 mg/ml (w/v)

appropriate GAG (either CSA or Heparin) as the sole carbon source to mid-exponential

phase in glass test tubes. Cells were harvested by centrifugation at 5,000 × g for 10 min at

room temperature and washed 2 x with 5 ml PBS (pH 7.2) before being resuspended in 1

mL PBS. Cells washed in PBS were lysed and the lysates assayed against 5 mM of the

appropriate substrate at 37 °C for up to 24 h. Assays were analysed by TLC, and 2 μL each
sample was spotted onto silica plates and resolved in butanol:acetic acid:water (2:1:1)
mobile phase. The plates were dried, and the sugars were visualized using diphenylamine
stain (DPA: 1 ml of 37.5% HCl, 2 ml of aniline, 10 ml of 85% H_3PO_3, 100 ml of ethyl acetate,
and 2 g of diphenylamine) by heating using a heat gun set to 450 °C on medium flow.
Arylsulfates and their pNP products were visualised using 1 M NaOH and then heated as
described for DPA stains.
546

High performance anion exchange chromatography (HPAEC)
Sonicated bacterial cell lysates grown on the glycosaminoglycans CSA and heparin were
treated with 1 mM of arylsulfamates 3 and 12 for 1 hour then challenged with CSA or
heparin mono- and disaccharides by mixing 1:1, to a final concentration of 5 mM, to see if
sulfatase activity was still present. Analysis of the sugar products was performed by HPAEC
using an ICS-6000 with pulsed amperometic detection using a gold work electrode and pDH
reference electrode with a standard quad carbohydrate waveform. Separation was done
using a Dionex PA-200 analytical column preceded by a PA-200 guard column. Samples
were resolved isocratically in 100 mM NaOH for 20 mins, then the column cleaned with 500
mM NaOH for 10 mins, and finally ran back into 100 mM NaOH for 5 mins, with a flow rate of
0.25 ml/min.
558

Synthesis of chemical compounds
560 Full detail of the synthetic procedures can be found in the supporting information.
561

Thermal proteome profiling
A 300 ml culture of minimal media (see 'Bacterial growth experiments' section)
supplemented with 5 mg/ml chondroitin sulfate A (CSA), stored in the anaerobic chamber
overnight, and inoculated with 10 ml of *B. thetaiotaomicron* VPI-5482 was grown overnight in
BHI media. This culture was grown to a target OD_{600nm} of ~0.5-0.6 at which point cells were
centrifuged at 5000 x g and washed with PBS, this process was repeated two more times.
Cells were then treated as in *Mateus et al*[27]; briefly, cells were then resuspended in lysis
buffer (50μg/ml lysozyme, 1×protease inhibitor(Roche), 250 U/ml benzonase, and 1 mM
$MgCl_2$ in PBS) to give an OD_{600nm} of 50. Samples were then sonicated on ice for 15 seconds
three times. The lysate was then split into three: one sample was a control (1% (v/v) DMSO)
and experimental samples contained 1 mM arylsulfamate **3** or 1 mM arylsulfamate **12** (1%
(v/v) DMSO). Next 20 μl of each condition was aliquoted into a 96 well PCR plate across 10
wells in quadruplicate. The plates were then subjected to a temperature gradient from 45 –
72 °C (45, 46.8, 49.1, 52.1, 55.9, 60.3, 64.4, 67.4, 69.9, 72°C) for 3 min, followed by 3 min at

room temperature. NP-40 (nonyl phenoxypolyethoxylethanol) was then added to samples to

a final concentration of 0.8% (v/v). Samples were transferred to a 0.22 µm 96 well filter plate

(Millipore: MSGVS2210) and centrifuged at 500 x g for 5 min to remove protein aggregates.

Following filtration, the flowthrough was combined 1:1 with 400 mM Tris pH 8.0 and 4% SDS

(w/v). The protein concentration for each sample was determined by Bradford assay, and a

volume corresponding to 10 µg for the lowest-temperature condition was used as input for all

other samples in a set for analysis. Protein digestion was performed using a modified solid-

phase-enhanced sample preparation (SP3) based method adapted to a 96 well format in

0.45µM filter plates (Millipore, MSRPN04)[42,43]. Premixed magnetic SpeedBeads (Cytiva,

(1:1, GE45152105050250 and GE 65152105050250)) were added to the samples,

combined with two volumes of ethanol and incubated for 10 min while mixing at 150 RPM.

Samples were transferred to the filter plates and washed four times with 200 µL of 70%

ethanol (v/v) with centrifugation for 2 min at 2000 x g between each wash. Beads were

resuspended in 40 µL 0.1 mM HEPES pH 8.5 containing trypsin/Lys-c (Thermo Scientific),

1.25 mM TCEP and 5 mM chloroacetamide (Sigma-Aldrich) and incubated at room

temperature overnight while mixing 500 rpm. Peptides were collected by centrifugation,

followed by a second elution with 2% DMSO. TMT 10-plex labelling (Thermo Scientific) was

performed at a 2:1 ratio for 1 hr while mixing at 500 RPM. The labelling reaction was

quenched with 0.4% hydroxylamine (v/v) for 15 min before each sample was combined and

dried under vacuum. Peptides were resolved in 0.1% trifluoroacetic acid (TFA) and

fractionated using 0.1% triethylamine with increasing concentrations of acetonitrile by high

pH reverse phase into 8 fractions using C18 desalting columns (Pierce, 89852) pooled into 4

fractions and dried under a vacuum.

Lyophilized samples were redissolved in 0.1% TFA and analysed by LC-MS/MS

using a nano HPLC system (EASY-nLC 1200, Thermo Scientific) coupled to a Q-Exactive

HF-X mass spectrometer (Thermo Scientific). Peptides were separated using in-house

packed columns (150 x 0.075 mm) packed with Reprosil-Pur C18-AQ 3 µm particles (Dr.

Maisch). Elution was performed with a 5 to 45% gradient (A: 0.1% formic acid, B: 0.1%

formic acid ,80% acetonitrile) in 135 minutes at 250 nl. Full mass spectra were acquired over

a mass range of minimum 400 m/z and maximum 1600 m/z, with a resolution of 60,000 at

200 m/z. The isolation window for fragmentation spectra collection was set to 1Da and a

fixed first mass of 100 m/z with a resolution of 30,000 at 200 m/z. The top 15 most intense

peaks with a charge state ≥2 – 5 were selected for fragmentation by HCD with a normalized

collision energy of 32% and subsequent excluded for selection for 30 seconds. Thermo

.RAW files were converted into MZml using msConvert (proteowizard) and FragPipe (v20.0)

configured with Msfragger (v3.8) and Philosopher (v5.0)[44,45] was used for protein

identification and TMT quantification. Database searches were performed against the

Bacteroides thetaiotaomicron reference proteome (4782 entries, UP000001414, 2023_10)

with the following settings, parent ion mass and fragment ion mass accuracy after

recalibration was set to 20 ppm. Enzyme specificity was set to trypsin, fixed modifications

were set for cysteine carbamidomethylation and TMT modified lysine, variable modifications

considered were methionine oxidation, protein n-terminal acetylation and TMT modification

on peptide n-terminal or serine. The identified proteins and peptides were both filtered at a

1% and only identifications based on ≥2 unique peptides with a minimum ion purity of 0.75

were considered for quantification and thermal proteome profiling. Further data analysis was

performed in R environment to combine the search outcome, normalize reporter ion

channels. Meltome curve fitting analysis and statistical analysis were executed using the

TPP-TR workflow in the TPP package[46].Briefly, protein candidates with altered melting

temperature were considered based on the following criteria. P-values for the two replicate

experiments are <0.05 and <0.1, the melting point shifts in the vehicle versus treatment

experiments have the same direction. Both melting point differences in the two pairs of

control versus treatment experiments are greater than the melting point difference between

the two vehicle controls and the minimum slope in each of the control versus treatment

experiments was <0.06. In addition, proteins were only considered for melting curve analysis

if the identification was based on a minimum of two unique peptide identification; see

Dataset 1.

Ligand docking experiments

All docking experiments were conducted using SwissDock[47] on the Expasy server using

default parameter; no binding site was specified. The Alphafold model of BT4322 (Q89ZQ4)

and the crystal structure of SphK1 (3VZB) were used, and where appropriate heteroatoms

removed, and uploaded to the SwissDock server as PDB files. Small molecules to be

docked were built in Jligand[48], the generated PDB or CIF files converted to MOL2 files using

the openbabel server, and uploaded to SwissDock as MOL2 files.

Lipidomic analysis

All solvents used were HPLC or LC-MS grade. To the cell pellet, 997 µL of 2:1 methanol :

chloroform (MeOH:CHCl$_3$) and 208 µL of 0.005 N HCl were added. The tubes were vortexed

briefly and incubated on ice for 5 minutes. Following centrifugation at 18,000 xg for 5

minutes at 4°C the supernatant was transferred to a fresh tube. CHCl$_3$ (332 µL)) and H$_2$O

(332 µL) were added and briefly vortexed to form two phases. After centrifugation at 3600

xg for 1 minute at 4°C, the lower organic phase was transferred to a LC-MS tapered vial and

dried under a gentle stream of nitrogen. The samples were resuspended in 80 µL of 7:3

acetonitrile : isopropanol (ACN:IPA) for LC-MS analysis.

Untargeted lipid analysis was performed on bacterial pellets and data processed
using R scripts as previously described[49], except that data was collected on a Thermo
Orbitrap Exploris 480 mass spectrometer with MS1 and HCD ddMS2 scans collected at
120,000 and 15,000 resolution (FWHM), respectively. The following parameters were used
to acquire MS data: spray voltage 3500 V (positive mode), 2500 V (negative mode), sheath
gas 50, auxiliary gas 10, sweep gas 1, ion transfer tube temperature 280°C, vaporizer
temperature 200°C, RF lens 30, scan range m/z 270-2000. For MS2 parameters, mass
selection range m/z 400-2000, ramped HCD collision energies 30, 50, 70%. MS1 m/z values
were searched against candidate ions from positive and negative ionization mode adduct
databases generated from the LipidMAPS structural database (https://www.lipidmaps.org/)
and the list of Bacteroides lipids described in Barone *et al* 2024[40]. Candidate lipids were
additionally annotated by searching MS2 spectra using LipidMatch[50], LipidBlast[51] and
SIRIUS (version 6.1.0)[52].

Annotations were manually curated using either Barone's names, or the sum
composition notation described in LipidMAPS. To distinguish between multiple isomers,
annotations were prepended with a unique label indicating MS1 m/z values ("M") and
retention time in s ("T"). The feature list was filtered to remove background contaminants
(cutoff = 3 standard deviations above average background feature area), and any feature
where relative standard deviation exceeded 30%. Features that could not be annotated as
lipids were removed and where there were redundant measurements across positive and
negative ionization modes or adduct type, only the most abundant representative was kept.
This resulted in a final feature list of 63 lipids, covering ceramides (CE), menaquinones,
triglycerides (TG) , phosphatidylethanolamines (PE), and various glycolipids. Feature area
data was further processed through MetaboAnalyst 6.0 (https://www.metaboanalyst.ca)
using the Statistical Analysis (one factor) workflow. Data was sample normalized by median
before calculating fold-changes and log10 transformed before performing t-tests to calculate
p-values.

Mice experiments
All experimental procedures involving animals were approved by the Swedish Laboratory
Animal Ethical Committee at the University of Gothenburg. 8-week-old C57BL/6N mice were
maintained under standardized specific pathogen free conditions with ad libitum access to
food and water. Mice were treated at day 0, 1 and 2 by gavage with 3.2 mM of Irosustat
dissolved in PEG 400:water:DMSO (6:3:1). The control group was gavaged with the vehicle
only. Fecal samples for all animals were collected at day 0, 3 and 7. The weight of the mice
was recorded at the days the mice were gavaged or during fecal collection. Bacterial
genomic DNA was extracted with QIAamp PowerFecalPro DNA kit (Qiagen®) and

quantification by qPCR as described previously[53]. The following primers were used for
relative quantification of phylum Bacteroidetes (Fw 5'-CRAACAGGATTAGATACCT; Rv 5'-
GGTAAGGTTCCTCGCGTAT) , phylum Firmicutes (Fw 5'-TGAAACTYAAAGGAATTGAGG;
Rv 5'-ACCATGCACCACCTGTC) and genus Bacteroides (Fw 5'-GAAGGTCCCCCACATTG;
Rv 5'-CGCKACTTGGCTGGTTCAG). The relative quantification normalized of specific
taxons was determined for each sample using the 16S rRNA gene as reference.
693

References:
1 Goodman, A. L. *et al.* Identifying genetic determinants needed to establish a human
gut symbiont in its habitat. *Cell host & microbe* **6**, 279-289 (2009).
https://doi.org:10.1016/j.chom.2009.08.003
2 Nysten, J. & Van Dijck, P. Can we microbe-manage our vitamin acquisition for better
health? *PLoS Pathog* **19**, e1011361 (2023).
https://doi.org:10.1371/journal.ppat.1011361
3 McNeil, N. I. The contribution of the large intestine to energy supplies in man. *The
American journal of clinical nutrition* **39**, 338-342 (1984).
https://doi.org:10.1093/ajcn/39.2.338
4 Troy, E. B. & Kasper, D. L. Beneficial effects of Bacteroides fragilis polysaccharides on
the immune system. *Front Biosci (Landmark Ed)* **15**, 25-34 (2010).
https://doi.org:10.2741/3603
5 Belkaid, Y. & Hand, T. W. Role of the microbiota in immunity and inflammation. *Cell*
157, 121-141 (2014). https://doi.org:10.1016/j.cell.2014.03.011
6 Damdindorj, L. *et al.* A Comparative Analysis of Constitutive Promoters Located in
Adeno-Associated Viral Vectors. *PloS one* **9**, e106472 (2014).
https://doi.org:10.1371/journal.pone.0106472
7 Carabotti, M., Scirocco, A., Maselli, M. A. & Severi, C. The gut-brain axis: interactions
between enteric microbiota, central and enteric nervous systems. *Ann Gastroenterol*
28, 203-209 (2015).
8 Hansson, G. C. Mucins and the Microbiome. *Annu Rev Biochem* **89**, 769-793 (2020).
https://doi.org:10.1146/annurev-biochem-011520-105053
9 Johansson, M. E., Larsson, J. M. & Hansson, G. C. The two mucus layers of colon are
organized by the MUC2 mucin, whereas the outer layer is a legislator of host-
microbial interactions. *Proceedings of the National Academy of Sciences of the
United States of America* **108 Suppl 1**, 4659-4665 (2011).
https://doi.org:10.1073/pnas.1006451107
10 Kawashima, H. Roles of the gel-forming MUC2 mucin and its O-glycosylation in the
protection against colitis and colorectal cancer. *Biological & pharmaceutical bulletin*
35, 1637-1641 (2012).
11 Luis, A. S. *et al.* A single sulfatase is required to access colonic mucin by a gut
bacterium. *Nature* **598**, 332-337 (2021). https://doi.org:10.1038/s41586-021-03967-
5
12 Hickey, C. A. *et al.* Colitogenic Bacteroides thetaiotaomicron Antigens Access Host
Immune Cells in a Sulfatase-Dependent Manner via Outer Membrane Vesicles. *Cell
host & microbe* **17**, 672-680 (2015). https://doi.org:10.1016/j.chom.2015.04.002

13 Van der Sluis, M. *et al.* Muc2-deficient mice spontaneously develop colitis, indicating
that MUC2 is critical for colonic protection. *Gastroenterology* **131**, 117-129 (2006).
https://doi.org:10.1053/j.gastro.2006.04.020
14 Tsai, H. H., Dwarakanath, A. D., Hart, C. A., Milton, J. D. & Rhodes, J. M. Increased
faecal mucin sulphatase activity in ulcerative colitis: a potential target for treatment.
Gut **36**, 570-576 (1995). https://doi.org:10.1136/gut.36.4.570
15 Velcich, A. *et al.* Colorectal cancer in mice genetically deficient in the mucin Muc2.
Science **295**, 1726-1729 (2002). https://doi.org:10.1126/science.1069094
16 Barbeyron, T. *et al.* Matching the Diversity of Sulfated Biomolecules: Creation of a
Classification Database for Sulfatases Reflecting Their Substrate Specificity. *PloS one*
11, e0164846 (2016). https://doi.org:10.1371/journal.pone.0164846
17 Stam, M. *et al.* SulfAtlas, the sulfatase database: state of the art and new
developments. *Nucleic Acids Res* **51**, D647-D653 (2023).
https://doi.org:10.1093/nar/gkac977
18 Hanson, S. R., Best, M. D. & Wong, C. H. Sulfatases: structure, mechanism, biological
activity, inhibition, and synthetic utility. *Angewandte Chemie* **43**, 5736-5763 (2004).
https://doi.org:10.1002/anie.200300632
19 Luis, A. S., Yates, E. A. & Cartmell, A. Functions and specificity of bacterial
carbohydrate sulfatases targeting host glycans. *Essays Biochem* **67**, 429-442 (2023).
https://doi.org:10.1042/EBC20220120
20 Luis, A. S. *et al.* Sulfated glycan recognition by carbohydrate sulfatases of the human
gut microbiota. *Nature chemical biology* **18**, 841-849 (2022).
https://doi.org:10.1038/s41589-022-01039-x
21 Bojarova, P. & Williams, S. J. Aryl sulfamates are broad spectrum inactivators of
sulfatases: effects on sulfatases from various sources. *Bioorg Med Chem Lett* **19**,
477-480 (2009). https://doi.org:10.1016/j.bmcl.2008.11.059
22 Palmieri, C., Januszewski, A., Stanway, S. & Coombes, R. C. Irosustat: a first-
generation steroid sulfatase inhibitor in breast cancer. *Expert Rev Anticancer Ther*
11, 179-183 (2011). https://doi.org:10.1586/era.10.201
23 Thomas, M. P. & Potter, B. V. Estrogen O-sulfamates and their analogues: Clinical
steroid sulfatase inhibitors with broad potential. *J Steroid Biochem Mol Biol* **153**,
160-169 (2015). https://doi.org:10.1016/j.jsbmb.2015.03.012
24 Palmieri, C. *et al.* IRIS study: a phase II study of the steroid sulfatase inhibitor
Irosustat when added to an aromatase inhibitor in ER-positive breast cancer
patients. *Breast Cancer Res Treat* **165**, 343-353 (2017).
https://doi.org:10.1007/s10549-017-4328-z
25 Potter, B. V. L. SULFATION PATHWAYS: Steroid sulphatase inhibition via aryl
sulphamates: clinical progress, mechanism and future prospects. *J Mol Endocrinol*
61, T233-T252 (2018). https://doi.org:10.1530/JME-18-0045
26 Copeland, R. A. Evaluation of enzyme inhibitors in drug discovery. A guide for
medicinal chemists and pharmacologists. *Methods Biochem Anal* **46**, 1-265 (2005).
27 Mateus, A. *et al.* Thermal proteome profiling in bacteria: probing protein state in
vivo. *Mol Syst Biol* **14**, e8242 (2018). https://doi.org:10.15252/msb.20188242
28 Martens, E. C. *et al.* Recognition and degradation of plant cell wall polysaccharides
by two human gut symbionts. *PLoS biology* **9**, e1001221 (2011).
https://doi.org:10.1371/journal.pbio.1001221

29 Arjes, H. A. *et al.* Construction and characterization of a genome-scale ordered
mutant collection of Bacteroides thetaiotaomicron. *BMC Biol* **20**, 285 (2022).
https://doi.org:10.1186/s12915-022-01481-2
30 Jaeger, T., Arsic, M. & Mayer, C. Scission of the lactyl ether bond of N-acetylmuramic
acid by Escherichia coli "etherase". *The Journal of biological chemistry* **280**, 30100-
30106 (2005). https://doi.org:10.1074/jbc.M502208200
31 Jerga, A., Lu, Y. J., Schujman, G. E., de Mendoza, D. & Rock, C. O. Identification of a
soluble diacylglycerol kinase required for lipoteichoic acid production in Bacillus
subtilis. *The Journal of biological chemistry* **282**, 21738-21745 (2007).
https://doi.org:10.1074/jbc.M703536200
32 Miller, D. J., Jerga, A., Rock, C. O. & White, S. W. Analysis of the Staphylococcus
aureus DgkB structure reveals a common catalytic mechanism for the soluble
diacylglycerol kinases. *Structure* **16**, 1036-1046 (2008).
https://doi.org:10.1016/j.str.2008.03.019
33 Wang, Z. *et al.* Molecular basis of sphingosine kinase 1 substrate recognition and
catalysis. *Structure* **21**, 798-809 (2013). https://doi.org:10.1016/j.str.2013.02.025
34 Gustin, D. J. *et al.* Structure guided design of a series of sphingosine kinase (SphK)
inhibitors. *Bioorg Med Chem Lett* **23**, 4608-4616 (2013).
https://doi.org:10.1016/j.bmcl.2013.06.030
35 Palmieri, C. *et al.* IPET study: an FLT-PET window study to assess the activity of the
steroid sulfatase inhibitor irosustat in early breast cancer. *Breast Cancer Res Treat*
166, 527-539 (2017). https://doi.org:10.1007/s10549-017-4427-x
36 Cartmell, A. *et al.* How members of the human gut microbiota overcome the
sulfation problem posed by glycosaminoglycans. *Proceedings of the National*
Academy of Sciences of the United States of America **114**, 7037-7042 (2017).
https://doi.org:10.1073/pnas.1704367114
37 Ndeh, D. *et al.* Metabolism of multiple glycosaminoglycans by Bacteroides
thetaiotaomicron is orchestrated by a versatile core genetic locus. *Nature*
communications **11**, 646 (2020). https://doi.org:10.1038/s41467-020-14509-4
38 Schelwies, M. *et al.* Glucosamine-6-sulfamate analogues of heparan sulfate as
inhibitors of endosulfatases. *Chembiochem : a European journal of chemical biology*
11, 2393-2397 (2010). https://doi.org:10.1002/cbic.201000401
39 Chiu, L. T. *et al.* Trisaccharide Sulfate and Its Sulfonamide as an Effective Substrate
and Inhibitor of Human Endo-O-sulfatase-1. *J Am Chem Soc* **142**, 5282-5292 (2020).
https://doi.org:10.1021/jacs.0c00005
40 Mirretta Barone, C. *et al.* Spatially resolved lipidomics shows conditional transfer of
lipids produced by Bacteroides thetaiotaomicron into the mouse gut. *Cell host &*
microbe **32**, 1025-1036 e1025 (2024). https://doi.org:10.1016/j.chom.2024.04.021
41 Byrne, D. P., London, J. A., Eyers, P. A., Yates, E. A. & Cartmell, A. Mobility shift-based
electrophoresis coupled with fluorescent detection enables real-time enzyme
analysis of carbohydrate sulfatase activity. *The Biochemical journal* **478**, 735-748
(2021). https://doi.org:10.1042/BCJ20200952
42 Moggridge, S., Sorensen, P. H., Morin, G. B. & Hughes, C. S. Extending the
Compatibility of the SP3 Paramagnetic Bead Processing Approach for Proteomics. *J*
Proteome Res **17**, 1730-1740 (2018).
https://doi.org:10.1021/acs.jproteome.7b00913

43   Smits, A. H. *et al.* Biological plasticity rescues target activity in CRISPR knock outs.
Nat Methods **16**, 1087-1093 (2019). https://doi.org/10.1038/s41592-019-0614-5
44   Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I.
MSFragger: ultrafast and comprehensive peptide identification in mass
spectrometry-based proteomics. *Nat Methods* **14**, 513-520 (2017).
https://doi.org/10.1038/nmeth.4256
45   da Veiga Leprevost, F. *et al.* Philosopher: a versatile toolkit for shotgun proteomics
data analysis. *Nat Methods* **17**, 869-870 (2020). https://doi.org/10.1038/s41592-020-
0912-y
46   Franken, H. *et al.* Thermal proteome profiling for unbiased identification of direct
and indirect drug targets using multiplexed quantitative mass spectrometry. *Nat
Protoc* **10**, 1567-1593 (2015). https://doi.org/10.1038/nprot.2015.101
47   Grosdidier, A., Zoete, V. & Michielin, O. Fast docking using the CHARMM force field
with EADock DSS. *J Comput Chem* **32**, 2149-2159 (2011).
https://doi.org/10.1002/jcc.21797
48   Lebedev, A. A. *et al.* JLigand: a graphical tool for the CCP4 template-restraint library.
Acta crystallographica. Section D, Biological crystallography **68**, 431-440 (2012).
https://doi.org/10.1107/S090744491200251X
49   Webb, J. *et al.* Systems Analyses Reveal the Resilience of Escherichia coli Physiology
during Accumulation and Export of the Nonnative Organic Acid Citramalate.
mSystems **4** (2019). https://doi.org/10.1128/mSystems.00187-19
50   Koelmel, J. P. *et al.* LipidMatch: an automated workflow for rule-based lipid
identification using untargeted high-resolution tandem mass spectrometry data.
BMC Bioinformatics **18**, 331 (2017). https://doi.org/10.1186/s12859-017-1744-3
51   Kind, T. *et al.* LipidBlast in silico tandem mass spectrometry database for lipid
identification. *Nat Methods* **10**, 755-758 (2013). https://doi.org/10.1038/nmeth.2551
52   Duhrkop, K. *et al.* SIRIUS 4: a rapid tool for turning tandem mass spectra into
metabolite structure information. *Nat Methods* **16**, 299-302 (2019).
https://doi.org/10.1038/s41592-019-0344-8
53   Birchenough, G. M. H. *et al.* Muc2-dependent microbial colonization of the jejunal
mucus layer is diet sensitive and confers local resistance to enteric pathogen
infection. *Cell Rep* **42**, 112084 (2023). https://doi.org/10.1016/j.celrep.2023.112084
855

**Figure 1. General example of the conserved S1 sulfatase fold, consensus active
site sequence, and tested library of aryl- and carbohydrate sulfamates.**
a. 3-D structure of human lysosomal sulfatase GALNS (PDB:4FDJ). The N-terminal domain
is coloured red and yellow for alpha helices and beta sheets, with the C-terminal subdomain
coloured cyan and magenta for alpha helices and beta sheets. Zoom shows the consensus
sequence of GALNS with a formylglycine residue, and with the general consensus pictured
to the right. **b.** Structures of aryl- and carbohydrate sulfamate inhibitors used in this study. **c.**
Structures of phenylvinyl sulfone, phenylsulfonyl fluoride, phenylvinyl sulfonate, and
carbohydrate vinyl sulfone.
865
**Figure 2. Growth of *Bacteroides thetaiotaomicron* on varied carbon sources in the
presence of arylsulfamate inhibitors**
Bacteroides thetaiotaomicron (*B. theta*) grown on varied carbon sources; two rich carbon
sources, brain heart infusion media (BHI) and glucose (Glc); two sulfated host glycan carbon
sources, chondroitin sulfate c (CSC) and Heparin (Hep), and two plant glycan carbon
sources, potato galactan (PG) and larch arabinogalactan (LAG), in the presence of
arylsulfamates inhibitors **1-18** (see **Figure 1**). Complex glycans and arylsulfamate inhibitors

were used at a concentration of 5 mg/ml and 1 mM, respectively. All carbon sources were
utilised in a minimal media except BHI, which was purchased from Sigma and dissolved in
water. The traces are the mean of 3 independent growth experiments, and errors bars have
been omitted for clarity; individual graphs with error bars are presented in supplemental
figures **S2-S7**. Severe effects were judged qualitatively on the basis no classically
identifiable features of a bacterial growth were able to be measured: a lag and mid-
exponential growth phase leading to robust increases in OD_{600nm}. For sublethal growth
curves that displayed the aforementioned classical features, and were measurable, a two-
tailed unpaired t-test was performed to check for significance at a threshold of $p < 0.05$ (Table
S1 and S2). A difference was called significant if it met this threshold in any one of three
criteria: lag phase, growth rate, and max O.D. reached.
884
Figure 3. The effectiveness of arylsulfamate inhibitors against S1 steroid and
carbohydrate sulfatases.
a. The steroid sulfatases *Pa*Asta and *Hp*Sulf, belonging to families S1_4 (green bar) and
S1_2 (yellow bar), respectively, assayed against the panel of arylsulfamates (compounds **1-**
18 listed in **Figure 1**). **b.** Activities of family S1_11 sulfatases (BT3177[6S-GlcNAc], BT4656[6S-
GlcNAc], Amuc_1033[6S-GlcNAc], Amuc_1074[6S-GlcNAc]; blue bar above), S1_20 sulfatases
(BT1622[3S-Gal/GalNAc], BT1636[3S-Gal], Amuc_0451[3S-Gal], Amuc_0491[3S-Gal]; pink bar above), and
the S1_16 sulfatases (BT3796[4S-Gal/GalNAc] and BT1755[4S-Gal]; orange bar above) against the
panel of arylsulfamates. For both a and b the right two panels display activity with 1 mM
arylsulfamate compound in the assay relative to untreated enzyme, whilst the left two panels
are overnight preincubation of the enzyme with selected 1 mM arylsulfamates then dilution
of this preincubation mix into the reaction. Con indicates enzyme control without
arylsulfamate. Kinetic curves from which the data are derived are shown in supplemental
figures S8-S31. The blue dotted line indicates 100% activity level and the red bars indicate
control compounds.
900
Figure 4. The carbohydrate sulfamate substrate mimics are ineffective against the S1
carbohydrate sulfatase BT1636[3S-Gal].
a. Structure of the carbohydrate sulfonate/sulfamates and their aryl equivalents**. b.** Plots
show the activities of carbohydrate sulfamates, carbohydrate sulfovinyl, and arylsulfonates
against BT1636[3S-Gal]. Left plot shows activity of enzyme treated with the compounds relative
to untreated enzyme, whilst right plot show the relative activity after the enzyme had been
preincubated with the compounds overnight then diluted into the reaction mixture. **c.**
Thermostability shift assays (TSA) using inactive BT1636[3S-Gal] against the substrate 3S-Gal,
and the carbohydrate sulfamates, carbohydrate vinylsulfonate, and arylsulfonates. All
compounds in **b** were used at a concentration of 1 mM.
911
Figure 5. Arylsulfamates broadly affect growth of *Bacteroides* spp. of the HGM
Bacteroides species were grown in BHI media in the presence, or absence, of 1 mM of the
listed arylsulfamate. The data plotted are the maximal OD_{600nm} observed for arylsulfamate
treated growths relative to the untreated cells, which were each run in triplicate and
randomly paired, then averages derived. For all the estradiol growth experiments limited
material meant this shows a single experiment, except for *B. theta* which was run in
triplicate.
919
Figure 6. Global protein thermal stability reveals off target interactions of
arylsulfamate inhibitors 2 and 17.
a. Table of proteins that display an altered Tm in the presence of arylsulfamates **2** and **17**. **b.**
Proteins identified with a significant change (see methods and Dataset 1) in melting point
upon treatment with inhibitor **2** (n=2). **c.** Proteins identified with a significant change in
melting point upon treatment with inhibitor **17** (n=2). The dashed box highlights BT4322 as
the only protein common to both groups. **d.** *Bacteroides* species were grown in BHI media in
the presence, or absence, of 0.125 mM or 0.0625 mM diacylglycerol kinase inhibitor I and II,

respectively. The data plotted are the maximal OD_{600nm} observed for inhibitor treated growths
relative to the untreated controls, which were each run in triplicate and randomly paired, then
averages derived.
931
Figure 7. Recombinantly expressed BT4322 binds arylsulfamates, diacylglycerol
kinase inhibitors, and ATP
Thermostability shift assays of BT4322 against arylsulfamate and diacylglycerol kinase
inhibitors, and the kinase substrate ATP. A final protein concentration of 5 μM, in 100 mM
Bis-Tris-Propane (BTP), pH 7.0, and 150 mM NaCl, supplemented with the appropriate
ligand concentration and 5 % DMSO. A minimum of three, and a maximum of six,
independent assays were performed for each protein and protein ligand combination and
standard errors of the mean derived.
940
941
Figure 8. The effect of arylsulfamate on the growth of colonic bacteria *in vitro* and *in*
vivo.
a. a volcano plot showing the alterations of membrane lipids when *B. theta* is grown in brain
heart infusion media in the presence of 0.2 mM arylsulfamate 2. The structures of the up and
down regulated lipids are shown to the right and left of the plot, respectively. **b.** The changes
in relative abundance of colonic bacteria in the mouse gut from animals that have been
orally gavaged with the phase II arylsulfamate Irosustat; nd indicates no significant
difference as determined from a two-tailed, unpaired, t-test with a threshold of p-value of
≤0.05.

Figure 1 General example of the conserved S1 sulfatase fold, consensus active site sequence, and tested library of aryl- and carbohydrate sulfamates.

a. The 3-D structure of the human lysosomal sulfatase GALNS (PDB:4FDJ), the N-terminal domain is coloured red and yellow for alpha helices and beta sheets, with the C-terminal subdomain coloured cyan and magenta for alpha helices and beta sheets; a zoom in of the consensus sequence of GALNS with a converted formylglycine, and with the general consensus pictured to the right; **b.** Chemical structures of aryl- and carbohydrate sulfamate inhibitors used in the study. **c**. Chemical structures of phenylvinyl sulfone, phenylsulfonyl fluoride, phenylvinyl sulfonate, and carbohydrate vinyl sulfone.

Figure 2 Growth of Bacteroides thetaiotaomicron on varied carbon sources in the presence of arylsulfamate inhibitors

Bacteroides thetaiotaomicron (B. theta) grown on varied carbon sources; two rich carbon sources, brain heart infusion media (BHI) and glucose (Glc); two sulfated host glycan carbon sources, chondroitin sulfate c (CSC) and Heparin (Hep), and two plant glycan carbon sources, potato galactan (PG) and larch arabinogalactan (LAG), in the presence of arylsulfamates inhibitors **1-18** (see Figure 1). Complex glycans and arylsulfamate inhibitors were used at a concentration of 5 mg/ml and 1 mM, respectively. All carbon sources were utilised in a minimal media except BHI, which was purchased from Sigma and dissolved in water. The traces are the mean of 3 independent growth experiments, and errors bars have been omitted for clarity; individual graphs with error bars are presented in supplemental figures S2-S7. Severe effects were judged qualitatively on the basis no classically identifiable features of a bacterial growth were able to be measured: a lag and mid-exponential growth phase leading to robust increases in OD600nm. For sublethal growth curves that displayed the aforementioned classical features, and were measurable, a two-tailed unpaired t-test was performed to check for significance at a threshold of $p<0.05$ (Table S1 and S2). A difference was called significant if it met this threshold in any one of three criteria: lag phase, growth rate, and max O.D. reached.

Figure 3 The effectiveness of arylsulfamate inhibitors against S1 steroid and carbohydrate sulfatases.
a. The steroid sulfatases PaAsta and HpSulf, belonging to families S1_4 (green bar) and S1_2 (yellow bar), respectively, assayed against the panel of arylsulfamates (compounds 1-18 listed in Figure 1). **b.** Activities of family S1_11 sulfatases (BT3177 6S-GlcNAc, BT4656 6S-GlcNAc, Amuc_1033 6S-GlcNAc, Amuc_1074 6S-GlcNAc; blue bar above), S1_20 sulfatases (BT1622 3S-Gal/GalNAc, BT1636 3S-Gal, Amuc_0451 3S-Gal, Amuc_0491 3S-Gal; pink bar above), and the S1_16 sulfatases (BT3796 4S-Gal/GalNAc and BT1755 4S-Gal; orange bar above) against the panel of arylsulfamates. For both a and b the right two panels display activity with 1 mM arylsulfamate compound in the assay relative to untreated enzyme, whilst the left two panels are overnight preincubation of the enzyme with selected 1 mM arylsulfamates then dilution of this preincubation mix into the reaction. Con indicates enzyme control without arylsulfamate. Kinetic curves from which the data are derived are shown in supplemental figures S8-S31. The blue dotted line indicates 100 % activity level and the red bars indicate control compounds.

Figure 4. The effectiveness of carbohydrate sulfamate substrate mimics against the S1 carbohydrate sulfatase BT1636[3S-Gal].

a. Structure of the carbohydrate sulfonate/sulfamates and their aryl equivalents. **b.** The graphs show the effectiveness of carbohydrate sulfamates, carbohydrate sulfovinyl, and arylsulfonates against BT16363S-Gal; the left graph shows relative activity to wildtype with the compounds included in the assay, whilst the right graph show the relative activity after the enzyme had been preincubated with the compounds overnight then diluted into the reaction mixture, con indicates enzyme control rate against 3S-Gal with no added compound; **c.** Thermostability shift assays (TSA) using inactive BT1636[3S-Gal] against the substrate 3S-Gal, and the putative inhibitor structures carbohydrate sulfamates, carbohydrate sulfovinyl, and arylsulfonates. All compounds in b were used at a concentration of 1 mM.

Figure 5. Arylsulfamates broadly affect Bacteroides spp. of the HGM

Bacteroidota species were grown in BHI media in the presence, or absence, of 1 mM of the listed arylsulfamate. The data plotted are the maximal OD_{600nm} observed for arylsulfamate treated growths relative to the untreated controls which were both ran in triplicate and randomly paired then averages derived. For all the estradiol growths a single experiment was conducted except for *B. theta* which was ran in triplicate; this was due to limited material. Shown errors are standard errors of the mean.

1=*B. caccae*   2=*B. cellulolyticus*   3=*B. clarus*   4=*B. ovatus*   5=*B. intestinalis*   6=*B. salyersiae*   7=*B. oleicipnenius*   8=*B. fragilis*   9=*B. xylanisolvens*   10=*B. thetaiotaomicron*

Figure 6. Global protein thermal stability reveals off target interations of arylsulfamate inhibitors 2 and 17.
a. Table of proteins which displayed an altered Tm in the presence of arylsulfamates 2 and 17. **b.** Proteins identified
with a significance change (see methods and Dataset 1) in melting point upon inhibitor 2 treatment (n=2).
c. Proteins identified with a significance change in melting point upon inhibitor 17 treatment (n=2). A dashed box
indicates BT4322 is the only protein common across both groups. **d.** Bacteroides species were grown in BHI media
in the presence, or absence, of 0.125 mM or 0.0625 mM diacylglycerol kinase inhibitor I and II, respectively. The data
plotted are the maximal OD$_{600nm}$ observed for inhibitor treated growths relative to the untreated controls which were
both ran in triplicate and randomly paired then averages derived.

Figure 7. Recombinantly expressed BT4322 binds arylsulfamates, diacylglycerol kinase inhibitors, and ATP

Thermostability shift assays of BT4322 against arylsulfamate and diacylglycerol kinase inhibitors, and the

kinase substrate ATP. A final protein concentration of 5 µM, in 100 mM Bis-Tris-Propane (BTP), pH 7.0, and

150 mM NaCl, supplemented with the appropriate ligand concentration and 5 % DMSO. A minimum of three,

and a maximum of six, independent assays were performed for each protein and protein ligand combination

and standard errors of the mean derived.

Figure 8. The effect of arylsulfamate on the growth of colonic bacteria in vitro and in vivo.

a. a volcano plot showing the alterations of membrane lipids when B. theta is grown in brain heart infusion media in the presence of 0.2 mM arylsulfamate 2. The structures of the up and down regulated lipids are shown to the right and left of the plot, respectively. b. The changes in relative abundance of colonic bacteria in the mouse gut from animals that have been orally gavaged with the phase II arylsulfamate Irosustat; nd indicates no significant difference as determined from a two-tailed, unpaired, t-test with a threshold of p-value of ≤0.05.