The interplay between light, arsenic and H_2O_2 controls oxygenic photosynthesis in a Precambrian analog cyanobacterial mat.

Short title: Light, arsenic, and H_2O_2 govern photosynthesis in a microbial mat

Andrea Castillejos Sepúlveda[1,*], Daan R. Speth[2,3§], Carolin Kerl[4], Daniel Doherty-Weason[5,6,7], Federico A. Vignale[8], Kinga Santha[1], Wiebke Mohr[2], Arjun Chennu[9], Elisa Merz[1], Dirk de Beer[1], Maria E. Farias[10], Judith M. Klatt[1,5,6,7*].

[1] Microsensor Group, Max Planck Institute for Marine Microbiology, Bremen, Germany.

[2] Biogeochemistry Group, Max Planck Institute for Marine Microbiology, Bremen, Germany.

[3] Division of Microbial Ecology, Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria.

[4] Environmental Geochemistry, Bayreuth Center for Ecology and Environmental Research (BayCEER), University of Bayreuth, D-95440 Bayreuth, Germany

[5] Microcosm Earth Center, University of Marburg & Max Planck Institute for Terrestrial Microbiology, 35032 Marburg, Germany

[6] Center for Synthetic Microbiology (SYNMIKRO), Karl-von-Frisch-Straße 10, 35032 Marburg, Germany

[7] Biogeochemistry Group, Department of Chemistry, University of Marburg

[8] European Molecular Biology Laboratory - Hamburg Unit, Notkestrasse 85, Hamburg 22607, Germany

[9] Leibniz Centre for Tropical Marine Research, Data Science and Technology, Fahrenheitstr. 6, 28359 Bremen, Germany

[10] PUNABIO S.A. Campus USP-T Av. Solano Vera y Camino a Villa Nougués, San Pablo, Tucumán, Argentina

*Correspondence: andreacasbio@gmail.com, Judith.Klatt@mpi-marburg.mpg.de

Abstract

The regulation of O_2 production by cyanobacteria is critical to understanding the co-evolution of oxygenic photosynthesis (OP) and Earth's redox landscape. This includes their response to electron donors for competitive anoxygenic photosynthesis, like arsenic. In this work, we assessed the effect of arsenic cycling on photosynthetic activity in a modern cyanobacterial mat thriving beneath an arsenate-rich (~9-10 µM) water column in the high-altitude central Andes, using biogeochemical and omics approaches. Microsensor measurements and hyperspectral imaging revealed two O_2-producing cyanobacterial layers. During the afternoon, OP ceased in the lower, Chl f-dominated layer. Ex-situ measurements revealed that the combination of high light and arsenic induced a prominent rise in local H_2O_2 concentration, which then coincided with the interruption of OP. Mat incubations suggested that after OP ceased, cyanobacteria transitioned to As(III)-driven anoxygenic photosynthesis using far-red light. Additional incubations and metatranscriptomics on in-situ samples reveal simultaneous As(V) reduction during the day, at sufficiently high rates to supply electron donors for anoxygenic photosynthesis. This study proposes that As(III) can serve as an alternative electron donor for cyanobacterial photosynthesis. It also reveals the crucial role of arsenic in moderating OP, a consequence of the interaction between arsenic and reactive oxygen species under high irradiance. Given the widespread abundance of arsenic in the Precambrian, we further discuss how this regulatory mechanism could have played an important role in the early evolution of OP.

keywords

1. Introduction

Oxygenic photosynthesis (OP) evolved in cyanobacteria during the Archean, a period referring to 3.7 to 2.5 billion years ago, within the Precambrian eon. The general consensus is that the evolution of OP was preceded by anoxygenic photosynthesis (AP) [1], in which reduced substances (e.g. sulfide, ferrous iron, arsenite) were used as electron donors for light-driven carbon fixation. These electron donors were widely available in the mostly reducing environment of the Archean oceans [2]. Intriguingly, extreme volcanic activity resulted in high arsenic concentrations, i.e. high reduced inorganic arsenite (As(III)) [3,4]. Given its prevalence, As(III) could have been a widespread electron donor for AP during the Archean [5], and may have sustained an active redox cycle, given that bacteria can metabolize both arsenate (As(V))) or As(III) for energetic gain [6].

It is widely accepted that ancient cyanobacteria, seeking protection from both the intense UV radiation and resulting reactive oxygen species (ROS), avoided surface layers with direct strong irradiation. Cyanobacteria could have instead positioned themselves under mineral crusts or, or deeper inside stratified microbial communities, known as microbial mats [7]. This represents a fragile compromise between harvesting photosynthetic light and preventing exposure to UV radiation. Consequently, early cyanobacteria might have developed diverse strategies to harness low-energy light and exploit a diverse set of spectral niches. One intriguing strategy is known as FaR-Red Light Adaptations [8–10], which involves the utilization of chlorophyll f (Chl f) to harvest light at approximately 710 nm. The use of Chl f could potentially signify an ancient trait developed by cyanobacteria[11].

In addition to optimizing light harvesting, modern cyanobacteria exhibit metabolic flexibility, enabling them to navigate complex environments. Certain strains demonstrate the remarkable capability of engaging in both OP and sulfide-driven AP [12–14]. Interestingly, light capture and type of photosynthesis might be linked through small differences in D1, a core protein of PSII, which is essential for OP. Chl f synthesis requires the exchange of D1 (in photosystem II), for a similar protein (super-rogue D1) that functions as a Chl f synthase [15]. As a result, OP is blocked, possibly favoring

PSI-based AP through the far-red light capture via Chl f [15,16]. To date, only sulfide has been

confirmed as an electron donor for cyanobacterial AP, with well-documented and highly species-

dependent responses to sulfide exposure [17,18]. Although sulfur and arsenic metabolism seem

evolutionarily interconnected [19], our understanding of how cyanobacteria manage arsenic

exposure remains limited.

Studying modern analogues of Precambrian environments, such as the isolated lakes and wetlands

in the central Andes with their extreme conditions, offers valuable insights into potential interactions

between OP and arsenic cycling on ancient Earth [20–24]. Among those ecosystems, Laguna Pozo

Bravo, located in the Salar de Antofalla, Argentina, represents an ideal study site as environmental

pressures there include high altitude, intense UV radiation, drastic shifts in air temperature, and high

arsenic concentrations due to the volcanic origin of the Andean mountain chain [25,26]. Laguna Pozo

Bravo harbors thriving microbialites and microbial mats despite the seemingly harsh environmental

conditions [25,27]. Similar microbial mats and microbialites were abundant in submerged

environments through the Precambrian [28].

In microbial mats, functional groups of microorganisms form distinct vertically-stratified layers

according to their metabolic niches. Due to the close proximity of different microorganisms,

metabolites diffusing across layers of the mat can be efficiently recycled. However, bacteria in mats

also encounter challenges from competition for resources, as well as the quick spread of hazardous

metabolites, such as sulfide, arsenic and ROS, from neighboring layers.

In this study, we combined incubation experiments with microsensor- and omics- based approaches

to assess the interactions between arsenic and diel dynamics of OP. We hypothesized that OP is

fundamentally influenced by local arsenic speciation, which is shaped by the co-inhabiting

microorganisms, and that the regulation is predominantly mediated via ROS and competition with

AP.

2. Results

2.1. Description of Laguna Pozo Bravo and their microbial mats

Laguna Pozo Bravo is a hypersaline lake at an altitude of 3330 m [25]. During our sampling in February 2019, the water column had a pH of ~8.00 and high salinity (137.25 g L^{-1}). Irradiance reached extremely high levels at noon (up to 2950 μmol photons m^{-2} s^{-1}). Air temperature fluctuated greatly through the day (4°C at night, 30°C at noon), while water temperature remained around 20°C.

The lake sediment was mostly covered by light pink microbial mats, interspersed with patches of dark gray lithifying mat zones and microbialites (Fig 1a; Lencina et al., 2021). The microbial mats displayed a vertical stratification into four distinct visible layers: a pink upper crust rich in calcium carbonate, a green layer, a red layer, and a brown layer (Fig 1b). The mats form cohesive sheets and can be easily separated from the underlying coarse dark brown sediment. Hyperspectral imaging showed that the green layer consisted of a zone dominated by Chl a (absorption max at 690 nm) and phycocyanin (absorption max at 625 nm) indicative of cyanobacteria [29,30], and a zone dominated by BChl a (absorption max at 810 nm [31]) indicative of purple anoxygenic phototrophs [32] (Fig 1d). The red layer showed a clear absorption maximum at 715 nm, characteristic of specific BChl e in green anoxygenic bacteria like Chloroflexi [33] or of Chl f in cyanobacteria [34]. No spectral signs of photosynthetic pigments were found in the pink and brown layers.

Fig 1. Overview of microbialites and microbial mat structures in Pozo Bravo. (a) Microbialites and microbial mats (pink and lithifying gray) found in Pozo Bravo, (b) Pink microbial mat cross section, with layers indicated. (c,d) Distribution of selected pigments in the top 7 mm of Pozo Bravo microbial mat, captured by hyperspectral imaging. c) Regular view, see Supp. Fig S1 for a different view of the same sample, with brighter color banding. The microbial mat surface is indicated by a white line. (d) Second derivative of reflectance spectra around 675 nm, 715 nm, and 805 nm as indicated by different colors. Pigments with in-vivo absorption maxima at the respective wavelength are shown in parentheses.

2.1.1. Microbial mat metagenome analysis

To investigate the vertical distribution of phyla in the mat, we performed metagenomics on slices of a microbial mat. The slices were obtained by separating a piece of mat with a scalpel into the four visible layers (pink, green, red, brown, Fig 1b), directly after sampling. An assembly was then made using the metagenomic reads from all layers. After assigning the assembled contigs to bins, between 62-69% of the reads from each layer were successfully mapped back to the bins, indicating that the bins give a good overview of the microbial community in each layer (Fig 2, Supp. Table S2). Among bins with taxonomic assignments, Pseudomonadota (Alphaproteobacteria and Gammaproteobacteria) was found to be the most abundant phylum across all layers of the microbial mat, followed by Bacteroidota (Bacteroidia and Rhodothermia) and Bacillota (Halanaerobiia, Bacilli, and Clostridia). Chloroflexota (Chloroflexia and Anaerolineae) were also abundant in all layers except for the upper crust, while Deinococcota (Deinococci) were present in a relatively large abundance only in the crust. With the exception of an incomplete Lokiarchaea bin in the red and brown layers, all bins were bacterial. Two cyanobacterial bins, *Halothece sp.* and ESFC-1 (of family *Spirulinaceae*), were recovered from a co-assembly made using all samples. A new assembly was conducted only using reads from the green microbial mat layer to specifically target cyanobacteria. The same cyanobacterial bins, plus RECHO1 (of family *Elainellaceae*), were recovered. From the abundance of reads mapped to each bin (Supp. Table S3), ESFC-1 and RECHO1 were likely located in the pink and green layers, and Halothece in the green and red layers.

Although pigment absorbance (Fig 1d) indicated a relatively high abundance of cyanobacteria in the green and red layers of the microbial mat, total cyanobacterial abundance in the mat appears to be underestimated. For possible reasons behind this discrepancy see Supplement to Results 2.2. The relative abundance between the cyanobacterial taxa per mat layer, however, likely reflects the real vertical distribution under the assumption that DNA extraction efficiency is similar for all cyanobacteria through the mat.

Fig 2. Taxonomic profile of phyla from metagenomic bins in a Pozo Bravo microbial mat. The bins

were selected from a metagenomic assembly made using reads from all vertically-stratified layers

(crust, green, red, brown). Only bins with over 55% completeness and less than 10% redundancy are

shown. Relative abundance was obtained by mapping the reads from each individual layer

metagenome to the total set of bins. Results were summarised by grouping bins by phylum.

2.1.2. Arsenic redox pathways in the metagenome

We then used our metagenome data to screen for genes involved in arsenic metabolism. A great

variety of genes directly related to arsenic metabolism are present in the Pozo Bravo microbial mat,

to the abundance of housekeeping genes(Fig 3). The roles of these genes are discussed in more detail

below and shown in Fig 2b, Supp. Results 2.3, and Supp. Table S1.

Several distinct routes for microbial arsenic detoxification exist, broadly divided into three steps.

Initially, arsenic gets imported into microbial cells via specific transporters, then it is chemically

modified by specific detoxification systems, and then finally exported using dedicated efflux systems.

First, we focused on arsenic import pathways. As(V) and As(III) can enter the cell through different

mechanisms. As(III) (in the protonated form $As(OH)_3$) can enter the cell by diffusion across the

membrane or through aquaglyceroporin channels (*aqps*), the genes for which were found in the

metagenome [35,36]. As(V) can be imported through phosphate channels [37]. Almost all

components (*pstSCAB*) of the high-affinity phosphate import channel were found. The pst system is

also highly selective for phosphate because the slightly larger size of As(V) affects crucial bonds in the

transporter, thus allowig the discrimination of phosphate over arsenate by 500- to 4500-fold [38].

Thus, the use of pst indicates an adaptation to limit As(V) intake (see Supp. Results 2.3) [37–39].

Once inside the cell, arsenic can be metabolized for energy generation or detoxified through a series

of redox reactions. The genes found in the Pozo Bravo metagenome were summarized into three

putative detoxification-related pathways (Fig 3): The first detoxification pathway is based on an

As(V)-transferring gap3/gadph, and arsJ, which were both found in the metagenome. The second

pathway involves reduction of As(V) to As(III) via arsC1, arsC2 and mrx1. The genes encoding for

arsC1 and arsC2 were? not recovered from the metagenome. Yet, their presence is required for

functionality of mrx1 (Fig 3, for the full Mrx1 cycle see Ordóñez et al. 2009 [40]. While this pathway

has thus far exclusively been assigned a detoxification function, the third pathway involving As(V)

reduction mediated by the product of the genes *arsRCDAB* and *acr3*, has been hypothesized to be

used in As(V) respiration [41]. In all other studies of genomes and metagenomes in the Puna region,

an environment similar to Pozo Bravo, *acr3* was the predominant As(V) reduction gene [23,42,43].

No other respiratory *arrA* or *arrB* As(V) reductases were found. As(III) oxidation is mediated by the

products of *aioA* and *aioB*, coupled to the reduction of either cytochrome c or azurin, as part of

processes such as AP [44]. Both *aioA* and *aioB* were found in reads from all layers of the

metagenome. Other arsenite oxidation genes (*arxA*) were not found in the metagenome. Overall, the

metagenomic analysis revealed a significant abundance of arsenic-related genes in all layers of the

Pozo Bravo microbial mat, thus enabling bacteria to thrive under these harsh conditions.

Fig 3. **Pathways of arsenic metabolism present ain Pozo Bravo microbial mats.** (a) Presence of genes

associated with arsenic cycling and housekeeping genes (DNA gyrase (*gyrA*, *gyrB*)) in the

metagenome of Pozo Bravo microbial mat. Shades of gray represent the number of gene copies

(normalized as Transcripts Per Million, TPM [45]) from each microbial mat layer that were mapped to

the metagenomic assembly, which was made using reads from all microbial mat layers. Almost as

many gene copies are dedicated to arsenic cycling as for housekeeping tasks, suggesting intense

metabolizing of arsenic species. (b) Overview on the pathways of redox arsenic transformation

detected in Pozo Bravo mats, based on previously reported functions of arsenic genes [6,40,46,47].

Genes were broadly classified in expressing transporters for arsenic uptake (blue) and export (red),

enzymes involved in reduction of As(V) (orange), oxidation of As(III) (green), and production and

export of organic arsenicals (pink). ArsC1 and ArsC2 are shown in gray since their presence could not

be determined in the metagenome). Details and a list of abbreviations can be found in Table S1.

2.2.   In-situ dynamics of oxygenic photosynthesis and dissolved arsenic speciation

To assess the dynamics of photosynthetic O_2 production under the hostile and dynamic conditions of

Pozo Bravo, we performed in-situ microsensor depth profiling. This revealed that OP occurred in two

distinct layers within the mat. An upper O_2 concentration peak was observed at approximately 1 mm

depth (Fig 4a & d, Supp. Fig S2) in the green Chl *a*-dominated layer (Fig 1d). An additional net OP

maximum was observed at approximately 2-3 mm depth (Fig 4a & d, Supp. Figs. S3-S4), coinciding with the depth of the red microbial mat layer. The production of O_2 in this layer strongly suggests that the dominant pigment with the absorption maximum of 715 nm is cyanobacterial and hence Chl *f*, instead of BChl e.

The rates of net OP in both layers exhibited unusual dynamics over a diel cycle. In the top cyanobacterial layer, net OP decreased at high light intensities after noon but recovered again in the late afternoon (Fig 4a & d, Supp. Figs S3-S4). This pattern was more pronounced in the deeper Chl *f*-cyanobacterial layer, where OP was completely interrupted at around 13:00 and only resumed around 18:00 in one of the two replicate spots (Fig 4a, Supp. Fig S2).

To assess co-occurring changes in the concentration of electron donors for AP, H_2S dynamics were measured using microsensors, and the speciation dynamics of iron and arsenic were determined through porewater analysis. The insubstantial dynamics of both iron and H_2S suggests that neither drive AP (see Suppl. Results, Supp. Figs. S5). However, arsenic speciation in the porewater underwent strong changes over the diel cycle. Although the depth resolution obtained was low, porewater measurements clearly showed the greatest changes in redox cycling of dissolved arsenic within the top centimeter of the microbial mat, i.e. in the photic zone (Fig 4b & c). Here, up to 98% of dissolved arsenic was oxidized during daylight hours (07:00-18:00), and reduced at 19:00 and in the predawn hours (00:30-06:00) (Supp. Fig S6). Measurements could not be obtained between approximately 19:00 – 00:30 due to strong winds, but we assume that dissolved arsenic were reduced during this time period. Thus, the interruption of OP in the deep cyanobacterial layer coincided with a maximum in As(V) and minimum in As(III) concentration.

Fig 4. Interruption of cyanobacterial photosynthesis coincides with strong redox dynamics in dissolved arsenic. Depth profiles of dissolved oxygen (a), and speciation of inorganic arsenic over a diel cycle in porewater from Pozo Bravo (b&c, n=1-3 per depth). The microbial mat surface is at 0 cm depth (horizontal white line in (a-c), black dotted line in (d)). Profiles selected from various points in time are shown in (d). The timepoints used for d) are indicated by rectangles in (a-c), following the numbering in (b). Profile 4 represents a single point in time (00:30). Colored background represents the estimated placement of the green and red microbial mat layers (Fig 1b). The chosen profiles exemplify the oxygen production during the morning (d1) and late afternoon (d3), with a period of interruption in between (d2). D4)shows a comparison of the dissolved arsenic speciation at night (00:30). The interruption in oxygen production coincides with a maximum in dissolved As(V). Net volumetric rates calculated based on the concentration depth profiles of oxygen measured in two replicate spots can be found in Supp. Fig S3, and depth integrated rates over a diel cycle in Supp. Fig S4.

2.3.   A combined effect of light and arsenic on H_2O_2 and oxygenic photosynthesis

To test if the interruption of OP at noon was related to arsenic or exclusively to light, depth profiles of O_2 were measured ex-situ in microbial mats before and after As(V) addition at increasing light intensities, from low to high light (~500 to ~1000 µmol photons s^{-1} m^{-2}) (Supp. Fig S7). Arsenic was added the evening prior to the measurements to allow for diffusion into the mat and reestablishment of internal redox cycling. The results revealed a combined effect of both arsenic and light intensity on OP. In the microbial mat without arsenic, local O_2 concentration increased until reaching a steady state under both high and low light conditions, as expected from normal cyanobacterial OP. However, O_2 production ceased after approximately 30 min in microbial mats treated with As(V) (to 10 µM final concentration) when exposed to high light. O_2 production resumed after lowering light intensity, showing that OP was hindered only in the presence of both arsenic and high light. Although most mat pieces did not exhibit the clearly separated double peaks observed in-situ, occasionally two net productivity maxima were observed. The inhibitory effect of light and arsenic occurred more rapidly and extensively in the deeper maximum. OP was also interrupted in As(V)-supplemented mats that were instantly exposed to high light and not to a stepwise increase. Thus, the interruption of OP appeared to be independent of exposure time to light and time of day, and exclusively driven by a combined effect of high light and arsenic.

To further elucidate the mechanism behind the inhibition of OP, we simultaneously measured H_2O_2

concentration in depth profiles (Fig 5). In the absence of arsenic, no net production of H_2O_2 was

observed across light intensities. In As(V)-treated mats, H_2O_2 concentration within the

photosynthetically active zone dramatically increased at 1000 µmol photons m^{-2} s^{-1}. Upon the

cessation of OP, H_2O_2 concentrations decreased to values comparable to dark conditions. Notably, a

transient increase of H_2O_2 at the oxic interface was observed in live mats treated with both arsenic

and DCMU, an inhibitor of OP (Fig 5f). Although the increase in H_2O_2 was much smaller when OP was

inhibited, the increase was nevertheless surprising as OP was expected to be the only source of H_2O_2.

Moreover, measurements following the addition of approximately 200 µM H_2O_2 to the water

column, demonstrated that OP remained unaffected and H_2O_2 was entirely consumed within the first

1-2 mm (Supp. Fig S8), as also observed in other mat systems [48]. In summary, these results suggest

three key findings: (1) H_2O_2 production is related to a light-driven mechanism in the presence of

dissolved arsenic. Due to the possibility of arsenic cycling within the mat, the arsenic species

triggering ROS production and OP interruption remains unclear. (2) The resultant local H_2O_2

production rate must be enormous, overwhelming the substantial sink strength. (3) Given that H_2O_2

production occurs in the presence of DCMU (01:00 in Fig 5f), but not in the phase of OP

discontinuation (15:00-17:00 in Fig 5f), the mat community likely transitions to an activity that

tempers ROS abundance when no OP occurs.

Fig 5. As(V) inhibits OP at high light and induces H_2O_2 evolution. O_2 and H_2O_2 measurements of

microbial mats before (a-c) and after As(V) addition (10 µM final concentration) (d-f)). Bars over each

graph represent light conditions over time, as either darkness, low light (500 µmol photons m^{-2} s^{-1}),

or high light (1000 µmol photons m^{-2} s^{-1}). Contour plots show concentration of O_2 (a,d) or H_2O_2 (c,e)

across the depth of the microbial mat over time. The solid white horizontal line represents the mat

surface at depth 0. The vertical dashed line in d-f indicates time of DCMU addition, an inhibitor of

oxygenic photosynthesis. (c) and (f) Depth-integrated rates of production (DIR) for O_2 (n=3 pieces of

mat) and H_2O_2 (n=2 pieces of mat).

2.4. Transition from oxygenic photosynthesis to light-driven As(III) oxidation in cyanobacteria using far-red light

To assess the interaction between dissolved arsenic speciation and photosynthesis, microbial mat pieces were incubated, with and without an OP inhibitor (DCMU), over a diel cycle using various light qualities targeting different phototrophs. This showed that As(V) reduction occurred under most treatments in the dark (Fig 6, Supp. Figs. S9-S12). Second, we found AP under all light conditions, including those specifically targeting cyanobacteria (Fig 6).

Incubations were initiated by the addition of As(V) or As(III) to enable differentiation between oxidation and reduction rates. Total dissolved arsenic remained relatively constant in all incubations, suggesting that sorption/desorption dynamics played a minor role. This also allowed us to calculate rates of both As(III) oxidation and As(V) reduction based only on As(V) dynamics, which was favorable due to the higher accuracy of the employed colorimetric method for As(V) [49]. The incubations showed a strong dependence of arsenic speciation on light (Fig 6, Supp. Figs. S9-S12). Most notably, As(V) was consistently reduced in the dark, indicating respiration with As(V) as the electron acceptor. Upon exposure to all types of light, rates of reduction consistently dropped (Supp. Figs. S9-S12). Full spectrum white light and 715 nm light even triggered net As(V) production, i.e. net As(III) oxidation, albeit with a delay under 715 nm light (Fig 6a & e, Supp. Figs. S9 & S10).

Given that As(V) respiration and photosynthetic As(III) oxidation likely co-occur, we conducted a detailed analysis of As(V) consumption (i.e., respiration) rates to estimate photosynthetic gross rates from net rates. Rates of As(V) consumption in the dark were linearly dependent on As(V) concentration across the complete measured range (Fig 4b). Assuming that As(V) respiration rates were unaffected by light and solely determined by the As(V) concentration, we were able to calculate light-driven gross As(III) oxidation as

$$G\_As(III)_{oxidation} = N\_As(V)_{prod} + As(V)_{resp}([As(V)]) \tag{1}$$

where $G\_As(III)_{ox}$ is the gross rate of As(III) oxidation (i.e. gross As(V) production, Fig 6b), $N\_As(V)_{prod}$ is the net rate of As(V) production, (i.e. net As(III) oxidation), and $As(V)_{resp}([As(V)])$ is the rate of light-

independent As(V) consumption as a function of concentration derived from the linear regression in

Fig 6d.

These considerations revealed light-driven gross As(III) oxidation under all light treatments. The rates

were highest in full-spectrum white light in the absence of DCMU, i.e. under highly oxic conditions

favoring aerobic oxidation of As(III) (Fig 6b). Rates under these white light conditions were

substantially lower in the presence of DCMU but not absent, suggesting that As(III) was also truly

photosynthetically oxidized in the absence of O_2 derived from OP. At 810 nm, purple bacteria

oxidized As(III) by anoxygenic photosynthesis as previously observed in various cultures [5,44] and in

mats [24]. Intriguingly, at 690 nm and at 715 nm, which specifically target cyanobacterial Chl a and f,

respectively, we also found evidence for As(III)-driven AP (Fig 6b). Most strikingly, gross As(III)

oxidation occurred in the presence of DCMU, strongly suggesting that As(III)-driven photosynthesis

indeed occurs in the cyanobacteria from Pozo Bravo.

The regulation of As(III)-driven photosynthesis differed between the 690 nm and 715 nm treatments.

In the 690 nm treatments without inhibitor, O_2 production was only triggered after As(III) depletion

indicating that the cyanobacteria transitioned from As(III)-driven anoxygenic photosynthesis to OP

upon depletion of the alternative electron donor (Supp. Fig S9, Fig 6c). For incubations under 715 nm,

the diel patterns in photosynthesis dynamics were similar to those observed in porewater extracted

from mats (Fig 6a,c,e, Supp. Fig S10): Net OP occurred within the first hours of light exposure (phase

1) and then ceased (phase 2) (Fig 6e). For treatments with and without DCMU, As(V) release was

absent in phase 1 (first hours of light exposure), but increased drastically in the afternoon (phase 2)

(Fig 6e). Due to the lack of oxygen in treatments with DCMU, the phases for this treatment were

estimated by the change in As(V) release partway through the experiment, rather than by net OP.

Only the 715 nm treatment exhibited these two clear phases. The similarity in gross As(V) production

with and without DCMU at 715 nm strongly suggests that As(III) is oxidized by cyanobacteria using

Chl f via anoxygenic photosynthesis after the interruption of OP.

Fig 6. Rates of arsenic cycling dependent on light quality and photosynthesis. (a) Net rates of As(V) production (Net_As(V)$_{prod}$) in incubations of mat slurries with and without DCMU, under white light, light at λmax = 690 nm to target Chl *a*, λmax = 715 nm to target Chl *f,* and light at λmax =810 nm to target BChl *a*. The gray bars overlaid in 715 nm represent phase 1 and phase 2 of As(V) production, as shown in (e). (b) Rate of As(V) consumption, i.e. As(V) respiration, in the dark dependent on the concentration of As(V). (c) Gross rates of As(III) oxidation (G_As(III)$_{oxidation}$) calculated from (a) by adding the expected rate of As(V) removal (As(V)$_{resp}$([As(V)])) from (b) according to Eq (1). (d) Net rates of oxygenic photosynthesis in the same incubations as in (a) and (c). In (e) an example for a time series of dissolved As(V) and O_2 in batch incubations is shown, based on which the net rates of arsenic production in (a) and net OP in (d) were calculated. The slurry was initially exposed to ~20 µM As(V) and, after a dark phase, exposed to light of λmax = 715 nm. Errors represent the standard deviation of concentration in separate vials (n=3). In phase 1, which represents the initial hours of exposure to light, net O_2 production occurred and arsenite was not photosynthetically oxidized. In phase 2, after sustained exposure to light, O_2 production ceased and As(III) was photosynthetically oxidized to As(V). A similar pattern was observed in the treatment with DCMU, which blocks OP. Namely, As(V) was reduced for an initial five hours (phase 1), after which As(III) was photosynthetically oxidized to As(V) (phase 2). Additional time series at different light qualities and with different initial arsenic conditions are in Supp. Figs. S9-S12.

2.5. Metatranscriptomic evidence for observed activity transitions

2.5.1. Concurrent arsenic oxidation and reduction

We used metatranscriptomes from Pozo Bravo to determine diel dynamics in transcription of genes related to arsenic metabolism. Specifically, we investigated photosynthetic gene transcription in the afternoon, and whether transcription of genes related to arsenite oxidation increased during the same period. For this purpose, we mapped transcriptomic reads from in-situ mat samples obtained over a diel cycle to the metagenomic bins (Fig 3a,7). This analysis generally supported our process rate observations of concomitant oxidation and reduction reactions and revealed mechanistic insights. Unfortunately arsenic cycling by cyanobacteria could not be assessed, except for *arsR*, and As(III) efflux encoders *arsA, arsB,* and *acr3;* and importers *pstSCAB* in the Halothece and ESFC-1 bins. The genes linked to arsenite oxidation in cyanobacteria may be currently undescribed, or highly divergent from sequences in databases. Also, few metatranscriptomic reads mapped to the cyanobacterial bins in general.

The highest read numbers mapped to arsenic genes were present in Alpha- and Gamma-
proteobacterial bins (Fig 7a), which stimulated in-depth analysis of diel transcription by these groups.
We found that As(III) oxidation was likely mainly driven by *aioAB* transcribed by taxa among the
Rhodobacterales and Chromatiales, prominent representatives of purple photosynthetic bacteria.
Specifically, *aioA* encoding for the large subunit of arsenite oxidase, was transcribed at all timepoints
in Rhodobacterales (Alphaproteobacteria) (Fig 7). The highest *aioA* transcription, and thus potentially
As(III) oxidation, happened during the day (9 am - 6 pm) although most of the arsenic present in
porewater was As(V) (Fig 7a). Transcription of *aioB*, which encodes for the small subunit of the same
As(III) oxidase, was also found during 9 am to 3 pm, in line with high *aioAB* activity during this portion
of the day. During daylight hours, Rhodobacterales expressing *aioAB* also transcribed *pufL* and *pufM*
(Supp. Fig S14), which encode for components of a type II photosynthetic reaction center, thus
supporting arsenite-driven anoxygenic photosynthesis. Similar patterns of *aioAB* and *pufLM*
transcription were observed in Rhizobiales, mainly during the morning and late afternoon (Supp. Fig
S14, Fig 7b).

Gammaproteobacteria were separated into three main orders: Chromatiales, Pseudomonadales and
Xanthomonadales (Fig 7b). Transcription of *pufLM* was found in both Pseudomonadales and
Xanthomonadales during the day as well as the night. However, unlike in Alphaproteobacteria,
transcription of *aioA* primarily took place during the night, except for Chromatiales that showed
transcription at 3 pm. Transcription of *aioB* was not found. This could suggest non-photosynthetic
As(III) oxidation at night, i.e.- coupled to nitrate reduction [50,51], in analogy to
chemolithoautotrophic sulfide oxidation by purple sulfur bacteria commonly observed in other mat
ecosystems [52].

The potential As(III)-oxidizing purple anoxygenic phototrophs among the Alphaproteobacteria
appeared to transition to As(V) reduction at low light and in the night. Namely, in Rhodobacterales,
the transcription of 1-arseno-3-phosphoglycerate via As(V)-transferring GAPDH and *arsJ* peaked at
night, and As(V) reduction and As(III) excretion via *arsC* and *arsB*, occurred in the morning. Most

Gammaproteobacteria did not show such pattern. Instead, *arsC*, *acr3* and *arsB* were continuously

expressed. Chromatiales, however, co-expressed *aioA* and genes involved in excretion of intracellular

As(III) (*arsA*, *arsB*) during the day. This general pattern of alternating key players in As(V) reduction

held true across the complete metatranscriptome. Most prominently, the relative contribution of

bacterial classes driving arsenate reduction via *arsC* changed per timepoint (Fig 7a). Transcription of

arsC by Gammaproteobacteria was high and relatively constant, while *arsC* transcription at 9 am was

highest by Alphaproteobacteria, and at noon by Bacteroidota and Chloroflexota. Spirochaetia was

the only known bacterial class to express *arsC* at night (1 am), but some transcription was also found

in unidentified bins. Transcription in unbinned reads was negligible. While the key players may

alternate over a diel cycle, intriguingly, the general *arsC* transcription was highest during daylight

hours (Fig 7a), consistent with pronounced cryptic redox cycling of arsenic.

Fig 7. Transcription of arsenic metabolism genes over a diel cycle in Pozo Bravo microbial mats. (a)
Sum of *aioA*, *arsC*, and *grxC* transcription, in TPM, over the metatranscriptome of all layers of Pozo
Bravo microbial mat. *aioA* is involved in arsenite oxidation, *arsC* is involved in arsenate reduction,
and *grxC* may be part of arsenate reduction or reactive oxygen species response pathways. (b)
Transcription of genes related to arsenic cycling in selected taxa, shown as the sum of TPM per gene,
normalized across all timepoints (i.e., per column). Normalization was performed due to the varying
number of gene copies found per gene. Higher gene transcription is represented in red, lower
transcription in blue. Transcription of *pufL* and *pufM*, which encode for photosynthetic reaction
center II, can be found in Supp. Fig S14. An overview of transcription of genes in cyanobacterial bins
can be found in Supp. Fig S15.

2.5.2. Elevated general stress and ROS response prior to the interruption of OP

To confirm the involvement of ROS in the interruption of OP, we focused on the transcription of

genes related to ROS response and stress. The transcription of genes involved in ROS repair peaked

at 9:00 and 12:30 (Fig 7b), with few exceptions in the early morning and late afternoon (see Supp.

Results 2.4). The transcription of ROS genes through the entire day may indicate that microbial mat

inhabitants plausibly have to deal with continuous ROS stress, despite no net production.

Only a few reads from the metatranscriptome sample at 12:30 mapped to cyanobacterial genes,

among which the stress proteins PspA and Hsp70 in RECH01 were of interest (Supp. Fig S15). *pspA* is

a component of a phage shock protein involved in lipid membrane repair, while Hsp70 is a highly

conserved heat shock protein found under both normal and stressful conditions in cyanobacteria. As

suggested in previous reports [53,54], it is likely that in this case, *pspA* transcription is a response to

membrane disturbance rather than an attack from phages. This is due to ROS production, from the

intense light, potentially disrupting the membrane and highlighting the stress that cyanobacteria face

at noon when OP is also interrupted.

420 *2.5.3. Changes in cyanobacterial photosystems*

Cyanobacterial bins showed clear differences in the gene transcription of the two photosystems over

a diel cycle. Most prominently, the majority of PSI components were expressed in the morning, while

those of PSII were expressed in the evening. The former could be consistent with the re-instatement

of photosystems after a period of darkness. The latter suggests fundamental changes, like the repair

or replacement of central photosystem components, when the cyanobacteria transition back to OP.

Interestingly, the transcription of *psbA*, a central component of PSII, was highest shortly before the

reinstatement of OP was observed (Supp. Fig S16). Diverse isoforms of D1, the protein encoded by

psbA, are used by cyanobacteria based on changing environmental conditions, such as irradiance and

O_2 concentration [55]. While it is known that D1 requires constant replacement due to ROS damage

by the neighboring oxygen-evolving complex during OP, it is possible that AP also leads to radical

formation in PSII when excitation energy cannot yield charge separation [56]. Thus, D1 replacement

via *psbA* would enable cyanobacteria performing AP to replace D1 due to the accumulating ROS

damage from AP, and/or express a different isoform of D1 better adapted to OP.

434 3. Discussion

As(III)-driven anoxygenic photosynthesis by bacteria has only been characterized in a small number

of organisms, mostly in Mono Lake, an arsenic-containing environment [5,44,57]. In the

cyanobacteria-dominated and arsenic-rich environment of Laguna Pozo Bravo, it appears to be the

dominant form of AP despite the low local concentration of As(III) in the mat. Omics and process

rates consistently suggest that AP was driven by a constant supply of As(III) from As(V) reduction in a

highly active, cryptic redox cycle of arsenic. The photosynthetic As(III) oxidation was performed by

various bacterial groups throughout the day. As expected, obligate anoxygenic phototrophs (purple

bacteria) utilized arsenic for photosynthesis (Fig 3&7). Surprisingly, As(III) appears to also be used by

cyanobacteria for AP. The regulation of the transition between OP and AP in the cyanobacteria is

complex, with ROS likely playing a key role.

In situ, the cyanobacteria of Pozo Bravo likely produce ROS during OP, as evidenced by the

transcription of oxidative stress proteins like thioredoxins and peroxiredoxin-5 (AHP1) (Supp. Fig

S13). Even though we observed no net H_2O_2 production, gross rates are likely very high, supported by

the large sink capacity of the mat revealed by the addition of H_2O_2 (Supp. Fig S8). Further ROS

formation from increasing light radiation through the day could be aggravated by 1) abiotic reactions

with iron and UV light [58], 2) singlet oxygen production during photosystem repair [59,60], and 3)

ROS produced due to arsenic oxidation, as proposed under physiological conditions by del Razo

(2001) [61]:

$H_3AsO_3 + 2H_2O + O_2 \rightarrow H_3AsO_4 + H_2O_2$ ($\Delta G° = -40.82$ kcal/mol)   (2)

At noon, during maximum illumination, the sum of H_2O_2 production from OP and plausibly from

abiotic As(III) oxidation (Eq. 2) exceeds sink strength and net production of H_2O_2 becomes

measurable.

It is well documented that ROS can lead to the inhibition or complete interruption of OP to avoid

further ROS formation [62]. In Laguna Pozo Bravo, OP by cyanobacteria in the bottom layer, which

contains Chl-f (red in Fig 1), ceases after H_2O_2 is produced. OP is then replaced by oxidation of As(III)

under light targeting these Chl f cyanobacteria – an intriguing strategy to circumvent two problems

simultaneously: intracellular ROS production linked to OP and extracellular ROS production linked to

the presence of As(III) (Eq. 2). Cyanobacterial As(III)-driven AP is supported by a line of indirect

evidence. Most prominently, As(III) was oxidized under exposure to wavelengths exclusively targeting

oxygenic phototrophs. Also, DCMU did not inhibit this light-driven As(III) oxidation. Thus, oxidation

with photosynthetically produced O_2 can be excluded. The lack of inhibition by DCMU provides

additional information about the mechanisms behind As(III) oxidation. Specifically, it suggests that

electrons derived from As(III) oxidation enter the photosynthetic electron transport chain

downstream of PSII (Supp. Fig S17). This is because DCMU inhibits electron transfer at the acceptor

side of PSII. The specific enzymes involved in As(III)-driven AP in cyanobacteria, however, remain

unclear from the metagenomic and -transcriptomic data.

Neither As(III) concentration nor the light available to PSI regulate the transitions between

photosynthetic modes in the Chl f cyanobacteria. Instead, the shift in cyanobacterial photosynthetic

mode occurs exclusively under conditions of high light and in the presence of arsenic, resulting in an

accumulation of local ROS concentrations, resembling a stress response. Notably, As(III)-driven AP in

the Chl a cyanobacteria is regulated fundamentally different, i.e., it is not linked to the abundance of

H_2O_2. OP in incubations targeting Chl a only began after As(III) was depleted, indicating a regulation

by electron donor availability in Chl a cyanobacteria, analogous to observations in cyanobacteria

performing sulfide-driven photosynthesis [14]. To confidently test this hypothesis, however, it would

also be necessary to show photosynthetic As(III) oxidation in cyanobacterial isolates, rather than in

microbial mats.

The transitions from OP to AP could only be measured by supplementation with surplus arsenic. In

situ, however, almost no As(III) was measurable at noon. Yet, the interruption of OP and in situ gene

transcription of As(V) reductases (*ars*C) and As(III) oxidases (*aioA, aioB*) indicate that As(III) oxidation

was likely still occurring – in an active, yet cryptic, redox cycle. The source of As(III) for AP can only be

the reduction of the highly abundant As(V). Accordingly, *arsC* transcription was high during daylight

hours in the metatranscriptome (Fig 7a), and process rate measurements clearly showed that rates

were in the range of AP rates. Further, As(V) reduction rates were linearly dependent on

concentration within the measured range (up to ~19 µM) and did not yet saturate according to

Michaelis-Menten kinetics (Fig 4d). Such first order process suggests low-affinity enzymes, which are

most efficient during the day at high As(V) levels.

To summarize, we propose that during the morning (dawn-10 am), both layers of cyanobacteria

perform OP, while some Alphaproteobacteria (e.g. Rhodobacterales) begin with As(III)-AP (Fig 8).

Through the mat, other Alphaproteobacteria, as well as Gammaproteobacteria, begin the reduction

of intracellular As(V), resulting in As(V) reduction throughout the entire mat. ROS damage response

appears to take place starting in the morning. During the late morning (10 am-12 pm), both layers of

cyanobacteria continue doing OP even though As(III) is still present in the porewater. Then ROS

accumulation is aggravated by increasing sunlight intensity combined with the presence of As(III).

This could be due to spontaneous oxidation (eq. 2) , and/or ROS production during As(III)-affected

OP. In the early afternoon (12 pm -3 pm), the increased ROS damage triggers cyanobacteria in the

deeper layer to stop OP and to transition to As(III)-driven AP. In the late afternoon (3pm to sunset),

less ROS is produced in the top layer due to lower OP, and the lower Chl f cyanobacteria switch back

to OP. It remains unclear why the upper cyanobacteria do not follow the same photosynthetic switch

as the lower zone, but instead keep performing OP during the afternoon, albeit at diminished rates

(green mat layer in Fig 1a). A plausible explanation could be the lower As(III) supply from the deeper

layers.

Fig 8. Proposed metabolic activities driving arsenic cycling through a diel cycle within Pozo Bravo
microbial mats. These are based on pooling insights from hyperspectral imaging, process rate
measurements and metatranscriptomic analyses. Reactive oxygen species (ROS) production during
the morning, afternoon and late afternoon is indicated with grey clouds.

These dynamics have implications for the regulation of OP throughout the Precambrian. Due to the

generally reducing environment in the Precambrian, and frequent input of arsenic from volcanic

activity, As(III) was abundant in ancient oceans and was likely to be an electron donor for ancient AP

[3,63,64]. Although some evidence pointing toward AP in *Oscillatoria* sp. arose in studies of microbial

mats from Mono Lake, California [5], As(III)-driven AP has not been conclusively reported in

cyanobacteria. This study yields valuable experimental data as another step in a long line of evidence

for the hidden metabolic capabilities of cyanobacteria [19,65,66]. While the exact metabolic

mechanisms behind the interruption of OP and transition to As(III)-driven AP are yet to be

understood, here we clearly show that arsenic has a fundamental effect on the diel O_2 budget of

cyanobacterial mats by tempering O_2 export. Additionally, the cryptic cycling of arsenic in Laguna

Pozo Bravo may showcase the hidden, yet quick and important, nature of arsenic metabolisms.

4. Methods

4.1. Field measurements and sampling

Field measurements and sampling were conducted in Laguna Pozo Bravo (25° 30' 58" S, 67° 34' 42"

W), in the Salar de Antofalla, Argentina in late February 2019 and January 2023 at an elevation of

3330 m. To assess diel dynamics of oxygen and H_2S within the mat, we continuously measured depth

profiles of concentration on two consecutive days, in two spots of the mat. The microsensors (H_2S

and Clark-type O_2 with <50 μM tip width) were constructed, calibrated and used as previously

described [67–70]. Vertical profiles were measured simultaneously by attaching the microsensors to

a multi-sensor holder, yielding a distance between tips of ~0.5 cm. Electricity was provided by a

portable generator. Using O_2 microsensor measurements, we calculated the net rates of OP as the

difference in the fluxes into and out of the two photosynthetically active zones.

In parallel with the microsensor measurements, subsamples of the mat were regularly taken for

porewater extraction and meta-omics. Cores for porewater extraction (2.5 cm inner diameter) were

sampled in triplicate every three hours from 1 am until 6 pm. Porewater was immediately extracted

from the cores (2.5 cm ID, up to 6 cm depth) at 1 cm intervals using rhizones (Rhizosphere Research

Products, NL). An aliquot of 500 μL porewater was instantly frozen with liquid nitrogen and stored

and transported in a dry shipper until arsenic speciation analysis using an ICP-MS as previously

described [49] (Supp. Methods 1.1).

Additional sediment cores for meta-omics (2.5 cm inner diameter, up to 4 cm depth) were sampled

in triplicates every three hours. The microbial mat was rapidly and carefully cut into four pieces per

core that were instantly frozen in liquid nitrogen for metagenomic and metatranscriptomic analyses.

A few additional cores were frozen intact for hyperspectral imaging (Supp. Methods 1.2). Further, 10

x 5 cm pieces of mat with ~0.5 cm underlying sediment were stored in plastic containers for

transport and maintained in mesocosms at ~700 μmol photons m^{-2} s^{-1} underneath a water column

with a salinity of ~137 g L^{-1} (TropicMarine artificial saltwater), constantly circulated to maintain a

temperature of 21°C, until ex-situ measurements.

Temperature of air and water above the mat were recorded using HOBO loggers (Onset Computer

Corp., MA). Measurements of light intensity failed due to rapid oversaturation of the sensor signal in

the morning. Occasionally, pH measurements in the water column were taken using a Mettler Toledo

pH meter. Salinity was measured with an Atago refractometer.

551   4.2.   Assessment of diel H_2O_2 dynamics

To assess the effect of light and arsenic on net O_2 production and H_2O_2 dynamics in the mat, we

subsampled microbial mat pieces (~ 2.5×2.5 cm) from the laboratory mesocosms and continuously

measured O_2 and H_2O_2 depth profiles during simulated diel light cycles, in the absence and presence

of dissolved arsenic. Mat pieces were cut and fixed with low-melting agarose (1%) to the bottom of

glass dishes. Circular flow was established with the same hypersaline water (TropicMarine artificial

saltwater approximately 137 g L^{-1}) used for regular maintenance of the mats, and adjusted using a

peristaltic pump (Minipuls 3 Abimed, Gilson). Diel cycles were simulated using halogen light sources

(KL 2500 and KL 1500, Schott). Microsensors (H_2O_2, and O_2 with 80-120 μM tip diameter) were

constructed, calibrated and used as previously described [67–70]. Concentration-depth profiles were

performed under three light conditions: dark, 500 and 1000 μmol photons m^{-2} s^{-1}. First, profiles were

measured without supplementing As(V), assuming that most arsenic had been washed out after four

months of lab maintenance. Then As(V) was added to the overlying water column to a final

concentration approximately 10 μM, mimicking the environmental As(V) maximum. After As(V)

addition, mats were left undisturbed overnight to enable the diffusion of arsenic into the mat, and

diel profiling began in the morning. Diel profiling was also performed in As(V)-supplemented mats

after addition of DCMU, an inhibitor of OP, and after autoclaving the mat as an abiotic control.

Depth-integrated rates of net O_2 and H_2O_2 production were calculated from the concentration depth

profiles by considering Fick's second law of diffusion. First, local volumetric net rates were calculated

as the second derivative of concentration multiplied by the diffusion coefficient corrected for

temperature and salinity. Volumetric depth rates were then integrated over the uppermost 5 mm of

the mat.

4.3.   Assessment of arsenic cycling rates in batch incubations

To assess the dependency of light quality on arsenic cycling, small subsamples of microbial mats were

incubated under As(V), As(III), and LED lights targeting distinct groups of phototrophs. Hypersaline

water (TropicMarine artificial saltwater ~137 g L^{-1}) was deoxygenated by bubbling with helium,

vigorously shaking, then removing gas bubbles, repeated five times. Subsamples cut from microbial

mats (~200 mg) using 8 mm width cores (surface area 0.5 cm^2) were inserted into 3 mL gastight glass

vials with a septum cap (exetainers, Labco) which were later filled completely with deoxygenated

hypersaline saltwater. Then they were placed into containers on shakers, with constant flow of

temperature-regulated water. The incubations were subjected to 8-hour cycles of light with various

different spectral quality, followed by 8 hours of darkness. Controls were placed under constant

darkness. All light treatments were repeated with the addition of 5 µL DCMU solution (5 mM) to

inhibit oxygenic photosynthesis.

The light treatments were chosen at specific wavelengths to target different groups of phototrophs.

LEDs with an emission maximum (λ_{max}) at 690 nm (hexagonal high-power LEDs from Roithner

Lasertechnik, Vienna; Supp. Fig S18) were used to specifically target Chl *a*, i.e., the light harvesting

pigment of most cyanobacteria for photosystem I and the central pigment in the reaction center of

both photosystems. Since an unexpected absorption maximum was found at 715 nm in the lower

cyanobacterial layer, we also used LEDs in the far-red range of the spectrum with λ_{max} = 715 nm to

target cyanobacterial Chl *f*, which can likely act as antenna in both photosystems [11,71].

Additionally, we used λ_{max} = 810 to target BChl *a* of purple anoxygenic bacteria, and full-spectrum

white light to target all phototrophs.

Every three hours, three incubation vials per treatment were removed for measurements. First,

oxygen in each exetainer was measured with a microsensor, then 2 mL of the overlying water were

filtered. From the filtered water, 1 mL was fixed with 4 mL 2% zinc acetate for total dissolved sulfide

measurements, and 1 mL was fixed with 15.53 μL fixing solution and flash frozen for arsenic

measurements. Fixing solution for arsenic measurements was made by mixing 27 mL of concentrated

HNO_3 diluted 1:2 with ultrapure water and 100 mL fresh DTPA solution. The DTPA solution was made

by mixing 6.75 mL ultrapure water, 6.75 mL concentrated HNO_3 and 50 mL of 40% premade

DTPA*Na$_5$ solution (Sigma Aldrich, CAS 140-01-2). Sulfide was later measured as previously described

[72], and arsenic was measured colorimetrically based on Castillejos Sepúlveda et al. (2022), with

modifications to calculations due to the high salinity and absence of phosphate (Supp. Methods 1.2).

4.4. Metagenomics

A piece of microbial mat collected in situ at 12:30 pm, following the procedure described above, was

sliced into the four distinct layers (crust, green, red, brown) and sent to the Max Planck Genome

Center (MPGC, Cologne, Germany) for processing. DNA extraction was performed using the DNeasy

PowerSoil Pro Kit (Qiagen). An Illumina-compatible library was prepared with NEB Ultra II DNA kit

(NEB) and each library was shotgun sequenced with Illumina HiSeq3000 (2 x 150 bp, paired-end read

mode). The crust and brown layers were sequenced at a depth of 6.8 million reads, while the green

and red layers were sequenced at a depth of 34 million reads to target cyanobacteria, which are

notoriously difficult to sequence.

Raw sequences were quality-trimmed with Trimmomatic v.0.38[73], and assembled with MEGAHIT

v.1.2.9[74]. Initially, a co-assembly of all samples was used, but due to low cyanobacterial read

depth, a second assembly was prepared using only reads from the green layer of the microbial mat.

Both assemblies were later processed using the same workflow, but the second assembly was only

used for analysis of cyanobacteria.

Samples were mapped onto the assembly with Bowtie2 v.2.4.5 [75]. Gene calling was performed

with Prodigal v.2.6.3 [76], and the resulting gene calls were annotated in anvio v.7.1 [77] using the

KEGG KOfam v.95.0 [78] and COG v.2020 [79] databases with DIAMOND v.0.9.14 [80] search.

Interproscan v.78 [81,82] was used for further annotations with the following databases: CATH-

Gene3D v.4.2.0 [83], CDD v.3.17 [84], HAMAP v.2020_01 [85], PANTHER v.14.1 [86], Pfam v.32.0 [87],

PIRSF v.3.02 [88], PRINTS v.42.0 [89], PROSITE Patterns and PROSITE Profiles v.2019_11 [90], SMART

v.7.1 [91], SFLD v.4 [92], SUPERFAMILY v.1.75 [93] and TIGRFAMs v.15.0 [94]. Genes related to

arsenic, photosynthesis, glutaredoxins, and thioredoxins were manually selected and annotated

based on consensus from the automatically-generated annotations, using BLASTx v.2.2.31 [95] to

resolve discrepancies.

Metagenomic analyses were initially focused on genes that could result in redox changes, before

binning was performed to assign genes to distinct bacterial taxa.

Automatic binning was done with MetaBAT2 v.2.15 [96] on the large co-assembly and MaxBin2 2.2.7

[97] on the second assembly. In both cases, bins were manually refined in anvio. Taxonomy was

assigned to bins using GTDB-TK v.95.0 [98,99] within anvio. The relative abundance of each bin was

calculated using coverM v. 0.4.0 [100] and plotted using ggplot2 [101] in R v. 4.0.3 [102].

4.5.   Metatranscriptomics

Microbial mat samples collected following the procedure described above in situ at 1:00 am, 9:40

am, 12:30 pm, 3:15 pm, and 6:30 pm were sent to Max Planck Genome Center (MPGC, Cologne,

Germany) for processing. RNA was isolated using Quick-DNA/RNA Miniprep Plus Kit (Zymo Research).

Next, an Illumina-compatible library was prepared with the kit Universal Prokaryotic RNA-Seq

incl. Prokaryotic AnyDeplete for rRNA depletion (Tecan Genomics). Each sample was shotgun-

sequenced with Illumina HiSeq3000 (2x150 bp, paired-end read mode) at depths ranging from 35.5

million to 51 million reads. Raw sequences were quality trimmed with Trimmomatic, then mapped

onto the metagenomic assemblies with BBmap v38.73 [103], and used featureCounts from Rsubread

v. 1.22.2 to obtain counts of each gene per metatranscriptomic sample. To compare gene

transcription across samples, counts of each gene call were normalized by calculating Transcripts Per

Million (TPM) as previously described [45]. The TPM of all gene calls annotated with the same

function (i.e.- the same gene name) were summed, and used to make heatmaps with Pheatmap

v.1.0.12 [104] in R.

Data availability

649 Data has been submitted to the EMBL Nucleotide Sequence Database (ENA). Metagenomic and

650 metatranscriptomic raw reads, as well as cyanobacterial metagenome-assembled genomes described

651 in Supp. Figure S15 and Supp. Table S4, can be found under project number PRJEB41764.

652 Metagenome-assembled genomes from the co-assembly (Figs 2,3,7, Supp. Fig S13-14 and Supp.

653 Table S3) can be found under project number PRJEB73824.

Acknowledgements

We would like to thank Secretaría de Medio Ambiente de la Provincia de Catamarca for providing the

research permit (Expte.EX-2022-02222431-CAT-DPB#SEAS), and Secretaría de Política Ambiental en

Recursos Naturales, Ministerio de Ambiente y Desarrollo Sostenible de la Nación Argentina for

providing the certificate of compliance (Title: IF-2023-65966642-APN-SPARN#MAD, UId: ABSCH-IRCC-

AR-264943-1) and the export certificate for genetic resources (CE-2023-69835755-APN-

SPARN#MAD). We would like to express our gratitude to Luis Ahumada for his assistance during the

field campaigns, and the native communities of Antofalla, El Peñón, and Antofagasta de la Sierra for

their valuable support. The authors also thank the laboratory technicians in the Microsensor working

group, the mechanical workshop in MPI Bremen, David Benito Merino, Marit van Erk, Britta Planer-

Friedrich, Martina Alisch, and Boran Kartal for their help with labwork, advice, and laboratory

equipment.

References

1. Nisbet EG, Fowler CMR. The Early History of Life. Treatise on Geochemistry. Elsevier; 2003. pp.
1–39. doi:10.1016/B0-08-043751-6/08166-4

2. Sleep NH. The Hadean-Archaean Environment. Cold Spring Harb Perspect Biol. 2010;2:
a002527–a002527. doi:10.1101/cshperspect.a002527

3. Oremland RS, Saltikov CW, Wolfe-Simon F, Stolz JF. Arsenic in the evolution of earth and
extraterrestrial ecosystems. Geomicrobiol J. 2009;26: 522–536.
doi:10.1080/01490450903102525

4. Smedley P., Kinniburgh D. A review of the source, behaviour and distribution of arsenic in
natural waters. Appl Geochemistry. 2002;17: 517–568. doi:10.1016/S0883-2927(02)00018-5

5. Kulp TR, Hoeft SE, Asao M, Madigan MT, Hollibaugh JT, Fisher JC, et al. Arsenic(III) Fuels
Anoxygenic Photosynthesis in Hot Spring Biofilms from Mono Lake, California. Science (80-).
2008;321: 967–970. doi:10.1126/science.1160799

6. Slyemi D, Bonnefoy V. How prokaryotes deal with arsenic. Environ Microbiol Rep. 2012;4:
571–586. doi:10.1111/j.1758-2229.2011.00300.x

7. Castenholz RW. Phototrophic Bacteria Under UV Stress. In: Seckbach J, editor. Origins.
Dordrecht: Kluwer Academic Publishers; 2004. pp. 445–461. doi:10.1007/1-4020-2522-X_27

8. Averina S, Velichko N, Senatskaya E, Pinevich A. Far-red light photoadaptations in aquatic
cyanobacteria. Hydrobiologia. Springer International Publishing; 2018. pp. 1–17.
doi:10.1007/s10750-018-3519-x

9. Gan F, Bryant DA. Adaptive and acclimative responses of cyanobacteria to far-red light.
Environmental Microbiology. John Wiley & Sons, Ltd; 2015. pp. 3450–3465.
doi:10.1111/1462-2920.12992

10. Gan F, Shen G, Bryant DA. Occurrence of far-red light photoacclimation (FaRLiP) in diverse
cyanobacteria. Life. 2015;5: 4–24. doi:10.3390/life5010004

11. Gisriel CJ, Shen G, Ho MY, Kurashov V, Flesher DA, Wang J, et al. Structure of a monomeric
photosystem II core complex from a cyanobacterium acclimated to far-red light reveals the
functions of chlorophylls d and f. J Biol Chem. 2022;298: 101424.
doi:10.1016/J.JBC.2021.101424/ATTACHMENT/E0177B24-B0ED-4877-814E-
AD8491135B45/MMC2.ZIP

12. Padan E. Facultative Anoxygenic Photosynthesis in Cyanobacteria. Annu Rev Plant Physiol.
1979;30: 27–40. doi:10.1146/annurev.pp.30.060179.000331

13. Grim SL, Dick GJ. Photosynthetic Versatility in the Genome of Geitlerinema sp. PCC 9228
(Formerly Oscillatoria limnetica 'Solar Lake'), a Model Anoxygenic Photosynthetic
Cyanobacterium. Front Microbiol. 2016;7. doi:10.3389/fmicb.2016.01546

14. Klatt JM, Al-Najjar MAA, Yilmaz P, Lavik G, de Beer D, Polerecky L. Anoxygenic Photosynthesis
Controls Oxygenic Photosynthesis in a Cyanobacterium from a Sulfidic Spring. Kostka JE,
editor. Appl Environ Microbiol. 2015;81: 2025–2031. doi:10.1128/AEM.03579-14

15. Trinugroho JP, Bečková M, Shao S, Yu J, Zhao Z, Murray JW, et al. Chlorophyll f synthesis by a
super-rogue photosystem II complex. Nature Plants. Nature Publishing Group; 2020. pp. 238–
244. doi:10.1038/s41477-020-0616-4

16. Cardona T, Murray JW, Rutherford AW. Origin and Evolution of Water Oxidation before the

Last Common Ancestor of the Cyanobacteria. Mol Biol Evol. 2015;32: 1310–1328.
doi:10.1093/molbev/msv024

17.   Cohen Y, Jørgensen BB, Revsbech NP, Poplawski R. Adaptation to Hydrogen Sulfide of
Oxygenic and Anoxygenic Photosynthesis among Cyanobacteria. Appl Environ Microbiol.
1986;51: 398–407. doi:10.1128/aem.51.2.398-407.1986

18.   Kelly DJ, Dow CS. Microbial Photosynthesis. Wang Q, editor. Compr Biotech, The Princip, Appl
and Regal of Biotechnol in Ind, Agric and Med. Singapore: Springer Singapore; 2020.
doi:10.1007/978-981-15-3110-1

19.   Nagy CI, Vass I, Rákhely G, Vass IZ, Tóth A, Duzs A, et al. Coregulated genes link
sulfide:quinone oxidoreductase and arsenic metabolism in Synechocystis sp. strain PCC6803. J
Bacteriol. 2014;196: 3430–40. doi:10.1128/JB.01864-14

20.   Rascovan N, Maldonado J, Vazquez MP, Eugenia Farías M. Metagenomic study of red biofilms
from Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea. ISME J. 2016;10:
299–309. doi:10.1038/ismej.2015.109

21.   Sancho-Tomás M, Somogyi A, Medjoubi K, Bergamaschi A, Visscher PT, van Driessche AES, et
al. Geochemical evidence for arsenic cycling in living microbialites of a High Altitude Andean
Lake (Laguna Diamante, Argentina). Chem Geol. 2020;549: 119681.
doi:10.1016/j.chemgeo.2020.119681

22.   Sancho-Tomás M, Somogyi A, Medjoubi K, Bergamaschi A, Visscher PT, Van Driessche AES, et
al. Distribution, redox state and (bio)geochemical implications of arsenic in present day
microbialites of Laguna Brava, Salar de Atacama. Chem Geol. 2018;490: 13–21.
doi:10.1016/j.chemgeo.2018.04.029

23.   Kurth D, Amadio A, Ordoñez OF, Albarracín VH, Gärtner W, Farías ME. Arsenic metabolism in
high altitude modern stromatolites revealed by metagenomic analysis. Sci Rep. 2017;7: 1024.
doi:10.1038/s41598-017-00896-0

24.   Visscher PT, Gallagher KL, Bouton A, Farias ME, Kurth D, Sancho-Tomás M, et al. Modern
arsenotrophic microbial mats provide an analogue for life in the anoxic Archean. Commun
Earth Environ. 2020;1: 1–10. doi:10.1038/s43247-020-00025-2

25.   Lencina AI, Soria MN, Gomez FJ, Gérard E, Farias ME. Composite microbialites: Thrombolite,
dendrolite, and stromatolite associations in a modern environment, Pozo Bravo lake, Salar de
Antofalla, Catamarca Puna, Argentina. J Sediment Res. 2021;91: 1305–1330.
doi:10.2110/jsr.2020.166

26.   Vignale FA, Lencina AI, Stepanenko TM, Soria MN, Saona LA, Kurth D, et al. Lithifying and Non-
Lithifying Microbial Ecosystems in the Wetlands and Salt Flats of the Central Andes. Microb
Ecol. 2022;83: 1–17. doi:10.1007/s00248-021-01725-8

27.   Farías ME, Villafañe PG, Lencina AI. Integral prospection of Andean microbial ecosystem
project. Microb Ecosyst Cent Andes Extrem Environ biofilms, Microb mats, microbialites
endoevaporites. 2020; 245–260.

28.   Nisbet EG, Fowler CMR. Archaean metabolic evolution of microbial mats. Proc R Soc B Biol Sci.
1999;266: 2375–2382. doi:10.1098/rspb.1999.0934

29.   Yacobi YZ, Köhler J, Leunert F, Gitelson A. Phycocyanin-specific absorption coefficient:
Eliminating the effect of chlorophylls absorption. Limnol Oceanogr Methods. 2015;13: 157–
168. doi:10.1002/lom3.10015

30.   Jørgensen BB, Cohen Y, des Marais DJ, Jorgensen BB, Cohen Y, des Marais DJ. Photosynthetic

action spectra and adaptation to spectral light distribution in a benthic cyanobacterial mat.
Appl Environ Microbiol. 1987;53: 879–886. doi:10.4319/lo.1992.37.8.1813

31. Stomp M, Huisman J, Stal LJ, Matthijs HCP. Colorful niches of phototrophic microorganisms
shaped by vibrations of the water molecule. ISME J. 2007;1: 271–282.
doi:10.1038/ismej.2007.59

32. Miller LC, Martin DS, Liu L-N, Canniffe DP, Miller LC, Canniffe · D P, et al. Composition,
Organisation and Function of Purple Photosynthetic Machinery. Microb Photosynth. 2020;
73–114. doi:10.1007/978-981-15-3110-1_4

33. Blankenship RE, Olson JM, Miller M. Antenna Complexes from Green Photosynthetic Bacteria.
Anoxygenic Photosynthetic Bacteria. Dordrecht: Kluwer Academic Publishers; 1995. pp. 399–
435. doi:10.1007/0-306-47954-0_20

34. Sawicki A, Chen M, Sawicki A, Chen · M. Molecular Mechanism of Photosynthesis Driven by
Red-Shifted Chlorophylls. Microb Photosynth. 2020; 3–42. doi:10.1007/978-981-15-3110-1_1

35. Rosen BP. Biochemistry of arsenic detoxification. FEBS Lett. 2002;529: 86–92.
doi:http://dx.doi.org/10.1016/S0014-5793(02)03186-1

36. Yang HC, Fu HL, Lin YF, Rosen BP. Pathways of Arsenic Uptake and Efflux. Current Topics in
Membranes. Academic Press Inc.; 2012. pp. 325–358. doi:10.1016/B978-0-12-394390-
3.00012-4

37. Willsky GR, Malamy MH. Effect of arsenate on inorganic phosphate transport in Escherichia
coli. J Bacteriol. 1980;144: 366–374. doi:10.1128/jb.144.1.366-374.1980

38. Elias M, Wellner A, Goldin-Azulay K, Chabriere E, Vorholt JA, Erb TJ, et al. The molecular basis
of phosphate discrimination in arsenate-rich environments. Nature. 2012;491: 134–137.
doi:10.1038/nature11517

39. Saona LA, Soria M, Durán-Toro V, Wörmer L, Milucka J, Castro-Nallar E, et al. Phosphate-
Arsenic Interactions in Halophilic Microorganisms of the Microbial Mat from Laguna
Tebenquiche: from the Microenvironment to the Genomes. Microb Ecol. 2021;81: 941–953.
doi:10.1007/s00248-020-01673-9

40. Ordóñez E, Van Belle K, Roos G, De Galan S, Letek M, Gil JA, et al. Arsenate reductase,
mycothiol, and mycoredoxin concert thiol/disulfide exchange. J Biol Chem. 2009;284: 15107–
15116. Available: https://pubmed.ncbi.nlm.nih.gov/19286650/

41. Jiang Z, Shen X, Shi B, Cui M, Wang Y, Li P. Arsenic Mobilization and Transformation by
Ammonium-Generating Bacteria Isolated from High Arsenic Groundwater in Hetao Plain,
China. Int J Environ Res Public Health. 2022;19: 9606. doi:10.3390/ijerph19159606

42. Rascovan N, Maldonado J, Vazquez MP, Eugenia Farías M. Metagenomic study of red biofilms
from Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea. ISME J. 2016;10:
299–309. doi:10.1038/ismej.2015.109

43. Ordoñez OF, Lanzarotti E, Kurth D, Cortez N, Farías ME, Turjanski AG. Genome comparison of
two Exiguobacterium strains from high altitude andean lakes with different arsenic resistance:
identification and 3D modeling of the Acr3 efflux pump. Front Environ Sci. 2015;3.
doi:10.3389/fenvs.2015.00050

44. Hoeft McCann S, Boren A, Hernandez-Maldonado J, Stoneburner B, Saltikov C, Stolz J, et al.
Arsenite as an Electron Donor for Anoxygenic Photosynthesis: Description of Three Strains of
Ectothiorhodospira from Mono Lake, California and Big Soda Lake, Nevada. Life. 2016;7: 1.
doi:10.3390/life7010001

45. Wagner GP, Kin K, Lynch VJ. Measurement of mRNA abundance using RNA-seq data: RPKM
measure is inconsistent among samples. Theory Biosci. 2012;131: 281–285.
doi:10.1007/s12064-012-0162-3

46. Páez-Espino D, Tamames J, De Lorenzo V, Cánovas D. Microbial responses to environmental
arsenic. BioMetals. 2009;22: 117–130. doi:10.1007/s10534-008-9195-y

47. Silver S, Phung LT. Genes and enzymes involved in bacterial oxidation and reduction of
inorganic arsenic. Appl Environ Microbiol. 2005;71: 599–608. doi:10.1128/AEM.71.2.599-
608.2005

48. de Beer D, Meyer V, Klatt J, Li T. Photosynthesis under very high oxygen concentrations in
dense microbial mats and biofilms. bioRxiv. 2018; 335299. doi:10.1101/335299

49. Castillejos Sepúlveda A, Gatti LM, Kerl CF, Chennu A, Klatt JM. Arsenic speciation analysis in
porewater by a novel colorimetric assay. Sci Total Environ. 2022;827: 154155.
doi:10.1016/j.scitotenv.2022.154155

50. Oremland RS, Hoeft SE, Santini JM, Bano N, Hollibaugh RA, Hollibaugh JT. Anaerobic oxidation
of arsenite in Mono Lake water and by a facultative, arsenite-oxidizing chemoautotroph,
strain MLHE-1. Appl Environ Microbiol. 2002;68: 4795–802. doi:10.1128/AEM.68.10.4795

51. Zhang J, Zhou W, Liu B, He J, Shen Q, Zhao FJ. Anaerobic arsenite oxidation by an autotrophic
arsenite-oxidizing bacterium from an arsenic-contaminated paddy soil. Environ Sci Technol.
2015;49: 5956–5964. doi:10.1021/es506097c

52. Franks J, Stolz JF. Flat laminated microbial mat communities. Earth-Science Rev. 2009;96:
163–172. doi:10.1016/J.EARSCIREV.2008.10.004

53. Flores-Kim J, Darwin AJ. The Phage Shock Protein Response. Annu Rev Microbiol. 2016;70: 83–
101. doi:10.1146/ANNUREV-MICRO-102215-095359

54. Junglas B, Huber ST, Heidler T, Schlösser L, Mann D, Hennig R, et al. PspA adopts an ESCRT-III-
like fold and remodels bacterial membranes. Cell. 2021;184: 3674-3688.e18.
doi:10.1016/J.CELL.2021.05.042

55. Sheridan KJ, Duncan EJ, Eaton-Rye JJ, Summerfield TC. The diversity and distribution of D1
proteins in cyanobacteria. Photosynth Res. 2020;145: 111–128. doi:10.1007/s11120-020-
00762-7

56. Lubitz W, Lendzian F, Bittl R. Radicals, radical pairs and triplet states in photosynthesis. Acc
Chem Res. 2002;35: 313–320.
doi:10.1021/AR000084G/ASSET/IMAGES/LARGE/AR000084GF00007.JPEG

57. Budinoff CR, Hollibaugh JT. Arsenite-dependent photoautotrophy by an Ectothiorhodospira-
dominated consortium. ISME J. 2008;2: 340–343. doi:10.1038/ismej.2007.115

58. Blokhina O, Virolainen E, Fagerstedt K V. Antioxidants, oxidative damage and oxygen
deprivation stress: A review. Annals of Botany. 2003. pp. 179–194. doi:10.1093/aob/mcf118

59. Hakala-Yatkin M, Sarvikas P, Paturi P, Mäntysaari M, Mattila H, Tyystjärvi T, et al. Magnetic
field protects plants against high light by slowing down production of singlet oxygen. Physiol
Plant. 2011;142: 26–34. doi:10.1111/j.1399-3054.2011.01453.x

60. Marais A, Sinayskiy I, Petruccione F, Van Grondelle R. A quantum protective mechanism in
photosynthesis. Sci Reports 2015 51. 2015;5: 1–8. doi:10.1038/srep08720

61. Del Razo LM, Quintanilla-Vega B, Brambila-Colombres E, Calderón-Aranda ES, Manno M,
Albores A. Stress proteins induced by arsenic. Toxicol Appl Pharmacol. 2001;177: 132–148.

doi:10.1006/TAAP.2001.9291

62.   Latifi A, Ruiz M, Zhang C-C. Oxidative stress in cyanobacteria. FEMS Microbiol Rev. 2009;33:
258–278. doi:10.1111/j.1574-6976.2008.00134.x

63.   Jackson CR, Langner HW, Donahoe-Christiansen J, Inskeep WP, McDermott TR. Molecular
analysis of microbial community structure in an arsenite-oxidizing acidic thermal spring.
Environ Microbiol. 2001;3: 532–542. doi:10.1046/j.1462-2920.2001.00221.x

64.   Lebrun E, Brugna M, Baymann F, Muller D, Lièvremont D, Lett M-C, et al. Arsenite oxidase, an
ancient bioenergetic enzyme. Mol Biol Evol. 2003;20: 686–93. doi:10.1093/molbev/msg071

65.   Grim SL, Dick GJ. Photosynthetic versatility in the genome of Geitlerinema sp. PCC 9228
(Formerly Oscillatoria limnetica 'solar lake'), a model anoxygenic photosynthetic
cyanobacterium. Front Microbiol. 2016;7: 1–14. doi:10.3389/fmicb.2016.01546

66.   Sforna MC, Philippot P, Somogyi A, Van Zuilen MA, Medjoubi K, Schoepp-Cothenet B, et al.
Evidence for arsenic metabolism and cycling by microorganisms 2.7 billion years ago. Nat
Geosci. 2014;7: 811–815. doi:10.1038/ngeo2276

67.   De Beer D, Schramm A, Santegoeds CM, Kühl M, Kuhl M. A nitrite microsensor for profiling
environmental biofilms. Appl Environ Microbiol. 1997;63: 973–7. Available:
http://www.ncbi.nlm.nih.gov/pubmed/16535560

68.   Jeroschewski P, Steuckart C, Kühl M. An Amperometric Microsensor for the Determination of
H 2 S in Aquatic Environments. Anal Chem. 1996;68: 4351–4357. doi:10.1021/ac960091b

69.   Klatt JM, Meyer S, Häusler S, Macalady JL, De Beer D, Polerecky L. Structure and function of
natural sulphide-oxidizing microbial mats under dynamic input of light and chemical energy.
ISME J. 2016;10: 921–33. doi:10.1038/ismej.2015.167

70.   Revsbech NP. An oxygen microsensor with a guard cathode. Limnology and Oceanography.
John Wiley & Sons, Ltd; 1989. pp. 474–478. doi:10.4319/lo.1989.34.2.0474

71.   Tros M, Mascoli V, Shen G, Ho MY, Bersanini L, Gisriel CJ, et al. Breaking the Red Limit:
Efficient Trapping of Long-Wavelength Excitations in Chlorophyll-f-Containing Photosystem I.
Chem. 2021;7: 155–173. doi:10.1016/J.CHEMPR.2020.10.024

72.   Cline JD. Spectrophotometric determination of hydrogren sulfide in natural waters. Limnol
Oceanogr. 1969;14: 454–458. doi:10.4319/lo.1969.14.3.0454

73.   Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30: 2114–2120. doi:10.1093/bioinformatics/btu170

74.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for
large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.
2015;31: 1674–1676. doi:10.1093/bioinformatics/btv033

75.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:
357–359. doi:10.1038/nmeth.1923

76.   Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: Prokaryotic gene
recognition and translation initiation site identification. BMC Bioinformatics. 2010;11: 1–11.
doi:10.1186/1471-2105-11-119

77.   Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: An advanced
analysis and visualization platformfor 'omics data. PeerJ. 2015;2015: e1319.
doi:10.7717/PEERJ.1319/SUPP-5

78. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al. KofamKOALA:
KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Valencia A,
editor. Bioinformatics. 2020;36: 2251–2252. doi:10.1093/bioinformatics/btz859

79. Tatusov RL, Galperin MY, Natale DA, Koonin E V. The COG database: a tool for genome-scale
analysis of protein functions and evolution. Nucleic Acids Res. 2000;28: 33.
doi:10.1093/NAR/28.1.33

80. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat
Methods. 2015;12: 59–60. doi:10.1038/nmeth.3176

81. Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The InterPro
protein families and domains database: 20 years on. Nucleic Acids Res. 2021;49: D344–D354.
doi:10.1093/nar/gkaa977

82. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, et al. InterProScan 5: Genome-scale
protein   function   classification.   Bioinformatics.   2014;30:   1236–1240.
doi:10.1093/bioinformatics/btu031

83. Sillitoe I, Dawson N, Lewis TE, Das S, Lees JG, Ashford P, et al. CATH: Expanding the horizons of
structure-based functional annotations for genome sequences. Nucleic Acids Res. 2019;47:
D280–D284. doi:10.1093/nar/gky1097

84. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE: the
conserved domain database in 2020. - PubMed - NCBI. Nucleic Acids Res. 2020;48: D265–
D268. Available: https://academic.oup.com/nar/article-abstract/48/D1/D265/5645006

85. Pedruzzi I, Rivoire C, Auchincloss AH, Coudert E, Keller G, De Castro E, et al. HAMAP in 2015:
Updates to the protein family classification and annotation system. Nucleic Acids Res.
2015;43: D1064–D1070. doi:10.1093/nar/gku1002

86. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: More genomes, a
new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res.
2019;47: D419–D426. doi:10.1093/nar/gky1038

87. Bateman A, Birney E, Durbin R, Eddy SR, Howe KL, Sonnhammer ELL. The Pfam protein
families database. Nucleic Acids Research. 2000. pp. 263–266. doi:10.1093/nar/28.1.263

88. Nikolskaya AN, Arighi CN, Huang H, Barker WC, Wu CH. PIRSF Family Classification System for
Protein Functional and Evolutionary Analysis. Evol Bioinforma. 2006;2: 117693430600200.
doi:10.1177/117693430600200033

89. Attwood TK, Coletta A, Muirhead G, Pavlopoulou A, Philippou PB, Popov I, et al. The PRINTS
database: A fine-grained protein sequence annotation and analysis resource-its status in
2012. Database. 2012;2012. doi:10.1093/database/bas019

90. Sigrist CCJA, Castro E De, … LC-N acids, 2012 undefined, de Castro E, Cerutti L, et al. New and
continuing developments at PROSITE. 2012;41: D344–D347. doi:10.1093/nar/gks1067

91. Letunic I, Bork P. 20 years of the SMART protein domain annotation resource. Nucleic Acids
Res. 2018;46: D493–D496. doi:10.1093/nar/gkx922

92. Brown S, Pegg S, Ojha S, Huang C, Ferrin TE, Babbitt PC. The Structure-Function Linkage
Database. FASEB J. 2006;20. doi:10.1096/fasebj.20.5.a899-c

93. Pandurangan AP, Stahlhacke J, Oates ME, Smithers B, Gough J. The SUPERFAMILY 2.0
database: A significant proteome update and a new webserver. Nucleic Acids Res. 2019;47:
D490–D494. doi:10.1093/nar/gky1130

94. Haft DH, Selengut JD, Richter RA, Harkins D, Basu MK, Beck E. TIGRFAMs and genome
properties in 2013. Nucleic Acids Res. 2013;41. doi:10.1093/nar/gks1234

95. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
Architecture and applications. BMC Bioinformatics. 2009;10: 1–9. doi:10.1186/1471-2105-10-
421/FIGURES/4

96. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: An adaptive binning
algorithm for robust and efficient genome reconstruction from metagenome assemblies.
PeerJ. 2019;2019: e7359. doi:10.7717/PEERJ.7359/SUPP-3

97. Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover
genomes from multiple metagenomic datasets. Bioinformatics. 2016;32: 605–607.
doi:10.1093/bioinformatics/btv638

98. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A
standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of
life. Nat Biotechnol. 2018;36: 996–1004. doi:10.1038/nbt.4229

99. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with
the   Genome   Taxonomy   Database.   Bioinformatics.   2020;36:   1925–1927.
doi:10.1093/BIOINFORMATICS/BTZ848

100. Woodcroft BJ, Newell R. wwood/CoverM: Read coverage calculator for metagenomics. Centre
for Microbiome Research, School of Biomedical Sciences, Faculty of Health, Queensland
University   of   Technology;   2021   [cited   22   Jul   2022].   Available:
https://github.com/wwood/CoverM#readme

101. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Cham: Springer-Verlag New York;
2016. doi:10.1007/978-3-319-24277-4

102. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria; 2020
[cited 3 May 2023]. Available: https://www.r-project.org/

103. Bushnell B. BBMap short-read aligner, and other bioinformatics tools. 2015.

104. Kolde R. CRAN - Package pheatmap. In: https://cran.r-project.org/package=pheatmap
[Internet].   2019   [cited   12   Jun   2022].   Available: https://cran.r-
project.org/web/packages/pheatmap/index.html

1.   Nisbet EG, Fowler CMR. The Early History of Life. Treatise on Geochemistry. Elsevier; 2003. pp.

1–39. doi:10.1016/B0-08-043751-6/08166-4

2.   Sleep NH. The Hadean-Archaean Environment. Cold Spring Harb Perspect Biol. 2010;2:

a002527–a002527. doi:10.1101/cshperspect.a002527

3.   Oremland RS, Saltikov CW, Wolfe-Simon F, Stolz JF. Arsenic in the evolution of earth and

extraterrestrial   ecosystems.   Geomicrobiol   J.   2009;26:   522–536.

doi:10.1080/01490450903102525

4.   Smedley P., Kinniburgh D. A review of the source, behaviour and distribution of arsenic in

natural waters. Appl Geochemistry. 2002;17: 517–568. doi:10.1016/S0883-2927(02)00018-5

5.   Kulp TR, Hoeft SE, Asao M, Madigan MT, Hollibaugh JT, Fisher JC, et al. Arsenic(III) Fuels

Anoxygenic Photosynthesis in Hot Spring Biofilms from Mono Lake, California. Science (80-).

2008;321: 967–970. doi:10.1126/science.1160799

6.   Slyemi D, Bonnefoy V. How prokaryotes deal with arsenic. Environ Microbiol Rep. 2012;4:

571–586. doi:10.1111/j.1758-2229.2011.00300.x

7.   Castenholz RW. Phototrophic Bacteria Under UV Stress. In: Seckbach J, editor. Origins.

Dordrecht: Kluwer Academic Publishers; 2004. pp. 445–461. doi:10.1007/1-4020-2522-X_27

8.   Averina S, Velichko N, Senatskaya E, Pinevich A. Far-red light photoadaptations in aquatic

cyanobacteria. Hydrobiologia. Springer International Publishing; 2018. pp. 1–17.

doi:10.1007/s10750-018-3519-x

9.   Gan F, Bryant DA. Adaptive and acclimative responses of cyanobacteria to far-red light.

Environmental Microbiology. John Wiley & Sons, Ltd; 2015. pp. 3450–3465.

doi:10.1111/1462-2920.12992

10.   Gan F, Shen G, Bryant DA. Occurrence of far-red light photoacclimation (FaRLiP) in diverse

cyanobacteria. Life. 2015;5: 4–24. doi:10.3390/life5010004

11.   Gisriel CJ, Shen G, Ho MY, Kurashov V, Flesher DA, Wang J, et al. Structure of a monomeric

photosystem II core complex from a cyanobacterium acclimated to far-red light reveals the

functions of chlorophylls d and f. J Biol Chem. 2022;298: 101424.

doi:10.1016/J.JBC.2021.101424/ATTACHMENT/E0177B24-B0ED-4877-814E-

AD8491135B45/MMC2.ZIP

12.   Padan E. Facultative Anoxygenic Photosynthesis in Cyanobacteria. Annu Rev Plant Physiol.

1979;30: 27–40. doi:10.1146/annurev.pp.30.060179.000331

13.   Grim SL, Dick GJ. Photosynthetic Versatility in the Genome of Geitlerinema sp. PCC 9228

(Formerly Oscillatoria limnetica 'Solar Lake'), a Model Anoxygenic Photosynthetic

Cyanobacterium. Front Microbiol. 2016;7. doi:10.3389/fmicb.2016.01546

14.   Klatt JM, Al-Najjar MAA, Yilmaz P, Lavik G, de Beer D, Polerecky L. Anoxygenic Photosynthesis

Controls Oxygenic Photosynthesis in a Cyanobacterium from a Sulfidic Spring. Kostka JE,

editor. Appl Environ Microbiol. 2015;81: 2025–2031. doi:10.1128/AEM.03579-14

15.   Trinugroho JP, Bečková M, Shao S, Yu J, Zhao Z, Murray JW, et al. Chlorophyll f synthesis by a

super-rogue photosystem II complex. Nature Plants. Nature Publishing Group; 2020. pp. 238–

244. doi:10.1038/s41477-020-0616-4

16.   Cardona T, Murray JW, Rutherford AW. Origin and Evolution of Water Oxidation before the

Last Common Ancestor of the Cyanobacteria. Mol Biol Evol. 2015;32: 1310–1328.

doi:10.1093/molbev/msv024

17.   Cohen Y, Jørgensen BB, Revsbech NP, Poplawski R. Adaptation to Hydrogen Sulfide of

Oxygenic and Anoxygenic Photosynthesis among Cyanobacteria. Appl Environ Microbiol.

1986;51: 398–407. doi:10.1128/aem.51.2.398-407.1986

18.   Kelly DJ, Dow CS. Microbial Photosynthesis. Wang Q, editor. Compr Biotech, The Princip, Appl

and Regal of Biotechnol in Ind, Agric and Med. Singapore: Springer Singapore; 2020.

doi:10.1007/978-981-15-3110-1

19.   Nagy CI, Vass I, Rákhely G, Vass IZ, Tóth A, Duzs A, et al. Coregulated genes link

sulfide:quinone oxidoreductase and arsenic metabolism in Synechocystis sp. strain PCC6803. J

Bacteriol. 2014;196: 3430–40. doi:10.1128/JB.01864-14

20.   Rascovan N, Maldonado J, Vazquez MP, Eugenia Farías M. Metagenomic study of red biofilms

from Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea. ISME J. 2016;10:

299–309. doi:10.1038/ismej.2015.109

21.   Sancho-Tomás M, Somogyi A, Medjoubi K, Bergamaschi A, Visscher PT, van Driessche AES, et

al. Geochemical evidence for arsenic cycling in living microbialites of a High Altitude Andean

Lake   (Laguna   Diamante,   Argentina).   Chem   Geol.   2020;549:   119681.

doi:10.1016/j.chemgeo.2020.119681

22.   Sancho-Tomás M, Somogyi A, Medjoubi K, Bergamaschi A, Visscher PT, Van Driessche AES, et

al. Distribution, redox state and (bio)geochemical implications of arsenic in present day

microbialites of Laguna Brava, Salar de Atacama. Chem Geol. 2018;490: 13–21.

doi:10.1016/j.chemgeo.2018.04.029

23.   Kurth D, Amadio A, Ordoñez OF, Albarracín VH, Gärtner W, Farías ME. Arsenic metabolism in

high altitude modern stromatolites revealed by metagenomic analysis. Sci Rep. 2017;7: 1024.

doi:10.1038/s41598-017-00896-0

24.   Visscher PT, Gallagher KL, Bouton A, Farias ME, Kurth D, Sancho-Tomás M, et al. Modern

arsenotrophic microbial mats provide an analogue for life in the anoxic Archean. Commun

Earth Environ. 2020;1: 1–10. doi:10.1038/s43247-020-00025-2

25.   Lencina AI, Soria MN, Gomez FJ, Gérard E, Farias ME. Composite microbialites: Thrombolite,

dendrolite, and stromatolite associations in a modern environment, Pozo Bravo lake, Salar de

Antofalla,   Catamarca   Puna,   Argentina.   J   Sediment   Res.   2021;91:   1305–1330.

doi:10.2110/jsr.2020.166

26.   Vignale FA, Lencina AI, Stepanenko TM, Soria MN, Saona LA, Kurth D, et al. Lithifying and Non-

Lithifying Microbial Ecosystems in the Wetlands and Salt Flats of the Central Andes. Microb

Ecol. 2022;83: 1–17. doi:10.1007/s00248-021-01725-8

27.   Farías ME, Villafañe PG, Lencina AI. Integral prospection of Andean microbial ecosystem

project. Microb Ecosyst Cent Andes Extrem Environ biofilms, Microb mats, microbialites

endoevaporites. 2020; 245–260.

28.   Nisbet EG, Fowler CMR. Archaean metabolic evolution of microbial mats. Proc R Soc B Biol Sci.

1999;266: 2375–2382. doi:10.1098/rspb.1999.0934

29.   Yacobi YZ, Köhler J, Leunert F, Gitelson A. Phycocyanin-specific absorption coefficient:

Eliminating the effect of chlorophylls absorption. Limnol Oceanogr Methods. 2015;13: 157–

168. doi:10.1002/lom3.10015

30.   Jørgensen BB, Cohen Y, des Marais DJ, Jorgensen BB, Cohen Y, des Marais DJ. Photosynthetic

action spectra and adaptation to spectral light distribution in a benthic cyanobacterial mat.

Appl Environ Microbiol. 1987;53: 879–886. doi:10.4319/lo.1992.37.8.1813

31.   Stomp M, Huisman J, Stal LJ, Matthijs HCP. Colorful niches of phototrophic microorganisms

shaped by vibrations of the water molecule. ISME J. 2007;1: 271–282.

doi:10.1038/ismej.2007.59

32.   Miller LC, Martin DS, Liu L-N, Canniffe DP, Miller LC, Canniffe · D P, et al. Composition,

Organisation and Function of Purple Photosynthetic Machinery. Microb Photosynth. 2020;

73–114. doi:10.1007/978-981-15-3110-1_4

33.   Blankenship RE, Olson JM, Miller M. Antenna Complexes from Green Photosynthetic Bacteria.

Anoxygenic Photosynthetic Bacteria. Dordrecht: Kluwer Academic Publishers; 1995. pp. 399–

435. doi:10.1007/0-306-47954-0_20

34.   Sawicki A, Chen M, Sawicki A, Chen · M. Molecular Mechanism of Photosynthesis Driven by

Red-Shifted Chlorophylls. Microb Photosynth. 2020; 3–42. doi:10.1007/978-981-15-3110-1_1

35.   Rosen BP. Biochemistry of arsenic detoxification. FEBS Lett. 2002;529: 86–92.

doi:http://dx.doi.org/10.1016/S0014-5793(02)03186-1

36.   Yang HC, Fu HL, Lin YF, Rosen BP. Pathways of Arsenic Uptake and Efflux. Current Topics in

Membranes. Academic Press Inc.; 2012. pp. 325–358. doi:10.1016/B978-0-12-394390-

3.00012-4

37.   Willsky GR, Malamy MH. Effect of arsenate on inorganic phosphate transport in Escherichia

coli. J Bacteriol. 1980;144: 366–374. doi:10.1128/jb.144.1.366-374.1980

38.   Elias M, Wellner A, Goldin-Azulay K, Chabriere E, Vorholt JA, Erb TJ, et al. The molecular basis

of phosphate discrimination in arsenate-rich environments. Nature. 2012;491: 134–137.

doi:10.1038/nature11517

39.   Saona LA, Soria M, Durán-Toro V, Wörmer L, Milucka J, Castro-Nallar E, et al. Phosphate-

Arsenic Interactions in Halophilic Microorganisms of the Microbial Mat from Laguna

Tebenquiche: from the Microenvironment to the Genomes. Microb Ecol. 2021;81: 941–953.

doi:10.1007/s00248-020-01673-9

40.   Ordóñez E, Van Belle K, Roos G, De Galan S, Letek M, Gil JA, et al. Arsenate reductase,

mycothiol, and mycoredoxin concert thiol/disulfide exchange. J Biol Chem. 2009;284: 15107–

15116. Available: https://pubmed.ncbi.nlm.nih.gov/19286650/

41.   Jiang Z, Shen X, Shi B, Cui M, Wang Y, Li P. Arsenic Mobilization and Transformation by

Ammonium-Generating Bacteria Isolated from High Arsenic Groundwater in Hetao Plain,

China. Int J Environ Res Public Health. 2022;19: 9606. doi:10.3390/ijerph19159606

42.   Rascovan N, Maldonado J, Vazquez MP, Eugenia Farías M. Metagenomic study of red biofilms

from Diamante Lake reveals ancient arsenic bioenergetics in haloarchaea. ISME J. 2016;10:

299–309. doi:10.1038/ismej.2015.109

43.   Ordoñez OF, Lanzarotti E, Kurth D, Cortez N, Farías ME, Turjanski AG. Genome comparison of

two Exiguobacterium strains from high altitude andean lakes with different arsenic resistance:

identification and 3D modeling of the Acr3 efflux pump. Front Environ Sci. 2015;3.

doi:10.3389/fenvs.2015.00050

44.   Hoeft McCann S, Boren A, Hernandez-Maldonado J, Stoneburner B, Saltikov C, Stolz J, et al.

Arsenite as an Electron Donor for Anoxygenic Photosynthesis: Description of Three Strains of

Ectothiorhodospira from Mono Lake, California and Big Soda Lake, Nevada. Life. 2016;7: 1.

doi:10.3390/life7010001

45.   Wagner GP, Kin K, Lynch VJ. Measurement of mRNA abundance using RNA-seq data: RPKM
1083   measure is inconsistent among samples. Theory Biosci. 2012;131: 281–285.
1084   doi:10.1007/s12064-012-0162-3

1085   46.   Páez-Espino D, Tamames J, De Lorenzo V, Cánovas D. Microbial responses to environmental
1086   arsenic. BioMetals. 2009;22: 117–130. doi:10.1007/s10534-008-9195-y

1087   47.   Silver S, Phung LT. Genes and enzymes involved in bacterial oxidation and reduction of
1088   inorganic arsenic. Appl Environ Microbiol. 2005;71: 599–608. doi:10.1128/AEM.71.2.599-
1089   608.2005

1090   48.   de Beer D, Meyer V, Klatt J, Li T. Photosynthesis under very high oxygen concentrations in
1091   dense microbial mats and biofilms. bioRxiv. 2018; 335299. doi:10.1101/335299

1092   49.   Castillejos Sepúlveda A, Gatti LM, Kerl CF, Chennu A, Klatt JM. Arsenic speciation analysis in
1093   porewater by a novel colorimetric assay. Sci Total Environ. 2022;827: 154155.
1094   doi:10.1016/j.scitotenv.2022.154155

1095   50.   Oremland RS, Hoeft SE, Santini JM, Bano N, Hollibaugh RA, Hollibaugh JT. Anaerobic oxidation
1096   of arsenite in Mono Lake water and by a facultative, arsenite-oxidizing chemoautotroph,
1097   strain MLHE-1. Appl Environ Microbiol. 2002;68: 4795–802. doi:10.1128/AEM.68.10.4795

1098   51.   Zhang J, Zhou W, Liu B, He J, Shen Q, Zhao FJ. Anaerobic arsenite oxidation by an autotrophic
1099   arsenite-oxidizing bacterium from an arsenic-contaminated paddy soil. Environ Sci Technol.
1100   2015;49: 5956–5964. doi:10.1021/es506097c

1101   52.   Franks J, Stolz JF. Flat laminated microbial mat communities. Earth-Science Rev. 2009;96:
1102   163–172. doi:10.1016/J.EARSCIREV.2008.10.004

1103   53.   Flores-Kim J, Darwin AJ. The Phage Shock Protein Response. Annu Rev Microbiol. 2016;70: 83–
1104   101. doi:10.1146/ANNUREV-MICRO-102215-095359

1105   54.   Junglas B, Huber ST, Heidler T, Schlösser L, Mann D, Hennig R, et al. PspA adopts an ESCRT-III-

1106   like fold and remodels bacterial membranes. Cell. 2021;184: 3674-3688.e18.

1107   doi:10.1016/J.CELL.2021.05.042

1108   55.   Sheridan KJ, Duncan EJ, Eaton-Rye JJ, Summerfield TC. The diversity and distribution of D1

1109   proteins in cyanobacteria. Photosynth Res. 2020;145: 111–128. doi:10.1007/s11120-020-

1110   00762-7

1111   56.   Lubitz W, Lendzian F, Bittl R. Radicals, radical pairs and triplet states in photosynthesis. Acc

1112   Chem   Res.   2002;35:   313–320.

1113   doi:10.1021/AR000084G/ASSET/IMAGES/LARGE/AR000084GF00007.JPEG

1114   57.   Budinoff CR, Hollibaugh JT. Arsenite-dependent photoautotrophy by an Ectothiorhodospira-

1115   dominated consortium. ISME J. 2008;2: 340–343. doi:10.1038/ismej.2007.115

1116   58.   Blokhina O, Virolainen E, Fagerstedt K V. Antioxidants, oxidative damage and oxygen

1117   deprivation stress: A review. Annals of Botany. 2003. pp. 179–194. doi:10.1093/aob/mcf118

1118   59.   Hakala-Yatkin M, Sarvikas P, Paturi P, Mäntysaari M, Mattila H, Tyystjärvi T, et al. Magnetic

1119   field protects plants against high light by slowing down production of singlet oxygen. Physiol

1120   Plant. 2011;142: 26–34. doi:10.1111/j.1399-3054.2011.01453.x

1121   60.   Marais A, Sinayskiy I, Petruccione F, Van Grondelle R. A quantum protective mechanism in

1122   photosynthesis. Sci Reports 2015 51. 2015;5: 1–8. doi:10.1038/srep08720

1123   61.   Del Razo LM, Quintanilla-Vega B, Brambila-Colombres E, Calderón-Aranda ES, Manno M,

1124   Albores A. Stress proteins induced by arsenic. Toxicol Appl Pharmacol. 2001;177: 132–148.

1125   doi:10.1006/TAAP.2001.9291

1126   62.   Latifi A, Ruiz M, Zhang C-C. Oxidative stress in cyanobacteria. FEMS Microbiol Rev. 2009;33:

1127   258–278. doi:10.1111/j.1574-6976.2008.00134.x

1128   63.   Jackson CR, Langner HW, Donahoe-Christiansen J, Inskeep WP, McDermott TR. Molecular

1129   analysis of microbial community structure in an arsenite-oxidizing acidic thermal spring.

1130   Environ Microbiol. 2001;3: 532–542. doi:10.1046/j.1462-2920.2001.00221.x

1131 64.   Lebrun E, Brugna M, Baymann F, Muller D, Lièvremont D, Lett M-C, et al. Arsenite oxidase, an

1132   ancient bioenergetic enzyme. Mol Biol Evol. 2003;20: 686–93. doi:10.1093/molbev/msg071

1133 65.   Grim SL, Dick GJ. Photosynthetic versatility in the genome of Geitlerinema sp. PCC 9228

1134   (Formerly Oscillatoria limnetica 'solar lake'), a model anoxygenic photosynthetic

1135   cyanobacterium. Front Microbiol. 2016;7: 1–14. doi:10.3389/fmicb.2016.01546

1136 66.   Sforna MC, Philippot P, Somogyi A, Van Zuilen MA, Medjoubi K, Schoepp-Cothenet B, et al.

1137   Evidence for arsenic metabolism and cycling by microorganisms 2.7 billion years ago. Nat

1138   Geosci. 2014;7: 811–815. doi:10.1038/ngeo2276

1139 67.   De Beer D, Schramm A, Santegoeds CM, Kühl M, Kuhl M. A nitrite microsensor for profiling

1140   environmental biofilms. Appl Environ Microbiol. 1997;63: 973–7. Available:

1141   http://www.ncbi.nlm.nih.gov/pubmed/16535560

1142 68.   Jeroschewski P, Steuckart C, Kühl M. An Amperometric Microsensor for the Determination of

1143   H 2 S in Aquatic Environments. Anal Chem. 1996;68: 4351–4357. doi:10.1021/ac960091b

1144 69.   Klatt JM, Meyer S, Häusler S, Macalady JL, De Beer D, Polerecky L. Structure and function of

1145   natural sulphide-oxidizing microbial mats under dynamic input of light and chemical energy.

1146   ISME J. 2016;10: 921–33. doi:10.1038/ismej.2015.167

1147 70.   Revsbech NP. An oxygen microsensor with a guard cathode. Limnology and Oceanography.

1148   John Wiley & Sons, Ltd; 1989. pp. 474–478. doi:10.4319/lo.1989.34.2.0474

1149 71.   Tros M, Mascoli V, Shen G, Ho MY, Bersanini L, Gisriel CJ, et al. Breaking the Red Limit:

1150   Efficient Trapping of Long-Wavelength Excitations in Chlorophyll-f-Containing Photosystem I.

1151   Chem. 2021;7: 155–173. doi:10.1016/J.CHEMPR.2020.10.024

72.   Cline JD. Spectrophotometric determination of hydrogren sulfide in natural waters. Limnol

Oceanogr. 1969;14: 454–458. doi:10.4319/lo.1969.14.3.0454

73.   Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data.

Bioinformatics. 2014;30: 2114–2120. doi:10.1093/bioinformatics/btu170

74.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for

large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.

2015;31: 1674–1676. doi:10.1093/bioinformatics/btv033

75.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:

357–359. doi:10.1038/nmeth.1923

76.   Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: Prokaryotic gene

recognition and translation initiation site identification. BMC Bioinformatics. 2010;11: 1–11.

doi:10.1186/1471-2105-11-119

77.   Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: An advanced

analysis and visualization platformfor 'omics data. PeerJ. 2015;2015: e1319.

doi:10.7717/PEERJ.1319/SUPP-5

78.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al. KofamKOALA:

KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Valencia A,

editor. Bioinformatics. 2020;36: 2251–2252. doi:10.1093/bioinformatics/btz859

79.   Tatusov RL, Galperin MY, Natale DA, Koonin E V. The COG database: a tool for genome-scale

analysis of protein functions and evolution. Nucleic Acids Res. 2000;28: 33.

doi:10.1093/NAR/28.1.33

80.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat

Methods. 2015;12: 59–60. doi:10.1038/nmeth.3176

81.   Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The InterPro

protein families and domains database: 20 years on. Nucleic Acids Res. 2021;49: D344–D354.

doi:10.1093/nar/gkaa977

82.   Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, et al. InterProScan 5: Genome-scale

protein   function   classification.   Bioinformatics.   2014;30:   1236–1240.

doi:10.1093/bioinformatics/btu031

83.   Sillitoe I, Dawson N, Lewis TE, Das S, Lees JG, Ashford P, et al. CATH: Expanding the horizons of

structure-based functional annotations for genome sequences. Nucleic Acids Res. 2019;47:

D280–D284. doi:10.1093/nar/gky1097

84.   Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE: the

conserved domain database in 2020. - PubMed - NCBI. Nucleic Acids Res. 2020;48: D265–

D268. Available: https://academic.oup.com/nar/article-abstract/48/D1/D265/5645006

85.   Pedruzzi I, Rivoire C, Auchincloss AH, Coudert E, Keller G, De Castro E, et al. HAMAP in 2015:

Updates to the protein family classification and annotation system. Nucleic Acids Res.

2015;43: D1064–D1070. doi:10.1093/nar/gku1002

86.   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: More genomes, a

new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res.

2019;47: D419–D426. doi:10.1093/nar/gky1038

87.   Bateman A, Birney E, Durbin R, Eddy SR, Howe KL, Sonnhammer ELL. The Pfam protein

families database. Nucleic Acids Research. 2000. pp. 263–266. doi:10.1093/nar/28.1.263

88.   Nikolskaya AN, Arighi CN, Huang H, Barker WC, Wu CH. PIRSF Family Classification System for

Protein Functional and Evolutionary Analysis. Evol Bioinforma. 2006;2: 117693430600200.

doi:10.1177/117693430600200033

89.   Attwood TK, Coletta A, Muirhead G, Pavlopoulou A, Philippou PB, Popov I, et al. The PRINTS

database: A fine-grained protein sequence annotation and analysis resource-its status in

2012. Database. 2012;2012. doi:10.1093/database/bas019

90.   Sigrist CCJA, Castro E De, … LC-N acids, 2012 undefined, de Castro E, Cerutti L, et al. New and

1202   continuing developments at PROSITE. 2012;41: D344–D347. doi:10.1093/nar/gks1067

1203   91.   Letunic I, Bork P. 20 years of the SMART protein domain annotation resource. Nucleic Acids

1204   Res. 2018;46: D493–D496. doi:10.1093/nar/gkx922

1205   92.   Brown S, Pegg S, Ojha S, Huang C, Ferrin TE, Babbitt PC. The Structure-Function Linkage

1206   Database. FASEB J. 2006;20. doi:10.1096/fasebj.20.5.a899-c

1207   93.   Pandurangan AP, Stahlhacke J, Oates ME, Smithers B, Gough J. The SUPERFAMILY 2.0

1208   database: A significant proteome update and a new webserver. Nucleic Acids Res. 2019;47:

1209   D490–D494. doi:10.1093/nar/gky1130

1210   94.   Haft DH, Selengut JD, Richter RA, Harkins D, Basu MK, Beck E. TIGRFAMs and genome

1211   properties in 2013. Nucleic Acids Res. 2013;41. doi:10.1093/nar/gks1234

1212   95.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

1213   Architecture and applications. BMC Bioinformatics. 2009;10: 1–9. doi:10.1186/1471-2105-10-

1214   421/FIGURES/4

1215   96.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: An adaptive binning

1216   algorithm for robust and efficient genome reconstruction from metagenome assemblies.

1217   PeerJ. 2019;2019: e7359. doi:10.7717/PEERJ.7359/SUPP-3

1218   97.   Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover

1219   genomes from multiple metagenomic datasets. Bioinformatics. 2016;32: 605–607.

1220   doi:10.1093/bioinformatics/btv638

1221   98.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A

1222   standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of

1223   life. Nat Biotechnol. 2018;36: 996–1004. doi:10.1038/nbt.4229

99.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with

the Genome Taxonomy Database. Bioinformatics. 2020;36: 1925–1927.

doi:10.1093/BIOINFORMATICS/BTZ848

100. Woodcroft BJ, Newell R. wwood/CoverM: Read coverage calculator for metagenomics. Centre

for Microbiome Research, School of Biomedical Sciences, Faculty of Health, Queensland

University of Technology; 2021 [cited 22 Jul 2022]. Available:

https://github.com/wwood/CoverM#readme

101. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Cham: Springer-Verlag New York;

2016. doi:10.1007/978-3-319-24277-4

102. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria; 2020

[cited 3 May 2023]. Available: https://www.r-project.org/

103. Bushnell B. BBMap short-read aligner, and other bioinformatics tools. 2015.

104. Kolde R. CRAN - Package pheatmap. In: https://cran.r-project.org/package=pheatmap

[Internet]. 2019 [cited 12 Jun 2022]. Available: https://cran.r-

project.org/web/packages/pheatmap/index.html

1239

a)

Microbialites

Pink mats

Lithifying dark gray mats

b)

Pink crust

Green layer

Red layer

Brown bottom sediment

c)

Pink crust

Green

Red

Brown

d)

Depth (mm)

675 nm (Chl a)

715 nm (BChl e/Chl f)

805 nm (Bchl a)