Disorganized Inhibitory Dynamics in Hippocampal area CA1 of 22q11.2 Deletion Mutant Mice

Stephanie A. Herrlinger[1,2,*], Bovey Y Rao[1,2,3,*], Margaret E. Conde Paredes[1,2,3,4], Anna L. Tuttman[1,2], Haroon Arain[1], Erdem Varol[4], Joseph A. Gogos[1,2,5,6,7], and Attila Losonczy[1,2]

[1] Department of Neuroscience, Columbia University, New York, NY, United States

[2] Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, United States

[3] Doctoral Program in Neurobiology and Behavior, Columbia University, New York, NY, United States

[4] Department of Computer Science & Engineering, Tandon School of Engineering, New York University, New York, NY, United States

[5] Department of Physiology, Columbia University, New York, NY, United States

[6] Department of Psychiatry, Columbia University, New York, NY, United States

[7] Stavros Niarchos Foundation Center for Precision Psychiatry and Mental Health, Columbia University, New York, NY, United States

Individuals with the 22q11.2 deletion syndrome, one of the strongest genetic risk factors for schizophrenia, demonstrate cognitive impairments including episodic memory dysfunction. Place cell activity of excitatory pyramidal neurons in the hippocampus supporting episodic memory are impaired in a mouse model for the 22q11.2 deletion ($Df(16)A^{+/-}$). While excitatory dynamics are under tight inhibitory control by multiple subtypes of GABAergic interneurons, previous studies have predominantly focused on a single subtype of PV-expressing interneurons; there have not yet been studies describing the functional relationships between molecularly identified inhibitory types in $Df(16)A^{+/-}$ mice. Here, we examined interneuron subtype-specific activity dynamics in the dorsal hippocampal area CA1 of $Df(16)A^{+/-}$ mice during random foraging and spatial reward learning tasks. Capitalizing on 3D acousto-optical deflector two-photon microscopy with *post hoc* immunohistochemical identification, we found that multiple interneuron types exhibit aberrant responses to reward locations and delayed reward enrichment extinction. $Df(16)A^{+/-}$ inhibitory interneurons also carry markedly reduced spatial information in a subtype-dependent manner. We observed task-dependent changes in the correlation structure and coactivity among multiple GABAergic subtypes, suggesting a broadly disorganized microcircuit functionality in mutant mice. Overall, we identify widespread and heterogeneous subtype-specific alterations in interneuron dynamics during spatial reward learning, reflecting impaired flexibility and organization in CA1 inhibitory microcircuits. Our study provides critical insights into how schizophrenia-risk mutations affect local-circuit interactions among diverse cell types in the mouse hippocampus during learning and spatial navigation.

INTRODUCTION

Adults and children with the 22q11.2 deletion syndrome (22q11.2DS) exhibit social and emotional impairments alongside a range of cognitive deficits[1,2]. The 22q11.2 deletion is one of the strongest genetic risk factors for schizophrenia, leading to a marked ~30-fold increase in the risk of developing the disorder[1,3,4]. Cognitive deficits represent core features of schizophrenia and are likely mediated by the same abnormal physiological processes that underlie other more prototypical symptoms[5–8]. Schizophrenia patients experience a spectrum of cognitive deficits, which are often treatment-resistant. Among these, episodic memory dysfunction[5,9,10] is both relatively common and highly debilitating. Episodic memory deficits, which have the largest effect sizes among cognitive impairments in schizophrenia[11], are often present prior to the onset of psychosis, and are frequently observed in relatives of affected individuals[11–13]. Episodic memory relies on the hippocampus, which exhibits both anatomical and functional deficits in schizophrenia and 22q11.2 deletion patients[14–22].

While mouse models cannot fully capture the symptoms associated with psychiatric diseases, they are valuable for modeling disease-associated mutations and assessing core aspects of a disorder that manifest in an equivalent manner in rodents[6,23,24]. The hippocampus in particular is one of the most evolutionarily conserved brain regions[25,26] and the mouse hippocampus recapitulates human features in both normal physiology[27–29] and pathophysiology of neurological disorders[30–33]. By using animal models of rare, highly penetrant risk mutations, and coupling these models with behavioral tasks that assess key cognitive features of learning and memory, researchers can apply high-resolution neuroscience techniques to gain unprecedented insights into the mechanisms underlying mental illness[34–36]. These approaches have introduced a new biological framework for studying psychiatric disease, emphasizing the disordered activity and plasticity of both local and long-range neuronal assemblies and circuits[34,35,37–47]. Such disruptions may reveal common points of convergence of genetic liability, leading to erratic neuronal activation patterns that trigger the constellation of cognitive symptoms, including episodic memory deficits[23].

Hippocampal place cells, a subset of hippocampal principal cells that are selectively active in specific environmental locations, provide cellular substrates for episodic memory and goal-directed spatial learning[48,49]. Place cell ensembles are both sparse and dynamic[43,50–57], forming rapidly in novel environments[58–60], and undergoing experience-dependent stabilization as the environment transitions from novel to familiar[61–63]. Place cells are also responsive to salient reinforcers, such as rewards[43,51,64–67], with distinct, sparse populations of place cells representing different contexts[52,63,67–69]. Our previous study uncovered behavioral and place cell correlates of episodic memory deficits in *Df(16)A^+/−* mice, an animal model for the 22q11.2 deletion, by using a spatial goal-oriented, hippocampal-dependent learning task[43]. *Df(16)A^+/−* mice exhibited a marked impairment in learning performance, which was disrupted by changes in task-context or reward

location. Furthermore, place cells in the dorsal hippocampal area CA1 of mutant mice exhibited reduced long-term stability, impaired context-related remapping, and a lack of reward-related reorganization.

45 These observations provide a crucial entry point for investigating how schizophrenia risk mutations lead to altered hippocampal microcircuit dynamics during learning. However,, the role of many hippocampal circuit components has remained uncharacterized, representing a major gap in our understanding of circuit mechanisms that drive disease-predisposing genetic lesions into clinical symptoms. In particular, GABAergic interneurons (IN) play a crucial role in shaping

50 the activity of excitatory neuron populations in the brain by regulating the timing, amplitude, and synchronization of neuronal firing[70–72]. CA1 pyramidal cell activity and plasticity are under stringent inhibitory control by a diverse array of IN subtypes, each with distinct morphological, molecular, and functional properties[73–75]. Different IN subtypes target specific subcellular compartments of excitatory neurons and have specialized roles in regulating microcircuit

55 function[74,76–81]. Dysfunction in inhibitory circuits has been implicated in numerous psychiatric disorders, particularly in schizophrenia pathophysiology[41,42,82–91]. Therefore, understanding the role of inhibitory neurons in microcircuit dysfunction is key to unraveling the complexity of neural circuits and their contribution to various brain functions and psychiatric conditions. Yet, the *in vivo* alterations in activity dynamics across most CA1 IN subtypes resulting from 22q11.2 deletions

60 remain completely unknown. Prior studies were limited to investigating IN subtypes in isolation ('one cell-type at a time')[85,92–94], which presents a major limitation to capturing a complete picture of inhibition. This piecemeal approach has hindered the identification of disease-associated inhibitory dysfunction, highlighting the need for holistic experimental and analytical strategies that can simultaneously assess functional interactions among multiple IN subtypes.

To overcome the previous studies' limitations, here we innovate both in terms of applying advanced imaging methods and designing new analysis approaches to yield novel insights about hippocampal circuit organization. Namely, we utilized a three-dimensional two-photon (2p) Ca^{2+} imaging strategy with multiplexed molecular characterization to capture a simultaneous functional

70 readout of a large volumetric population of GABAergic IN subtypes in the dorsal hippocampal CA1 area[71,72]. We then introduce a new computational model to quantify the interplay between molecular readouts of INs and their pairwise functional correlation patterns. This analysis enables us to approximate circuit co-activation changes amongst INs from different genotypes to make many-cell-types-at-time inferences about circuit differences (as opposed to existing one-cell-at-

75 a-time approaches). These techniques were applied on *Df(16)A*[+/–] mice and wild-type (WT) littermates to address key gaps in our understanding of the local circuit alterations associated with the 22q11.2 deletion and to dissect the relationships between cellular components that contribute to the pathophysiology of schizophrenia-related memory deficits.

80 **RESULTS**

Interneuron subtype number or distribution in dorsal CA1 is unaltered in *Df(16)A*[+/-] mice.

To assess interneuron dynamics within the hippocampus during navigation in *Df(16)A*$^{+/-}$ mice, we employed an unbiased, large-scale, 3D method for *in vivo* 2p Ca^{2+} imaging of dorsal CA1 INs with random access acousto-optical deflector (3D-AOD) microscopy (Fig. 1)[95,96]. We performed 3D-AOD 2p Ca^{2+} imaging in *Vgat-Cre::Df(16)A*$^{+/-}$ mice and WT littermates injected with a Cre-dependent adeno-associated virus for GCaMP7f expression in dorsal hippocampal area CA1 INs during a Random Foraging behavior paradigm (see methods, Fig. 1A,B)[95,96]. After taking a Z-stack of the dorsal CA1 field of view (Fig. 1C), we manually selected individual IN somata distributed across the 3D volume (Fig. 1D). This approach enables simultaneous, longitudinal monitoring of activity dynamics in large IN populations (up to ~250 cells per mouse) across all CA1 sublayers (Fig. 1E). We imaged the same INs for two consecutive days while the head-fixed mice navigated a spatially enriched treadmill belt for randomly deposited water rewards. We performed *post hoc* immunohistochemical (IHC) identification of imaged INs and manually registered them to their *in vivo* cell somas based on their GCaMP7f signal (Fig. 1F). We identified five major IN subtypes: axo-axonic cells (AACs), parvalbumin-expressing basket cells (PVBCs), somatostatin-expressing cells (SomCs), ivy/neurogliaform cells (IvC/NGFCs), and cholecystokinin (CCKCs) cells (Fig. 1G, see methods, Table 2)[73,74,80,95]. Other IN populations including vasoactive intestinal polypeptide-expressing and calretinin-expressing interneurons were not examined in this study due to the limitations in the number of IHC channels and antibody combinations available for use within the same animal. Our analysis revealed that the subtype-specific distribution of imaged INs across CA1 sublayers was not noticeably altered in the mutants, and there were no significant differences in the number of IN subtypes recorded per mouse or their proportional distribution within each CA1 layer (WT n=6 mice, 1127 total cells, 187.83 ± 15.22 cells per mouse; Df(16)A$^{+/-}$ n = 5,775 total cells, 155 ± 30.21 cells per mouse)(Suppl. Fig. 1A-C). Using the markers in our subtyping strategy (see methods), we quantified the total number of cells expressing Parvalbumin (PV+), Somatostatin (SST+), Neuropeptide Y (NPY+), Satb1+, and Cholecystokinin (CCK+) in dorsal CA1 and assessed labeled cell soma density by layer (Suppl. Fig. 1D-I). Due to limitations in imaging depth and resolution, fewer INs were recovered in deeper layers of the hippocampus. Unidentified INs, which represent a highly heterogeneous VGAT population with variable functional properties, were excluded from subsequent analyses. Consistent with findings from other mouse models of the 22q11.2 deletion[97], none of these quantifications revealed significant differences in the number of immuno-positive cells for each marker.

Figure 1. Df(16)A$^{+/-}$ interneurons exhibit altered subtype-specific reward related responses. (A) Schematic of the *in vivo* imaging setup using acousto-optical deflector (*AOD*) microscopy and viral GCaMP7f expression strategy. *Vgat-Cre::Df(16)A$^{+/-}$* mice and WT littermates were injected with AAV1(*syn-FLEX-GCaMP7f*) in dorsal hippocampal area CA1 to express genetically encoded Ca^{2+} indicator, GCaMP7f, in interneurons (INs). (B) Schematic of the Random Foraging behavior paradigm. Water-restricted mice were trained to run for randomly deposited water rewards across a 2-meter burlap belt with distinct features sewed across the belt. (C) 3-dimensional reconstruction of *in vivo* z-stack of

INs in CA1. (D) Schematic of the 3D imaging targeting strategy on the AOD, coronal view. *s.o.: stratum oriens, s.p.: stratum pyramidale, s.r.: stratum radiatum, s.l.m.: stratum lacunosum moleculare*. (E) Example time averages of the chessboard scan of 100 simultaneously imaged INs. *ROI*: regions of interest. (F) Example pair of *in vivo* and *ex vivo* images manually registered between imaging modalities. (G) Example images for each channel demonstrating the immunohistochemical (IHC) targeting strategy for IN cell-type characterization. Immunolabels are on the X-axis: *CCK*: cholecystokinin; *SST*: somatostatin; *NPY*: neuropeptide Y; *Pvalb*: parvalbumin. Cell-type labels used based on IHC strategy on the Y-axis: *AAC*: Axo-Axonic cells; *PVBC*: Parvalbumin Basket cell; *SomC*: Somatostatin-expressing cell; *IvC/NGFC*: Ivy/Neurogliaform cell; *CCKC*: Cholecystokinin-expressing cell. (H) Example relative GCaMP7f fluorescence *(ΔF/F)* traces from molecularly identified interneurons during running and immobility bouts from 1 animal for each genotype. WT Cell Numbers: AAC = 62 cells, SomC = 132 cells, PVBC = 46 cells, CCKC = 32 cells, IvC/NGFC = 73 cells; *Df(16)A$^{+/-}$* cell numbers: AAC = 34 cells, SomC = 125 cells, PVBC = 51 cells, CCKC = 14 cells, IvC/NGFC = 47 cells. (I) Example relative GCaMP7f fluorescence *(ΔF/F)* select traces from interneurons highlighting rewarded areas during random foraging. (J) Individual cell responses to reward during random foraging task. Left: WT; Right: *Df(16)A$^{+/-}$*. (K) Averaged z-scored reward-onset peri-event time averages (PETAs) between -1.5 and 1.5 z-scores by cell type across all cells within cell type during the Random Foraging task for 3 seconds pre- and 3 seconds post-event. (L) Post - Pre PETA responses per cell type before and after reward onset comparing the 3 seconds pre- to 3 seconds post-event. n.s.: not significant. *: P < 0.05. **: P < 0.01. ***: P < 0.001. See Suppl. Table 1 for detailed statistics.

Unaltered locomotion-related interneurons dynamics in *Df(16)A$^{+/-}$* mice

To assess *in vivo* functional activity dynamics of molecularly identified CA1 interneurons (INs), *Vgat-Cre::Df(16)A$^{+/−}$* mice and their wild-type (WT) littermates were water-restricted and trained to forage for randomly distributed water rewards on a spatially cued belt during 3D-AOD imaging (Fig. 1B). Licking and average running velocity were consistent between *Df(16)A$^{+/-}$* mice and WTs during this task (Suppl. Fig. 2A,B). Additionally, *Df(16)A$^{+/-}$* mice ran the same number of laps without any changes in belt occupancy (Suppl. Fig. 2C,D). Most INs exhibited positive activity correlation with locomotion (Fig. 1H), which remained true in the described cell types in the mutant mice (Suppl. Fig. 2E)[95]. Similarly, CCKCs exhibited a bimodal correlation profile in both genotypes, with a subset of CCKCs showing negative velocity correlation[77,95]. Peri-event time average (PETA) analysis of subtype-specific IN activity during *Run-Start* and *Run-Stop* events showed no significant differences between genotypes (Suppl. Fig. 2F,G). Further analysis using a generalized linear model confirmed that velocity remained a significant predictor of IN activity, with no observable differences between the genotypes (Suppl. Fig. 2H). Thus, locomotion-related activity dynamics of INs remain largely unaltered in mutant mice.

Df(16)A$^{+/-}$ interneurons exhibit subtype-specific alterations in reward-related inhibitory dynamics during random foraging

Locomotion equally drives interneuron activity in both genotypes (Fig. 1H, Suppl. Fig. 2). However, closer examination of activity traces suggested potential differences in IN responses to reward delivery between the genotypes (Fig. 1I). To isolate these differences, we regressed out the velocity component from the calcium traces to remove the highly correlated IN activity related to movement speed. We then investigated inhibitory responses to random rewards by analyzing neuronal activity during the random foraging task focusing on reward-onset PETAs. Our analysis revealed that more *Df(16)A$^{+/-}$* INs exhibited stronger post-reward onset responses overall compared to wildtype INs (Fig. 1J). This post-reward enhanced activity was subtype-specific. For instance, SomCs, which typically showed increased post-reward activity in wildtype mice,

170   exhibited an even greater recruitment in *Df(16)A*[+/-]mice. Similarly, AACs and PVBCs which typically exhibit decreased post-reward activity in WT mice, displayed a significant shift towards higher post-reward responses (Fig. 1K/L) in the *Df(16)A*[+/-] group. These findings indicate a substantial alteration in the inhibitory reward response to random rewards, in Df(16)A+/- mice

175   ***Df(16)A*[+/-] interneurons exhibit enhanced reward zone modulation and attenuated plasticity during a spatial navigation task.**

Considering our findings that inhibition is enhanced in response to random reward presentation and that *Df(16)A*[+/-] mice display decreased principal neuron enrichment near known reward
180   zones[43,98], we next aimed to characterize the spatial reward-related activity of INs, which is known to be cell type-specific[95,99,100]. To achieve this, we designed a simple spatially guided reward translocation task to explore how IN activity is shaped under two conditions: 1) when the reward locations are known and 2) when reward zones are translocated. We trained the mice on a fixed reward task for 3 days prior to imaging with a non-operantly delivered reward (Fig. 2A). On the
185   fourth day, CA1 IN Ca^{2+} activity was imaged during a 10-min trial in the familiar condition (Fig. 2B). On the next day, the location of the non-operant reward was translocated to another position on the treadmill belt and was repeated at that new position on the third day (Reward Translocation task, Fig. 2B). Both wildtype and *Df(16)A*[+/-] mice learned this task well over the course of the three days, as evidenced by increased licking leading up to and inside the reward zone during all three
190   recording days (Suppl. Fig. 3A). Mice displayed a quick adjustment to the translocated reward zone based on their licking behavior on the first translocation day (Suppl. Fig. 3A, middle). *Df(16)A*[+/-] mice exhibited more frequent licking than their WT counterparts (Suppl. Fig. 3B), particularly in the reward zones, suggesting that they continued licking even after reward delivery ceased (after 2 seconds) (Suppl. Fig. 3A). This phenotype appeared to be specific to fixed reward
195   tasks, as there were no differences in licking behavior during the Random Foraging task (Suppl. Fig. 2A), indicating that hyperactivity in this mouse line[98,101,102] was unlikely to explain these findings. When comparing all three days together, we additionally found that the mutant mice spent a greater proportion of time in the reward zones (Suppl. Fig. 3C/D) despite running the same number of laps (Suppl. Fig. 3E) and having equivalent average speed (Suppl. Fig. 4F/G).
200   Overall, *Df(16)A*[+/-] mice learned the spatial navigation task while exhibiting perseverant behavior at reward zones.

We next interrogated distinct reward-related inhibitory responses during a Reward Translocation task (Fig. 2B) and observed subtype-specific changes in reward-related activity of multiple IN
205   subtypes between genotypes. By comparing post-pre reward responses from reward-onset PETAs we identified increased pre-reward responses from SomCs and a stronger post-reward response from PVBCs (Suppl. Fig. 4A/B, Fig. 2C). We next examined the spatially defined average activity of each IN subtype preceding known reward locations. As expected, SomC activity was enriched leading up to both the familiar (Fig. 2D) and translocated reward zones in
210   both the wildtypes and *Df(16)A*[+/-] INs (Suppl. Fig. 2C)[95]. However, in addition to SomCs, other IN subtypes, including PVBCs and AACs, also showed increased activity leading up to reward zones in *Df(16)A*[+/-] INs (Fig. 2D, Suppl. Fig. 2C, right). After reward translocation, we observed that SomCs and PVBCs maintained elevated activity levels preceding the old reward zone,

indicating a persistent, inflexible response to prior reward locations in the mutant mice (Fig. 2E-H). To quantify these findings we computed a reward zone modulation index (RMI, as previously described[95]) which normalized pre-reward zone activity to activity levels across the remainder of the belt (Fig. 2G/I, Suppl. Fig. 4D/E). This analysis confirmed sustained SomC, PVBC, and IvC/NGFC activity in the old reward zones of *Df(16)A*$^{+/-}$ mice compared to their WT littermates after reward translocation (Fig. 2G,I,K). Interestingly, AACs uniquely exhibited aberrant reward-related activity only in the active reward zones (Fig. 2D, J, Suppl. Fig. 4C, F), whereas IvC/NGFCs showed aberrant activity only after the reward had been translocated (Fig. 2J/K). These findings suggest that task-dependent reward-related alterations in the *Df(16)A*$^{+/-}$ mice behavior are accompanied by subtype-specific IN activity, reflecting aberrant and inflexible local inhibition during spatial navigation.

Figure 2.

Figure 2. Aberrant subtype-specific activity during spatial reward learning task. (A) Reward translocation task timeline schematic. Orange: familiar task trial days, Blue: reward translocation days. (B) Reward translocation task (*RZ*: reward zone) schematic for the three days of imaging, the familiar reward zone days (top), and the two reward translocation days (bottom). (C) Post - Pre cell responses centered around the reward onset for reward translocation task observed across all three days. (D) Average responses of each cell type in the preceding 40 cm of the belt to the reward location in the WT (left) and *Df(16)A+/-* (right) on Day 1. (E) Average responses of each cell type in the preceding 40 cm of the belt to the prior reward location in the WT (left) and *Df(16)A+/-* (right) on Day 2 after reward translocation. (F) Average responses of SomCs in the preceding 40 cm of the belt to the old reward location in the WT (red) and

Df(16)A[+/-] interneurons represent reduced spatial information across tasks

We previously identified impaired place cell stability in CA1 and fewer place fields of place cells in mutant mice[43]. Given that place field formation and stability are under strong local inhibitory control[64,103,104], and a subset of INs themselves exhibit positive or negative place fields in WT mice[81,95,96,103,105,106], it is plausible that the aberrant reward-related activity exhibited by *Df(16)A*[+/-] INs (Fig. 1, 2) could fail to adequately support spatial representations and contribute to the spatial learning deficits observed in *Df(16)A*[+/-] mice[43]. To investigate this, we specifically examined spatial IN activity by asking how well we could decode the mouse position during the Random Foraging task based solely on IN activity (Fig. 3A). We trained a support vector machine based on all recorded IN activity (see methods) and found a significant increase in decoding error in mutant mice INs compared to controls. This deficit persisted even after correcting for the number of cells recorded per animal and the number of laps ran by mice (Fig. 3B,C). In addition, we also observed a non-significant decrease in spatial information encoded in mutant IN activity as measured by the Skaggs Information content (Fig. 3D)[107,108].

Next, we calculated spatial tuning curves for spatial selectivity in all IN subtypes during random foraging trials using the peak method (Fig. 3E)[95,109], and found that IN place field widths did not differ between genotypes. Most place fields were detected within 30 bins of the track (Suppl. Fig. 5A). However, we detected fewer INs with place fields in *Df(16)A*[+/-] INs compared with controls (Fig. 3F) and this deficit became more pronounced when cells with only negative place fields were excluded (Fig. 3G). Notably, negative place cells did not show any reduction between genotypes (Suppl. Fig. 5B), suggesting a specific loss of positive place field information in mutant IN activity. When considering all detected place field types, mutant INs appear to generate fewer place fields per cell (Suppl. Fig. 3C). This deficit was particularly pronounced when again focusing solely on positive place fields (Fig. 3H), indicating a selective loss in positive place field enrichment in the mutant INs. Further analysis of cell types involved in this reduction revealed that AACs, PVBCs, and SomCs exhibited significant reductions in the proportion of positive place cells, with IvC/NGFCs potentially showing decreased place preference as well l (Fig. 3I). Overall, fewer INs exhibit spatial selectivity in mutant mice (Fig. 3).

Finally, we assessed if the fixed reward zones in the Reward Translocation task resulted in increased spatial representation between the genotypes. When accounting for spatial representations of reward, there was no significant difference in the number of place fields detected using the peak method during this task (Suppl. Fig. 5D). However, when we examined the spatial relationship of these fields regarding reward by computing the distance between the center of each positive place field and the reward zone, we found that the positive IN place fields were significantly closer to the reward in mutant mice when compared to littermate controls (Fig.

3J). Furthermore, consistent with our findings in the Random Foraging task when comparing all three days, there was still a significant decrease in spatial information represented in mutant IN activity during the Reward Translocation task (Fig. 3K) as measured using the Skaggs Information content method[108]. This suggests that despite the aberrant reward-related activity observed in PVBCs, AACs, and IvC/NGFCs, this altered activity did not contribute to an increase in the overall spatial information retained from the treadmill belt. Instead, it led to severely altered and persistent reward-related activity (Fig. 2).

Figure 3.

Figure 3: Reduced spatial information in *Df(16)A⁺/⁻* interneuron activity. (A) All recorded interneuron (IN) activity was used to train a support vector machine (*SVM*) to decode discrete position bins (*D.E.*: decoding error) across the belt. Each bin is equivalent to 2 cm. (B) SVM results iterated by number of cells used with constant laps per genotype per bin. (C) SVM results iterated by the number of laps used with constant number of cells per genotype per bin. (D) Skaggs information content of IN activity by genotype. Dots represent individual cell information content. (E) Example traces of positive place, negative place, both, and neither cells, based on the Peak Place method. Blue lines: lower bound to define significant negative place fields, red lines define significant C.I.: confidence interval. (F) Place field proportion between genotypes. (G) Proportion of positive place fields detected. (H) Number of positive place fields

detected per cell. (I) Place field proportions by interneuron cell type. (J) Distance from the center of the detected positive place field (P.F.), surpassing the 95th percentile shuffle distribution) to the center of the reward zone. (K) Skaggs information content of interneuron activity by genotype during the reward translocation task. n.s.: not significant. *: P < 0.05. **: P < 0.01. ***: P < 0.001. See Suppl. Table 1 for detailed statistics.

Subtype correlation structure is weakened in the *Df(16)A^+/-* CA1

The diffuse IN subtype-specific alterations that we observed during spatial navigation for rewards
305   suggest a potential lack of coordinated inhibition in the *Df(16)A^+/-* hippocampal CA1. Microcircuit coordination is necessary for appropriate information flow and network stability[103,104]. Leveraging our method to image multiple IN subtypes simultaneously, we investigated whether the coordinated activity patterns in dorsal CA1 INs are impaired during spatial navigation tasks. We compared activity between subtypes within each genotype using Pearson's cross-correlation
310   analysis of cells to the average activity of a given cell class during a given trial for each task and genotype. In wildtype comparisons, IN subtypes exhibit highly correlated activity (even after normalizing for mouse velocity) and reflected known relationships between subtypes (Fig. 4A/B, left). For example, the CCKC subtype is comprised of both perisomatic-targeting basket cells (CCKBCs) and dendrite-targeting CCK+ INs[77,80,110]. CCKBCs exhibit uniquely anticorrelated
315   activity with velocity[77,95] , and as expected, we found that CCKCs in WT showed the lowest within-group and outside-group correlations of any IN type characterized in this study (Fig. 4A, left). Across both behavior tasks, most cell-type comparisons between genotypes suggested an overall decrease in correlated activity in *Df(16)A^+/-* INs (Fig. 4A/B, right). Particularly during the Reward Translocation task significant reductions in correlations between CCKC, PVBC, AAC, and SomC
320   subtypes portray a global disorganization of microcircuit activity, rather than implicating a specific subtype (Fig. 4B).

Figure 4.

Figure 4: Coordinated activity is diminished in the Df(16)A⁺/⁻ CA1 microcircuit. (A) Pairwise Pearson's correlation coefficient value matrix between cell types of ($\Delta F/F$) traces across the trial during the Random Foraging paradigm from WT (left), *Df(16)A⁺/⁻* (center) and the comparative differences between genotypes (right). (B) Pairwise Pearson's correlation coefficient value matrix between cell types of ($\Delta F/F$) traces across the Reward Translocation trials from WT (left), *Df(16)A⁺/⁻* (center) and the comparative differences between genotypes (right). *: P < 0.05. **: P < 0.01. ***: P < 0.001.

A probabilistic graphical model uncovers altered coactivity patterns between interneuron subtypes in *Df(16)A⁺/⁻* mice

A loss of coordinated activity in *Df(16)A⁺/⁻* INs hints at alterations in cell-cell connectivity or responses to a common input (hereafter referred to as *coactivity*). We analyzed the coactivity patterns between IN subtypes across trials, animals and behaviors to infer whether there are consistencies across biological replicates. Further, differences in consistency between different genotypes in the coactivity patterns between INs was used to suggest an increase or decrease in organization of the circuits that these INs were part of. For example, if the coactivity between SomC's and all other INs was consistent and predictable across most of the WTs but no longer predictable in *Df(16)A⁺/⁻ mice*, we would deduce that circuit organization in *Df(16)A⁺/⁻* mice was not driven by these cell types, and therefore relatively disorganized. To quantify all of these effects and signatures of coactivity between INs, we devised a probabilistic graphical model (Suppl. Fig. 6A, see methods) to model cell-types as a function of their co-activation patterns with all other IN subtypes in CA1 (Fig. 5A). The graphical representation of this model (Suppl. Fig. 6A) illustrates the hierarchical nature of the model, highlighting the dependencies between the different levels of interaction. For example, we model that each cell has a correlation value with all other cells in our observed dataset. Then we capture the coactivation of this cell with different cell-types as the

maximum absolute correlation value amongst all cells of the specified type. This captures the most likely coactivation partner of a cell within each of the IN subtypes (thus each cell is represented by a vector of K coactivations for K IN subtypes). We term this cell-to-cell-type coactivation (e.g. single cell vs. all IN subtypes). We then take averages of these cell-to-cell-type values for all cells within a cell-type to deduce the expected *cell-type-to-cell-type* co-activations (e.g. SomC vs. all IN subtypes). All these interactions, both *cell-to-cell-type* and *cell-type-to-cell-type* are modelled under a hierarchical Bayesian probabilistic model where we infer parameters of these distributions (e.g. mean and covariance of *cell-type-to-cell-type* co-activation patterns) to make quantitative and probabilistic statements about observed data. Using this model, we aimed to achieve two goals: 1) assess how co-activation profiles changed in mutant mice to infer connectivity alterations and 2) determine the predictability of the molecular identity of INs based on their interaction profiles with known IN subtypes.

To probe cell-type activity relationships during behavior, we visualized the average cell-type interactions (Ω) to reveal the structural-functional correlations within the hippocampal network. This notably recapitulated known mutual anticorrelated interactions such as those between CCKC and PVBC type INs during Random Foraging (Fig. 5B, upper left)[77,95,96]. In WT mice, most cell type relationships tended to be more coactive (or anti-coactive) than expected under the assumption that their activity was completely random during both behavioral tasks (Fig. 5B, upper and lower left). When we compared the relative coactivity patterns between genotypes, we observed that the directionality of coactivity changes was consistent between the behavior tasks suggesting a microcircuit-wide perturbation of cell-cell coordination or connectivity in the mutant mice (Fig. 5C). Notably, coactivity patterns were largely diminished during the Reward Translocation task (Fig. 5B, lower right) indicating that microcircuit coordination may be more severely compromised when there are higher demands on the circuit.

Altered cell type predictability infers altered circuit organization in *Df(16)A$^{+/-}$*

Neuronal cell-type is hypothesized to be a key determinant of neural function within a circuit[77,95,111]. We investigated whether neuronal identity could be inferred from the activity patterns of IN subtypes in local hippocampal CA1 circuits of WT and mutant mice. The separation of INs along principal component space (i.e. SomC and AAC along PC1, CCKC against all other INs along PC2) in the low dimensional projection of interaction parameters (Suppl. Fig. 6B) suggest that our model effectively captures features of cell interactions that facilitate cell type classification. To further assess the nature of these coordinated activity signatures, we attempted to decode IN cell types based on their coherence; if a cell type is predictive, we can infer that their coordinated activity is reliable. As stated before, we used the relative predictability of cell-types from their coactivation patterns as a surrogate of how organized a circuit was. Our reasoning is that an organized circuit has consistent coactivation patterns across biological replicates (e.g. different animals) and thus we can make accurate predictions of cell types from their coactivation patterns. In contrast, if a circuit is disorganized, then there may be low amount of consistency of co-activation patterns across different animals, and thus predictive models will suffer in out-of-sample generalizability. Our probabilistic model demonstrated robust classification for all five IN types tested (AACs, CCKCs, PVBCs, SomCs, IvC/NGFCs) in the WT with accuracy significantly above random chance during both Random Foraging (Fig. 5D, left) and Reward Translocation

390   tasks (Fig. 5E, left). In contrast, our results revealed both task-dependent and cell type-specific alterations in IN predictability in *Df(16)A$^{+/-}$* mutants. During Random Foraging, neither AACs nor CCKCs were predictable, although the latter could be attributed to lower numbers of CCKCs recovered in the mutants (Fig. 5D, right). Meanwhile, during the Reward Translocation task, all cell types remained significantly predictable, with PVBCs and IvC/NGFCs showing enhanced

395   predictability. This is expected for PVBCs, as we identified a reliable enhanced reward response unique to the mutants. The increased predictability in IvC/NGFCs, however, suggests a task-dependent functional change specific to their activity. The fact that cell types remain predictable in the mutants implies that the alterations in coactivation are consistent across animals, rather than arising from random circuit disorganization.

400   In conclusion, our coactivity analysis using probabilistic graphical modeling supports task-dependent alterations in coordinated activity across multiple IN subtypes (Fig. 4), indicating a broad reorganization of IN microcircuit dynamics in mutant mice.

Figure 5.

A

For cell 'A₁' coactivity with celltype 'B', we identify the max correlated (+ or -) 'B':

We then average all maximum correlation pairs for all 'A' cells with 'B':

B   *Wildtype*   *Df(16)A⁺ᐟ⁻*

C   *Df(16)A⁺ᐟ⁻ - Wildtype*

D   **Random Foraging**   *Wildtype*   *Df(16)A⁺ᐟ⁻*

E   **Reward Translocation Task**   *Wildtype*   *Df(16)A⁺ᐟ⁻*

Figure 5: Coactivity alterations are genotype and task dependent. (A) Schematic depicting how the model identifies the average maximum coactivation between a given pair of cells to estimate coactivity motifs. (B) Circuit representation of the cell-to-cell-type interactions of Wildtype (left) and *Df(16)A⁺ᐟ⁻* (right) interneurons and by behavior paradigm (top:

Random Foraging, bottom: Reward Translocation task), as determined by the probabilistic graphical model. Nodes represent different interneuron types, with colors corresponding to AAC (purple), CCKC (green), IvC/NGFC (blue), PVBC (yellow), and SomC (red). Edges connecting the nodes reflect the z-scores of the differences between the average correlations, with green edges indicating a correlation greater than expected by chance and red edges indicating less than expected. (C) Circuit representation of the difference in cell-to-cell-type interactions between mutant *Df(16)A*[+/-] and WT genotypes by behavior paradigm (top: Random Foraging, bottom: Reward Translocation task), as determined by the probabilistic graphical model. (D) Cell type classification accuracy using inferred cell types from *A* using our IN activity data for WT and *Df(16)(A)*[+/-] during Random Foraging (D) and reward translocation task (E). Light boxes indicate the null distribution of classification accuracy upon shuffling cell-type labels. n.s.: not significant. *: P < 0.05. **: P < 0.01. ***: P < 0.001. See Suppl. Table 1 for detailed statistics.

DISCUSSION

Our study provides the first comprehensive characterization of inhibitory local circuit dynamics in the murine hippocampus of mutant mice carrying a schizophrenia-predisposing genetic lesion. We discovered that mice carrying the 22q11.2 deletion, one of the strongest genetic risk factors for cognitive dysfunction and schizophrenia, exhibit widespread disruptions in hippocampal IN dynamics and coactivation patterns. By examining IN subtype-specific responses during distinct spatial navigation tasks, our study reveals distributed alterations at the microcircuit level (Table 1). These changes are more holistically represented through our approach and could not be fully captured by focusing solely on individual IN subtypes.

By monitoring IN activity dynamics during Random Foraging, we confirmed an overall positive velocity modulation of IN activity during locomotion in *both* WT and *Df(16)A*[+/-] mice. Despite the relative conservation of these fundamental functional characteristics of hippocampal IN activity[76,77,79,95,112–114], our findings identified significant alterations in the reward encoding of broad classes of INs within the *Df(16)A*[+/-] mice when compared to the WT littermates. Prior studies have implicated the role of specific IN subtypes, including SomCs and PVBCs[64,95,96,99,100], in reward modulation. Indeed, we identified PVBCs and SomCs as having particularly differential reward modulation during both random foraging and goal-oriented learning tasks, indicating pervasive alterations in the hippocampal reward processing circuit. Previously, we found that *Df(16)A*[+/-] CA1 pyramidal neurons exhibit reduced enrichment in the reward zone during a goal-oriented learning task with a fixed reward zone, and that these mice struggle to extinguish licking behavior at the old reward zone location licking behavior compared to WT littermates[43]. Thus, this enhanced and inflexible subtype-specific IN activity regulating reward responses may contribute to memory deficits and impaired reward-related representational dynamics of pyramidal cells observed in the *Df(16)A*[+/-] mice.

Further, our results also uncovered a population-level deficit in the spatial information encoded by INs in the *Df(16)A*[+/-] mice when compared to the WT littermates. Specifically, IN activity in the *Df(16)A*[+/-] mice carries significantly less spatial information implying a reduced spatially localized inhibitory influence over CA1 pyramidal cell ensembles[95,104,115–119]. While positive place field detection in INs was reduced, negative place fields were preserved, indicating differential roles for dynamic IN activity[96].

We identified large-scale relational networks of INs in *both* WT and *Df(16)A*[+/-] mice revealing complex, correlational relationships between IN subtypes with specifically reduced correlations in the mutants. Through conceiving a probabilistic graphical model for coactivity, we were able to take these observations a step further and uncovered significant alterations in the coactivity patterns among IN subtypes in *Df(16)A*[+/-] mice that implicate an uncoordinated microcircuit. Our model was also able to effectively predict several IN subtypes, highlighting a distinct and discernible interaction pattern among cell types in the WT mouse hippocampus. However, the predictability of AACs and CCKCs during Random Foraging was absent in *Df(16)A*[+/-] mice, suggesting a drift in microcircuit activity. This modeling approach lays the groundwork for future research to further explore the structural correlates of inferred connectivity patterns, potentially leading to a more comprehensive mapping of neural circuitry.

In the present study, we focused on major IN subtypes that directly target pyramidal cells in CA1. It remains to be determined whether the observed alterations in inhibitory subtype dynamics are the primary drivers of the disrupted pyramidal cell activity we previously reported[43] or, alternatively, these changes are a consequence of altered local recurrent or long-range excitatory inputs to INs[120–122]. Likewise, it will also be important to determine if disinhibitory circuits that inhibit other INs[64,95,123–125], including the subtypes investigated in the present study, also exhibit altered dynamics, perhaps playing a causal role in the observed alterations.

Studies in mouse models for schizophrenia and human patients have predominantly focused on PV-expressing INs, implicating reduced prevalence and activity of these cells in disorganized network activity[41,86,88,89,97,102,126–129]. Along these lines, destabilization in hippocampal neuronal dynamics analogous to the one described in Zaremba et al., 2017[43] has been subsequently recapitulated in hippocampal *in vitro* preparations from another 22q11.2 deletion model (*Lgdel/+* mice) and correlated with deficits in the synchronized CA1 neuronal network activity, including diminished spike correlation between neuronal pairs and reduced theta frequency oscillations[102]. Importantly, these network and hippocampal-dependent behavioral deficits were partially or fully restored by acute pharmacological or chemogenetic manipulations of PV+ IN excitability, underscoring the role of inhibitory regulation in maintaining local circuit stability in the hippocampus. However, recent studies also implicate dysfunction in other IN types, including SomCs, Neuropeptide-Y-expressing INs, and AACs in various disease-related cortical circuits[7,45,87,125,130–136]. Importantly, these subtypes directly interact with one another, influencing microcircuit functional output. In agreement with this body of work, our results also show that PV+ INs deficits alone do not account for all IN dysfunction associated with this high-risk schizophrenia locus. Diverse GABAergic INs express multiple genes encompassed within the *Df(16)A*[+/-] deletion (Supp. Fig. 7A). The pattern of enrichment differs from the one observed for glutamatergic neurons, suggesting that distinct genetic pathways may be affected in glutamatergic and GABAergic cell types (Supp. Fig. 7B/C). Based on expression profiles, a network-wide genetic perturbation in INs could cause widespread dysfunction and altered excitation-inhibition balance rather than causing purely subtype-specific effects.

Overall, while further research is needed to fully elucidate the precise mechanisms by which 22q11.2 deletions and other risk mutations contribute to cognitive dysfunction and schizophrenia,

it is increasingly clear that these effects are distributed across multiple IN subtypes within the brain (Table 1). This distribution affects both individual INs and ensembles of interacting IN cell types during cognitive operations in patients. Our findings suggest that genetic manipulation, modeling key aspects of schizophrenia risk such as 22q11.2 deletions, induces changes in the coactivity of local IN ensembles that are markedly more pronounced than alterations in individual INs. This implies that even seemingly small functional changes in individual INs can culminate in significant systems-level alterations when integrated into functional microcircuits.

Assessment	AAC	PVBC	SomC	IvC/NGFC	CCKC
Reward Response:	aberrant	aberrant	enhanced		
Spatial information:	reduced	reduced	reduced		
Reward response extinction:		delay	delay	aberrant	
Correlated activity:	task-dependent de-correlated activity				
Inferred connectivity:	task-dependent loss of co-activity patterns				
Cell type decoding:	task-dependent altered functionally inferred cell types				

Table 1. Summary of interneuron subtype-specific deficits.

Our study provides critical insights into how schizophrenia-predisposing mutations affecting cognitive dysfunction disrupt complex interactions among IN cell types in the mouse hippocampus during learning. It highlights a sophisticated interplay between disease-predisposing genetic lesions, the brain's dynamic functional patterns, and, eventually, the manifestation of cognitive symptoms. Moreover, our findings have broader implications for the predictive modeling of neuron types based on their functional interactions. This approach offers a novel framework for understanding how the brain's molecular structure and dynamic activity patterns are altered under pathological conditions. Future investigations will aid in expanding the molecular and structural correlates of the observed altered IN coactivity patterns, potentially leading to a more comprehensive understanding of how genetic liability leads to distorted maps of neural circuitry.

METHODS

RESOURCE AVAILABILITY

Lead Contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Attila Losonczy (al2856@columbia.edu).

Materials Availability
The study did not generate new unique reagents or mouse lines.

Data and Code Availability
Data and code reported in this paper are available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All experiments were conducted in accordance with NIH guidelines and with the approval of the Columbia University Institutional Animal Care and Use Committee. Experiments were performed with healthy, 2-5 month old, heterozygous adult male and female *Vgat-IRES-Cre* mice (Jackson Laboratory, Stock No: 016962, referred to as Vgat-Cre mice) that were crossed with hemizygous *Df(16)A$^{+/-}$* mice generated as previously described[98]. Mice were held in a 12-hour light/dark cycle vivarium and housed 2-5 mice in each cage. All experiments were conducted during the light portion of the cycle. Most mice used in the experiment were male (n = 9), with (n =5 WT, n = 4 *Df(16)A$^{+/-}$*) with n = 2 females (n = 1 WT, n = 1 *Df(16)A$^{+/-}$*). All wild-type mice were littermate controls of the *Df(16)A$^{+/-}$* mice.

METHOD DETAILS

Viruses
Cre-dependent recombinant adeno-associated virus (rAAV) for GCaMP7f (rAAV1-syn-FLEX-jGCaMP7f-WPRE, Addgene #1944820-AAV1, titer: ≥1:10^{13} vg/mL) were used to express GCaMP7f in the hippocampus of *Vgat-Cre* mice. The Cre-dependent recombinase enabled expression of Ca^{2+} sensor GCaMP7f in Vgat+ interneurons (IN)).

Viral injections and hippocampal window/headpost implant
During isoflurane anesthesia, *Df(16)A$^{+/-}$* Vgat-Cre mice were placed onto a stereotaxic surgery instrument. Meloxicam and bupivacaine were administered subcutaneously to reduce discomfort. The skin was cut along the midline to expose the skull, and a craniotomy was performed over the right hemisphere with a dental drill. A Nanoject injector was loaded with a sterile glass capillary filled with mineral oil and rAAV1-syn-FLEX-jGCaMP7f-WPRE and lowered into dorsal CA1 of the right hemisphere. Injections were made at (in mm): AP -2.2, ML -1.75, DV - 1.8, -1.6, -1.4, -1.2, -1.1, -1.0, -0.8 relative to bregma with 64 nl of virus injected at each DV level. The pipette was held within the brain for 5-10 minutes after the last injection and slowly raised from the brain. The

skin was sutured, and a topical antibiotic was applied to the suture site. Four days after viral injection, a circular cannula with a glass window was inserted into the cavity produced by craniotomy as previously described in ref. [120]. Similarly, as described above, mice were anesthetized with isoflurane and placed on a stereotaxic surgery instrument. Meloxicam and bupivacaine were subcutaneously given to the mice. The skull was exposed, and a 3 mm craniotomy was performed over the right hippocampus at AP -2.2, ML -1.75 relative to bregma, centered around the injection site. After removal of the dura, the cortex was slowly aspirated with negative pressure and the consistent application of ice-cold cortex buffer. When the horizontal fibers over CA1 were exposed, aspiration was stopped, and a sterile collagen membrane was applied to stop bleeding. A 3 mm circular cannula with a glass window was placed into the craniotomy and glued into the site with Vetbond. A headpost was then attached to the posterior of the skull with layers of dental cement. 1.0 mL of saline was subcutaneously given to each mouse, while the mice recovered in their home cage with a heat pad. All mice were monitored for 3 days of post-operative care until behavior training began.

Behavioral training and paradigms
Following the post-op recovery period, mice were acclimated to handling and head-fixation for a few days, while being water restricted to 85-90% of their original weight. Mice were trained to run on a two meter fabric, cue-free belt with random rewards that were automatically dispensed at a decreasing number of random locations. Mice were trained for ~1 week on the cue-free belt and then were transitioned to the fixed reward task.

Random Foraging Task
Mice were imaged on a two-meter familiar context belt for 1 day with 3 random rewards distributed across the length of the belt, randomized every lap. On day 2 of imaging, a novel belt was introduced with the same random reward paradigm. The belt consisted of four fabric ribbons with different textures and a variety of tactile cues: colored tape, foam strips, cotton pom poms, etc. The novel belt had distinct fabric materials and distinct tactile cues.

Reward Translocation Task
Mice were trained on a two meter cue-rich belt with a fixed reward zone at the end of the belt for 3 or 4 days leading up to imaging. The belt consisted of four fabric ribbons with different textures and a variety of tactile cues: colored tape, foam strips, cotton pom poms, etc. The following day the mice were imaged with the familiar reward zone location. On the second and third day of imaging, the reward zone was relocated to a new reward zone location at the center of the belt. The reward size was larger in this task than in random foraging task to account for the decreased number of rewards per lap.

Two-photon Ca^{2+} imaging
After the mice were acclimated and trained in the respective tasks, a high-resolution structural Z-stack was captured 24 hours prior to imaging, while the mouse was anesthetized with isoflurane. The reference Z-stack was used to identify the 3D position of the GCaMP-positive neurons for access by the custom-modified AOD (acousto-optic deflector) microscope (Femto3D-ATLAS). The AOD microscope was used as previously described [95,96,103].

A water immersion objective (x16 Nikon CF175) was used to focus at the dorsal CA1 pyramidal layer and then fixed in position. A tunable two-photon laser (Coherent Ultra II) was adjusted to wavelength = 920 nm. The high-resolution structural Z-stack was captured in CA1 from ~100um
610   below to glass to ~600 μm to span the entirety of CA1 from the *stratum oriens* to the *stratum lacunosum-moleculare* with an 800x800 pixel capture with a resolution of 1.25 μm per pixel with a 4 μm step size. Laser power and photomultiplier tube (PMT) detectors (GaAsP, H10770PA-40, Hamamatsu) were compensated in power and gain throughout the Z-stack capture. After the Z-stack was completed, the mouse was returned to its home cage and given 24 hours to recover
615   before functional imaging.

Using the integrated software from Femtonics (MES), the Z-stack was examined, and 100-150 INs were manually selected to generate a spreadsheet of the x-y-z positions at the center of each respective cell. These x-y-z positions served as the regions of interest (ROIs) that were then used
620   on subsequent days by the AOD microscope to find the position of the corresponding interneurons. For each day of functional imaging in each mouse, the identical field of view was identified within the CA1 pyramidal layer, and then the ROIs were loaded using the software for AOD 3D imaging. Each imaging session was taken for 10 minutes, with 5-10 Hz for each recording experiment (frame rate depended on the number of ROIs, ROI size, imaging resolution, and
625   scanning speed).

Immunohistochemistry

Mice were perfused with 20mL of Phosphate Buffered-Saline followed by 20 mL 4%
630   paraformaldehyde (PFA, Electron Microscopy Sciences). The brains were stored in 4% PFA overnight and washed 3x5 min with PBS the next day. The brains were then sectioned for 75 μm transverse sections of the hippocampus on a vibratome and rinsed 3x15 min with PBS prior to immunohistochemical staining procedure. Histological tissue prep was performed as previously described[95,96]. In brief: sections were permeabilized 2x20 min with 0.3% Triton X-100 (Sigma-
635   Aldrick), and subsequently blocked with 10% Normal Donkey Serum (Jackson ImmunoResearch) in PBST (PBS with 0.3% Triton X-100). After 45 minutes, the sections were incubated in PBS (containing 3 first-round primary antibodies, see below) for one hour at room temperature and then 2 days at 4°C with agitation. The sections were rinsed 3x15 min with PBS to wash out excess primary antibodies, and then, for two hours at room temperature, incubated in a PBS solution of
640   secondary antibodies tagged for fluorescence (Table 2). Sections were then rinsed 5x15 min in PBS at room temperature and mounted onto rectangular glass slides with ThermoFisher Scientific aqueous mounting medium. After 1 hour of drying at 4°C, the slides were imaged. After imaging, PBS was used to remove coverslips, and sections were rinsed 3x15 min in PBS for round two of staining. The sections were blocked with 10% Normal Donkey Serum in PBST for 45 minutes
645   before the addition of the round two primary antibodies. After, secondary staining protocol (now using round two secondary antibodies) was followed as completed in the first round. staining for CCK, SOM and NPY first led to stronger expression of fluorescence for those markers, opposed to staining for any of the three markers second (Table 2).

650 *Antibody information and staining strategy*

Cell Marker	Round	Primary Antibody (dilution)	Secondary Antibody (dilution)	Catalog #
Cholecystokinin (CCK)	1	rabbit anti-proCCK (1:500)	donkey anti-rabbit DyLight 405 (1:300)	Frontier Institute CCK-pro-Rb-Af350
Somatostatin (SST)	1	rat anti-somatostatin (1:500)	donkey anti-rat Alexa 568 (1:300)	Millipore MAB354
Neuropeptide-Y (NPY)	1	sheep anti-NPY (1:500)	donkey anti-sheep F(ab)2 Alexa 647 (1:300)	Abcam ab6173
Parvalbumin (Pvalb)	2	chicken anti-PV (1:5000)	donkey anti-chicken DyLight 405 (1:300)	*Gift from Susan Brenner-Morton*
SATB1	2	rabbit anti-SATB1 (1:1000)	donkey anti-rabbit Alexa 647 (1:300)	Abcam ab70004

Table 2. Antibodies used in this study

Assignment of subtype identity based on immunostaining

655 Interneuron subtypes were assigned based on our previously published classification strategy[95].

Axo-axonic cells (AACs):
Immunopositive for Pvalb and immunonegative for SATB1, SST, NPY, & CCK.

660 *Parvalbumin Basket Cells (PVBCs):*
Immunopositive for Pvalb and SATB1, immunonegative for SST, NPY, & CCK.

Somatostatin-expressing cells (SomCs):
Immunopositive for SST, with or without Neuropeptide Y (NPY) and SATB1, and immunonegative
665 for Pvalb and CCK.

Ivy / Neurogliaform cells (Ivy/NGFCs):
Immunopositive for Neuropeptide Y, with or without SATB1, and immunonegative for SST, Pvalb, and CCK.
670
Cholecystokinin-expressing cells (CCKCs):
Immunopositive for proCCK, immunonegative for SST, PV, NPY and SATB1.

Confocal imaging
675 The NikonA1 confocal microscope used to take multi-channel fluorescence images (405 nm, 488 nm, 561 nm, and 640 nm excitation for the four far-red, blue, green and red colored channels) of labeled tissue sections. Image capturing yielded 2048 x 2048 pixel z-stacks of the entire depth of

the section (with 3 microns between each image of a given stack). Z-stacks were viewable via Fiji (Image J).

Registration of confocal images to *in vivo* z-stacks and identification of immunopositivity/negativity

Unique groupings of cells from *in vivo* z-stacks were used as visual fiducial markers to manually register *ex vivo* 75 μm z-stack images in ImageJ, section by section. Care was taken to note the

685 orientation of the *ex vivo* sections regarding the *in vivo* z-stack images. The coordinates taken from the AOD software when selecting ROIs for imaging against the *in vivo* z-stack were used to identify ROI ID numbers in the registered immunohistochemically stained images.

Immunohistochemical marker density quantification with CATT

690 A novel Cell Analysis/Typing Tool (CATT) designed by E.V., was used to identify positive cells in confocal fluorescence microscopy images. The code uses the entry of pixel dimensions to generate an image that combines the four fluorescence color channels of an original section. A region of interest (ROI) was drawn to delineate distinct CA1 layers for the program to focus on counting cells within a certain hippocampal layer. Given the ROI and pixel dimensions, the

695 program outputs the volume, as well as an image displaying all the IHC markers across the channels relative to the green channel (GCaMP7+ signal). Each positive cell marker was manually counted and divided by the ROI volume to derive positive marker density for a given ROI.

Behavior analysis

Licking behavior

For each animal during each experiment, the two meter belt was divided into 100 bins (~2cm per bin). The number of licks per bin per experiment were visualized to create licking curves. The proportion of licks that occur within the reward zone was also computed.

Reward zone occupancy and latency post-reward

Each animal's position per experiment was also digitized into 100 equal spatial bins representing the 2-meter belt. The occupancy time within the reward zone was quantified and compared between the genotypes. Furthermore, the time spent in the reward zone after the reward was no

710 longer dispensed was computed and defined as the latency.

Imaging analysis

Pre-processing of Ca^{2+} imaging data

715 Raw movies of each individual cell were independently motion-corrected using the whole-frame cross-registration implemented in the SIMA package[137]. ROIs were then hand-drawn over each imaged cell. Fluorescence was extracted from each drawn ROI using SIMA. $\Delta F/F$ was then computed based on a 1st percentile baseline value over a 30-second rolling window. The $\Delta F/F$ traces were then smoothed using an exponential filter. For analyses involving velocity-regressed

720 traces, each animal's velocity was smoothed with a Hanning filter, and then covariate linear

regression was applied to each fluorescence trace where the smoothed velocity was regressed out.

Velocity correlation

725 For each cell, velocity correlation was computed by taking Pearson's correlation coefficient between the ΔF/F trace and animal's smoothed velocity with a -5 sec to +5 sec rolling lag. The highest absolute value of the correlation coefficients in this time window was taken as the correlation coefficient for the respective cell.

730 ### *Run-Start/Run-Stop*

Based on a velocity cut-off of 0.2 cm/s, run-start and run-stop events were observed throughout the imaging session. Each run-start and run-stop event was at least 10 imaging frames (1-2 seconds) and nearby events within 20 frames (3-5 seconds) were merged. Peri-event time averages were generated for each animal. The response magnitude for each cell was computed

735 by taking the difference between the mean post-event ΔF/F and the mean pre-event ΔF/F.

Peri-Event Time Averages (PETA)

For salient stimuli such as run-start/run-stop events and reward distribution events, a PETA was constructed around each event respectively. These events were z-score and then averaged for

740 each respective cell, and the z-scored averages were visualized using a heatmap or an averaged line-plot. A Post-Pre metric was calculated based on the z-scored averages as a measure for event modulation.

Position decoding

745 A linear classifier support vector machine (SVM) was trained on the tuning curves of each respective cell, and the decoding error was visualized. The SVM was cross-validated by training the classifier on (n-1) laps and evaluating on the held-out lap, and this process was repeated such that every lap was utilized as a test lap at least once. Furthermore, the SVM was bootstrapped with different numbers of input cells and input laps averaged by animal to compare the position

750 decoding efficacy between genotypes.

Spatial tuning curve

The 2-meter belt was divided into 100 bins (~2cm per bin). Spatial tuning curves for each cell during an experiment were generated by averaging the velocity regressed ΔF/F (as described

755 above in preprocessing of Ca2+ imaging data) in each bin when the animal was running over 5 cm/s.

Peak place field calculation

The spatial tuning curves were shuffled 1000 times by circularly rotating the velocity-regressed

760 ΔF/F tuning curves around the 100 spatial bins. Cells were classified as being spatially selective (i.e., having a place field) if the cell exceeded the 95th percentile (or was under the 5th percentile for negatively selective cells) with at least 10 sequential bins. The number of bins over or under the 95th and 5th percentile curves were visualized and a cutoff of 30 consecutive bins was used for the figures as it represented the approximate 75th quartile for place field width.

Spatial information

Using the spatial tuning curve, a spatial information score was computed following the methodology described by Skaggs et al 1992[108]. While the method was originally conceived for spiking data, we applied it onto the spatial tuning curve from Ca^{2+} data as a proxy for information
770   content.

Reward Modulated Activity

For goal-directed experiments, the peri-event time averages were constructed around each reward event. The 2-meter belt was divided into 100 bins (~2cm per bin). Tuning curves were
775   generated by averaging the $\Delta F/F$ in each bin for both regular $\Delta F/F$ and velocity-regressed $\Delta F/F$.

Reward Modulation Index

A reward modulation index (RMI) was computed as described in refs. [95,96]. Briefly, we compared the difference of the average activity 10 cm preceding the reward zone and the average activity
780   on the remainder of the belt, normalized by the sum of the average activity over the two segments.

Pairwise Pearson's correlation analysis

Pairwise Pearson's correlation coefficients were computed for every pair of cells within a given experiment/trial. The average coefficient value for each subtype was then calculated per trial pair
785   and averaged across mice.

Multivariate ridge regression analysis

A multivariate ridge regression model was trained on the $\Delta F/F$ trace of Ca^{2+} activity (velocity not regressed) for each cell with three predictors: velocity, position, licking. Each predictor was
790   normalized to a 0-1 scale by using a minimum-maximum scaler. Given the inherent relationship between these variables, we used a ridge regression model with a cross-validated alpha value to limit the interaction between the predictors.

Probabilistic Graphical Model Framework
795

Model design

We designed a probabilistic graphical model to evaluate functional interactions between neurons, represented by the absolute correlation between any two cells i and j, denoted as $x_{i,j}$. Each cell i was associated with a cell type c_i, drawn from K distinct categories. The prior probability of cell i
800   belonging to cell type k was represented by $p(c_i = k) = \pi_k$. For each experimental session, the recorded cells were randomly partitioned into two subsets: 80% were allocated to the training set and the remaining 20% to the test set. This split was performed to develop and assess the predictive model in a controlled manner. The labels for the test cells were concealed during the model training phase and were only revealed during the validation stage to verify the accuracy of
805   the model's identification predictions.

Parameterization of cell interactions

Interactions between each cell and others of a given type k were modeled using a gamma distribution with parameters $\theta_{i,k}$, comprised of shape $\alpha_{i,k}$ and rate $\beta_{i,k}$. Here shape $\alpha_{i,k}$ and rate $\beta_{i,k}$ quantify how correlations between a cell of type k to individual cell i are distributed. For example, if all (or most) cells from type k is highly correlated with cell I both shape $\alpha_{i,k}$ and rate $\beta_{i,k}$ are going to be high values. Conversely, if most cells from cell type k do not interact with cell i, the shape parameter $\alpha_{i,k}$ is going to be close to 0. This parameterization was chosen to reflect the unique dynamics of interaction between cell i and cell type k, capturing the diversity and intensity of these inter-cell-type interactions and modeling the interactions as a random distribution to make the model fully Bayesian.

Multivariate distribution of interaction parameters

The full interaction profile of cell i across all K cell types was modeled using a multivariate Gaussian distribution, conditional on the cell type c_i. This distribution is defined by a mean vector μ_a and a covariance matrix Σ_a, specific to cell type a, and is denoted as $p(\theta_{i,1},...,\theta_{i,K}|c_i = a) \sim N(\Omega_a)$. This approach captures the ensemble of interactions that a cell of type a typically engages in with the spectrum of other cell types within the circuit.

Inference and Bayesian approach

We infer model parameters using Bayesian inference and computing the maximum a posteriori estimates of cell type c_i for a subset of cells $\{c_j\}_{j \in L}$, where L is the set of indices corresponding to known cell types. This process leveraged both the prior probabilities encoded in the model and the observed cell-to-cell interactions to probabilistically assign cell types to the remaining cells in the circuit. These analyses were performed for $Df16(A)^{+/-}$ and WT mice, separately.

Neuronal identity prediction

Classification validation

Our model was validated through its ability to classify cell identity based on functional correlation patterns. Model validation was performed by comparing the predictive accuracy of our probabilistic model against a null distribution. The null distribution was generated by shuffling cell type labels, thereby preserving the overall structure of the data while eliminating any true correlation between cell types and their functional interaction patterns. This approach allowed us to control for the possibility that any predictive success of the model could be due to chance. Predictive accuracies of our model were then contrasted with those from the null distribution to affirm the model's capability to identify cell types beyond random assignment. To assess the robustness of our probabilistic model, we conducted 31 independent runs, each initialized with a different random seed.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses

Subtype comparisons were conducted using one-way ANOVA tests with Tukey's range tests with correction for multiple testing if appropriate. For comparisons across genotypes, unpaired t tests were utilized if the data followed a normal distribution, assessed using the Shapiro-Wilk test.

Otherwise, for non-normally distributed comparisons, the Mann-Whitney U test was used. The Kolmogorov–Smirnov test was used when comparing CDFs of continuous data. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$. n.s. is not significant. All data analysis and visualization were done with custom software built on Python version 2.7.15, 3.8.11, and MATLAB R2023b.

Acknowledgements

S.A.H. is supported by the Burroughs Wellcome Fund. A.L. is supported by NIMHR01MH124047, NIMHR01MH124867, NINDSR01NS121106, NINDSU01NS115530, NINDSR01NS133381, NINDSR01NS131728, NIARF1AG080818. E.V. is supported by NIMHK99MH128772. J.A.G. is supported by NIMHR01MH124047 and NIMH R01MH097879. We thank Tristan Geiller, Bert Vancura for consultation on the analysis, and Erica Rodriguez for the mouse schematic in Fig. 1A.

Author Contributions

S.H., B.R., J.A.G., and A.L. conceived the study and wrote the manuscript. S.H. and B.R. performed experiments and analyzed the data with assistance from M.E.C.P., A.T., and H.A.. S.H. and E.V. conceived the coactivity and CATT analyses, which was developed and implemented by E.V., M.E.C.P., and H.A..

Supplementary Table 1: Statistical detail summary.

Figure	Statistics
1L	*AAC*: axo-axonic cells Mann-Whitney U test: P = 0.0017, *PVBC*: parvalbumin basket cells Mann-Whitney U test: P = 1.45e-5, *SomC*: Somatostatin-expressing cells Mann-Whitney U test: P = 0.012, *IvC/NGFC*: Ivy/Neurogliaform cells Mann-Whitney U test: P = 0.492, *CCKC*: cholecystokinin cells Mann-Whitney U test: P = 0.306. For Mann Whitney U tests: median ± inter-quartile range.
2C	AAC: Mann-Whitney U test: P = 0.136, PVBC: Mann-Whitney U test: P = 0.002, SomC: Mann-Whitney U test: P = 0.015, IvC/NGFC: Mann-Whitney U test: P = 0.068, CCKC: Mann-Whitney U test: P = 0.175.
2G	KS test: P = 1.156e-11.
2H	KS test: P = 2.055e-4.
2I	AAC: Mann-Whitney U test: P = 0.175, PVBC: Mann-Whitney U test: P = 2.052e-5, SomC: Mann-Whitney U test: P = 4.944e-12, IvC/NGFC: Mann-Whitney U test: P = 4.35e-6, CCKC: Mann-Whitney U test: P = 0.102.
2K	AAC: Mann-Whitney U test: P = 0.175, PVBC: Mann-Whitney U test: P = 2.052e-5, SomC: Mann-Whitney U test: P = 4.944e-12, IvC/NGFC: Mann-Whitney U test: P = 4.35e-6, CCKC: Mann-Whitney U test: P = 0.102. For Mann Whitney U tests: median ± inter-quartile range.
3A	WT: 16.4 ± 0.7 bins, n = 6; *Df(16)A$^{+/-}$*: 20.7 ± 0.9 bins, n = 5; Mann-Whitney U test: P = 0.004.
3B	10 cells: Mann-Whitney U test: p = 2.48e-13, 30 cells: Mann-Whitney U test: p = 3.90e-33, 50 cells: Mann-Whitney U test: p = 5.22e-34, 70 cells: Mann-Whitney U test: p = 2.07e-34, 90 cells: Mann-Whitney U test: p = 1.64e-31.
3C	5 laps: WT: Mann-Whitney U test: p = 4.12e-28, 10 laps: WT: Mann-Whitney U test: p = 1.18e-32, 15 laps: WT: Mann-Whitney U test: p = 4.36e-34.
3D	Comparison across animals: WT: 0.039 ± 0.011, n = 6; *Df(16)A$^{+/-}$*: 0.022 ± 0.017, n = 5; Mann Whitney U test: P = 0.0549.
3F	WT: 0.89 ± 0.13, n = 11 trials across 6 mice; *Df(16)A$^{+/-}$*: 0.72 ± 0.21, n = 8 trials across 5 mice; Mann-Whitney U test: P = 0.04.
3G	WT: 0.39 ± 0.14, n = 11 trials across 6 mice; *Df(16)A$^{+/-}$*: 0.16 ± 0.11, n = 8 trials across 5 mice; Mann-Whitney U test: P = 0.001.
3H	0 fields: Mann-Whitney U test: P = 0.0071, 1 field: Mann-Whitney U test P = 0.0071, 2 fields: Mann-Whitney U test: P = 0.0549.
3I	*AAC*: Axo-axonic cells Mann-Whitney U test: P = 0.00027, *PVBC*: parvalbumin basket cells Mann-Whitney U test: P = 0.122, *SomC*: Somatostatin-expressing cells Mann-Whitney U test: P = 0.0028, *IvC/NGFC*: Ivy/Neurogliaform cells Mann-Whitney U test: P = 0.058, *CCKC*: cholecystokinin cells Mann-Whitney U test: P = 0.285. WT Cell Numbers: AAC = 62 cells, SomC = 132 cells, PVBC = 46 cells, CCKC = 32 cells, IvC/NGFC = 73 cells; *Df(16)A$^{+/-}$* cell numbers: AAC = 34 cells, SomC = 125 cells, PVBC = 51 cells, CCKC = 14 cells, IvC/NGFC = 47 cells
3J	Day 1: Mann Whitney U test: statistic = P = 2.327e-7, Day 2: Mann Whitney U test: P = 0.172, Day 3: Mann Whitney U test: P = 0.010
3K	WT: 0.061 ± 0.038; *Df(16)A*: 0.022 ± 0.019; Mann-Whitney U test: P = 0.041.
4D/E	P-values of WT for Random Foraging task: AAC (P = 3.48e-27***), PVBC (P = 3.91e-13***), SomC (P = 1.71e-43***), IvC/NGFC (P = 4.27e-07***), CCKC (p = 2.56e-23***). P-values of *Df(16)A$^{+/-}$* for Random Foraging task: AAC (P = 3.94e-01, n.s.), PVBC (P = 1.47e-20***), SomC (P = 1.25e-32***), IvC/NGFC (P = 2.27e-04***), CCKC (P = 1.43e-01, n.s.). P-values of Wildtype for Reward Translocation Task: AAC (P = 8.76e-49***), PVBC (P = 1.58e-23***). SomC (P = 3.46e-53***). IvC/NGFC (P = 8.23e-15***). CCKC (P = 2.16e-40***). P-values of *Df(16)A$^{+/-}$* for Reward Translocation Task: AAC (P = 2.52e-22***). PVBC (P = 7.21e-31***). SomC (P = 5.72e-41***). IvC/NGFC (P = 2.66e-29***). CCKC (P = 1.49e-24***).

Boxplots show the 25th, 50th (median), and 75th quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles, respectively. Outliers are defined as values extending beyond the whisker ranges. For Welch's t tests: mean ± sem. For Mann Whitney U tests: median ± inter-quartile range. n.s.: not significant. *: P < 0.05. **: P < 0.01. ***: P < 0.001.

Supplemental Figure 1. 3D-AOD imaging and molecular characterization strategy for interneuron subtypes. (A) Summary distribution plots of all cells imaged in this study for each genotype based on imaging coordinates superimposed across all animals (n = 6 WT, n = 5 *Df(16)A*$^{+/-}$ mice) Dimensions are in microns (800 μm x 800 μm x 400 μm). (B) Cell type number totals averaged per genotype (n = 6 WT, *n* = 5 *Df(16)A*$^{+/-}$ mice) (Mann-Whitney U Test, p > 0.05). (C) Distribution of total cells identified, ordered by cell type, across all layers of CA1. *s.l.m.: stratum lacunosum moleculare; s.r.: stratum radiatum; s.p.: stratum pyramidale; s.o.: stratum oriens.* Black and gray bars represent cells that could not be identified (D) Quantification of total marker density for all immunohistochemical markers used (unpaired t-tests with Welch's correction (P > 0.05)). (E) Quantification of the density of cholecystokinin+ cells within each layer of CA1. *CCK*: Cholecystokinin. 6-7 serial 75 μm sections were stained and quantified for CCK+/GCaMP+ cell signal across the four layers of CA1 in n = 6 WT n = 6 *Df(16)A*$^{+/-}$ mice (unpaired t-tests with Welch's correction (P > 0.05)). (F) Quantification of the density of SATB1+ cells within each layer of CA1. 6-7 serial 75 μm sections were stained and quantified for SATB1+/GCaMP+ cell signal across the four layers of CA1 in n = 6 WT, n = 6 *Df(16)A*$^{+/-}$ mice (unpaired t-tests with Welch's correction (P > 0.05)). (G) Quantification of the density of parvalbumin+ cells within each layer of CA1. *PV*: parvalbumin. 6-7 serial 75 μm sections were stained and quantified for PV+/GCaMP+ cell signal across the four layers of CA1 in n = 6 WT, n = 6 *Df(16)A*$^{+/-}$ mice (unpaired t-tests with Welch's correction (P > 0.05)). (H) Quantification of the density of somatostatin+ (*Som*) cells within each layer of CA1. 6-7 serial 75 μm sections were stained and quantified for Som+/GCaMP+ cell signal across the four layers of CA1 in n = 6 WT n = 5 *Df(16)A*$^{+/-}$ mice (unpaired t-tests with Welch's correction (P > 0.05)). (I) Quantification of the density of NPY+ cells within each layer of CA1. 6-7 serial 75 μm sections were stained and quantified NPY+/GCaMP+ cell signal across the four layers of CA1 in n = 6 WT, n = 6 *Df(16)A*$^{+/-}$ mice (unpaired t-tests with Welch's correction (P > 0.05)). *NPY*: Neuropeptide Y. *n.s.*= not significant.

Supplemental Figure 2.

Supplemental Figure 2. Locomotion related interneuron activity is preserved in Df(16)A$^{+/-}$ mice. (A) Licking proportion of WT (black) and *Df(16)A$^{+/-}$* (brown). There is no significant difference in lick location (left) or total licks (right) across the belt during Random Foraging in WT (*n* = 6) or *Df(16)A$^{+/-}$* (*n* = 5) depicted by box-and-whisker plot, WT: 1534 ± 190 licks, *Df(16)A$^{+/-}$*: 1608 ± 291 licks, Welch's t-test, p = 0.838. B) Average velocity in cm per second plotted by location on the belt of WT (black) and *Df(16)A$^{+/-}$* (brown) mice (left). There is no significant difference in average velocity during Random Foraging in WT (n = 6) or *Df(16)A$^{+/-}$* (n = 5) depicted by box-and-whisker plot (right), WT: 13.7 ± 0.7 cm/s, *Df(16)A$^{+/-}$*: 15.2 ± 1.6 cm/s, Welch's t-test, P = 0.405. (C) Total laps across mice within genotype (WT: 22.6 ± 1.5 laps, *Df(16)A$^{+/-}$*: 20.6 ± 2.2 laps, Welch's t test, t = 0.74, P = 0.47). (D) Average percent occupancy on the belt by position per genotype (Mann-Whitney U test, P = 0.3). (E) Pearson correlation coefficients of interneuron activity and velocity for all tested interneuron subtypes (WT: n = 97 AACs, 53 PVBCs, 194 SomCs, 115 IvCs/NGFCs, and 44 CCKCs from n = 6 mice, *Df(16)A$^{+/-}$*: 46 AACs, 87 PVBCs, 181 SomCs, 72 IvCs/NGFCs, and 14 CCKCs from n = 5 mice). WT: one-way ANOVA (p < 10^{26}), *Df(16)A$^{+/-}$*: one-way ANOVA (P < 10^{10}) with post hoc Tukey's range test corrected for multiple testing. (F) Peri-event time averages (PETAs) of averaged z-scored traces with a range between -1.75 and 1.75 z-scores during Run-Start and Run-Stop events. Left: WT; Right: *Df(16)A$^{+/-}$*. (G) Average Post-Pre PETA responses to Run-Start (Left) and Run-Stop (Right) events. (H) Ridge Regression Model shows velocity as a significant predictor of IN GCaMP-Ca^{2+} activity. Wilcoxon: WT: 0.257 ± 0.007, P = 4.68e-141, *Df(16)A$^{+/-}$*: 0.240 ± 0.007, P = 3.60e-117. Boxplots show the 25th, 50th (median), and 75th quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles, respectively. Outliers are defined as values extending beyond the whisker ranges. For Welch's t tests and Wilcoxon test: mean ± sem. For Mann Whitney U tests: median ± inter-quartile range. n.s.: not significant. ****: P < 0.0001.

Supplemental Figure 3. Mutant mice exhibit protracted reward response behavior. (A) Total lick distribution across position bins on familiar trial Day 1 (top), translocation Day 2 (middle) and translocation Day 3 (bottom) of recording. (B) Total licks across trials (WT: 959 ± 415 licks; *Df(16)A⁺/⁻*: 1895 ± 1364 licks; Mann-Whitney U test: P = 0.0064). (C) Proportion of time spent across position bins on familiar trial Day 1 (top), translocation Day 2 (middle), and translocation Day 3 (bottom) of recording. (D) Proportion of time spent in the reward zone across trials (averaged across trials WT: 0.228 ± 0.188; *Df(16)A⁺/⁻*: 0.328 ± 0.108; Mann-Whitney U test: P = 0.02). (E) Average laps run by mouse per trial averaged across animals (WT: 25.5 ± 16.5 laps; *Df(16)A⁺/⁻*: 25.5 ± 15.5 laps; Mann-Whitney U test: P = 0.40). (F) Average velocity run by mouse per trial averaged across animals: WT: 14.84 ± 10.01 cm/s; Df(16)A⁺/⁻: 18.61 ± 4.32 cm/s; Mann-Whitney U test: P = 0.24. (G) Average velocity by position on the belt for Day 1 (top), Day 2 after translocation of the reward (middle), and Day 3 (bottom). Boxplots show the 25th, 50th (median), and 75th quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles, respectively. Outliers are defined as values extending beyond the whisker ranges. For Welch's t tests: mean ± sem. For Mann Whitney U tests: median ± inter-quartile range. n.s.: not significant. *: P < 0.05. **: P < 0.01. ***: P < 0.001.

Supplemental Figure 4: Multiple interneuron subtypes exhibit enhanced reward related activity in mutants. (A) Individual cell responses to reward during reward translocation task, Left: WT, Right: $Df(16)A^{+/-}$ (B) Averaged z-scored cell responses with a range between -1.5 and 1.5 z-scores to reward onset during reward translocation task, Left: WT, Right: $Df(16)A^{+/-}$ (C) Average responses of each cell type in the preceding 40 cm of the belt to the *new* reward location in the WT (left) and $Df(16)A^{+/-}$ (right) on Day 2. (D) Reward zone modulation index (*RMI,* see methods under Reward Modulation Index) for SomCs and PVBCs at the old reward zone during the first day. (SomCs: Kolmogorov-Smirnov (KS) test: P = 0.282, PVBCs: KS test: P = 0.076.) (E) RMI for SomCs and PVBCs at the old reward zone on second translocation day (Day 3). (SomCs: KS test: P = 3.812e-6, PVBCs: KS test: P = 6.888e-4.) (F) Reward zone modulation index for all cells at the new reward zone on the first day of reward translocation. (AAC: Mann-Whitney U test: P = 7.742e-4, PVBC: Mann-Whitney U test: P = 0.153, SomC: Mann-Whitney U test: P =0.049, IvC/NGFC: Mann-Whitney U test: P = 0.254, CCKC: Mann-Whitney U test: P = 0.409) *: P < 0.05. **: P < 0.01. ***: P < 0.001. *n.s.* = not significant.

Supplemental Figure 5.

950   **Supplemental Figure 5: Negative place fields are not reduced in *Df(16)A⁺/⁻* compared with WT.** (A) Positive place
field widths of detected place fields using Peak Place method. Averaged across animals: Mann-Whitney U test: P =
0.464. (B) Negative place fields detected between genotypes. WT: 0.79 ± 0.19, *Df(16)A⁺/⁻*: 0.61 ± 0.29 Mann-Whitney
U test:, P = 0.087. (C) Number of all detected place fields per cell by genotype. 0 place fields: Mann-Whitney U test: P
= 0.0829, 1 place field: Mann-Whitney U test: P = 0.4576, 2 place fields: Mann-Whitney U test: P = 0.1205, 3 place
955   fields: Mann-Whitney U test: P = 0.0126, 4 place fields: Mann-Whitney U test: P = 0.0549. (D) Proportion of positive
place cells during reward translocation task. (Student's T-test: P = 0.6288). Boxplots show the 25th, 50th (median), and
75th quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles,
respectively. Outliers are defined as values extending beyond the whisker ranges. For Welch's t tests: mean ± sem.
For Mann Whitney U tests: median ± inter-quartile range. *n.s.* = not significant.

Supplemental Figure 6: A probabilistic graphical model to infer coactivation patterns. (A) A schematic of the probabilistic graphical model of cell type-to-cell type interactions. The model begins with a cell type prior (π_k) for each of the K possible types denoting probability of assignment by chance. Each cell (i) is then probabilistically assigned a cell type (c_i) using maximum a posteriori estimation, weighed by the prior. Parameters $\theta_{i,k}$ reflect the unique interaction

965 dynamics of cell i with cells of type k and are derived from cell-to-cell correlations ($x_{i,j}$). These parameters are informed by both the observed cell types ($c_{ij} = k$) and the overarching cell-type-to-cell-type parameters (Ω_{ci}). See Probabilistic Graphical Model Framework section for more detail in methods section. (B) Left: projection of cell-to-cell-type interaction parameters ($\theta_{i,k}$) for all cells with cell types labeled in the training set. Cell types are denoted with the following colors: AAC (purple), CCKC (green), IvC/NGFC (blue), PVBC (yellow), and SomC (red). Right: projection of cell-to-cell-type

970 interaction parameters ($\theta_{i,k}$) into a low-dimensional subspace for five different hippocampal IN subtypes in the testing set.

Supplemental Figure 7: Expression profile of genes residing within the deleted locus in *Df(16)A+/-* mice in defined hippocampal (A) GABAergic interneuron subtypes and (B) Glutamatergic subtypes from SMART-seq single cell RNAseq data from the Allen Brain Atlas[138]. (C) Expression profile of genes used for transgenic and immunohistochemical targeting for cell subtyping in Allen Brain Atlas defined hippocampal GABAergic cell-types.

References

1. Karayiorgou, M., Simon, T. J. & Gogos, J. A. 22q11.2 microdeletions: linking DNA structural variation to brain dysfunction and schizophrenia. *Nature Reviews Neuroscience 2010 11:6* **11**, 402–416 (2010).
2. Jonas, R. K., Montojo, C. A. & Bearden, C. E. The 22q11.2 deletion syndrome as a window into complex neuropsychiatric disorders over the lifespan. *Biol. Psychiatry* **75**, 351–360 (2014).
3. Xu, B. *et al.* Strong association of de novo copy number mutations with sporadic schizophrenia. *Nat. Genet.* **40**, 880–885 (2008).

4. Marshall, C. R. *et al.* Contribution of copy number variants to schizophrenia from a genome-wide study of 41,321 subjects. *Nat. Genet.* **49**, 27–35 (2017).
5. Green, M. & Walker, E. *Neuropsychological Performance and Positive and Negative Symptoms in Schizophrenia.* vol. 94 460–469 (1985).
6. Young, J. W. & Geyer, M. A. Developing treatments for cognitive deficits in schizophrenia: The challenge of translation. *J. Psychopharmacol.* **29**, 178–196 (2015).

7. Starc, M. *et al.* Schizophrenia is associated with a pattern of spatial working memory deficits consistent with cortical disinhibition. *Schizophr. Res.* **181**, 107–116 (2017).
8. Andreasen, N. C. *et al.* Defining the phenotype of schizophrenia: Cognitive dysmetria and its neural mechanisms. *Biol. Psychiatry* **46**, 908–920 (1999).

1000

9. Bowie, C. R., Reichenberg, A., Patterson, T. L., Heaton, R. K. & Harvey, P. D. Determinants of real-world functional performance in schizophrenia subjects: Correlations with cognition, functional capacity, and symptoms. *Am. J. Psychiatry* **163**, 418–425 (2006).
10. Green, M. F. What are the functional consequences of neurocognitive deficits in schizophrenia? *https://doi.org/10.1176/ajp.153.3.321* **153**, 321–330 (2006).

1005

11. Aleman, A., Hijman, R., De Haan, E. H. F. & Kahn, R. S. Memory impairment in schizophrenia: A meta-analysis. *Am. J. Psychiatry* **156**, 1358–1366 (1999).
12. Danion, J.-M., Huron, C. & Vidailhet, P. Functional Mechanisms of Episodic Memory. *Can. J. Psychiatry* **52**, 693–701 (2007).
13. Fusar-Poli, P. *et al.* Neurofunctional correlates of vulnerability to psychosis: A systematic review and meta-analysis. *Neurosci. Biobehav. Rev.* **31**, 465–484 (2007).

1010

14. Harrison, P. J. & Eastwood, S. L. Neuropathological studies of synaptic connectivity in the hippocampal formation in schizophrenia. *Hippocampus* **11**, 508–519 (2001).
15. Tamminga, C. A., Stan, A. D. & Wagner, A. D. The hippocampal formation in schizophrenia. *Am. J. Psychiatry* **167**, 1178–1193 (2010).

1015

16. Bobilev, A. M., Perez, J. M. & Tamminga, C. A. Molecular alterations in the medial temporal lobe in schizophrenia. *Schizophr. Res.* (2019) doi:10.1016/j.schres.2019.06.001.
17. Flahault, A., Schaer, M., Ottet, M.-C., Debbané, M. & Eliez, S. Hippocampal volume reduction in chromosome 22q11.2 deletion syndrome (22q11.2DS): A longitudinal study of morphometry and symptomatology. *Psychiatry Research: Neuroimaging* **203**, 1–5 (2012).

1020

18. Deboer, T., Wu, Z., Lee, A. & Simon, T. J. Hippocampal volume reduction in children with chromosome 22q11.2 deletion syndrome is associated with cognitive impairment. *Behav. Brain Funct.* **3**, 54 (2007).
19. Debbané, M., Schaer, M., Farhoumand, R., Glaser, B. & Eliez, S. Hippocampal volume reduction in 22q11.2 deletion syndrome. *Neuropsychologia* **44**, 2360–2365 (2006).
20. Nelson, M. D., Saykin, A. J., Flashman, L. A. & Riordan, H. J. Hippocampal volume reduction in schizophrenia as assessed by magnetic resonance imaging: a meta-analytic study. *Arch. Gen. Psychiatry* **55**, 433–440 (1998).

1025

21. Wright, I. C. *et al.* Meta-analysis of regional brain volumes in schizophrenia. *Am. J. Psychiatry* **157**, 16–25 (2000).
22. Lieberman, J. A. *et al.* Hippocampal dysfunction in the pathophysiology of schizophrenia: a selective review and hypothesis for early detection and intervention. *Mol. Psychiatry* **23**, 1764–1772 (2018).
23. Gogos, J. A., Crabtree, G. & Diamantopoulou, A. The abiding relevance of mouse models of rare mutations to psychiatric neuroscience and therapeutics. *Schizophr. Res.* **217**, 37–51 (2020).

1030

24. Arguello, P. A. & Gogos, J. A. Modeling madness in mice: one piece at a time. *Neuron* **52**, 179–196 (2006).
25. Broglio, C. *et al.* Hippocampal pallium and map-like memories through vertebrate evolution. *J. Behav. Brain Sci.* **05**, 109–120 (2015).

26. Rodríguez, F. *et al.* Spatial memory and hippocampal pallium through vertebrate evolution: insights from reptiles and teleost fish. *Brain Res. Bull.* **57**, 499–503 (2002).

27. Pelkey, K. A. *et al.* Evolutionary conservation of hippocampal mossy fiber synapse properties. *Neuron* **111**, 3802-3818.e5 (2023).

28. Buzsáki, G. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning: HIPPOCAMPAL SHARP WAVE-RIPPLE. *Hippocampus* **25**, 1073–1188 (2015).

29. Buzsáki, G. Theta oscillations in the hippocampus. *Neuron* **33**, 325–340 (2002).

30. Seyfried, T. N. & Glaser, G. H. A review of mouse mutants as genetic models of epilepsy. *Epilepsia* **26**, 143–150 (1985).

31. Spellman, T. J. & Gordon, J. A. Synchrony in schizophrenia: a window into circuit-level pathophysiology. *Curr. Opin. Neurobiol.* **30**, 17–23 (2015).

32. Elder, G. A., Gama Sosa, M. A. & De Gasperi, R. Transgenic mouse models of Alzheimer's disease: G. A. Elder ET Al.: Transgenic models and Alzheimer's disease. *Mt. Sinai J. Med.* **77**, 69–81 (2010).

33. Wegrzyn, D., Juckel, G. & Faissner, A. Structural and functional deviations of the hippocampus in schizophrenia and schizophrenia animal models. *Int. J. Mol. Sci.* **23**, 5482 (2022).

34. Mukai, J., Cannavò, E., Sun, Z., Crabtree, G. & Diamantopoulou, A. Recapitulation and reversal of schizophrenia-related phenotypes in Setd1a - deficient mice. (2019).

35. Sigurdsson, T., Stark, K. L., Karayiorgou, M., Gogos, J. A. & Gordon, J. A. Impaired hippocampal–prefrontal synchrony in a genetic mouse model of schizophrenia. *Nature* **464**, 763–767 (2010).

36. Goff, K. M., Liebergall, S. R., Jiang, E., Somarowthu, A. & Goldberg, E. M. VIP interneuron impairment promotes in vivo circuit dysfunction and autism-related behaviors in Dravet syndrome. *Cell Rep.* **42**, 112628 (2023).

37. Lisman, J. Excitation, inhibition, local oscillations, or large-scale loops: what causes the symptoms of schizophrenia? *Curr. Opin. Neurobiol.* **22**, 537–544 (2012).

38. Uhlhaas, P. J. Dysconnectivity, large-scale networks and neuronal dynamics in schizophrenia. *Curr. Opin. Neurobiol.* **23**, 283–290 (2013).

39. Krystal, J. H. *et al.* Impaired Tuning of Neural Ensembles and the Pathophysiology of Schizophrenia: A Translational and Computational Neuroscience Perspective. *Biol. Psychiatry* **81**, 874–885 (2017).

40. Crabtree, G. & Gogos, J. A. Synaptic plasticity, neural circuits, and the emerging role of altered short-term information processing in schizophrenia. *Front. Synaptic Neurosci.* **6**, (2014).

41. Ferguson, B. R. & Gao, W. J. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. *Front. Neural Circuits* **12**, (2018).

42. Heckers, S. & Konradi, C. GABAergic mechanisms of hippocampal hyperactivity in schizophrenia. *Schizophr. Res.* **167**, 4–11 (2015).

43. Zaremba, J. D. *et al.* Impaired hippocampal place cell dynamics in a mouse model of the 22q11.2 deletion. *Nat. Neurosci.* **20**, 1612–1623 (2017).

44. Mukai, J. *et al.* Molecular substrates of altered axonal growth and brain connectivity in a mouse model of schizophrenia. *Neuron* **86**, 680–695 (2015).

45. Hamm, J. P., Peterka, D. S., Gogos, J. A. & Yuste, R. Altered Cortical Ensembles in Mouse Models of Schizophrenia. *Neuron* **94**, 153-167.e8 (2017).

46. Exposito-Alonso, D. & Rico, B. Mechanisms underlying circuit dysfunction in neurodevelopmental disorders. *Annu. Rev. Genet.* **56**, 391–422 (2022).

47. Farsi, Z. *et al.* Brain-region-specific changes in neurons and glia and dysregulation of dopamine signaling in Grin2a mutant mice. *Neuron* **111**, 3378-3396.e9 (2023).

48. Buzsáki, G. & Moser, E. I. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature Neuroscience 2013 16:2* **16**, 130–138 (2013).

49. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34**, 171–175 (1971).

50. Knierim, J. J., Lee, I. & Hargreaves, E. L. Hippocampal place cells: Parallel input streams, subregional processing, and implications for episodic memory. *Hippocampus* **16**, 755–764 (2006).

51. Danielson, N. B. *et al.* Sublayer-Specific Coding Dynamics during Spatial Navigation and Learning in Hippocampal Area CA1. *Neuron* **91**, 652–665 (2016).

52. Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory engrams in the hippocampus. *Nature* **558**, 292–296 (2018).

53. Mizuseki, K., Royer, S., Diba, K. & Buzsáki, G. Activity dynamics and behavioral correlates of CA3 and CA1 hippocampal pyramidal neurons. *Hippocampus* **22**, 1659–1680 (2012).

54. Rao-Ruiz, P. *et al.* Engram-specific transcriptome profiling of contextual memory consolidation. *Nat. Commun.* **10**, 1–14 (2019).

55. Rashid, T. Opposing gradients of ephrin-As and EphA7 in the superior colliculus are essential for topographic mapping in the mammalian visual system. *Neuron* **47**, 57–69 (2005).

56. Sosa, M., Gillespie, A. K. & Frank, L. M. Neural activity patterns underlying spatial coding in the hippocampus. *Curr. Top. Behav. Neurosci.* **37**, 43–100 (2018).

57. Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nature Neuroscience 2013 16:3* **16**, 264–266 (2013).

58. Frank, L. M., Stanley, G. B. & Brown, E. N. Hippocampal Plasticity across Multiple Days of Exposure to Novel Environments. *Journal of Neuroscience* **24**, 7681–7689 (2004).

59. Hill, A. J. First occurrence of hippocampal spatial firing in a new environment. *Exp. Neurol.* **62**, 282–297 (1978).

60. Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M. B. & Moser, E. I. Distinct ensemble codes in hippocampal areas CA3 and CA1. *Science* **305**, 1295–1298 (2004).

61. Cacucci, F., Wills, T. J., Lever, C., Giese, K. P. & O'Keefe, J. Experience-Dependent Increase in CA1 Place Cell Spatial Information, But Not Spatial Reproducibility, Is Dependent on the Autophosphorylation of the α-Isoform of the Calcium/Calmodulin-Dependent Protein Kinase II. *Journal of Neuroscience* **27**, 7854–7859 (2007).

62. Karlsson, M. P. & Frank, L. M. Network Dynamics Underlying the Formation of Sparse, Informative Representations in the Hippocampus. *Journal of Neuroscience* **28**, 14271–14281 (2008).

63. Wilson, M. A. & McNaughton, B. L. Dynamics of the hippocampal ensemble code for space. *Science* **261**, 1055–1058 (1993).

64. Turi, G. F. *et al.* Vasoactive Intestinal Polypeptide-Expressing Interneurons in the Hippocampus Support Goal-Article Vasoactive Intestinal Polypeptide-Expressing Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning. *Neuron* 1–16 (2019).

65. Gauthier, J. L. & Tank, D. W. A Dedicated Population for Reward Coding in the Hippocampus. *Neuron* **99**, 179-193.e7 (2018).

66. Hollup, S. A., Molden, S., Donnett, J. G., Moser, M. B. & Moser, E. I. Accumulation of Hippocampal Place Fields at the Goal Location in an Annular Watermaze Task. *Journal of Neuroscience* **21**, 1635–1644 (2001).

67. Muller, R. U. & Kubie, J. L. The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *Journal of Neuroscience* **7**, 1951–1968 (1987).

68. Colgin, L. L., Moser, E. I. & Moser, M. B. Understanding memory through hippocampal remapping. *Trends Neurosci.* **31**, 469–477 (2008).

69. Kubie, J. L., Levy, E. R. J. & Fenton, A. A. Is hippocampal remapping the physiological basis for context? *Hippocampus* **30**, 851 (2020).

70. Kepecs, A. & Fishell, G. Interneuron cell types are fit to function. *Nature* **505**, 318–326 (2014).

71. Tzilivaki, A. *et al.* Hippocampal GABAergic interneurons and memory. *Neuron* **111**, 3154–3175 (2023).

72. Cardin, J. A. Inhibitory interneurons regulate temporal precision and correlations in cortical circuits. *Trends Neurosci.* **41**, 689–700 (2018).

73. Harris, K. D. *et al. Classes and Continua of Hippocampal CA1 Inhibitory Neurons Revealed by Single-Cell Transcriptomics.* 1–37 (2018).

74. Pelkey, K. A. *et al.* Hippocampal GABAergic inhibitory interneurons. *Physiol. Rev.* **97**, 1619–1747 (2017).

75. Freund, T. F. & Buzsáki, G. Interneurons of the Hippocampus. *Hippocampus* **6**, 347–470 (1996).

76. Dudok, B. *et al.* Recruitment and inhibitory action of hippocampal axo-axonic cells during behavior. *Neuron* **109**, 3838-3850.e8 (2021).

77. Dudok, B. *et al.* Alternating sources of perisomatic inhibition during behavior. *Neuron* **109**, 997-1012.e9 (2021).

78. Chamberland, S. *et al.* Functional specialization of hippocampal somatostatin-expressing interneurons. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2306382121 (2024).

79. Varga, C., Golshani, P. & Soltesz, I. Frequency-invariant temporal ordering of interneuronal discharges during hippocampal oscillations in awake mice. *Proc. Natl. Acad. Sci. U. S. A.* **109**, E2726-34 (2012).

80. Somogyi, P. & Klausberger, T. Defined types of cortical interneurone structure space and spike timing in the hippocampus. *J. Physiol.* **562**, 9–26 (2005).

81. Grienberger, C., Milstein, A. D., Bittner, K. C., Romani, S. & Magee, J. C. Inhibitory suppression of heterogeneously tuned excitation enhances spatial coding in CA1 place cells. *Nat. Neurosci.* **20**, 417–426 (2017).

82. Juarez, P. & Martínez Cerdeño, V. Parvalbumin and parvalbumin chandelier interneurons in autism and other psychiatric disorders. *Front. Psychiatry* **13**, 913550 (2022).

83. Beasley, C. L. & Reynolds, G. P. Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics. *Schizophr. Res.* **24**, 349–355 (1997).

84. Lewis, D. A., Curley, A. A., Glausier, J. R. & Volk, D. W. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends in Neurosciences* Preprint at https://doi.org/10.1016/j.tins.2011.10.004 (2012).

85. Nakazawa, K. *et al.* GABAergic interneuron origin of schizophrenia pathophysiology. *Neuropharmacology* **62**, 1574–1583 (2012).

86. Chung, D. W., Fish, K. N. & Lewis, D. A. Pathological basis for deficient excitatory drive to cortical parvalbumin interneurons in schizophrenia. *Am. J. Psychiatry* **173**, 1131–1139 (2016).

87. Fung, S. J., Fillman, S. G., Webster, M. J. & Shannon Weickert, C. Schizophrenia and bipolar disorder show both common and distinct changes in cortical interneuron markers. *Schizophr. Res.* **155**, 26–30 (2014).

88. Gonzalez-Burgos, G., Cho, R. Y. & Lewis, D. A. Alterations in Cortical Network Oscillations and Parvalbumin Neurons in Schizophrenia. *Biol. Psychiatry* **77**, 1031–1040 (2015).

89. Lewis, D. A. Inhibitory neurons in human cortical circuits: substrate for cognitive dysfunction in schizophrenia. *Curr. Opin. Neurobiol.* **26**, 22–26 (2014).

90. Dienel, S. J. *et al.* Diagnostic specificity and association with cognition of molecular alterations in prefrontal somatostatin neurons in schizophrenia. *JAMA Psychiatry* **80**, 1235–1245 (2023).

91. Dienel, S. J., Fish, K. N. & Lewis, D. A. The nature of prefrontal cortical GABA neuron alterations in schizophrenia: Markedly lower somatostatin and parvalbumin gene expression without missing neurons. *Am. J. Psychiatry* **180**, 495–507 (2023).

92. Schmidt, M. J. & Mirnics, K. Modeling interneuron dysfunction in schizophrenia. *Dev. Neurosci.* **34**, 152–158 (2012).

93. Al-Absi, A.-R. *et al.* Layers II/III of prefrontal cortex in Df(h22q11)/+ mouse model of the 22q11.2 deletion display loss of parvalbumin interneurons and modulation of neuronal morphology and excitability. *Mol. Neurobiol.* **57**, 4978–4988 (2020).

94. Jahangir, M., Zhou, J.-S., Lang, B. & Wang, X.-P. GABAergic system dysfunction and challenges in schizophrenia research. *Front. Cell Dev. Biol.* **9**, 663854 (2021).

95. Geiller, T. *et al.* Large-Scale 3D Two-Photon Imaging of Molecularly Identified CA1 Interneuron Dynamics in Behaving Mice. *Neuron* **108**, 968-983.e9 (2020).

96. Vancura, B., Geiller, T., Grosmark, A., Zhao, V. & Losonczy, A. Inhibitory control of sharp-wave ripple duration during learning in hippocampal recurrent networks. *Nat. Neurosci.* **26**, 788–797 (2023).

97. Mukherjee, A., Carvalho, F., Eliez, S. & Caroni, P. Long-Lasting Rescue of Network and Cognitive Dysfunction in a Genetic Schizophrenia Model. *Cell* **178**, 1387-1402.e14 (2019).

98. Stark, K. L. *et al.* Altered brain microRNA biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. *Nat. Genet.* **40**, 751–760 (2008).

99. Forro, T. & Klausberger, T. Differential behavior-related activity of distinct hippocampal interneuron types during odor-associated spatial navigation. *Neuron* **111**, 2399-2413.e5 (2023).

100. Michon, F.-X., Laplante, I., Bosson, A., Robitaille, R. & Lacaille, J.-C. mTORC1-mediated acquisition of reward-related representations by hippocampal somatostatin interneurons. *Mol. Brain* **16**, 55 (2023).

101. Diamantopoulou, A. *et al.* Loss-of-function mutation in *Mirta22/Emc10* rescues specific schizophrenia-related phenotypes in a mouse model of the 22q11.2 deletion. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E6127–E6136 (2017).

102. Marissal, T. *et al.* Restoring wild-type-like CA1 network dynamics and behavior during adulthood in a mouse model of schizophrenia. *Nat. Neurosci.* **21**, 1412–1420 (2018).

103. Geiller, T. *et al.* Local circuit amplification of spatial selectivity in the hippocampus. *Nature* **601**, 105–109 (2021).

104. Rolotti, S. V. *et al.* Local feedback inhibition tightly controls rapid formation of hippocampal place fields. *Neuron* **110**, 783-794.e6 (2022).

105. Hangya, B., Li, Y., Muller, R. U. & Czurkó, A. Complementary spatial firing in place cell-interneuron pairs. *J. Physiol.* **588**, 4165–4175 (2010).

106. Ego-Stengel, V. & Wilson, M. A. Spatial selectivity and theta phase precession in CA1 interneurons. *Hippocampus* **17**, 161–174 (2007).

107. Markus, E. J., Barnes, C. A., McNaughton, B. L., Gladden, V. L. & Skaggs, W. E. Spatial information content and reliability of hippocampal CA1 neurons: effects of visual input. *Hippocampus* **4**, 410–421 (1994).

108. Skaggs, W., McNaughton, B. & Gothard, K. An information-theoretic approach to deciphering the hippocampal code. *Adv. Neural Inf. Process. Syst.* 1030–1037 (1992).

109. Grijseels, D. M., Shaw, K., Barry, C. & Hall, C. N. Choice of method of place cell classification determines the population of cells identified. *PLoS Comput. Biol.* **17**, e1008835 (2021).

110. Somogyi, J. *et al.* GABAergic basket cells expressing cholecystokinin contain vesicular glutamate transporter type 3 (VGLUT3) in their synaptic terminals in hippocampus and isocortex of the rat. *Eur. J. Neurosci.* **19**, 552–569 (2004).

111. Schneider, A. *et al.* Transcriptomic cell type structures in vivo neuronal activity across multiple timescales. *Cell Rep.* **42**, 112318 (2023).

112. Fuhrmann, F. *et al.* Locomotion, theta oscillations, and the speed-correlated firing of hippocampal neurons are controlled by a medial septal glutamatergic circuit. *Neuron* **86**, 1253–1264 (2015).

113. Arriaga, M. & Han, E. B. Dedicated hippocampal inhibitory networks for locomotion and immobility. *J. Neurosci.* **37**, 9222–9238 (2017).

114. Lapray, D. *et al.* Behavior-dependent specialization of identified hippocampal interneurons. *Nat. Neurosci.* **15**, 1265–1271 (2012).

115. Gridchyn, I., Schoenenberger, P., O'Neill, J. & Csicsvari, J. Optogenetic inhibition-mediated activity-dependent modification of CA1 pyramidal-interneuron connections during behavior. *Elife* **9**, (2020).

116. Schoenenberger, P., O'Neill, J. & Csicsvari, J. Activity-dependent plasticity of hippocampal place maps. *Nat. Commun.* **7**, 11824 (2016).

117. Priestley, J. B., Bowler, J. C., Rolotti, S. V., Fusi, S. & Losonczy, A. Signatures of rapid plasticity in hippocampal CA1 representations during novel experiences. *Neuron* **110**, 1978-1992.e6 (2022).

118. Hainmueller, T., Cazala, A., Huang, L.-W. & Bartos, M. Subfield-specific interneuron circuits govern the hippocampal response to novelty in male mice. *Nat. Commun.* **15**, 714 (2024).

119. Rangel Guerrero, D. K. *et al.* Hippocampal cholecystokinin-expressing interneurons regulate temporal coding and contextual learning. *Neuron* (2024) doi:10.1016/j.neuron.2024.03.019.

120. Lovett-Barron, M. *et al.* Dendritic inhibition in the hippocampus supports fear learning. *Science* **343**, 857–863 (2014).

121. Varga, V. *et al.* Fast synaptic subcortical control of hippocampal circuits. *Science* **326**, 449–453 (2009).

122. Basu, J. *et al.* Gating of hippocampal activity, plasticity, and memory by entorhinal cortex long-range inhibition. *Science* **351**, aaa5694 (2016).

123. Chamberland, S. & Topolnik, L. Inhibitory control of hippocampal inhibitory neurons. *Front. Neurosci.* **0**, 165 (2012).

124. Szabo, G. G. *et al.* Ripple-selective GABAergic projection cells in the hippocampus. *Neuron* **110**, 1959-1977.e9 (2022).

125. Batista-Brito, R. *et al.* Developmental Dysfunction of VIP Interneurons Impairs Cortical Circuits. *Neuron* **95**, 884-895.e9 (2017).

126. Piskorowski, R. A. *et al.* Age-Dependent Specific Changes in Area CA2 of the Hippocampus and Social Memory Deficit in a Mouse Model of the 22q11.2 Deletion Syndrome. *Neuron* **89**, 163–176 (2016).

127. Brown, J. A. *et al.* Inhibition of parvalbumin-expressing interneurons results in complex behavioral changes. *Mol. Psychiatry* **20**, 1499–1507 (2015).

128. Sauer, J.-F. & Bartos, M. Disrupted-in-schizophrenia-1 is required for normal pyramidal cell-interneuron communication and assembly dynamics in the prefrontal cortex. *Elife* **11**, (2022).

129. Del Pino, I. *et al.* Erbb4 deletion from fast-spiking interneurons causes schizophrenia-like phenotypes. *Neuron* **79**, 1152–1168 (2013).

130. Schmidt, M. J. *et al.* Modulation of behavioral networks by selective interneuronal inactivation. *Molecular Psychiatry 2013 19:5* **19**, 580–587 (2013).

131. Morris, H. M., Hashimoto, T. & Lewis, D. A. Alterations in Somatostatin mRNA Expression in the Dorsolateral Prefrontal Cortex of Subjects with Schizophrenia or Schizoaffective Disorder. *Cereb. Cortex* **18**, 1575–1587 (2008).

132. Morris, H. M., Stopczynski, R. E. & Lewis, D. A. NPY mRNA expression in the prefrontal cortex: Selective reduction in the superficial white matter of subjects with schizoaffective disorder. *Schizophr. Res.* **115**, 261–269 (2009).

133. Koukouli, F. *et al.* Nicotine reverses hypofrontality in animal models of addiction and schizophrenia. *Nature Medicine 2017 23:3* **23**, 347–354 (2017).

1250

134. Rocco, B. R., DeDionisio, A. M., Lewis, D. A. & Fish, K. N. Alterations in a Unique Class of Cortical Chandelier Cell Axon Cartridges in Schizophrenia. *Biol. Psychiatry* **82**, 40–48 (2017).

135. Wang, Y., Zhang, P. & Wyskiel, D. R. Chandelier Cells in Functional and Dysfunctional Neural Circuits. *Front. Neural Circuits* **10**, 33 (2016).

136. Van Derveer, A. B. *et al.* A role for somatostatin-positive interneurons in neuro-oscillatory and information processing deficits in schizophrenia. *Schizophr. Bull.* **47**, 1385–1398 (2021).

1255

137. Kaifosh, P., Zaremba, J. D., Danielson, N. B. & Losonczy, A. SIMA: Python software for analysis of dynamic fluorescence imaging data. *Front. Neuroinform.* **8**, 80 (2014).

138. Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex and hippocampal formation. *bioRxiv* (2020) doi:10.1101/2020.03.30.015214.