GLP-1R associates with VAPB and SPHKAP at ERMCSs to regulate β-cell mitochondrial remodelling and function

Gregory Austin[1,*], Affiong I. Oqua[1,*], Liliane El Eid[1], Mingli Zhu[1], Yusman Manchanda[1], Priyanka Peres[2], Helena Coyle[2], Yelyzaveta Poliakova[3], Karim Bouzakri[4], Alex Montoya[5], Dominic J. Withers[5,6], Michele Solimena[7,8,9], Ben Jones[10], Steven J. Millership[1], Steffen Burgold[11], David C.A. Gaboriau[12], Endre Majorovits[11], Evelyn Garlick[13], Maria Lima[13], Inga Prokopenko[14,15], Jonathon Nixon-Abell[2], Andreas Müller[7,8,9] and Alejandra Tomas[1,#]

[1] Section of Cell Biology and Functional Genomics, Division of Diabetes, Endocrinology, and Metabolism, Department of Metabolism, Digestion and Reproduction, Imperial College London, London, United Kingdom.
[2] Cambridge Institute for Medical Research, Department of Clinical Neurosciences, University of Cambridge, Cambridge, United Kingdom.
[3] School of Public Health, Imperial College London, London, United Kingdom.
[4] Centre Européen d'Étude du Diabète, Strasbourg, France.
[5] MRC Laboratory of Medical Sciences, London, United Kingdom.
[6] Institute of Clinical Sciences, Faculty of Medicine, Imperial College London, London, United Kingdom.
[7] Molecular Diabetology, University Hospital and Faculty of Medicine Carl Gustav Carus, TU Dresden, Dresden, Germany.
[8] Paul Langerhans Institute Dresden, Helmholtz Center Munich, University Hospital Carl Gustav Carus and Faculty of Medicine, TU Dresden, Dresden, Germany.
[9] German Center for Diabetes Research, Neuherberg, Germany.
[10] Section of Endocrinology and Investigative Medicine, Division of Diabetes, Endocrinology, and Metabolism, Department of Metabolism, Digestion and Reproduction, Imperial College London, London, United Kingdom.
[11] Carl Zeiss Microscopy GmbH, ZEISS Gruppe, Oberkochen, Germany.
[12] Facility for Imaging by Light Microscopy, National Heart and Lung Institute, Imperial College London, London, United Kingdom.
[13] Abberior Instruments GmbH, Göttingen, Germany.
[14] Department of Clinical and Experimental Medicine, University of Surrey, Guildford, United Kingdom.
[15] People-Centred Artificial Intelligence Institute, University of Surrey, Guildford, United Kingdom.

*Joint first authors

#Corresponding author: a.tomas-catala@imperial.ac.uk

Abstract

Glucagon-like peptide-1 receptor (GLP-1R) agonists (GLP-1RAs) ameliorate mitochondrial health by increasing mitochondrial turnover in metabolically relevant tissues. Mitochondrial adaptation to metabolic stress is crucial to maintain pancreatic β-cell function and prevent type 2 diabetes (T2D) progression. While the GLP-1R is well-known to stimulate cAMP production leading to Protein Kinase A (PKA) and Exchange Protein Activated by cyclic AMP 2 (Epac2) activation, there is a lack of understanding of the molecular mechanisms linking GLP-1R signalling with mitochondrial and β-cell functional adaptation. Here, we present a comprehensive study in β-cell lines and primary islets that demonstrates that, following GLP-1RA stimulation, GLP-1R-positive endosomes associate with the endoplasmic reticulum (ER) membrane contact site (MCS) tether VAPB at ER-mitochondria MCSs (ERMCSs), where active GLP-1R engages with SPHKAP, an A-kinase anchoring protein (AKAP) previously linked to T2D and adiposity risk in genome-wide association studies (GWAS). The inter-organelle complex formed by endosomal GLP-1R, ER VAPB and SPHKAP triggers a pool of ERMCS-localised cAMP/PKA signalling via the formation of a PKA-RIα biomolecular condensate which leads to changes in mitochondrial contact site and cristae organising system (MICOS) complex phosphorylation, mitochondrial remodelling, and β-cell functional adaptation, with important consequences for the regulation of β-cell insulin secretion and survival to stress.

Introduction

Maintenance of mitochondrial function and adaptation to changes in nutritional state is vital for the preservation of pancreatic β-cell regulated secretion of insulin[1], the main hormone in charge of lowering blood glucose concentrations, with disruptions in this process underlying the development and/or progression of type 2 diabetes (T2D)[1, 2] – an uncontrolled pandemic affecting ~500 million people worldwide, and causing ~1.5 million deaths per year[3]. The incretin glucagon-like peptide-1 (GLP-1), released from the gut during food intake, amplifies glucose-stimulated insulin release by binding to and activating its cognate G protein-coupled receptor (GPCR), the GLP-1 receptor (GLP-1R), triggering cAMP generation and downstream Protein Kinase A (PKA) and Exchange Protein Activated by cyclic AMP 2 (Epac2) signalling[4], a process that has been exploited to develop pharmacological GLP-1R agonists (GLP-1RAs) as T2D therapies. There is extensive data highlighting the beneficial role of GLP-1RAs in the maintenance and/or restoration of mitochondrial function in β-cells and beyond, with reports emphasising their capacity to induce mitochondrial remodelling/turnover and functional adaptation[5, 6, 7, 8], two closely intertwined processes essential for the preservation of optimal metabolic activity[9]. While a range of GLP-1R-induced pathways have been suggested, there is no clear description of the molecular mechanism engaged by the receptor to modulate mitochondrial remodelling and improve mitochondrial function. Here, we present new evidence in β-cells and primary mouse and human islets which demonstrates that, following agonist binding and internalisation, endosomal GLP-1R binds to endoplasmic reticulum (ER)-localised ER-mitochondria membrane contact site (ERMCS) organising factor VAPB to engage SPHKAP, a PKA-RIα-specific A-kinase anchoring protein (AKAP) whose gene coding variants have previously been linked to T2D and high BMI risk in genome-wide association studies (GWAS). We further show that SPHKAP itself localises to ERMCSs via its direct interaction with VAPB through a pFFAT motif present in SPHKAP. Establishment of a membrane contact site (MCS) between endosomal GLP-1R, ERMCS-localised VAPB and SPHKAP leads to the generation of a highly-localised cAMP / PKA signalling hub that triggers the PKA-dependent phosphorylation of the mitochondrial contact site and cristae organising system (MICOS) complex, mitochondrial remodelling and improved mitochondrial function, leading to the potentiation of insulin secretion and survival to ER stress downstream of GLP-1R action. This is, to our knowledge, the first description of a three-way contact between an endosomal GPCR, the ERMCS tether VAPB and a PKA-RIα-recruiting AKAP, which uncovers the assembly of a GLP-1R-induced ERMCS-localised PKA biomolecular condensate that controls mitochondrial turnover and function. Our data not only unveils a previously unknown pathway of ERMCS-localised GPCR signalling into mitochondria, potentially applicable to other GPCRs beyond the GLP-1R, but also opens the door to further roles of the GLP-1R in the regulation of mitochondrial lipid metabolism, a process tightly linked to changes in mitochondrial morphology governed by MICOS complex activity[10]. In summary, this study has uncovered the molecular mechanism employed by the GLP-1R to achieve mitochondrial regeneration leading to optimised β-cell function, with important repercussions for our understanding of the role of GLP-1RAs in disorders such as T2D, obesity, and neurodegeneration.

Results

Human GLP-1R β-cell interactome reveals binding of agonist-stimulated GLP-1R to signalling effectors, with strong prevalence of ERMCSs-localised proteins.

To gain a comprehensive understanding of the specific downstream partners and signalling mediators engaged by active GLP-1Rs in pancreatic β-cells, we performed an interactome analysis of proteins that bind to the receptor under vehicle *versus* 5-minute stimulation with the GLP-1RA exendin-4 in a subline of rat INS-1 832/3 β-cells stably expressing

SNAP/FLAG-tagged human GLP-1R (SNAP/FLAG-hGLP-1R). Mass spectrometry (MS) analysis of GLP-1R co-immunoprecipitants revealed a list of proteins enriched in agonist-stimulated *versus* unstimulated conditions (Fig. 1a,b, Supplementary Fig. 1a). Gene ontology analysis of the identified factors indicated that these were involved in pathways related to endomembranes and the ER (Supplementary Fig. 1b). Amongst these factors, we found a high number of proteins localised to ERMCSs (Fig. 1c), which are points of close interaction, but not fusion, between the ER and mitochondria that fulfil specific functions including cross-organelle ion and lipid sensing and transfer, and signal transmission to coordinate mitochondrial remodelling[11]. While known GLP-1R interactors such as $G\alpha_S$, AP2, Rab5, and β-arrestin 2 were reassuringly identified in our interactomics list, the most prominently enriched factor interacting with active GLP-1R was VAMP Associated Protein B (VAPB), followed by its homolog VAPA, integral ER proteins whose main role is the establishment and maintenance of organelle-ER interactions, or membrane contact sites (MCSs)[12, 13], including those involving endosomes and/or mitochondria. We next validated the active GLP-1R binding to VAPB inferred from our MS results by co-immunoprecipitating both EGFP-VAPB (Fig. 1d) and endogenous VAPB (Fig. 1e, Supplementary Fig. 1c) with SNAP/FLAG-hGLP-1R in vehicle *versus* exendin-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells. Confocal microscopy analysis of vehicle *versus* exendin-4-stimulated SNAP/FLAG-hGLP-1R co-localisation with EGFP-VAPB in these cells revealed that, upon exendin-4 stimulation, the GLP-1R traffics from the plasma membrane to endosomes intimately associated with VAPB-positive ER regions (Supplementary Fig. 1d,e), with the ER often surrounding a central core of GLP-1R-positive signal (Fig. 1f), suggesting that GLP-1R-containing endosomes directly contact VAPB-positive ER membranes, likely via MCSs. This hypothesis was validated by transmission electron microscopy (TEM) localisation of SNAP/FLAG-hGLP-1R (labelled at the cell surface with SNAP-biotin plus streptavidin-gold particles in living cells prior to exendin-4 stimulation) to endosome-ER MCSs (Fig. 1g). Further microscopy experiments performed to visualise GLP-1R engagement with EGFP-VAPB in response to exendin-4 stimulation include confocal imaging of serial optical sections covering the entire cell thickness after SNAP/FLAG-hGLP-1R surface labelling with a membrane impermeable SNAP-tag probe followed by exendin-4 stimulation, represented as a maximum intensity projection (MIP) in Fig. 1h, and time-lapse confocal microscopy analysis of exendin-4-induced SNAP/FLAG-hGLP-1R trafficking from the plasma membrane to endosomes in close proximity to EGFP-VAPB-positive ER regions (Supplementary Fig. 1f, Supplementary Movie 1). Two-colour nanometre-scale resolution imaging by Minimal Photon Fluxes (MINFLUX), an ultra-high resolution microscopy technique that combines aspects of single-molecule localisation microscopy (SMLM) and stimulated emission depletion (STED) microscopy[14], was employed to determine the average distance between endosomal SNAP/FLAG-hGLP-1R and VAPB, estimated at around 28 nm (Fig. 1i,j), compatible with the expected distance of an ER-endosome MCS[15].

The interaction of active GLP-1R with VAPB was deemed functionally relevant, as it was no longer present when VAPB P56S, a self-aggregating loss-of-function VAPB mutant associated with amyotrophic lateral sclerosis (ALS)[16, 17], was expressed instead of its wildtype (WT) counterpart (Supplementary Fig. 1g-i), and as both VAPB knockdown by RNAi (Fig. 1k, Supplementary Fig. 1j,k), and overexpression of VAPB P56S, but not of WT VAPB (Supplementary Fig. 1l), significantly reduced the capacity of GLP-1R to potentiate insulin secretion from INS-1 832/3 cells.

GLP-1R binding to VAPB requires GLP-1R internalisation and is differentially modulated by GLP-1RAs with varying GLP-1R internalisation propensities.

To elucidate whether active GLP-1R requires its prior internalisation to engage VAPB, we co-transfected INS-1 832/3 SNAP/FLAG-hGLP-1R cells with dominant negative (K44A) mutants of dynamin 1 and 2 to block GLP-1R endocytosis triggered by exendin-4 (Supplementary Fig. 2a). Under these conditions, increased binding of the receptor to VAPB in stimulated

versus vehicle conditions was abrogated (Fig. 2a), demonstrating that active GLP-1R
interacts with VAPB following its agonist-induced internalisation. Supporting this notion, MS
interactomic analyses of GLP-1R co-immunoprecipitates in INS-1 832/3 SNAP/FLAG-hGLP-
1R cells after 5-minute stimulation with the endogenous agonist GLP-1, as well as with the
exendin-4-based biased GLP-1RAs exendin-F1 and exendin-D3, previously shown to trigger
differing degrees of GLP-1R internalisation[18, 19], resulted in marked differences in the
propensity of the receptor to associate with VAPB (Supplementary Fig. 3), with agonists that
trigger robust receptor internalisation (GLP-1, exendin-4, exendin-D3) showing a stronger
degree of GLP-1R-VAPB interaction compared to slow GLP-1R-internalising exendin-F1.
Marked kinetic differences in GLP-1R-VAPB interaction following exendin-4 *versus* exendin-
F1 stimulation were also demonstrated in INS-1 832/3 cells by a novel NanoBRET assay,
with Nanoluciferase-tagged GLP-1R co-expressed with Venus-tagged VAPB (Fig. 2b,
Supplementary Fig. 2b). We extended our analysis to a panel of seven GLP-1RAs, including,
as well as exendin-4, exendin-F1 and exendin-D3, fatty acid-modified semaglutide[20] and
tirzepatide[21], as well as small molecules orforglipron[22] and danuglipron[23]. We first tested
these GLP-1RAs for their capacity to induce GLP-1R internalisation in purified mouse islets
transduced with SNAP/FLAG-hGLP-1R-expresssing adenoviruses (Fig. 2c). As previously
shown, acute (5-minute) exposure to exendin-4 and exendin-D3 triggered noticeable GLP-
1R internalisation, while this was much reduced with the biased agonist exendin-F1.
Additionally, and as previously published[24], semaglutide triggered a much more robust GLP-
1R internalisation than tirzepatide, an agonist that shows a similar degree of GLP-1R $G_{\alpha S}$
bias to exendin-F1. We also detected a clear difference in GLP-1R internalisation triggered
by orforglipron *versus* danuglipron, with the latter showing a similar level of receptor
internalisation to that of exendin-4, while the former resulted in very reduced receptor
internalisation 5 minutes post-agonist exposure. VAPB co-immunoprecipitation using the
panel of GLP-1RAs described above unveiled a correlation between the level of GLP-1R–
VAPB binding and the degree of receptor internalisation elicited by each of the tested GLP-
1RAs (Fig. 2d, Supplementary Fig. 2c).
247
248 **Exendin-4-stimulated GLP-1R triggers VAPB-localised cAMP generation and PKA**
249 **activity at the outer mitochondrial membrane.**
250
As stated above, besides VAPB, our interactome analysis highlighted several factors
enriched for their association with active GLP-1R localised at the ER, mitochondria and/or
ERMCSs (Fig. 1c), a location where VAPB plays a prominent organising role[17], suggesting
that ERMCSs could be a potential hub for endosomal GLP-1R signalling. To determine if
GLP-1R induces cAMP generation specifically at VAPB locations, we modified a
fluorescence resonance energy transfer (FRET)-based system of biosensors targeted to
endogenous proteins (FluoSTEPs)[25], so that a FRET cAMP sensor would fully reconstitute
only in the presence of GFP_{11} fragment fused to VAPB, resulting in functional biosensor
assembly specifically at VAPB loci (Supplementary Fig. 2d). This modified FluoSTEP system
was used to measure VAPB-localised cAMP generation in response to either exendin-4 or
exendin-F1 stimulation (Fig. 2e). Results in WT INS-1 832/3 cells (expressing endogenous
levels of GLP-1R) show VAPB-localised cAMP responses with both agonists, which are less
pronounced for the slow internalising exendin-F1 compared to exendin-4, demonstrating that
GLP-1R actively signals at VAPB loci. Next, we evaluated whether GLP-1R signalling would
also trigger a pool of PKA activity at the surface of mitochondria, a process known to be
important for the control of mitochondrial function[26]. To this end, we employed either a global
(whole cell) or a mitochondrially-restricted PKA biosensor expressed at the outer
mitochondrial membrane (OMM)[27] (Supplementary Fig. 2e). Exendin-4 stimulation of WT
INS-1 832/3 cells triggered, as expected, a robust global PKA response, which was also
evident with the OMM-localised PKA biosensor, demonstrating that GLP-1R stimulation
triggers PKA activity at the mitochondrial surface (Supplementary Fig. 2f). Furthermore,
utilising the same biosensors as above, we detected reduced OMM-localised but no change
in global PKA activity with slow internalising exendin-F1 *versus* exendin-4 (Fig. 2f), leading to

increased mitochondrial over global PKA signalling propensity with exendin-4 at this acute
(5-minute) stimulation time-point (Fig. 2g). This, together with the VAPB-localised cAMP
FluoSTEP results from above, indicates that GLP-1R internalisation is important for GLP-1R-
induced cAMP generation at VAPB loci and PKA signalling at OMMs, at least acutely.
278
279 **Exendin-4-stimulated GLP-1R associates with ERMCSs.**
280
To examine the potential docking of GLP-1R-positive endosomes to ERMCSs, we employed
a split-GFP-based contact site sensor (SPLICS)[28] to specifically visualise ERMCSs in
conjunction with SNAP/FLAG-hGLP-1R in vehicle *versus* exendin-4-stimulated INS-1 832/3
SNAP/FLAG-hGLP-1R cells (Fig. 3a-c, Supplementary Movie 2). This approach allowed us
to observe an intimate association between GLP-1R-positive endosomes and ERMCSs over
time, validating our hypothesis. Additional cell labelling with fluorescent MitoTracker prior to
exendin-4 exposure showed SNAP/FLAG-hGLP-1R extensively colocalised with EGFP-
VAPB in proximity to mitochondria, an organelle with which VAPB is known to interact (Fig.
3d).
290
Further high-resolution 3D imaging to confirm endosomal GLP-1R localisation to ERMCSs
included *in situ* Correlative Light and Electron Microscopy (CLEM) at room temperature to
correlate ultrastructural features with positive fluorescent signals for SNAP/FLAG-hGLP-1R,
EGFP-VAPB and mitochondria. To this end, INS-1 832/3 SNAP/FLAG-hGLP-1R cells were
labelled at the cell surface with a membrane impermeable fluorescent SNAP-tag probe prior
to exendin-4 stimulation, with correlation of the confocal slice to the corresponding TEM
tomogram (Fig. 3e) revealing two mitochondrial volumes in close contact with EGFP-VAPB-
positive ER, itself docked to a vesicle positive for SNAP/FLAG-hGLP-1R indicative of a GLP-
1R-positive endosome (Fig. 3f, Supplementary Movie 3). Similar results were obtained using
a cryo-CLEM approach, with a SNAP/FLAG-hGLP-1R endosome docked to the EGFP-
VAPB-positive ER network, itself in contact with a nearby mitochondrion, following
correlation of confocal and FIB-SEM data (Supplementary Fig. 4a-c).
303
These results were further replicated using an immuno-EM approach in primary mouse islets
transduced with a SNAP/FLAG-hGLP-1R-expressing adenovirus, by labelling cell surface
SNAP/FLAG-hGLP-1R in living islets with membrane-impermeable SNAP-biotin plus 5-nm
gold-conjugated streptavidin prior to exendin-4 stimulation and EM processing. TEM imaging
of ultrathin (70-nm) sections from the resin-embedded islets from above revealed 5-nm gold-
positive endosomes (indicative of the presence of SNAP/FLAG-hGLP-1R) forming contacts
with the ER, itself contacting nearby mitochondria (Fig. 3g).
311
312 **Active GLP-1R interacts with the PKA-RI-recruiting AKAP SPHKAP at ERMCSs via**
313 **VAPB.**
314
Amongst the interactors enriched for active GLP-1R binding in our MS interactome analysis,
we identified both PKA-RIα/β and the A-kinase anchoring protein (AKAP) SPHKAP, a
paralog of AKAP11 (also known as AKAP220) that binds to two RIα subunits of the PKA
holoenzyme[29]. Both SPHKAP and AKAP11 harbour a FFAT (two phenylalanines in an acidic
tract) motif[30, 31], a specific short protein sequence that binds to the Major Sperm Protein
(MSP) domain of VAPs (Supplementary Fig. 5a), which, in the case of SPHKAP, can be
regulated by phosphorylation (phosphoFFAT, or pFFAT)[31]. Agreeably, SPHKAP has
previously been shown to interact with VAP proteins in neurons and assemble functionally
relevant PKA-RI biomolecular condensates at VAPA/B-positive ER locations[32, 33]. SPHKAP is
also expressed in metabolically relevant tissues, including pancreatic islets and intestinal K-
and L-cells, and has been previously shown to play important roles in both incretin secretion
from the gut and incretin-dependent glucoregulation[34, 35]. Additionally, we found SPHKAP
mRNA expression to be decreased in a published RNA Seq database of islets from *db/db*
versus WT mice (Supplementary Fig. 5b), and meta-analyses from four separate genome-

wide association studies (GWAS) in humans uncovered genome-wide significant associations between *SPHKAP* gene variants, T2D risk, and high BMI [36, 37, 38, 39] (Supplementary Fig. 5c,d). We found the effect of *SPHKAP* gene variants in T2D *versus* high BMI to be tightly correlated (Supplementary Fig. 5e), suggesting a shared mechanism of action for SPHKAP in both disorders where GLP-1RAs are also known to be effective. Our own genetic analysis also demonstrated a significant association between *SPHKAP* gene coding variants and random blood glucose levels in whole exome sequencing (WES) data from UK Biobank (UKBB) individuals without diabetes (Supplementary Fig. 5f), strongly supporting a role for SPHKAP in the regulation of β-cell function.

We therefore next investigated the potential involvement of SPHKAP/PKA-RI in GLP-1R signal transduction from ERMCSs. We first confirmed active GLP-1R – SPHKAP protein-protein interaction, inferred from our MS interactome analysis, by measuring increased co-immunoprecipitation of SPHKAP-EGFP (Fig. 4a,b, Control RNAi conditions; Supplementary Fig. 6a), as well as of endogenous SPHKAP (Supplementary Fig. 6b), with SNAP/FLAG-hGLP-1R in exendin-4-stimulated *versus* vehicle-treated INS-1 832/3 SNAP/FLAG-hGLP-1R cells. Interestingly, we also detected a similar association between the other incretin receptor, glucose-dependent insulinotropic polypeptide receptor (GIPR), and endogenous SPHKAP following GIP stimulation of INS-1 832/3 cells stably expressing SNAP/FLAG-hGIPR (Supplementary Fig. 6c), suggesting that this localised PKA signalling module is shared between both incretin receptors.

We next determined that the exendin-4-induced GLP-1R – SPHKAP interaction requires VAPB, as it is abrogated by VAPB knockdown (Fig. 4a,b, VAPB RNAi conditions; Supplementary Fig. 6a). Confocal analysis of SPHKAP-EGFP localisation in WT INS-1 832/3 cells revealed that, in contrast with a previous report suggesting mitochondrial localisation[40], SPHKAP localises to an intracellular network occasionally in contact with, but not inside, mitochondria (Fig. 4c, Control RNAi panels; Supplementary Fig. 6d). Indeed, labelling of ER with the fluorescent dye ER-Tracker revealed the punctate localisation of SPHKAP-EGFP along ER membranes (Fig. 4d,e), in agreement with recent reports which also indicate SPHKAP localisation and PKA-RI targeting to the ER[32, 33]. Also in agreement with these reports, the localisation of SPHKAP-EGFP to ER loci required VABP, as it was lost in VAPB RNAi-treated cells (Fig. 4c, VAPB RNAi panels). We next generated a mutant version of SPHKAP-EGFP harbouring a point mutation in its pFFAT motif (SPHKAP-EGFP A215E, DFLT**A**SE to DFLT**E**SE). This strategy was based on previous structural data showing that position 5 of the FFAT core motif is positioned within a hydrophobic pocket in the MSP domain of VAP[41], with previous studies indicating that introducing a negative charge in this position causes a steric hindrance which effectively disrupts FFAT motif-VAP interaction[42]. Agreeably, our SPHKAP-EGFP pFFAT mutant lost both its binding to exendin-4-stimulated SNAP/FLAG-hGLP-1R (Supplementary Fig. 6e) and its localisation to the ER network (Supplementary Fig. 6f). We additionally demonstrated the co-existence of SNAP/FLAG-hGLP-1R, VAPB and SPHKAP in mitochondria-associated membranes (MAMs) purified from exendin-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells (Supplementary Fig. 6g), and a protein-protein interaction between SPHKAP and EGFP-VAPB in the same cells (Supplementary Fig. 6h). Taken as a whole, these results demonstrate the association of SPHKAP with active GLP-1R at ERMCSs in a VAPB-dependent manner via its pFFAT motif.

ERMCS-localised GLP-1R-dependent PKA signalling requires SPHKAP.

We next tested the effect of SPHKAP downregulation on GLP-1R-dependent downstream signalling. SPHKAP knockdown efficiency in INS-1 832/3 cells was validated against both endogenous SPHKAP (Supplementary Fig. 6i) and SPHKAP-EGFP (Supplementary Fig. 6j). Loss of SPHKAP did not affect the interaction between the receptor and VAPB (Supplementary Fig. 7a-c), indicating that GLP-1R is recruited to VAPB-positive ERMCSs independently of SPHKAP. Loss of SPHKAP also did not affect the capacity for the receptor

to generate cAMP following exendin-4 stimulation of WT INS-1 832/3 cells (Fig. 4f). In
contrast, while global PKA responses to exendin-4 were not affected (Fig. 4g), exendin-4-
induced OMM-localised PKA activity was blunted by SPHKAP downregulation (Fig. 4h).
Finally, we employed a similar modified FluoSTEP system to the one shown in Fig. 2, this
time adapted to assess VAPB-localised PKA activity in response to exendin-4 stimulation in
Control *versus* SPHKAP RNAi-treated WT INS-1 832/3 cells (see Supplementary Fig. 7d for
a schematic of this method). Results showed that exendin-4 stimulation triggers VAPB-
localised PKA activity in WT INS-1 832/3 cells, and that this activity requires SPHKAP (Fig.
4i), presumably via SPHKAP binding to PKA-RI and assembly of a PKA signalling hub at
ERMCSs. Consequently, we could observe the appearance of PKA-RIα-EGFP-positive
puncta, indicative of liquid-liquid phase separation (LLPS) and biomolecular condensate
assembly[25], in exendin-4-stimulated WT INS-1 832/3 cells (Supplementary Fig. 7e).
396
397 **Control of mitochondrial and β-cell function by SPHKAP-dependent GLP-1R**
398 **signalling.**
399
We next assessed the effect of ERMCS-localised GLP-1R-induced PKA signalling in the
control of mitochondrial function in INS-1 832/3 cells. Exendin-4 stimulation triggered a small
increase in mitochondrial membrane potential under 6 mM glucose conditions, with this
response further enhanced under increased (20 mM) glucose concentrations. Both
membrane potential rises were significantly reduced in SPHKAP RNAi-treated *versus*
Control WT INS-1 832/3 cells (Fig. 5a). Accordingly, we observed a loss of ATP tone in
exendin-4-stimulated INS-1 832/3 cells following SPHKAP knockdown (Fig. 5b,
Supplementary Fig. 7f), suggesting a defect in GLP-1R-dependent mitochondrial respiration.
We further determined the impact of ERMCS-localised GLP-1R signalling on β-cell function
by analysing acute insulin secretion responses and β-cell survival to ER stress downstream
of GLP-1R stimulation in SPHKAP RNAi-treated *versus* Control WT INS-1 832/3 cells.
Exendin-4-stimulated insulin secretion was significantly reduced in SPHKAP RNAi *versus*
control conditions (Fig. 5c, Supplementary Fig. 7g), without any effect of SPHKAP over-
expression in this parameter (Supplementary Fig. 7h). Additionally, the anti-apoptotic effect
of exendin-4 in cells exposed to thapsigargin (to trigger ER stress) was blunted following
SPHKAP knockdown (Fig. 5d). Finally, we measured exendin-4-induced mitochondrial and
ER Ca^{2+} responses in Control *versus* SPHKAP RNAi-treated WT INS-1 832/3 cells.
Increases in both ER and mitochondrial Ca^{2+} in response to exendin-4 stimulation were
apparent in these cells. However, while ER responses were not affected by SPHKAP
knockdown, we detected a delay in the mitochondrial Ca^{2+} response peak time in SPHKAP
RNAi *versus* Control RNAi-treated WT INS-1 832/3 cells (Supplementary Fig. 7i,j).
421
As SPHKAP has previously been shown to bind to the mitochondrial cristae junction
regulator and MICOS complex component MIC19[40], a prominent PKA substrate involved in
both cristae assembly during mitochondrial remodelling[43] and in regulation of ERMCSs, we
next tested whether exendin-4 stimulation would trigger MIC19 phosphorylation in WT INS-1
832/3 cells using a MIC19 immunoprecipitation strategy coupled with Western blotting with
an antibody raised against PKA phosphosites (RRxpS/T)[44]. Exendin-4 exposure resulted in a
~5-fold increase in PKA-specific MIC19 phosphorylation, an effect that was potentiated by
the adenylate cyclase activator forskolin (Fig. 5e). Dual knockdown of SPHKAP and VAPB in
INS-1 832/3 cells led to widespread changes in PKA phosphorylation of several MICOS
complex components downstream of GLP-1R activation, suggesting that ERMCS-localised
GLP-1R signalling might modulate the activity of the MICOS complex (Supplementary Fig.
7k).
434
435 **Control of mitochondrial morphology by SPHKAP-dependent ERMCS-localised GLP-**
436 **1R signalling.**
437

As the main role of the MICOS complex is to orchestrate mitochondrial cristae assembly[45],
and MIC19 plays an important role in the transfer of lipid precursors across ERMCSs during
mitochondrial remodelling[10], we next assessed whether GLP-1R stimulation would play a
role in mitochondrial remodelling processes in INS-1 832/3 cells, and whether this would be
affected by SPHKAP knockdown. We first investigated if acute exendin-4 stimulation triggers
changes in mitochondrial turnover. Despite no detectable changes in mitochondrial DNA
copy number after 30 minutes of exendin-4 exposure in Control *versus* SPHKAP RNAi-
treated, or in GLP-1R KO INS-1 832/3 cells (Supplementary Fig. 7l), acute (10-minute)
stimulation of WT INS-1 832/3 cells with exendin-4 triggered increased mitophagy, assessed
using an mKeima-based assay, an effect not present following SPHKAP downregulation
(Fig. 5f). Further evidence of exendin-4-induced mitophagy was obtained by CLEM analysis
of INS-1 832/3 SNAP/FLAG-hGLP-1R cells, where we could find instances of SNAP/FLAG-
hGLP-1R-positive endolysosomes surrounded by mitochondria (Supplementary Fig. 8a,b).
We also detected increased recruitment of the mitochondrial fission protein mCherry-Drp1 to
mito-BFP-expressing mitochondria in exendin-4-stimulated *versus* vehicle-exposed WT INS-
1 832/3 cells (Fig. 5g, Supplementary Fig. 8c, Supplementary Movies 4 and 5). Drp1
phosphorylation at Serine 616, which, unlike Serine 637[46], is important for regulating
mitochondrial fission and is not directly PKA-dependent[47, 48], was, as expected, not modified
by acute exendin-4 exposure in Control RNAi-treated cells, but was however progressively
lost 15 minutes post-exendin-4 stimulation in SPHKAP RNAi-treated WT INS-1 832/3 cells
(Fig. 5h).
459
We next directly assessed changes in mitochondrial morphology by confocal microscopy
imaging of MitoTracker-labelled mitochondria in vehicle *versus* exendin-4-stimulated WT
INS-1 832/3 cells. Here, we employed a similar approach to [49] to manually score cells by
mitochondrial phenotype into three categories: long tubular (referred to as "hyperfused"),
intermediate/short tubular (referred to as "tubular") and fragmented. Results showed that 10
minutes of exendin-4 exposure completely remodels the mitochondrial network from a
predominantly hyperfused to a fragmented phenotype (Fig. 6a). As expected, and serving as
validation of our classification approach, this remodelling did not take place in INS-1 832/3
cells lacking GLP-1R (GLP-1R KO) (Fig. 6b). We further confirmed our results by re-
analysing the data from above using the automated Mitochondria Analyzer plugin in Fiji to
determine changes in mitochondrial form factor, which provides information on mitochondrial
shape, as well as mitochondrial mean area (Supplementary Fig. 8d). Agreeably, we found a
significant reduction on mitochondrial form factor and mean area, indicative of remodelling,
upon exendin-4 stimulation of INS-1 832/3, but not of INS-1 832/3 GLP-1R KO cells.
474
Next, we performed mitochondrial morphology analysis of WT INS-1 832/3 cells treated with
Control *versus* SPHKAP RNAi prior to vehicle or exendin-4 exposure (Fig. 6c). While
exendin-4 triggered a similar degree of mitochondrial network remodelling as above in
Control RNAi-treated cells, SPHKAP knockdown resulted in disrupted mitochondrial
morphology at vehicle conditions, which somewhat resembled that of exendin-4-stimulated
Control RNAi-treated cells in terms of tubularity but had further visual abnormalities.
Moreover, exendin-4 stimulation had no effect on the mitochondrial morphology of SPHKAP
RNAi-treated cells. These results were validated with the Mitochondria Analyzer plugin in Fiji,
which showed reduced mitochondrial form factor and mean area following exendin-4
exposure of Control RNAi-treated cells, with SPHKAP RNAi-treated cells under vehicle
conditions showing similar values to those of exendin-4-stimulated control cells
(Supplementary Fig. 8e). However, with this analysis method, we also detected an apparent
increase in form factor and mitochondrial mean area in exendin-4-stimulated *versus* vehicle-
treated SPHKAP RNAi-treated cells, an effect opposite to that triggered by exendin-4 in
control conditions.
490
To better visualise the changes in mitochondrial morphology under these different conditions,
we performed z-series analysis of confocal sections across whole cells, represented as MIPs

in Fig. 6d, as well as Stimulated Emission Depletion (STED) microscopy analysis of
mitochondrial morphologies using the fixable STED-compatible mitochondrial dye PKmito
ORANGE FX (Fig. 6e), with Control *versus* SPHKAP RNAi-treated cells showing similar
changes in mitochondrial morphology as those described above.
497
Finally, we also performed high-resolution TEM imaging for precise quantification of
mitochondrial area and number of cristae per mitochondria in ultrathin (70-nm) sections on
resin-embedded Control *versus* SPHKAP RNAi-treated WT INS-1 832/3 cells, as well as
INS-1 832/3 GLP-1R KO cells, both under vehicle conditions and following exendin-4
stimulation. In agreement with our previous results, we measured an exendin-4-induced
reduction in both parameters in Control RNAi-treated cells, but no change in response to
exendin-4 in either SPHKAP RNAi-treated or in GLP-1R KO cells (Fig. 7a,b). Interestingly,
the mitochondrial morphology changes triggered by exendin-4 in Control RNAi-treated cells
were accompanied by a significant reduction in ERMCS length, which appeared already
reduced under vehicle conditions without further effect of exendin-4 following SPHKAP
downregulation or in GLP-1R KO cells, with no apparent effect of SPHKAP-EGFP
overexpression in INS-1 832/3 cells in this parameter (Fig. 7c). This ERMCS length
reduction was accompanied by a reduced number of ERMCSs per mitochondrial area in
exendin-4-stimulated *versus* vehicle-exposed WT INS-1 832/3 cells, assessed using a
splitFAST approach as in [50] (Fig. 7d,e).
513
514   **GLP-1R localisation and GLP-1R-induced PKA signalling at ERMCSs in primary**
515   **mouse and human islets.**
516
To validate the interaction of GLP-1R with VAPB and SPHKAP at ERMCSs in a
physiologically relevant setting with endogenous levels of receptor expression, proximity
ligation assays (PLAs) were performed in purified mouse primary islets using one of the few
existing validated mouse GLP-1R antibodies from the Developmental Studies Hybridoma
Bank[51, 52, 53], with GLP-1R–VAPB and GLP-1R–SPHKAP interactions readily observed in
exendin-4-stimulated islets (Fig. 8a, Supplementary Fig. 9a). To verify that these interactions
are present within pancreatic β-cells, we counterstained both vehicle and exendin-4-
stimulated islets for insulin (Supplementary Fig. 9b). Next, to analyse the functional impact of
SPHKAP downregulation on islet function, we employed a lentiviral shRNA approach to
knock down SPHKAP on intact islets, as most functional outputs from primary islets cannot
be optimally recapitulated with dispersed islet cells due to the loss of important intra-islet
cell-cell interactions. Lentiviral transduction of cells within intact islets is restricted to the
outermost islet cell layers so that only a small fraction of cells (which we estimate at ~25% of
the total) is effectively transduced. However, those represent a higher functional cell pool
due to necrosis of the islet core in non-vascularised isolated islets[54], so that it is still possible
to assess potential functional differences from a relatively small percentage of transduced
cells. Accordingly, we could only detect a tendency for reduced *SPHKAP* mRNA levels in
whole islets transduced with SPHKAP *versus* Control shRNA lentiviruses (Supplementary
Fig. 9c), which nevertheless suggests a good level of downregulation from the pool of
transduced cells. We next analysed some key signalling outcomes from these islets. As for
INS-1 832/3 cells, ATP levels in exendin-4-stimulated islets were reduced with SPHKAP
shRNA (Fig. 8b). Exendin-4 exposure also led to a significant reduction in the level of
reactive oxygen species (ROS) in Control shRNA islets exposed to thapsigargin, but this
effect was no longer present in SPHKAP shRNA-transduced islets, which presented with
basally higher ROS levels than the corresponding control islets (Fig. 8c). Finally, while the
effect of SPHKAP knockdown in the acute potentiation of insulin secretion triggered by
exendin-4 in these islets was mild, we found a small but significant increase in insulin
secretion at 11 mM glucose alone in SPHKAP *versus* Control shRNA-transduced islets (Fig.
8d, left), which led to a near-significant reduction in exendin-4-induced insulin secretion fold
(Fig. 8d, right), an effect which agrees with previously published results in a SPHKAP KO

mouse model, with increased insulin secretion under glucose alone conditions but
dampened incretin responses in islets from these mice[34].
549
We next assessed changes in mitochondrial morphology in intact mouse islets by confocal
imaging of mitochondria from the outermost layer of islet cells, to maximise our chances of
recording phenotypes from lentivirally transduced cells. We used a similar approach as in
INS-1 832/3 cells to segment and classify islet cells as predominantly presenting a
hyperfused, tubular or fragmented mitochondrial phenotype (see Supplementary Fig. 9d for
an example of our method of islet segmentation and cell classification). Cells in WT mouse
islets presented a similar distribution of mitochondrial morphologies as in INS-1 832/3 cells,
with mitochondria primarily hyperfused or tubular under vehicle conditions but remodelling
towards a more fragmented phenotype following exendin-4 exposure (Fig. 9a). Islets
extracted from GLP-1R KO mice presented a higher level of hyperfused mitochondria that
did not undergo remodelling with exendin-4 (Fig. 9b). As in INS-1 832/3 cells, transduction of
WT islets with SPHKAP shRNA-expressing lentiviruses triggered a change in mitochondrial
morphology under vehicle conditions, with a higher degree of fragmentation *versus* control
islets and no effect of exendin-4 exposure following SPHKAP downregulation (Fig. 9c).
564
We also performed mitochondrial morphology assessments in dispersed islet cells from mice
transfected with Control *versus* SPHKAP siRNA, which responded similarly to intact islets in
terms of increased mitochondrial fragmentation following exendin-4 exposure in control
conditions, but showing disturbed mitochondrial morphology under vehicle conditions and no
effect of exendin-4 exposure in SPHKAP siRNA-treated cells (Fig. 9d). These phenotypic
changes were still present when assessed as changes in mitochondrial form factor and
mean area using the Mitochondria Analyser Fiji plugin with data from these dispersed islet
cells (Supplementary Fig. 9e).
573
Lasty, we repeated some of our assessments with human donor islets. PLA assays again
demonstrated interactions between endogenous GLP-1R, VAPB and SPHKAP in exendin-4-
stimulated human islets (Fig. 10a,b). As in mouse islets, assessment of mitochondrial
morphology by confocal microscopy analysis in intact human islets revealed predominantly
hyperfused mitochondria under vehicle conditions which was rapidly remodelled by acute
exendin-4 exposure to a more fragmented morphology (Fig. 10c,d).
580
581 **Discussion**
582
Mitochondrial malfunction is a prominent feature of diabetes, encompassing defects in
mitochondrial structure and dynamics, ROS overproduction and apoptosis, breakdown of
nutrient sensing and metabolic uncoupling[55, 56], with strong backing evidence from human
genetic studies[57, 58, 59]. Mitochondrial quality control mechanisms have been recently found to
control cellular identity and maturity in metabolically relevant tissues[60], including in
pancreatic β-cells, highlighting the importance that mitochondria have in the control of β-cell
function. Strategies to correct the detrimental effect of T2D in affected tissues are therefore
likely to involve amelioration of mitochondrial function; GLP-1RAs are no exception, with
numerous reports highlighting their capacity to potentiate mitochondrial quality control
mechanisms and improve mitochondrial health[5, 6, 7, 8], with mitochondrial regulation
potentially key to explain their beneficial effects in metabolic tissues.
594
Despite the accumulating evidence around pathways engaged by the GLP-1R and other
GPCRs in target tissues, there is still substantial uncertainty about the precise molecular
mechanisms engaged, particularly with regards to the spatiotemporal organisation of their
signalling and their presence in spatially restricted subcellular (and even suborganelle)
signalling hubs or nanodomains[61]. In this context, the study of MCSs has become an area of
considerable importance to understand the spatial architecture of signalling, with recent
reports recognising their prominent role in pancreatic β-cells for key processes ranging from

insulin granule biogenesis to calcium homeostasis[62, 63, 64]. In this study, we took an unbiased
approach to unveil new molecular partners that might work in concert with the GLP-1R to
control β-cell function, finding not only that the receptor engages with ER-endosome MCSs
to organise its signalling, but that the main constituents of these inter-organelle contacts,
namely the VAPs, are by a considerable margin the most prominent intracellular binding
partners of active GLP-1R. This suggests that engagement of VAPs, and subsequent
generation of ER-localised GLP-1R signalling, is not just one of many pathways employed
by the receptor, but the main site of action of endosomal GLP-1R following its internalisation
from the plasma membrane.

This also suggests that GLP-1RAs with distinct rates of receptor internalisation and
lysosomal targeting might differentially regulate the kinetics of this engagement, so that slow
GLP-1R internalising compounds are expected to trigger initially reduced VAPB interactions,
which might however paradoxically result in prolonged VAPB engagement over time, and
therefore potentially sustained ER signalling, as these biased compounds are also
associated with reduced lysosomal degradation from the pool of internalised receptors[18, 24].
Of note, we have found here a similar protein-protein interaction between SPHKAP and GIP-
stimulated GIPR, suggesting that compounds such as tirzepatide, which target the GLP-1R
as a biased but the GIPR as a balanced agonist, might present with an optimal signalling
profile by means of maximising prolonged plasma membrane signalling from the GLP-1R
while at the same time engaging in GIPR-dependent acute ER signalling.

In the present study, we have not only unveiled the interaction of endosomal GLP-1R with
the ER, but also its localisation to VAPB-containing ERMCSs, from where the receptor
assembles a newly described PKA signalling nanodomain, enabling cAMP-dependent PKA
activation at the interface between the ER and mitochondria. This ERMCS-localised GLP-1R
signalling hub requires the engagement of the AKAP SPHKAP, known to assemble PKA-RIα
aggregates at the ER[32, 33], with VAPB playing a central role as the tethering factor
responsible for the separate recruitments of GLP-1R and SPHKAP to ERMCSs, the latter via
the direct interaction of VAPB with the pFFAT motif of SPHKAP. It is important to highlight
that SPHKAP is one of the few AKAPs known to recruit PKA-RI instead of RII subunits[40].
This is highly relevant as PKA-RIα aggregates undergo LLPS to form biomolecular
condensates[65] which spatially restrict PKA activity, leading to signal compartmentalisation[25],
explaining why the presence of endosomal GLP-1R at ERMCSs is required to assemble this
signalling hub, and uncovering a new modality of GLP-1R signal transduction which involves
the local assembly of a biomolecular condensate containing not only signalling mediators
such as SPHKAP/PKA, and also presumably adenylate cyclase, but downstream targets for
PKA phosphorylation such as the MICOS complex and other ERMCS components, some
likely to be amongst the factors identified as active GLP-1R interactors in our interactome.
An additional level of regulation could be provided by kinases and/or phosphatases
controlling VAPB – SPHKAP – PKA-RIα assembly/disassembly, such as DYRK1A, recently
shown to be recruited to VAPB/SPHKAP/ PKA-RIα condensates in neurons[33] and previously
identified in a kinome study to be regulated by GLP-1R activity[66], as well as by the
autophagic degradation of the VAPB – SPHKAP – PKA-RIα complex via the interaction of
VAPB/SPHKAP with its paralog AKAP11 and GSK3α/β, a pathway recently identified in
neurons as linked to the control of neurotransmission, though to be defective in AKAP11-
associated psychiatric diseases[33].

The abovementioned signalling hub appears to be equally important for the control of β-cell
function, a notion supported by human genetic data of genome-wide significance unveiling
an association between *SPHKAP* gene variants, random blood glucose levels and T2D, and
high BMI. As we observe a correlation between *SPHKAP* variant effects in T2D and high
BMI (as a proxy for obesity), we propose that SPHKAP might play a pleiotropic role in both
disorders by the engagement of common signalling pathways in different tissues. These

could involve the GLP-1R, known to signal in anorectic neurons to regulate food intake as
well as in β-cells to potentiate insulin secretion and β-cell survival[67]. SPHKAP might likely
additionally engage with other metabolically relevant GPCRs via VAPB in tissues where this
AKAP is expressed, including the pancreas, brain and heart[29, 32], amongst others. For
example, this mechanism might potentially regulate liver mitochondrial function, a possibility
supported by the previously published interaction between the closely related glucagon
receptor and VAPB in hepatocytes[68]. The significance of the ERMCS-localised signalling hub
identified here is therefore likely to extend well beyond the GLP-1R to other GPCRs, for
example GIPR, as shown here. Accordingly, beyond the effect observed in exendin-4-
induced responses, we find a significant impact of SPHKAP downregulation in mitochondrial
morphology and other parameters already present under vehicle conditions, suggesting
additional effects of ER-localised PKA signalling in β-cells.
668
Another outcome from the present study is the identification of the MICOS complex protein
MIC19 as a downstream target of ERMCS-localised GLP-1R-induced PKA phosphorylation.
MIC19 is not only a key member of mitochondrial cristae junctions, responsible for the
organisation of mitochondrial inner and outer membrane contacts, as well as for the
architecture of respiratory complexes and the proton pump, but also plays a prominent role
in the transfer of lipid precursors across ERMCSs for the biosynthesis of the key
mitochondrial lipid cardiolipin[69], therefore connecting mitochondrial cristae organisation with
the generation of membrane curvature and control of mitochondrial function afforded by
cardiolipin[10]. We therefore hypothesise that the GLP-1R-dependent phosphorylation of
MIC19 at ERMCSs might not only highlight a direct role for the GLP-1R in the control of
mitochondrial dynamics, but also in the regulation of lipid precursor transfer across ERMCSs
to sustain mitochondrial turnover. This is particularly relevant as alterations in cardiolipin
profiles have been reported in several diseases where GLP-1RAs play beneficial roles,
including Alzheimer's, Parkinson's, obesity, and T2D[70]. Interestingly, our MS interactomic
analysis identifies TAMM41, an essential enzyme for the biosynthesis of cardiolipin[71], as an
active GLP-1R interactor.
685
Amongst the mitochondrial processes modulated by SPHKAP-dependent ERMCS-localised
GLP-1R cAMP/PKA signalling, we have found acute increases in mitochondrial fission and
mitophagy, two key quality control mechanisms which enable the clearance of damaged,
unhealthy, or old mitochondria, and hold a prominent role in the maintenance of β-cell
function[56] and identity[60]. Loss of these functions can lead to impaired insulin release and
glucose intolerance[72], with the present study indicating that controlling these processes is an
important mechanism engaged by the GLP-1R to exert its beneficial action on β-cell function
and survival under oxidative stress conditions.
694
Amongst the major metabolic triggers of β-cell dysfunction is glucose and lipid-induced
cytotoxicity. In the context of obesity, elevated glucose and fatty acids increase insulin
demand and contribute to β-cell oxidative stress. Additionally, excess free fatty acids
accumulated in β-cells induce further ROS generation, contributing to increase ER stress
and β-cell dysfunction. β-cell mitochondria therefore need to appropriately adapt to
glucolipotoxic conditions to avoid progression towards overt T2D[73]. Changes in
mitochondrial morphology have been previously associated with modified nutrient utilisation,
with a strong correlation between mitochondrial fragmentation and increased fatty acid
oxidation (FAO)[74], a process normally inhibited in β-cells under high glucose conditions to
favour metabolic coupling towards insulin secretion[73, 75], but that might be differentially
modulated by the GLP-1R under excess lipid conditions to enable lipid clearance and
detoxification[73]. In support of this hypothesis, our MS interactome indicates the association
of active GLP-1R with a range of factors involved in lipid metabolism, including lipases and
regulators of lipase transfer to lipid droplets such as Mest[76] and Copb1[77], lipid metabolising
enzymes including TMEM120A[78], Acsl4[79], Fads1[80], Chp1[81], Cds2[82], and Apgat4[83], and
mitochondrial FAO and ketone body enzymes (HADHA/B[84], Bdh1[85]), as well as the ER-lipid

droplet small GTPase Rab18[86], several subunits of the mitochondrial ATP synthase, the
respiratory chain subunit SDHA, and the FAO-fuelled plasma membrane Na^+/K^+ pump[87].
Interestingly, a recent study has identified ERMCSs as required for lipid radical transfer from
mitochondria to the ER to reduce mitochondrial oxidative stress[88], which could represent
another mechanism engaged by ERMCS-localised GLP-1R to enhance β-cell survival under
glucolipotoxic conditions, and the SPHKAP paralog AKAP11 as playing an important role in
regulating lipid metabolism in astrocytes[89].
718
Beyond its interaction with lipid enzymes, our MS interactome analysis also indicates that
the GLP-1R binds to ERMCS-localised Sarcoendoplasmic Reticulum Calcium ATPase
(SERCA) pump in the ER and voltage-dependent anion-selective channel protein 2 (VDAC2)
in the OMM, both involved in calcium/ion transfer across ERMCSs[90], suggesting a prominent
role for the receptor in the control of intra-organelle ion homeostasis, with important
implications for the regulation of mitochondrial function, ER stress responses and insulin
secretion[91]. Overall, all these MS-identified interactions suggest that the effect of GLP-1R
PKA signalling at ERMCS is likely to extend well beyond the MICOS complex into a more
comprehensive programme of ER, mitochondria and lipid metabolic pathway regulation.
728
In summary, this investigation has unveiled the existence of a newly characterised inter-
organelle signalling hub in pancreatic β-cells which involves active endosomal GLP-1R
localisation to ERMCSs via VAPB binding, and engagement of the AKAP SPHKAP to enable
ERMCS-restricted cAMP/PKA activity via the assembly of PKA-RIα biomolecular
condensates, leading to PKA phosphorylation of the MICOS complex, enhanced
mitochondrial remodelling and improved mitochondrial function, with important implications
for the modulation of β-cell function and survival, and the potential to serve as a mechanism
for GLP-1R-induced mitochondrial adaptation during metabolic stress and excess nutrient
exposure in β-cells. Further studies to investigate this later possibility, including examination
of the effect of GLP-1R in β-cell lipid metabolism and mitochondrial nutrient utilisation under
normal and glucolipotoxic stress conditions are warranted, as well as the investigation of the
in vivo role of SPHKAP in GLP-1R-dependent glucoregulation.
741
742 **Methods**
743
744 *Cell culture*
745
INS-1 832/3 rat insulinoma cells were maintained in RPMI-1640 at 11 mM D-glucose,
supplemented with 10% fetal bovine serum (FBS), 10 mM HEPES, 2 mM L-glutamine, 1 mM
pyruvate, 50 µM β-mercaptoethanol and 1% penicillin/streptomycin. INS-1 832/3 cells stably
expressing SNAP/FLAG-tagged human GLP-1R (hGLP-1R) or human GIPR (hGIPR) were
generated by transfection of pSNAP-GLP-1R or pSNAP-GIPR (Cisbio) into INS-1 832/3 cells
lacking endogenous GLP-1R or GIPR expression (INS-1 832/3 GLP-1R or GIPR KO, a gift
from Dr Jacqueline Naylor, AstraZeneca[92]) followed by G418 (1 mg/mL) selection.
Mycoplasma testing was performed biannually. Cell culture reagents were obtained from
Sigma or Thermo Fisher Scientific.
755
756 *Peptides and fluorescent probes*
757
GLP-1 was purchased from Bachem, semaglutide was from Imperial College London
Healthcare NHS Trust pharmacy, and tirzepatide was a gift from Sun Pharmaceuticals.
Custom analogues were from WuXi AppTec, China, and danuglipron and orforglipron were
purchased from MedChemExpress. SNAP-Surface fluorescent probes were from New
England Biolabs and MitoTracker™ Red CMXRos and ER-Tracker Red were from Thermo
Fisher Scientific.
764

GLP-1R β-cell interactome

INS-1 832/3 SNAP/FLAG-hGLP-1R cells were stimulated in 6-cm dishes with vehicle or 100
nM GLP-1RA (GLP-1, exendin-4, exendin-F1 or exendin-D3) for 5 minutes. Following
stimulation, cells were washed in PBS and osmotically lysed in lysis buffer (50 mM Tris-HCl,
pH 7.4, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA) with 1% protease inhibitor (Roche)
and 1% phosphatase inhibitor cocktail (Sigma-Aldrich) at 4°C. Cell lysates were centrifuged
at 12,000 g and 4°C for 10 minutes, prior to addition of 40 µL anti-FLAG M2 beads (A2220,
Sigma-Aldrich) for overnight rotating incubation at 4°C to increase binding efficiency.
Samples were centrifuged at 5,000 g for 30 seconds, and bead pellets washed 3x in 500 µL
wash buffer (50 mM Tris-HCl, 150 mM NaCl, pH 7.4) before storage at -80°C until further
analysis.

Beads were resuspended in 150 µL 1 M urea in 20 mM HEPES pH 8.0 with 1 mM DTT,
followed by addition of 1.5 µg Trypsin gold (Promega) with overnight digestion at 37°C.
Supernatants were subsequently acidified with 0.5% trifluoroacetic acid (TFA), followed by
desalting using reversed-phase spin tips (Glygen Corp) and drying using vacuum
centrifugation. Dried peptides were resuspended in 0.1% TFA and Liquid Chromatography
Tandem Mass Spectrometry (LC-MS/MS) performed as previously described[93] using an
Ultimate 3000 nano-HPLC coupled to a Q-Exactive mass spectrometer (Thermo Scientific)
via an EASY-Spray source. Peptides were loaded onto a trap column (Acclaim PepMap 100
C18, 100 µm × 2 cm) at 8 µL/minute in 2% acetonitrile and 0.1% TFA. Peptides were then
eluted on-line to an analytical column (Acclaim Pepmap RSLC C18, 75 µm × 75 cm). A 90-
minute stepped gradient separation was used with 4-25% acetonitrile 80%, formic acid 0.1%
for 60 minutes, followed by 25-45% for 30 minutes. Eluted peptides were analysed by Q-
Exactive operating in positive polarity and data-dependent acquisition mode. Ions for
fragmentation were determined from an initial MS1 survey scan at 70,000 resolution (at m/z
200), followed by higher-energy collisional dissociation of the top 12 most abundant ions at a
resolution of 17,500. MS1 and MS2 scan AGC targets were set to 3e6 and 5e4 for a
maximum injection time of 50 and 75 miliseconds, respectively. A survey scan range of 350 –
1800 m/z was used, with a normalised collision energy set to 27%, minimum AGC of 1e3,
charge state exclusion enabled for unassigned and +1 ions and a dynamic exclusion of 45
seconds.

Data was processed using the MaxQuant software platform (v1.6.10.43), with database
searches carried out by the in-built Andromeda search engine against the Swissprot Rattus
Norvegicus database (Downloaded – 21st May 2022, entries: 8,132) concatenated with the
human GLP-1R protein sequence. A reverse decoy database approach was used at a 1%
FDR for peptide spectrum matches and protein identifications. Search parameters included:
maximum missed cleavages set to 2, variable modifications of methionine oxidation, protein
N-terminal acetylation, asparagine deamidation and cyclisation of glutamine to
pyroglutamate.

Label-free quantification was enabled with an LFQ minimum ratio count of 1. Proteins were
filtered for those containing at least one unique peptide and identified in all 4 biological
replicates. For normalisation, proteins were ranked by fold change increase in abundance
compared with control (vehicle-treated) immunoprecipitates. The resulting data was pre-
processed by elimination of proteins that are non-specifically bound and residual background
proteins, such as common contaminants and reverse database proteins. To further facilitate
the selection of GLP-1R interactors, the following criteria were established; i) identified at
least twice from four runs; ii) not nuclear proteins; and (iii) previously identified in pancreatic
β-cells.

Untargeted label-free quantitation (LFQ) intensity values, normalised to hGLP-1R levels in
each immunoprecipitated sample, were analysed by LFQ Analyst[94] to identify differences in

levels of individual protein-GLP-1R interactions between vehicle and GLP-1RA-treated
conditions. Paired t-tests adjusted for multiple comparisons using Benjamin-Hochberg false
discovery rate correction and Perseus type imputations were used for statistical analysis. To
generate novel information about the functional pathways represented by these potential
GLP-1R interactors, we performed pathway enrichment analysis with GLP-1R binding
partners exhibiting changes between vehicle and exendin-4-stimulated conditions using
gProfiler.
827
828 *cDNA and siRNA transfection*
829
830 INS-1 832/3 cells were transfected with relevant amounts of plasmid cDNA outlined in Table
831 1, or siRNA outlined in Table 2, using Lipofectamine 2000 (Life Technologies) in media
832 without antibiotics according to the manufacturer's instructions. Experiments were carried
833 out 24-48 hours after plasmid cDNA, or 72 hours after siRNA transfection.
834
835

Name	Origin	Assay
EGFP-VAPB	Gift from Dr Emily Eden, UCL	Co-Immunoprecipitation, confocal microscopy
Venus-VAPB	Generated *in house*	NanoBRET
EGFP-VAPB P56S	Generated *in house*	Co-immunoprecipitation, insulin secretion
GLP-1R-NanoLuc	Generated *in house*	NanoBRET
pcDNA3.1+		Control empty plasmid
Dynamin1 K44A-EGFP	Addgene #34681	Co-Immunoprecipitation, confocal microscopy
Dynamin2 K44A-mRFP	Addgene #128153	Co-Immunoprecipitation, confocal microscopy
SPHKAP-EGFP	Gift from Prof John D. Scott, Howard Hughes Medical Institute[40]	Co-Immunoprecipitation, confocal microscopy
SPHKAP pFFAT mut (DFLTESE)-EGFP	Generated *in house*	Co-Immunoprecipitation, confocal microscopy
pcDNA3-mitoExRai-AKAR2	Addgene #161757	Mitochondrial PKA
pcDNA3.1(+)-ExRai-AKAR2	Addgene #161753	Global PKA
cyto-Ruby3-iATPSnFR1.0	Addgene #102551	ATP
MIC19-3xHA	Gift from Prof Toshihiko Oka, Rikkyo University[95]	Immunoprecipitation, PKA-dependent phosphorylation
SPLICS Mt-ER Long P2A	Addgene #164107	Confocal microscopy of ERMCSs
Proinsulin-NLuc	Addgene #62057	Insulin secretion
mKeima-Red-Mito-7	Addgene #56018	Mitophagy
pCAG G-CEPIA1er	Addgene #105012	ER calcium
pCAG mito-RCaMP1h	Addgene #105013	Mitochondrial calcium
pEGFPC1-hVAP-B	Addgene #104448	Cloning of pcDNA3-GFP11(x7)-VAPB

pcDNA3-GFP11(x7)-Actin	Addgene #181967	Cloning of pcDNA3-GFP11(x7)-VAPB
pcDNA3-GFP11(x7)-VAPB	Generated *in house*	FluoSTEPs assays
mito-BFP	Addgene #49151	Mitochondrial fission
mCh-Drp1	Addgene #49152	Mitochondrial fission
pcDNA3.1(+)-FluoSTEP-ICUE	Addgene #181845	FluoSTEP cAMP
pcDNA3.1(+)-FluoSTEP-AKAR	Addgene #181844	FluoSTEP PKA
4xMTS-Halo	Gift from the Avezov group, UK DRI	Mitochondrial imaging for splitFAST data
ER-RspA-NFAST and OMM-short-RspA-CFAST	Gifts from the Filadi lab, CNR Institute of Neuroscience[50]	ERMCS imaging for splitFAST data
pcDNA3.1(+)-RIα-EGFP	Addgene #181840	Visualisation of PKA-RIα-EGFP biomolecular condensates

836
837 **Table 1:** Plasmid cDNA origin and use.
838

Name	Origin
Control siRNA	Ambion™ Silencer™ Select Negative Control #1 siRNA Cat#:4390843
VAPB siRNA	Ambion™ Silencer™ Select Pre-Designed siRNA Cat#:4390771, s133396
SPHKAP siRNA	Ambion™ Silencer™ Select Pre-Designed siRNA Cat#:4390771, s164929

839
840 **Table 2:** siRNA origin.
841
842 *Isolation and culture of pancreatic islets*
843
WT or GLP-1R KO mice (*in house*) were humanely euthanised before inflating the pancreas
through the common bile duct with RPMI-1640 medium containing 1 mg/mL collagenase
(Nordmark Biochemicals). The pancreas was then dissected, incubated at 37°C for 10
minutes, and digestion halted with cold RPMI-1640 supplemented with 10% FBS (F7524,
Sigma-Aldrich) and 1% (v/v) penicillin/streptomycin solution (15070-063, Invitrogen). Mouse
islets were subsequently washed, purified using a Histopaque gradient (Histopaque-1119
and Histopaque-1083, Sigma-Aldrich) and rested overnight at 37°C in a 95% O_2 / 5% CO_2
incubator before performing experiments in mouse islet culture medium (RPMI-1640 with 11
mM D-glucose, supplemented with 10% FBS and 1% penicillin/streptomycin).
853
Human islets were received from the isolation unit at the Centre Européen d'Etude du
Diabète (CEED), Strasbourg, or from isolation centres ascribed to the Integrated Islet
Distribution Program (IIDP), National Institute of Diabetes and Kidney Diseases (NIDDK),
National Institutes of Health (NIH), USA, with all ethical approvals in place. Human islets
were cultured on arrival in RPMI-1640 medium supplemented with L-glutamine, 10% FBS,
1% penicillin/streptomycin solution, 0.25 µg/mL amphotericin B, and 5.5 mM glucose at 37°C
in a 95% O_2 / 5% CO_2 incubator. Donor characteristics are indicated in Table 3.
861

DATE OF ARRIVAL	SEX	AGE	BMI	SOURCE	C.O.D.

14.12.23	M	34	24.4	CEED	Anoxia
29.01.24	M	71	29.75	CEED	Stroke
15.03.24	M	61	30	CEED	Stroke
23.10.24	F	59	38.8	CEED	Stroke
01.11.24	F	64	23.8	IIDP	Head trauma
15.11.24	F	23	32.5	IIDP	Head trauma
16.12.24	F	55	29.5	IIDP	Stroke

Table 3: Details of human islet donors.

Islet shRNA transduction

Mouse islets were transduced with Control or SPHKAP shRNA-expressing lentiviral particles (MISSION® shRNA, Target Clone ID TRCN0000088765 mouse *SPHKAP*, Sigma Aldrich) at an MOI of 1 and cultured for 72 hours before experiments.

Co-immunoprecipitation

INS-1 832/3 cells previously transfected with the relevant plasmid, or INS-1 832/3 SNAP/FLAG-hGLP-1R or SNAP/FLAG-hGIPR cells seeded in 6-well plates at 1.2×10^6 cells per well and incubated overnight. Cells were stimulated in normal media with 100 nM exendin-4 or vehicle for 5 minutes at 37°C, washed once with ice cold PBS and lysed with 1 mL lysis buffer supplemented with 1% phosphatase inhibitor cocktail (Sigma-Aldrich) and 1% complete EDTA-free protease inhibitor cocktail (Sigma-Aldrich). Cell lysates were centrifuged at 12,000 g for 10 minutes, and supernatants added to 40 µL anti-FLAG M2 Magnetic beads (Sigma Aldrich), anti-HA Magnetic beads (Thermo Fisher Scientific), or ChromoTek GFP-Trap® Magnetic Particles (Proteintech), and incubated at 4°C for 1 hour. Beads were then washed 3x in 500 µL wash buffer and proteins eluted with urea buffer (200 mM Tris-HCl, pH 6.8; 5% w/v SDS, 8 M urea, 100 mM dithiothreitol, 0.02% w/v bromophenol blue) at 37°C for 10 minutes before SDS-PAGE and Western blot analysis.

SDS-PAGE and Western blotting

Samples were loaded onto 10% SDS-polyacrylamide gels then transferred to 0.45-µm PVDF membranes and blocked with 5% skimmed milk [or 5% bovine serum albumin (BSA) for anti-phospho antibodies] in TBS buffer supplemented with 0.1% Tween 20 (TBS-T). Membranes were incubated with primary antibody overnight diluted in TBS-T/milk (or TBS-T/BSA for anti-phospho antibodies) at 4°C, washed in TBS-T and incubated for 1 hour at RT with HRP-conjugated secondary antibody in TBS-T/milk. Following a further 3x washes in TBS-T, membranes were developed with Clarity Western ECL substrate (BioRad) and imaged on a Biorad ChemiDoc or in a Xograph Compact X5 processor with Fuji medical X-ray films. Specific band densities were quantified in Fiji. The following antibodies were used (source, catalogue number, dilution): rabbit anti-VAPB (Antibodies.com, A14703, 1:5,000); rabbit anti-SNAP tag (New England Biolabs, P93105, 1:5,000); rabbit anti-SPHKAP (kind gift from Dr Nicholas Vierra, University of Utah[32], 36690, 1:1,000); mouse anti-Tubulin (Sigma Aldrich, T-5168, 1:2,000); rabbit anti-HA tag (Abcam, Ab9110, 1:1,000); rabbit anti-GFP (Proteintech, 50430-2-AP, 1:2,000); rabbit anti-phospho-PKA Substrate (RRXS*/T*) (Cell Signaling, 9624, 1:1,000); rabbit anti-Drp1 (D6C7) (Cell Signaling, 8570S, 1:1,000); rabbit anti-phospho-Drp1 Ser616 (D9A1) (Cell Signaling, 4494S, 1:1,000); mouse anti-human Tom20 (F-10) (Santa Cruz Biotechnology, sc-17764, 1:1,000); goat anti-mouse IgG-HRP (Abcam, ab205719, 1:5,000); goat anti-rabbit IgG HRP (Abcam, ab205718, 1:2,000).

907 *SNAP/FLAG-hGLP-1R localisation by confocal microscopy*
908
INS-1 832/3 SNAP/FLAG-hGLP-1R cells seeded onto 35-mm glass-bottom dishes (MatTek)
were transfected with the indicated constructs. SNAP-FLAG-hGLP-1R was labelled for 15
minutes at 37°C with SNAP-Surface Alexa Fluor 546 or 647 in full media prior to stimulation
with 100 nM exendin-4 and imaging in a Leica Stellaris 8 inverted confocal microscope with
a 63x/1.40 oil immersion objective with Lightning confocal super resolution modality from the
Facility for Imaging by Light Microscopy (FILM) at Imperial College London. Mouse islets
were transduced 24 hours prior to imaging with an adenoviral SNAP/FLAG-hGLP-1R vector
(custom made by VectorBuilder) at a MOI of 1 and subsequently labelled for 10 minutes at
37°C with SNAP-Surface Alexa Fluor 647 before 5-minute stimulation with indicated agonists
and imaging of whole islets in the same microscope as above. Images were analysed in Fiji.
919
920 *MINFLUX microscopy*
921
INS-1 832/3 SNAP/FLAG-hGLP-1R cells were seeded onto 18-mm coverslips, labelled with
SNAP-Surface Alexa Fluor 647 and stimulated with 100 nM exendin-4 as above. Cells were
then fixed in 4% paraformaldehyde and processed for immunofluorescence with primary
rabbit anti-VAPB antibody (Sigma Aldrich, HPA013144, 1:150) followed by secondary anti-
rabbit FLUX 680 antibody (custom-made at Abberior, 1:200). Directly before MINFLUX
imaging, samples were incubated with 150 nm gold fiducials (BBI solutions) for 5 minutes.
After three brief washes with PBS, the samples were incubated with 10 mM $MgCl_2$ in PBS
for a further 5 minutes and then mounted on a well slide in freshly prepared GLOX buffer
(100 mM Tris-HCl pH 8, 10% w/v glucose, 0.4 mg/mL glucose oxidase, 64 ug/mL catalase)
supplemented with 10 mM MEA and sealed with two part silicon (eco-sil speed, Picodent).
932
MINFLUX imaging was performed on an Abberior instruments 3D MINFLUX microscope[14].
Briefly, the microscope is built on a motorized inverted microscope body (IX83, Olympus)
with a CoolLED pE-4000 for epifluorescence illumination. The system is equipped with 405
nm, 485 nm, 561 nm, and 642 nm laser lines, as well as a 980 nm IR laser and xyz piezo
(Piezoconcept) for active sample stabilisation. Images were acquired with a 100X, 1.45 NA
UPLXAPO oil objective (Olympus). During acquisition, the pinhole was set to 0.83 AU.
Detection was performed with two APDs in spectral windows Cy5 near (650 – 685 nm) and
Cy5 far (685 – 740 nm). MINFLUX imaging was performed with the default 2D imaging
sequence supplied with the system. In brief, the sequence defines parameters used in the
iterative localising process of MINFLUX, culminating in a final iteration with a targeted
coordinate pattern (TCP) diameter of 40 nm and minimum photon threshold of 150 photons.
944
Two-colour spectral separation was performed based on detection channel ratio (DCR)
values accounting for background inhomogeneity. DCR values from the HeadStart iterations
were weighted averaged by the corresponding effective photon counts at the MINFLUX
pattern offset. A two-component Gaussian Mixture Model was then fitted to the resulting
values and the assigned probability of belonging to distribution 1 or 2 was retrieved. A
localisation was then assigned to one colour if its probability was greater than 95% or
otherwise set to a 'not defined' category. All localisations associated with the same molecule
were subsequently assigned to the colour category that occurred most frequently within that
molecule's trace. Molecules for which most localisations were not defined were not
accounted for in the two-colour data. After two-colour separation, localisations were filtered
by efo (<80kHz) and number of localisations per trace (>3). The centroid of each remaining
trace was calculated and DBSCAN cluster analysis used to identify centroid clusters
(minimum points = 10, epsilon = 150 nm). Nearest neighbour analysis was performed
between FLUX 680 and Alexa Fluor 647 clusters using the K-D Tree neighbour search
algorithm.
960
961 *Transmission electron microscopy (TEM)*

962
Cells and islets were prepared for TEM and imaged as previously described[18]. For
SNAP/FLAG-hGLP-1R gold labelling, INS-1 832/3 SNAP/FLAG-hGLP-1R cells grown on
Thermanox coverslips (Agar Scientific), or mouse islets transduced with SNAP/FLAG-hGLP-
1R adenoviruses, were labelled with 2 µM cleavable SNAP-Surface biotin probe (kind gift of
Dr Ivan Corrêa Jr, New England Biolabs) in full media, followed by incubation with 5 µg/mL
NaN_3-free Alexa Fluor 488 Streptavidin, 5-nm or 10-nm colloidal gold conjugate (Thermo
Fisher Scientific) to label SNAP/FLAG-hGLP-1R prior to stimulation with 100 nM exendin-4,
fixation and processing for conventional TEM. Cells or islets were fixed at room temperature
for 20 minutes in 2% PFA + 2% glutaraldehyde in 0.1 M sodium cacodylate, washed 2x with
0.1 M sodium cacodylate and 1x with ddH_2O, and then post-stained on ice with 2% osmium
tetroxide for 1 hour followed by 1% tannic acid at room temperature for 45 minutes, and
washed 2x in 0.05 M sodium cacodylate before stepwise dehydration in 70%, 90%, and
100% EtOH and Epon resin embedding. Cells mounted on Epon stubs and resin-embedded
islets in moulds were baked at 65°C overnight. Samples were trimmed and cut *en face* with
a diamond knife (DiATOME) in a Leica Ultracut UCT 6 ultramicrotome before examination on
an FEI Tecnai G2-Spirit TEM. Images were acquired in a charge-coupled device camera
(Eagle) and analysed in Fiji. Mitochondrial area, number of cristae per mitochondria and
ERMCS length was determined as in [96] by manually drawing mitochondria and ERMCSs
visible from the TEM micrographs.
982
983 *GLP-1R–VAPB interaction assay by NanoBRET*
984
INS-1 832/3 cells were co-transfected with 1 µg Venus-VAPB (generated *in house* from
EGFP-VAPB) and 0.5 µg GLP-1R-NanoLuc[97]. NanoBRET assays were carried out as
previously described[97]. Briefly, 24 hours after transfection, cells were washed with PBS,
detached, and resuspended in NanoGlo Live Cell Reagent (Promega) with furimazine (1:40
dilution in HBSS) and seeded into white 96-well half-area plates. A baseline reading was
taken for 5 minutes at 37°C at 460 nm for NanoLuc emission and at 535 nm for Venus
emission in a Flexstation 3 microplate reader. Cells were then stimulated with or without 1
µM of the indicated agonist and luminescent signal read for 30 minutes. Ratio (535/460 nm)
BRET readings were normalised to baseline and to average vehicle readings for agonist-
specific responses. Area under the curve (AUC) was calculated per agonist response using
GraphPad Prism 10.2.1.
996
997 *Global and mitochondrial PKA assays*
998
INS-1 832/3 cells were seeded onto 35-mm glass-bottom dishes (MatTek) and transfected
with enhanced excitation-ratiometric PKA biosensor pcDNA3.1(+)-ExRai-AKAR2 (for global
PKA) or pcDNA3-mitoExRai-AKAR2 (for mitochondrial PKA)[27]. Cells were washed with PBS
24 hours after transfection and imaged in Krebs-Ringer bicarbonate-Hepes (KRBH) buffer
(140 mM NaCl, 3.6 mM KCl, 1.5 mM $CaCl_2$, 0.5 mM $MgSO_4$, 0.5 mM NaH_2PO_4, 2 mM
$NaHCO_3$, 10 mM Hepes, 0.1% BSA, saturated with 95% O_2 / 5% CO_2, pH 7.4) supplemented
with 11 mM glucose and imaged ratiometrically at excitation wavelengths of 405 and 488 nm
and emission of 530 nm using a Zeiss LSM-780 inverted confocal laser-scanning
microscope with a 63x oil immersion objective, with images taken every 6 seconds, in a
37°C heated chamber. A baseline reading was taken for 1 minute followed by stimulation
with 1 µM exendin-4 and imaging for 5 minutes before addition of 10 µM forskolin (FSK) +
100 µM isobutyl methylxanthine (IBMX) for the last 2 minutes of acquisition to record
maximal responses. AUCs were calculated using GraphPad Prism 10.2.1.
1012
1013 *cAMP assays*
1014
1015 INS-1 832/3 cells were seeded onto 35-mm glass-bottom dishes (MatTek) and incubated
1016 overnight with the Green Up cADDis biosensor in a BacMam vector (Montana Molecular)

according to the manufacturer's instructions. Cells were washed with PBS and imaged in 11 mM glucose KRBH buffer using a Nikon Eclipse Ti microscope with an ORCA-Flash 4.0 camera (Hamamatsu) and Metamorph software (Molecular Devices) with a 20x air objective and 488 nm laser, with images taken every 6 seconds, on a 37°C stage. A baseline reading was taken for 1 minute followed by stimulation with 1 µM exendin-4 for 5 minutes, and 10 µM FSK + 100 µM IBMX addition for the last 2 minutes of the acquisition for maximal responses. Raw fluorescent intensities were extracted in Fiji to plot an average intensity trace and AUCs calculated in GraphPad Prism 10.2.1.

Mitochondrial morphology assays

INS-1 832/3 cells were seeded at 50,000 cells per well in 96-well glass-bottom imaging plates (Sigma Aldrich) coated with poly-L-lysine and laminin (1:50). Primary mouse islets were dispersed by trituration with trypsin-EDTA 0.25% for 3 minutes and seeded as above, or imaged whole without dispersion. Both cells and islets were stimulated with 100 nM exendin-4 or vehicle for 10 minutes and subsequently labelled with MitoTracker Red CMXRos (Thermo Fisher Scientific), washed 1x with PBS and imaged in phenol red free RPMI-1640 in a Leica Stellaris 8 inverted confocal microscope with 63x/1.40 (islets) or 100x/1.40 (cells) oil immersion objectives. Five images were collected per condition and analysed using the Mitochondria Analyzer plugin for Fiji to determine mitochondrial mean area and form factor. Cells were also classified by mitochondrial phenotype as in [49, 98] into a) hyperfused (highly interconnected and elongated), b) tubular (mixed) or c) fragmented mitochondrial types. For INS-1 832/3 and dispersed islet cell experiments, 30 to 130 cells were analysed from *n*=4-5 independent repeats per experiment; for intact islet experiments, all visible cells in a total of 15 to 30 islets from *n*=5-6 independent repeats were analysed per experiment.

Imaging of mitochondria by STED microscopy

INS-1 832-3 cells were seeded in glass-bottom dishes (MatTek), labelled with 500 nM PKmito Orange FX (Spirochrome) for 30 minutes, washed 3x in PBS and incubated in complete media for 30 minutes to recover. Cells were then treated with vehicle or 100 nM exendin-4 in complete media for 10 minutes prior to fixation in 4% PFA and imaging in PBS using a Leica Stellaris 8 microscope in TauSTED mode with a 100x/1.40 oil immersion objective. Cells were excited at 591 nm and pulsed depletion performed at 775 nm with 750 picosecond delay at 50% intensity.

Mitochondrial membrane potential assay

INS-1 832/3 cells were seeded 35-mm glass-bottom dishes (MatTek), incubated in 3 mM glucose KRBH for 1 hour, followed by incubation with 50 µM tetramethylrhodamine (TMRE) in 3 mM glucose KRBH for 15 minutes to label active mitochondria[99]. Cells were then washed with KRBH and imaged in 3 mM glucose KRBH with images taken every 12 seconds in a 60x/1.4 oil immersion objective with a 561 nm laser in a Nikon Eclipse Ti microscope with an ORCA-Flash 4.0 camera (Hamamatsu) and Metamorph software (Molecular Devices). A baseline reading was recorded for 1 minute followed by stimulation with 1 µM exendin-4 in 6 mM glucose KRBH for 2 minutes. Glucose levels were then increased to 20 mM and cells imaged for a further 5 minutes, followed by mitochondrial hyperpolarisation using 20 µg/ml oligomycin for 5 minutes and depolarisation using 20 µM carbonyl cyanide 4-trifluoromethoxyphenylhydrazone (FCCP) for 5 minutes. Raw fluorescent intensities were extracted in Fiji to obtain average intensity traces and AUCs calculated using GraphPad Prism 10.2.1.

ATP assays

By time-lapse microscopy: INS-1 832/3 cells were seeded in 35-mm glass-bottom dishes
(MatTek) and transfected with the ATP biosensor cyto-Ruby3-iATPSnFR1.0[100]. Cells were
imaged 24 hours after DNA transfection in 11 mM glucose KRBH buffer in a Zeiss LSM-780
inverted confocal laser-scanning microscope using a 63x/1.4 oil immersion objective with a
488 nm laser, with images taken every 6 seconds. A baseline reading was recorded for 1
minute followed by stimulation with 1 µM exendin-4 for 5 minutes. Raw fluorescent
intensities were extracted using Fiji to obtain average intensity traces and AUCs calculated
using GraphPad Prism 10.2.1.
1080
Biochemical assay: INS-1 832/3 cells seeded in 12-well plates, or mouse islets, were pre-
incubated with KRBH buffer supplemented with 3 mM glucose for 30 minutes followed by
stimulation with KRBH supplemented with 11 mM glucose +/- 100 nM exendin-4 for 15
minutes. Cells were lysed in KRBH + 0.1% Triton X-100 and sonicated, while islets were
lysed in acidic ethanol (75% ethanol, 15 mM HCl). Samples were centrifuged at 13,200 rpm
and stored at -20°C until further analysis. ATP concentrations from islets and cells were
determined using the ATP Determination Kit (A22066, Thermo Fisher Scientific).
1088
1089   *Insulin secretion assays*
1090
INS-1 832/3 cells were seeded in 6-well plates at 1.2×10^6 cells per well. For VAPB WT
versus P56S experiments, cells were co-transfected with 40 ng proinsulin-NLuc and 200 ng
of either pEGFP-VAPB, pEGFP-VAPB-P56S (generated *in house* by site directed
mutagenesis from pEGFP-VAPB), or pcDNA3.1+ as a control. For SPHKAP-EGFP
experiments, cells were co-transfected as above with proinsulin-NLuc and either empty
EGFP or SPHKAP-EGFP. For RNAi experiments, cells were transfected with Control *versus*
Target (VAPB or SPHKAP) siRNA. Cells were incubated overnight in INS-1 832/3 media at 3
mM glucose 24 hours after cDNA, and 48 hours after siRNA transfections and then pre-
incubated for 30 minutes in KRBH buffer supplemented with 3 mM glucose before
stimulation in KRBH buffer supplemented with 11 mM glucose +/- 100 nM exendin-4 for 60
minutes. Cell supernatants were collected to assess secreted insulin and cells lysed in
KRBH + 0.1% Triton X-100 with sonication for total insulin extraction. For cDNA-transfected
cells, NanoGlo Live Cell Reagent with furimazine (1:40 dilution in HBSS) was added and
relative luciferase units (RLU) recorded at 460 nm using a Flexstation 3 microplate reader.
1105
Mouse islets were incubated in KRBH buffer supplemented with 3 mM glucose for 1 hour
followed by stimulation in KRBH buffer supplemented with 11 mM glucose +/- 100 nM
exendin-4 for 30 minutes. Supernatants were collected and centrifuged at 1,000 rpm for 3
minutes. Islets were lysed in acidic ethanol (75% ethanol, 15 mM HCl), sonicated 3x10
seconds in a water bath and centrifuged at 13,200 rpm for 10 minutes to determine
intracellular insulin concentrations.
1112
1113   For RNAi-transfected cells and shRNA-transduced islets, sample insulin concentration was
1114   determined using the Insulin Ultra-Sensitive HTRF Assay kit (Cisbio) according to
1115   manufacturer's instructions.
1116
1117   *TUNEL assays*
1118
INS-1 832/3 SNAP/FLAG-hGLP-1R cells seeded onto coverslips were incubated with or
without 5 µM thapsigargin in vehicle or 100 nM exendin-4-supplemented media for 8 hours
before fixation and terminal deoxynucleotidyl transferase dUTP nick-end labelling (TUNEL)
assay performed using the DeadEnd™ Fluorometric TUNEL system (Promega) following the
manufacturer's instructions. Coverslips were mounted with ProLong™ Diamond Antifade
Mountant with DAPI (Thermo Fisher Scientific) and imaged for fluorescein and DAPI (to
assess total cell number) in a Zeiss LSM-780 inverted confocal laser-scanning microscope

with a 20x/0/8 air objective. The percentage of TUNEL positive cells was calculated in Fiji
from 5 regions per coverslip.
1128
1129 *Proximity ligation assays (PLAs)*
1130
Mouse and human islets were stimulated with vehicle or 100 nM exendin-4 in full media for
10 minutes at 37°C. Islets were washed with PBS and fixed for a minimum of 1 hour. Islets
were washed again with PBS and processed by PLA with the NaveniFlex Tissue MR Red kit
(Navinci) following the manufacturer's instructions, with an extra permeabilisation step.
Antibody pairs for mouse islets were mouse anti-mouse GLP-1R Ab Mab 7F38 (DSHB) and
either rabbit anti-VAPB (A14703, Antibodies.com), or rabbit anti-mouse SPHKAP (36690,
kind gift from Dr Nicholas Vierra, University of Utah[32]); antibody pairs for human islets were
mouse anti-human GLP-1R Mab 3F52 (DSHB) and either rabbit anti VAPB (A14703,
Antibodies.com), or rabbit anti-human SPHKAP (HPA042499, Atlas Antibodies). For insulin
co-staining, islets were further incubated with anti-insulin antibody (IR002, Dako Agilent
Technologies) overnight followed by secondary anti-guinea pig Alexa Fluor 647 (A21450, Life
Technologies) for 1 hour. Islets were imaged using the confocal settings on a Leica Stellaris
STED FALCON microscope and a 40x/0.8 air objective.
1144
1145 *Room temperature Correlative Light and Electron Microscopy (CLEM)*
1146
The CLEM protocol was adapted from [101]. Briefly, INS-1 832/3 SNAP/FLAG-hGLP-1R cells
were seeded on gridded coverslips (MatTek) and transfected with EGFP-VAPB before
labelling with MitoTracker Red CMXRos and SNAP-Surface Alexa Fluor 647 and stimulation
with 100 nM exendin-4 for 5 minutes. Cells were initially fixed for confocal microscopy with
4% PFA + 0.1% glutaraldehyde, and confocal stacks acquired with a C2+ confocal
microscope (Nikon) and a 60x oil immersion objective. After fluorescence imaging, cells were
additionally fixed for EM with 4% PFA + 2.5 % glutaraldehyde in sodium phosphate buffer.
Samples were further processed according to a protocol modified from [102] using osmium
tetroxide (OsO_4) and thiocarbohydrazide (TCH). In brief, samples were incubated in a 2%
aqueous OsO_4 solution containing 1.5% potassium ferrocyanide and 2 mM $CaCl_2$
(30 minutes on ice), washed in 1% TCH, rinsed in ddH_2O and incubated again in 2% OsO_4.
After washing, samples were stained *en bloc* with 1% uranyl acetate (UA), rinsed in ddH_2O
and dehydrated in a graded series of ethanol. Samples were embedded into epoxy resin
Embed812 (Science Services). After resin curing at 60°C, the coverslip was removed by
dipping into liquid nitrogen. The block was trimmed to a region identified during confocal
imaging and serial sections with a thickness of 300 nm cut. Section post-staining was
performed with 2% UA in methanol and 1% lead citrate in ddH_2O. After application of 15-nm
colloidal gold beads, sections were imaged in a 300 kV Thermo Fisher F30 electron
microscope. Tilt series were acquired at steps of 1° ranging from -63° to +63°. Tomograms
were reconstructed with the IMOD software package[103]. Correlation of confocal images and
tomogram slices was done with the EC-CLEM plugin[104] in Icy[105]. Segmentation of the
tomograms was performed manually with Microscopy Image Browser[106]. 3D rendering of the
segmentation was performed with ORS Dragonfly
(www.theobjects.com/dragonfly/index.html).
1171
1172 *Cryo-CLEM*
1173
INS-1 832/3 SNAP/FLAG-hGLP-1R cells previously transfected with EGFP-VAPB were
plated onto glow discharged (30 seconds, 20 mAmp) poly-L-lysine-coated Quantifoil R2/2
200 mesh gold EM grids (Electron Microscopy Sciences). Cells were labelled with 500 nM
PKmito Orange FX (Spirochrome) and SNAP-Surface Alexa Fluor 647 before stimulation
with 100 nM exendin-4 in complete media for 10 minutes. Grids were subsequently plunge-
frozen in condensed ethane using a Vitrobot and stored in liquid N_2 for further analysis.
1180

The clipped plunge-frozen grids were loaded into the Zeiss correlative cryo-holder which can be transferred into the cryo-LSM and the cryo-FIB-SEM without further removal from the holder. For cryo-confocal data acquisition, the cryo-holder was transferred onto the cryo-Linkam stage (Linkam Scientific) and mounted onto a ZEISS Axio Imager.Z2 fitted with a LSM900 confocal scan head equipped with 3 confocal channels, including the Airyscan 2 detector for super resolution. Low magnification objectives (C Epiplan-Apochromat 5x/0.2 DIC, C Epiplan-Apochromat 10x/0.4 DIC) were used for overviews, and a LD EC Epiplan-Neofluar 100x/0.75 DIC used for high resolution. Images were acquired using the following settings: 488 nm, 561 nm and 640 nm excitation with emission bands of 493-551 nm, 553-612 nm and 613-700 nm. Reflection and brightfield channels were additionally acquired for correlation of light microscopy with SEM images. Confocal images were processed with LSM Plus (linear Wiener deconvolution). For highest resolution, Airyscan detection was used with the following settings: 488 nm, 561 nm and 640 nm excitation with emission bands of 492-549 nm, 564-611 nm and 644-700 nm and images processed using the Airyscan joint deconvolution algorithm. For cryo FIB-SEM data acquisition, the correlative cryo-holder was transferred into the Zeiss Crossbeam 550 equipped with a Quorum cryo stage. The transfer is done through the Quorum Prep Chamber attached to the FIB-SEM chamber. Inside the chamber, the sample surface was sputter-coated with platinum with a current of 5 mA for 45 seconds to make the surface conductive for SEM imaging and FIB milling. After insertion, the FIB-SEM coordinate system was aligned with respect to the LSM dataset using the ZEN software (Zeiss) and ROIs acquired in the LSM relocated and imaged in the FIB-SEM. The sample was then tilted to 54° (perpendicular to the FIB) and a cold platinum deposition applied to minimize the curtaining effect (vertical stripes in the SEM images that result from uneven milling of the sample due to roughness of the sample surface). For this purpose, the gas injection system (GIS) nozzle was inserted at ~2.5 mm from the sample surface and the unheated GIS opened for about 60 seconds resulting in a protection layer with a thickness of ~0.5 μm. Trenches were milled at the ROIs with a FIB current of 3 nA (Gallium ion energy 30 keV) to generate cross-sections. FIB-SEM volume imaging was performed with a milling current of 300 pA (Gallium ion energy 30 keV) using a slice thickness of 30 nm to obtain SEM image stacks representing a volume of 16 μm x 12 μm x 9 μm (XYZ). The SEM images were acquired with an electron energy of 2.3 keV and an electron current of 30 pA with a pixel size of 8 nm using the InLens SE detector. The resulting FIB-SEM datasets were imported into ZEN project. After alignment of the image stack, images were processed using "destriping" (ZEN) based on the VSNR algorithm[107]. Processed image stacks were cropped to an appropriate size and a 3D alignment of the SEM and LSM stacks performed. A selected area in the aligned region of the FIB-SEM data was manually segmented, slice by slice, in Fiji, using the freehand selection tool. Outlines were saved as masks, and 3D reconstruction performed in Icy[105].

Reactive oxygen species (ROS) assays

ROS was determined using the ROS-Glo™ H_2O_2 Assay (G8820, Promega) according to the manufacturer's instructions. Briefly, islets were incubated overnight in full media containing 1 μM thapsigargin +/- 100 nM exendin-4 at 37°C and subsequently incubated in 25 μM H_2O_2 Substrate diluted in H_2O_2 Substrate Dilution Buffer plus 1 μM thapsigargin +/- 100 nM exendin-4 in a final volume of 100 μL for 3 hours at 37°C in a 5% CO_2 incubator. Media (50 μL) was combined with 50 μL ROS-Glo™ Detection Solution in a 96-well opaque plate and incubated for 20 minutes at room temperature. Relative luminescence units (RLU) were recorded in a Flexstation 3 plate reader.

Mitophagy (keima) assays

INS-1 832/3 cells were transfected with the dual-excitation ratiometric biosensor mKeima-Red-Mito-7 and seeded onto 35-mm glass-bottom dishes (MatTek). Cells were imaged 24 hours after seeding in RPMI-1640 buffer without phenol red +/- 100 nM exendin-4 by time-

1236 lapse microscopy every 12 seconds for 11 minutes in a 37°C heated chamber using a Leica
1237 Stellaris 8 inverted confocal microscope with a 40x/1.2 oil immersion objective with excitation
1238 wavelengths of 440 and 580 nm and emission wavelength of 620 nm. Ratio fluorescence
1239 intensities for the two excitation wavelengths (580/440 nm) was calculated to obtain a
1240 response trace as in [108].
1241
1242 *ER and mitochondrial calcium assays*
1243
INS-1 832/3 cells were transfected with pCAG G-CEPIA1er (for ER calcium) or pCAG mito-
RCaMP1h (for mitochondrial calcium)[109]. 24 hours after transfection, cells were washed 1x
with PBS and imaged by time-lapse microscopy in RPMI-1640 without phenol red in a
40x/1.2 oil immersion objective Nikon Eclipse Ti microscope with an ORCA-Flash 4.0
camera (Hamamatsu) and Metamorph software (Molecular Devices). Mitochondrial calcium
was imaged with 561 nm, and ER calcium with 488 nm excitation wavelength, with images
taken every 3 seconds. A baseline reading was recorded for 1 minute followed by 1 µM
exendin-4 stimulation for 3 minutes. Raw fluorescent intensities were extracted using Fiji to
obtain average intensity traces and AUCs calculated using GraphPad Prism 10.2.1.
1253
1254 *qPCR assays*
1255
<u>SPHKAP knock-down efficiency:</u> Mouse islets transduced with either Control or SPHKAP
shRNA lentiviral particles were lysed using Trizol and RNA extracted using 200 µL
chloroform per 1 mL Trizol and then precipitated in 500 µL isopropanol and 5 mg glycogen
per mL of Trizol. RNA was washed with 75% EtOH before resuspension in ddH$_2$O. 250 ng of
purified RNA was reverse transcribed into cDNA using the high-capacity cDNA reverse
transcription kit (Applied Biosystems) and relative mRNA expression determined using FAST
SYBR green chemistry qPCR and Applied Biosystems 7500 Real time PCR system.
1263
<u>Mitochondrial DNA copy number:</u> INS1 832/3 cells were seeded at 300,000 cells per well on
a 24-well plate. Cells were stimulated +/- 100 nM exendin-4 for 30 minutes at 37°C. Total
DNA was extracted using 20 mM Tris pH 8.0, 5 mM EDTA, 400 mM NaCl, 1% SDS and 40
µg/mL proteinase K, as previously described[110], and used at 3 ng/mL for qPCR analysis of
mitochondrial DNA (mtDNA) relative to genomic DNA (gDNA) as in [111]. Primers were
designed using NCBI Primer Blast and validated *in silico* and then using a 1:2 standard
curve of DNA ranging from 12 - 0.75 ng/mL, resulting in reaction efficiencies of 97 to 102%.
1271
1272 qPCR primer sequences are shown in Table 4.
1273

Gene	Species	Fwd (5'→3')	Rev (5'→3')
HK1	rat	CCAAGCGTCTCCATAAGGCA	GTTGGTCAGCCAGACGGTAA
mitochondrial tRNA[Trp]	rat	GTGAAATCAACAACCCGCCC	ATAGTCACCCCCAGGACGAA
SPHKAP	mouse	CCAAACTCTGCTACCTGCCTT	GTGATTCCACGTTGCTGGATT
Actb	mouse	CACTGTCGAGTCGCGTCC	TCATCCATGGCGAACTGGTG

1274
1275 **Table 4:** qPCR primer sequences.
1276
1277 *Mitochondria-associated membranes (MAMs) purification*
1278
1279 2x10[7] INS-1 832/3 SNAP/FLAG-hGLP-1R cells were seeded in a T75 flask in complete
1280 media and cultured overnight. Cells were then stimulated in complete media containing 100
1281 nM exendin-4 for 10 minutes. The crude mitochondrial fraction containing ER-associated
1282 membranes was isolated using the mitochondria isolation kit for cultured cells (89874,

Thermo Fisher Scientific) according to the manufacturer's protocol. MAMs were subsequently separated from mitochondria by ultracentrifugation at 95,000 × g for 30 minutes in an OptiPrep™ Density Gradient Medium (Sigma Aldrich). The MAM fraction was diluted 1:10 in mitochondrial resuspension buffer (250 mM mannitol, 5 mM HEPES, 0.5 M EGTA, pH7.4), pelleted by ultracentrifugation at 100,000 × g for 1 hour and prepared for Western blot analysis as described above.

pcDNA3-GFP11(x7)-VAPB cloning

pcDNA3-GFP11(x7)-VAPB was cloned *in house* from pEGFP-C1-hVAPB and pcDNA3-GFP11(x7)-Actin by PCR amplification of human VAPB with addition of 5'-BglII and 3'-XhoI restriction sites. The second BglII site at position 7107 in pcDNA3-GFP11(x7)-Actin was removed by site directed mutagenesis using the Quick-Change Lightning kit (Agilent) prior to in frame cloning of human VAPB using T4 DNA ligase (Promega).

Fluorescent Sensors Targeted to Endogenous Proteins (FluoSTEPs) assays of VAPB-localised cAMP and PKA signalling

The FluoSTEP system was used to determine VAPB-localised cAMP and PKA signalling as in [25]. INS1 832/3 cells were transfected with pcDNA3-GFP11(x7)-VAPB and pcDNA3.1(+)-FluoSTEP-ICUE (for cAMP) or pcDNA3.1(+)-FluoSTEP-AKAR (for PKA) at a 1:1 ratio. Cells were seeded onto glass-bottom dishes (MatTek) 24 hours before imaging in RPMI-1640 without phenol red + 10% FBS. Cells were imaged by time-lapse microscopy using a Zeiss LSM-780 inverted confocal laser-scanning microscope with a 63x/1.4 oil immersion objective and a 488 nm excitation wavelength and emissions at 508 and 583 nm for 2 minutes to record a baseline, followed by agonist stimulation at 100 nM for 5 minutes and 2 minutes exposure to 10 µM FSK + 100 µM IBMX to generate a maximal response. FRET emission ratios were calculated and normalised to mean baseline values using GraphPad Prism 10.2.1.

Quantification of ERMCSs per mitochondrial area using splitFAST

INS-1 832/3 cells were plated onto Matrigel (Gibco) coated glass-bottom imaging chambers (Ibidi) and transfected with 4xMTS-Halo, ER-RspA-NFAST and OMM-short-RspA-CFAST[50]. After 18 hours, cells were labelled with 10 nM JFX646 Halo ligand (kind gift from the Lavis lab, Janelia) in optiMEM (Gibco) for 3 minutes, followed by 3x washes in mKRB buffer (140 mM NaCl, 2.8 mM KCl, 2 mM $MgCl_2$, 1 mM $CaCl_2$, 10 mM HEPES, 10 mM glucose, pH 7.4) at 37°C. The Lime splitFAST ligand (Twinkle Factory) was prepared at 5 µM in mKRB buffer and passed through a 0.22-µm filter before being added to cells. Exendin-4 was used at 1 µM in Lime splitFAST ligand in mKRB and added to cells for 10 minutes prior to imaging with an LSM980/Airyscan2 point-scanning confocal microscope (Zeiss) in LSMPlus mode. Simultaneous excitation was achieved using solid-state diode lasers at 488 nm/10 mW (0.9% max power) and 639 nm/7.5 mW (1.0% max power), with imaging conditions experimentally selected to minimise crosstalk. The resulting fluorescence was collected using a 63x/1.4 oil immersion objective and detected on a 34-channel spectral GaAsP array detector with the gain set at 650 V and split into two ranges of 490–632 nm and 641–694 nm. Samples were imaged at room temperature in a humidified imaging chamber with a Definite Focus module (Zeiss) employed for thermal drift correction and ZEN Blue v3.2 (Zeiss) software used for acquisition and deconvolution (LSMPlus function). The number of contact sites per mitochondrial area was quantified in Fiji. Firstly, the acquired images were gaussian blurred (sigma radius 1.5) to aid in the generation of reliable downstream masks and puncta detection. An automated Otsu threshold was used to generate mitochondrial masks and quantitate mitochondrial area. These same masks were then applied to the splitFAST channel to ensure that only true contact sites (i.e. puncta on mitochondria) were

counted in the analysis. A blinded count of total number of ERMCSs in each cell was taken and normalised to the mitochondrial area.

Statistical analyses for cellular experiments

All statistical analyses and graph generation was performed using GraphPad Prism 10.2.1 (GraphPad Software Inc). Statistical tests used are indicated in the corresponding figure legends. The number of replicates per comparison represents biological replicates and technical replicates within biological replicates are averaged before statistical testing. Data are presented as mean ± SEM. The p value threshold for statistical significance was set at 0.05.

Statistical analyses in human datasets

To compare the effects of *SPHKAP* gene region variants on T2D *versus* BMI, we used GWAS meta-analyses summary statistics from DIAMANTE consortium (http://www.diagram-consortium.org) for T2D by Mahajan *et al*[36], and from GIANT (https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files) for BMI by Locke *et al*[39]. We report SNPs from the region surrounding *SPHKAP* gene (Chr2:228,844,666-229,046,361, NCBI human genome built37). Data pre-processing involved merging and aligning variant effects to T2D risk-increasing allele. We extracted beta values for BMI and T2D signals and calculated respective odds ratios (OR) for T2D using the formula OR = exp(beta). The plot contains SNPs with at least nominally significant (p<0.05) effects on both BMI and T2D. A scatter plot was generated using RStudio with the ggplot2 package.

Data availability

Proteomics data is deposited in the ProteomeXchange (PRIDE) repository with accession number PXD056782.

Acknowledgements

The authors thank Dr Nicholas Vierra, University of Utah, for the SPHKAP antibodies; Prof Toshihiko Oka, Rikkyo University, for the MIC19-3xHA construct; Prof John D. Scott, Howard Hughes Medical Institute, for the SPHKAP-EGFP construct; Dr Emily Eden, UCL, for the EGFP-VAPB construct; the Filadi lab, CNR Institute of Neuroscience, for the ER-RspA-NFAST and OMM-short-RspA-CFAST splitFAST constructs; and the Avezov lab, UK Dementia Research Institute, for the 4xMTS-Halo construct. We thank the Imperial College Facility for Imaging by Light Microscopy (FILM) for technical support on light microscopy experiments, the Electron Microscopy Centre at the Centre of Structural Biology, Imperial College London, for support during TEM data acquisition, and Dr Ricardo Aramayo at the Electron Microscopy Facility of the MRC Laboratory of Medical Sciences (LMS) for support during cryo-CLEM sample preparation. The room temperature CLEM was supported by the Electron Microscopy and Histology Facility, a Core Facility of the CMCB Technology Platform at TU Dresden. We thank the EM facility of the Max Planck Institute of Molecular Cell Biology and Genetics for their services. We additionally thank the Cambridge Institute for Medical Research (CIMR) Microscopy Facility for their support with acquiring the splitFAST imaging data. The CIMR microscopy LSM980/Airyscan2 was funded by the MRC (MR/Y002172/1). The FILM Facility at Imperial College London is part-supported by funding from the Wellcome Trust (grant 104931/Z/14/Z) and BBSRC (grant BB/L015129/1). The NIDDK-funded Integrated Islet Distribution Program (IIDP) (RRID:SCR_014387) at City of Hope is supported by NIH Grant # 2UC4DK098085. The AT lab is funded by grants from Diabetes UK (19/0006094), the MRC (MR/X021467/1), and the Wellcome Trust (301619/Z/23/Z). GA is supported by a President's PhD Scholarship from Imperial College

London and AIO by a Commonwealth PhD Scholarship. MS and AM are supported by the German Center for Diabetes Research (DZD e.V.) and the German Ministry for Education and Research (BMBF). The JNA group is funded by a Wellcome Trust Career Development Award (227745/Z/23/Z).

References

1. Rivera Nieves AM, Wauford BM, Fu A. Mitochondrial bioenergetics, metabolism, and beyond in pancreatic beta-cells and diabetes. *Front Mol Biosci* **11**, 1354199 (2024).

2. Haythorne E*, et al.* Diabetes causes marked inhibition of mitochondrial metabolism in pancreatic beta-cells. *Nat Commun* **10**, 2474 (2019).

3. Ahmad E, Lim S, Lamptey R, Webb DR, Davies MJ. Type 2 diabetes. *Lancet* **400**, 1803-1820 (2022).

4. Mayendraraj A, Rosenkilde MM, Gasbjerg LS. GLP-1 and GIP receptor signaling in beta cells - A review of receptor interactions and co-stimulation. *Peptides* **151**, 170749 (2022).

5. Boland ML*, et al.* Resolution of NASH and hepatic fibrosis by the GLP-1R/GcgR dual-agonist Cotadutide via modulating mitochondrial function and lipogenesis. *Nat Metab* **2**, 413-431 (2020).

6. Timper K*, et al.* GLP-1 Receptor Signaling in Astrocytes Regulates Fatty Acid Oxidation, Mitochondrial Integrity, and Function. *Cell Metab* **31**, 1189-1205 e1113 (2020).

7. Wu P*, et al.* Liraglutide Regulates Mitochondrial Quality Control System Through PGC-1alpha in a Mouse Model of Parkinson's Disease. *Neurotox Res* **40**, 286-297 (2022).

8. Carlessi R*, et al.* GLP-1 receptor signalling promotes beta-cell glucose metabolism via mTOR-dependent HIF-1alpha activation. *Sci Rep* **7**, 2661 (2017).

9. Ng MYW, Wai T, Simonsen A. Quality control of the mitochondrion. *Dev Cell* **56**, 881-905 (2021).

10. Dong J*, et al.* Mic19 depletion impairs endoplasmic reticulum-mitochondrial contacts and mitochondrial lipid metabolism and triggers liver disease. *Nat Commun* **15**, 168 (2024).

11. Prinz WA, Toulmay A, Balla T. The functional universe of membrane contact sites. *Nat Rev Mol Cell Biol* **21**, 7-24 (2020).

12. Dudas EF, Huynen MA, Lesk AM, Pastore A. Invisible leashes: The tethering VAPs from infectious diseases to neurodegeneration. *J Biol Chem* **296**, 100421 (2021).

13. Kors S, Costello JL, Schrader M. VAP Proteins - From Organelle Tethers to Pathogenic Host Interactors and Their Role in Neuronal Disease. *Front Cell Dev Biol* **10**, 895856 (2022).

14. Schmidt R*, et al.* MINFLUX nanometer-scale 3D imaging and microsecond-range tracking on a common fluorescence microscope. *Nat Commun* **12**, 1478 (2021).

15. Di Mattia T, Tomasetto C, Alpy F. Faraway, so close! Functions of Endoplasmic reticulum-Endosome contacts. *Biochim Biophys Acta Mol Cell Biol Lipids* **1865**, 158490 (2020).

16. Suzuki H*, et al.* ALS-linked P56S-VAPB, an aggregated loss-of-function mutant of VAPB, predisposes motor neurons to ER stress-related death by inducing aggregation of co-expressed wild-type VAPB. *J Neurochem* **108**, 973-985 (2009).

17. Obara CJ*, et al.* Motion of VAPB molecules reveals ER-mitochondria contact site subdomains. *Nature* **626**, 169-176 (2024).

18. Jones B*, et al.* Targeting GLP-1 receptor trafficking to improve agonist efficacy. *Nat Commun* **9**, 1602 (2018).

19. Fang Z*, et al.* Ligand-Specific Factors Influencing GLP-1 Receptor Post-Endocytic Trafficking and Degradation in Pancreatic Beta Cells. *Int J Mol Sci* **21**, (2020).

20. Madsbad S, Holst JJ. Glycaemic control and weight loss with semaglutide in type 2 diabetes. *Lancet Diabetes Endocrinol* **5**, 315-317 (2017).

21. Nauck MA, D'Alessio DA. Tirzepatide, a dual GIP/GLP-1 receptor co-agonist for the treatment of type 2 diabetes with unmatched effectiveness regrading glycaemic control and body weight reduction. *Cardiovasc Diabetol* **21**, 169 (2022).

22. Dutta D, Nagendra L, Anne B, Kumar M, Sharma M, Kamrul-Hasan ABM. Orforglipron, a novel non-peptide oral daily glucagon-like peptide-1 receptor agonist as an anti-obesity medicine: A systematic review and meta-analysis. *Obes Sci Pract* **10**, e743 (2024).

23. Fatima H*, et al.* Evaluating Glycemic Control Efficacy and Safety of the Oral Small Molecule Glucagon-Like Peptide 1 Receptor Agonist Danuglipron in Type 2 Diabetes Patients: A Systemic Review and Meta-Analysis. *Diabetes Metab Syndr Obes* **16**, 3567-3578 (2023).

24. Novikoff A*, et al.* Spatiotemporal GLP-1 and GIP receptor signaling and trafficking/recycling dynamics induced by selected receptor mono- and dual-agonists. *Mol Metab* **49**, 101181 (2021).

25. Zhang JZ*, et al.* Phase Separation of a PKA Regulatory Subunit Controls cAMP Compartmentation and Oncogenic Signaling. *Cell* **182**, 1531-1544 e1515 (2020).

26. Akabane S, Oka T. Insights into the regulation of mitochondrial functions by protein kinase A-mediated phosphorylation. *J Biochem* **175**, 1-7 (2023).

27. Zhang JF*, et al.* An ultrasensitive biosensor for high-resolution kinase activity imaging in awake mice. *Nat Chem Biol* **17**, 39-46 (2021).

28. Vallese F*, et al.* An expanded palette of improved SPLICS reporters detects multiple organelle contacts in vitro and in vivo. *Nat Commun* **11**, 6069 (2020).

29. Kovanich D, van der Heyden MA, Aye TT, van Veen TA, Heck AJ, Scholten A. Sphingosine kinase interacting protein is an A-kinase anchoring protein specific for type I cAMP-dependent protein kinase. *Chembiochem* **11**, 963-971 (2010).

1501 30. Neefjes J, Cabukusta B. What the VAP: The Expanded VAP Family of Proteins
1502 Interacting With FFAT and FFAT-Related Motifs for Interorganellar Contact. *Contact*
1503 *(Thousand Oaks)* **4**, 25152564211012246 (2021).
1504
1505 31. Di Mattia T*, et al.* FFAT motif phosphorylation controls formation and lipid transfer
1506 function of inter-organelle contacts. *EMBO J* **39**, e104369 (2020).
1507
1508 32. Vierra NC*, et al.* Neuronal ER-plasma membrane junctions couple excitation to
1509 Ca(2+)-activated PKA signaling. *Nat Commun* **14**, 5231 (2023).
1510
1511 33. Lee YK*, et al.* Bipolar and schizophrenia risk gene AKAP11 encodes an autophagy
1512 receptor coupling the regulation of PKA kinase network homeostasis to synaptic
1513 transmission. *Res Sq*, (2025).
1514
1515 34. Wang Y, Harashima SI, Liu Y, Usui R, Inagaki N. Sphingosine kinase 1-interacting
1516 protein is a novel regulator of glucose-stimulated insulin secretion. *Sci Rep* **7**, 779
1517 (2017).
1518
1519 35. Liu Y*, et al.* Sphingosine kinase 1-interacting protein is a dual regulator of insulin and
1520 incretin secretion. *FASEB J* **33**, 6239-6253 (2019).
1521
1522 36. Mahajan A*, et al.* Multi-ancestry genetic study of type 2 diabetes highlights the power
1523 of diverse populations for discovery and translation. *Nat Genet* **54**, 560-572 (2022).
1524
1525 37. Suzuki K*, et al.* Genetic drivers of heterogeneity in type 2 diabetes pathophysiology.
1526 *Nature* **627**, 347-357 (2024).
1527
1528 38. Pulit SL*, et al.* Meta-analysis of genome-wide association studies for body fat
1529 distribution in 694 649 individuals of European ancestry. *Hum Mol Genet* **28**, 166-174
1530 (2019).
1531
1532 39. Locke AE*, et al.* Genetic studies of body mass index yield new insights for obesity
1533 biology. *Nature* **518**, 197-206 (2015).
1534
1535 40. Means CK*, et al.* An entirely specific type I A-kinase anchoring protein that can
1536 sequester two molecules of protein kinase A at mitochondria. *Proc Natl Acad Sci U S*
1537 *A* **108**, E1227-1235 (2011).
1538
1539 41. Kaiser SE, Brickner JH, Reilein AR, Fenn TD, Walter P, Brunger AT. Structural basis
1540 of FFAT motif-mediated ER targeting. *Structure* **13**, 1035-1045 (2005).
1541
1542 42. Kors S, Schrader M, Costello JL. Multiple Ways to Keep FFAT Under Control!
1543 *Contact (Thousand Oaks)* **5**, 1-4 (2022).
1544
1545 43. Friedman JR, Mourier A, Yamada J, McCaffery JM, Nunnari J. MICOS coordinates
1546 with respiratory complexes and lipids to establish mitochondrial inner membrane
1547 architecture. *Elife* **4**, (2015).
1548
1549 44. Pearson RB, Kemp BE. Protein kinase phosphorylation site sequences and
1550 consensus specificity motifs: tabulations. *Methods Enzymol* **200**, 62-81 (1991).
1551
1552 45. Bock-Bierbaum T*, et al.* Structural insights into crista junction formation by the Mic60-
1553 Mic19 complex. *Sci Adv* **8**, eabo4946 (2022).
1554

1555   46.   Yu R*, et al.* The phosphorylation status of Ser-637 in dynamin-related protein 1
1556   (Drp1) does not determine Drp1 recruitment to mitochondria. *J Biol Chem* **294**,
1557   17262-17277 (2019).
1558

1559   47.   Kashatus JA*, et al.* Erk2 phosphorylation of Drp1 promotes mitochondrial fission and
1560   MAPK-driven tumor growth. *Mol Cell* **57**, 537-551 (2015).
1561

1562   48.   Breitzig MT, Alleyn MD, Lockey RF, Kolliputi N. A mitochondrial delicacy: dynamin-
1563   related protein 1 and mitochondrial dynamics. *Am J Physiol Cell Physiol* **315**, C80-
1564   C90 (2018).
1565

1566   49.   von der Malsburg A*, et al.* Structural mechanism of mitochondrial membrane
1567   remodelling by human OPA1. *Nature* **620**, 1101-1108 (2023).
1568

1569   50.   Garcia Casas P*, et al.* Simultaneous detection of membrane contact dynamics and
1570   associated Ca(2+) signals by reversible chemogenetic reporters. *Nat Commun* **15**,
1571   9775 (2024).
1572

1573   51.   Ast J*, et al.* Revealing the tissue-level complexity of endogenous glucagon-like
1574   peptide-1 receptor expression and signaling. *Nat Commun* **14**, 301 (2023).
1575

1576   52.   Ast J*, et al.* Expanded LUXendin Color Palette for GLP1R Detection and
1577   Visualization In Vitro and In Vivo. *JACS Au* **2**, 1007-1017 (2022).
1578

1579   53.   Ast J, Broichhagen J, Hodson DJ. Reagents and models for detecting endogenous
1580   GLP1R and GIPR. *EBioMedicine* **74**, 103739 (2021).
1581

1582   54.   Komatsu H*, et al.* Oxygen environment and islet size are the primary limiting factors
1583   of isolated pancreatic islet survival. *PLoS One* **12**, e0183780 (2017).
1584

1585   55.   Veluthakal R*, et al.* Mitochondrial Dysfunction, Oxidative Stress, and Inter-Organ
1586   Miscommunications in T2D Progression. *Int J Mol Sci* **25**, (2024).
1587

1588   56.   Rutter GA, Sidarala V, Kaufman BA, Soleimanpour SA. Mitochondrial metabolism
1589   and dynamics in pancreatic beta cell glucose sensing. *Biochem J* **480**, 773-789
1590   (2023).
1591

1592   57.   Koeck T*, et al.* A common variant in TFB1M is associated with reduced insulin
1593   secretion and increased future risk of type 2 diabetes. *Cell Metab* **13**, 80-91 (2011).
1594

1595   58.   Reardon W*, et al.* Diabetes mellitus associated with a pathogenic point mutation in
1596   mitochondrial DNA. *Lancet* **340**, 1376-1379 (1992).
1597

1598   59.   Mulder H. Transcribing beta-cell mitochondria in health and disease. *Mol Metab* **6**,
1599   1040-1051 (2017).
1600

1601   60.   Walker EM*, et al.* Retrograde mitochondrial signaling governs the identity and
1602   maturity of metabolic tissues. *Science*, eadf2034 (2025).
1603

1604   61.   Anton SE*, et al.* Receptor-associated independent cAMP nanodomains mediate
1605   spatiotemporal specificity of GPCR signaling. *Cell* **185**, 1130-1142 e1111 (2022).
1606

1607   62.   Xie B, Nguyen PM, Idevall-Hagren O. Feedback regulation of insulin secretion by
1608   extended synaptotagmin-1. *FASEB J* **33**, 4716-4728 (2019).
1609

63. Xie B, Panagiotou S, Cen J, Gilon P, Bergsten P, Idevall-Hagren O. The endoplasmic reticulum-plasma membrane tethering protein TMEM24 is a regulator of cellular Ca2+ homeostasis. *J Cell Sci* **135**, (2022).

64. Panagiotou S*, et al.* OSBP-mediated PI(4)P-cholesterol exchange at endoplasmic reticulum-secretory granule contact sites controls insulin secretion. *Cell Rep* **43**, 113992 (2024).

65. Hardy JC*, et al.* Molecular determinants and signaling effects of PKA RIalpha phase separation. *Mol Cell* **84**, 1570-1584 e1577 (2024).

66. Xiao J*, et al.* Control of human pancreatic beta cell kinome by glucagon-like peptide-1 receptor biased agonism. *Diabetes Obes Metab*, (2023).

67. Chen Z*, et al.* GLP-1R-positive neurons in the lateral septum mediate the anorectic and weight-lowering effects of liraglutide in mice. *J Clin Invest* **134**, (2024).

68. Han J*, et al.* The Identification of Novel Protein-Protein Interactions in Liver that Affect Glucagon Receptor Activity. *PLoS One* **10**, e0129226 (2015).

69. Paradies G, Paradies V, Ruggiero FM, Petrosillo G. Role of Cardiolipin in Mitochondrial Function and Dynamics in Health and Disease: Molecular and Pharmacological Aspects. *Cells* **8**, (2019).

70. Zhang J, Shi Y. In Search of the Holy Grail: Toward a Unified Hypothesis on Mitochondrial Dysfunction in Age-Related Diseases. *Cells* **11**, (2022).

71. Thompson K*, et al.* Biallelic variants in TAMM41 are associated with low muscle cardiolipin levels, leading to neonatal mitochondrial disease. *HGG Adv* **3**, 100097 (2022).

72. Kaufman BA, Li C, Soleimanpour SA. Mitochondrial regulation of beta-cell function: maintaining the momentum for insulin release. *Mol Aspects Med* **42**, 91-104 (2015).

73. Prentki M, Peyot ML, Masiello P, Madiraju SRM. Nutrient-Induced Metabolic Stress, Adaptation, Detoxification, and Toxicity in the Pancreatic beta-Cell. *Diabetes* **69**, 279-290 (2020).

74. Ngo J*, et al.* Mitochondrial morphology controls fatty acid utilization by changing CPT1 sensitivity to malonyl-CoA. *EMBO J* **42**, e111901 (2023).

75. Prentki M, Corkey BE, Madiraju SRM. Lipid-associated metabolic signalling networks in pancreatic beta cell function. *Diabetologia* **63**, 10-20 (2020).

76. Nikonova L, Koza RA, Mendoza T, Chao PM, Curley JP, Kozak LP. Mesoderm-specific transcript is associated with fat mass expansion in response to a positive energy balance. *FASEB J* **22**, 3925-3937 (2008).

77. Wang T*, et al.* OSBPL2 Is Required for the Binding of COPB1 to ATGL and the Regulation of Lipid Droplet Lipolysis. *iScience* **23**, 101252 (2020).

78. Xue J*, et al.* TMEM120A is a coenzyme A-binding membrane protein with structural similarities to ELOVL fatty acid elongase. *Elife* **10**, (2021).

1664 79. Lin J*, et al.* A positive feedback loop between ZEB2 and ACSL4 regulates lipid
1665 metabolism to promote breast cancer metastasis. *Elife* **12**, (2023).
1666
1667 80. Gromovsky AD*, et al.* Delta-5 Fatty Acid Desaturase FADS1 Impacts Metabolic
1668 Disease by Balancing Proinflammatory and Proresolving Lipid Mediators. *Arterioscler*
1669 *Thromb Vasc Biol* **38**, 218-231 (2018).
1670
1671 81. Obaseki E, Adebayo D, Bandyopadhyay S, Hariri H. Lipid droplets and fatty acid-
1672 induced lipotoxicity: in a nutshell. *FEBS Lett* **598**, 1207-1214 (2024).
1673
1674 82. Xu Y*, et al.* CDP-DAG synthase 1 and 2 regulate lipid droplet growth through distinct
1675 mechanisms. *J Biol Chem* **294**, 16740-16755 (2019).
1676
1677 83. Yamashita A*, et al.* Glycerophosphate/Acylglycerophosphate acyltransferases.
1678 *Biology (Basel)* **3**, 801-830 (2014).
1679
1680 84. Miklas JW*, et al.* TFPa/HADHA is required for fatty acid beta-oxidation and cardiolipin
1681 re-modeling in human cardiomyocytes. *Nat Commun* **10**, 4671 (2019).
1682
1683 85. Williams AS*, et al.* Ketone flux through BDH1 supports metabolic remodeling of
1684 skeletal and cardiac muscles in response to intermittent time-restricted feeding. *Cell*
1685 *Metab* **36**, 422-437 e428 (2024).
1686
1687 86. Hugenroth M, Bohnert M. Come a little bit closer! Lipid droplet-ER contact sites are
1688 getting crowded. *Biochim Biophys Acta Mol Cell Res* **1867**, 118603 (2020).
1689
1690 87. Briant LJB*, et al.* CPT1a-Dependent Long-Chain Fatty Acid Oxidation Contributes to
1691 Maintaining Glucagon Secretion from Pancreatic Islets. *Cell Rep* **23**, 3300-3311
1692 (2018).
1693
1694 88. Shiiba I*, et al.* ER-mitochondria contacts mediate lipid radical transfer via
1695 RMDN3/PTPIP51 phosphorylation to reduce mitochondrial oxidative stress. *Nat*
1696 *Commun* **16**, 1508 (2025).
1697
1698 89. Liu X-M*, et al.* Abnormal Lipid Metabolism and Altered Neuronal Support by
1699 Astrocytes Lacking Akap11, a Risk Gene for Schizophrenia and Bipolar
1700 Disorder. *bioRxiv*, 2025.2004.2025.650548 (2025).
1701
1702 90. Raffaello A, Mammucari C, Gherardi G, Rizzuto R. Calcium at the Center of Cell
1703 Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes.
1704 *Trends Biochem Sci* **41**, 1035-1049 (2016).
1705
1706 91. Klec C, Ziomek G, Pichler M, Malli R, Graier WF. Calcium Signaling in ss-cell
1707 Physiology and Pathology: A Revisit. *Int J Mol Sci* **20**, (2019).
1708
1709 92. Naylor J*, et al.* Use of CRISPR/Cas9-engineered INS-1 pancreatic beta cells to
1710 define the pharmacology of dual GIPR/GLP-1R agonists. *Biochem J* **473**, 2881-2891
1711 (2016).
1712
1713 93. Yu V*, et al.* Differential CpG methylation at Nnat in the early establishment of beta
1714 cell heterogeneity. *Diabetologia*, (2024).
1715
1716 94. Shah AD, Goode RJA, Huang C, Powell DR, Schittenhelm RB. LFQ-Analyst: An
1717 Easy-To-Use Interactive Web Platform To Analyze and Visualize Label-Free
1718 Proteomics Data Preprocessed with MaxQuant. *J Proteome Res* **19**, 204-211 (2020).

95. Akabane S*, et al.* PKA Regulates PINK1 Stability and Parkin Recruitment to Damaged Mitochondria through Phosphorylation of MIC60. *Mol Cell* **62**, 371-384 (2016).

96. Lam J*, et al.* A Universal Approach to Analyzing Transmission Electron Microscopy with ImageJ. *Cells* **10**, (2021).

97. Manchanda Y*, et al.* Enhanced Endosomal Signaling and Desensitization of GLP-1R vs GIPR in Pancreatic Beta Cells. *Endocrinology* **164**, (2023).

98. Landoni JC, Kleele T, Winter J, Stepp W, Manley S. Mitochondrial Structure, Dynamics, and Physiology: Light Microscopy to Disentangle the Network. *Annu Rev Cell Dev Biol* **40**, 219-240 (2024).

99. Perry SW, Norman JP, Barbieri J, Brown EB, Gelbard HA. Mitochondrial membrane potential probes and the proton gradient: a practical usage guide. *Biotechniques* **50**, 98-115 (2011).

100. Lobas MA*, et al.* A genetically encoded single-wavelength sensor for imaging cytosolic and cell surface ATP. *Nat Commun* **10**, 711 (2019).

101. Polishchuk RS*, et al.* Correlative light-electron microscopy reveals the tubular-saccular ultrastructure of carriers operating between Golgi apparatus and plasma membrane. *J Cell Biol* **148**, 45-58 (2000).

102. Seligman AM, Wasserkrug HL, Hanker JS. A new staining method (OTO) for enhancing contrast of lipid--containing membranes and droplets in osmium tetroxide--fixed tissue with osmiophilic thiocarbohydrazide(TCH). *J Cell Biol* **30**, 424-432 (1966).

103. Kremer JR, Mastronarde DN, McIntosh JR. Computer visualization of three-dimensional image data using IMOD. *J Struct Biol* **116**, 71-76 (1996).

104. Paul-Gilloteaux P*, et al.* eC-CLEM: flexible multidimensional registration software for correlative microscopies. *Nat Methods* **14**, 102-103 (2017).

105. de Chaumont F*, et al.* Icy: an open bioimage informatics platform for extended reproducible research. *Nat Methods* **9**, 690-696 (2012).

106. Belevich I, Joensuu M, Kumar D, Vihinen H, Jokitalo E. Microscopy Image Browser: A Platform for Segmentation and Analysis of Multidimensional Datasets. *PLoS Biol* **14**, e1002340 (2016).

107. Fehrenbach J, Weiss P, Lorenzo C. Variational algorithms to remove stationary noise: applications to microscopy imaging. *IEEE Trans Image Process* **21**, 4420-4430 (2012).

108. Sun N, Malide D, Liu J, Rovira, II, Combs CA, Finkel T. A fluorescence-based imaging method to measure in vitro and in vivo mitophagy using mt-Keima. *Nat Protoc* **12**, 1576-1587 (2017).

109. Hirabayashi Y*, et al.* ER-mitochondria tethering by PDZD8 regulates Ca(2+) dynamics in mammalian neurons. *Science* **358**, 623-630 (2017).

1774 110. Sambrook J, Russell DW. Purification of nucleic acids by extraction with
1775   phenol:chloroform. *CSH Protoc* **2006**, (2006).
1776
1777 111. Rooney JP*, et al.* PCR based determination of mitochondrial DNA copy number in
1778   multiple species. *Methods Mol Biol* **1241**, 23-38 (2015).
1779
1780 112. Neelankal John A, Ram R, Jiang FX. RNA-Seq Analysis of Islets to Characterise the
1781   Dedifferentiation in Type 2 Diabetes Model Mice db/db. *Endocr Pathol* **29**, 207-221
1782   (2018).

1783
1784
1785 **Figure Legends**
1786

Fig. 1. Human GLP-1R β-cell interactome and validation of VAPB-hGLP-1R interaction.
a, Heatmap of GLP-1R interactor enrichment from anti-FLAG co-immunoprecipitates of
vehicle (Veh) *versus* 5-minute exendin-4 (Ex-4)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-
1R cells; blue, decreased; red, increased binding; LC-MS/MS data analysed in LFQ-Analyst
and normalised to hGLP-1R levels for each experimental sample; colour scale centred
around overall median value of the experiment; n=4. **b**, Volcano plot of data from (**a**)
depicting $\log_2$ fold change to vehicle (x-axis) *versus* $-\log10_q$value (y-axis). **c**, List of ER,
mitochondria or ERMCS-localised hGLP-1R interactors enriched in exendin-4-stimulated
versus vehicle conditions. **d**, Co-immunoprecipitation (co-IP) of EGFP-VAPB with
SNAP/FLAG-hGLP-1R in Veh *versus* Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R
cells. Representative blots and quantification of EGFP-VAPB levels per condition normalised
to SNAP shown; n=3, p value as indicated by paired t-test. **e**, As for (**d**) for endogenous
VAPB; n=4, p value as indicated by paired t-test. **f**, Confocal microscopy analysis of
SNAP/FLAG-hGLP-1R (red) co-localisation with EGFP-VAPB (green) in Ex-4-stimulated
INS-1 832/3 SNAP/FLAG-hGLP-1R cells; box, magnification inset; size bar, 5 µm. **g**,
Electron microscopy micrograph depicting gold-labelled SNAP/FLAG-hGLP-1R (arrows)
localised to an endosome–ER MCS in an Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-
1R cell; endosome highlighted in gold and ER in blue; size bar, 100 nm. **h**, Maximum
intensity projection (MIP) of a confocal z-stack of EGFP-VAPB (green) and SNAP/FLAG-
hGLP-1R (red) from an Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cell; size bar, 10
µm. **i**, Confocal detection of FLUX 680-labelled VAPB (magenta) and SNAP-Surface Alexa
Fluor 647-labelled SNAP/FLAG-hGLP-1R (green), in Ex-4-stimulated INS-1 832/3
SNAP/FLAG-hGLP-1R cells, showing crosstalk across detector channels (Cy5 near: 650 –
685 nm, Cy5 far: 685 – 720 nm); inset, region selected for MINFLUX imaging; size bar, 5
µm. **j**, Two-colour MINFLUX analysis of selected region from (**i**) after spectral separation,
including nearest neighbour calculation of average distance between FLUX 680 and Alexa
Fluor 647 clusters within the region indicated by dashed box. **k**, Ex-4-induced insulin
secretion (fold over 11 mM glucose) in non-targeting Control *versus* VAPB RNAi-treated INS-
1 832/3 cells; n=4, p value as indicated by paired t-test.

1816
Fig. 2. Internalisation-dependent hGLP-1R-VAPB binding and VAPB-localised GLP-1R
signalling. a, EGFP-VAPB co-IP with SNAP/FLAG-hGLP-1R in vehicle (Veh) *versus*
exendin-4 (Ex-4)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells co-expressing control
pcDNA3.1+ *versus* dominant negative dynamin (Dyn)1/2 K44A. **b**, Veh-subtracted Ex-4-
versus exendin-F1 (Ex-F1)-induced BRET kinetic traces (left) and corresponding AUCs
(right) of SNAP/FLAG-hGLP-1R-NLuc–Venus-VAPB interactions in INS-1 832/3 cells; n=3, p
value as indicated by paired t-test. **c** (top), Confocal microscopy analysis of SNAP/FLAG-
hGLP-1R localisation in mouse primary islets transduced with SNAP/FLAG-hGLP-1R
adenoviruses under Veh conditions or following 5-minute stimulation with the indicated
agonists [Ex-4, Ex-F1, exendin-D3 (Ex-D3), tirzepatide (Tirz) and semaglutide (Sema) used
at 100 nM; orforglipron (Orf) and danuglipron (Dan) used at 5 µM]; size bars, 20 µm; **c**
(bottom), Quantification of residual surface SNAP/FLAG-hGLP-1R levels in islets from

above; *n*=3. **d**, Quantification of EGFP-VAPB co-IP with SNAP/FLAG-hGLP-1R in INS-1 832/3 SNAP/FLAG-hGLP-1R cells stimulated for 5 minutes with the indicated agonists; *n*=4. **e**, VAPB-localised cAMP generation (measured by FluoSTEP) in response to stimulation with 100 nM Ex-4 *versus* Ex-F1 in INS-1 832/3 cells expressing pcDNA3-GFP11(x7)-VAPB + pcDNA3.1(+)-FluoSTEP-ICUE; maximal responses (to FSK+IBMX) shown; kinetic traces (left) and corresponding AUCs for the agonist-stimulated period (right) shown; *n*=5, p value as indicated by paired t-test. **f** (top), Ex-4 *versus* Ex-F1-induced global PKA activity, measured with the global PKA biosensor pcDNA3.1(+)- ExRai-AKAR2 in INS-1 832/3 cells; kinetic traces (left) and corresponding AUCs (right) shown; *n*=3; ns, non-significant by paired t-test. **f** (bottom), Ex-4 *versus* Ex-F1-induced mitochondrial PKA activity, measured with the mitochondrial PKA biosensor pcDNA3-mitoExRai-AKAR2 in INS-1 832/3 cells; kinetic traces (left) and corresponding AUCs (right) shown; *n*=3, p value as indicated by paired t-test. **g**, Mitochondrial over global PKA activity, calculated as AUC ratio from data in (**f**); *n*=3, p value as indicated by paired t-test.

Fig. 3. Human GLP-1R localisation to ERMCSs. a, Confocal microscopy analysis of SNAP/FLAG-hGLP-1R (red) co-localisation with ERMCSs (SPLICS Mt-ER Long P2A, green) in vehicle (Veh, left) *versus* exendin-4 (Ex-4, right)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells; size bars, 10 µm. **b**, SNAP/FLAG-hGLP-1R - SPLICS Mt-ER Long P2A co-localisation (Mander's coefficient) in Veh *versus* Ex-4-stimulated conditions from data in (**a**); *n*=5, p value as indicated by unpaired t-test. **c**, Time-lapse confocal microscopy analysis of SNAP/FLAG-hGLP-1R–ERMCS association, imaged as in (**a**); arrows indicate co-localised spots; time post-Ex-4 addition indicated; size bar, 10 µm. **d**, Confocal microscopy analysis of EFGP-VAPB (blue), mitochondria (MitoTracker Red, green) and SNAP/FLAG-hGLP-1R (SNAP-Surface Alexa Fluor 647, red) in an Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cell; size bar, 10 µm. **e**, Single confocal slice from an Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cell, with signals for EFGP-VAPB (green), SNAP/FLAG-hGLP-1R (yellow), mitochondria (magenta), and DAPI (blue) shown; white arrow points to a SNAP/FLAG-hGLP-1R-positive endosome used for CLEM analysis; size bar, 10 µm; also included CLEM overlay of aligned confocal slice with corresponding single slice of an EM tomogram (size bar, 10 µm) and magnification of EM tomogram slices (size bar, 1 µm) with indicated SNAP/FLAG-hGLP-1R-positive endosome (black arrow). **f**, 3D rendering of the segmentation of data from (**e**), with VAPB-positive ER (purple), mitochondria (orange) and SNAP/FLAG-hGLP-1R-positive endosome (yellow) shown. **g**, TEM micrograph of a 70-nm ultrathin section from resin embedded mouse islets transduced with SNAP/FLAG-hGLP-1R adenovirus, with SNAP/FLAG-hGLP-1R labelled with SNAP-Surface biotin + Alexa Fluor 488 Streptavidin, 5-nm colloidal gold prior to Ex-4 stimulation and fixation; ER (blue), mitochondria (yellow) and SNAP/FLAG-hGLP-1R-positive endosome (pink); inset shows magnified endosome area containing 5-nm gold particles; size bars, 100 nm.

Fig. 4. Human GLP-1R–SPHKAP association via shared VAPB binding and SPHKAP-dependent GLP-1R-induced VAPB-localised PKA activity. a, SPHKAP-EGFP co-IP with SNAP/FLAG-hGLP-1R in Control *versus* VAPB RNAi-treated INS-1 832/3 SNAP/FLAG-hGLP-1R cells under vehicle (Veh) *versus* exendin-4 (Ex-4)-stimulated conditions. **b**, Quantification of SPHKAP-EGFP over SNAP/FLAG-hGLP-1R levels from (**a**); *n*=4, p values as indicated by one-way ANOVA with Šidák post-hoc test. **c**, Confocal microscopy analysis of SPHKAP-EGFP *versus* mitochondria (imaged with MitoTracker Red) localisation in INS-1 832/3 cells treated with Control *versus* VAPB RNAi; size bars, 5 µm. **d**, Confocal microscopy analysis of SPHKAP-EGFP *versus* ER (imaged with ER-Tracker Red) localisation in INS-1 832/3 cells; size bar, 10 µm. **e**, SPHKAP-EGFP co-localisation (Mander's coefficient) with mitochondria (Mito) *versus* ER; *n*=5, p value as indicated by unpaired t-test. **f**, Ex-4-stimulated cAMP responses in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells, measured with cADDis cAMP biosensor; left, cAMP traces; right, AUC quantification; *n*=3; p value as indicated by paired t-test. **g**, Ex-4-stimulated global PKA responses in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells; left, PKA traces; right, AUC quantification;

n=4; p value as indicated by paired t-test. **h**, Ex-4-stimulated mitochondrial PKA responses
in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells; left, PKA traces; right, AUC
quantification; *n*=4, p value as indicated by paired t-test. **i**, VAPB-localised PKA activity
(measured by FluoSTEP) in response to 100 nM Ex-4 stimulation in Control *versus* SPHKAP
RNAi-treated INS-1 832/3 cells expressing pcDNA3-GFP11(x7)-VAPB + pcDNA3.1(+)-
FluoSTEP-AKAR; maximal responses (to FSK+IBMX) shown; kinetic traces (left) and
corresponding AUCs for the agonist-stimulated period (right) shown; *n*=5, p value as
indicated by paired t-test.
1892
Fig. 5. Regulation of GLP-1R function by SPHKAP. a, Exendin-4 (Ex-4)-stimulated
changes in mitochondrial membrane potential in Control *versus* SPHKAP RNAi-treated INS-
1 832/3 cells, measured with TMRE; left, membrane potential traces; G20, 20 mM glucose;
right, quantification of AUCs during the Ex-4 and G20 periods; *n*=5, p value as indicated by
paired t-test. **b**, Ex-4-stimulated ATP level in Control *versus* SPHKAP RNAi-treated INS-1
832/3 cells, measured with the cyto-Ruby3-iATPSnFR1.0 biosensor; left, ATP traces; right,
AUC quantification; *n*=3, p value as indicated by paired t-test. **c**, Ex-4-induced insulin
secretion (fold over 11 mM glucose) in Control *versus* SPHKAP RNAi-treated INS-1 832/3
cells; *n*=3, p value as indicated by paired t-test. **d**, Percentage of apoptosis +/- thapsigargin
exposure in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells under vehicle (Veh)
versus Ex-4-stimulated conditions; *n*=3; p values as indicated by one-way ANOVA with Šidák
post-hoc test. **e**, PKA-dependent phosphorylation of MIC19-3xHA in INS-1 832/3 cells under
Veh *versus* Ex-4-stimulated conditions, determined by Western blotting of anti-HA
immunoprecipitates with an anti-RRx-pS/T antibody; forskolin (FSK), positive control for PKA
activation; left, representative blots; right, quantification of phospho-MIC19 levels; *n*=3-4, p
value as indicated by one-way ANOVA with Šidák post-hoc test. **f**, Ex-4-induced mitophagy,
measured as changes in ratio fluorescence intensities (580/440 nm) over time with the dual-
excitation ratiometric biosensor mKeima-Red-Mito-7 in Control *versus* SPHKAP RNAi-
treated INS-1 832/3 cells; *n*=6. **g**, Time-lapse confocal microscopy imaging showing Drp1
localisation to mitochondria in Veh *versus* Ex-4-stimulated INS-1 832/3 cells; mitochondria
(mito-BFP, blue) and Drp1 (mCherry-Drp1, red); arrows indicate Drp1-positive spots on
mitochondria; time post-Veh or Ex-4 exposure indicated; size bars, 5 µm. **h**, Changes in
Drp1 phosphorylation at Ser 616 in Control *versus* SPHKAP RNAi-treated Ex-4-stimulated
INS-1 832/3 cells; left, representative blots; right, quantification; *n*=4; p values as indicated
by two-way ANOVA with Tukey post-hoc test.
1918
1919 **Fig. 6. GLP-1R/SPHKAP-dependent mitochondrial remodelling. a**, Confocal microscopy
1920 analysis of mitochondrial morphology in vehicle (Veh) *versus* exendin-4 (Ex-4)-stimulated
1921 WT INS-1 832/3 cells; top, representative images with mitochondria labelled with
1922 MitoTracker Red; size bar, 10 µm; bottom, percentage of cells classified as presenting
1923 hyperfused, tubular or fragmented mitochondria per condition; *n*=4, p values as indicated by
1924 two-way ANOVA with Tukey post-hoc test. **b**, as in (**a**) for INS-1 832/3 GLP-1R KO cells;
1925 *n*=4; p values as indicated by two-way ANOVA with Tukey post-hoc test. **c**, as in (**a**) for
1926 Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells under Veh *versus* Ex-4-stimulated
1927 conditions; *n*=5, p values as indicated by two-way ANOVA with Tukey post-hoc test. **d**,
1928 Maximum intensity projections (MIP) from confocal z-stacks of mitochondria in vehicle (Veh)
1929 *versus* Ex-4-stimulated Control and SPHKAP RNAi-treated INS-1 832/3 cells; size bars, 10
1930 µm. **e**, Representative images of mitochondrial morphology in Control *versus* SPHKAP
1931 RNAi-treated INS-1 832/3 cells under Veh *versus* Ex-4-stimulated conditions, imaged by
1932 super-resolved STED microscopy; size bars, 2 µm.
1933
Fig. 7. Exendin-4-induced changes in mitochondrial morphology and number of
ERMCSs per mitochondria. a, Representative TEM micrographs of resin-embedded
ultrathin sections depicting mitochondrial morphology in Control *versus* SPHKAP RNAi-
treated INS-1 832/3 cells, and INS-1 832/3 GLP-1R KO cells, under vehicle (Veh) *versus*
exendin-4 (Ex-4)-stimulated conditions; size bars, 100 nm. **b**, Quantification of mitochondrial

area and number of cristae per mitochondria from ~200 TEM images collected from n=3
independent experiments, p values as indicated by one-way ANOVA with Šidák post-hoc
test. **c**, Quantification of ERMCS length in Control *versus* SPHKAP RNAi-treated INS-1
832/3 cells and INS-1 832/3 GLP-1R KO cells from (**a**), as well as SPHKAP-EGFP-
expressing WT INS-1 832/3 cells; p values as indicated by one-way ANOVA with Šidák post-
hoc test. **d**, Representative confocal images of WT INS-1 832/3 cells co-expressing the
mitochondrial marker 4xMTS-Halo (green) and splitFAST reporters ER-RspA-NFAST and
OMM-short-RspA-CFAST, labelled with a Lime splitFAST ligand to detect ERMCSs
(magenta) under Veh *versus* Ex-4-stimulated conditions; size bars, 5 µm. **e**, Quantification of
the number of ERMCSs per mitochondrial area from data in (**d**); results per cell (left) and per
independent experiment (right) shown; 50-57 cells imaged in n=3 independent experiments,
p values as indicated by unpaired t-tests.
1951
1952 **Fig. 8. SPHKAP control of GLP-1R function in primary mouse islets. a**, PLAs showing
1953 interactions between endogenous GLP-1R and VAPB (left) or SPHKAP (right) in exendin-4
1954 (Ex-4)-stimulated primary mouse islets; size bars, 20 µm. **b**, Ex-4-induced changes in ATP
1955 level (normalised to total protein) in Control *versus* SPHKAP shRNA-transduced primary
1956 mouse islets; n=5, p value as indicated by one-way ANOVA with Šidák post-hoc test. **c**, Ex-
1957 4-induced changes in thapsigargin-triggered ROS accumulation in Control *versus* SPHKAP
1958 shRNA-transduced primary mouse islets; n=6, p value as indicated by one-way ANOVA with
1959 Šidák post-hoc test. **d**, Ex-4-induced potentiation of insulin secretion in Control *versus*
1960 SPHKAP shRNA-transduced primary mouse islets; n=9, p values as indicated by one-way
1961 ANOVA with Šidák post-hoc test; fold over 11 mM glucose (G11) also shown, p value as
1962 indicated by paired ratio t-test.
1963
1964 **Fig. 9. SPHKAP control of GLP-1R-dependent mitochondrial remodelling in primary**
1965 **mouse islets. a**, Confocal microscopy analysis of mitochondrial morphology in vehicle (Veh)
1966 *versus* exendin-4 (Ex-4)-stimulated WT primary mouse islets; top, representative images
1967 with MitoTracker Red-labelled mitochondria; size bar, 20 µm; bottom, percentage of islet
1968 cells classified as presenting hyperfused, tubular or fragmented mitochondria per condition;
1969 n=5 islet preps from separate mice, p values as indicated by two-way ANOVA with Tukey
1970 post-hoc test. **b**, as in (**a**) for islets from GLP-1R KO mice; n=5 islet preps from separate
1971 mice; p values as indicated by two-way ANOVA with Tukey post-hoc test. **c**, as in (**a**) for
1972 Control *versus* SPHKAP shRNA-transduced mouse islets under Veh *versus* Ex-4-stimulated
1973 conditions; n=6 islet preps from separate mice, p values as indicated by two-way ANOVA
1974 with Tukey post-hoc test. **d**, as in (**a**) for Control *versus* SPHKAP RNAi-treated dispersed
1975 primary mouse islet cells under Veh *versus* Ex-4-stimulated conditions; size bars, 5 µm; n=4
1976 islet preps from separate mice, p values as indicated by two-way ANOVA with Tukey post-
1977 hoc test.
1978
1979 **Fig. 10. GLP-1R–VAPB/SPHKAP interactions and GLP-1R-dependent mitochondrial**
1980 **remodelling in primary human islets. a**, PLA showing interactions between endogenous
1981 GLP-1R and VAPB (left) or SPHKAP (right) in exendin-4 (Ex-4)-stimulated primary human
1982 islets; size bars, 20 µm. **b**, Negative control (no primary antibody) for PLA data shown in (**a**);
1983 nuclei (DAPI), blue; size bar, 20 µm. **c**, Confocal microscopy analysis of mitochondrial
1984 morphology in vehicle (Veh) *versus* Ex-4-stimulated primary human islets; representative
1985 images with mitochondria labelled with MitoTracker Red; size bars, 20 µm; inset,
1986 magnification areas. **d**, Percentage of human islet cells classified as presenting hyperfused,
1987 tubular or fragmented mitochondria per condition; n=5 islet preps from separate human
1988 donors, p values as indicated by two-way ANOVA with Tukey post-hoc test.
1989
1990 **Supplementary Figure Legends**
1991
1992 **Supplementary Fig. 1. Human GLP-1R β-cell interactome and validation of VAPB-**
1993 **hGLP-1R interaction – Extra data. a**, Principal component analysis (PCA) of biological

repeats from interactome data from Fig. 1a,b. **b**, Pathway enrichment analysis using gProfiler; results show 10 most significant gene ontology terms represented in the interactome data from Fig.1a,b. **c**, SNAP/FLAG-hGLP-1R levels in input *versus* output supernatants from co-IP experiments from Fig. 1e. **d**, Confocal microscopy analysis of SNAP/FLAG-hGLP-1R (red) co-localisation with EGFP-VAPB (green) in vehicle (Veh, left) *versus* exendin-4 (Ex-4, right)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells; size bars, 10 μm. **e**, SNAP/FLAG-hGLP-1R - EGFP-VAPB co-localisation (Mander's coefficient) in Veh *versus* Ex-4-stimulated conditions from data in (**d**); n=7, p value as indicated by unpaired t-test. **f**, Time-series confocal microscopy analysis of SNAP/FLAG-hGLP-1R (red) and EGFP-VAPB (green) localisation in response to exendin-4 (Ex-4) in INS-1 832/3 SNAP/FLAG-GLP-1R cells; time post-Ex-4 addition indicated; size bar, 5 μm. **g**, EGFP-VAPB WT *versus* P56S mutant co-IP with SNAP/FLAG-hGLP-1R in vehicle (Veh) *versus* Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells. **h**, Confocal microscopy analysis of SNAP/FLAG-hGLP-1R (red) co-localisation with EGFP-VAPB WT or P56S (green) in Veh *versus* Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells; size bars, 5 μm. **i**, SNAP/FLAG-hGLP-1R - EGFP-VAPB P56S co-localisation (Mander's coefficient) in Veh *versus* Ex-4-stimulated conditions from data in (**h**) (left), and comparison of SNAP/FLAG-hGLP-1R co-localisation with EGFP-VAPB WT *versus* P56S following Ex-4 stimulation (right); n=5-7, p values as indicated by unpaired t-test. **j**, Western blot analysis of endogenous VAPB levels in Control *versus* VAPB RNAi-treated INS-1 832/3 cells; representative blots and quantification of VAPB knockdown shown; n=3, p value as indicated by paired t-test. **k**, Ex-4-induced potentiation of insulin secretion in Control *versus* VAPB RNAi-treated INS-1 832/3 cells; n=4, p values as indicated by one-way ANOVA with Šidák post-hoc test. **l**, Ex-4-induced insulin secretion in pcDNA3.1+, EGFP-VAPB WT and P56S-transfected INS-1 832/3 cells; fold over pcDNA3.1+ also shown, n=5; p values as indicated by one-way ANOVA with Šidák post-hoc test, or by paired t-test.

Supplementary Fig. 2. Internalisation-dependent hGLP-1R-VAPB binding and VAPB-localised GLP-1R signalling – Extra data. a, Confocal microscopy analysis of SNAP/FLAG-hGLP-1R localisation in exendin-4 (Ex-4)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells co-expressing dominant negative dynamin (Dyn)1/2 K44A; white arrow indicates plasma membrane SNAP/FLAG-hGLP-1R localisation with Dyn1/2 K44A co-expression; yellow arrow indicates endosomal SNAP/FLAG-hGLP-1R localisation in a non-Dyn1/2 K44A expressing cell; size bar, 5 μm. **b**, Schematic of the hGLP-1R-Nanoluciferase (NLuc)–Venus-VAPB interaction NanoBRET assay. **c**, Representative blots from EGFP-VAPB co-IP with SNAP/FLAG-hGLP-1R in INS-1 832/3 SNAP/FLAG-hGLP-1R cells stimulated for 5 minutes with: vehicle (Veh), Ex-4, exendin-F1 (Ex-F1), exendin-D3 (Ex-D3), tirzepatide (Tirz), semaglutide (Sema) all used at 100 nM; orforglipron (Orf) and danuglipron (Dan) used at 5 μM. **d**, Schematic of the FluoSTEP method used to quantify GLP-1R-induced VAPB-localised cAMP generation; Epac1 (149-881), exchange protein activated by cAMP, amino acids 149-881; RFP, red fluorescent protein; $sfGFP_{1-10}$, superfolder green fluorescent protein helices 1-10; GFP_{11}, 11th helix of sfGFP; VAPB, vesicle-associated membrane protein-associated protein B. **e**, Localisation of the excitation-ratiometric global PKA biosensor pcDNA3.1(+)-ExRai-AKAR2 and mitochondrial PKA biosensor pcDNA3-mitoExRai-AKAR2 in INS-1 832/3 cells; size bars,10 μm. **f**, Ex-4-stimulated global and mitochondrial PKA responses in INS-1 832/3 cells, measured with the biosensors from (**e**); Forskolin (FSK) + IBMX added to record maximal responses.

Supplementary Fig. 3. Human GLP-1R β-cell interactome analysis with further agonists. Heatmap of GLP-1R interactor enrichment analysis from anti-FLAG co-immunoprecipitates from INS-1 832/3 SNAP/FLAG-hGLP-1R cells stimulated for 5 minutes with the indicated agonists; blue, decreased; red, increased enrichment; LC-MS/MS data analysed in LFQ-Analyst and normalised to hGLP-1R levels for each experimental sample; colour scale centred around overall median value; n=4 for exendin-4 (Ex-4), exendin-F1 (Ex-F1) and exendin-D3 (Ex-D3); n=3 for GLP-1.

2049
2050 **Supplementary Fig. 4. Human GLP-1R localisation to ERMCSs – Extra data. a**, Single
2051 cryo-confocal slice from an exendin-4 (Ex-4)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R
2052 cell, with signals for EGFP-VAPB (cyan), SNAP/FLAG-hGLP-1R (magenta), and
2053 mitochondria (yellow); inset, magnified area; arrow points to SNAP/FLAG-hGLP-1R-positive
2054 endosome used for cryo-CLEM analysis; size bar, 5 µm. **b**, Overlay of a cryo-confocal slice
2055 aligned with the corresponding single slice of a resliced 3D cryoFIB/SEM volume. **c**, 3D
2056 rendering of the segmentation of the FIB-SEM volume data from (**b**), with ER (cyan),
2057 mitochondria (yellow) and SNAP/FLAG-hGLP-1R-positive endosome (magenta) shown.
2058
Supplementary Fig. 5. Gene expression data and genetic association of *SPHKAP*
variants with T2D and BMI. a, Schematic of the AKAP SPHKAP protein primary structure.
AKB, A-kinase binding domain; pFFAT, phospho-FFAT motif. Alignment of SPHKAP FFAT
motif to the canonical FFAT sequence depicted; Alcoholic residues are highlighted in blue,
acidic residues in green, and aromatic residues in red; all other residues are in black. **b**,
Analysis of *SPHKAP* mRNA levels from primary islets isolated from WT *versus* heterozygous
and homozygous *db/db* mice; data obtained from RNA Seq dataset GSE107489[112]; *n*=3, p
values as indicated by one-way ANOVA with Dunnett post-hoc test. **c**, *SPHKAP* gene region
variant associations from T2D European ancestry[36] and multi-ancestry[37] GWAS; y-axes
represent significance (-$\log_{10}$ p-value) of phenotype–variant associations and x-axes
genomic location of the variants (NCBI human genome build37). Secondary y-axes
represent recombination fraction from 1000 Genomes Project and horizontal dashed lines
highlight association threshold for significance (p=5×10^{-8}). Lead variant with the highest
significance is shown as a purple diamond and correlation (LD, r^2) of other SNPs in the lead
variant region represented with colours as per box legend. Variants previously reported for
associations with phenotypes in GWAS catalog (www.ebi.gwas.org) are indicated and
respective phenotypes listed. **d**, As for (**c**) for BMI GWAS meta-analyses[38, 39]. **e**, Comparison
of the effects of *SPHKAP* gene region variants showing at least nominal association (p<0.05)
with both T2D risk and high adiposity (BMI) from GWAS meta-analyses from (**c**,**d**). X-axis
[odds ratio (OR) with 95% confidence interval] corresponds to associations with T2D, while
y-axis (beta with standard error) corresponds to associations with BMI. Variant points are
scaled in size to their significance with T2D (z-score from association test). Seven *SPHKAP*
single nucleotide polymorphisms (SNPs), 2 for T2D and 5 for high BMI, identified for T2D /
BMI from GWAS Catalog (www.ebi.gwas.org) as being in high linkage disequilibrium (LD)
with each other, are indicated. **f**, *SPHKAP* coding variant effects on random glucose (RG)
from UKBB whole-exome sequencing (WES) data for ~450,000 individuals; y-axis
represents significance (-$\log_{10}$ p-value) of variant association and x-axis represents the
genomic location of variants (NCBI human genome build37). Secondary y-axis represents
recombination fraction from 1000 Genomes Project and horizontal dashed line highlights the
association threshold for significance (p=5×10^{-8}). RG levels in mmol/L are natural logarithm
transformed and adjusted for time since last meal (t) as t+t^2+t^3, age, sex, and six principal
components (derived from UKBB GWAS data).
2091
Supplementary Fig. 6. Human GLP-1R–SPHKAP association via shared VAPB binding
– Extra data. a, SNAP/FLAG-hGLP-1R levels in input *versus* output supernatants from co-IP
experiments from Fig. 4a,b. **b**, Endogenous SPHKAP co-IP with SNAP/FLAG-hGLP-1R in
vehicle (Veh) *versus* exendin-4 (Ex-4)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells.
c, Endogenous SPHKAP co-IP with SNAP/FLAG-hGIPR in Veh *versus* GIP-stimulated INS-1
832/3 SNAP/FLAG-hGIPR cells. **d**, z-stack series of confocal microscopy images of
SPHKAP-EGFP and mitochondria (imaged with MitoTracker Red) in INS-1 832/3 cells; size
bar, 10 µm. **e**, SPHKAP-EGFP pFFAT motif mutant (DFLTESE) co-IP with SNAP/FLAG-
hGLP-1R in Veh *versus* Ex-4-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells. **f**,
Confocal microscopy analysis of SPHKAP-EGFP WT *versus* pFFAT motif mutant in INS-1
832/3 cells; size bars, 5 µm. **g**, Western blot analysis of SNAP/FLAG-hGLP-1R, VAPB, and
SPHKAP in Tom20-positive MAM fraction purified from Ex-4-stimulated INS-1 832/3

SNAP/FLAG-hGLP-1R cells. **h**, Endogenous SPHKAP - EGFP-VAPB co-IP in INS-1 832/3 cells, including input and output fractions. **i**, Western blot analysis of endogenous SPHKAP levels in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells. Quantification of SPHKAP knockdown also shown; n=3, p value as indicated by paired t-test. **j**, Western blot analysis of SPHKAP-EGFP levels in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells.

Supplementary Fig. 7. Regulation of GLP-1R function by SPHKAP – Extra data. a, EGFP-VAPB co-IP with SNAP/FLAG-hGLP-1R in Control *versus* SPHKAP RNAi-treated INS-1 832/3 SNAP/FLAG-hGLP-1R cells under vehicle (Veh) *versus* exendin-4 (Ex-4)-stimulated conditions. **b**, Confocal microscopy analysis of EGFP-VAPB co-localisation with SNAP/FLAG-hGLP-1R in Veh *versus* Ex-4-stimulated conditions in Control *versus* SPHKAP RNAi-treated INS-1 832/3 SNAP/FLAG-hGLP-1R cells; size bars, 10 µm. **c**, SNAP/FLAG-hGLP-1R - EGFP-VAPB co-localisation (Mander's coefficient) in Ex-4-stimulated Control *versus* SPHKAP RNAi-treated INS-1 832/3 SNAP/FLAG-hGLP-1R cells from data in (**b**); n=5, p value as indicated by unpaired t-test. **d**, Schematic of the FluoSTEP method used to quantify GLP-1R-induced VAPB-localised PKA activity; FHA1, forkhead-associated 1 phospho-peptide binding protein; PKA, protein kinase A; RFP, red fluorescent protein; sfGFP$_{1-10}$, superfolder green fluorescent protein helices 1-10; GFP$_{11}$, 11th helix of sfGFP; VAPB, vesicle-associated membrane protein-associated protein B. **e**, Confocal microscopy analysis of PKA-RIα-EGFP localisation in Veh *versus* Ex-4-stimulated INS-1 832/3 cells, size bars, 10 µm; arrows indicate newly formed PKA-RIα-EGFP puncta. **f**, Ex-4-induced changes in ATP level in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells; n=3, p values as indicated by one-way ANOVA with Šidák post-hoc test. **g**, Ex-4-induced potentiation of insulin secretion in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells; n=3, p value as indicated by one-way ANOVA with Šidák post-hoc test. **h**, Ex-4-induced potentiation of insulin secretion in INS-1 832/3 cells transfected with EGFP *versus* SPHKAP-EGFP; n=3; p values as indicated by one-way ANOVA with Šidák post-hoc test; fold over 11 mM glucose (G11) also shown; p value as indicated by paired t-test. **i**, Ex-4 induced changes in ER Ca^{2+}, measured with pCAG G-CEPIA1er, in Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells; n=4. **j**, As in (**i**) for mitochondrial Ca^{2+}, measured with pCAG mito-RCaMP1h; peak response times indicated per condition; n=4. **k**, PKA-dependent phosphorylation of MICOS complex components isolated by MIC19-3xHA co-IP in Control *versus* VAPB+SPHKAP RNAi-treated INS-1 832/3 cells under Veh *versus* Ex-4-stimulated conditions, determined by Western blotting with an anti-RRx-pS/T antibody; arrows indicate proteins showing changes in PKA-dependent phosphorylation. **l**, Mitochondrial DNA (mtDNA) copy number, measured as a ratio of mitochondrial tRNA^{Pro} over hexokinase (HK) in Veh *versus* Ex-4-stimulated Control and SPHKAP RNAi-treated INS-1 832/3 cells, as well as INS-1 832/3 GLP-1R KO cells; n=7; p values as indicated by two-way ANOVA with Tukey post-hoc test.

Supplementary Fig. 8. GLP-1R and SPHKAP-dependent mitochondrial remodelling – Extra data. a, Overlay of a single confocal slice, with signals for VAPB (green), GLP-1R (yellow), mitochondria (magenta), and DAPI (blue) shown, and corresponding single slice of an EM tomogram following CLEM analysis of exendin-4 (Ex-4)-stimulated INS-1 832/3 SNAP/FLAG-hGLP-1R cells; size bar, 1 µm; magnified confocal slice region with arrow pointing to SNAP/FLAG-hGLP-1R positive signal also shown; size bar, 1 µm. **b**, Magnification of tomogram slices corresponding to magnified confocal slice from (**a**); arrow points to the same SNAP/FLAG-hGLP-1R-positive lysosome as in (**a**) embedded in a mitochondrion, indicating mitophagy; size bar, 1 µm. **c**, Single channel images corresponding to mito-BFP (blue) and mCherry-Drp1 (red) signals from Fig. 5g. **d**, Quantification of mitochondrial mean form factor and mean area in Veh *versus* Ex-4-stimulated WT and GLP-1R KO INS-1 832/3 cells; n=5, p values as indicated by one-way ANOVA with Šidák post-hoc test. **e**, As in (**d**) in Veh *versus* Ex-4-stimulated Control *versus* SPHKAP RNAi-treated INS-1 832/3 cells; n=5, p values as indicated by one-way ANOVA with Šidák post-hoc test.

Supplementary Fig. 9. GLP-1R–VAPB/SPHKAP interactions and GLP-1R-dependent mitochondrial remodelling in primary islets – Extra data. a, Negative control (no primary antibody) for PLA data shown in Fig. 8a; nuclei (DAPI), blue; size bar, 20 µm. **b**, PLA showing interactions between endogenous GLP-1R and SPHKAP (green) in vehicle (Veh) and exendin-4 (Ex-4)-stimulated primary mouse islets with insulin co-staining (red); nuclei (DAPI), blue; size bars, 20 µm. **c**, Quantification of SPHKAP mRNA levels by qPCR in Control *versus* SPHKAP shRNA-transduced intact primary mouse islets; *n*=3, p value as indicated by paired t-test. **d**, Example of cell segmentation into hyperfused (blue), tubular (purple) or fragmented (pink) mitochondrial phenotypes in a primary mouse islet. **e**, Quantification of mitochondrial mean form factor and mean area in Control *versus* SPHKAP RNAi-treated dispersed mouse islet cells under Veh *versus* Ex-4-stimulated conditions; *n*=4, p values as indicated by one-way ANOVA with Šidák post-hoc test.

Supplementary Movie 1. Time-lapse confocal microscopy imaging of SNAP/FLAG-hGLP-1R (red) and EGFP-VAPB (green) localisation in response to exendin-4 stimulation in INS-1 832/3 SNAP/FLAG-hGLP-1R cells; images taken every 6 seconds, shown at 5 frames per second.

Supplementary Movie 2. Time-lapse confocal microscopy imaging of SNAP/FLAG-hGLP-1R (red) and ERMCS (SPLICS Mt-ER Long P2A, green) localisation in response to exendin-4 stimulation in INS-1 832/3 SNAP/FLAG-hGLP-1R cells; images taken every 6 seconds, shown at 5 frames per second; acquisition starts at 60 seconds post-exendin-4 exposure.

Supplementary Movie 3. Z-stack EM tomogram slices from CLEM data in Fig. 3e, used for image segmentation and 3D rendering shown in Fig. 3f.

Supplementary Movie 4. Time-lapse confocal microscopy imaging of Drp1 (mCherry-Drp1, red) and mitochondria (mito-BFP, blue) localisation in vehicle-exposed INS-1 832/3 cells; images taken every 6 seconds, shown at 5 frames per second; acquisition starts at 60 seconds post-vehicle exposure.

Supplementary Movie 5. Time-lapse confocal microscopy imaging of Drp1 (mCherry-Drp1, red) and mitochondria (mito-BFP, blue) localisation in exendin-4-stimulated INS-1 832/3 cells; images taken every 6 seconds, shown at 5 frames per second; acquisition starts at 60 seconds post-vehicle exposure.

c

Name	Localisation	Function
VAPB	ER-Mitochondria contacts	lipid and ion transport
VAPA	ER-Mitochondria contacts	lipid and ion transport
HADHB	Mitochondria	β-oxidation of long chain fatty acids
FAAH	ER-Mitochondria contacts	fatty acid amide hydrolysis
HADHA	Mitochondria	β-oxidation of long chain fatty acids
APGAT4	ER / Mitochondria	phosphatidic acid biosynthesis
DHCR7	ER	cholesterol biosynthesis
VDAC2	ER-Mitochondria contacts	metabolite transport, mPTP component
CDS2	ER	phosphatidylinositol biosynthesis
ATP2A3	ER-Mitochondria contacts	SERCA pump, calcium transport
ATP5B	Mitochondria	ATP synthase subunit
ATP5F1	Mitochondria	ATP synthase subunit
PGRMC1	ER / Mitochondria	cholesterol metabolism
PTDSS1	ER	phosphatidylserine biosynthesis
Rab18	ER	lipid droplet biogenesis, lipolysis
SPHKAP	Mitochondria	PKA anchoring protein
TAMM41	Mitochondria	phosphatidylinositol biosynthesis
CNP	Mitochondria	calcium-induced opening of mPTP
FADS1	Mitochondria	fatty acid unsaturation
ATP5D	Mitochondria	ATP synthase subunit
ATP5A1	Mitochondria	ATP synthase subunit
BDH1	Mitochondria	fatty acid catabolism
POR	ER	cholesterol biosynthesis
REEP6	ER-Mitochondria contacts	GPCR regulation of mitochondrial function
HMGCR	ER	cholesterol biosynthesis
SDHA	Mitochondria	mitochondrial respiratory chain complex

a

Merged	SPLICS Mt-ER Long P2A	SNAP/FLAG hGLP-1R		Merged	SPLICS Mt-ER Long P2A	SNAP/FLAG hGLP-1R
Veh				Ex-4		

b

Co-localisation (Manders' coefficient)

0.0068

Veh   Ex-4

c

60"   96"   126"   156"   186"

d

EGFP-VAPB   MitoTracker Red   SNAP/FLAG hGLP-1R

e

MitoTracker Red
EGFP-VAPB
SNAP/FLAG hGLP-1R
DAPI

f

VAPB ER
Mitochondria
GLP-1R Endosome
Mitochondria
VAPB ER

g SNAP/FLAG-hGLP-1R, 5-nm gold

Mitochondria
ER
Endosome
Mitochondria

Figure 5