Heterogeneity of the human immune response to malaria infection and vaccination

driven by latent cytomegalovirus infection

Reena Mukhiya[1,2,3], Wim A. Fleischmann[4], Jessica R. Loughland[1,3], Jo-Anne Chan[1,5,6],

Fabian de Labastida Rivera[3], Dean Andrew[3], James G. Beeson[1, 5, 6], James S McCarthy[3, 7],

Bridget E Barber[3], J. Alejandro Lopez [2,3], Christian Engwerda[2,3], Richard Thomson-Luque [8,9],

Michelle J Boyle[1,2,3,5,6 #]

Author affiliations

1. Burnet Institute, Melbourne, Australia

2. School of Environmental Sciences, Griffith University, Brisbane Australia

3. QIMR Berghofer Medical Research Institute, Brisbane, Australia

4. Heidelberg University, Medical Faculty, University Hospital Heidelberg, Center for

Infectious Diseases, Virology

5. Department of Infectious Diseases, University of Melbourne

6. Department of Microbiology and School of Translational Medicine, Monash University

7. Walter and Eliza Hall Institute of Medical Research, Melbourne, Australia

8. Sumaya-Biotech GmbH & Co. KG,

9. Heidelberg University, Medical Faculty, University Hospital Heidelberg, Centre for

Infectious Diseases, Parasitology

corresponding author, michelle.boyle@burnet.edu.au

Abstract

Human immune responses to infection and vaccination are heterogenous, driven by multiple

factors including genetics, environmental exposures and personal infection histories. For

malaria caused by *Plasmodium falciparum* parasites, host factors that impact on humoral

immunity are poorly understood. We investigated the role of latent cytomegalovirus (CMV)

on the host immune response to malaria using blood stage *P. falciparum* Controlled Human

Malaria Infection (CHMI) and in a MSP1 vaccine Phase 1a clinical trial. CMV seropositivity

was associated with reduced induction of parasite specific antibodies following malaria

infection and vaccination. During infection, reduced antibody induction was associated with

modifications to the T -follicular helper (Tfh) cell compartment. CMV seropositivity was

associated with a skew towards Tfh1 cell subsets before and after malaria infection, and

reduced activation of Tfh2 cells. Protective Tfh2 cell activation was only associated with

antibody development in CMV seronegative individuals, and a higher proportion of Tfh1

cells was associated with lower antibody development in CMV seropositive individuals.

During MSP1 vaccination, reduced antibody induction in CMV seropositive individuals was

associated with CD4 T cell expression of terminal differentiation marker CD57. These

findings are particularly relevant for malaria endemic regions where CMV infection is

acquired early in life and may modify immunity to malaria gained during infection or

vaccination.

44
45
46

Introduction

The human immune response to infection and vaccination is heterogenous. Responses range from robust responses that resolve infection, to sub-optimal responses that may lead to severe disease or fail to generate long-term protection. This heterogeneity reflects the interplay of genetics, environment, and personal exposure histories. For malaria caused by *P. falciparum* parasites, infection in endemic areas ranges from asymptomatic parasitemia to severe disease and death [1]. At the population level, the largest disease burden occurs in children under the age of five, who develop immunity after repeated infection throughout childhood. However, the rate of immune development is heterogeneous, with some children rapidly gaining protection while others experience numerous symptomatic episodes, despite similar exposure [2]. Heterogeneity is also seen in responses to malaria vaccination, both between individuals within malaria endemic areas, and between populations across geographical locations [3–5]. Of note, responses to *P. falciparum* malaria vaccines are consistently lower in low- and middle- income countries (LMIC) compared to those reported during Phase 1 trials in high income countries. For example, antibodies to circumsporozoite protein (CSP) following experimental vaccination with radiation-attenuated whole sporozoite parasite vaccine (PfSPZ) were significantly lower in Tanzanian and Malian adults, compared to adults in the USA [6]. Further, CSP antibodies induced in Kenyan adults with the licenced malaria subunit vaccine RTS,S are generally lower than those seen following vaccination of adults in the USA [7–10]. As such, dissecting factors that modify immune responses to *P. falciparum* may inform development of second-generation vaccines with better protection, or other avenues for malaria control.

Immunity to malaria generated by infection or vaccination is mediated by antibodies that prevent parasite replication, largely via Fc mediated mechanisms [11,12]. Antibody induction

is supported by the CD4[+] T cell compartment, particularly T follicular CD4[+] T (Tfh) cells,

which activate B cells to drive germinal centre activation and antibody development [13],

including during malaria [14]. In human infection, Tfh cell subsets can be identified based

on CXCR3 and CCR6 chemokine expression, and specific Tfh cell subsets are associated

with the development of antibodies following infection and vaccination in a context

dependent manner [15]. We have previously evaluated the role of Tfh cells in antibody

development during *P. falciparum* malaria, using the powerful controlled human malaria

infection (CHMI) model that allows the longitudinal analysis of the immune response to a

single infection without the confounding effects of prior malaria exposure and heterogeneity

in timing of infection that occur when investigating immunity in naturally exposed

individuals [16]. We showed that early activation of Tfh2 cells was associated with induction

of functional antibodies, and Tfh1 cells were associated with short lived plasmablasts which

have negative roles in germinal centre formation [17,18]. While age has been shown to

influence Tfh2 and Tfh1 proportions and activation [19,20], other host factors that influence

Tfh cells during malaria are unknown.

One of the multiple factors known to modify the human immune response is cytomegalovirus

(CMV) infection, a ubiquitous beta-herpes virus which establishes a life-long persistence and

results in major remodelling of the immune response [21]. In monozygotic twins with

discordant CMV serostatus, over 50% of immune parameters tested were influenced by CMV

seropositivity [22], with impacts across the immune landscape, including monocytes, NK

cells, and T cells [23]. Within the T cell compartment, CMV seropositivity has been

associated with the expansion of memory cells that lose expression of costimulatory receptors

CD27 and CD28, and gain expression of markers of terminal differentiation including CD57

[24]. While these changes are most striking in CD8[+] T cells, they also occur within CD4[+] T

cells [25]. This immune remodelling has consequences for subsequent pathogenesis of a

broad range of diseases and induction of antibody following infection with other pathogens

[26]. The global seroprevalence of CMV is 83%, which ranges from less than 50% in high

income countries to up to 100% in LMIC [27], where the majority of infants are infected in

the first year of life [28]. These geographical differences in CMV seroprevalence have

implications for vaccine efficacy for pathogens where CMV can modify the immune response

[29]. Whether CMV can modulate the host immune response to malaria infection or

vaccination is unknown.

To investigate the impact of CMV serostatus on antibody induction during malaria infection

and after vaccination, we investigated antibody induction in CMV seronegative and

seropositive individuals enrolled in CHMI studies [17] and in a phase 1a clinical trial of a

full-length MSP1 vaccine candidate [30,31]. We found that CMV seropositivity was

associated with reduced antibody induction following both CHMI and vaccination, and that

these reductions were associated with remodelling of the CD4+ T cell compartment. These

findings have implications for the development of malaria immunity, either induced by

infection or vaccination, in children in endemic areas where CMV infection occurs in

infancy, and CMV seropositivity reaches 100% [28,32].

Results

Latent CMV infection is associated with reduced antibody induction following controlled

human malaria infection in adults

To investigate the impact of latent CMV infection (quantified as CMV seropositive) on the

immune response to *P. falciparum* infection, we analysed antibody and Tfh responses in 40

malaria-naïve adults during blood stage CHMI (median age 25.5, range 18-52 years) [17].

Within the cohort, 21/40 (52%) individuals were seropositive for CMV. Sex, Epstein-Barr

virus (EBV) seropositivity and age, were comparable between the CMV negative and

positive individuals (Table 1).

As previously reported, the induced antibody response was measured as an antibody score,

that captured the breadth, magnitude and functionality of antibodies to the merozoite parasite

stage, and major merozoite antigen, merozoite surface protein 2 (MSP2) [17]. Antibody score

was significantly lower in CMV seropositive individuals (Fig. 1 A). There was no difference

in antibody score with EBV status, and antibody score was not associated with age (Fig. 1 B-

D).

Table 1: Study populations demographics

	CMV serostatus	
	Negative	Positive
Total, n (%)	19, (48%)	21, (52%)
Sex, male, n (%)	18 (95%)	18 (86%)
EBV, positive, n (%)	18 (95%)	17 (81%)
Age years, median [IQR]	26 [20.25-31.75]	25 [21-29]

Figure 1. **Antibodies induced by controlled human malaria infection are reduced in adults**
with latent CMV infection
IgM, IgG and functional antibodies to merozoite and major merozoite antigen MSP2 were
quantified and antibody score used to capture the total magnitude, breadth and functionality
of the responses. Antibody score stratified by (A) CMV serostatus (CMV- n=19, CMV+
n=21), (B) EBV serostatus (EBV– n=5, EBV+ n=35), (C) Sex (female n=3, male n=37), and
(D) correlated with age. A-C data is Tukey boxplots with the median, 25th and 75th
percentiles. The upper and lower hinges extend to the largest and smallest values,
respectively but no further then 1.5XIQR from the hinge. P are Mann-Whitney U test. D, rho
and P are Spearman's correlations.

CMV seropositive adults have reduced induction of antibodies following controlled human

malaria infection

To assess the antibodies that were contributing to reduced antibody score in CMV positive

individuals, IgM, IgG (total and subclasses) and functional antibodies targeting the merozoite

were analysed. In CMV positive individuals, IgG to the merozoite was lower at end of study

(EOS, Day 27-36), but there were no differences in the magnitude of induced IgM (Fig. 2 A).

The reduced IgG in CMV positive individuals was driven by a reduced IgG1 response. The

magnitude of IgG2 and IgG3 were also lower, however differences were not statistically

significant (Fig. 2 B). IgG1 is a cytophilic antibody which has important functional capacity

to fix complement and interact with Fc receptors on phagocytes. These functional antibody

responses have essential roles in immunity to malaria and can target the merozoite to block

invasion of the red blood cell and mediate protection [11,33,34]. Consistent with reduced

IgG1 induction in CMV positive individuals, CMV positive individuals had reduced

induction of antibodies that could fix complement (measured by C1q fixation, the first step in

the classical complement cascade [33]), and reduced binding of FcγRII and FcgγIII, which

are involved in phagocytosis of parasites by neutrophils and other immune cells [35] (Fig. 2

C/D). However, antibodies that could mediate opsonic phagocytosis by the THP-1 pro-

monocytic cell line, which primarily involves FcγRI [35], did not differ between the two

groups (Fig. 2 E). There was no difference in the antibody response to MSP2 (Supplementary

Fig. 1). Together these data show that CMV is an important modulator of the primary

immune response to malaria infection.

Figure 2. **IgG1 and functional antibodies to the merozoite induced by controlled human**
malaria infection are reduced in CMV infected adults
*(**A**) IgG and IgM antibodies, (**B**) IgG subclass antibodies and levels of functional antibodies*
*that can (**C**) fix C1q complement component, (**D**) cross link FcγRII and FcγRIII, and (**E**)*
drive opsonic phagocytosis by THP1 cells, targeting merozoite stage parasites, stratified by
CMV serostatus. CMV- white bars (n=19), CMV+ grey bars (n=21), data is Tukey boxplots
with the median, 25th and 75th percentiles. The upper and lower hinges extend to the largest
and smallest values, respectively but no further then 1.5XIQR from the hinge. P are Mann-
Whitney U test. See also Supplementary Figure 1.
189

CMV seropositive adults have expansion of Tfh1 cells, and reduced proportions of

activated Tfh2 cells during infection

We have previously shown that in this cohort of CHMI participants, early activation of Tfh2

cells was associated with anti-parasitic antibody induction [17]. In contrast, activation of

Tfh1 cells was not associated with antibody levels, but was associated with increased

antibody secreting cell development which may impair germinal centres by acting as a

nutrient sink during infection [18]. To assess if CMV seropositivity impacted Tfh cell

differentiation either before or during CHMI, we analysed the same data set, and stratified by

CMV serostatus. Tfh cells were defined as all CXCR5$^+$ cells, subsets identified based on

CXCR3 and CCR6 expression, and activation measured by expression of PD1, CD38 and

ICOS. Non-Tfh effector CD4$^+$ cells (CXCR5$^-$ CD4$^+$ T cells) were also analysed

(Supplementary Fig. 2 A). Within the Tfh cell population, CMV seropositive individuals had

a significantly higher proportion of Tfh1 cells, and a significantly decreased proportion of

Tfh2 cells, before and during CHMI (Fig. 3 B). This expansion towards Th1-like cells was

not seen in non-Tfh effectors, suggesting that the expansion of Tfh1 cells in CMV

seropositive individuals was not due to a systemic inflammatory phenotype (Supplementary

Fig. 2 C). Importantly, amongst activated Tfh cells (analysed as either PD1$^+$, ICOS$^+$ or

CD38$^+$ Tfh cells), there was a significantly increased proportion of Tfh1 cells (PD1$^+$) and a

reduced proportion of Tfh2 cells (ICOS$^+$ and CD38$^+$) at day 14/15 (Fig. 3 C). Within the

CD4[+] T cell population, there was no difference in the proportion of Tfh cells, nor any major

differences in CD4[+] effector (CXCR5-) Th1-, Th2- and Th17- like populations between

CMV infected and uninfected individuals (Supplementary Fig. 2 B-C). Further, there was no

difference in magnitude of activation in Tfh cells, nor non-Tfh effector cells during CHMI,

aside from an increase in Th2 cell activation at day 14 post-inoculation (Fig. 3 A,

Supplementary Fig. 2 D-E). While previous studies have shown that CMV antigen specific

Tfh cells are dominated by Tfh1 cell subsets [36], these findings show that latent CMV

infection modulates the entire Tfh cell compartment.

*Figure 3. **Tfh cell response is skewed to Tfh1 subsets in CMV seropositive adults***
*(**A**) Activation of Tfh cells as measured by PD1+, ICOS+, and CD38+ cells (% of Tfh CD4 T*
*cells) in CHMI stratified by CMV status. (**B**) Tfh cell subsets (% of Tfh CXCR5+ CD4 T*
*cells), stratified by CMV status. (**C**) Tfh cell subsets as a proportion of activated Tfh cells,*
stratified by CMV serostatus at day 14/15. CMV- white bars (n=19 for day 0, 8, 15, n=17 for
day EOS), CMV+ grey bars (n=21 for day 0, 8, 15 and n=20 for EOS), data is Tukey
boxplots with the median, 25th and 75th percentiles. The upper and lower hinges extend to the
largest and smallest values, respectively but no further then 1.5XIQR from the hinge. P are
Mann-Whitney U test. See also Supplementary Figure 2.
229

The relationship between Tfh cell subsets and antibody induction is modulated by CMV

serostatus

Different subsets of Tfh cells have specific functions within the germinal centre to drive B

cell activation [15]. To assess if the CMV mediated bias of the Tfh cell compartment towards

Tfh1 cells may impact antibody development in malaria, we assessed the correlations

between Tfh cell subsets and antibody score in our cohort, stratified by CMV serostatus. The

activation of Tfh2 cells, previously shown to be associated with antibody induction [17], was

only associated with antibody score in CMV seronegative individuals (Fig. 4 A). Further,

amongst CMV seropositive individuals, the proportion of Tfh1 cells in the Tfh cell

compartment negatively correlated with antibody score (Fig. 4 B). Together, these data show

that latent CMV infection modulates the structure of the Tfh cell compartment, which

negatively impacts antibody production during malaria.

Figure 4. **Correlation between Tfh cell subsets and antibody development based on CMV serostatus**
(A) Correlation between activated Tfh2 cells (ICOS+ Tfh2 (% of Tfh CXCR5+ CD4T cells)
and **(B)** the proportion of Tfh1 (% of Tfh CXCR5+ CD4 T cells) at day 0, 8, 14/15, EOS and
Antibody score measured at EOS. CMV seronegative green (n=19 for day 0, 8, 15, n=17 for
day EOS), CMV seropositive orange (n=21 for day 0, 8, 15 and n=20 for EOS). Spearman's
Rho and p are indicated.

Latent CMV infection reduces antibody response to malaria vaccination

To assess if latent CMV infection also affects malaria vaccine responses, we analysed

antibody levels in 24 individuals vaccinated with full-length MSP1 formulated with GLA-SE

in a Phase 1a clinical trial [30,31]. Within this study 12 individuals were CMV seronegative

and 12 were CMV seropositive. Sex, age and EBV status were comparable between CMV

seropositive and seronegative individuals (Supplementary Table 1). We analysed induced

MSP1 antibodies at day 85 following vaccination, which has been shown to be when

antibody response peak [30,31]. IgG and IgM to MSP1, C1q fixation, opsonic phagocytosis

by THP1 and neutrophils, and neutrophil antibody dependent respiratory burst (ADRB)

responses were assessed. Additionally, IgG subclasses and NK IFNγ production and

degranulation (measured by CD107α expression), were available for 11-13 participants.

Unbiased clustering of all antibody responses aggregated the cohort into two clusters, low

responders who were 8/12 (66%) CMV seropositive, and high responders who were 4/12

(33%) CMV seropositive, consistent with a negative impact of latent CMV infection on

vaccine induced antibodies (Fig. 5A). To evaluate whether there was a difference in the

antibody response between CMV positive and negative individuals, we used a MANOVA

test which includes all antibody data in a single analysis. This found that there was a

significant difference between CMV serostatus groups based on antibody parameters

(MANOVA p=0.027, Fig. 5B). MSP1 targeted IgG, IgM, C1q and neutrophil ADRB were all

reduced in CMV seropositive individuals. Differences were most pronounced for C1q

fixation (Wilcox p=0.045, Fig. 5B). For antibody responses where only partial data was

available, there was reduced IgG1, IgG3, IgG4, along with reduced NK IFNγ and

degranulation in CMV seropositive individuals (Fig. 5C).

To assess if CMV mediated cellular changes contributed to the reduced MSP1 antibody

induction, we analysed the cellular compartment in vaccinated individuals at time of

vaccination (day 0) and peak antibody response (day 85) (Supplementary Figure 3A). Unlike

in the CHMI cohort, we did not detect any significant differences in subset distribution based

on CMV infection status within Tfh and non-Tfh CD4$^+$ T cells (Supplementary Figure 3B).

Previous studies have reported CMV to be associated with down regulation of co-stimulatory

marker CD28, and upregulation of terminally differentiation marker CD57 [25], both of

which were included in cellular analysis. Consistent with previous studies, CMV infected

individuals had increased frequencies of CD28$^-$ CD4$^+$ T cells, and increased frequencies of

CD57$^+$ CD4$^+$ T cells within our cohort, both at day 0 and day 85 (Fig. 5D). These differences

were significant within CXCR5$^-$ CD4$^+$ T cells, but not within the Tfh cell compartment

(Supplementary Figure 3C). The frequency of CD57$^+$ CD4 T cells at both day 0 and day 85

was negatively associated with the levels of C1q fixation, consistent with a role of CMV

mediated terminally differentiation in reduced antibody responses to vaccination. Taken

together, the data shows that latent CMV infection can modulate both the Tfh and non-Tfh

CD4$^+$ T cell compartments and that these changes negatively impact antibody production

during malaria or following vaccination.

*Figure 5. **Impact of CMV serostatus on antibody responses to MSP1 vaccination.***
*(**A**) Clustering of individuals within vaccination cohort based on MSP1 vaccine inducted*
antibodies. Antibody responses were normalised and z-scores to IgG, IgM and IgG1, 2, 3,
and 4 subclasses, along with C1q complement fixation, opsonic phagocytosis (OPA) from
THP1 cells and neutrophils, neutrophil antibody dependent respiratory burst (ADRB), and
NK production of IFNγ and degranulation (CD107α). Individuals aggregated into low and
*high responders. CMV status is shown (n=24). (**B**) Impact of CMV on antibody responses for*
IgG, IgM, C1q, THP1 OPA, neutrophil OPA and neutrophil ADRB (n=24). Manova p is
*shown. (**C**) IgG subclasses, and NK responses between CMV negative and positive*
*individuals (n=11-13). (**D**) Proportion of CD28- and CD57+ non-Tfh CD4 T cells (CXCR5-)*
*at day 0 and day 85 stratified by CMV infection status. (**E**) Correlation between frequency of*

CD57+ non-Tfh CD4 T cells at day 0 and day 85, and C1q antibodies at day 85 following
vaccination. Spearman's Rho and p are indicated. See also Supplementary Figure 3.
308

Discussion

Here, we show that CMV serostatus has a substantial negative impact on antibody

development during malaria infection and following vaccination. Within our CHMI cohort,

CMV seropositive individuals had a bias in Tfh cell compartment towards Tfh1 cells, with

the proportion of Tfh1 cells being negatively associated with antibody development. In our

vaccination cohort, CMV seropositive individuals had increased frequencies of CD28$^-$ and

CD57$^+$ CD4$^+$ T cells, which was negatively associated with antibody development. These

findings have significant implications for immune development during malaria via infection

or vaccination in endemic areas, where there is typically a very high CMV seroprevalence

[28,32].

CHMI studies are a powerful platform to evaluate mechanisms of immunity to malaria. In

previous studies using this model, we have identified that expression of specific miRNAs

prior to CHMI can predict individual control of parasite growth and antibody development

following drug treatment [40], suggesting that preexisting host factors have major roles in the

immune response to malaria. Here, we show that the protective role of Tfh2 cells in antibody

induction [17] is significantly modulated by CMV serostatus, thus identifying a host factor

underpinning immune heterogeneity. The impact of CMV on antibody induction was linked

to a bias towards Tfh1 cells within the Tfh cell compartment. This link between CMV and

modified Tfh cell compartment may explain previous reports that indicate that the impact of

CMV on immune development in response to vaccination and infection is highly pathogen

and/or context dependent. For example, while the impact of CMV serostatus on antibody

induction to influenza vaccination is well studied, the findings are highly variable. Some

studies have reported a negative impact of CMV seropositivity [41–48], others no impact

[44,49–51], and others reporting a positive impact of CMV on antibody induction [52–54].

The impact of CMV on immune induction in the context of infection is also mixed, with

antibodies induced by H1N1 influenza infection higher in CMV positive individuals [55], but

no CMV mediated differences in antibodies induced following SARS-CoV-2 infection [56].

The role of Tfh cells in antibody induction is pathogen and context dependent [15]. In

malaria, where Tfh2 cells have been associated with antibody development in response to

infection [17], including in children in endemic areas [20], or after vaccination with both the

licenced RTS,S [37] and experimental blood stage malaria vaccines [38,39], latent CMV

infection may inhibit antibody induction. In contrast, following influenza vaccination, where

Tfh1 cells are consistently positively associated with antibody induction [57,58], CMV

seropositive individual may have enhance antibody development, as seen in some studies

[52–54].

Highlighting the important role of our findings with CHMI to vaccine development for

malaria, we also show that antibodies induced by full-length MSP1 vaccination are reduced

in CMV seropositive individuals within a Phase 1 a clinical study. Despite the limited cohort

size within this study, significantly reduced induction of complement fixing antibodies was

shown in CMV seropositive individuals. While within the vaccine cohort we did not detected

differences in the phenotype of Tfh cells, instead reduced vaccine induced antibodies were

associated with increased frequencies of $CD57^+$ $CD4^+$ T cells, a cell subset thought to be

terminally differentiated. The absences of detectible CMV mediated differences in the Tfh

compartment may be due to the small cohort size, technical differences in Tfh subset staining,

or may reflect genetic, environmental and past infection distinct between the CHMI and

vaccine cohorts. Nevertheless, our findings are consistent with a recent study of immune

responses induced by Ebola vaccination in UK and Senegalese participants [29]. In that study

latent CMV infection was also associated with the reduced induction of antibodies following

vaccination, and similarly changes to CD28 and CD57 expression within the T cell

compartment were reported. Differences in CMV seroprevalence between recipients of this

Ebola vaccine in the UK (50% CMV positive) and Senegal (100% CMV positive) accounted

for the previously reported reduced vaccine responsiveness in the Senegalese cohort, with

CMV infected UK participants having comparable responses to Senegalese individuals [59].

These differences in vaccine responsiveness in UK compared to Senegalese participants

mimic differences in responsiveness to malaria vaccines seen between cohorts in the US and

malaria endemic countries [6–10]. As such, studies to investigate the role of latent CMV

infection on malaria vaccine responsiveness are required, particularly in target populations

were CMV infection occurs early in life.

Concluding remarks

Taken together, these data identify that latent CMV infection is an important driver of

heterogeneity in the host immune response to malaria, with a striking negative effect on the

development of malaria-specific antibodies induced by infection or vaccination. During

malaria infection, this reduced antibody induction was linked to the skewing of Tfh cells to

the Tfh1 cell subsets, and away from Tfh2 cells, which have previously been linked to

antibody induction in CHMI [17]. In vaccination, reduced antibodies were associated with

increased frequencies of $CD57^+$ $CD4^+$ T cells, highlighting a potential role of CMV mediated

immune senescence in reduced antibody induction. These findings have important

implications for understanding heterogeneity in immune responses to malaria infection and

vaccination, particularly as CMV is typically highly prevalent and acquired in early life in

areas of malaria transmission [28,32]. Further studies are required to assess if latent CMV

infection contributes to the slow acquisition of protective functional antibodies in children in

malaria endemic areas. Knowledge of the major impact of CMV on malaria immune

responses may open new avenues for malaria control, such as targeting CMV infection or the

immune changes induced by CMV infection to improve immunity to malaria and thus

protection.

Materials and Methods

Study populations

Written informed consent was obtained from all participants. Ethics approval for the use of

human samples in the relevant studies was obtained from the Alfred Human Research and

Ethics Committee for the Burnet Institute (#225/19), the Human Research and Ethics

Committee of the QIMR-Berghofer Medical Research Institute (P1479, P3444 and P3445)

and the Ethics Committee of the Medical Faculty of Heidelberg (AFmo-538/2016) and the

relevant regulatory authority (Paul Ehrlich Institute, Langen, Germany)

Controlled human malaria infection (CHMI) studies were performed as previously described

using the induced blood stage malaria model [60]. Malaria naïve individuals were inoculated

by intravenous injection of 2800 *P. falciparum* infected red blood cells and monitored for

parasite growth with qPCR [61]. Here, blood samples were collected at baseline (day 0), day

of treatment (day 8) and at 14 or 15 days and at end of study (EOS), 27-36 days after

inoculation (4 studies, across 6 independent infection cohorts). Clinical trials were registered

at ClinicalTrials.gov NCT02867059 [62], NCT02783833[63] , NCT02431637 [64],

NCT02431650 [64]. Whole blood was stained and analysed immediately by flow cytometry

and plasma was collected from lithium heparin collection tubes.

Full-length MSP1 vaccination was performed as previously described in a Phase 1a clinical

trial of full-length MSP1 adjuvanted with GLA-SE [30,31] registered with EudraCT (No.

2016-002463-33. Volunteers received three vaccinations at intervals of 29±3 days at 25, 50

or 150μg. For the current study, antibody responses at day 85 following first vaccination, and

cellular responses at day 0 and day 85 were considered. Antibody responses were measured

from sera collected in S-Monovetter serum-gel, and PMBCs were collected from sodium-

citrate tubes.

CMV and EBV serostatus

For CHMI cohorts CMV and EBV seroprevalence was assessed using baseline plasma

samples by commercially available ELISA kits (ab108724 and ab108730), according to

manufacturer's instructions. CMV ELISA kit has a reported sensitivity and specificity of

98% and 97.5% respectively. For the vaccination cohort, CMV seroprevalence was assessed

using day 0 plasma samples by commercially available CMV IgG ELISA (EUROIMMUN EI

2570-9601 G). One individual fell into a CMV intermediate status in the ELISA. For this

sample, an immunblot (MIKROGEN DIANOSTIK *recom*Line CMV IgG 5572) was

performed which was positive. The EBV seroprevalence was established using the Anti-EBV

CA_ELISA (IgG) Assay (EUROIMMUN EI 2791-9601 G).

Induced antibody responses

For CHMI studies, antibody response to intact merozoites and merozoite surface antigen

MSP2 were quantified as previously described [17]. To isolate merozoites, *P. falciparum*

3D7 parasites were maintained in continuous culture in RPMI-HEPES medium supplemented

with hypoxanthine(370 mM), gentamicin (30 mg/ml), 25 mM sodium bicarbonate and 0.25%

AlbuMAX II (GIBCO) or 5% heat-inactivated human sera in O+ RBCs from malaria-naive

donors (Australian Red Cross blood bank). Cultures were incubated at 37°C in 1% O_2, 5%

CO_2, 94% N_2 and schizont stage parasites were purified by MACS separation (Miltenyl

Biotec). To isolate merozoites, magnet purified schizonts were incubation with the protease

inhibitor E64 (10 mg/ml), and following complete development, merozoites were isolated by

membrane filtration (1.2 mm). 50 ml of *P. falciparum* 3D7 merozoites (2.5×10^5

merozoites/ml) or 50 ml of 0.5 mg/ml MSP2 recombinant antigen [65] in PBS were coated to

96-well flat bottom MaxiSorb plates (Nunc) overnight at 4°C. Plates were blocked with 10%

skim milk for merozoites, or 1% casein (Sigma-Aldrich) for MSP2 for 2 hours at 37°C.

Plasma was diluted in 0.1% casein in PBS1/100 for IgG, 1/250 for IgG subclasses and IgM,

1/100 for C1q, 1/100 for FcγR targeting merozoites or 1/50 for FcγR targeting MSP2) and

incubated for 2 hours at room temperature. For total antigen specific IgG detection, plates

were incubated with goat polyclonal

anti-human IgG HRP- conjugate (1/1000; Thermo Fisher Scientific) for 1 hour at room

temperature. For detection of IgG subclasses and IgM, plates were incubated with a mouse

anti-human IgG1 (clone HP6069), mouse anti-human IgG3 (HP6050) or mouse anti-human

IgM (clone HP6083) at 1/1000 (Thermo Fisher Scientific) for 1 hour at room temperature.

This was followed by detection with a goat polyclonal anti-mouse IgG HRP-conjugate

(1/1000; Millipore). For all ELISAs, plates were washed three times with PBS (for merozoite

ELISAs) or PBS-Tween 0.05% (for MSP2) between antibody incubation steps. For detection

of complement fixing antibodies, following incubation with human sera, plates were

incubated with purified C1q (10 mg/ml; Millipore) as a complement source, for 30 min at

room temperature. C1q fixation was detected with rabbit anti- C1q antibodies (1/2000; in-

house) and a goat anti-rabbit-HRP (1/2500; Millipore). For FcγR assays, 100ul of biotin-

conjugated rsFcγRIIa H131 or rsFcγRIIIa V158 ectodomain dimer (0.2ug/ml) was incubated

at 37°C for 1 hour followed by 3 washes with PBS-Tween. The binding was detected with

horseradish peroxidase (HRP)-conjugated streptavidin antibody (1:10,000) in PBS-BSA at

37°C for 1 hour.

TMB liquid substrate (Life Technologies) was added for 1 hour at room temperature and the

reaction was stopped using 1M sulfuric acid. The optical density (OD) was read at 450 nm.

For opsonic phagocytosis, THP1 cells were incubated with intact merozites (stained with

Ethidium Bromide and opsonised with plasma diluted 1/100) or latex beads coated with

MSP2 (opsonised with plasma diluted 1/10) for 20 min at 37°C and cells washed with FACS

buffer. The proportion of THP-1 cells containing fluorescent-positive beads was evaluated by

flow cytometry (FACS CantoII, BD Biosciences), analyzed using FlowJo software and

presented as phagocytosis index (the percentage of THP-1 monocytes with ingested

merozoites or beads).

To calculate antibody score antibody responses below positive cut-off threshold were set as

negative, and remaining positive responses were used to calculate median and used to

categorise responses into low (below median) and high (above median) responses. Antibody

score was calculated by giving categories zero/low/high a numerical score of 0/1/2 and then

summing across all antibody responses.

For full-length MSP1 vaccination studies, antibodies were measured as previously reported

[30,31]. Total IgG and IgM antibody levels were determined by ELISA using MaxiSorp

plates (Thermo Fisher Scientific) coated with 100nM recombinant full-length MSP1

(Glycotope Biotechnology GmbH, Heidelberg). Sera were titrated in two-fold dilutions and

incubated for 2h. A secondary antibody goat anti-human IgG and IgM alkaline phosphatase

conjugate (Sigma-Aldrich) was used at a dilution of 1:20,000 for 1h. The substrate p-

nitrophenyl-phosphate was added and incubated for 1h in the dark, and stopped with 1M

NaOH. The absorbance at 405 nm was determined using the plate reader Cytation 3

(BioTek). For the IgG subtype ELISA, the same protocol was applied, with the exception that

subclass-specific peroxidase-conjugated secondary antibodies were used (The Binding Site

GmbH). The substrate 1-step turbo TMB (Thermo Fisher) was added for 20 min in the dark,

then stopped by adding 1 M HCl. Optical density was recorded at 450nm. For the C1q

fixation ELISA assay, full-length MSP1 coated plates were also incubated overnight at 4°C.

The next day, the plates were washed four times with 1x PBS containing 0.05% Tween 20

and blocked with 1% casein/PBS at 37°C for 2 h. After washing, the plates were incubated

with 50µl of purified IgG at 1mg/ml blocking buffer, respectively at 37°C for 1 h. After

incubation, the plates were washed and recombinant C1q (Abcam) at 10µg/ml in blocking

buffer was added for 30 min at 37°C. To detect C1q binding, anti-C1q horse radish

peroxidase (HRP)-conjugated secondary antibodies (Abcam) was added at a 1:100 dilution in

blocking buffer for 1h at 37°C. Then SigmaFAST OPD (Sigma-Aldrich) was added for 30

min to 1h in the dark at room temperature for development. The reaction was stopped with

1M HCL and the signal intensity was measured at 492nm using the Biotek Cytation 3 plate.

For the Antibody-dependent respiratory burst (ADRB) assay full length MSP1 protein was

coated at 100nM in PBS in opaque 96-well Lumitrac microplates (Greiner Bio-One), blocked

with 1% casein/PBS and incubated for 1h at 37°C with 50µ/well of IgG at 1mg/ml. The

plates were washed with PBS and 50µl/well of luminol at 0.04mg/ml were added to each

well.   Next, freshly isolated neutrophils were added at 10^7 cells/ml and absorption was

immediately read at 450nm using the Biotek Cytation 3 reader for every 2 min over 1 h. For

the Opsonic phagocytosis activity assay, full-length MSP1 coupled to fluorescent beads were

performed as previously reported [30,31]   using THP1 cells and neutrophils). Briefly, an

antigen coupled bead suspension was added to 96-well to U-bottomed plates and opsonized

for 1h at 37°C with 50µl/well of purified IgG. The plates were centrifuged at 2000x g for 7

min and washed with PBS. Opsonized beads were resuspended in culture medium (RPMI

1640 media with 2mM L-glutamine, 10% FCS and 1% penicillin-streptomycin) before

incubation with 5 x 10^4 phagocytes at 37°C. Phagocytosis was arrested by centrifugation at

1200 rpm for 7 min at 4°C and washing with ice-cold FACS buffer (0.5% BSA and 2mM

EDTA in PBS). Cells were resuspended in 2% formaldehyde/PBS and the proportion of

phagocytes containing PE fluorescent beads was on a FACS Canto II (BD biosciences). NK

activity assays, full-length MSP1-coated plates were washed PBS and blocked with 1%

casein/PBS for 4 h at 37°C. The plates were washed and incubated for 2h at 37°C with IgG at

1mg/ml. Then 5 x 10^4 freshly isolated human NK cells together with anti-human CD107a PE

(BD biosciences,1:70), brefeldin A (Sigma-Aldrich, 1:200) and monensin (Sigma, 1:200) was

added for 18 h at 37°C. Afterwards, the cells were transferred to 96-well V-bottomed plates,

centrifuged at 1500 rpm for min at 4°C and washed with ice-cold FACS buffer. Cell viability

was determined by adding dye eFluor™520 (Thermo Fischer). NK cell surface markers were

labelled with an antibody cocktail of anti-CD56 APC (BD biosciences, 1:17) and anti-CD3

PE-Cy5 (BD biosciences, 1:33) for 30 min at 4°C in the dark. After washing, NK cells were

fixed with CellFIX (BD biosciences) at 4°C and permeabilized with permwash (BD

biosciences) for 10 min at 4°C. Intracellular IFN-γ was measured by adding anti- IFN-γ PE-

Cy7 (BD biosciences, 1:33) for 1h at 4°C. NK cells were washed with permwash,

resuspended in FACS buffer and Ab-NK activity (proportion of NK cells with CD107a

and/or IFN-γ staining was assessed by a FACS ContoII (BD biosciences).

Quantification of T cells

For CHMI studies, whole blood was analysed as described previously [17]. 200 mL of whole

blood were stained with the following conjugated antibodies, all from BD Biosciences; anti-

CD20-BUV395 (2H7, 1:150, Cat# 563782), anti-CXCR5-BV421 (RF8B2, 1:50,

Cat#562747), anti-CD4 (V500, 1:30, Cat# 560768), anti-CCR6-BV650 (11A9, 1:200,

Cat#563922), anti-CD38-BV786 (HIT2, 1:400, Cat#563964), anti-CXCR3-APC (1C6, 1:25,

Cat#550967), anti-CD27-APC-R700 (M-T271, 1:100, Cat#565116), anti-CD8-APC-Cy7

(SK1, 1:150, Cat#557834), anti-CD19-FITC (HIB19, 1:20, Cat#55412), anti-CD45-PerCP-

Cy5.5 (2D1, 1:50, Cat#340953), anti-ICOS-PE (DX29, 1:10, Cat#557802), anti-CD3-PE-

CF594 (UCHT1, 1:600, Cat#562280), anti-PD1-PE-Cy7 (EH12.1, 1:100, Cat#561272).

RBCs were lysed with FACS lysing solution (BD) and re-suspended in 2% FBS/PBS.

Samples were acquired on the BD LSR Fortessa TM 5 laser cytometer (BD Biosciences).

These data were analyzed using FlowJo version 10.8 software (Tree Star, San Carlos, CA,

USA).

For full-length MSP1 vaccine cohorts, cellular responses were analysed as previously

reported [31]. The phenotyping of T- and B-lymphocytes was conducted in whole blood

samples within 24 hours. To monitor the dynamics of lymphocyte subpopulations,

comprehensive T-cell and B-cell panels were employed, based on modified recommendations

from the "Human Immunophenotyping Consortium" [66].

Data availability statement

The data underlying all figures are available in the published article and its online

supplemental material.

Funding and financial conflicts of interest

Authors declare no conflicts of interest.

RTL is employee from Sumaya GmbH & Co. KG.

Acknowledgements

RBC and human serum were provided by the Australian Red Cross Blood Bank (Melbourne).

We thank the participants involved in the controlled human malaria infection studies and all

study clinicians and support staff at QPharm and Medicine for Malaria Venture for funding

the CHMI studies. This work was supported by the National Health and Medical Research

Council of Australia (program grant 1132975 to J.S.M. and C.R.E.; Practitioner Fellowship

1135955 to J.S.M., Senior Research Fellowship 1154265 to C.R.E., Career Development

Award 1141278, Project Grant 1125656, and Ideas Grant 1181932 to M.J.B.; Program Grant

290208, Senior Research Fellowship 1077636 to J.G.B.; Emerging Leadership 2 Fellowship

to BEB, Australian Centre for Research Excellence in Malaria Elimination 1134989 to J.G.B.

and J.S.M.); the Jim and Margaret Beever Fellowship to J.A.C; the CSL Centenary

Fellowship to M.J.B, and the Snow Medical Foundation Fellowship to M.J.B. The Burnet

Institute is supported by the NHMRC for Independent Research Institutes Infrastructure

Support Scheme and the Victorian State Government Operational Infrastructure Support.

Special thanks to the volunteers that participated in the first-in-human phase 1a clinical trial

with full-length MSP1 in Heidelberg and to all the team at the Parasitology Unit as well as

the Clinical Pharmacology and Pharmacoepidemiology Unit at the University Hospital

Heidelberg, for their enthusiastic work in running the trial and ulterior analysis of samples.

This work was funded by Sumaya Biotech GmbH & Co. KG, Heidelberg, Germany, via a

venture loan provided by the EU Malaria Fund. We are very thankful to the entire team at

Sumaya-Biotech for constant support.

References:

[1] Geneva: World Health Organisation. World malaria report 2023. 2023.

[2] Holla P, Bhardwaj J, Tran TM. Mature beyond their years: young children who escape detection of parasitemia despite living in settings of intense malaria transmission. Biochem Soc Trans 2024. https://doi.org/10.1042/bst20230401.

[3] White MT, Verity R, Griffin JT, Asante KP, Owusu-Agyei S, Greenwood B, et al. Immunogenicity of the RTS,S/AS01 malaria vaccine and implications for duration of vaccine efficacy: secondary analysis of data from a phase 3 randomised controlled trial. The Lancet Infectious Diseases 2015;15:1450–8. https://doi.org/10.1016/s1473-3099(15)00239-x.

[4] Bejon P, White MT, Olotu A, Bojang K, Lusingu JP, Salim N, et al. Efficacy of RTS,S malaria vaccines: individual-participant pooled analysis of phase 2 data. Lancet Infect Dis 2013;13:319–27. https://doi.org/10.1016/s1473-3099(13)70005-7.

[5] van Dorst MMAR, Pyuza JJ, Nkurunungi G, Kullaya VI, Smits HH, Hogendoorn PCW, et al. Immunological factors linked to geographical variation in vaccine responses. Nat Rev Immunol 2024;24:250–63. https://doi.org/10.1038/s41577-023-00941-2.

[6] Jongo SA, Shekalaghe SA, Church LWP, Ruben AJ, Schindler T, Zenklusen I, et al. Safety, Immunogenicity, and Protective Efficacy against Controlled Human Malaria Infection of Plasmodium falciparum Sporozoite Vaccine in Tanzanian Adults. The American Journal of Tropical Medicine and Hygiene 2018;99:338–49. https://doi.org/10.4269/ajtmh.17-1014.

[7] Stoute JA, Slaoui M, Heppner DG, Momin P, Kester KE, Desmons P, et al. A Preliminary Evaluation of a Recombinant Circumsporozoite Protein Vaccine against Plasmodium falciparum Malaria. N Engl J Med 1997;336:86–91. https://doi.org/10.1056/nejm199701093360202.

[8] Kester KE, McKinney DA, Tornieporth N, Ockenhouse CF, Heppner DG, Hall T, et al. Efficacy of Recombinant Circumsporozoite Protein Vaccine Regimens against Experimental Plasmodium falciparum Malaria. J Infect Dis 2001;183:640–7. https://doi.org/10.1086/318534.

[9] Kester KE, McKinney DA, Tornieporth N, Ockenhouse CF, Heppner DG, Hall T, et al. A phase I/IIa safety, immunogenicity, and efficacy bridging randomized study of a two-dose regimen of liquid and lyophilized formulations of the candidate malaria vaccine RTS,S/AS02A in malaria-naïve adults. Vaccine 2007;25:5359–66. https://doi.org/10.1016/j.vaccine.2007.05.005.

[10] Polhemus ME, Remich SA, Ogutu BR, Waitumbi JN, Otieno L, Apollo S, et al. Evaluation of RTS,S/AS02A and RTS,S/AS01B in Adults in a High Malaria Transmission Area. PLoS ONE 2009;4:e6465. https://doi.org/10.1371/journal.pone.0006465.

[11] Opi DH, Kurtovic L, Chan J-A, Horton JL, Feng G, Beeson JG. Multi-functional
antibody profiling for malaria vaccine development and evaluation. Expert Rev Vaccines
2021;20:1257–72. https://doi.org/10.1080/14760584.2021.1981864.

[12] Nkumama IN, Ogwang R, Odera D, Musasia F, Mwai K, Nyamako L, et al. Breadth of
Fc-mediated effector function correlates with clinical immunity following human malaria
challenge. Immunity 2024. https://doi.org/10.1016/j.immuni.2024.05.001.

[13] Crotty S. T Follicular Helper Cell Biology: A Decade of Discovery and Diseases.
Immunity 2019;50:1132–48. https://doi.org/10.1016/j.immuni.2019.04.011.

[14] Soon MSF, Nalubega M, Boyle MJ. T-follicular helper cells in malaria infection and
roles in antibody induction. Oxf Open Immunol 2021;2:iqab008-.
https://doi.org/10.1093/oxfimm/iqab008.

[15] Olatunde AC, Hale JS, Lamb TJ. Cytokine-skewed Tfh cells: functional consequences
for B cell help. Trends Immunol 2021;42:536–50. https://doi.org/10.1016/j.it.2021.04.006.

[16] Engwerda CR, Minigo G, Amante FH, McCarthy JS. Experimentally induced blood
stage malaria infection as a tool for clinical research. Trends in Parasitology 2012;28:515–21.
https://doi.org/10.1016/j.pt.2012.09.001.

[17] Chan J-A, Loughland JR, Rivera F de L, SheelaNair A, Andrew DW, Dooley NL, et al.
Th2-like T Follicular Helper Cells Promote Functional Antibody Production during
Plasmodium falciparum Infection. Cell Reports Medicine 2020;1:100157.
https://doi.org/10.1016/j.xcrm.2020.100157.

[18] Vijay R, Guthmiller JJ, Sturtz AJ, Surette FA, Rogers KJ, Sompallae RR, et al.
Infection-induced plasmablasts are a nutrient sink that impairs humoral immunity to malaria.
Nat Immunol 2020;21:790–801. https://doi.org/10.1038/s41590-020-0678-5.

[19] Oyong DamianA, Loughland JessicaR, Soon MeganSF, Chan J-A, Andrew D, Wines
BD, et al. Adults with Plasmodium falciparum malaria have higher magnitude and quality of
circulating T-follicular helper cells compared to children. Ebiomedicine 2022;75:103784.
https://doi.org/10.1016/j.ebiom.2021.103784.

[20] Chan J-A, Loughland JR, Parte L de la, Okano S, Ssewanyana I, Nalubega M, et al. Age-
dependent changes in circulating Tfh cells influence development of functional malaria
antibodies in children. Nat Commun 2022;13:4159. https://doi.org/10.1038/s41467-022-
31880-6.

[21] Chidrawar S, Khan N, Wei W, McLarnon A, Smith N, Nayak L, et al.
Cytomegalovirus⬜seropositivity has a profound influence on the magnitude of major
lymphoid subsets within healthy individuals. Clin Exp Immunol 2009;155:423–32.
https://doi.org/10.1111/j.1365-2249.2008.03785.x.

[22] Brodin P, Jojic V, Gao T, Bhattacharya S, Angel CJL, Furman D, et al. Variation in the
Human Immune System Is Largely Driven by Non-Heritable Influences. Cell 2015;160:37–
47. https://doi.org/10.1016/j.cell.2014.12.020.

[23] Picarda G, Benedict CA. Cytomegalovirus: Shape-Shifting the Immune System. J
Immunol 2018;200:3881–9. https://doi.org/10.4049/jimmunol.1800171.

[24] Klenerman P, Oxenius A. T cell responses to cytomegalovirus. Nat Rev Immunol
2016;16:367–77. https://doi.org/10.1038/nri.2016.38.

[25] Pourgheysari B, Khan N, Best D, Bruton R, Nayak L, Moss PAH. The Cytomegalovirus-
Specific CD4 + T-Cell Response Expands with Age and Markedly Alters the CD4 + T-Cell
Repertoire. J Virol 2007;81:7759–65. https://doi.org/10.1128/jvi.01262-06.

[26] Moseley P, Klenerman P, Kadambari S. Indirect effects of cytomegalovirus infection:
Implications for vaccine development. Rev Méd Virol 2023;33:e2405.
https://doi.org/10.1002/rmv.2405.

[27] Zuhair M, Smit GSA, Wallis G, Jabbar F, Smith C, Devleesschauwer B, et al. Estimation
of the worldwide seroprevalence of cytomegalovirus: A systematic review and
meta☐analysis. Rev Méd Virol 2019;29:e2034. https://doi.org/10.1002/rmv.2034.

[28] Kaye S, Miles D, Antoine P, Burny W, Ojuola B, Kaye P, et al. Virological and
Immunological Correlates of Mother☐to☐Child Transmission of Cytomegalovirus in The
Gambia. J Infect Dis 2008;197:1307–14. https://doi.org/10.1086/586715.

[29] Bowyer G, Sharpe H, Venkatraman N, Ndiaye PB, Wade D, Brenner N, et al. Reduced
Ebola vaccine responses in CMV+ young adults is associated with expansion of
CD57+KLRG1+ T cells. J Exp Med 2020;217:e20200004.
https://doi.org/10.1084/jem.20200004.

[30] Blank A, Fürle K, Jäschke A, Mikus G, Lehmann M, Hüsing J, et al. Immunization with
full-length Plasmodium falciparum merozoite surface protein 1 is safe and elicits functional
cytophilic antibodies in a randomized first-in-human trial. Npj Vaccines 2020;5:10.
https://doi.org/10.1038/s41541-020-0160-2.

[31] Rosenkranz M, Fürle K, Hibbert J, Ulmer A, Ali A, Giese T, et al. Multifunctional
IgG/IgM antibodies and cellular cytotoxicity are elicited by the full-length MSP1
SumayaVac-1 malaria vaccine. Npj Vaccines 2023;8:112. https://doi.org/10.1038/s41541-
023-00701-2.

[32] Miles DJC, Sanneh M, Holder B, Crozier S, Nyamweya S, Touray ES, et al.
Cytomegalovirus infection induces T☐cell differentiation without impairing antigen☐specific
responses in Gambian infants. Immunology 2008;124:388–400.
https://doi.org/10.1111/j.1365-2567.2007.02787.x.

[33] Boyle MJ, Reiling L, Feng G, Langer C, Osier FH, Aspeling-Jones H, et al. Human
antibodies fix complement to inhibit Plasmodium falciparum invasion of erythrocytes and are
associated with protection against malaria. Immunity 2015;42:580–90.
https://doi.org/10.1016/j.immuni.2015.02.012.

[34] Osier FH, Feng G, Boyle MJ, Langer C, Zhou J, Richards JS, et al. Opsonic
phagocytosis of Plasmodium falciparum merozoites: mechanism in human immunity and a

correlate of protection against malaria. BMC Medicine 2014;12:108.
https://doi.org/10.1186/1741-7015-12-108.

[35] Feng G, Wines BD, Kurtovic L, Chan J-A, Boeuf P, Mollard V, et al. Mechanisms and targets of Fcγ-receptor mediated immunity to malaria sporozoites. Nat Commun 2021;12:1742. https://doi.org/10.1038/s41467-021-21998-4.

[36] Niessl J, Baxter AE, Morou A, Brunet-Ratnasingham E, Sannier G, Gendron-Lepage G, et al. Persistent expansion and Th1-like skewing of HIV-specific circulating T follicular helper cells during antiretroviral therapy. EBioMedicine 2020;54:102727. https://doi.org/10.1016/j.ebiom.2020.102727.

[37] Bowyer G, Grobbelaar A, Rampling T, Venkatraman N, Morelle D, Ballou RW, et al. CXCR3+ T Follicular Helper Cells Induced by Co-Administration of RTS,S/AS01B and Viral-Vectored Vaccines Are Associated With Reduced Immunogenicity and Efficacy Against Malaria. Front Immunol 2018;9:1660. https://doi.org/10.3389/fimmu.2018.01660.

[38] Nielsen CM, Ogbe A, Pedroza-Pacheco I, Doeleman SE, Chen Y, Silk SE, et al. Protein/AS01B vaccination elicits stronger, more Th2-skewed antigen-specific human T follicular helper cell responses than heterologous viral vectors. Cell Reports Medicine 2021:100207. https://doi.org/10.1016/j.xcrm.2021.100207.

[39] Minassian AM, Silk SE, Barrett JR, Nielsen CM, Miura K, Diouf A, et al. Reduced blood-stage malaria growth and immune correlates in humans following RH5 vaccination. Med 2021. https://doi.org/10.1016/j.medj.2021.03.014.

[40] Burel JG, Apte SH, Groves PL, Boyle MJ, Langer C, Beeson JG, et al. Dichotomous miR expression and immune responses following primary blood-stage malaria. JCI Insight 2017;2. https://doi.org/10.1172/jci.insight.93434.

[41] Turner JE, Campbell JP, Edwards KM, Howarth LJ, Pawelec G, Aldred S, et al. Rudimentary signs of immunosenescence in Cytomegalovirus-seropositive healthy young adults. Age 2014;36:287–97. https://doi.org/10.1007/s11357-013-9557-4.

[42] Frasca D, Diaz A, Romero M, Landin AM, Blomberg BB. Cytomegalovirus (CMV) seropositivity decreases B cell responses to the influenza vaccine. Vaccine 2015;33:1433–9. https://doi.org/10.1016/j.vaccine.2015.01.071.

[43] Trzonkowski P, Myśliwska J, Szmit E, Wieckiewicz J, Łukaszuk K, Brydak LB, et al. Association between cytomegalovirus infection, enhanced proinflammatory response and low level of anti-hemagglutinins during the anti-influenza vaccination—an impact of immunosenescence. Vaccine 2003;21:3826–36. https://doi.org/10.1016/s0264-410x(03)00309-8.

[44] Derhovanessian E, Theeten H, Hähnel K, Damme PV, Cools N, Pawelec G. Cytomegalovirus-associated accumulation of late-differentiated CD4 T-cells correlates with poor humoral response to influenza vaccination. Vaccine 2013;31:685–90. https://doi.org/10.1016/j.vaccine.2012.11.041.

[45] Reed RG, Greenberg RN, Segerstrom SC. Cytomegalovirus serostatus, inflammation,
and antibody response to influenza vaccination in older adults: The moderating effect of beta
blockade. Brain, Behav, Immun 2017;61:14–20. https://doi.org/10.1016/j.bbi.2016.09.025.

[46] Arias RA, Moro-García MA, Echeverría A, Solano-Jaurrieta JJ, Suárez-García FM,
López-Larrea C. Intensity of the Humoral Response to Cytomegalovirus Is Associated with
the Phenotypic and Functional Status of the Immune System. J Virol 2013;87:4486–95.
https://doi.org/10.1128/jvi.02425-12.

[47] Moro-García MA, Alonso-Arias R, López-Vázquez A, Suárez-García FM, Solano-
Jaurrieta JJ, Baltar J, et al. Relationship between functional ability in older people, immune
system status, and intensity of response to CMV. AGE 2012;34:479–95.
https://doi.org/10.1007/s11357-011-9240-6.

[48] Wald A, Selke S, Magaret A, Boeckh M. Impact of human cytomegalovirus (CMV)
infection on immune response to pandemic 2009 H1N1 influenza vaccine in healthy adults. J
Méd Virol 2013;85:1557–60. https://doi.org/10.1002/jmv.23642.

[49] Haq K, Fulop T, Tedder G, Gentleman B, Garneau H, Meneilly GS, et al.
Cytomegalovirus Seropositivity Predicts a Decline in the T Cell But Not the Antibody
Response to Influenza in Vaccinated Older Adults Independent of Type 2 Diabetes Status. J
Gerontol Ser A: Biomed Sci Méd Sci 2017;72:1163–70.
https://doi.org/10.1093/gerona/glw216.

[50] den Elzen WPJ, Vossen ACMT, Cools HJM, Westendorp RGJ, Kroes ACM, Gussekloo
J. Cytomegalovirus infection and responsiveness to influenza vaccination in elderly residents
of long-term care facilities. Vaccine 2011;29:4869–74.
https://doi.org/10.1016/j.vaccine.2011.03.086.

[51] Strindhall J, Ernerudh J, Mörner A, Waalen K, Löfgren S, Matussek A, et al. Humoral
response to influenza vaccination in relation to pre-vaccination antibody titres, vaccination
history, cytomegalovirus serostatus and CD4/CD8 ratio. Infect Dis 2016;48:436–42.
https://doi.org/10.3109/23744235.2015.1135252.

[52] Furman D, Jojic V, Sharma S, Shen-Orr SS, Angel CJL, Onengut-Gumuscu S, et al.
Cytomegalovirus infection enhances the immune response to influenza. Sci Transl Med
2015;7:281ra43-281ra43. https://doi.org/10.1126/scitranslmed.aaa2293.

[53] McElhaney JE, Garneau H, Camous X, Dupuis G, Pawelec G, Baehl S, et al. Predictors
of the antibody response to influenza vaccination in older adults with type 2 diabetes. BMJ
Open Diabetes Res Care 2015;3:e000140. https://doi.org/10.1136/bmjdrc-2015-000140.

[54] Berg SPH van den, Wong A, Hendriks M, Jacobi RHJ, Baarle D van, Beek J van.
Negative Effect of Age, but Not of Latent Cytomegalovirus Infection on the Antibody
Response to a Novel Influenza Vaccine Strain in Healthy Adults. Front Immunol 2018;9:82.
https://doi.org/10.3389/fimmu.2018.00082.

[55] Nielsen CM, White MJ, Bottomley C, Lusa C, Rodríguez-Galán A, Turner SEG, et al.
Impaired NK Cell Responses to Pertussis and H1N1 Influenza Vaccine Antigens in Human

Cytomegalovirus-Infected Individuals. J Immunol 2015;194:4657–67.
https://doi.org/10.4049/jimmunol.1403080.

[56] Freeman ML, Oyebanji OA, Moisi D, Payne M, Sheehan ML, Balazs AB, et al.
Association of Cytomegalovirus Serostatus With Severe Acute Respiratory Syndrome
Coronavirus 2 Vaccine Responsiveness in Nursing Home Residents and Healthcare Workers.
Open Forum Infect Dis 2023;10:ofad063. https://doi.org/10.1093/ofid/ofad063.

[57] Bentebibel S-E, Lopez S, Obermoser G, Schmitt N, Mueller C, Harrod C, et al.
Induction of ICOS+CXCR3+CXCR5+ TH Cells Correlates with Antibody Responses to
Influenza Vaccination. Sci Transl Med 2013;5:176ra32.
https://doi.org/10.1126/scitranslmed.3005191.

[58] Bentebibel S-E, Khurana S, Schmitt N, Kurup P, Mueller C, Obermoser G, et al.
ICOS+PD-1+CXCR3+ T follicular helper cells contribute to the generation of high-avidity
antibodies following influenza vaccination. Sci Rep 2016;6:26494.
https://doi.org/10.1038/srep26494.

[59] Venkatraman N, Ndiaye BP, Bowyer G, Wade D, Sridhar S, Wright D, et al. Safety and
Immunogenicity of a Heterologous Prime-Boost Ebola Virus Vaccine Regimen in Healthy
Adults in the United Kingdom and Senegal. J Infect Dis 2018;219:1187–97.
https://doi.org/10.1093/infdis/jiy639.

[60] McCarthy JS, Sekuloski S, Griffin PM, Elliott S, Douglas N, Peatey C, et al. A pilot
randomised trial of induced blood-stage Plasmodium falciparum infections in healthy
volunteers for testing efficacy of new antimalarial drugs. PLoS ONE 2011;6:e21914.
https://doi.org/10.1371/journal.pone.0021914.

[61] Rockett RJ, Tozer SJ, Peatey C, Bialasiewicz S, Whiley DM, Nissen MD, et al. A real-
time, quantitative PCR method using hydrolysis probes for the monitoring of Plasmodium
falciparum load in experimentally infected human volunteers. Malaria Journal 2011;10:48.
https://doi.org/10.1186/1475-2875-10-48.

[62] Gaur AH, McCarthy JS, Panetta JC, Dallas RH, Woodford J, Tang L, et al. Safety,
tolerability, pharmacokinetics, and antimalarial efficacy of a novel Plasmodium falciparum
ATP4 inhibitor SJ733: a first-in-human and induced blood-stage malaria phase 1a/b trial. The
Lancet Infectious Diseases 2020. https://doi.org/10.1016/s1473-3099(19)30611-5.

[63] McCarthy JS, Donini C, Chalon S, Woodford J, Marquart L, Collins KA, et al. A phase
1, placebo controlled, randomised, single ascending dose study and a volunteer infection
study to characterize the safety, pharmacokinetics and antimalarial activity of the
Plasmodium phosphatidylinositol 4-kinase inhibitor MMV390048. Clinical Infectious
Diseases 2020. https://doi.org/10.1093/cid/ciaa368.

[64] Collins KA, Wang CY, Adams M, Mitchell H, Rampton M, Elliott S, et al. A controlled
human malaria infection model enabling evaluation of transmission-blocking interventions.
The Journal of Clinical Investigation 2018;128:1551–62. https://doi.org/10.1172/jci98012.

[65] McCarthy JS, Marjason J, Elliott S, Fahey P, Bang G, Malkin E, et al. A phase 1 trial of
MSP2-C1, a blood-stage malaria vaccine containing 2 isoforms of MSP2 formulated with

Montanide® ISA 720. PLoS ONE 2011;6:e24413.
https://doi.org/10.1371/journal.pone.0024413.

[66] Maecker HT, McCoy JP, Nussenblatt R. Standardizing immunophenotyping for the
Human Immunology Project. Nat Rev Immunol 2012;12:191–200.
https://doi.org/10.1038/nri3158.

815