Rhizobial effector NopM ubiquitinates Nod factor receptor NFR5 and

promotes rhizobial infection in *Lotus japonicus*

Yanan Wang[1], Hanbin Bao[1], Yutao Lei[1], Lifa Yuan[1], Haoxing Li[1], Hui Zhu[1],

Dawei Xin[2], Christian Staehelin[3], and Yangrong Cao[1*]

[1] National Key Lab of Agricultural Microbiology, Hubei Hongshan Laboratory,

Huazhong Agricultural University, Wuhan, Hubei, 430070, China.

[2] Key Laboratory of Soybean Biology of Chinese Ministry of Education, Key

Laboratory of Soybean Biology and Breeding/Genetics of Chinese Agriculture

Ministry, Northeast Agricultural University, Harbin, Heilongjiang, 150038, China.

[3] State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant

Resources, School of Life Sciences, Sun Yat-sen University, Guangzhou,

510275, China.

*Corresponding author: Yangrong Cao

Email: yrcao@mail.hzau.edu.cn

Abstract

Bacterial pathogens and nitrogen-fixing rhizobia employ type III protein

secretion system (T3SS) effectors as potent tools to manipulate plant signaling

pathways, thereby facilitating infection and host colonization. However, the

molecular mechanisms underlying effects of rhizobial effectors on legume

infection remain largely elusive. In this study, we investigated the symbiotic role

of T3SS effectors in the interaction between *Sinorhizobium fredii* NGR234 and

Lotus japonicus. Mutants deficient in the T3SS genes *Tts1* or *NopA* showed

enhanced rhizobial infection of *L. japonicus* roots. Further mutant analysis

showed that the NopT effector negatively affects infection, while the NopM

effector in the absence of NopT promotes infection. Notably, NopM interacts

with the Nod factor receptors LjNFR1 and LjNFR5. NopM ubiquitinates LjNFR5

with ten ubiquitinated lysine residues in LjNFR5 identified using mass

spectrometry. Expression of *NopM* in *L. japonicus* resulted in an approximately

twofold increase in LjNFR5 protein levels and enhanced rhizobial infection. Our

findings indicate that NopM directly interferes with the symbiotic signaling

pathway through interaction and ubiquitination of Nod factor receptors, which

likely benefits rhizobial infection of *L. japonicus*. Our research contributes to the

intricate interplay between the Nod factor signaling pathway and rhizobial T3SS

effectors delivered into host cells.

Keywords: Legume–rhizobium symbiosis; rhizobial effector; NopM; Nod factor

receptors.

Introduction

The legume-rhizobium symbiosis leads to the formation of nodules, where

rhizobia reside and reduce nitrogen into ammonia, nutrilizing the plants. Nodule

formation depends on complicated regulations and mutual signal exchanges

between hosts and symbionts. Legume roots release flavonoids, which might

activate NodD transcriptional regulators, initiating the expression of genes for

type III protein secretion system (T3SS) and biosynthesis of lipo-

chitooligosaccharides known as Nod factors (NFs) (Yang et al., 2022). These

NFs are recognized by Nod factor receptors (NFRs) in host plants, such as

LjNFR1 and LjNFR5 in *Lotus japonicus*, activating symbiotic signaling

pathways crucial for successful nodulation (Radutoiu et al., 2003).

Similar to gram-negative bacterial pathogens whose type III protein secretion

system (T3SS) effectors are vital for interacting with host pathways, rhizobia

inject effector proteins into plant cells to promote rhizobial symbiosis and/or

suppress plant immunity (Teulet et al., 2022). The expression of rhizobial genes

related to T3SS and effectors is induced by flavonoid-NodD complex and TtsI

(Wassem et al., 2008). These T3SS components and effectors, termed

nodulation outer proteins (Nops), have been implicated in infection, nodulation,

and host specificity (Deakin & Broughton, 2009; Teulet et al., 2022). The

symbiosis promoting effects of translocated effectors are often attributed to the

suppression of host defense reactions, while negative (asymbiotic) effects may

be the result of direct or indirect recognition of effectors by intracellular

resistance proteins (Teulet et al., 2022).

Several Nops have been characterized biochemically to date. For instance,

Sinorhizobium fredii NGR234, a strain with broad-host range, has been well-

studied. NopA (Deakin et al., 2005) and NopB (Saad et al., 2005) are major

components of the T3SS complex and interact physically with NopX, facilitating

the translocation of effector proteins into plant cells (Büttner et al., 2002). NopL,

known to delay nodule senescence in bean, acts as a mitogen-activated protein

kinase (MAPK) substrate and reduces levels of pathogenesis-related (PR)

proteins in plants (Ge et al., 2016, Bartsev et al., 2004). NopP synergistically

promotes nodulation with NopL and is phosphorylated by unknown kinases

(Skorpil et al., 2005). NopT, a cysteine protease, regulates symbiosis between

NGR234 and various hosts (Dai et al., 2008), triggering rapid cell death

dependent on its protease activity (Dai et al., 2008). Additionally, NopT cleaves

both LjNFR5 and GmPBS1, regulating plant symbiotic signaling and defense

responses, respectively (Khan et al., 2022, Li et al., 2023). NopM, an E3

ubiquitin ligase, dampens flagellin-induced immunity and interferes with MAPK

signaling (Xu et al., 2018).

While effectors of bacterial pathogens in plant pathogenesis are known to target

plant components of pattern-triggered immunity pathway to favor pathogenic

infection, it remains unknown whether rhizobial effectors function similarly

regulating symbiotic signaling pathway. Here, we investigated the function of

T3SS effectors from *S. fredii* NGR234 in the interaction with *L. japonicus* and

found that the T3SS negatively affects rhizobial infection in this plant. Further

mutant analysis showed that NopT suppresses rhizobial infection, whereas

NopM (in the absence of NopT) promoted infection. NopM interacts with

LjNFR1 and LjNFR5. But NopM selectively ubiquitinates LjNFR5, increasing

LjNFR5 protein levels. Transgenic plants expressing NopM showed increased

infection events and a higher number of nodule primordia. These findings

establish a direct link between rhizobial effectors and the symbiotic signaling

pathway, providing crucial insights into the mechanisms of legume-rhizobium

symbiosis.

Materials and Methods

Plant Materials and Inoculation with Rhizobia

Lotus japonicus Gifu B–129 ecotype was provided by the Center for

Carbohydrate Recognition and Signaling (https://lotus.au.dk/), and used for

phenotype analysis. *L. japonicus* Gifu *nfr 5* mutant (Radutoiu et al., 2003) and

Gifu *proNIN:GUS* transgenic line (Heckmann et al., 2011). Wild type plants of

L. japonicus 'Miyakojima' MG20 were used for phenotype analysis and stable

transformation. Transgenic *L. japonicus* MG20 plants overexpressing *LjNFR5*

from a *Ubiquitin* promoter (encoding HA-tagged LjNFR5) were constructed in

our laboratory and used for hairy root transformation (*proUbiquitin:LjNFR5-HA*

line).

All *L. japonicus* seeds were surface-sterilized with H_2SO_4 for 6 min, and then

with 2 % (w/v) NaClO for 6 min. After synchronization of germination at 4 °C in

the dark for 2 days, the seeds were placed on Murashige & Skoog (MS) medium

(PhytoTech, M519) containing 1.5 % (w/v) agar for germination (22°C for 2 days

in the dark and 2 days in the light). Seedlings were then transferred to growth

pots containing vermiculite supplied with ½ B&D (Broughton & Dilworth)

medium (Broughton and Dilworth, 1971) without nitrogen. Plants were kept

under long–day conditions (16 h light/ 8 h dark) at 22 °C. For infection

experiments, seven-day-old seedlings were inoculated with *S. fredii* NGR234

and NGR234 mutants or *M. loti* MAFF303099 (OD_{600}=0.02).

Plasmid Construction

All plasmids used in this study were generated using MultiF Seamless

Assembly Mix (ABclonal, China, RK21020). Plasmids pG5XX–HA, pG5XX–

Myc, pG5XX–FLAG, pG5XX–nLUC, pG5XX–cLUC, pG5XX–GFP (Bao et al.,

2024), point mutant *LjNFR1^{K351E}* and point mutant *NopM^{C338A}* using overlapping

PCR were generated. To detect the interaction between *LjNFR1*, *LjNFR5*, and

NopM^{C338A} in *Nicotiana benthamiana* in split-luciferase complementation

experiments, the full–length coding sequences of *LjNFR1*, *LjNFR1^{K351E}*,

LjNFR5, *NopM^{C338A}*, and *AtFLS2* were cloned into *Xba*I-*Kpn*I digested pG5XX–

nLUC (Wang et al., 2021) and pG5XX–cLUC (Wang et al., 2021), respectively.

For BIFC experiments, the full–length coding sequences of *LjNFR1*, *LjNFR5*,

NopM^{C338A}, and *AtFLS2* were cloned into pSPYCE (MR) (Waadt et al., 2008)

and pSPYNE (R)173 (Waadt et al., 2008), respectively, between *Xba*I and

*Bam*HI. For the co-IP assay, the full–length *LjNFR1*, *LjNFR5*, *NopM^{C338A}* coding

sequences were cloned into pG5XX–GFP, pG5XX–FLAG, respectively,

between *Xba*I and *Kpn*I. For the pull-down assay with *E. coli,* the sequence

encoding the cytosolic domain of *LjNFR5* fused to a Myc tag was cloned into

plasmid pMAL-c2X (Addgene, 75286) between *Bam*HI and *Hind*III and the

NopM coding sequence was cloned into plasmid pACYCDuet ((Han et al., 2017)

between *Sgs*I and *Not*I. The *Myc* tag was cloned from pG5XX–Myc and inserted

into plasmid pMAL-c2X between *Bam*HI and *Hind*III, generating plasmid pMAL-

c2X-Myc used as control. For *in vitro* ubiquitin assay in *E. coli*, *NopM* or

NopM^{C338A}, and Myc tagged cytosolic domain (CD) of NFR5 (NFR5^{CD}-Myc/

LjNFR5[10KR]-Myc) were inserted into plasmid pACYDuet and pCDFDuet ((Han

et al., 2017) between *Sgs*I and *Not*I respectively. His and FLAG tagged ubiquitin

(His-FLAG-Ub[K0]) was generated by GenScript Biotech. For hairy roots

transformation of *L. japonicus*, the plasmid pUB-RFP (Feng et al., 2021) was

digested with *Kpn*I and *Xba*I, and was inserted fragment *NopM* and *NopM^{C338A}*.

For stable transformation in *L. japonicus NopM* was cloned into *Kpn*I-*Xba*I

digested pUB-Hyg-FLAG (Feng et al., 2021), generating plasmid pUB-Hyg-

NopM-FLAG with a *Ubiquitin* promoter. A 3-kb promoter fragment upstream of

the start codon of *NIN* was then cloned from genomic *L. japonicus* MG20 DNA

and inserted into *Xba*I- *Pst*I digested pUB-Hyg-NopM-FLAG to replace the

Ubiquitin promoter. The resulting plasmid was named proNIN-NopM-Hyg-FLAG.

All primers used are listed in the Supplementary Table S1.

Stable Transformation in *L. japonicus*

The plasmid proNIN-NopM-Hyg-FLAG was transformed into *L. japonicus* MG20

seedlings using *Agrobacterium tumefaciens* strain EHA105 as described by

Handberg and Stougaard (Handberg & Stougaard, 1992). 50 mg/ml

hygromycin B (Sangon Biotech, China, A600230) was used for selection of

transgenic plants. The final positive transgenic plants were detected by PCR

analysis.

Transient Expressing in *N. benthamiana* Leaves

A. tumefaciens EHA105 strains carrying various constructs were cultivated

overnight and resuspended with an *Agrobacterium* culture harboring the P19

suppressor with the ratio 1:1 in infiltration buffer (10 mM $MgCl_2$, 10 mM MES

pH=5.8 and 100 µM acetosyringone). The bacterial suspensions were infiltrated

into leaves of 4–week–old *N. benthamiana* plants. Leaves were harvested for

two days post infiltration.

BiFC and Split Luciferase Assays

In the BiFC assay with transformed *N. benthamiana* leaves, the eYFP fluorescence of fusion proteins was analyzed 2–3 days after infiltration using a confocal microscope (Leica TCS SP8).

In the split–luciferase assay, the plasmids were transformed into *N. benthamiana* leaves using *Agrobacterium* strain EHA105. Three days after infection, leaves were sprayed with 1 mM luciferin (Promega, E1603) and 0.02 % (v/v) Triton X–100. The LUC activity was detected using the Tanon Imaging system (Tanon, China, 4600).

Co-IP and Pull-down Assays

For the co-IP assay, total proteins were extracted from the *N. benthamiana* leaves in IP buffer (50 mM Tris-HCl, pH=7.4, 150 mM NaCl, 1 mM EDTA, 1 % TritonX-100, 5 % glycerol) supplemented with a protease inhibitor cocktail (Yuanye, China, S25910). Samples were incubated at 4 °C for 10 min and centrifuged at 12,000 × g for 10 min at 4 °C. The supernatant was added to 10 μL anti-GFP Nanobody Agarose Beads (Alpa Life Bio, KTSM1301) and incubated at 4 °C for 3 h. The beads were washed at least five times with washing buffer (50 mM Tris-HCl, pH=7.4, 150 mM NaCl, 1 mM EDTA). Western blotting was detected by anti-GFP (ABclonal, China, AB_2770402) and anti-FLAG (Sigma, F1804) antibodies.

For the pull-down assay with *E. coli* BL21 (DE3), protein synthesis of co-expressed MBP-LjNFR5^{CD}-Myc or MBP-Myc and His-NopM was induced with 0.5 mM IPTG and incubation at 28 °C for 18 h. MBP-LjNFR1^{CD}-Myc, or MBP-Myc and His-NopM were co–expressed in a similar way but bacteria were kept

at 18 °C for 20 h. Total proteins were extracted from *E. coli* in column buffer (20

mM Tris-HCl, pH=7.4, 0.2 M NaCl, 1 mM EDTA). Samples were centrifuged at

12,000 × g for 10 min at 4 °C. The supernatant was added to 20 μL amylose

resin beads (New England Biolabs, E8022S) and incubated at 4 °C for 3 h. The

beads then were washed at least five times with column buffer. Western blotting

was detected by anti-Myc (Bio Legend, 626808) antibody and anti-NopM

antibody kindly provided by Dr. Xin from North East Agriculture University.

In Vitro Ubiquitination Assay

The *in vitro* ubiquitination assay was performed using genes encoding AtUBA1

(E1), AtUBC8 (E2), NopM (or NopM^{C338A}), ubiquitin (or His-FLAG-Ub^{K0}) and the

cytosolic domain of LjNFR5 (or LjNFR1) as described previously (Han et al.,

2017). Different combinations of vectors were transformed into the *E. coli* BL21

(DE3). Recombinant protein expression was induced with 0.5 mM IPTG and

cells were grown at 28 °C for 20 h. The cell lysates were subjected to Western

blotting analysis using anti-Myc (Bio Legend, 626808) antibody, anti-FLAG

(Sigma, F1804) antibody, and anti-NopM antibody.

Generation of Hairy Roots Expressing NopM and NopM^{C338A} in *L. japonicus*

The pUB-RFP vectors carrying *NopM* and *NopM^{C338A}* were transformed into

proUbiquitin:LjNFR5-HA seedlings and Gifu seedlings using *A. rhizogenes*

LBA1334 (Feng et al., 2021). The *proLjNFR5:LjNFR5-GFP* and

proLjNFR5:LjNFR5^{10KR}-GFP vectors were transformed into Gifu *nfr5* seedlings

using *A. rhizogenes* LBA1334. The expression of RFP and GFP markers

enabled selection of transgenic hairy roots using a fluorescence stereo

microscope (Nikon, SMZ18).

Analysis of proteins expressed in *L. japonicus*

Total proteins were extracted from transgenic hairy roots using IP buffer (20 mM

Tris–HCl pH=7.5, 50 mM NaCl, 1 mM EDTA, 0.1 % SDS, 1 % Triton X-100)

supplemented with a protease inhibitor cocktail (Yuanye, China, S25910).

Western blotting was performed with anti-HA–peroxidase (Sigma, clone 3F10),

anti-actin (ABclonal, China, AC009), and anti-NopM antibodies.

NGR234 mutants Construction

S. fredii NGR234 derivatives with mutated *NopL (*NGRΩ*nopL*) (Marie et al.,

2003), and *NopT* (NGRΔ*nopT*) (Dai et al., 2008) have been constructed

previously. Additional mutations in the symbiotic plasmid of NGR234 were

generated by homologous recombination using the suicide vector pJQ200SK

(Quandt & Hynes, 1993). *S. fredii* NGR234 derivatives with mutated *NopJ*

(NGRΩ*nopJ*) was provided by Christian Staehelin.

NGR234 derivatives with mutated *TtsI*, *NopA*, *NopM*, *NopP* or *NopZ* (y4yJ)

genes, as well as double mutants with mutated *NopJ*, *NopL*, *NopM, NopP* or

NopZ genes in the *ΔnopT* background, were generated as follows (Fig. S1).

Firstly, a homologous recombination plasmid was constructed. A kanamycin

fragment cloned from plasmid pENTR/D-TOPO (Invitrogen, K240020SP) was

inserted into pJQ200SK digested with *Pst*I and *Bam*HI, yielding plasmid

pJQ200SK-Kan. A *Paph* promoter fragment cloned from plasmid pRJPaph-

bjGFP (Addgene, 67032) was then inserted into pJQ200SK-Kan digested with

*Pst*I, yielding plasmid pJQ200SK-*proKan:Kan* with a kanamycin resistance

interposon (ΩKan). A PCR-amplified 500-bp genomic NGR234 sequence

upstream of nucleotide position 18-21 of a given coding sequence and a

corresponding 800-bp genomic NGR234 sequence downstream of this position

were chosen as homologous arms. Plasmid pJQ200SK-*proKan:Kan* was

digested with *Pst*I, *Xho*I, *Bam*HI, and *Xba*I. The 500-bp upstream sequence

was cloned into the *Pst*I-*Xho*I site, and the 800-bp downstream sequence was

cloned into the *Bam*HI-*Xba*I site, generating a pJQ200SK-Ω plasmid with

homologous arms. Finally, the constructed plasmids were mobilized from *E. coli*

DH5a into NGR234 or the *ΔnopT* mutant by triparental mating using the

pRK2013 helper plasmid (Figurski and Helinski, 1979). Gene replacement was

forced by selecting for the resistance of the Ω interposon marker and for growth

on 5 % (w/v) sucrose. Double-crossover events at homologous sites were

confirmed by PCR analysis using genomic DNA extracted by a sodium dodecyl

sulfate (SDS) method (Fig. S2). Primers used are listed in Supplementary Table

S1 and constructed mutants in Supplementary Table S2.

To observe infection events, GFP-labeled strains were constructed. A GFP-

containing derivative of plasmid pHC60 (NovoPro, V005592) was mobilized into

NGR234 and the mutants by bacterial conjugation (Gross & Caro, 1966). GFP-

labeled colonies were detected by a fluorescence microscope (Nikon SMZ18).

GUS Staining

Roots were stained overnight in X-Gluc solution (50 mM sodium phosphate

pH=7.0, 50 mM EDTA pH=8.0, 0.5 mM K_3Fe (CN)$_6$, 0.5 mM K_4Fe (CN)$_6$, 0.1%

Triton X-100, 1 mM X-Gluc, 20 % methanol) at 37 °C. The chlorophyll was

removed by incubating the samples in 70 % ethanol. Images of roots were

taken using a fluorescence stereo microscope (Nikon, SMZ18).

Gene Expression Analysis

Total RNA was isolated using Plant RNA Kit (Yeasen, China, 19291ES50) from

transgenic hairy roots in plate selected by 50 mg/ml hygromycin B (Sangon

Biotech, China, A600230) inoculated *M. loti* MAFF303099 (OD_{600}=0.02) for 48

h. The total RNA (1 μg) was used for gDNA removal and cDNA synthesis using

the ABScript III RT Master Mix for qPCR with gDNA Remover (ABclonal, China,

RK20429). The cDNA was used as template for RT-PCR using 2X Universal

SYBR Green Fast qPCR Mix (ABclonal, China, RK21203). ATPase was used

as the reference gene. The primer sequences used for RT-PCR were listed in

Supplementary Table S1.

Sequence Alignment and Phylogenetic Analysis of NopM Orthologs

The protein sequences of NopM orthologs were retrieved using the BLASTP

program at the National Center for Biotechnology Information (NCBI)

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). The phylogenetic trees for the NopM

orthologs were constructed using MEGA5. The protein sequence alignment

was performed using MultAlin (Multiple sequence alignment by Florence Corpet

(http://multalin.toulouse.inra.fr/multalin/multalin.html).

Identification of Ubiquitination Sites through Mass Spectrometry

His-LjNFR5-Myc protein obtained from the ubiquitination assay with

Escherichia coli was purified using High Affinity Ni-NTA Resin (GenScript, China,

L00250). The protein and the beads supplemented with a protease inhibitor

cocktail (Yuanye, China, S25910) at 4 °C overnight. The beads were washed

at least five times with washing buffer (20 mM NaH_2PO_4, pH=8.0, 300 mM NaCl,

50 mM imidazole). Finally, the protein was were eluted by elution buffer (20 mM

NaH_2PO_4, pH=8.0, 300 mM NaCl, 250 mM imidazole). The protein was then

analyzed by Center for Protein Research (CPR) at Huazhong Agricultural

University. The LjNFR5 amino acid sequence and identified and non-identified

lysine residues are shown in Supplementary Table S3, Fig.S6 and Fig.S7.

  Results

NopM in a Δ*nopT* mutant positively regulates rhizobial infection of *L. japonicus*

To investigate the involvement of effectors secreted by the T3SS in the

regulation of rhizobial infection, we generated two T3SS knockout mutants of

S. fredii NGR234, namely *ΩttsI* and *ΩnopA*. The former lacked the

transcriptional regulator gene *TtsI* (*ΩttsI*), and the latter lacked a component of

the T3SS pilus (*ΩnopA*). Surprisingly, both mutants induced a significant

increase in infections of *L. japonicus* (Gifu) compared to the wild-type strain

(Fig. 1A, Fig. 1B and Fig. S3A). These findings suggested that T3SS in NGR234

negatively influences rhizobial infection.

To explore the specific contribution of individual effectors to rhizobial infection,

we examined mutants for six putative or *bona fide* effector genes (*ΩnopJ*,

ΩnopL, *ΩnopP*, *ΩnopZ*, *ΩnopM*, Δ*nopT*) (Kimbrel et al., 2013). Compared to

the parent strain NGR234, the Δ*nopT* mutant showed increased numbers of

infection (Fig. 1C and Fig. S3B), however, no significant differences were

observed for other effector mutants (Fig. 1D, 1E and Fig. S3B). This suggested

that the impact of effectors on symbiosis was not dependent on a single effector

but rather on the combined consequences of their activities, acting

synergistically, redundantly, or antagonistically.

To identify potential effectors that positively regulate rhizobial infection in the

Δ*nopT* mutant, we constructed five double mutants (Δ*nopTΩnopJ*,

Δ*nopTΩnopL*, Δ*nopTΩnopP*, Δ*nopTΩnopZ*, Δ*nopTΩnopM*) and inoculated

them onto *L. Japonicus*. The Δ*nopTΩnopJ* and Δ*nopTΩnopL* double mutants

induced massive infections similar to Δ*nopT* (Fig. 1F and Fig. S3C), indicating

that NopJ and NopL in the Δ*nopT* mutant might not affect rhizobial infection. In

contrast, Δ*nopTΩnopP*, Δ*nopTΩnopZ*, and Δ*nopTΩnopM* double mutants

showed reduced infection numbers similar to the NGR234 WT strain (Fig. 1G),

suggesting that NopM, NopP and NopZ might act antagonistically with NopT in

rhizobial infection. Despite previous reports on NopM being an E3 ligase

involved in symbiosis regulation (Xu et al., 2018, Xin et al., 2012), its host

targets have not been reported. We then focused on the identification of NopM

targets in plants.

NopM interacts with LjNFR1 and LjNFR5

Given that NopT negatively regulates rhizobial infection by cleaving LjNFR5

(Bao et al., 2024) and that NopM dimerization occurs at the plasma membrane

(Xu et al., 2018), we hypothesized that NopM might also associate with NFRs.

To test this, we first conducted a luciferase complementationg assay in

Nicotiana benthamiana expression system. A ubiquitin ligase inactive

NopM^{C338A} was used in these experiments, because enzymatically active NopM

induces cell death in *N. benthamiana* cells (Xu et al., 2018). The results showed

that NopM^{C338A} was indeed associated with both LjNFR1 and LjNFR5, as

indicated by strong luciferase activity when proteins were co-expressed in *N.*

benthamiana leaves (Fig. 2A and 2B). Further validation through a bimolecular

fluorescence complementation (BiFC) assay confirmed the interaction between

NopM^{C338A} and LjNFR1 or LjNFR5 (Fig. 2C). A co-immunoprecipitation (co-IP)

assay provided additional evidence supporting the interaction between

NopM^{C338A} and LjNFR1 or LjNFR5 (Fig. 2D, 2E). For a subsequent *in vitro* pull-

down assay, the cytosolic domain (CD) of LjNFR1 or LjNFR5 was co-expressed

with NopM in *Escherichia coli* cells and it was found that NopM indeed

physically interacted with Both LjNFR1^{CD} and LjNFR5^{CD} (Fig. 2F, 2G).

NopM ubiquitinates LjNFR5

Since NopM is an E3 ligase (Xin et al., 2012), we wondered whether NopM can

ubiquitinate NFRs. To test this, a ubiquitination assay was carried out using a

reconstituted ubiquitination cascade system (E1, E2, E3 enzymes and ubiquitin)

expressed in *E. coli* cells (Han et al., 2017). As shown in Fig. 3A, NopM, but not

NopM^{C338A}, exhibited autoubiquitination activity, which is consistent with

previous studies (Xu et al., 2018). Next, NopM was co-expressed with LjNFR5

in *E. coli* cells.The result showed that NopM could ubiquitinate LjNFR5^{CD} as

strong smear protein bands were visible on immunoblots. The strong smear

bands of LjNFR5^{CD} suggested that LjNFR5^{CD} might be poly-ubiquitinated (Fig.

3A). Interestingly, NopM did not ubiquitinate LjNFR1^{CD} (Fig. 3B). These findings

indicated that NopM selectively ubiquitinates LjNFR5.

Next, we wondered whether the ubiquitination of LjNFR5 by NopM is a

conserved mechanism that also occurs in other symbiotic interactions. We first

phylogenetically analyzed NopM sequences from different rhizobium species.

Amino acid sequence comparisons showed that the NopM proteins can be

divided into three clades (Fig. S4). NopMs of *Bradyrhizobium diazoefficiens*

USDA110 and *S. fredii* HH103 were located in Clade I and Clade III,

respectively. NopM of *S. fredii* NGR234 was located in Clade II. Two *NopM*

genes, named NopM1$_{USDA110}$ and NopM2$_{USDA110}$ were present in the USDA110.

S. fredii HH103 possesses two NopM genes with identical sequences. A

multiple sequence alignment indicated that C-terminal E3 ligase domain of

NopM was conserved whereas the length of the N-terminal Leucine-Rich

Repeats domain was different among three NopM clades (Fig. S5). We then

examined the ubiquitination of soybean GmNFR5^{CD} by NopM1$_{USDA110}$ and

NopM2$_{USDA110}$ (Clade I) and NopM$_{HH103}$ (clade III). The results showed that

these three NopMs could ubiquitinate GmNFR5^{CD} (Fig. 3C). These findings

suggested that NopM homologs from different rhizobium species exhibited a

conserved activity in ubiquitinating NFR5 proteins.

Expression of NopM in *L. japonicus* increases LjNFR5 protein levels

Ubiquitination is an important post-translational modification that controls

protein levels by mediating protein degradation or stability (Smalle and Vierstra,

2004). To assess whether NopM-mediated ubiquitination regulates LjNFR5

protein levels *in vivo*, we expressed *NopM* and *NopM^C338A* under the control of

a *Ubiquitin* promoter in hairy roots of stable transgenic *L. japonicus* expressing

LjNFR5-HA. LjNFR5 was detected at higher levels in NopM-expressing roots

than those in roots expressing NopM^{C338A} (Fig. 4A and 4B). These findings

indicated that the ubiquitination of LjNFR5 by NopM did not cause degradation,

but rather led to increased LjNFR5 protein levels.

Given that ubiquitination of LjNFR5 lead to its stability, we proposed that NopM-

mediated ubiquitination might form mono-ubiquitination chains. To investigate

the type of ubiquitin-chain linkage on LjNFR5, we utilized the Ub^{K0} variant (all

lysine residues in ubiqutin were substituted to arginine) (Ma et al., 2020).

Interestingly, the ubiquitination pattern of LjNFR5 in presence of Ub^{K0} remained

similar to that with ubiquitin, suggesting that LjNFR5 might be mono-

ubiquitinated by NopM (Fig. 4C). Accordingly, mass spectrometry analysis of

LjNFR5 identified ten potential ubiquitinated lysine residues (Fig. S6 and Fig.

S7). LjNFR5^{10KR}, a mutant version of LjNFR5 with ten lysine residues

substituted to arginine residues, still interacts with NopM (Fig. S8A and S8B)

and did not abolish the ubiquitination by NopM (Fig. 4D). However, when using

the Ub^{K0} variant, LjNFR5 ubiquitination was dramatically reduced (Fig. 4D),

suggesting that LjNFR5 still contains additional lysine residues that can be

either mono- or poly-ubiquitinated.

To assess the impact of ubiquitinated lysine residues on LjNFR5 function, a cell

death test was performed within *N. benthamiana* leaves that depends on co-

expression of NFRs. Co-expression of LjNFR1 with LjNFR5 iinduces a rapid

cell death as reported previously (Madsen et al., 2011). Co-expression of

LjNFR1 with LjNFR5^{10KR} caused a similar cell death response (Fig. S8C). We then generated transgenic hairy roots of *L. japonicus* (Gifu) *nfr5* mutant plants that expressed *LjNFR5* or *LjNFR5^{10KR}* driven by the native promoter (*proLjNFR5:LjNFR5*, *proLjNFR5:LjNFR5^{10KR}*). When inoculated with GFP-labelled NGR234, hairy roots transformed with the *proLjNFR5: LjNFR5^{10KR}* construct showed barely infection events whereas numerous infections were observed in roots expressing the LjNFR5 (Fig. S8D). These findings indicate that specific lysine residues required for ubiquitinatio in LjNFR5 are essential for LjNFR5 to activate symbiotic signaling.

NopM promotes rhizobial infection and nodule primordia formation

To examine the role of NopM in symbiotic response, we transformed *L. japonicus* MG20 plants with *NopM* driven by *CaMV 35S* or *Ubiquitin* promoters. Unfortunately, we could not obtain stable transgenic plants with these constructs, suggesting that constitutive NopM expression might have a negative effect on plant growth. We therefore generated *L. japonicus* MG20 plants expressing *NopM* driven by *LjNIN* promoter (*proNIN:NopM*), whose activity is induced during symbiosis   (Fig. S9 A-C). Two independent lines were obtained and inoculated with a GFP-tagged *Mesorhizobium loti* MAFF303099 which does not contain *NopT* and *NopM* genes in its genome. As shown in Fig. 5A, 5D and Fig. S9D, increased infection foci and infection threads were observed in these transgenic plants compared to control plants. Consistent with this, transgenic plants also formed more nodule primordia than with wild type plants at 14 dpi (Fig. 5B). Finally, to corroborate these results, we generated transgenic roots expressing *NopM* driven by *Ubiquitin* promoter. Upon inoculation with GFP-tagged *M. loti*, the number of rhizobial infections in these *NopM*-expressing roots was significantly higher than in control roots transformed with the control vector (Fig. 5C). Taken together, these data indicate that NopM positively regulates rhizobial infection and nodule primordia

formation in *L. japonicus*.

Discussion

The study of type III secretion system (T3SS) effectors in gram-negative

bacteria has primarily focused on their role in promoting pathogenic infections

in plants. While there are parallels between the processes of rhizobial infection

and colonization and those of pathogenic bacteria, the specific functions of

rhizobial effectors in manipulating the plant symbiotic pathway have remained

elusive. This overall aim of study was to unravel the mechanisms by which

T3SS effectors regulate rhizobium-legume symbiosis. This research

demonstrated that the rhizobial effector NopM can associate with two key

receptors, LjNFR1 and LjNFR5, essential for activating symbiotic signaling in

plants. NopM selectively ubiquitinates LjNFR5, leading to increased protein

levels and enhanced rhizobial infection. These findings suggested a regulatory

model in which NopM targets NFR5 in legumes to promote the symbiotic

signaling transduction by altering the stability of NFR5.

Root infection by rhizobia often depends on a mixture of T3SS effectors which

may act redundantly, synergistically, or antagonistically within host cells.

Accordingly, mutations of effector genes show positive, negative or no effects

on symbiosis with a given host plant. Analysis of rhizobial double mutants may

reveal the symbiotic role of an effector that would not have been visible by

mutating a single effector gene. While mutation of *nopT* in NGR234 had a

positive effect on infections of *L. japonicus* roots, mutation of other effector

genes did not result in an altered phenotype. However, mutation of *nopM, nopP*

and nopZ in the *nopT* mutant resulted in reduced infection numbers, indicating

that the symbiosis promoting effect of these effector genes was only observed

in double mutants. This suggests that NopT and NopM work antagonistically,

with a possiblity sharing the same target protein from hosts, to precisely

regulate rhizobial infection. The observed asymbiotic role of NopT may be due

to interactions with NFRs (Bao et al., 2024) or protein kinases related to

effector-triggered immunity (Khan et al., 2022, Li et al., 2023). Similar to NopT,

we report here that NopM can interact with the LjNFR1 and LjNFR5 of *L.*

japonicus. The findings suggest that NopT and NopM play antagonistically

redundant roles to fine-tune rhizobial infection in host plants.

LjNFR1 and LjNFR5 play vital roles in perceiving rhizobial NFs, activating

symbioti signaling in *L. japonicus* (Radutoiu et al., 2003, Madsen et al., 2011).

Once NF signaling is initiated, protein turnover and signaling activity of NFRs

may be critical for maintaining proper symbiotic signaling and subsequent

rhizobial infection. Previous research demonstrated that mutation in the

LjPUB13 gene suppresses nodulation in *L. japonicus* (Tsikou et al., 2018).

LjPUB13 is an E3 ligase that interacts with and ubiquitinates LjNFR5 *in vitro*

(Tsikou et al., 2018). Thus, the biological role of LiPUB13 in promoting rhizobial

symbiosis may be similar to that of NopM, as both proteins can ubiquitinate

LjNFR5. In general, proteins with K48-linked polyubiquitin chains are rapidly

degraded, whereas proteins with other polyubiquitin chains and mono-

ubiquitinated proteins are not degraded and often exhibit altered properties

(Zhou & Zeng, 2017). In our study, we demonstrate that NopM can mono-

ubiquitinate LjNFR5 to stabilize LjNFR5, promoting rhizobial infection in *L.*

*japonicu*s. Hairy roots of *LjNFR5-HA* transgenic plants expressing

proUbiquitin:NopM showed increased LjNFR5 protein levels when compared to

those transformed with *proUbiquitin:NopM^{C338A}*). Although poly-ubiquitination of

LjNFR5 by NopM was detected at weak levels, the mono-ubiquitination of

LjNFR5 might be major reason leading to increased protein levels. It is therefore

tempting to speculate that NopM preferentially monoubiquitinates LjNFR5

and/or that ubiquitinated LjNFR5 contains polyubiquitin chains without K48-

linkages.

Pathogens utilize effectors to facilitate infection by suppressing host immunity,
exemplified by the E3 ubiquitin ligase effector AvrPtoB in *Pseudomonas*
syringae. AvrPtoB regulates flagellin receptor FLS2 (Goehre et al., 2008) and
chitin receptor CERK1 (Gimenez-Ibanez et al., 2009) protein stability in
Arabidopsis thaliana, leading to suppression of PTI. Similarly, NopM may
employ a similar strategy to support infection during symbiosis to avoid PTI.
NopM dampens the flg22-induced ROS burst in *N. benthamiana* (Xin et al.,
2012) and induces cell death in tobacco *via* its E3 ligase activity (Xu et al., 2018).
549

On the other hand, NopM activity in certain plant cells appears to induce
responses that are related to ETI: (i) The difficulties we observed in the
production of MG20 plants constitutively expressing NopM indicate negative
effects of NopM on plant development under certain physiological conditions.
(ii) NopM of NGR234 expressed in tobacco induces rapid cell death, which
depends on the E3 ubiquitin ligase activity (Xu et al., 2018). (iii) Mutant analysis
showed that NopM of *Bradyrhizobium elkanii* USDA61 triggers ETI-like early
nodule senescence in MG20 (Kusakabe et al., 2020). (iv) Hairy roots of MG20
transformed with a construct containing *NopM* of NGR234 and a tandem *CaMV*
35S promoter formed fewer nodules than plants expressing cnotrol vectors or
a *NopM*^{C338A} construct (Xu et al., 2018). Thus, although NopM promotes
rhizobial infection and nodule primordia formation shown in this study, the
ubiquitin ligase activity of NopM appears to have negative effects on nodulation
at later stages of the symbiosis. Host plants may use ETI as a mechanism to
avoid unfavorable rhizobia strains.
565

566 Furthermore, another hypothetical aspect is worth mentioning. In NGR234
567 cultures treated with host flavonoids, genes related to the TTSS are later
568 expressed than *nod* genes required for Nod factor synthesis (Perret et al., 1999).
569 Thus, it can be assumed that effectors are translocated into host cells in which
570 symbiotic signaling might have already started. Because Nod factor receptors

can be viewed as gatekeepers controlling symbiotic signaling transduction, the precise regulation on Nod factor receptors could be the most effective mean by rhizobium to fine-tune symbiotic signaling in host cells. NopT of NGR234 has been found to proteolyze LjNFR5 and to inhibit symbiotic infection (Bao et al., 2024), while in our study NopM positively affected LjNFR5 levels and promoted infection by rhizobia. Mutual regulation on NFR5 by two effectors from the same rhizobium might lead to an optimized or suboptimal infection depending on the host plant.

Acknowledgments

We thank Jens Stougaard (Aarhus University) for providing *nfr5* mutant and *proNIN:GUS* transgenic seeds. We also thank Dongping Lv (Shanghai Jiaotong University) for providing bacterial ubiquitination system strains. This work was supported by the National Key R&D Program of China (2019YFA0904700), the National Natural Science Foundation of China (32090063). The qPCR and microscopy data were acquired from the Core Facility Center run by the National Key Lab of Agricultural Microbiology. Mass spectrometry analysis was performed by the Center for Protein Research at Huazhong Agricultural University.

Competing Interests

None declared.

Author Contributions: Y.W., D.X., C.S., and Y.C. conceived the idea and designed the research, Y.W., H.B., Y.L., and L.Y. performed experiments; Y.W., H.B., H.Z., and Y.C. analyzed the data, H. L. performed the bioinformatic analysis; C.S. provided some rhizobium mutants; Y.W. and Y.C wrote the manuscript; C.S. provided precious ideas and revised the manuscript.

Data availability

All data in this study are included in the main text or in the Supporting
Information.

References

Büttner, D., Nennstiel, D., Klüsener, B., and Bonas, U. 2002. Functional
analysis of HrpF, a putative type III translocon protein from *Xanthomonas
campestris* pv. *vesicatoria*. *J Bacteriol* **184**: 2389-2398.

**Bartsev, A.V., Deakin, W.J., Boukli, N.M., McAlvin, C.B., Stacey, G., Malnoe,
P., Broughton, W.J., and Staehelin, C. 2004.** NopL, an effector protein of sp
NGR234, thwarts activation of plant defense reactions. *Plant Physiol* **134**: 871-
879.

Broughton, W.J., and Dilworth, M.J. 1971. Control of leghaemoglobin
synthesis in snake beans. *Biochem J* **125**: 1075-1080.

Dai, W.J., Zeng, Y., Xie, Z.P., and Staehelin, C. 2008. S Symbiosis-promoting
and deleterious effects of NopT, a novel type 3 effector of Rhizobium sp strain
NGR234. *J Bacteriol* **190**: 5101-5110.

Deakin, W.J., and Broughton, W.J. 2009. Symbiotic use of pathogenic
strategies: rhizobial protein secretion systems. *Nat. Rev. Microbiol* **7**: 312-320.

**Deakin, W.J., Marie, C., Saad, M.M., Krishnan, H.B., and Broughton, W.J.
2005.** NopA is associated with cell surface appendages produced by the type
III secretion system of Rhizobium sp. strain NGR234. *Mol Plant Microbe*

Interact **18**: 499-507.

Feng, Y., Wu, P., Liu, C., Peng, L.W., Wang, T., Wang, C., Tan, Q., Li, B.X.,

Ou, Y.J., Zhu, H., et al. 2021. Suppression of LjBAK1-mediated immunity by

SymRK promotes rhizobial infection in *L. japonicus*. *Mol. Plant* **14**: 1935-1950.

Figurski, D.H., and Helinski, D.R. 1979. Replication of an origin-containing

derivative of plasmid Rk2 dependent on a plasmid function provided in trans.

Proc Natl Acad Sci USA **76**: 1648-1652.

Ge, Y.Y., Xiang, Q.W., Wagner, C., Zhang, D., Xie, Z.P., and Staehelin, C.

2016. The type 3 effector NopL of Sinorhizobium sp. strain NGR234 is a

mitogen-activated protein kinase substrate. *J Exp Bot* **67**: 2483-2494.

Gimenez-Ibanez, S., Hann, D.R., Ntoukakls, V., Petutschnig, E., Lipka, V.,

and Rathjen, J.P. 2009. AvrPtoB targets the LysM receptor kinase CERK1 to

promote bacterial virulence on plants. *Curr Biol* **19**: 423-429.

Goehre, V, Spallek, T., Haeweker, H., Mersmann, S., Mentzel, T., Boller, T.,

de Torres, M., Mansfield, J.W., and Robatzek, S. 2008. Plant pattern-

recognition receptor FLS2 is directed for degradation by the bacterial ubiquitin

ligase AvrPtoB. *Curr Biol* **18**: 1824-1832.

Gross, J.D., and Caro, L.G. 1966. DNA transfer in bacterial conjugation. *J*

Mol Biol **16**: 269-284.

Han, Y.F., Sun, J.H., Yang, J., Tan, Z.Y., Luo, J.J., and Lu, D.P. 2017.

Reconstitution of the plant ubiquitination cascade in bacteria using a synthetic

biology approach. *Plant J* **91**: 766-776.

Handberg, K., and Stougaard, J. 1992. *Lotus japonicus*, an autogamous,
diploid legume species for classical and molecular-genetics. *Plant J* **2**: 487-496.
663

664 **Heckmann, A.B., Sandal, N., Bek, A.S., Madsen, L.H., Jurkiewicz, A.,**
665 **Nielsen, M.W., Tirichine, L., and Stougaard, J. 2011.** Cytokinin induction of
666 root nodule primordia in *Lotus japonicus* is regulated by a mechanism operating
667 in the root cortex. *Mol Plant Microbe Interact* **24**: 1385-1395.
668

669 **Khan, A., Wadood, S.F., Chen, M., Wang, Y., Xie, Z.P., and Staehelin, C.**
670 **2022.** Effector-triggered inhibition of nodulation: A rhizobial effector protease
671 targets soybean kinase GmPBS1-1. *Plant Physiol* **189**: 2382-2395.
672

673 **Kimbrel, J.A., Thomas, W.J., Jiang, Y., Creason, A.L., Thireault, C.A.,**
674 **Sachs, J.L., and Chang, J.H. 2013.** Mutualistic co-evolution of type III effector
675 genes in *Sinorhizobium fredii* and *Bradyrhizobium japonicum*. *PLoS Pathog* **9**:
676 e1003204.
677

678 **Kusakabe, S., Higasitani, N., Kaneko, T., Yasuda, M., Miwa, H., Okazaki, S.,**
679 **Saeki, K., Higashitani, A., and Sato, S. 2020.** Accessions possess multiple
680 checkpoints triggered by different type III secretion system effectors of the wide-
681 host-range symbiont USDA61. *Microbes Environ* **35**: 2020.
682

683 **Li, D.D., Zhu, Z.K., Deng, X.M., Zou, J.N., Ma, C., Li, C.D., Yin, T., Liu, C.Y.,**
684 **Wang, J.H., Chen, Q.S., et al. 2023.** A hub gene interacting with rhizobial type-
685 III effectors NopT and NopP, regulates soybean nodulation. *Agronomy-Basel*
686 **13**: 1242.
687

688 **Ma, X.Y., Claus, L.A.N., Leslie, M.E., Tao, K., Wu, Z.P., Liu, J., Yu, X., Li, B.,**
689 **Zhou, J.G., Savatin, D.V., et al. 2020.** Ligand-induced monoubiquitination of
690 BIK1 regulates plant immunity. *Nature* **581**: 199-203.

Madsen, E.B., Antolín-Llovera, M., Grossmann, C., Ye, J.Y., Vieweg, S., Broghammer, A., Krusell, L., Radutoiu, S., Jensen, O.N., Stougaard, J., et al. 2011. Autophosphorylation is essential for the function of the Nod factor receptor 1 and receptor-mediated signaling in cooperation with Nod factor receptor 5. *Plant J* **65**: 404-417.

Marie, C., Deakin, W.J., Viprey, V., Kopciñska, J., Golinowski, W., Krishnan, H.B., Perret, X., and Broughton, W.J. 2003. Characterization of Nops, nodulation outer proteins, secreted via the type III secretion system of NGR234. *Mol. Plant Microbe Interact* **16**: 743-751.

Perret, X., Freiberg, C., Rosenthal, A., Broughton, W.J., and Fellay, R. 1999. High-resolution transcriptional analysis of the symbiotic plasmid of Rhizobium sp. NGR234. *Mol Microbiol* **32**: 415-425.

Quandt, J., and Hynes, M.F. 1993. Versatile suicide vectors which allow direct selection for gene replacement in gram-negative bacteria. *Gene* **127**: 15-21.

Radutoiu, S., Madsen, L.H., Madsen, E.B., Felle, H.H., Umehara, Y., Gronlund, M., Sato, S., Nakamura, Y., Tabata, S., Sandal, N., et al. 2003. Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature* **425**: 585-592.

Saad, M.M., Kobayashi, H., Marie, C., Brown, I.R., Mansfield, J.W., Broughton, W.J., and Deakin, W.J. 2005. NopB, a type III secreted protein of Rhizobium sp. strain NGR234, is associated with pilus-like surface appendages. *J Bacteriol* **187**: 1173-1181.

Skorpil, P., Saad, M.M., Boukli, N.M., Kobayashi, H., Ares-Orpel, F.,

Broughton, W.J., and Deakin, W.J. 2005. NopP, a phosphorylated effector of

Rhizobium sp. strain NGR234, is a major determinant of nodulation of the

tropical legumes *Flemingia congesta* and *Tephrosia vogelii*. *Mol Microbiol* **57**:

1304-1317.

Smalle, J., and Vierstra, R.D. 2004. The ubiquitin 26S proteasome proteolytic

pathway. *Annu Rev Plant Biol* **55**: 555-590.

Teulet, A., Camuel, A., Perret, X., and Giraud, E. 2022. The versatile roles of

type III secretion systems in rhizobium-legume symbioses. *Annu Rev Microbiol*

76: 45-65.

Tsikou, D., Ramirez, E.E., Psarrakou, I.S., Wong, J.E., Jensen, D.B., Isono,

E., Radutoiu, S., and Papadopoulou, K.K. 2018. A E3 ligase interacts with

the Nod Factor Receptor 5 and positively regulates nodulation. *BMC Plant Biol*

18: 217.

Waadt, R., Schmidt, L.K., Lohse, M., Hashimoto, K., Bock, R., and Kudla,

J. 2008. Multicolor bimolecular fluorescence complementation reveals

simultaneous formation of alternative CBL/CIPK complexes. *Plant J* **56**: 505-

516.

Wang, L., Yu, G., Macho, A.P., and Lozano-Duran, R. 2021. Split-luciferase

complementation imaging assay to study protein-protein interactions in

Nicotiana benthamiana. *Bio Protoc* **11**: e4237.

Wassem, R., Kobayashi, H., Kambara, K., Le Quere, A., Walker, G.C.,

Broughton, W.J., and Deakin, W.J. 2008. TtsI regulates symbiotic genes in

Rhizobium species NGR234 by binding to tts boxes. *Mol Microbiol* **68**: 736-748.

Xin, D.W., Liao, S., Xie, Z.P., Hann, D.R., Steinle, L., Boller, T., and Staehelin,

C. 2012. Functional analysis of NopM, a novel E3 ubiquitin ligase (NEL) domain

effector of Rhizobium sp. strain NGR234. *PLoS Pathog* **8**: e1002707.

Xu, C.C., Zhang, D., Hann, D.R., Xie, Z.P., and Staehelin, C. 2018.

Biochemical properties and in planta effects of NopM, a rhizobial E3 ubiquitin

ligase. *J Biol Chem* **293**: 15304-15315.

Yang, J., Lan, L., Jin, Y., Yu, N., Wang, D., and Wang, E. 2022. Mechanisms

underlying legume-rhizobium symbioses. *J Integr Plant Biol* **64**: 244-267.

Zhou, B.J., and Zeng, L.R. 2017. Conventional and unconventional

ubiquitination in plant immunity. *Mol Plant Pathol* **18**: 1313-1330.

Supporting Information

Fig. S1. Schematic of NGR234 Mutants.

Fig. S2. PCR Analysis of NGR234 Mutants.

Fig. S3. GUS Staining Pictures for Roots of *L. japonicus* Inoculated with *S. fredii*

NGR234 and Mutant Strains.

Fig. S4. Phylogenetic Tree of NopM Homologs in Different Rhizobial Species.

Fig. S5. Amino Acid Sequence Alignment of NopM Homologous in Different

Rhizobial Species.

Fig. S6. Amino Acid Sequence of LjNFR5 with Identified Lysine Ubiquitination

Sites.

Fig. S7. *In Vitro* Ubiquitination Sites of LjNFR5 Identified by Mass Spectrometry.

Fig. S8. LjNFR5[10KR] Does Not Affect Interactions with Nod Factor Receptors

but Is Symbiotically Inactive.

Fig. S9. Detection of *NopM* DNA, *NopM* Expression and NopM Protein in

Transgenic Plants

Table S1. Primers Used in This Study.

Table S2. Mutants of *S. fredii* NGR234 Constructed in This Study.

Table S3. LjNFR5 *In Vitro* Ubiquitination Sites Identified by Mass Spectrometry.

Figures

Figure 1. Infection Numbers of *L. japonicus* Roots Inoculated with NGR234 and Mutant Strains.

The number of rhizobial infections formed on *L. japonicus* (Gifu) roots inoculated with *S. fredii* NGR234 wild-type and mutants at 6 dpi, (A) NGR234 (n=8) and *ΩttsI* (n=8); (B) NGR234 (n=8) and *ΩnopA* (n=8); (C) NGR234 (n=8) and *ΔnopT* (n=8); (D) NGR234 (n=10) and *ΩnopM* (n=10); (E) NGR234 (n=8), *ΩnopJ* (n=7), *ΩnopL* (n=7), *ΩnopP* (n=8), and *ΩnopZ* (n=8); (F) NGR234 (n=8), *ΔnopT* (n=5), *ΔnopTΩnopJ* (n=5), and *ΔnopTΩnopL* (n=5); (G) NGR234 (n=5), *ΔnopT* (n=5), *ΔnopTΩnopM* (n=5), *ΔnopTΩnopP* (n=5), and *ΔnopTΩnopZ* (n=5). All the rhizobia were labeled with GFP markers. Data are mean ± SD (Student's t-test, $P < 0.01$; ns, not significant).

Figure 2. NopM Interacts with LjNFR1 and LjNFR5.

(A) and (B) NopM interacts with LjNFR5 or LjNFR1 in split-luciferase assay. NopM^{C338A}-nLUC was co-expressed with LjNFR5-cLUC and LjNFR1-cLUC in *N. benthamiana* leaves. NopM^{C338A}-nLUC and AtFLS2-cLUC are negative controls, whileLjNFR5-cLUC and LjNFR1^{K351E}-nLUC (kinase-dead) served as a positive control. The combination NopM^{C338A}-nLUC with NopM^{C338A}-cLUC was also analyzed.Luminescence signals indicate protein–protein interactions.

(C) NopM interacts with LjNFR5 or LjNFR1 in BiFC assay. NopM^{C338A}-nYFP was co-expressed with LjNFR5-cYFP or LjNFR1-cYFP in *N. benthamiana*, and fluorescence was visualized by confocal microscopy. Negative controls included NopM^{C338A}-nYFP with AtFLS2-cYFP. The combination NopM^{C338A}-nYFP with NopM^{C338A}-cYFP was also examined.   Bar=25 μm.

(D) and (E) NopM interacts with LjNFR5 or LjNFR1 inCo-immunoprecipitation (co-IP) assay. LjNFR5-GFP and LjNFR1-GFP were immunoprecipitated with anti-GFP antibody and detected by immunoblotting with anti-FLAG antibody.

(F) and (G) NopM physically interacts LjNFR5 or LjNFR1 in pull-down assay. MBP-LjNFR5-Myc, MBP-LjNFR1-Myc, and NopM were co-expressed in *E. coli* cells. Pull-down was performed with amylose resin and detected by immunoblotting with anti-NopM and anti-Myc antibodies.

Figure 3. NopM Ubiquitinates LjNFR5.

(A) NopM , but not NopM^{C338A}, ubiquitinates LjNFR5 in *E. coli.*. NopM or

NopM^{C338A}, AtUBA1 (E1), AtUBC8 (E2), His-FLAG-Ub, and Myc tagged

cytosolic domain (CD) of LjNFR5 (LjNFR5^{CD}-Myc) were co-expressed in *E.*

coli. Bacterial lysates were subjected to immunoblotting using anti-FLAG, anti-

Myc, and anti-NopM antibodies. Asterisks indicated non- specific band.

(B) NopM ubiquitinates LjNFR5, but not LjNFR1. LjNFR5^{CD}-Myc or LjNFR1^{CD}-

Myc were co-expressed with NopM, AtUBA1 (E1), AtUBC8 (E2) and His-

FLAG- Ub. Western blotting was performed with bacterial lysates using anti-

Myc and anti-FLAG antibodies.

(C) Different NopM proteins from different rhizobium species ubiquitinate

LjNFR5 or GmNFR5. The Myc-tagged cytosolic domain of both NFR5

(NFR5^{CD}-Myc), NopM from indicated strains, AtUBA1 (E1), AtUBC8 (E2), and

His and FLAG-tagged ubiquitin (His-FLAG- Ub) were co-expressed in *E. coli.*

The bacterial lysates were subjected to Western blotting using anti-FLAG and

anti-Myc antibodies.

Figure 4. NopM Expression in *L. japonicus* Roots Increases LjNFR5

Levels and NopM-mediated Multi-monoubiquitination of LjNFR5 with

Ub^{K0}.

(A) NopM expression in *L. japonicus* results in increased LjNFR5 protein

levels. NopM-FLAG driven by the *Ubiquitin* promoter, was expressed in hairy

roots of *proUbiquitin:LjNFR5-HA* plants. Protein levels detected by

immunoblotting with anti-HA, anti-NopM, and anti-actin antibodies.(B)

Quantification of signals of Western blot bands shown in panel (A). Protein

expression levels of LjNFR5 are shown as fold change relative to actin. **$p <$

0.05 (Student's t-test).

(C) NopM-mediated multi-monoubiquitination of LjNFR5 with Ub^{K0} The

ubiquitination assay with *E. coli* cells was performed using His-Flag-tagged

Ub^{K0}, a ubiquitin variant with all lysine residues substituted to arginine (His-

Flag- Ub^{K0}) Ubiquitinated proteins were detected by immunoblotting with anti-

FLAG and anti-Myc. Asterisks indicated non- specific bands.

(D) A similar ubiquitination experiment was performed with $LjNFR5^{10KR}$, in

which ten lysine residues are substituted to arginine

Figure 5. *L. japonicus* Plants Transformed with *proNIN:NopM* and

Inoculated with *M. loti* Show Increased Infection and Nodule Primordia

Formation.

(A) The number of rhizobial infections per plant was significantly greater in

transgenic *L. japonicus* MG20 lines carrying the *proNIN:NopM* construct than

in the MG20 control plants. Roots were inoculated with GFP-labelled *M. loti*

303099 and microscopically analyzed at 5 dpi. Error bars represent ±SD. ***p

< 0.01 (Student's t-test, n=8).

(B) Nodule primordia number per plant were counted in *proNIN:NopM* lines

inoculated with GFP-labelled *M. loti* when analyzed at 14 dpi with *M. loti*.

Data are mean ±SD (***p < 0.01, Student's t-test; MG20, n=12; *proNIN:*

NopM-1, n=11; *proNIN: NopM*-2, n=9).

(C) The number of rhizobial infections per plant increased significantly in hairy

roots of *L. japonicus* MG20 carrying the *proUbiquitin:NopM* construct as

compared to the empty vector control (EV). Roots were microscopically

analyzed at 5 dpi. Error bars represent ±SD. ***p < 0.01 (Student's t-test;

MG20, n=8; *proUbiquitin:NopM*, n=5).

(D) Microscopic images illustrating rhizobial root infection events (the upper

images were infection foci and the lower images were infection threads) in

MG20 wild type and *proNIN:NopM* lines (5 dpi). Fluorescent, GFP-

expressing *M. loti* bacteria are shown in the upper images and corresponding

brightfield images are presented below. Bar=200 μm. White asterisks indicate

bacteria in the upper images.

909