Deep Quantitative Proteomics Identifies Conserved Proteome Alterations in Low and High-Grade Serous Ovarian Cancers

Christopher M. Tarney[1,2*], Paulette Mhawech-Fauceglia[1], Jonathan Ogata[1,3], Julie Oliver[1,3], Tamara Abulez[1,3], Brian L. Hood[1,3], Kelly A. Conrads[1,3], Kendal Rosalik[4], Kwong-Kwok Wong[5], David M. Gershenson[5], Sanghoon Lee[5], Anil K. Sood[5], Robert C. Bast, Jr[6], Kathleen M. Darcy[1-3], Neil T. Phippen[1,2], G. Larry Maxwell[1,2,7], Thomas P. Conrads[1,2, 7], Nicholas W. Bateman[1-3*]

[1]Gynecologic Cancer Center of Excellence, Department of Gynecologic Surgery and Obstetrics, Uniformed Services University and Walter Reed National Military Medical Center, Bethesda, MD 20889, USA.

[2]The John P. Murtha Cancer Center, Uniformed Services University and Walter Reed National Military Medical Center, MD 20889, USA

[3]Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc. Bethesda, MD 20817, USA.

[4]Department of Gynecologic Surgery and Obstetrics, Madigan Army Medical Center, Joint Base Lewis-McChord, WA 98431, USA.

[5]Department of Gynecologic Oncology and Reproductive Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

[6]Department of Experimental Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

[7]Women's Health Integrated Research Center, Women's Service Line, Inova Health System, VA 22003, USA.

*Author's contributed equally

Keywords: high-grade serous ovarian cancer, low-grade serous ovarian cancer, HGSOC, LGSOC, quantitative proteomics, mucin-16, MUC16, cancer antigen 125, CA125, hepatocyte growth factor receptor, focal adhesion kinase

Address correspondence to: Nicholas W. Bateman, email: batemann@whirc.org or Thomas P. Conrads, email: conrads@whirc.org, 3289 Woodburn Rd, Suite 375, Annandale, VA 22003; Tel: 703-207-3357; Fax: 703-560-0919.

Abstract

Low-grade serous ovarian carcinoma (LGSOC) is a rare and largely chemoresistant subtype of epithelial ovarian cancer. Unlike treatment for high-grade serous ovarian cancer (HGSOC), management options for LGSOC patients are limited, in part, due to a lack of deep molecular characterization of this disease. To address this limitation, we performed quantitative mass spectrometry-based proteomic analyses of whole tissue collections of tumors harvested from LGSOC (n=12) and HGSOC (n=24) patients or normal fallopian tube tissues (n=12) from women with benign disease. We quantified 7,043 proteins across all patient samples and unsupervised analyses revealed proteome alterations distinctly stratify fallopian tube tissue, LGSOC, and HGSOC tumors. Proteins elevated in LGSOC compared to HGSOC tumors (LIMMA adjusted p < 0.05) were enriched for pathways regulating cilium assembly and included a pathway regulating activation of FAK1 by MET, a therapeutic target in LGSOC. Using data from an independent proteomic analysis of LGSOC (n=14) and HGSOC (n=30) tumors, we identified 330 co-altered proteins between LGSOC and HGSOC tumors exhibiting high quantitative correlation (Spearman Rho = 0.803, $P < 2.2 \times 10^{-16}$). Among these, MUC16 (CA125) was identified as significantly elevated (>1.7 $\log_2$-fold) in LGSOC versus HGSOC. Immunohistochemistry analysis of MUC16 verified the increased abundance in LGSOC tissues and identified that MUC16 staining patterns in LGSOC are uniquely apical compared to HGSOC (Fisher's Exact p = 0.001). Our deep proteomic analyses identifies highly-conserved proteome alterations distinguishing LGSOC from HGSOC tumors, including candidates regulating FAK1 signaling as well as a novel identification that MUC16 is elevated and exhibits an apical staining pattern in LGSOC tumors. These findings deepen our molecular understanding of LGSOC and provide unique insights into the regulation of MUC16 in LGSOC tumors.

Introduction

Despite overall improvements in mortality, ovarian cancer is one of the most lethal gynecologic malignancies, with fewer than half of women surviving five years beyond initial diagnosis[1]. In 2024, the American Cancer Society estimates that 19,680 women diagnosed with ovarian cancer and 12,740 (64%) will die from their disease[2]. Epithelial ovarian cancer is the most common histologic subtype of ovarian cancer and includes high-grade serous, low-grade serous, mucinous, clear cell, endometrioid, and carcinosarcoma. High-grade serous ovarian cancer (HGSOC) is the most common histologic subtype and is treated with a combination of cytoreductive surgery and platinum-taxane chemotherapy. Low-grade serous ovarian cancer (LGSOC) is responsible for 2-5% of ovarian cancer and is an overall challenging disease to treat given the high rate of chemoresistance of this disease compared to HGSOC[3].

Historically, the hypothesis was that LGSOC ultimately progressed to HGSOC, but this hypothesis has been challenged and current opinion suggests these two disease processes to be independent[4]. From a molecular standpoint, both HGSOC and LGSOC vary by morphology and somatic mutations. HGSOC is associated with *TP53* mutations as opposed to LGSOC, which often harbors mutations in Kirsten rat sarcoma virus (KRAS), the proto-oncogene B-Raf (BRAF), receptor tyrosine-protein kinase erbB-2 (ERBB2), beta-catenin (CTNNB1), as well as phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform (PIK3CA)[4]. However, co-existing high-grade and low-grade tumors often exhibit the same mutation and lack a TP53 mutation, suggesting there may be a stepwise, non-TP53 related pathogenic process[4]. More importantly, an understanding of genomic differences between these disease subtypes has also introduced new treatments for LGSOC patients [5].

Although the genomic characterization of HGSOC and LGSOC is well established, there

is currently limited proteomic data characterizing these tumors. Some hypotheses suggest that

fallopian tube epithelium may serve as a site of origin for both LGSOC and HGSOC [3,6], which

provides a rationale for its incorporation in the proteomic evaluation of these tumors. As a result,

we sought to perform a deep proteomic investigation evaluating tumors from patients diagnosed

with LGSOC and HGSOC as well as normal fallopian tube tissues from women with benign

disease to better understand the differences in molecular alterations between these tumors and to

determine whether there may be any additional pathways and therapeutic targets to help guide

further treatment paradigms.

Methods

Patient cohort and tissue collections

Formalin-fixed, paraffin-embedded (FFPE) LGSOC and HGSOC tumor and normal fallopian tube tissues were obtained from Inova Fairfax Hospital (Falls Church, VA) under a less than minimal-risk WCG IRB-approved protocol #20122048 with waivers for consent and HIPAA authorization and evaluated under protocol #14-1679 with an exempt determination by the WCG IRB in accordance with the use of de-identified data under US Federal regulation 45 CFR 46.102(f). Benign fallopian tube tissues (FT) (n=12), as well as tumor tissues from LGSOC (n=12) and HGSOC (n=24) patients, were sectioned onto 8 μm polyethylene naphthalate membrane slides and hematoxylin-eosin stained. Equivalent areas of whole tissue from FT and tumor epithelium were collected using laser microdissection, avoiding areas of necrosis and hemorrhage, directly into 50 μL of Optima^{TM} LC/MS Grade water (Fisher Chemical) in microcentrifuge tubes and stored at -80ºC until sample preparation.

Sample preparation for quantitative proteomic analysis

Protein from FFPE tissues was prepared for quantitative mass spectrometry (MS)-based proteomics as previously described[7]. Briefly, tissues were transferred to MicroTubes (Pressure BioSciences, Inc., Easton, MA, USA) including 20 μL of 100 mM TEAB (pH 8.0)/10% acetonitrile and underwent pressure-assisted trypsin digestion employing a barocycler (2320EXT, Pressure BioSciences, Inc.) and a heat-stable form of trypsin (SMART Trypsin, Thermo Fisher Scientific, Inc., Waltham, MA, USA). Peptide digest concentrations were determined using the bicinchoninic acid assay (BCA; Thermo Fisher Scientific, Inc.). Peptides were labeled with isobaric tandem mass tag (TMT) reagents according to the manufacturer's instructions (TMT11^{TM}

Isobaric Label Reagent Set, Thermo Fisher Scientific). Sample multiplexes were reversed-phase fractionated (basic pH) on a 1260 Infinity II offline liquid chromatography system (Agilent Technologies, Inc., Santa Clara, CA, USA) into 96 fractions using a linear gradient of acetonitrile (0.69% min^{-1}) followed by concatenation into 24 pooled fractions. Pooled fractions were lyophilized in a Speed-Vac and stored at -20 ºC until resuspended in 25 mM NH_4HCO_3 and analyzed by liquid chromatography-tandem mass spectrometry (LC-MS/MS).

LC-MS/MS Analysis

Peptide identifications and quantitation for TMT multiplexes were performed as recently described[8]. Liquid chromatography-tandem mass spectrometry (LC-MS/MS) analyses of TMT11 multiplexes were performed on a nanoflow high-performance LC system (EASY-nLC 1200, Thermo Fisher Scientific, Inc.) coupled online with an Orbitrap mass spectrometer (Q Exactive HF-X, Thermo Fisher Scientific, Inc.). Samples were loaded on a reversed-phase trap column (Acclaim^{TM} PepMap^{TM} 100 Å, C-18, 20 mm length, nanoViper Trap column, Thermo Fisher Scientific, Inc.) and eluted on a heated (50 °C) reversed-phase analytical column (Acclaim^{TM} PepMap^{TM} RSLC C-18, 2 μm, 100 Å, 75 μm × 500 mm, nanoViper, Thermo Fisher Scientific, Inc.) by developing a linear gradient from 2% mobile phase A (2% acetonitrile, 0.1% formic acid) to 32% mobile phase B (95% acetonitrile, 0.1% formic acid) over 120 min at a constant flow rate of 250 nL/min. Full scan mass spectra (MS) were acquired using a mass range of *m/z* 400-1600, followed by selection of the top 12 most intense molecular ions in each MS scan for high-energy collisional dissociation (HCD). Instrument parameters were as follows: Full MS: AGC, $3 \times 10e6$; resolution, 60 k; S-Lens RF, 40%; max IT, 45 ms; MS2: AGC, 10e5; resolution, 45 k; max IT, 95 ms; quadrupole isolation, 1.0 *m/z*; isolation offset, 0.2 *m/z*; NCE, 34; fixed first mass, 100 *m/z*;

intensity threshold, $2 \times 10e5$; charge state, 2–4; dynamic exclusion, 20 s with 10 ppm tolerance,

TMT optimization. Global protein-level abundances were generated from peptide spectral matches

identified by searching .raw data files against a publicly available, non-redundant human proteome

database (http://www.uniprot.org/, SwissProt, Homo sapiens, downloaded 12-01-2017 using

Mascot (Matrix Science, v2.6.0), Proteome Discoverer (v2.2.0.388, Thermo Fisher Scientific,

Inc.), and in-house tools using identical parameters as previously described[8].

Bioinformatic Analysis

Differential analyses of TMT-11 data matrixes were performed using the LIMMA package

(version 3.8)[9] in R (version 3.5.2) and Mann-Whitney U rank sum testing in MedCalc (version

19.0.3). Pathway analysis was performed using MSigDB and GSEA[10] considering hallmark and

canonical pathways using default enrichment settings, i.e. significantly enriched pathways

satisfied a false discovery rate < 0.05 or STRING (https://string-db.org/) using default parameters

for Homo Sapiens. Visualizations were generated using SRplot[11] , GraphPad Prism (v.10.0.3) or

MedCalc. Global proteome data for LGSOC and HGSOC validation cohorts were previously

reported[8,12] and HGSOC data prioritized for this analysis was focused on tumors where tissue

source site was designated as primary.

Immunohistochemical analysis of mucin-16 (MUC16)/ cancer antigen 125 (CA125)

LGSOC and HGSOC tissues were sectioned onto charged glass slides, fixed with 100% methanol,

incubated at 5 min at ambient temperature then rinsed with phosphate-buffered saline (PBS).

Tissue was permeabilized by incubating with 0.05% triton-PBS for 15 min and 2.5% normal goat

serum. The slides were incubated overnight at 4 °C with anti-CA125 mouse monoclonal antibody

(Clone M11, Agilent Dako, Santa Clara, CA, USA 1:1000). Dako's Envision diaminobenzidine

(DAB) detection system was used to label bound antibody complexes. Normal lung tissue was

used as the positive tissue control. Following detection, slides were counterstained with

hematoxylin then dehydrated and cover slipped. Stained tissue sections were scanned on an Aperio

ScanScope XT slide scanner (Leica Microsystems, Wetzlar, Germany). Digital images underwent

expert pathology review (PMF).

Results

Quantitative proteomic analyses of whole tissue collections from normal fallopian tube as well as low and high-grade serous ovarian cancers.

We performed multiplexed, quantitative proteomic analyses of whole tissue equivalents collected from low-grade (LGSOC, n=12) or high-grade serous ovarian cancers (HGSOC, n=24) tumors and fallopian tube tissues (FT, n=12) from women with benign disease (Figure 1, Supplementary Table 1). Global proteomic analyses resulted in the quantitation of 7,043 proteins across all patient samples (Supplementary Table 2). Principal component analysis of top variably abundant proteins quantified across all patient samples explained 40.1% (PC1) and 9.4% (PC2) of the variance between FT, LGSOC and HGSOC tissue collections (Figure 2A). Differential analysis of proteins quantified between HGSOC or LGSOC and FT collections identified both co-altered as well as unique protein alterations between these comparisons (Figure 2B, Supplementary Table 3 and 4). Pathway analysis of co-altered proteins elevated in HGSOC and LGSOC compared to FT identified enrichment of top pathways correlating with regulation of protein translation (i.e., 40S/ 60S ribosome and eukaryotic translation elongation) as well as protein metabolism (i.e., response of EIF2AK4 to amino acid deficiency and selenoamino acid metabolism) in tumor samples (Figure 2C, Supplementary Table 5). Pathways enriched among proteins elevated in FT tissues largely correlated with regulation of cilium development as well as organelle biogenesis (Figure 2C, Supplementary Table 5). Pathway analysis of proteins uniquely elevated in HGSOC identified pathways largely associated with cell cycle regulation while pathways enriched for proteins uniquely elevated in LGSOC tumors correlated with pathways regulating epithelial to mesenchymal transition and hematopoietic related signaling (Figure 2D, Supplementary Table 6).

Characterization and validation of proteome alterations between low and high-grade serous ovarian cancers

Differential analysis of proteins quantified in LGSOC and HGSOC tumors identified 785 proteins as significantly altered between these tumor types (Figure 3A). Top protein alterations included proliferation marker protein Ki-67 (MKI67) elevated in HGSOC and multiple collagen isoforms in LGSOC tumors (Figure 3A, Supplementary Table 7). Comparison of these protein alterations with global proteome data recently described for an independent cohort of tumor tissues collected from women diagnosed with LGSOC[12] (n=14) or HGSOC[8] (n=30) tumors identified 330 protein features as co-altered between these disease subtypes that further exhibit high quantitative correlation (Figure 3B, Supplementary Table 8). Pathway analysis of 313 proteins within this subset exhibiting concordant abundance trends showed proteins elevated in LGSOC tumors are enriched for pathways regulating cilium development as well as focal adhesions and, specifically, regulation of focal adhesion kinase 1 (FAK1/ PTK2) by hepatocyte growth factor receptor (MET) (Figure 3C). Proteins elevated in HGSOC tumors showed enrichment of pathways largely regulating cell cycle and cellular mitosis (Figure 3C, Supplementary Table 9). Investigation of putative drug targets among proteins co-altered between LGSOC and HGSOC tumors in both discovery and validation cohorts identified 34 candidates (Supplementary Table 8). Putative drug targets elevated in HGSOC tumors included several proteins known to regulate DNA replication[13], i.e. DNA topoisomerase IIα (TOP2A) and polymerase delta 1 (POLD1) while those elevated in LGSOC tumors included diverse candidates like hepatocyte growth factor receptor (MET), targeted by diverse receptor tyrosine kinase inhibitors[14] as well as mucin-16 (MUC-16), also known as cancer antigen 125 (CA125, Figure 3D). Circulating MUC16/CA125 levels are assessed as part of routine diagnosis and disease monitoring in the clinical management of ovarian cancer[3,6].

To uncover whether other putatively conserved protein candidates elevated in LGSOC tumors are

predicted to interact with MUC16, we investigated these protein features using the STRING

interaction network (string-db.org). We identified nesprin-1 (SYNE1), mucin-5B (MUC5B) and

galecin-3 (LGALS3) as putative interactome partners of MUC16 and all significantly elevated in

LGSOC tumors (Supplementary Table 8). Considering the importance of MUC16 in ovarian

cancer disease biology, we were motivated to investigate this protein alteration in LGSOC and

HGSOC tissues further.

Mucin-16 (MUC16)/ cancer antigen 125 (CA125) is elevated and exhibits a predominantly

apical staining pattern in low versus high-grade serous ovarian cancers

To further investigate elevation of MUC16 protein as a putatively conserved protein alteration in

LGSOC versus HGSOC tumors, we performed immunohistochemistry (IHC) analysis of tissues

for a subset of LGSOC (n=11) and HGSOC (n=24) tumors characterized within our discovery

cohort. A comparison of tumor cells stained positive for MUC16 between LGSOC and HGSOC

tumors showed HGSOC tumors overall have fewer MUC16-positive cells than LGSOC cells

(Figure 4A, Supplementary Table 10). Further analysis of MUC16 staining identified discrete

localization patterns correlating with disease subtypes, where HGSOCs exhibited a more

membranous or combined apical and membranous staining pattern while LGSOCs exhibited a

predominantly apical MUC16 staining pattern (Figure 4B). We further include MUC16 IHC

images for two representative HGSOC cases demonstrating apical and membranous staining as

well as for two LGSOC cases, which exhibit a predominant and intense, apical membrane staining

pattern (Figure 4C).

Discussion

Our deep proteomic analysis of benign fallopian tube tissues as well as low (LGSOC) and high (HGSOC) grade serous ovarian cancers shows these samples can be stratified by variably abundant protein alterations suggesting distinct proteome abundance profiles are characteristic of these tissue types. Differential analysis of tumor (LGSOC and HGSOC) and benign fallopian tube tissue identified that proteins elevated in fallopian tubes were largely enriched for pathways regulating cilium development. Recent evidence from single cell studies has identified several subpopulations of epithelial cell types exist within benign fallopian tube, including ciliated cell types[15,16]. This cell population is considered to reflect a more differentiated subpopulation of fallopian tube epithelial cells emerging from non-ciliated, secretory epithelial cell types[15,16]. The loss of proteins regulating cilium development that we observe in LGSOC and HGSOC tissues is consistent with evidence suggesting secretory epithelial cell types often exhibit greater neoplastic potential than other cell types in fallopian tube epithelium[15]. We also find that proteins elevated in both LGSOC and HGSOC tumors exhibit enrichment of pathways regulating protein translation, specifically ribosome and translational elongation, as well as amino acid metabolism in comparison with benign fallopian tube. Increased ribosome biogenesis and protein translation has been described as being characteristic of ovarian cancer cells and has further been suggested as a possible therapeutic vulnerability in both LGSOC and HGSOC[17,18]. Proteins uniquely elevated in HGSOC tumors compared to benign fallopian tube were largely enriched for pathway regulating cell cycle and mitosis, consistent with the high-grade phenotype of this disease[19]. Proteins uniquely elevated in LGSOC were enriched for pathways regulating epithelial to mesenchymal transition including extracellular matrix organization, beta1 integrin cell surface interactions and hematopoietic-related signaling, i.e. complement and coagulation. Beta1 integrin signaling has been implicated in

regulating migration of ovarian cancer cells in the context of interactions with extracellular matrix

proteins[20]. Our comparison with fallopian tube tissues has identified proteome alterations largely

consistent with the malignant phenotypes of LGSOC and HGSOC tumors in comparison to benign

fallopian tube.

Comparison of LGSOC and HGSOC revealed marked protein alterations between these

disease subtypes, including elevation of proteins correlating with proliferation and DNA

replication in HGSOC. We also observe elevation of diverse extracellular matrix proteins in

LGSOC tumors, including multiple collagen isoforms which is consistent with the enrichment of

epithelial to mesenchymal signaling pathways[21] we also observed in these tumors. Our team

recently described global proteome analysis of tumors collected from women diagnosed with

LGSOC[12] or with HGSOC[8] and we were motivated to compare proteome alterations we have

observed between LGSOC and HGSOC tumors from this study with these independent cohorts to

identify and validate proteome alterations that are highly conserved between these disease

subtypes. This comparison resulted in the identification of a subset of protein alterations exhibiting

high quantitative correlation between LGSOC and HGSOC tumors (Spearman Rho > 0.8). These

features demonstrated that proteins elevated in HGSOC tumors largely correlate with enrichment

of pathways regulating cell cycle and cell mitosis signaling, while those elevated in LGSOC

tumors correlated with intercellular and extracellular matrix interactions, i.e. focal adhesion and

laminin interactions, but also included enrichment of pathways regulating cilium development as

well as regulation of focal adhesion kinase 1 (FAK/ PTK2) by hepatocyte growth factor receptor

(MET). Investigation of protein alterations mapping to this latter pathway shows that MET, but

not FAK, is significantly elevated in LGSOC versus HGSOC tumors (Supplementary Table 8).

MET is a receptor tyrosine kinase that can signal through phosphoinositide 3-kinase (PI3K) or

mitogen activated protein kinase (MAPK) pathways resulting in increased survival and invasive

potential of tumor cells[22,23]. MET has been shown to be elevated in ovarian cancer cells and

activated by increased hepatocyte growth factor production from cancer associated fibroblasts that

is further correlated with a chemoresistant phenotype[24]. FAK has been found to promote

oncogenesis in diverse cancers including ovarian by increasing the invasive potential of tumor

cells through extracellular matrix remodeling as well as by modulating PI3K and MAPK signaling

pathways to promote tumor cell aggressivity[25,26]. Notably, MAPK signaling has been shown to be

hyperactivated in LGSOC tumors[3,26] and targeting MAPK signaling using clinical inhibitors of

MAPK signaling, i.e. MEK inhibitors has led to multiple clinical trials focused on exploiting this

therapeutic vulnerability in LGSOC[3,26]. Recent evidence has shown that activation of FAK can

promote resistance to MEK inhibitors[27] and this has led to clinical trials combining MEK inhibitors

with the FAK inhibitor Defactinib to treat LGSOC, including a phase 2 trial focused on leveraging

this combination therapy for the treatment of recurrent LGSOC (ENGOT-ov60/GOG-3052/RAMP

201, i.e. NCT04625270)[26]. Our findings provide further supportive evidence that elevation of MET

protein abundance as well as enrichment of pathways correlating with FAK signaling is highly

conserved in LGSOC tumors.

Additional investigation of putatively conserved protein alterations between LGSOC and

HGSOC tumors identified mucin-16 (MUC16)/ cancer antigen 125 (CA125) as significantly

elevated in LGSOC versus HGSOC tumor tissues. MUC16 is a transmembrane glycoprotein that

is highly abundant in the female reproductive tract and can be actively shed from ovarian cancer

cells into blood plasma[6]. Circulating MUC16/ CA125 protein levels thus represents a biomarker

that can be measured to assess evidence of disease and is a facet of routine diagnosis and disease

monitoring in the clinical management of ovarian cancer[3,6]. Shedding of MUC16 has been shown

to occur as a product of proteolytic cleavage of the carboxy-terminal ectodomain in the acidifying

Golgi/ post-Golgi[28], possibly by membrane-type 1 matrix metalloproteinase[29] followed by release

of the N-terminal CA125 region[29] into interstitial fluid and ultimately into circulating blood[6]. As

we quantify membrane-type 1 matrix metalloproteinase (MT1-MMP/ MMP-14/ P50281) protein

in our discovery cohort, we compared levels between LGSOC &HGSOC tumors, but we do not

observe this protein as significantly altered between these disease subtypes (MWU p = 0.48).

Recent evidence investigating mechanisms regulating cell surface retention of MUC16 have

implicated N-glycosylation mediated by N-acetylglucosaminyltransferase I (MGAT1) that

promotes increased retention of MUC16 at the cell surface[30]. This study further identified N-

glycosylation of MUC16 promotes interaction with the protein galectin-3 (LGALS3) at the cell

surface[30]. Notably, we identify LGALS3 as significantly elevated in LGSOC versus HGSOC

tumor tissues and, although we quantify MGAT1, we do not see this protein as significantly altered

between LGSOC versus HGSOC tissues (Supplementary Table 7). Considering this evidence in

the context of our findings suggests LGSOC tumor cells perhaps exhibit increased retention of

MUC16 at the cell surface. Circulating CA125 levels have been observed in both LGSOC and

HGSOC patients, although it has been noted that these levels are more commonly lower in LGSOC

patients[3,31]. We also find that circulating CA125 levels were lower in our LGSOC versus HGSOC

patients (Table 1), providing further evidence in support of this hypothesis. A possible mechanism

underlying altered dynamics of MUC16 retention may be due to differential N-glycosylation of

MUC16 in LGSOC tumor cells, possibly due to altered enzymatic activity of MGAT1.

Alternatively, differences in MUC16 expression and shedding could relate to greater cleavage of

the extracellular domain in HGSOC, accounting for the greater shedding of CA125 and decreased

membrane expression. Future studies will focus on investigation of these hypothesis including

comparison of cleaved versus non-cleaved MUC16 levels in LGSOC and HGSOC tumors.

We also observe that LGSOC tumor cells exhibit a predominantly apical staining pattern

for MUC16 in comparison to HGSOC tumor cells. Similar findings were observed in murine

ovarian carcinomas where more intense apical expression of MUC16 was observed in grade 1

tumors[32]. Recent evidence has shown that MUC16 is maintained within apical membranes of

epithelial cells, such as in ocular epithelium[33], and to be dependent on glycosylation[30], which when

considered relative to our findings may explain why we observe increased apical staining patterns

for MUC16 in LGSOC tissues. Beyond its role as a biomarker, MUC16 has been shown to promote

cellular proliferation and invasion of ovarian cancer cells[34] and has previously been shown to

discourage immune surveillance and to be a potential target for immunotherapy[35]. Future efforts

will be focused on not only investigating mechanisms of membrane retention of MUC16 in

LGSOC, but further on improving our understanding of how this characteristic impacts LGSOC

tumor cell biology.

A limitation for the current study includes small sample sizes for normal fallopian tube as

well as LGSOC and HGSOC tissues in our discovery cohort. We aimed to address this limitation

by prioritizing protein alterations between LGSOC and HGSOC tumors that were further validated

within global proteomic data generated for independent cohorts of LGSOC and HGSOC patients.

One may also claim another limitation is that samples were obtained by formalin-fixed, paraffin-

embedded tissue instead of fresh frozen tissue. Nevertheless, comparison of fresh tissue samples

and fixed frozen paraffin embedded samples has not demonstrated differences in peptide

identification and quantification[36]. Strengths of this study include the foundation provided for

future proteomic research, and the contributions of our findings towards improving our

understanding of molecular alterations underlaying ovarian cancer pathogenesis.

Conclusion

Our study contributes to our understanding of global proteome dynamics between normal fallopian

tube tissues and LGSOC and HGSOC tumors and provide insights into the relative abundance and

localization of the ovarian cancer biomarker MUC16/ CA125 in LGSOC and HGSOC tumors that

may have bearing on clinical interpretation of circulating levels of CA125 in these disease

subtypes. Future study is needed to determine the significance of protein alterations in the course

of this disease, and which—if any—proteins may serve as additional therapeutic targets for disease

treatment.

Table 1: Patient and tumor clinicopathological characteristics

Characteristic	High Grade Serous Ovarian Cancer (N = 24)	Low Grade Serous Ovarian Cancer (N = 12)	Benign Fallopian Tube (N = 12)
Mean age (range) –yr	58.1 (30-90)	52.0 (30-73)	45.2 (41-49)
Menopausal status – no. (%)			
Premenopausal	5 (20.8)	6 (50.0)	12 (100.0)
Postmenopausal	19 (79.2)	6 (50.0)	0 (0.0)
Mean BMI (range)	25.6 (22.5 – 30)	25.4 (20.7 – 30.8)	30.1 (19 – 44.1)
Race – no. (%)			
White	13 (54.2)	11 (91.7)	4 (33.3)
Black	7 (29.2)	0 (0.0)	4 (33.3)
Asian	4 (16.7)	1 (8.3)	2 (16.7)
Hispanic	0 (0.0)	0 (0.0)	2 (16.7)
FIGO stage – no. (%)			
I or II	2 (8.3)	3 (25.0)	-
IIIA	3 (12.5)	4 (33.3)	-
IIIB	6 (25.0)	1 (8.4)	-
IIIC	7 (29.2)	4 (33.3)	-
IV	6 (25.0)	0 (0.0)	-
Debulking			
RD0	12 (50.0)	7 (58.3)	-
RD1	9 (37.5)	4 (33.3)	-
Suboptimal	2 (8.3)	1 (8.4)	-
Unknown	1 (4.2)	0 (0.0)	-
Platinum Sensitivity – no (%)			
Sensitive	14 (58.3)	8 (66.6)	-
Resistant	4 (16.7)	1 (8.4)	-
Unknown	6 (25.0)	3 (25.0)	-
Mean CA125 (range)	4917.1 (329 – 50,000)	672.7 (70 – 1576)	-

Data Availability

Proteomic data generated in this study is accessible at the ProteomeXChange under accession PXD050642.

Author Contributions

C.M.T., N.W.B., T.P.C. lead the study. C.M.T, P.F., J.O., G.L.M. performed identification of clinical specimens, C.M.T., G.L.M., K.M.D performed clinical data analysis. K.A.C., J.O., and C.M.T. performed sample collections and molecular extraction. N.W.B., J.O., T.A., B.L.H, generated and analyzed proteomics data. P.M.F. performed pathology review. C.M.T., N.W.B., wrote the manuscript, P.M.F., T.A., B.L.H, K.R., K.K.W, D.G., S.L., A.K.S., R.C.B, K.M.D., N.T.P, G.L.M, T.P.C reviewed the manuscript. All authors reviewed and approved the final manuscript.

Declaration of Interests

Nothing to disclose.

Acknowledgements

Funding for this project was provided from the Uniformed Services University of the Health Sciences from the Defense Health Program to the Henry M Jackson Foundation (HJF) for the Advancement of Military Medicine Inc. Gynecologic Cancer Center of Excellence Program, including awards HU0001-20-2-0033, HU0001-21-2-0027, HU0001-22-2-0016, and HU0001-23-2-0038 (PI: Neil T Phippen, Co-PI: G. Larry Maxwell).

Disclaimer

The views expressed herein are those of the authors and do not reflect the official policy of the Uniformed Services University of the Health Sciences, the Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc., Inova Health System, the Department of Army/Navy/Air Force, Department of Defense, or U.S. Government. Mention of trade names, commercial products, or organizations does not imply endorsement by the U.S. Government.

References

Torre, L. A. *et al.* Ovarian cancer statistics, 2018. *CA Cancer J Clin* **68**, 284-296, doi:10.3322/caac.21456 (2018).

Siegel, R. L., Giaquinto, A. N. & Jemal, A. Cancer statistics, 2024. *CA Cancer J Clin* **74**, 12-49, doi:10.3322/caac.21820 (2024).

Slomovitz, B. *et al.* Low-grade serous ovarian cancer: State of the science. *Gynecol Oncol* **156**, 715-725, doi:10.1016/j.ygyno.2019.12.033 (2020).

Zarei, S. *et al.* Clinicopathologic, Immunohistochemical, and Molecular Characteristics of Ovarian Serous Carcinoma With Mixed Morphologic Features of High-grade and Low-grade Serous Carcinoma. *Am J Surg Pathol* **44**, 316-328, doi:10.1097/PAS.0000000000001419 (2020).

Gershenson, D. M. *et al.* Trametinib versus standard of care in patients with recurrent low-grade serous ovarian cancer (GOG 281/LOGS): an international, randomised, open-label, multicentre, phase 2/3 trial. *Lancet* **399**, 541-553, doi:10.1016/S0140-6736(21)02175-9 (2022).

Giamougiannis, P., Martin-Hirsch, P. L. & Martin, F. L. The evolving role of MUC16 (CA125) in the transformation of ovarian cells and the progression of neoplasia. *Carcinogenesis* **42**, 327-343, doi:10.1093/carcin/bgab010 (2021).

Burdett, N. L. *et al.* Multiomic analysis of homologous recombination-deficient end-stage high-grade serous ovarian cancer. *Nat Genet* **55**, 437-450, doi:10.1038/s41588-023-01320-2 (2023).

Lee, S. *et al.* Molecular Analysis of Clinically Defined Subsets of High-Grade Serous Ovarian Cancer. *Cell Rep* **31**, 107502, doi:10.1016/j.celrep.2020.03.066 (2020).

Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47, doi:10.1093/nar/gkv007 (2015).

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550, doi:10.1073/pnas.0506580102 (2005).

Tang, D. *et al.* SRplot: A free online platform for data visualization and graphing. *PLoS One* **18**, e0294236, doi:10.1371/journal.pone.0294236 (2023).

Wong, K. K. *et al.* Integrated multi-omic analysis of low-grade ovarian serous carcinoma collected from short and long-term survivors. *J Transl Med* **20**, 606, doi:10.1186/s12967-022-03820-x (2022).

Ganai, R. A. & Johansson, E. DNA Replication-A Matter of Fidelity. *Mol Cell* **62**, 745-755, doi:10.1016/j.molcel.2016.05.003 (2016).

Huang, X. *et al.* Targeting the HGF/MET Axis in Cancer Therapy: Challenges in Resistance and Opportunities for Improvement. *Front Cell Dev Biol* **8**, 152, doi:10.3389/fcell.2020.00152 (2020).

Dinh, H. Q. *et al.* Single-cell transcriptomics identifies gene expression networks driving differentiation and tumorigenesis in the human fallopian tube. *Cell Rep* **35**, 108978, doi:10.1016/j.celrep.2021.108978 (2021).

Ghosh, A., Syed, S. M. & Tanwar, P. S. In vivo genetic cell lineage tracing reveals that oviductal secretory cells self-renew and give rise to ciliated cells. *Development* **144**, 3031-3041, doi:10.1242/dev.149989 (2017).

Yan, S. *et al.* The Potential of Targeting Ribosome Biogenesis in High-Grade Serous Ovarian Cancer. *Int J Mol Sci* **18**, doi:10.3390/ijms18010210 (2017).

18   Kang, J. *et al.* Ribosomal proteins and human diseases: molecular mechanisms and
targeted therapy. *Signal Transduct Target Ther* **6**, 323, doi:10.1038/s41392-021-00728-8
(2021).
19   Lopez-Reig, R. & Lopez-Guerrero, J. A. The hallmarks of ovarian cancer: proliferation
and cell growth. *EJC Suppl* **15**, 27-37, doi:10.1016/j.ejcsup.2019.12.001 (2020).
20   Casey, R. C. & Skubitz, A. P. CD44 and beta1 integrins mediate ovarian carcinoma cell
migration toward extracellular matrix proteins. *Clin Exp Metastasis* **18**, 67-75,
doi:10.1023/a:1026519016213 (2000).
21   Yi, B. R., Kim, T. H., Kim, Y. S. & Choi, K. C. Alteration of epithelial-mesenchymal
transition markers in human normal ovaries and neoplastic ovarian cancers. *Int J Oncol*
46, 272-280, doi:10.3892/ijo.2014.2695 (2015).
22   Kim, H. J. Therapeutic Strategies for Ovarian Cancer in Point of HGF/c-MET Targeting.
Medicina (Kaunas) **58**, doi:10.3390/medicina58050649 (2022).
23   Organ, S. L. & Tsao, M. S. An overview of the c-MET signaling pathway. *Ther Adv Med*
Oncol **3**, S7-S19, doi:10.1177/1758834011422556 (2011).
24   Di Renzo, M. F. *et al.* Overexpression of the Met/HGF receptor in ovarian cancer. *Int J*
Cancer **58**, 658-662, doi:10.1002/ijc.2910580507 (1994).
25   Wu, Y. *et al.* Focal adhesion kinase inhibitors, a heavy punch to cancer. *Discov Oncol* **12**,
52, doi:10.1007/s12672-021-00449-y (2021).
26   Zwimpfer, T. A. *et al.* Low grade serous ovarian cancer - A rare disease with increasing
therapeutic options. *Cancer Treat Rev* **112**, 102497, doi:10.1016/j.ctrv.2022.102497
(2023).
27   Taylor, K. N. & Schlaepfer, D. D. Adaptive Resistance to Chemotherapy, A Multi-FAK-
torial Linkage. *Mol Cancer Ther* **17**, 719-723, doi:10.1158/1535-7163.MCT-17-1177
(2018).
28   Das, S. *et al.* Membrane proximal ectodomain cleavage of MUC16 occurs in the
acidifying Golgi/post-Golgi compartments. *Sci Rep* **5**, 9759, doi:10.1038/srep09759
(2015).
29   Bruney, L., Conley, K. C., Moss, N. M., Liu, Y. & Stack, M. S. Membrane-type I matrix
metalloproteinase-dependent ectodomain shedding of mucin16/ CA-125 on ovarian
cancer cells modulates adhesion and invasion of peritoneal mesothelium. *Biol Chem* **395**,
1221-1231, doi:10.1515/hsz-2014-0155 (2014).
30   Taniguchi, T. *et al.* N-Glycosylation affects the stability and barrier function of the
MUC16 mucin. *J Biol Chem* **292**, 11079-11090, doi:10.1074/jbc.M116.770123 (2017).
31   But, I. & Gorisek, B. Preoperative value of CA 125 as a reflection of tumor grade in
epithelial ovarian cancer. *Gynecol Oncol* **63**, 166-172, doi:10.1006/gyno.1996.0301
(1996).
32   Wang, Y. *et al.* MUC16 expression during embryogenesis, in adult tissues, and ovarian
cancer in the mouse. *Differentiation* **76**, 1081-1092, doi:10.1111/j.1432-
0436.2008.00295.x (2008).
33   Mantelli, F. & Argueso, P. Functions of ocular surface mucins in health and disease. *Curr*
Opin Allergy Clin Immunol **8**, 477-483, doi:10.1097/ACI.0b013e32830e6b04 (2008).
34   Felder, M. *et al.* MUC16 (CA125): tumor biomarker to cancer therapy, a work in
progress. *Mol Cancer* **13**, 129, doi:10.1186/1476-4598-13-129 (2014).

35   Zhai, Y. *et al.* MUC16 affects the biological functions of ovarian cancer cells and induces
an antitumor immune response by activating dendritic cells. *Ann Transl Med* **8**, 1494,
doi:10.21037/atm-20-6388 (2020).
36   Bateman, N. W. & Conrads, T. P. Recent advances and opportunities in proteomic
analyses of tumour heterogeneity. *J Pathol* **244**, 628-637, doi:10.1002/path.5036 (2018).

Figure Legends

Figure 1: Workflow supporting quantitative proteomic analyses and validation of proteome alterations in low (LGSOC) and high-grade (HGSOC) serous ovarian cancers. Whole tissue collections from benign fallopian tube, LGSOC and HGSOC tissues was performed using laser microdissection. Tissue collections underwent multiplexed, quantitative proteomic analysis using isobaric tagging, i.e. tandem mass tags (TMT-11), followed by liquid chromatography, tandem mass spectrometry (LC-MS/MS) analysis on an Orbitrap mass spectrometer. Proteome alterations identified in discovery cohorts of LGSOC and HGSOC tissues were validated using global proteome data for independent cohorts of LGSOC[12] (n=14) and HGSOC[8] (n=30) tumors and included the finding that mucin-16 (MUC16) is elevated in LGSOC tumors which was further verified by immunohistochemistry analysis of LGSOC and HGSOC tissues.

Figure 2: Differential proteomic analysis of whole tissue collections from normal fallopian tube with low (LGSOC) and high-grade (HGSOC) serous ovarian cancers. A: Principal component analysis (PCA) of the top 500 most variably abundant proteins quantified across samples; proteins prioritized based on median absolute deviation. B: Comparison of proteins significantly altered between HGSOC or LGSOC and benign fallopian tube collections; proteins reflect LIMMA adjusted p-value < 0.05, fold-change ± 2.0. C: Pathway analysis of proteins co-altered between HGSOC and LGSOC vs benign fallopian tube collections; top hallmark and canonical pathways (gsea-msigdb.org) enriched among proteins elevated (84) or decreased (457) in HGSOC and LGSOC versus benign fallopian tube collections. D: Pathway analysis of proteins uniquely elevated in HGSOC or LGSOC vs normal fallopian tube collections; top hallmark and

canonical pathways (gsea-msigdb.org) enriched among proteins elevated in HGSOC (300) or

LGSOC (158) tumors versus benign fallopian tube collections.

Figure 3: Differential proteomic analysis of low (LGSOC) and high-grade (HGSOC) serous

ovarian cancers and validation of proteome alterations in independent LGSOC and HGSOC

cohorts. A: Volcano plot reflects proteins significantly altered between LGSOC and HGSOC

tumors; proteins reflect LIMMA adjusted p-value < 0.05, fold-change ± 1.5. B: Proteins co-altered

between LGSOC and HGSOC tumors in discovery and validation cohorts; 330 proteins exhibiting

LIMMA adjusted p-value < 0.05, fold-change ± 1.5. C: Pathway analysis of proteins co-altered

between LGSOC and HGSOC tumors in discovery and validation cohorts; top hallmark and

canonical pathways (gsea-msigdb.org) enriched among proteins elevated (144) or decreased (169)

between LGSOC and HGSOC tumors. D: Mucin-16 (MUC16) protein abundance in LGSOC and

HGSOC tumors from discovery or validation cohorts; * = LIMMA adjusted p-value < 0.05, ****

= LIMMA adjusted p-value < 1E-4.

Figure 4: Immunohistochemical analysis of Mucin-16 (MUC16) in low (LGSOC) and high-

grade (HGSOC) serous ovarian cancers. A: Comparison of cells stained positive for MUC16

following immunohistochemical analysis of LGSOC (n=11) and HGSOC (n=24) tissues; p-value

reflects Mann Whitney Rank Sum testing. B: Comparison of MUC16 staining patterns in LGSOC

(n=11) and HGSOC (n=24) tissues; tumor cell populations were identified to exhibit

predominantly apical, membranous or both apical and membranous patterns of MUC16 staining.

C: MUC16 immunohistochemistry stains in tissue sections from two representative LGSOC and

HGSOC patient tumors.

Supplementary Tables

1) Supplementary Table 1 (S1): Discovery cohort details

2) Supplementary Table 2 (S2): Global proteome matrix

3) Supplementary Table 3 (S3): Differential analysis of LGSOC vs Fallopian Tube

4) Supplementary Table 4 (S4): Differential analysis of HGSOC vs Fallopian Tube

5) Supplementary Table 5 (S5): Pathway analysis of proteins altered between HGSOC and

LGSOC vs Fallopian Tube.

6) Supplementary Table 6 (S6): Pathway analysis of proteins uniquely elevated in HGSOC or

LGSOC tumors vs Fallopian Tube.

7) Supplementary Table 7 (S7): Differential analysis of LGSOC vs HGSOC

8) Supplementary Table 8 (S8): Proteins co-altered in LGSOC vs HGSOC between discovery

and validation cohorts.

9) Supplementary Table 9 (S9): Pathway analysis of proteins altered between HGSOC and

LGSOC tumors.

10) Supplementary Table 10 (S10): MUC16 immunohistochemistry analysis

Normal Fallopian Tube, Low (LGSOC) & High-Grade (HGSOC) Serous Ovarian Cancer

Collection of Tissues by Laser Microdissection Followed by Quantitative Proteomics

Sample Multiplexing

Quant.   Ident.

m/z

Conserved Proteome Alterations in LGSOC & HGSOC Tissues

Validation

$\rho = 0.803$, $p < 2.2 \times 10^{-16}$

LGSOC vs HGSOC (Validation, protein logFC)

LGSOC vs HGSOC (Discovery, protein logFC)

LGSOC

HGSOC

MUC-16 IHC

A

B

C   Patient#1   Patient#2