1 **Nanomotif: Leveraging DNA Methylation Motifs for Genome Recovery and Host**
2 **Association of Plasmids in Metagenomes from Complex Microbial Communities**
3

4 Søren Heidelbach[1], Sebastian Mølvang Dall[1], Jeppe Støtt Bøjer[1], Jacob Nissen[2], Lucas N.L.
5 van der Maas[3], Mantas Sereika[1], Rasmus H. Kirkegaard[1], Sheila I. Jensen[3], Sabrina Just
6 Kousgaard[4,5], Ole Thorlacius-Ussing[4,5], Katja Hose[6], Thomas Dyhre Nielsen[2], Mads
7 Albertsen[1]*
8
[1]Center for Microbial Communities, Aalborg University, Denmark
[2]Department of Computer Science, Aalborg University, Denmark
[3]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark,
Denmark
[4]Department of Gastrointestinal Surgery, Aalborg University Hospital, Denmark
[5]Department of Clinical Medicine, Aalborg University, Denmark
[6]Institute of Logic and Computation, TU Wien, Austria
16
*corresponding author.

Abstract

DNA methylation is found across all domains of life but is a rarely used feature in recovery of
metagenome-assembled genomes (MAGs). Recently, Oxford Nanopore introduced all context
methylation detection models. We leveraged this to develop Nanomotif, which identifies and
exploits methylation motifs for enhanced MAG recovery. We demonstrate how Nanomotif
enables database-independent contamination removal from high-quality MAGs and host
association of plasmids directly from Nanopore sequencing data in complex metagenomes.

Main

In all domains of life, genomes are subjected to epigenetic modifications, which directly
influences gene expression, replication, and repair processes[1]. In bacteria, the most common
epigenetic modification is DNA methylation, which primarily acts as a host-defense
mechanism against phages[1]. DNA methylation is facilitated by DNA methyltransferases
(MTases), which recognizes specific DNA sequences, called motifs, and adds a methyl group
to the DNA[1,2]. MTases often appear in restriction-modification systems, where a restriction
enzyme recognizes a specific motif and cleaves the DNA if it lacks methylation. All DNA in the
host must therefore have the correct methylation pattern for it to persist, including mobile
genetic elements[2,3]. This evolutionary arms race has given rise to a great diversity of MTase
recognition sequences[4]. Historically, DNA methylations have been identified using bisulfite
conversions followed by short-read sequencing[1]. In recent years, Pacific Biosciences (PacBio)
and Oxford Nanopore Technologies (ONT) have enabled direct detection of DNA methylations
without the need for pre-treatment[5]. The most common methylations in bacteria are 5-
methylcytosine (5mC), N6-methyladenine (6mA), and N4-methylcytosine (4mC). PacBio was
first to demonstrate *de novo* detection of DNA methylation[5], but currently has a low sensitivity
for 5mC which requires a high sequencing coverage (250x)[6,7]. During 2023-24, ONT
introduced all context methylation detection models making 4mC, 5mC and 6mA methylation
calls readily available with high sensitivity (https://github.com/nanoporetech/dorado). Despite
this, only few efforts have been made to utilize ONT methylation calls for methylation motif
discovery in bacteria[8–10], but none which scales or extends motif discovery to metagenome
sequencing of microbial communities.

In metagenomics, DNA methylation motifs are directly applicable in binning by clustering
contigs, assess contamination in bins, and associate mobile genetic elements to specific
microbial hosts. Previous studies have utilized methylation motif information for metagenomic
binning and association of plasmids[2]. However, these methodologies suffer from the low
PacBio sensitivity for 5mC[2,11] or require whole genome amplification for detection of motifs
using ONT[12]. Building on the recent methylation calling capabilities of ONT sequencing, we
developed Nanomotif, a fast, scalable, and sensitive tool for identification and utilization of
methylation motifs in metagenomic samples. Nanomotif offers *de novo* methylated motif
identification, metagenomic bin contamination detection, association of unbinned contigs to
existing bins, and linking of restriction-modification systems to methylation motifs (Fig. 1A).
57
Nanomotif finds methylated motifs in individual contigs by first extracting windows of 20 bases
upstream and downstream of highly methylated positions (>80% methylated). Motif
candidates are then built iteratively using a beta-Bernoulli model, which evaluates whether the
new candidate is more methylated relative to its originating candidate motif. The motif
candidate search is directed using the KL-divergence from a non-methylated background,
which rapidly guides Nanomotif through the motif search space, greatly decreasing search
time compared to other algorithms (Supplementary note 1, Tab. S1).
65
We investigated a total of 28 monocultures, including 11 REBASE gold standard strains with
known methylation motifs. The 11 monocultures with 75 expected methylation motifs were
further split into 6 strains (29 motifs) for training and 5 strains (46 motifs) for testing. Firstly,
the 29 motifs were used for parameter tuning (Fig. S1). We then benchmarked Nanomotif
against Modkit[10], MicrobeMod[8] and MotifMaker[13] on both training and test monocultures. Only
Nanomotif and Modkit performed satisfactory and were therefore included in further
benchmarks (Fig. S2 & S3). Nanomotif achieved a high recall rate and precision across all
monocultures, identifying 68 out of the 75 expected motifs and 15 other motifs (Fig. 1B). Nine
of the other motifs were closely related to a non-identified expected motif (Fig. 1B-#1-4). In *M.*
ruber, RGAT**4mC**Y was missed, as it is a sub-motif of G**6mA**TC (Fig. 1B-#5). In *A. variabilis*,
4mCYCGRG and ATGC**6mA**T were missed due to only 36 and 74 occurrences, respectively
(Fig. 1B-#6). Lastly, four rare motifs were identified in *M. ruber* (57-474 counts) that likely
represent noise due to increased 5mC false positive rate in high GC% organisms (Fig. 1B-#7
and Fig. S4).
80
To simulate metagenomic conditions, we further benchmarked motif identification by
segmenting the test monoculture genomes to a varying number of fragment sizes and
coverages (Fig. 1C). Nanomotif and Modkit perform similarly on palindromic motifs, which can
be confidently identified in 25 kbp fragments even at 25x coverage. Palindromic motifs are
generally shorter and therefore easier to detect because of their higher frequency and
simplicity. For non-palindromic and bipartite motifs Nanomotif and Modkit have similar
performance on 10-100 kbp, however on 1000 kbp fragments Modkit finds more false-
positives leading to a drop in precision and hence F1-score (Fig. S5 & S6). Lastly, we
benchmarked scalability, where Modkit used 23-40 times more total time compared to
Nanomotif (Tab. S1). Modkit is limited to a single bin per instance, which requires all input files
to be split into single bins beforehand (see Methods). The time spent on preprocessing input
files was not included in the benchmark results.
93
A unique feature of Nanomotif is the scalability to complex metagenomic samples. We
therefore used Nanomotif on five increasingly complex metagenomic samples (Fig. 2A).

Across all metagenomic samples, at least one motif was identified within 87% of metagenome-
assembled genomes (MAGs) above 10x coverage, and within these the median number of
identified motifs was 3. This is more than previously reported in small-scale meta-epigenomic
studies, which only identified methylation motifs in approximately 50% of MAGs using
PacBio[14,15].
101
102 Building on the motif discovery algorithm, we developed three modules for Nanomotif, which
103 uses the motif methylation pattern; MAG contamination detection, inclusion of unbinned
104 contigs, and linking of motifs to the responsible methyltransferases.
105
Current MAG contamination evaluation tools rely on lineage-specific markers derived from
genome databases[16–18], however, it is a difficult task as the databases are far from complete,
and exceptions exist even within closely related organisms. To enable de novo contamination
detection in MAGs, we leveraged Nanomotif to identify methylation motifs and then used
ensemble clustering on the methylation pattern of bins (see methods). The 28 monocultures
were used to benchmark the contamination module by fragmenting the monocultures into one
1600 kbp fragment and several 20 kbp fragments and then randomly assigning 20 kbp
fragments to other bins (Fig. 2B, see methods). Nanomotif was able to achieve high sensitivity
and precision with a mean of 89% and 91%, respectively. Most monocultures had near perfect
contamination removal across all benchmarks (Fig. S7). We then applied the contamination
detection module to the five real metagenomes of increasing complexity. The median number
of contaminants in MAGs where at least one contaminant was detected was 1-2 for HQ MAGs.
For example, bin 1.169 (HQ MAG) from the anaerobic digester (Fig. 2C) included contig_6001
(80 kbp) that completely lacked CAAAA**6mA** and G**6mA**TC methylation, despite the remaining
bin being methylated at >75% in these motifs. In total, 196 contaminants were removed across
90 MAGs from the complex communities (Fig. S8). Each putative contaminant was manually
reviewed, and in 84 out of 90 MAGs, the removal appeared accurate based on the methylation
pattern, which matches the precision observed in the benchmark. This indicates a high
potential for methylation to serve as a powerful post-binning cleanup, especially as this
information is directly available for all new Nanopore sequencing projects.
126
The Nanomotif contig inclusion module assigns unbinned contigs to existing bins by training
a linear discriminant analysis model, random forest, and k-neighbors classifier on the
decontaminated bins (see methods). In case all three classifiers agree with a joint mean
probability >0.8, the contig is assigned to the bin. Nanomotif achieved a high precision of 96%
and moderate recall of 66% across the 28 fragmented monocultures described above (Fig.
S9). In the five real complex metagenomic samples, the include module added a median of
1–4 contigs per bin for HQ MAGs. Associating mobile genetic elements with MAGs is of major
importance as these can carry vital functionality[19]. This can be very difficult for traditional
binners due to large variation in coverage or GC-content from the host but should be possible
if a unique methylation motif signal is present. For example, using methylation motifs three
contigs from the Simple Fecal sample were assigned to bin 1.7 and classified as two plasmids
and a virus (Fig. 2d-e). It should be noted that the inclusion module is not a binner and
assignments should be considered putative as methylation motif patterns can be shared
across MAGs, which is also reflected in our efforts to prioritize precision over recall.
141
142 Restriction-modification (RM) systems are often substantial obstacles to genetic
143 transformation, which pose a significant barrier for the implementation of novel bacteria as cell
144 factories[20]. Circumventing these systems through RM system evasion or through heterologous

expression of the methyltransferases in the cloning host (RM system mimicking) has shown to increase transformation efficiency significantly[20,21]. Therefore, we developed the Nanomotif MTase-linker module, which links methylation motifs to their corresponding MTase and, when present, their entire RM system (Fig. S10, supplementary note 2 & 3, and supplementary data 2). Across 11 monocultures, 52 orphan MTases and 29 RM-systems (exclusive type IV) were identified. 19 RM-systems were associated with an active methylation motif, and a total of 42 out of 71 detected motifs could be linked to a single MTase gene or RM-system with high confidence. Across 549 recovered HQ MAGs from five metagenomic samples, nanomotif identified 3,123 MTase genes, of which 1,297 were associated with RM-systems. For 76% of the detected motifs, at least one candidate MTase with matching methylation characteristics was identified within the same genome. Additionally, Nanomotif, successfully generated a high-confidence set of target motif annotations for 232 MTases. Hence, Nanomotif has the potential to drastically increase the number of putative links between motifs and MTase genes, thereby vastly improving the molecular toolbox and the RM-system databases.

With Nanomotif, *de novo* motif discovery is now seamlessly possible with standard Nanopore sequencing, even for short and low coverage contigs from complex metagenomes. We provide simple implementations that utilize these motifs for robust identification of putative contamination in MAGs, association of mobile genetic elements to hosts, and linking of motifs to restriction-modification systems. This greatly enhances the resolution of metagenomic investigations, opening the possibility of linking extrachromosomal DNA elements to the host.

Data availability

Sequencing data generated during the current study is available in the European Nucleotide Archive (ENA) repository, under the accession number PRJEB74343. Assemblies, bins, and output from Nanomotif are available at https://doi.org/10.5281/zenodo.10964193.

Code availability

Nanomotif is available at https://github.com/MicrobialDarkMatter/nanomotif. Code for reproducing figures and supplementary resources can be found at https://github.com/SorenHeidelbach/nanomotif-article.

Acknowledgements

The study was funded by grants from VILLUM FONDEN (130690, 50093), the Poul Due Jensen Foundation (Microflora Danica) and the European Research Council (101078234). We further acknowledge the Novo Nordisk Foundation within the framework of the Fermentation-based Biomanufacturing Initiative (FBM), (Grant no. NNF17SA0031362), and the Novo Nordisk Foundation (Grant no. NNF20CC0035580).

Ethics

The simple fecal sample was collected as part of a study registered at ClinicalTrials.gov (Trial number NCT04100291). The study adhered to the Good Clinical Practice requirements and the Revised Declaration of Helsinki. The participant provided signed written informed consent to participate and allowed for the sample to be used in scientific research. Consent could be withdrawn at any time during the study period. Conduction of the study was approved by the Regional Research Ethics Committee of Northern Jutland, Denmark (project number N-20150021). The complex fecal sample was collected at Aalborg University with consent from the provider to be used in this study.

189
Fig. 1: Nanomotif overview and benchmark. a, Overview of Nanomotif functionality. White
boxes on the top row are required inputs for Nanomotif, colored boxes are Nanomotif modules.
b, Motif identification using Nanomotif in 11 monocultures with 75 known motifs
(*monocultures used only for testing). For each motif three values are shown: the number of
occurrences in the genome, the percent of motif positions with methylation >70% and the
median methylation of motif positions. Motifs are grouped by whether or not they were
expected to be found or not (See supplementary data 1.a). Seven cases of faulty motif
identification have been annotated 1-7. **c,** Benchmarking of palindrome, bipartite, and non-
palindromic motif identification with Nanomotif and Modkit[7] across different coverages and
reference lengths. The number of expected motifs for each motif type is given in the header.
200

Fig. 2: Nanomotif MAG contamination detection and association of mobile genetic
elements. a, Sample stats from binning and the Nanomotif modules. Only values for bins
above 10x coverage are reported. *The median is reported for bins with at least one motif
identified. **The median is reported for bins where contaminants were removed or contigs
included. **b,** Nanomotif detect_contamination benchmark metrics from 50 benchmark
datasets. 28 monocultures at 40x coverage were fragmented into one 1600 kbp fragment and
several 20 kbp fragments. One randomly selected fragment was randomly assigned to another
bin in each benchmark dataset. **c,** Example of contamination removal (red) from a HQ MAG
from an anaerobic digester. **d,** GC% and coverage of bin.1.7 in the simple fecal sample (blue).
Three contigs, predicted as two plasmids and one virus, are assigned to the bin with the
Nanomotif include_contigs module. **e,** Methylation profile of the HQ bins in the simple fecal
sample and highlighted plasmid & viral contigs assigned to bin.1.7.

Materials And Methods

Sampling

Escherichia coli K-12 MG1655 (labcollection), *Meiothermus ruber* 21 (DSM 1279), and *Parageobacillus thermoglucosidasius* DSMc 2542 were grown overnight in LB, DSMZ 256 Thermus ruber medium, and SPY medium, respectively. Genomic DNA for *Desulfobacca acetoxidans* ASRB2 (DSM 11109), *Sphaerobacter thermophilus* S 6022 (DSM 20745), *Thermanaerovibrio acidaminovorans* Su883 (DSM 6589), *Kangiella aquimarina* SW-154 (DSM 16071), *Anabaena variabilis* PCC 7120 (DSM 107007), *Pelobacter carbinolicus* Bd1 (DSM 2380) and *Salmonella bongori* 1224.72 (DSM 13772) was ordered from Leibniz-Institute DSMZ. Raw pod5 files for *Shewanella oneidensis* MR-1*, Cellulophaga lytica* Cy l20, LIM -21 (DSM 7489), *Kangiella aquimarina SW-154* (DSM 16071)*, Zymomonas mobilis* subsp. pomaceae Barker I and the remaining 15 monocultures used for contamination and inclusion benchmarks were acquired from s3://cultivarium-sequencing/MICROBEMOD-DATA-NOV2023/pod5/[8]. ZymoHMW (D6322), ZymoOral (D6332), ZymoGut (D6331) and ZymoFecal (D6323) were ordered from ZymoBIOMICS. The simple fecal sample was collected at Aalborg University Hospital at the Department of Gastrointestinal Surgery as part of a clinical trial (ClinicalTrials.gov NCT04100291)[22]. The complex fecal sample was collected at Aalborg University with consent from the provider. Sampling of the anaerobic digester sludge has been described elsewhere[23]. Sampling of soil has been described elsewhere[24].

DNA Extraction

DNA from cell pellets of overnight grown cultures of *E. coli* K-12 MG1655 and *M. ruber* 21 was extracted with the PureLink Genomic DNA mini kit (Invitrogen, Thermo Fisher Scientific, USA) following manufacturer's instructions with final elution in DNAse/RNAse free water. DNA from cell pellets of *P. thermoglucosidasius (*DSM 2542) was extracted with the MasterPure Gram positive DNA purification kit (Biosearch Technologies (Lucigen)), according to manufacturer's instructions with a 60 min incubation step and final elution in DNAse/RNAse free water. DNA from ZymoOral (D6332), ZymoGut (D6331) and ZymoFecal (D6323) was extracted with the DNeasy PowerSoil Pro kit according to manufacturer's instructions and suppliers suggestions. DNA from the simple fecal sample was extracted with the DNeasy PowerSoil Pro kit as described previously[25]. DNA from Complex fecal sample was extracted using DNeasy PowerSoil Pro kit according to manufacturer's instructions. DNA was extracted from the anaerobic digester as described previously[23].

Sequencing

All samples were sequenced on the Promethion24 using the R10.4.1 nanopore. Anaerobic digester, complex fecal and ZymoFecal (D6323) were prepared with SQK-LSK114. Monocultures, ZymoOral (D6332), ZymoGut (D6331) were prepared with SQK-RBK114.24. ZymoHMW (D6322) was prepared with the SQK-NBD114-24 ligation kit. Sequencing of simple fecal is described elsewhere[22]. Sequencing of soil is described elsewhere[24]. All samples were basecalled with Dorado v0.8 using the dna_r10.4.1_e8.2_400bps_sup@v5.0.0 model and DNA methylation was called with the respective v2 methylation models for 4mC_5mC and 6mA.

Assembly and binning

All samples were assembled and binned using the mmlong2-lite v1.1.0 pipeline available at [26]. Briefly, metaFlye (v2.9.4)[27] is used for assembly and eukaryotic contigs are removed with

Tiara (v1.0.3)[28] before assembly coverage is calculated with read mapping via minimap2
(v2.28)[29]. Binning is performed iteratively as an ensemble using SemiBin2 (v2.1.0)[30],
MetaBat2 (v2.15)[31], VAMB (v3.0.3)[32], and COMEbin (v1.0.4)[33] whereafter the best bin is
chosen with DAS tool (v1.1.3)[34]. MAG quality was classified according to the MIMAG definition
(Bowers et al., 2017). Completeness and contamination were evaluated with CheckM2 while
rRNA and tRNA genes were found with barrnap (v0.9, https://github.com/tseemann/barrnap)
and tRNAscan-SE (v2.0.16)[35], respectively.

Methylation pileup

267   Reads with methylation calls were mapped to the assembly using minimap2 v2.24[29] using
268   default settings. Nanopore's modkit v0.4.0 (https://github.com/nanoporetech/modkit) was
269   used to generate the methylation pileup from mapped reads using default settings.

Motif identification

MicrobeMod v1.0.3 with default settings was used for all motif identification experiments.
motifMaker (smartlink v13.1.0) with default settings was used for all motif identification
experiments. Modkit pileup is not directly compatible with motifMaker and had to be converted
to the same format as the output of ipdSummary. As the goal was a comparison of the motif
identification algorithm, we extracted generally methylated positions (>70% methylated) and
generated a GFF formatted file similar to the output of ipdSummary, marking all extracted
positions with high Q-score and IPDRatio. Modkit v0.4.0 was used for all motif identification
experiments. Default parameters were used for full genomes and scalability experiment. The
setting --min-sites 20 was used for benchmark with lowered coverage and fragmentation of
reference. This was done for fair comparison to Nanomotif minimum motif count of 20.
281
Nanomotif v0.5.0 was used for all experiments. Nanomotif motif discovery algorithm has two
main submodules, "find-motifs" and "bin-consensus". All subcommands are gathered in a
parent command "motif_discovery", which was executed with the following arguments for all
samples: threshold_methylation_confident=0.8,   threshold_methylation_general=0.7,
search_frame_size=41,   threshold_valid_coverage=5,   minimum_kl_divergence=0.05,
min_motif_score=0.2. "find-motifs" identifies motifs in contigs, referred to as directly identified
motifs. This is done using a greedy search and candidates are selected based on a Beta-
Bernoulli model, where each motif occurrence is Bernoulli trial, being a success if the fraction
of methylation of reads at the position is above a predefined threshold (default 0.70). The Beta-
Bernoulli was chosen in order to include uncertainty in the motif scoring process, instead of a
point estimate. The exact steps performed for motif identification is outlined in the pseudo
code below. For full details see supplementary note 1.
294

```
295    INPUT:
296        assembly - List of contigs sequences
297        modkitPileup - Methylation pileup data
298    OUTPUT:
299        Identified methylated motifs
300    BEGIN
301    |  Initialize identifiedMotifs as an empty list
302    |  FOR sequence IN assembly DO
303    |  |  FOR sequencePosition IN sequence DO
304    |  |  |  IF coverage(sequencePosition) < coverageThreshold (default: 5) THEN
305    |  |  |  |  Mark position as NA
306    |  |  |  ELSE IF fractionModified(sequencePosition) < methylatedThreshold (default: 0.7) THEN
307    |  |  |  |  Mark position as not methylated
308    |  |  |  ELSE
309    |  |  |  |  Mark position as methylated
310    |  |  |  |  IF fractionModified(position) >= confidentlyMethylatedThreshold (default: 0.8) THEN
311    |  |  |  |  |  Mark position as confidently methylated
312    |  |  Initialize seedMotif based on methylation type (C for 5mC, C for 4mC, A for 6mA)
313    |  |  currentMotif = seedMotif
314    |  |  sampledSeqs = Extract n sequences randomly from sequence (default n: 10,000)
315    |  |  sampledPPM = Positional nucleotide probability matrix of sampledSeqs
316    |  |  WHILE stopping criteria NOT met DO
317    |  |  |  methylatedSeqs = Extract sequences at confidently methylated positions (default window: 41)
318    |  |  |  Remove from methylatedSeqs any sequences that match any motif in identified motifs
319    |  |  |  WHILE stopping criteria NOT met DO
320    |  |  |  |  Initialize motifCandidates as an empty list
321    |  |  |  |  methylatedPPM = Calculate positional nucleotide probability matrix of methylatedSeqs
322    |  |  |  |  FOR position IN methylatedPPM DO
323    |  |  |  |  |  Compute KL-divergence from methylatedPPM[, position]  to sampledPPM[, position]
324    |  |  |  |  |  IF KL-divergence > threshold (default: 0.05) THEN
325    |  |  |  |  |  |  Identify valid bases WHERE
326    |  |  |  |  |  |  |  - methylatedPPM[, position] > 25%
327    |  |  |  |  |  |  |  - methylatedPPM[, position] > sampledPPM[, position]
328    |  |  |  |  |  |  FOR baseCombination IN validBases DO
329    |  |  |  |  |  |  |  newMotif = expand currentMotif with baseCombination at position
330    |  |  |  |  |  |  |  add newMotif to motifCandidates
331    |  |  |  |  FOR motifCandidate IN motifCandidates DO
332    |  |  |  |  |  Compute Beta-Bernoulli posterior parameters
333    |  |  |  |  |  Compute score and priority
334    |  |  |  |  Update currentMotif to lowest-priority motifCandidate
335    |  |  |  |  IF highestScoringMotif.score < currentMotif.score THEN
336    |  |  |  |  |  highestScoringMotif = currentMotif
337    |  |  |  |  Subset methylatedSeqs to sequences matching new currentMotif
338    |  |  |  |  Stopping criteria:
339    |  |  |  |  |  highestScoringMotif.score not improved for n round (default: 10)
340    |  |  |  IF highestScoringMotif.score > 0.2 THEN
341    |  |  |  |  Add highestScoringMotif to identifiedMotifs
342    |  |  |  |  Exclude sequences containing identified motifs
343    |  |  Stopping criteria:
344    |  |  |  - 25 low-scoring motifs dropped
345    |  |  |  - <1% methylation sequences remain
346    |  |  Remove motifs contained within more generic motifs
347    |  |  Remove motifs with isolated bases
348    |  |  mergeableMotifs  = motifs with similar sequences (Hamming distance ≤ 2):
349    |  |  FOR motifPair IN mergeableMotifs DO
350    |  |  |  IF methylation of merged motif ≥ methylation pre-merge motifs THEN
351    |  |  |  |  Accept merged motif
352    |  |  |  ELSE
353    |  |  |  |  Retain pre-merge motifs
354    |  |  Identify complementary motifs
355    |  |  Add processed motifs for contig to identifiedMotifs
356    |  RETURN identifiedMotifs
357    END
358
```

359   After identification of motifs in each contig "bin-consensus" evaluates the full set of identified

360   motifs in the contigs belonging to the bin. It performs motif merging like the post processing

361 steps in the motif identification algorithm, but for the bin motif set. Then motifs which are not
362 methylated more than 25% methylated in 75% of the bin are removed.

Motif Identification Benchmark

Motif identification was benchmarked using motifs identified in 11 monocultures with known
methylated motifs. Full set of expected motif and motif evidence is available in supplementary
data 1.a. Six monocultures were used for parameter justification of the "find-motifs" algorithm;
Shewanella oneidensis MR-1*, Kangiella aquimarina* DSM 16071*, Anabaena variabilis* ATCC
27893*, Escherichia coli* K-12 substr. MG1655*, Meiothermus ruber* DSM 1279*, Zymomonas
mobilis* subsp. pomaceae. A grid search was performed over the three most important
parameters, to justify the final parameter settings (Fig. S1). The algorithm performance is
stable across the selected parameters, and the final set parameters were chosen to increase
sensitivity in metagenomic settings. Five monocultures were used for testing; *Desulfobacca
acetoxidans* DSM 11109*, Salmonella bongori* NCTC 12419, *Sphaerobacter thermophilus*
DSM 20745, *Pelobacter carbinolicus, Thermanaerovibrio acidaminovorans* DSM 6589.
Testing monocultures were not seen during tuning.

Benchmark metrics were calculated by comparing identified motif with expected motif in
monocultures. If an identified motif matches an expected motif exactly, it counts as a true
positive. False positives are counted as motifs not in the expected motif set. False negatives
are counted as motifs in the expected set which are not identified. Both forward and reverse
complement are counted as a motif for motifs which are not palindromic. For benchmarks,
precision, recall and F1-score is reported.

Reduced information benchmarking was performed across two parameters; read coverage
(10, 25, 50, 100x) and contig size (10, 25, 50, 100, 1000 kbp). Read coverage affects false
positive and false negative in calling of generally methylated positions, as lower coverage is
more sensitive to non-systematic false positive and false negative calls at the reads level.
Lower coverage was achieved using Rasusa[28] by subsetting the total length of reads to a
multiple of the assembly length of the respective benchmarking organisms. As contig size is
proportional with motif occurrences, smaller contigs will have fewer motif observations,
thereby less information for motif identification. Differing contig sizes were created by chunking
the reference genome of the monoculture to fix sized windows using SeqKit2 (v2.5.1)[36]
(windows were not allowed to overlap). Up to 20 chunks from the procedure for each
monoculture were included in the benchmark. To fairly compare across fragmentation lengths,
we reduced the minimum required motif observations for modkit to 20, the same requirements
Nanomotif utilises for motif identification in contigs.

To benchmark execution time of Modkit on metagenomic samples, we split the pileup file into
separate files, each containing the information related to the contigs of a single bin. Then the
assembly was split into fasta files, each containing the contig sequences related to a single
bin. Execution time for preprocessing of files was not included in the reported run time or CPU
hours. To identify motifs in a bin, we executed Modkit on the files corresponding to a single
bin. Modkit was executed with a single thread, and parallelly executed in 144 instances.

Contamination detection

Contamination is evaluated using "nanomotif detect_contamination" which uses ensemble
clustering. In case all four clustering methods, HDBSCAN, Gaussian Mixture Model,

Agglomerative Clustering, and spectral clustering marks a contig as an outlier, the contig is
marked as contamination.
Firstly, motif methylation is calculated as follows: The mean read methylation for each motif
observation is calculated for motif observations with at least 3 read mappings. The median
value of these is then reported as the motif methylation. This methylation value was more
robust for smaller contigs compared to the mean of means. Before clustering, motif
methylation is filtered for each contig if the product of number of motif observations and the
mean read coverage is less than 24. This way the methylation value of a contig with a one
motif observation is trusted if the read coverage is at least 24 or the contig has at least 8 motif
observations. After filtering, contigs with missing motif values are imputed with the bin mean
and PCA is performed to reduce dimensions while retaining 90% of variance. Contigs are then
clustered with HDBSCAN (min_samples=3, min_cluster_size=2), Gaussian mixture model
(n_components = n_bins, covariance_type="full"), Agglomerative clustering (n_clusters =
n_bins, affinity = "nearest_neighbors") and spectral clustering (n_clusters = n_bins). For each
clustering method, the bin cluster is the cluster with the largest fraction of the bin length which
must constitute at least 85% of the total bin length. In case all methods agree a contig does
not belong to the bin, the contig is flagged as a contaminant. Contamination contigs are then
assigned to the "unbinned" pool.

Include contigs

The "nanomotif include_contigs" module will attempt to assign unbinned contigs to the bin
after decontamination. The include_contigs module uses an ensemble of supervised machine
learning techniques; random forest (n_estimators=100), linear discriminant analysis (solver =
"svd"), and k-nearest neighbors classifier (n_neighbors = 3) classifier to assign unbinned
contigs to bins. Firstly, the three classifiers are trained on the binned contigs after
dimensionality reduction (see contamination detection). Missing motif observations in
unbinned contigs are then imputed with a pseudo value randomly chosen between 0 and 0.15,
and projected into the lower dimensional space using the binned conversion matrix. A contig
is assigned to a bin if all three classifiers agree and the mean probability of the three classifiers
is above 0.80.

Contamination and inclusion benchmark

A synthetic benchmark dataset was constructed for developing and evaluating the
contamination and inclusion module. 28 monocultures (see Fig S6) were sampled to 40x
coverage and fragmented into 20 kbp fragments. For each fragmented monoculture, a single
long 1600 kbp contig was reconstructed by stitching together 80 consecutive 20 kbp
fragments, while the remaining fragments were retained unaltered. Nanomotif were then used
to find motifs anew and contamination and inclusion were evaluated using the found motifs.
To evaluate the contamination module 50 datasets were created where one randomly chosen
contig from each monoculture was randomly added to another. For the inclusion benchmark
50 datasets were created where a random contig was removed from each monoculture. The
1600 kbp contig was not shuffled or removed.

MTase-Linker

The Nanomotif MTase-linker module initially uses Prodigal v.2.6.3[37] for protein-coding gene
prediction (default settings) followed by DefenseFinder v1.2.0[38] to predict MTases and related
RM-system genes. The output file defense_finder_hmmer.tsv is filtered for all RM-related
MTase hits. When a single gene has several model hits, the model that yields the highest

score is selected. The output file defense_finder_systems.tsv is used to determine whether
the identified MTase hit is part of a complete defense system. MTase hits associated with non-
methylation-mediated defense systems are excluded. Additionally, RM type IIG MTase hits
not identified by DefenseFinder as part of a complete RM system are also removed.
Using hmmer (with parameter –cut_ga) the predicted MTase protein sequences are queried
against a set of hidden markov models (PF01555.22, PF02384.20, PF12161.12, PF05869.15,
PF02086.19, PF07669.15, PF13651.10, PF00145.21) from the PFAM database[39], to predict
the modification type (5mC or 6mA/4mC). Furthermore, to infer the probable target recognition
motif, the MTase protein sequences are queried using BLASTP (Blast v.2.14.1) against a
custom database of methyltransferases with known target recognition motif from REbase[40].
We employ a threshold of 80% sequence identity and 80% query coverage to confidently
predict the target recognition motif. Lastly, the RM system, RM sub-type, mod-type, and
predicted motif information for each methyltransferase gene are used to link methylation motifs
to the genes. The pipeline identifies high confidence MTase-motif matches, labeled as "linked",
through either a precise match between the predicted motif and the detected motif or when a
single gene and a single motif share a similar combination of methylation features, which are
unique within a MAG. When a high confidence match cannot be elucidated, the MTase-Motif-
linker assigns feasible candidate genes, with the corresponding motif type and modification
type, for each motif.

MGE classification
Contig were labeled as Mobile genetic elements (MGE) when classified as a viral or plasmid
by geNomad (v1.7.0) with a score >0.75, had a mean coverage above 10x and a minimum
length of 10kbp.
475
476
477

References

1. Seong, H. J., Han, S.-W. & Sul, W. J. Prokaryotic DNA methylation and its functional roles. *J. Microbiol.* **59**, 242–248 (2021).

2. Beaulaurier, J. *et al.* Metagenomic binning and association of plasmids with bacterial host genomes using DNA methylation. *Nat. Biotechnol.* **36**, 61–69 (2018).

3. Seong, H. J., Roux, S., Hwang, C. Y. & Sul, W. J. Marine DNA methylation patterns are associated with microbial community composition and inform virus-host dynamics. *Microbiome* **10**, 157 (2022).

4. Oliveira, P. H. & Fang, G. Conserved DNA Methyltransferases: A Window into Fundamental Mechanisms of Epigenetic Regulation in Bacteria. *Trends Microbiol.* **29**, 28–40 (2021).

5. Clark, T. A. *et al.* Characterization of DNA methyltransferase specificities using single-molecule, real-time DNA sequencing. *Nucleic Acids Res.* **40**, e29–e29 (2012).

6. Liu, J.-H. *et al.* Bacmethy: A novel and convenient tool for investigating bacterial DNA methylation pattern and their transcriptional regulation effects. *Imeta* **3**, e186 (2024).

7. Tue Kjærgaard Nielsen *et al.* Detection of nucleotide modifications in bacteria and bacteriophages; strengths and limitations of current technologies and software. *Mol. Ecol. (2022).*

8. Crits-Christoph, A., Kang, S. C., Lee, H. H. & Ostrov, N. MicrobeMod: A computational toolkit for identifying prokaryotic methylation and restriction-modification with nanopore sequencing. *bioRxiv* (2023) doi:10.1101/2023.11.13.566931.

9. Tidwell, A. K., Faust, E., Eckert, C. A., Guss, A. M. & Alexander, W. Discovering methylated DNA motifs in bacterial nanopore sequencing data with MIJAMP. *bioRxiv* 2024.08.14.607972 (2024) doi:10.1101/2024.08.14.607972.

10. *Modkit: A Bioinformatics Tool for Working with Modified Bases*. (Github).

11. Wilbanks, E. G. *et al.* Metagenomic methylation patterns resolve bacterial genomes of unusual size and structural complexity. *ISME J.* **16**, 1921–1931 (2022).

12. Tourancheau, A., Mead, E. A., Zhang, X.-S. & Fang, G. Discovering multiple types of

DNA methylation from bacteria and microbiome using nanopore sequencing. *Nat.*

Methods **18**, 491–498 (2021).

13. Li, T., Zhang, X., Luo, F., Wu, F.-X. & Wang, J. MultiMotifMaker: A Multi-Thread Tool for

Identifying DNA Methylation Motifs from Pacbio Reads. *IEEE/ACM Trans. Comput. Biol.*

Bioinform. **17**, 220–225 (2020).

14. Hiraoka, S. *et al.* Diverse DNA modification in marine prokaryotic and viral communities.

Nucleic Acids Res. **50**, 1531–1550 (2022).

15. Hiraoka, S. *et al.* Metaepigenomic analysis reveals the unexplored diversity of DNA

methylation in an environmental prokaryotic community. *Nat. Commun.* **10**, 159 (2019).

16. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

17. Orakov, A. *et al.* GUNC: detection of chimerism and contamination in prokaryotic

genomes. *Genome Biol.* **22**, 178 (2021).

18. Chklovski, A., Parks, D. H., Woodcroft, B. J. & Tyson, G. W. CheckM2: a rapid, scalable

and accurate tool for assessing microbial genome quality using machine learning. *Nat.*

Methods **20**, 1203–1212 (2023).

19. Frost, L. S., Leplae, R., Summers, A. O. & Toussaint, A. Mobile genetic elements: the

agents of open source evolution. *Nat. Rev. Microbiol.* **3**, 722–732 (2005).

20. Johnston, C. D. *et al.* Systematic evasion of the restriction-modification barrier in

bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 11454–11459 (2019).

21. Yasui, K. *et al.* Improvement of bacterial transformation efficiency using plasmid artificial

modification. *Nucleic Acids Res.* **37**, e3 (2009).

22. Kousgaard, S. J. *et al.* The effect of non-pooled multi-donor faecal microbiota

transplantation for inducing clinical remission in patients with chronic pouchitis: Results

from a multicentre randomised double-blinded placebo-controlled trial (MicroPouch). *J.*

Crohns. Colitis (2024)

23. Sereika, M. *et al.* Oxford Nanopore R10.4 long-read sequencing enables the generation

of near-finished bacterial genomes from pure cultures and metagenomes without short-read or reference polishing. *Nat. Methods* **19**, 823–826 (2022).

24. Singleton, C. M. *et al.* Microflora Danica: the atlas of Danish environmental microbiomes. *bioRxiv* (2024) doi:10.1101/2024.06.27.600767.

25. Jensen, T. B. N., Dall, S. M., Knutsson, S., Karst, S. M. & Albertsen, M. High-throughput DNA extraction and cost-effective miniaturized metagenome and amplicon library preparation of soil samples for DNA sequencing. *PLoS One* **19**, e0301446 (2024).

26. Sereika, M. Recovery of novel microbial genomes. (Aalborg University, 2024).

27. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads using repeat graphs. *Nat. Biotechnol.* **37**, 540–546 (2019).

28. Karlicki, M., Antonowicz, S. & Karnkowska, A. Tiara: deep learning-based classification system for eukaryotic sequences. *Bioinformatics* **38**, 344–350 (2022).

29. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).

30. Pan, S., Zhao, X.-M. & Coelho, L. P. SemiBin2: self-supervised contrastive learning leads to better MAGs for short- and long-read sequencing. *Bioinformatics* **39**, i21–i29 (2023).

31. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

32. Nissen, J. N. *et al. Binning Microbial Genomes Using Deep Learning.* http://biorxiv.org/lookup/doi/10.1101/490078 (2018).

33. Wang, Z. *et al.* Effective binning of metagenomic contigs using contrastive multi-view representation learning. *Nat. Commun.* **15**, 585 (2024).

34. Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat. Microbiol.* **3**, 836–843 (2018).

35. Chan, P. P., Lin, B. Y., Mak, A. J. & Lowe, T. M. tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes. *Nucleic Acids Res.* **49**, 9077–9096 (2021).

36. Shen, W., Sipos, B. & Zhao, L. SeqKit2: A Swiss army knife for sequence and alignment

processing. *Imeta* **3**, e191 (2024).

37. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

38. Tesson, F. *et al.* Systematic and quantitative view of the antiviral arsenal of prokaryotes.

Nat. Commun. **13**, 2561 (2022).

39. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**,

D412–D419 (2021).

40. Roberts, R. J., Vincze, T., Posfai, J. & Macelis, D. REBASE: a database for DNA

restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res.* **51**,

D629–D630 (2023).

41. Blow, M. J. *et al.* The Epigenomic Landscape of Prokaryotes. *PLoS Genet.* **12**,

e1005854 (2016).

Supplementary Figures

Dataset	Tool	CPU Time [h]	Total Time [h:m:s]	Time difference [fold]
Fecal, simple	modkit	6.7	6:32:03	120x
	nanomotif	0.8	0:03:14	1x
Fecal, complex	modkit	18.0	1 day, 2:11:43	35x
	nanomotif	17.0	0:44:15	1x
ZymoFecal	modkit	14.0	14:28:34	38x
	nanomotif	7.6	0:22:17	1x
Anaerobic Digester	modkit	86.0	1 day, 23:51:30	23x
	nanomotif	65.0	2:02:50	1x

Tab. S1: Benchmark of Nanomotif and Modkit motif identification performance. Running modkit was infeasible for the soil sample and is hence not included. Total time benchmark was performed using 144 AMD EPYC 7H12 CPUs.

585
586
Fig. S1: Parameter sweep over minimum KL-divergence, minimum motif score, and general
methylation threshold used the "motif_identification" module. The parameter sweep was
performed with expected motifs of the six training monocultures. Recall of expected motifs is
primarily affected by the minimum KL-divergence parameter, whereas precision is reversely
affected by the minimum KL-divergence. Precision increases with higher general methylation
threshold and higher minimum motif score. Motif identification is generally stable across
parameters.

594
595 **Fig. S2:** Full genome motif identification benchmark on 46 expected motifs from the five test
596 monocultures.
597

599
600 **Fig. S3:** Full genome motif identification benchmark on 29 expected motif from the six training
601 monocultures.
602

604
Fig. S4: High 5mC false positives in high GC% species. Left panel; On the x-axis is the fraction
of methylated positions occurring explained by a motif and the GC% of the organism's
genome. As the GC% increases, the degree of explained methylated positions decreases.
Additionally, large degrees of false positive 5mC motif are identified in the high GC%
organisms. Right panels; two example organismns, 1) *S. albidoflavus*, with a GC% of 74%,
where 5mC has a continuous drop off in fraction of read modified at C positions, while none
of these positions are explained by a 5mC motif. 2) *P. thermoglucosidasius*, with a GC% of
44%, where 5mC distribution is split into two groups; a low fraction group with no motif
associated and a high fraction group, all explained by a motif.
614

Precision

Recall

618
619 **Fig. S6:** Recall of benchmark presented in Fig. 1C.

620
621 **Fig. S7:** Methylation pattern of monocultures included in the benchmark dataset along with mean sensitivity, specificity, precision, accuracy, and
622 F1 score across the 50 datasets where a contig from each bin has been assigned to another.

623
Fig. S8: Completeness and contamination of MQ and HQ bins before and after removal of
putative contamination. Dashed boxes mark completeness >= 90%, contamination <5%. One
contaminant was removed in Fecal Simple but is not shown.

627
628 **Fig. S9:** Number of correct, incorrect and unassigned classification from nanomotif include
629 module using the monoculture benchmark.
630

632
Fig. S10: Summary of MTase annotations and motif assignments in 11 monocultures and
recovered HQ MAGs from five metagenomes. **a**, Distribution of detected motifs and
annotated MTase genes per. genome. **b**, Percentage of motifs with a high confidence linked
to an annotated MTase gene (orange), or motifs for which one or multiple candidate genes
have been found (blue). **c**, Breakdown of annotated MTases by modification type and RM-
system type. NA indicates unidentified modification type. **d**, Proportion of annotated MTase
involved in RM-systems.
640

 Supplementary Note 1

Direct motif identification in contigs

The assembly sequence and the methylation pileup from "modkit pileup" are used to identify
methylated motifs.

Motifs are identified in each contig sequence separately from other contigs in an assembly.
We use the "fraction modified" value in the modkit pileup output to determine if a position on
the contig is methylated. "Fraction modified" corresponds to the number of mapped reads
modified at the position divided by the number of valid bases at the position, which is the
number of reads with the same canonical base as the respective modification type (A for 6mA
and C for 5mC/4mC).

Each position with a valid base is classified as; NA, if the coverage below the threshold (default
5), not methylated, if the fraction of read methylation is below threshold (default 0.7) or
methylated. We further define two ways in which a position can be methylated; generally
methylated positions, where fraction of read methylation is above the threshold (default 0.7)
and confidently methylated position, where fraction of read methylation is above the threshold
(default 0.8).

Motif search is initiated at a seed motif (the default is the respective base to the evaluated
methylation type, C for 5mC & 4mC and A for 6mA). To determine which position to expand
from the motif, we extract sequences in a window around all confidently methylated positions
(default window size is 41, 20 bases upstream and 20 bases downstream of the methylated
position). These sequences are aligned with respect to the methylation, generating a
methylated sequence set, S. A positional probability matrix (PPM) is then calculated from the
set S.

$$p_{ij} = \frac{1}{S} \sum_{s \in S} \begin{cases} 1 \text{ if } s[i] = j \\ 0 \text{ if } s[i] \neq j \end{cases} , j \in \{A,T,G,C\}$$

This generates a 4x41 table, where the 41 columns correspond to the relative position with
respect to the methylation and the 4 rows correspond to the nucleotide. Next, 10,000
sequences of the same window size are sampled with replacement from the contig, S_{sample},
and a positional nucleotide frequency table of the same dimensions is calculated. The 10,000
sampled windows serve only to generate a background positional frequency table.
Resampling only reinforces the true nucleotide frequency of the background, hence
resampling is non problematic.

$$q_{ij} = \frac{1}{S} \sum_{s \in S_{sample}} \begin{cases} 1 \text{ if } s[i] = j \\ 0 \text{ if } s[i] \neq j \end{cases} , j \in \{A,T,G,C\}$$

679 For each relative position, i, the KL-divergence is calculated from the four frequencies of the
680 methylated sequence frequency table to the four frequencies of the sampled sequence
681 frequency table.

$$D_{KL}^{(i)}(\mathbf{P} \parallel \mathbf{Q})) = \sum_{j \in \{A, T, G, C\}} p_{ij} \log(\frac{p_{ij}}{q_{ij}})$$

This generates a vector of size 41, where each entry corresponds to a KL-divergence value.
Positions are, per default, only considered for expansion if the KL-divergence is greater than
0.05. After selecting which position to expand, we select which bases to incorporate at each
of these positions by two criteria; 1. the frequency of a base in the methylation sequence
frequency table must be above 25% and the frequency of a base must be above the frequency
in the sampled sequence frequency table. If more than one base at a position meets this
criteria, we keep both of them and combinations of them a, e.g. accepting A and G at relative
position 2 with seed NNANN would give rise to NNANA, NNANG and NNANR.
693
694 Each new motif candidate after the expansion is evaluated using a beta-Bernoulli model,
695 treating each motif occurrence as a Bernoulli trial, being a success if it is a generally
696 methylated position and a failure if not a methylated position. We use a Beta(α=0, β=1) as a
697 prior, which means the posterior is also a Beta distribution with the parameters:
698
699 $$\alpha = \alpha_{prior} + n_{methylated} \, , \beta = \beta_{prior} + n_{non-methylated}$$
700
The posterior distribution is used to score each motif using the mean, standard deviation, and
mean difference from the preceding motif. The mean represents the degree of motif
methylation, a value expected to increase as the motif is refined. The standard deviation is
used to penalize when few observations are present. Mean difference is expected to be high,
when a desirable nucleotide addition is made, as it keeps the N highly methylated motif
variants and disregards 4-N non-methylated motif variants, and is approximately zero for
nucleotide insertion which contributes nothing to the recognition sequence.
708
709 $$score = mean_{diff} \cdot mean \cdot -log_{10}(standard\ deviation)$$
710
After scoring each of the new motifs, the highest scoring motif is stored. Next, one of the motifs
is selected for propagation to the new set of motifs. The objective of the search is to converge
on the motif candidate contributing the most positive methylation sites. The search heuristic is
therefore formulated to minimize the proportion of generally methylated positions removed
and maximize the proportion of non-methylated positions removed with respect to the seed
motif. Concretely, the heuristic is calculated using the α and β parameters of the beta-
Bernoulli posterior of the current motif and the seed motif, as they represent the number of
methylated and non-methylated motif sites.
719
720 $$priority = (1 - (\alpha_{current}/\alpha_{seed})) \cdot (\beta_{current}/\beta_{seed})$$
721
The formulation of the priority results in the motif with the lowest priority score to be chosen
for the next iteration, as this decreases non-methylated sites while retaining maximum
methylated sites. For the next iteration, the methylation sequences extracted initially are
subsetted to those only containing the motif picked for expansion. After this the positional
frequency table and KL-divergence is recalculated and the same procedure as before follows.
The algorithm expands and scores following the steps described above, until the maximum
score of a motif has not increased for 10 rounds or no more motif candidates are left to explore.
The best scoring motif is then kept and saved to candidate motifs if its score is >0.2, otherwise
dropped. The whole procedure is then repeated from the same seed, but removing sequences
containing previously identified candidate motifs from methylated sequences. This is
continued until 25 candidate motifs with insufficient score have been dropped or only 1% of
methylation sequences remain.

734
After all candidate motifs have been identified in a contig, they are subjected to a series of
post-processing steps to improve final motifs. First, motifs which are a sub motif of other motifs
are removed, which is the case if the sequence of any other motif is contained within the
sequence of the current motif, e.g. C**5mC**WGG would give rise to removal of **6mA**CCWGG,
as CCWGG is contained within ACCWGG. This step was added to mitigate false positive
motifs resulting from 5mC methylations in close proximity to adenine can result in 6mA
methylation calls, which subsequently produce a sufficiently strong signal to "detect" 6mA
motifs. In this case we accept the possibility of removing similar motifs with different
methylation types. Next, we remove motifs which have isolated bases, defined as a non-N
position with at least 2 Ns on both sides. Next, we merge motifs whose sequences are similar,
which can be the case for more generic motifs such as CCWGG, where CCAGG and CCTGG
were found as separate motifs, but should constitute one motif. Motif merging is done by
constructing a distance graph between all motifs, where motifs are only connected if the
hamming distance is 2 or less. Motifs are then defined to be part of the same cluster in the
graph if they are mutually reachable. All motifs within the same cluster are merged into a single
motif, representing all motifs contained within the cluster. The merged motif is only accepted
if the mean degree of methylation is not less than 0.2 of the mean methylation of the pre merge
motifs, otherwise the pre-merge motifs are kept as is. Finally, motifs are queried for motif
complements. If another motif is the complementary sequence of the motif, it gets removed
and added as a complementary motif instead. Palindromic motifs are always considered as
the complementary of itself.
756
Indirect motif detection
Direct motif identification is performed on one contig without any information from other contigs
in an assembly. To detect potentially missed motifs in contigs, we perform what we term
indirect detection of motifs in contigs, so called as they are only detected because the motif
was directly detected with high confidence in another contig. To get indirectly identified motifs,
we take the complete set of all motifs identified in all contigs and calculate α and β of the Beta
posterior of the beta-Bernoulli model for all contigs. We report the α and beta parameters as
the number of motif methylations and non-methylations, respectively.
765
Bin consensus
Bin consensus is evaluated by taking the complete set of motifs for a bin and checking if a
motif meets a set of criteria. Firstly, a motif has to have been directly detected in at least one
of the contigs in the bin. Next, we remove motifs that are not methylated in at least 75% of the
contigs in the bin. We estimate this by counting the number of motif occurrences in contigs
with a mean methylation of a motif above 25% and dividing by the total number of motif
occurrences in the bin; if the fraction of motif occurrences present in methylated contigs is
above 0.75, they are kept. Lastly, of the kept motifs, sub-motifs are removed as described in
the post-processing step in the direct motif identification section. The remaining motifs are
considered bin consensus motifs.
776
777

Supplementary Note 2

Annotation of gold standard proteins and their methylation motifs

We initially analyzed 11 prokaryotic strains known to encode gold-standard (GS) RM system
proteins using Nanomotif (Fig. S9a, supplementary data 1 & 2). GS proteins are those whose
biochemical functions have been experimentally validated with their exact coding DNA
sequences identified[40]. The MTase-linker module of Nanomotif successfully annotated 43 out
of 44 gold-standard MTase enzymes and linked the associated motifs, if active, across the 11
monocultural genomes. The single missing annotation is likely a false negative, as
DefenseFinder had multiple HMM-profile hits for this gene, but ultimately assigned the
corresponding gene as part of a non-methylation mediated defense system. Among the 44
GS MTases, 40 were found to be active. Notably, the motifs identified for all active GS MTases
precisely   match   the   specificities   reported   for   the   GS   MTases   in   REBASE.
For the non-GS epigenetic systems in these 11 prokaryotic strains, our gene annotations and
motif assignments are generally aligned with existing REBASE entries. However, in *S.
oneidensis*, *K.aquimarina,* and *D. acetoxidans*, the MTase-linker module annotated additional
type II MTases beyond those previously reported in REBASE. While some may represent false
positives, for example, *contig_2_3607* in *S. oneidensis*, others are supported by active motifs
without any alternative gene assignments, for example *contig_1_1589* in *K. aquimarina*
(supplementary data 2).

In total, 42 out of 71 detected motifs were confidently linked to annotated RM systems or
orphan MTases. For 18 additional motifs, candidate genes with matching methylation features
were identified. Notably, the remaining motifs include several from *Anabaena variabilis,* that
may represent variants of fewer distinct motifs, and *M.ruber,* representing noise motifs
attributed to the increased false positive rate of ONT's methylation models in high GC contexts
(Fig. S3). Apart from these unassigned motifs and two motifs in *D. acetoxidans*, all detected
motifs were either supported by REBASE entries or corroborated by previous PacBio
sequencing data[41], further validating the accuracy and reliability of the detected motifs.

Supplementary Note 3

Discovery of epigenetics systems in diverse metagenomes

We next analyzed a diverse set of prokaryotic communities using nanomotif (Fig. S10). Using the MTase-linker, 3123 MTase genes were detected across 549 HQ MAGs, resulting in a median of 6 MTases genes per genome, and 3 RM systems encompassing an MTase per genome. Type II MTases were the most abundant (67%). This was followed by type I (16%), IIG (9%), and III (7%). Type II MTases were also the most prevalent of all types not associated with an RM-system. Only 24% of type II MTases were co-located with a cognate restriction enzyme. Previous studies have also reported the frequent presence of orphan MTases in a wide range of bacterial genomes[41]. However, it is important to acknowledge that some associated REase genes may have gone undetected due to sequence divergence, especially in the complex samples with high novelty. In such cases, there may be an overestimation of orphan type II MTase genes. Despite this, many examples of orphan MTases are indeed genuine, and they represent a large group of MTases with non-RM functions. Similarly, 24% and 39% of Type I and Type III MTases, respectively, were unexpectedly identified without corresponding restriction enzymes or Type IS subunits. In the soil sample, a significantly lower number of MTase genes per genome compared to the other metagenomes was observed as well. This discrepancy is likely due to the limited sensitivity of the HMM models in complex samples.

For 76% of detected motifs, at least one candidate MTase with similar methylation characteristics was found within the same genome. In 232 cases, a single motif could be confidently linked to a specific MTase gene or RM system, resulting in a high-confidence set of MTase target motif annotations. Notably, 65 of these motifs involved 5mC modifications, which are notoriously difficult to detect with SMRT sequencing. This highlights the potential of Nanomotif to accurately annotate MTase target recognition motifs in metagenomes, including those with 5mC modifications.

Supplementary Data

Supplementary data 1: a) expected motifs in REBASE gold standard monocultures. b) summary stats for recovered MAGs in metagenomes. c) contigs annotated as mobile genetic elements. d) Contamination and completeness before and after Nanomotif contamination removal. e) Motifs identified with Nanomotif in monocultures. f) Motifs identified with Nanomotif in Zymo reference samples. g) Motifs identified with Nanomotif in metagenomes.

Supplementary data 2: MTase-linker results from monocultures and metagenomic samples