1 **SLC35G1: A highly chloride-sensitive transporter responsible for the basolateral membrane**
2 **transport in intestinal citrate absorption**
3

4 Yoshihisa Mimura[1,†], Tomoya Yasujima[1,*], Katsuhisa Inoue[2], Shogo Akino[1], Chitaka Namba[1], Hiroyuki
5 Kusuhara[3], Yutaro Sekiguchi[1], Kinya Ohta[4], Takahiro Yamashiro[1], Hiroaki Yuasa[1]
6
[1] *Department of Biopharmaceutics, Graduate School of Pharmaceutical Sciences, Nagoya City University,*
3-1 Tanabe-dori, Mizuho-ku, Nagoya 467-8603, Japan
[2] *Department of Biopharmaceutics, School of Pharmacy, Tokyo University of Pharmacy and Life Sciences,*
1432-1 Horinouchi, Hachioji, Tokyo 192-0392, Japan
[3] *Laboratory of Molecular Pharmacokinetics, Graduate School of Pharmaceutical Sciences, The University*
of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
[4] *College of Pharmacy, Kinjo Gakuin University, 2-1723 Omori, Moriyama-ku, Nagoya 463-8521, Japan*
14
* Corresponding author
Tomoya Yasujima, Ph.D.
Department of Biopharmaceutics, Graduate School of Pharmaceutical Sciences, Nagoya City University,
3-1 Tanabe-dori, Mizuho-ku, Nagoya 467-8603, Japan.
Telephone/Fax: +81-52-836-3425
E-mail address: yasujima@phar.nagoya-cu.ac.jp
21
22 † Present address: *Department of Pharmacy, Nagoya City University Hospital, 1 Kawasumi Mizuho-cho,*
23 *Mizuho-ku, Nagoya, 467-8602, Japan*

Abstract

The intestinal absorption of essential nutrients, especially those not readily biosynthesized, is a critical physiological process for maintaining homeostasis. Numerous studies have indicated that intestinal absorption is mediated by various membrane transporters. Citrate, a crucial bioactive compound produced as an intermediate in the Krebs cycle, is absorbed in the small intestine through carrier-mediated systems because of its high hydrophilicity. While the luminal absorption of citrate is mediated by Na^+-dicarboxylate cotransporter 1 (NaDC1), the mechanism governing the release of the transported citrate into the bloodstream remains unknown. Here, we explored the transporters responsible for intestinal citrate absorption at the basolateral membrane, focusing on highly expressed orphan transporters in the small intestine as candidates. Consequently, SLC35G1, originally identified as a partner of stromal interaction molecule 1, a cell surface transmembrane glycoprotein, was found to play a role in the intestinal absorption of citrate at the basolateral membrane. Furthermore, our results revealed that SLC35G1-mediated citrate transport was diminished by chloride ions at extracellular concentrations. This suggests that SLC35G1, to our best knowledge, is the first transporter identified to be extremely sensitive to chloride ions among those functioning on the basolateral membrane of intestinal epithelial cells. This study provides valuable insights into the intestinal absorption of citrate and significantly contributes to elucidating the poorly understood molecular basis of the intestinal absorption system.

Introduction

Citrate plays a crucial role as a tricarboxylic acid (TCA) cycle intermediate, contributing to the generation of metabolizable energy. Since citrate is obtained not only through biosynthesis but also from the diet, the existence of intestinal absorption system has long been acknowledged (*Browne et al. , 1978*). Luminal absorption is predominantly mediated by Na^+-dependent dicarboxylate transporter 1 (NaDC1/SLC13A2) (*Pajor, 2006*); however, the transporter responsible for the release of citrate and its metabolites across the basolateral membrane remain unidentified (*Pajor, 2014*). To elucidate the molecular mechanism underlying citrate absorption at the basolateral membrane, we explored the transporters implicated in the process. Our study provides novel insights into basolateral membrane transporters, addressing a previously poorly understood aspect in intestinal epithelial cells.

Results and discussion

Our investigation revealed that SLC35G1, a member of the solute carrier (SLC) 35 family and originally identified as a partner of stromal interaction molecule 1 (*Krapivinsky et al. , 2011*), is capable of transporting citrate. While the functions of nucleoside-sugar transporters on the endoplasmic reticulum membrane have been identified for certain SLC35 family members, including SLC35A, SLC35B, SLC35C, and SLC35D (*Ishida and Kawakita, 2004*), the roles of SLC35E, SLC35F, and SLC35G have not yet been fully elucidated, as shown in *Figure 1A*. For the functional analysis, we first established Madin-Darby canine kidney (MDCKII) cells stably expressing SLC35G1. Using these cells, we observed a linear

increase in SLC35G1-mediated citrate uptake with time, specifically during the first 10 min in Hanks' solution mimicking the cytosolic ion content by eliminating Cl⁻ (*Figure 1B*). The acidic condition (pH 5.5) was used to gain sufficient uptake for functional analysis. Therefore, the uptake at 10 min was evaluated in subsequent studies. A kinetic analysis revealed that the SLC35G1-specific uptake of citrate exhibited saturation kinetics, with a V_{max} of 1.10 nmol/min/mg protein and a K_m of 519 μM (*Figure 1C*). Notably, the specific uptake of citrate by SLC35G1 was extensively inhibited by extracellular Cl⁻ (*Figure 1D*), with an IC_{50} value of 6.7 mM (*Figure 1E*). As previously reported (*Melkikh and Sutormina, 2008*), the Cl⁻ concentration is lower in the cytosol (5 mM) than in the extracellular conditions (120 mM), suggesting that SLC35G1 is a potential citrate exporter. Additionally, we examined the effect of various compounds, including other TCA cycle intermediates and typical inhibitors of organic anion/cation transporters at a concentration of 200 μM, to explore the potential role of SLC35G1 in their transport (*Figure 1F*). Among the tested compounds, certain anionic compounds significantly inhibited SLC35G1-specific citrate uptake, indicating that they are also recognized by SLC35G1. SLC35G1-specific citrate uptake showed pH dependence (*Figure 1G*). To investigate whether SLC35G1 utilizes an H⁺ gradient as a driving force for citrate transport, we examined the effect of the protonophore nigericin in the uptake solution at pH 5.5 to dissipate the H⁺ gradient across the plasma membrane. However, treatment with nigericin did not affect the specific uptake of citrate (*Figure 1H*), excluding this possibility. Given that one carboxylate can be protonated with a pK_a of 5.4 (*Contreras-Trigo et al. , 2018*) (*Figure 1I*), SLC35G1 may preferably transport dianionic citrate, like NaDC1 (*Pajor and Sun, 1996*). Finally, we determined the role of SLC35G1 in the transcellular transport of citrate at the basolateral membranes using SLC35G1-transfected polarized MDCKII cells on Transwell permeable membrane filters. As shown in *Figure 1J*, SLC35G1 predominantly localized to the basolateral membrane of polarized MDCKII cells. Transport studies were performed under pH gradient conditions (apical pH 5.5 and basal pH 7.4) in the presence of Cl⁻ to mimic intestinal physiological conditions. SLC35G1-transfected cells showed no significant change in intracellular citrate accumulation from the apical side, compared to mock cells (*Figure 1K*). However, apical-to-basolateral transcellular transport was significantly higher in SLC35G1-transfected cells (*Figure 1L*). On the other hand, SLC35G1 overexpression affected neither intracellular accumulation nor transcellular transport from the basolateral side. This is reasonable since the basolaterally localized SLC35G1 operates for efflux but not for uptake in the presence of Cl⁻ in the extracellular environment. These results suggest that SLC35G1 plays a role in the intestinal absorption of citrate at the basolateral membrane as an exporter under physiological conditions.

   To further investigate the physiological role of SLC35G1, we examined its expression in human tissues (*Figure 2A*). Quantitative real-time PCR analysis revealed that SLC35G1 is highly expressed in the digestive tract, especially in the upper part of the small intestine, including the duodenum and jejunum, followed by the testis and pancreas. Additionally, immunohistochemical staining of the human jejunum clearly showed that SLC35G1 is localized on the basolateral membrane, as indicated by the co-localization of ATP1A1, a marker protein on the basolateral membrane (*Figure 2B*). To investigate the physiological impact of SLC35G1 on intestinal citrate absorption, we examined the effect of SLC35G1 knockdown on

citrate uptake in human colorectal adenocarcinoma Caco-2 cells, a commonly used model for studying
intestinal absorbency (*Yee, 1997*) (*Figure 2C*), in the absence of Cl⁻. As shown in *Figure 2C*, citrate uptake
was significantly reduced following RNA interference with siRNAs targeting SLC35G1 mRNA, providing
evidence of the involvement of SLC35G1 in intestinal citrate absorption. The reduction in citrate uptake
was not indicative of off-target effects, as multiple siRNAs with distinct nucleic acid sequences yielded
similar reductions. Moreover, the expression of SLC35G1 mRNA was reduced by these siRNAs,
confirming successful silencing of SLC35G1 (*Figure 2D*). The correlation between the decrease in citrate
uptake and mRNA levels suggests that SLC35G1 is primarily involved in citrate uptake.
Immunohistochemical staining revealed that endogenous SLC35G1 in polarized Caco-2 cells cultured on
Transwell membranes was also localized on the basolateral membranes (*Figure 2E*). Taken together, these
results indicate that intestinal citrate transport at the basolateral membrane is primarily mediated by
SLC35G1. Additionally, it acts cooperatively with NaDC1, which is reported to be highly expressed in the
jejunum (*Pajor, 1995*), for citrate absorption on the apical membrane, as illustrated in *Figure 2F*.

Our findings suggest that SLC35G1 is localized on the basolateral side of intestinal epithelial cells
and, to our knowledge, is the first transporter identified with chloride-sensitivity, which is a unique
characteristic to render with the directional transport function for an SLC transporter. These findings pave
the way for several follow-up directions related to the molecular basis of the intestinal absorption of
various compounds, including nutrients and drugs.
119

Materials and Methods
Materials
[^{14}C]citrate (116.4 mCi/mmol) was obtained from PerkinElmer Life and Analytical Sciences (Boston, MA,
USA). Unlabeled citrate and Dulbecco's modified Eagle's medium (DMEM) were sourced from Wako Pure
Chemical Industries (Osaka, Japan), and fetal bovine serum (FBS) was obtained from Sigma Aldrich (St.
Louis, MO, USA). All other reagents were of analytical grade and commercially acquired.
126

127 **Cell culture**
128 Madin-Darby canine kidney II (MDCKII) cells and Caco-2 cells were cultured at 37°C and 5% CO_2 in
129 DMEM supplemented with 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin, as previously
130 described (*Mimura et al. , 2017*).
131

Preparation of plasmids
The cDNA for human SLC35G1 was cloned using a reverse transcription (RT)-PCR-based method, as
previously described (*Mimura et al. , 2017*). Briefly, an RT reaction was performed to obtain a cDNA
mixture, using 1 µg of the human intestine total RNA (BioChain Institute, Newark, CA), an oligo(dT)
primer, and the highly efficient reverse transcriptase ReverTra Ace (Toyobo, Osaka, Japan). The SLC35G1
cDNA was then amplified by PCR, using KOD plus polymerase (Toyobo) and the following primers:
forward primer, 5′-GAGATGCGGCCTCAGGACAG-3′, and reverse primer,

5′-TAGCCTCCCCACCCATATCCC-3′. These primers were designed based on the GenBank sequences for

SCL35G1 (accession number: NM_001134658.2). The second PCR was performed using the initial PCR

product as a template, with a forward primer containing an EcoRI restriction site (underlined),

5′-AGAATTCCGAGATGCGGCCTCAGGAC-3′, and a reverse primer containing an XbaI restriction site

(underlined), 5′-GTTCTAGATGTGTGCGTATGCTG-3′. The resulting cDNA product was inserted into the

pCI-neo vector (Promega, Madison, WI, USA) between the EcoRI and XbaI sites, and the sequence of the

final product was determined using an automated sequencer.

Uptake study

First, we established MCDKII cells stably expressing SLC35G1 as previously described (*Yamamoto et al.*,

2010). Briefly, MDCKII cells were transfected with the plasmid carrying the SLC35G1 cDNA, using

Lipofectamine 2000 as a transfection reagent. The cells were then cultured in DMEM supplemented with

10% FBS and 800 μg/mL G418 for 2–3 weeks. Subsequently, G418-resistant clones were selected and

assessed for the transport of [^{14}C]citrate. MDCKII cells stably expressing SLC35G1 with high citrate

transport activity were used in subsequent studies.

MDCKII cells stably expressing SLC35G1 (1.5×10^5 cells/mL, 1 mL/well) were grown in 24-well

plates for 48 h until they reached confluence. For standard transport assays, cells were preincubated in

substrate-free uptake buffer, which was Hanks' solution modified to reflect cytosolic conditions (142.06

mmol/L K-gluconate, 0.812 mmol/L $MgSO_4$, 0.379 mM K_2HPO_4, 0.441 mM KH_2PO_4, 0.952 mM

Ca-gluconate, 25 mM D-glucose) and supplemented with 10 mM 2-(N-morpholino)-ethanesulfonic acid

(MES; pH 5.5), for 5 min. Subsequently, uptake assays were initiated by replacing the substrate-free uptake

buffer with an uptake buffer containing a [^{14}C]citrate (0.25 mL). For initial uptake analysis experiments,

the uptake period was set to 10 min in the initial phase, in which uptake was in proportion to time. Other

experiments were conducted in a time-dependent manner, as specified. When examining the effects of

various compounds on citrate uptake, the test compounds were added to the buffer only during the uptake

period. All procedures were performed at 37°C. Assays were stopped by adding 2 mL of ice-cold

substrate-free uptake buffer, followed by two washes with an additional 2 mL of the same buffer. The cells

were solubilized in 0.5 mL of 0.2 mol/L NaOH solution containing 0.5% sodium dodecyl sulfate (SDS),

and the associated radioactivity was measured by liquid scintillation counting for uptake evaluation. The

cellular protein content was determined by the bicinchoninic acid (BCA) method, using bovine serum

albumin as the standard (3). In experiments to examine the effect of ionic conditions, K-gluconate in the

control uptake solution was replaced as indicated. Uptake solutions were supplemented with 10 mM MES

(pH 6.5 and below) or 10 mM HEPES (pH 7.0 and above) in experiments to examine the effect of pH. In

experiments to examine the effect of nigericin as an agent for intracellular acidification, a mannitol-based

buffer (250 mM mannitol, 1.2 mM $MgSO_4$, 2 mM KH_2PO_4, 5 mM D-glucose, 10 μM nigericin, and 20 mM

MES) was used and uptake assays were conducted for 10 min at pH 5.5 and 37°C in the presence (10 μM)

or absence (control) of nigericin after pretreatment for 5 min with or without nigericin under the same

conditions. To estimate nonspecific uptake, uptake assays were conducted in mock cells transfected with an

empty pCI-neo vector. The specific uptake of citrate by SLC35G1 was estimated by subtracting the uptake

in mock cells from that in SLC35G1-transfected cells.

Transcellular transport study

MDCKII cells were seeded at a density of 2×10^5 cells on each polycarbonate membrane insert in a

12-well Transwell plate and cultured for 5 days. After removing the culture medium from both sides of the

inserts, cells were preincubated for 5 min at 37°C in Hanks' solution (136.7 mM NaCl, 5.36 mM KCl,

0.952 mM $CaCl_2$, 0.812 mM $MgSO_4$, 0.441 mM KH_2PO_4, 0.385 mM Na_2HPO_4, and 25 mM D-glucose)

supplemented with 10 mM MES (pH 5.5) in the apical chamber and 10 mM HEPES (pH 7.4) in the

basolateral chamber. To initiate transcellular transport from the apical side to the basolateral side, 0.5 mL of

the Hanks' solution (pH 5.5) containing [^{14}C]citrate was replaced in the apical chamber. For transport in the

opposite direction (basolateral side to apical side), 1.5 mL of Hanks' solution (pH 7.4) containing

[^{14}C]citrate was replaced in the basal chamber. To monitor transcellular transport, samples (100 μL from

basal chamber and 50 μL from the apical chamber) were periodically collected, replenishing each collected

volume with an equal volume of fresh buffer. At the end of the transport study, the assays were stopped by

adding ice-cold Hanks' solution (1 mL for the apical chamber and 3 mL for the basolateral chamber),

followed by washing the cells twice with the same buffer. The cells were then solubilized in 0.5 mL of 0.2

M NaOH solution containing 0.5% SDS at room temperature for 1 h. The radioactivity associated with the

cells was measured by liquid scintillation counting for uptake evaluation.

Quantification of SLC35G1 mRNA by quantitative real-time PCR

Total RNA samples from various human tissues (BioChain Institute) and Caco-2 cells were used to prepare

cDNA using the ReverTra Ace reverse transcriptase. A quantitative real-time PCR was performed using

SsoFast EvaGreen Supermix with Low ROX (Bio-Rad Laboratories, Hercules, CA, USA) on a 7300 Fast

Real-time PCR System (Applied Biosystems, Foster City, CA, USA) with the following primers: forward

primer for SLC35G1, 5′- AGAGCCCACTGAGAAAAGGA -3′; reverse primer for SLC35G1,

5′-GTAGGTGCTGGCTCTGCCT-3′;   forward   primer   for   GAPDH,

5′-CGGAGTCAACGGATTTGGTCGTAT-3′;   reverse   primer   for   GAPDH,

5′-AGCCTTCTCCATGGTGGTGAAGAC-3′. GAPDH was employed as an internal control for

normalization of the mRNA expression levels of SLC35G1.

Knockdown study

Caco-2 cells (5.0×10^4 cells/mL, 1 mL/well) were grown on 24-well–coated plates for 6 h, transfected with

5 pmol/well of the Silencer Selected siRNAs specific to the mRNA of SLC35G1 (Thermo Fisher Scientific,

Waltham, MA), using 1.5 mL/well of Lipofectamine RNAi MAX (Thermo Fisher Scientific), and cultured

for 5 days for silencing of the designated transporter by RNAi. The sequences of the siRNAs are following:

sense sequence of #1 siRNA, GGAGUGAUCCUUAUCGUGATT; antisense sequence of #1 siRNA,

UCACGAUAAGGAUCACUCCAG, sense sequence of #2 siRNA, CUUGCUUAAUAUACAGAAATT;

antisense sequence of #2 siRNA, UUUCUGUAUAUUAAGCAAGGG, sense sequence of #3 siRNA,

CAUUUGGUAUUAUGUAGUATT;   antisense   sequence   of   #3   siRNA,

UACUACAUAAUACCAAAUGCT. For control, negative control Silencer Selected siRNA (Thermo

Fisher Scientific), was used. Citrate uptake assays were conducted as described above for

SLC35G1-transfected MDCKII cells.

Immunofluorescence staining

To examine the localization of SLC35G1, MDCKII cells stably expressing SLC35G1 were seeded at a

density of 2×10^5 cells/insert on a 12-well Transwell plate and cultured for 5 days, and Caco-2 cells were

seeded at a density of 5×10^5 cells/insert on a 12-well Transwell plate and cultured for 21 days. The cells

were washed thrice with ice-cold PBS and fixed in 5% paraformaldehyde for 15 min. After washing thrice

with PBS, the cells were incubated for 1 h at room temperature with an anti-human SLC35G1 goat

polyclonal antibody at a dilution of 1:100 in Can Get Signal Immunostain Immunoreaction Enhancer

Solution A (Toyobo). The cells were washed with PBS, and the primary antibody was probed with

anti-goat IgG Alexa Fluor 488 (Jackson ImmunoResearch Laboratories, West Grove, PA, USA) at a

dilution of 1:400 in Can Get Signal Immunostain Immunoreaction Enhancer Solution B (Toyobo) for 1 h at

room temperature. After three washes with PBS, the cells were mounted on a glass slide in 9:1

glycerol/PBS containing 1 μM DAPI.

A specimen of normal human adult small intestine frozen tissue sections (US Biomax, Derwood, MD,

USA) was blocked in G-Block (GenoStaff, Tokyo, Japan) for 1 h and then incubated for 1 h at room

temperature with an anti-human SLC35G1 goat polyclonal antibody at a dilution of 1:100 in Can Get

Signal Immunostain Immunoreaction Enhancer Solution A. After washing thrice with PBS, the primary

antibody was probed with anti-goat IgG Alexa Fluor 488 by incubating for 1 h at room temperature.

Additionally, we stained ATP1A1 with an anti-human ATP1A1 mouse monoclonal antibody (clone

BXP-21, Abcam, Cambridge, UK) labeled with the Ab-10 Rapid Hilyte Fluor 647 labeling kit (Dojindo

Laboratories, Kumamoto, Japan) according to the manufacturer's instructions. After washing thrice with

PBS, the tissues were mounted in 9:1 glycerol/PBS containing 1 μM DAPI.

The localization of immunofluorescently labeled proteins and nuclei was visualized using a confocal

laser scanning microscope (LSM510; Zeiss, Jena, Germany).

Data Analysis

The saturable transport of citrate by SLC35G1 was analyzed by assuming Michaelis-Menten-type

carrier-mediated transport, as represented by the following equation: $v = V_{max} \times s/(K_m + s)$. The apparent

parameters of the maximum transport rate (V_{max}) and Michaelis constant (K_m) were estimated by fitting this

equation to the experimental profile of the uptake rate (v) versus the concentration (s) of citrate as the

substrate, using a nonlinear least-squares regression analysis program, WinNonlin (Pharsight, Mountain

View, CA, USA), with v^{-2} as the weight.

When s is considerably smaller than K_m ($s << K_\mathrm{m}$), v in the presence of an inhibitor can be expressed

as $v = v_0/(1 + (i/IC_{50})^{n})$. The half-maximal inhibitory concentration (IC_{50}) was estimated, along with the

Hill coefficient (n) and v in the absence of inhibitors (v_0), by fitting this equation to the experimental profile

of v versus the inhibitor concentration (i).

ACKNOWLEDGEMENTS

This study was supported by research grant from Nakatomi Foundation.

Figure 1. Functional characteristics of SLC35G1 stably expressed in MDCKII cells. (A) Phylogenetic tree of the SLC35 members and identified functions. (B) Time course of [^{14}C]citrate (1 μM) uptake in SLC35G1-transfected MDCKII cells (open circles) and mock cells (closed circles), evaluated at pH 5.5 and 37°C. (C) Concentration-dependent [^{14}C]citrate uptake by SLC35G1, evaluated at various concentrations for 10 min at pH 5.5 and 37°C. The estimated values of V_{max} and K_m were 1.11 ± 0.16 nmol/min/mg protein and 519 ± 86 μM, respectively. (D) Effect of ionic conditions on [^{14}C]citrate (1 μM) uptake by SLC35G1, evaluated for 10 min at pH 5.5 and 37°C. K-gluconate in the control uptake solution was replaced as indicated. (E) Concentration-dependent chloride inhibition of [^{14}C]citrate (1 μM) uptake by SLC35G1, evaluated at various chloride concentrations for 10 min at pH 5.5 and 37°C. The estimated value of IC_{50} was 6.7 ± 1.4 mM. (F) Effect of various compounds on [^{14}C]citrate (1 μM) uptake by SLC35G1, evaluated for 10 min at pH 5.5 and 37°C in the presence (200 μM) or absence (control) of a test compound. BSP, bromosulfophthalein; DIDS, 4,4′-diisothiocyano-2,2′-stilbenedisulfonic acid. (G) Effect of pH on [^{14}C]citrate (1 μM) uptake in SLC35G1-transfected MDCKII cells (open circles) and mock cells (closed circles), evaluated for 10 min at 37°C, using the uptake solutions supplemented with 10 mM MES (pH 6.5 and below) or 10 mM HEPES (pH 7.0 and above). (H) Effect of the protonophore nigericin on [^{14}C]citrate (1 μM) uptake by SLC35G1, evaluated for 10 min at pH 5.5 and 37°C in the presence (10 μM) or absence (control) of nigericin after pretreatment for 5 min with or without nigericin under the same conditions. (I) Structural formula illustrating the chemical equilibrium of citrate. (J) Cellular localization of SLC35G1 stably expressed in polarized MDCKII cells. The confocal laser-scanning microscopic image shows SLC35G1 (green) and nuclei stained with 4′,6-diamidino-2-phenylindole (DAPI; blue). Scale bar, 50 μm. (K) Intracellular accumulation of citrate in polarized MDCKII cells stably expressing SLC35G1. Intracellular accumulation of [^{14}C]citrate (1 μM) was evaluated for 90 min at pH 5.5 on the apical side (a) and pH 7.4 on the basolateral side (b) of intestinal epithelial cells. (L) Time course of transcellular [^{14}C]citrate (1 μM) transport in polarized MDCKII cells stably expressing SLC35G1 and mock cells, evaluated at pH 5.5 in the apical and pH 7.4 in the basolateral chamber. Data represent the mean ± SD of three biological replicates (B-H, K, L). Statistical differences were assessed using analysis of variance (ANOVA) followed by Dunnett's test (D, F, L) or using Student's t-test (G, H, K). *, $p < 0.05$ compared with the control (F), or compared with the mock at each pH (G) or time point (L).

Figure 2. (A) Analysis of SLC35G1 mRNA expression in various human tissues using quantitative real-time PCR. (B) The immunofluorescent images show the subcellular localization of SLC35G1 (green), breast cancer resistance protein (BCRP; red), and 4′,6-diamidino-2-phenylindole (DAPI; blue) in the human jejunum. Scale bar, 100 μm. (C-D) Effect of SLC35G1 silencing on citrate uptake and on SLC35G1 mRNA expression in Caco-2 cells. (C) The uptake rate of [^{14}C]citrate (1 μM) was evaluated for 10 min at pH 5.5 and 37°C in Caco-2 cells transfected with three different siRNAs specific to SLC35G1 mRNA or control siRNA. (D) SLC35G1 mRNA levels were assessed using quantitative real-time PCR analysis. (E) Cellular localization of endogenous SLC35G1 in polarized Caco-2 cells. The confocal laser-scanning microscopic image shows SLC35G1 (green) and nuclei stained with DAPI (blue). Scale bar, 50 μm. (F)

298   Schematic model showing intestinal citrate absorption mediated by NaDC1 and SLC35G1. Data represent

299   the mean ± SD of three technical repeats (A) or biological repeats (C-D). Statistical differences were

300   assessed using analysis of variance (ANOVA) followed by Dunnett's test (C-D).

References

Browne JL, Sanford PA, Smyth DH. 1978. Transfer and metabolism of citrate, succinate, alpha-ketoglutarate and pyruvate by hamster small intestine. *Proc R Soc Lond B Biol Sci* **200**:117-135. DOI: 10.1098/rspb.1978.0010, PMID: 24848.

Contreras-Trigo B, Diaz-Garcia V, Guzman-Gutierrez E, Sanhueza I, Coelho P, Godoy SE, Torres S, Oyarzun P. 2018. Slight pH Fluctuations in the Gold Nanoparticle Synthesis Process Influence the Performance of the Citrate Reduction Method. *Sensors (Basel)* **18**. DOI: 10.3390/s18072246, PMID: 6068536.

Ishida N, Kawakita M. 2004. Molecular physiology and pathology of the nucleotide sugar transporter family (SLC35). *Pflugers Arch* **447**:768-775. DOI: 10.1007/s00424-003-1093-0, PMID: 12759756.

Krapivinsky G, Krapivinsky L, Stotz SC, Manasian Y, Clapham DE. 2011. POST, partner of stromal interaction molecule 1 (STIM1), targets STIM1 to multiple transporters. *Proc Natl Acad Sci U S A* **108**:19234-19239. DOI: 10.1073/pnas.1117231108, PMID: 3228472.

Melkikh AV, Sutormina MI. 2008. Model of active transport of ions in cardiac cell. *J Theor Biol* **252**:247-254. DOI: 10.1016/j.jtbi.2008.02.006, PMID: 18353373.

Mimura Y, Yasujima T, Ohta K, Inoue K, Yuasa H. 2017. Functional Identification of Plasma Membrane Monoamine Transporter (PMAT/SLC29A4) as an Atenolol Transporter Sensitive to Flavonoids Contained in Apple Juice. *J Pharm Sci* **106**:2592-2598. DOI: 10.1016/j.xphs.2017.01.009, PMID: 28089688.

Pajor AM. 1995. Sequence and functional characterization of a renal sodium/dicarboxylate cotransporter. *J Biol Chem* **270**:5779-5785. DOI: 10.1074/jbc.270.11.5779, PMID: 7890707.

Pajor AM. 2006. Molecular properties of the SLC13 family of dicarboxylate and sulfate transporters. *Pflugers Arch* **451**:597-605. DOI: 10.1007/s00424-005-1487-2, PMID: 1866268.

Pajor AM. 2014. Sodium-coupled dicarboxylate and citrate transporters from the SLC13 family. *Pflugers Arch* **466**:119-130. DOI: 10.1007/s00424-013-1369-y, PMID: 24114175.

Pajor AM, Sun N. 1996. Functional differences between rabbit and human Na(+)-dicarboxylate cotransporters, NaDC-1 and hNaDC-1. *Am J Physiol* **271**:F1093-1099. DOI: 10.1152/ajprenal.1996.271.5.F1093, PMID: 8946005.

Yamamoto S, Inoue K, Murata T, Kamigaso S, Yasujima T, Maeda JY, Yoshida Y, Ohta KY, Yuasa H. 2010. Identification and functional characterization of the first nucleobase transporter in mammals: implication in the species difference in the intestinal absorption mechanism of nucleobases and their analogs between higher primates and other mammals. *J Biol Chem* **285**:6522-6531. DOI: 10.1074/jbc.M109.032961, PMID: 2825448.

Yee S. 1997. In vitro permeability across Caco-2 cells (colonic) can predict in vivo (small intestinal) absorption in man--fact or myth. *Pharm Res* **14**:763-766. DOI: 10.1023/a:1012102522787, PMID: 9210194.

344
345

Figure 1.

Figure 2.

A

mRNA level
(fold increase in brain expression)

Brain, Thymus, Salivary glands, Lung, Heart, Liver, Kidney, Spleen, Pancreas, Stomach, Duodenum, Jejunum, Ileum, Colon, Skeletal muscle, Placenta, Testis, Epididymis, Prostate

B

SLC35G1 ATP1A1 DAPI Merge

C

Uptake rate (pmol/min/mg protein)

Control #1 #2 #3
SLC35G1

D

SLC35G1 mRNA level (% of control)

Control #1 #2 #3
SLC35G1

E

F

Luminal side Blood side

$[Na^+]_o$: 150 mM $[Na^+]_i$: 5 mM $[Cl^-]_i$: 5 mM $[Cl^-]_o$: 120 mM

NaDC1   SLC35G1

Concentrative accumulation   Cl⁻-sensitive facilitative transport

pH: 5.5   pH: 7.4   pH: 7.4