Human CD29+/CD56+ myogenic progenitors display tenogenic differentiation potential

and facilitate tendon regeneration

Xiexiang Shao[1†], Xingzuan Lin[3†], Hao Zhou[1†], Minhui Wang[6†], Lili Han[5], Xin Fu[1], Sheng

Li[1], SiyuanZhu[1], Shenao Zhou[5], Wenjun Yang[1*], Jianhua Wang[1*], Ping Hu[1, 2, 4, 5*]

Affiliations:

1. Xinhua Hospital affiliated to Shanghai Jiao Tong University School of Medicine.

Shanghai,20023, China.

2. Guangzhou Laboratory, Guangzhou, 510005, China.

3. Department of Sports Medicine, Peking University Third Hospital, Institute of Sports

Medicine of Peking University, Beijing Key Laboratory of Sports Injuries, Beijing, China.

4. Institute for Stem Cell and Regeneration, Chinese Academy of Sciences, Beijing ,

100101, China.

5. State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell

Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences,

Shanghai 200031, China.

6. Department of Bioinformatics, Fujian Key Laboratory of Medical Bioinformatics. School

of Medical Technology and Engineering. Fujian Medical University, Fuzhou 350122, China

*Corresponding author: Wenjun Yang, Email: wjyang@sibcb.ac.cn; Jianhua Wang, Email:

wangjianhua@xinhuamed.com.cn; PingHu, Email: hup@sibcb.ac.cn;

[†]These authors contributed equally to the current study.

Abstract

Tendon injury occurs at high frequency and is difficult to repair. Identification of

human stem cells being able to regenerate tendon will greatly facilitate the development of

regenerative medicine for tendon injury. We identified human CD29+/CD56+ myogenic

progenitors having tendon differentiation potential both in vitro and in vivo. Transplantation

of human CD29+/CD56+ myogenic progenitors contributes to injured tendon repair and

thus improves locomotor function. Interestingly, the tendon differentiation potential in

mouse muscle stem cells is minimal and the higher TGFβ signaling level may be the key

for the distinct feature of human CD29+/CD56+ myogenic progenitors. These findings

reveal that human CD29+/CD56+ myogenic progenitors are bi-potential adult stem cells

and can serve as a new source for tendon regeneration.

Introduction

Skeletal muscle is a tissue with great regeneration ability due to the existence of

muscle stem cells (MuSCs). MuSCs are adult stem cells located at the periphery of

myofibers between the basal lamina and the plasmalemma of the myofibers and

responsible for muscle regeneration[1,2]. MuSCs have been considered to be unipotent

stem cells to have a sole differentiation potential to myofibers[3]. MuSCs undergo

expansion and differentiate to multinuclei myofibers after injury in vivo. MuSCs have

remarkable abilities to support muscle regeneration. After transplantation, the engrafted

mouse MuSCs go through active expansion and regenerate myofibers[4]. Isolation by

FACS and expansion of MuSCs have been reported from several species including

mouse, pig, and human[5,6]. Single cell sequencing analysis from human skeletal muscles

have also revealed the existence of Pax7+ myogenic progenitors [6,7]. Due to the different

motion patterns, the regeneration capacity may be different between human and rodents.

Current investigations have suggested that human myogenic progenitor cells didn't share

same markers as that in mice[8], and the expression pattern of oxidative enzymes and

cytokines between these two species are also different[9], suggesting that human muscle

progenitor cells may have distinct features from mice.

FACS method to isolate cells with myogenic differentiation potential from human

muscle biopsies has been established. It has been reported that CD56 (NCAM)+ and

CD56+ CD29+ cells from skeletal muscle display myogenic differentiation potential[9,10,11,12].

The CD56+ CD34+ progenitor cells isolated from human skeletal muscle biopsies have

been reported to have chondrogenic, adipogenic, and osteogenic potentials besides

myogenic potential[13-15]. CD56+CD34- progenitor cells are free of adipogenic potential[11].

The differentiation potential of human myogenic progenitors remains to be further

explored.

Skeletal muscle directly connects to tendons which is responsible for transmitting

forces from skeletal muscle to bone to generate active movement. Tendinopathy affects

more than 10% of the population under 45 and compromises the tendon functions[16,17].

Tendon injury healing is slow and incomplete due to the low number of cells in tendon and

the hypovascular and anaerobic environment[18,19]. Tendon stem/progenitor cells (TDSCs)

is a cell population derived from tendon and considered to be a subgroup of mesenchymal

stem cells that have abilities to improve tendon injury healing[20,21]. However, the number of

TDSCs in tendon is low and retrieving TDSCs is invasive and exacerbates tendon injury.

Morphology and proliferation ability loss in the culture system also hampers the efforts to

obtain sufficient amounts of active TDSCs. To find more cell types supporting tendon

regeneration will facilitate the development of regenerative medicine to treat tendon injury.

The activity of tenogenesis is tightly regulated by many signaling pathways,

especially for TGFβ signaling[22]. TGFβ signaling is indispensable for tendon

development[23], and it could systematically promote the tenogenic differentiation of stem

cells[24,25]. The downstream effectors SMAD2 and SMAD3 are able to activate the

transcription of tendon-specific genes which further facilitates tendon development and

tenogenic differentiation[22,26,27]. After tendon injury, TGFβ signaling promotes the

proliferation, migration and differentiation of TDSCs[28], increases tendon collagen

synthesis[29], and contributes to matrix anabolism for tendon remodeling[30].

Here we found that human CD29+/CD56+ myogenic progenitors can be

differentiated to tendon cells both in vitro and in vivo. Transplantation of human

CD29+/CD56+ myogenic progenitors to injured tendon in mice improved the tendon

regeneration, suggesting its bipotential ability. Interestingly, the tendon differentiation

potential in mouse muscle stem cells is minimal and the higher TGFβ signaling level may

be the key for the distinct feature of human CD29+/CD56+ myogenic progenitors.

Results

The CD29+/CD56+ cells isolated from human muscle biopsies display robust

features of myogenic progenitors.

To analyze the cell components of human skeletal muscle, single cell sequencing

analysis was performed using skeletal muscle biopsy. A total of 57,193 cells were included

for analysis. The cells were grouped to 9 cell populations (Fig. 1a and b). Consistent with

the previous single cell sequencing results from human muscles[31], Fibro/Adipogenic

progenitors (FAPs), endothelial cells, myocytes, and myogenic progenitors were among

the identified cell types (Fig. 1a and b). Especially, myogenic progenitors accounted for

12.4% of all the mononuclear cells, while tenocytes only accounted for 0.06% (Fig. 1c). To

investigate the identity of CD29+/CD56+ cells, joint expression analysis was performed

(Fig. 1d). The scRNA-seq data revealed that all the CD29+/CD56+ cells were myogenic

progenitors, which occupied 19.3% of all the myogenic progenitors (Fig. 1e). However,

there existed no tenocytes with CD29+/CD56+ (Fig. 1d). Combined, the scRNA-seq data

revealed human CD29+/CD56+ cells were myogenic progenitors.

To confirm the single cell analysis results, we first isolated CD29+/CD56+ myogenic

progenitors from human muscle biopsy using FACS as described previously[6,12].

CD31-/CD45-/CD29+/CD56+ cells were collected from the single cell suspension

dissociated from muscle tissue. The collected cells were stained with anti-PAX7,

anti-MYOD1 and anti-MYF5 antibody to confirm the purity of isolated CD29+/CD56+ cells,

and nearly all of the obtained cells were positively stained with these myogenic

progenitors markers (Fig. 1f and g). The proliferation assay indicated high in vitro

expansion ability of these human CD29+/CD56+ myogenic progenitors (Fig. 1h). When

induced to differentiate by 0.4% Ultroser G, human CD29+/CD56+ myogenic progenitors

differentiate to myotubes robustly. Approximately 90% of nuclei were present in Myosin

heavy chain (MyHC) positive myotubes (Fig. 1i and j). Consistently, the expression of

PAX7, MYF5 and MYOD1 were enriched in the myogenic progenitors, and their

expression decreased after differentiation as shown by RT-qPCR assays (Fig. 1k). In

contrast, the expression of genes marking differentiation such as Myogenin (MYOG),

Myosin heavy chain 1 (MYH1), and Myosin heavy chain 3 (MYH3) were significantly

up-regulated in differentiated myotubes (Fig. 1l). Taken together, these results suggest

that the CD29+/CD56+ cells isolated from human muscle biopsies have robust features of

myogenic progenitors.

Human CD29+/CD56+ myogenic progenitors display tenogenic differentiation potential in vitro

Since previous studies indicated that there could be tenogenic differentiation ability for muscle-derived cells[32-34], we next explored the tenogenic differentiation potential of CD29+/CD56+ myogenic progenitors. The isolated primary human CD29+/CD56+ myogenic progenitors were induced to differentiate to tenocytes by 100ng/ml GDF5, 100ng/ml GDF7 and 0.2mM ascorbic acid for 12 days. After 12 days of tendon differentiation, the morphology of cells displayed dramatic differences from those undergoing myogenic differentiation (SFig. 1a). Furthermore, the expression of tendon markers such as TNC and SCX was significantly increased (Fig. 2a, b, and c). Moreover, some other tendon related genes, such as COL I, MKX, THBS4, and COMP, were also enriched after tendon differentiation (Fig. 2c). In contrast, the expression of these genes was not enriched upon myogenic differentiation (Fig. 2a, b, and c). The expression of genes marking myogenic differentiation such as MYOG and MyHC were only detected in a small portion of cells after tenogenic differentiation (SFig. 1b and c). Compared to the over 90% of differentiation efficiency upon myogenic differentiation, only about 20% of cells showed MYOG and MyHC expression after 12 days of tenogenic induction (SFig. 1b and c). The genes marking differentiated myotubes such as MYH1, MYH3, DESMIN, and MYL1 showed moderate elevation after tenogenic differentiation, while dramatic upregulation of these genes was observed after myogenic differentiation (SFig. 1d and e). These results combined suggest that human CD29+/CD56+ myogenic progenitors are capable of tendon differentiation in vitro.

To further confirm the results, we next performed clonal analysis. The freshly isolated primary human CD29+/CD56+ myogenic progenitors were seeded to 96 well plate with the concentration of 1 cell/well. The wells of single cell were allowed to proliferate for 10 days followed by tenogenic or myogenic induction. The plates with myogenic induction were differentiated for 4 days and immunofluorescence staining of MyHC was performed to determine myogenic differentiation. The plates with tenogenic induction were differentiated for 12 days and immunofluorescence staining of SCX was performed to

determine tenogenic differentiation (Fig. 2d). The number of wells showing positive MyHC

staining was counted and the myogenic differentiation efficiency was calculated (Fig. 2e

and g). MyHC+ myotubes was observed in over 95% of wells with alive cells (Fig. 2g).

Similarly, the number of wells displaying positive SCX staining were counted and the

tenogenic differentiation efficiency was calculated (Fig. 2f and g). Approximately 40% of

CD29+/CD56+ myogenic progenitors displayed tenogenic differentiation ability (Fig. 2g),

suggesting that human CD29+/CD56+ myogenic progenitors have tenogenic

differentiation potential. Taken together, these results suggest that human CD29+/CD56+

myogenic progenitors have dual differentiation potentials towards muscle or tendon in

vitro.

Tenocytes differentiated from human CD29+/CD56+ myogenic progenitors display

similar expression profile to primary tenocytes

RNA sequencing was then performed to further determine the lineage of the

tenocytes differentiated from human CD29+/CD56+ myogenic progenitors. Human

CD29+/CD56+ myogenic progenitors were induced towards myogenic or tenogenic

differentiation respectively. We also isolated primary human tenocytes as described

previously[35]. These cells were subjected for RNA sequencing analysis. The differentiated

myotubes and tenocytes displayed distinct expression patterns (Fig. 3a). The

differentiated tendon cells share high similarity to fresh isolated tenocytes from human

tendons (Fig. 3a), suggesting that human CD29+/CD56+ myogenic progenitors are

capable of tendon differentiation. Consistently, two distinct sets of genes representing

myogenic markers and tenogenic markers were up-regulated after myogenic induction

and tenogenic induction, respectively (Fig. 3b and c). These results suggest that human

CD29+/CD56+ myogenic progenitors are capable of dual direction differentiation.

Further GO analysis also displayed the activation of two distinct sets of cell features.

Upon the skeletal muscle differentiation, terms related to skeletal muscle functions such

as skeletal muscle thin filament assembly, skeletal muscle contraction, muscle organ

development, and sarcomere organization were enriched (Fig. 3d), suggesting the

muscular identity of the differentiated cells. In contrast, terms related to tendon formation,

tendon development, and tendon cell differentiation were enriched after tenogenic

differentiation (Fig. 3e), suggesting that tendon identity is achieved in the differentiated

cells. Together, these results suggest that human CD29+/CD56+ myogenic progenitors

are capable of both myogenic and tenogenic differentiation in vitro.

Murine MuSCs display poor tenogenic differentiation ability

We next investigated whether the tenogenic differentiation ability is present in rodent

muscle stem cells (MuSCs). Mouse MuSCs were isolated by positive marker of Vcam1 as

described previously[6] and induced for tenogenic differentiation. In sharp contrast to

human CD29+/CD56+ myogenic progenitors, murine MuSCs failed to be induced to

tendon cells upon the same induction condition as that for human CD29+/CD56+

myogenic progenitors, though the myogenic differentiation is as efficient as the human

CD29+/CD56+ myogenic progenitors. After myogenic differentiation, 93.9% of nuclei were

present in MyHC+ myotubes (Fig. 4a and b). In sharp contrast, no Scx+ cells were

observed after 12 days of induction for tenogenic differentiation (Fig. 4a). Consistent with

the immunofluorescence staining results, RT-qPCR results revealed that myogenic

differentiation marker genes such as MyoG, Myh1, and Myh3 were significantly

up-regulated under both myogenic and tenogenic differentiation conditions (Fig. 4c),

suggesting that murine MuSCs predominantly commit myogenic differentiation under

induction. Different from human CD29+/CD56+ myogenic progenitors, the expression of

genes indicating tendon cell fate such as Scx, Tnc, Col I, Mkx, and Thbs4 did not increase

after tenogenic differentiation (Fig. 4d), suggesting the failure to induce tenogenic cell fate

from murine MuSCs. Together, these results suggest that murine MuSCs display almost

no tenogenic differentiation potential in vitro.

We next performed the lineage tracing experiments in mice to further examine the

tenogenic potential of mouse MuSCs in vivo (Fig. 4e). Pax7CreERT2 mice were crossed

to flox-Stop-flox-tdTomato mice. MuSCs and the descendants of MuSCs will be labeled by

tdTomato (SFig. 2a). Tendon injury in mice was generated by mimicking the peroneus

longus tendon removal surgery in human (SFig. 2b). In human, it has been reported that

the tendon could be regenerated to some extent after the peroneus longus tendon

removal surgery based on MRI imaging[32]. In this surgery, injury of the skeletal muscle

adjacent to the removed tendon was inevitable. The accompanied skeletal muscle injury

could activate MuSCs and make them available for tendon regeneration. To further

activate MuSCs to guarantee that sufficient amount of activated MuSCs were available

around the tendon injury site, we also injected 15µl of 10µM cardiotoxin (CTX) at the

muscle adjacent to the sites where the tendon was removed to induce more muscle

injuries and further activated MuSCs. If MuSCs can participate tendon regeneration,

tdTomato+ tendon cells would be observed after the repair of tendon injury.

Expectedly, large amount of tdTomato+ myofibers were observed after muscle injury

(SFig. 2c-d), suggesting that the tracing system works well. Nevertheless, less than 0.2%

tendon cells originated from mouse MuSCs were observed even four months after tendon

removal (Fig. 4f-g). These results suggest that murine MuSCs have poor tendon

differentiation abilities.

224   Transplantation of human CD29+/CD56+ myogenic progenitors facilitates tendon

225   regeneration

We next went on to investigate whether the transplantation of human CD29+/CD56+

myogenic progenitors to mouse can improve tendon regeneration. An approximately

1.5mm long and 0.5mm width transverse incision was performed at 5mm from the

calcaneus in Achilles tendon for NOD/SCID immunodeficient mice. Total 50,000 human

CD29+/CD56+ myogenic progenitors packed in hydrogel were planted at the incision site

(Fig. 5a). As a control, mixture of PBS and hydrogel was transplanted in the SCID

recipient mice undergoing the same tendon injury at the cleavage sites. Packing the cells

with hydrogel concentrated the transplanted cells at the local injury sites. Two months

after transplantation, the tendons carrying transplanted with or without human

CD29+/CD56+ myogenic progenitors were harvested, respectively (Fig. 5a).

In mice transplanted with human CD29+/CD56+ myogenic progenitors, continuous

cryosections containing muscle and tendon tissues were generated. Immunofluorescence

staining was performed to detect tendon and muscle markers with two cryosections

adjacent to each other, respectively. The two sets of images obtained on continuous

cryosections were superimposed on each other to pinpoint the position of tendons.

Immunofluorescence staining of antibody specifically recognizing human TNC indicated

the position of regenerated tendon-like tissue from human cells in the harvested tissues

(Fig. 5b). The presence of human cells was also illustrated by the immunofluorescence

staining with the antibody specifically recognizing human Lamin A/C. In the control PBS

injection group, where the mixture of PBS and hydrogel instead human CD29+/CD56+

myogenic progenitors was injected to the incision sites, no human Lamin A/C was

detected. These results confirmed the specificity of human TNC and Lamin A/C antibody.

Immunofluorescence staining revealed that over 75% of the human cells showed TNC

expression (Fig. 5b and d), suggesting that the majority of the transplanted human

CD29+/CD56+ myogenic progenitors differentiate to tendon cells in vivo. The

immunofluorescence staining of MyHC and human Lamin A/C were performed to detect

the muscle cells originated from the transplanted human CD29+/CD56+ myogenic

progenitor cells. Only about 12.8% of the human cells detected expressed MyHC (Fig. 5c

and d). Moreover, the human cells were predominantly enriched at the tendon region (Fig.

5b). Only a few MyHC+ cells originated from the transplanted human CD29+/CD56+

myogenic progenitors scattered in the muscle region (Fig. 5c). To further confirm the

tendon differentiation potential of the transplanted human CD29+/CD56+ myogenic

progenitors, immunofluorescence staining of SCX and TNMD was also performed. The

majority (80.0% and 74.6%) of transplanted human CD29+/CD56+ myogenic progenitors

also expressed SCX and TNMD (SFig. 3a, b and c). Furthermore, Col I was

predominantly expressed in the regenerated tendon-like tissue rather than Col III after

human cells transplantation (Fig. 5e), indicating human CD29+/CD56+ myogenic

progenitors could contribute to structural repair of injured tendon and facilitate the healing

process. Taken together, these results suggest that human CD29+/CD56+ myogenic

progenitors are capable of tendon differentiation in vivo and contributing to tendon

regeneration.

In sharp contrast to human CD29+/CD56+ myogenic progenitors, when 50,000

murine MuSCs constitutively expressing tdTomato were transplanted to pre-injured

tendon in NOD/SCID mice under the same condition, less than 0.3% of tdTomato+ TNC+

cells were detected (Fig. 5f and g). However, the myogenic differentiation potential of

human CD29+/CD56+ myogenic progenitors and mouse MuSCs was similar. Muscle

injury was induced by muscular injection of CTX in tibialis anterior (TA) muscle in

NOD/SCID mice. TA muscles were irradiated to kill the local MuSCs as described

previously[36]. Transplantation of human CD29+/CD56+ myogenic progenitors and mouse

MuSCs to the irradiated pre-injured recipient mice was performed, respectively. TA

muscles were harvested after 28 days. Transplantation of both human CD29+/CD56+

myogenic progenitors and murine MuSC displayed robust engraftment efficiency (SFig.

3d-i). These results suggest that human CD29+/CD56+ myogenic progenitors and murine

MuSCs have similar myogenic differentiation potential.

Combined, the above results suggest that human CD29+/CD56+ myogenic

progenitors have dual differentiation potentials towards myogenesis or tenogenesis in vivo,

while murine MuSCs predominantly commit myogenesis.

284 Transplantation of human CD29+/CD56+ myogenic progenitors improves locomotor

285 function after tendon injury

We next checked whether transplantation of human CD29+/CD56+ myogenic

progenitors could improve tendon functions. First, transmission electron microscope

analysis was used to evaluate microstructure of injured tendon two months after

transplantation. Larger and denser collagen fibrils of the tendons were identified in

transplanted group than control group with PBS injection, although the maturation level of

collagen fibrils could still not reach to those in uninjured tendon (Fig. 6a and b). We next

examined whether transplantation of human CD29+/CD56+ myogenic progenitors could

improve the biomechanical property of tendon. The max failure load and stiffness of the

tendons from the transplantation group were significantly better than those from PBS

injection control group (Fig. 6c). These results combined suggest that transplantation of

human CD29+/CD56+ myogenic progenitors improves the collagen fibril maturation and

biomechanical property of the injured tendon.

Whether the improved tendon regeneration could facilitate the whole organism

locomotor function was next investigated. Since the Achilles tendon transmits the

plantarflexion force from gastrocnemius muscle to calcaneus, the plantarflexion force of

involved leg was also performed two months after tendon injury. Expectedly,

transplantation of human CD29+/CD56+ myogenic progenitors for injured tendon also

contributed to improving both twitch and tetanus plantarflexion force when compared with

PBS injection control group, although the plantarflexion force could still not reach to the

level of uninjured leg (Fig. 6d). Consistent with the improved plantarflexion force, the

endurance time and max fatigue speed of the mice transplanted with human

CD29+/CD56+ myogenic progenitors were better in treadmill test when compared to PBS

group (Fig. 6e). Since immunofluorescence staining with human Lamin A/C revealed that

only a small number of human CD29+/CD56+ myogenic progenitors engrafted in muscle

sporadically (Fig. 5c and d), these results suggest that the transplanted human

CD29+/CD56+ myogenic progenitors can improve locomotor function by directly repairing

injured tendon.

TGFβ signaling pathway contributes to tenogenic differentiation of human

CD29+/CD56+ myogenic progenitors

Since the human CD29+/CD56+ myogenic progenitors and murine MuSCs shared

the same strain of recipient mice and the same tendon injury while being transplanted,

they had the similar microenvironment. Therefore, the distinct differentiation potentials are

due to the cell intrinsic differences between species. We further compared the expression

profiles of human CD29+/CD56+ myogenic progenitors and murine MuSCs. Interestingly,

TGFβ signaling was identified in KEGG enrichment analysis of upregulated genes in

human CD29+/CD56+ myogenic progenitors when compared with murine MuSCs (Fig.

7a), indicating that TGFβ signaling could be the key node for maintaining the tendon

differentiation potential. Especially, SMAD2 and SMAD3 were identified in upregulated

gene set which was enriched in TGFβ signaling pathway (Fig. 7b). Since

TGFβ/SMAD2/SMAD3 axis plays a crucial role in tendon development and tenogenic

differentiation[22,23,26], these data further indicated potential crucial role of TGFβ signaling

for tenogenic differentiation ability of human CD29+/CD56+ myogenic progenitors. Then

TGFβ signaling inhibitor SB-431542 was used to investigate the biological effect of TGFβ

signaling during tenogenic induction. The immunofluorescent staining, western blot assay

and RT-qPCR results showed SB-431542 significantly suppressed expression level of

tendon related genes SCX, TNC, COL I, MKX, and THBS4 (Fig. 7c, d, e). On the contrary,

myogenic differentiation ability of human CD29+/CD56+ myogenic progenitors was

increased after treatment of SB-431542 during tenogenic induction (Fig. 7f and g). Taken

together, these data indicated that TGFβ signaling pathway contributes to tenogenic

differentiation of human CD29+/CD56+ myogenic progenitors.

Discussion

Here we show that human CD29+/CD56+ myogenic progenitors have dual

differentiation potentials towards muscle and tendon. Transplantation of human

CD29+/CD56+ myogenic progenitors contributes to injured tendon repair and thus

improves locomotor function. Thus, the human CD29+/CD56+ myogenic progenitors

could be served as a new source for tendon regeneration.

Tendon disorders widely occur in people of all ages[22]. It disrupts the stability and

mobility of joint, which deeply affects their locomotor function and quality of life. However,

the natural healing of injured tendon is very slow due to hypocellularity and

hypovascularity of tendon. The biomechanical property and structural integrity could be

hardly completely recovered even with surgical treatment[37,38]. It is still a great challenge in

clinical work to treat tendon injury.

The relative inefficient outcome of routine therapy for tendon injury sparked the

exploration of stem cell treatment. Seed cells with the ability to differentiate into tenocytes

and secrete paracrine factors to repair tendon injury are preferred. Thus, tendon derived

stem cells, embryonic stem cells, induced pluripotent stem cells and mesenchymal stem

cells have been introduced as seed cells to treat tendon injury[39]. However, inadequate

sources of tendon derived stem cells, ethical issue and risk of teratoma formation of

embryonic stem cells or induced pluripotent stem cell, and heterogeneity of mesenchymal

stem cells limit the development of these seed cells for tendon injury treatment. As for

muscle progenitors, it is high in proliferation and abundant in sources, and donor site

morbidity after muscle harvest was low. Thus, myogenic progenitors might be a promising

candidate as seed cells for tendon repair.

Here we also find that the differentiation potential of muscle stem cells is species

dependent. Human CD29+/CD56+ myogenic progenitors are bipotential adult stem cells.

In contrast, murine muscle stem cells barely have tendon differentiation potential. The

species difference might be due to the higher TGFβ signaling level in human

CD29+/CD56+ myogenic progenitors. It has been reported that TGFβ/SMAD2/SMAD3

axis plays a crucial role in tendon development and tenogenic differentiation[22,23,26].

SMAD2/3-dependent TGFβ signaling also acts as a crucial molecular brake for

myogenesis. It could suppress myogenic regulatory factors Myod1 and Myogenin[40,41], as

well as inhibit myotube fusion and muscle regeneration[42,43]. Thus, the elevated TGFβ

signaling may help inhibit the myogenic differentiation ability and stimulate the tenogenic

differentiation potential of human CD29+/CD56+ myogenic progenitors under specific

microenvironment.

Studies using mouse models contributed to the majority of our knowledge about

muscle stem cells and laid the foundation for our understanding of mammalian muscle

stem cells. However, human is dramatically different from mouse in many aspects such as

size, life span, and manners of motions. It is more demanding for humans to maintain the

homeostasis of the locomotion system and the whole organism locomotion ability in much

longer life span and bigger body size. Though our knowledge about human muscle

regeneration is limited, the current studies have revealed multiple differentiation potentials

of Pax7+ human myogenic progenitors in skeletal muscles. CD56+ CD34+ progenitor

cells in human skeletal muscle have been reported to have myogenic, osteogenic, and

adipogenic activity[14,15]. The CD56+ CD34- progenitor cells in human skeletal muscle have

been shown to be free of adipogenic potential[11]. Our results suggest that the

CD29+/CD56+ myogenic progenitors in skeletal muscles also have tenogenic

differentiation ability besides their myogenic differentiation ability. It seems that there are

multiple subpopulations of myogenic progenitors in skeletal muscle. They are all capable

of muscle regeneration, while with potentials to regenerate other components of the

motion system such as bone, tendon, and adipocytes. This could be an economic method

to maintain the functions of the motion system for the longer life span and more

complicated motion manner in human beings.

Methods

Animals

Animal care and use were in accordance with the guidelines of the animal facility
hosted by Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of
Sciences, and the operations were approved by the ethical committee of Shanghai
Institute of Biochemistry and Cell Biology. All mice were maintained in specific
pathogen-free (SPF) animal facility in individually ventilated cages (IVC) with controlled
temperature (22±1℃) and light (12h light/dark cycle). NOD/SCID mice were purchased
from Animal Model Research Center of Nanjing University. Pax7CreERT2 and
Rosa26-Flox-Stop-Flox-tdTomato mice were purchased from Jackson Laboratory. All
experiments were conducted on 3-month-old adult male mice.

Human samples

405   Peroneal longus muscle and remanent tendon were obtained from the wastes of
406 patients who underwent full-thickness peroneal longus tendon autograft for knee ligament
407 reconstructive procedures. The study was approved by the ethical committee of Xinhua
408 Hospital Affiliated to Shanghai Jiao Tong University, School of Medicine (Approval No.
409 XHEC-D-2019-043) and written informed consents were obtained from all donors.

Isolation of human CD29+/CD56+ myogenic progenitors

Human skeletal muscle from peroneal longus were dissected and digested as
described previously[6]. Briefly, muscle tissues were cut into small pieces and digested by
collagenase II (Worthington biochemical, 700-800U/ml, cat#LS004177) for 60min followed
by 30min of digestion with the mixtures of collagenase II and dispase (Life
Technologies,11U/ml, cat#17105-041). Digested cells were passed 10 times through a
20-gauge needle. Cell suspension was filtered through a 40μm cell strainer (BD Falcon,
cat#352340). Erythrocytes were removed by red blood cell lysis buffer (Thermo Fisher
Scientific, cat#00-433-57). The single cell suspension obtained from human muscle was
stained with a cocktail containing PE-Cy5 anti-human CD45 (BD Pharmingen,
cat#555484), Percp-Cy5.5 anti-human CD31 (BioLegend, cat#303132), AF-488
anti-human CD29 (BioLegend, cat#303016) and PE anti-human CD56 (BioLegend,
cat#304606) for 45 min at 4℃. CD45- CD31- CD29+ CD56+ myogenic progenitors were
sorted by BD Influx sorter (BD Biosciences).

Isolation of mouse muscle stem cells

For mouse muscle stem cells isolation, dissected TA muscles were first digested with

10ml muscle digestion buffer (DMEM containing 1% penicillin/streptomycin, 0.125mg/ml
Dispase II (Roche, 04942078001), and 10mg/ml Collagenase D (Roche, 11088866001))
for 90 minutes at 37°C. The digestion was stopped by adding 2ml of FBS. The digested
cells were filtered through 70µm strainers. Red blood cells were lysed by 7ml RBC lysis
buffer (0.802% NH4Cl, 0.084% NaHCO3, 0.037% EDTA in ddH2O, pH7.2-7.4) for 30s,
then filter through 40µm strainers. After staining with antibody cocktails
(AF700-anti-mouse Sca-1, PerCP/Cy5.5-anti-mouse CD11b, PerCP/Cy5.5-anti-mouse
CD31, PerCP/Cy5.5-anti-mouse CD45, FITC anti-mouse CD34, APC-anti-mouse Integrin
α7+), the mononuclear cells were subjected for FACS analysis using Influx (BD
Biosciences). The population of PI-CD45-CD11b-CD31-Sca1- CD34+ Integrin α7+ cells
was collected.

Primary human tenocytes isolation

Tendon tissues were obtained from the discarded materials of tendon autograft
surgery. They were washed with PBS. Epi- and peri-tendon sheath were completely
removed. Tenocytes were isolated as described previously[20]. Briefly, the tendons were
minced to 1mm^3 pieces and digested with 3 mg/ml collagenase I (Worthington
biochemical, cat# LS004194) in DMEM (Gibco, cat#11965118) at 37°C for 3hrs with
gentle agitation. The digested tissue was filtered through a 40µm cell strainer (BD Falcon,
cat#352340) and the isolated cells were plated for subsequent analysis.

Cell culture and differentiation

Primary human CD29+/CD56+ myogenic progenitors were plated in F10 basal
medium (Gibco, cat#11550043) containing 20% FBS (Gibco, cat#10-013-CV), 2.5ng/ml
bFGF (R&D, cat#233-FB-025) and 1% Penicillin-Streptomycin (Gibco, cat#15140-122) on
collagen-coated dishes. Mouse muscle stem cells were plated in F10 basal medium
(Gibco, cat#11550043) containing 20% FBS (Gibco, cat#10-013-CV), 2.5ng/ml bFGF
(R&D, cat#233-FB-025) and IL-1α, IL-13, IFN-γ and TNF-α as described previously[36].
DMEM (Gibco, cat#11965118) containing 0.4% Ultroser G (Pall Corporation,
cat#15950-17) and 1% Penicillin-Streptomycin were used to differentiate human muscle
stem/progenitor cells[5]. DMEM containing 2% horse serum (HyClone,
cat#HYCLSH30074.03HI) and 1% Penicillin-Streptomycin were used to differentiate
mouse muscle stem cells as described previously[44,45]. DMEM containing 10% FBS,
100ng/ml GDF5 (R&D, cat#8340-G5-050), GDF7 (R&D, cat#8386-G7-050), 0.2mM
ascorbic acid (Sigma-Aldrich, cat#A4403) and 1% Penicillin-Streptomycin were used to
induce tendon differentiation [46-48]. Primary tenocytes were cultured in DMEM containing
20% FBS, 2.5ng/ml bFGF (R&D, cat#233-FB-025) and 1% Penicillin-Streptomycin.

Single-cell RNA sequencing

The single cell suspension of mononuclear cells from human skeletal muscles was

firstly prepared. PI (Sangon Biotech, cat#E607328) and Hoechst (Sangon Biotech,

cat#A601112) were used to sort live cells by FACS. Then the sored cells were washed

twice with PBS containing 0.04% BSA, followed by library preparation with Chromium

Single Cell 3' Reagent Kits (10X genomics, cat# 1000121-1000157). The sequencing was

performed on Illumina Novaseq 6000 platform (Illumina).

Single cell RNA-seq data were analyzed by Seurat R (Version 3.2.0) package. Cells

with less than 200 genes, more than 6,000 genes detected, and more than 10%

mitochondrial genes were excluded. Three samples with a total of 57,193 cells were

included for subsequent analysis. Sequencing reads for each gene were normalized to

total UMIs in each cell to obtain normalized UMI values by "NormalizeData" function. The

"ScaleData" function was used to scale and center expression levels in the data set for

dimensional reduction. Total cell clustering was performed by "FindClusters" function at a

resolution of 0.1 and dimensionality reduction was performed with "RunUMAP" function[49].

Immunofluorescence staining

Cryosections were fixed in 4% formaldehyde for 15min, permeabilized in 0.5% Triton

X-100 for 15min, and stained with anti-Pax7 (Developmental Studies Hybridoma Bank),

anti-Laminin (Abcam, cat#ab11575), anti-Lamin A/C (Abcam, cat#ab108595;

cat#ab190380), anti-TNC (Abcam, cat#ab108930; cat#ab3970), anti-Scx (Abcam,

cat#ab58655), anti-Tnmd (Abcam, cat#ab203676), anti-Col I (Abcam, cat#ab260043),

anti-Col III (ABclonal, cat#A0817) or anti-MyHC (Millipore, cat#05-716) at 4°C overnight,

and incubated with Alexa 488-, Alexa 594 or Alexa 647-labeled anti-mouse or anti-rabbit

secondary antibodies (Invitrogen, 1:1000) at room temperature for one hour. The nuclei

were stained with 4,6-diamidino-2-phenylindole (DAPI, Vector Laboratories, cat#H-1200).

All images were acquired by Leica SP8 confocal microscope (Leica).

Gene expression analysis

Total RNA was isolated using TRIzol Reagent (Invitrogen, cat#15596-018) according

to the manufacturer's instruction. GAPDH served as internal control. The primers for

RT-qPCR are listed as below:

Human GAPDH-F   5'-CAAGGCTGAGAACGGGAAGC-3'

Human GAPDH-R   5'-AGGGGGCAGAGATGATGACC-3'

Human SCX-F	5'-AGCGATTCGCAGTTAGGAGG-3'
Human SCX-R	5'-GTCTGTACGTCCGTCTGTCC-3'
Human COL I-F	5'-GGCTCCTGCTCCTCTTAGCG-3'
Human COL I-R	5'-CATGGTACCTGAGGCCGTTC-3'
Human TNC-F	5'-GGTGGATGGATTGTGTTCCTGAGA-3'
Human TNC-R	5'-CTGTGTCCTTGTCAAAGGTGGAGA-3'
Human THBS4-F	5'-TGCTGCCAGTCCTGACAGA-3'
Human THBS4-R	5'-GTTTAAGCGTCCCATCACAGTA-3'
Human MKX-F	5'-TTCAAGGCAATGCTGAACGG-3'
Human MKX-R	5'-CTCCCGCTTTGATGACCGAA-3'
Human COMP-F	5'-GATCACGTTCCTGAAAAACACG-3'
Human COMP-R	5'-GCTCTCCGTCTGGATGCAG-3'
Human PAX7-F	5'-ACCCCTGCCTAACCACATC-3'
Human PAX7-R	5'-GCGGCAAAGAATCTTGGAGAC-3'
Human MYF5-F	5'-AATTTGGGGACGAGTTTGTG-3'
Human MYF5-R	5'-CATGGTGGTGGACTTCCTCT-3'
Human MYOD1-F	5'-GACGGCATGATGGACTACAG-3'
Human MYOD1-R	5'-AGGCAGTCTAGGCTCGACAC-3'
Human MYOG-F	5'-GCTCAGCTCCCTCAACCA-3'
Human MYOG-R	5'-GCTGTGAGAGCTGCATTCG-3'
Human MYH1-F	5'-GGGAGACCTAAAATTGGCTCAA-3'
Human MYH1-R	5'-TTGCAGACCGCTCATTTCAAA-3'
Human MYH3-F	5'-ATTGCTTCGTGGTGGACTCAA-3'
Human MYH3-R	5'-GGCCATGTCTTCGATCCTGTC-3'
Human DESMIN-F	5'-TCGGCTCTAAGGGCTCCTC-3'
Human DESMIN-R	5'-CGTGGTCAGAAACTCCTGGTT-3'
Human MYL1-F	5'-GTTGAGGGTCTGCGTGTCTTT-3'
Human MYL1-R	5'-ACCCAGGGTGGCTAGAACA-3'
Mouse Gapdh-F	5'-ACCCAGAAGACTGTGGATGG-3'
Mouse Gapdh-R	5'-ACACATTGGGGGTAGGAACA-3'
Mouse Col I-F	5'-CCAGCGAAGAACTCATACAGC-3'
Mouse Col I-R	5'-GGACACCCCTTCTACGTTGT-3'
Mouse Scx-F	5'-GAGAACACCCAGCCCAAAC-3'
Mouse Scx-R	5'-TCACCCGCCTGTCCATC-3'
Mouse Tnc-F	5'-GAGCCCCTTTGCCTCAACAA-3'
Mouse Tnc-R	5'-CTTCGCCCGTGAAACCTTCTT-3'
Mouse Mkx-F	5'-TGGTTTCCTGGACAATCCACA-3'
Mouse Mkx-R	5'-CGCTTATGCCTTACCTTCCCTC-3'

Mouse Thbs4-F	5'-GCCACAAGCACAGGAGACTTT-3'
Mouse Thbs4-R	5'-TGACCTGCTGCCTCAGAAGA-3'
Mouse MyoG-F	GAGACATCCCCCTATTTCTACCA
Mouse MyoG-R	GCTCAGTCCGCTCATAGCC
Mouse Myh1-F	5'-GCGAATCGAGGCTCAGAACAA-3'
Mouse Myh1-R	5'-GTAGTTCCGCCTTCGGTCTTG-3'
Mouse Myh3-F	5'-ATGAGTAGCGACACCGAGATG-3'
Mouse Myh3-R	5'-ACAAAGCAGTAGGTTTTGGCAT-3'

Cloning assay

Primary human CD29+/CD56+ myogenic progenitors were first sorted in 96 well

plates with density of single cell per well. Tenogenic induction or myogenic induction was

performed after proliferation for each well. After induction, the immunofluorescence

staining of SCX or MyHC was performed in each well. The differentiation efficiency was

determined by calculating the ratio of total wells with positive fluorescence signal to total

wells with alive cells.

RNA-sequencing

Total RNA was isolated using TRIzol Reagent (Invitrogen, cat#15596-018) according

to the manufacturer's instruction. mRNA was enriched with magnetic oligo (dT) beads

(New England Biolabs, cat#S1419S). The cDNA library was constructed with mean inserts

of 200bp with non-stranded library preparation using NEBNext Ultra RNA Library Prep Kit

for Illumina (New England Biolabs, cat#E7530L). Sequencing was performed by a

paired-end 125 cycles rapid run on the Illumina HiSeq2500. Sequencing data were filtered

by SolexaAQ (Q > 20 and length ≥ 25 bp)[50]. The adapter sequences and low quality

segments (Phred Quality Score<20) were trimmed using Cutadapt. Paired-end clean

reads were then mapped to the reference genome GRCh38.98 using HISAT2.

Htseq-count was used to quantify the gene expression value[51]. Read count of each gene

was normalized using FPKM (Fragments Per Kilo bases per Million fragments).

Differential expression (DE) analysis was performed using DESeq and significant DE

genes were defined as those with absolute log2FoldChange>1 and p< 0.05. Heatmap and

volcano plot of DE genes was generated by R package Pheatmap

(https://cran.r-project.org/web/packages/pheatmap/index.html) and EnhancedVolcano

(https://bioconductor.org/packages/release/bioc/vignettes/EnhancedVolcano/inst/doc/Enh

ancedVolcano.html), respectively. Gene enrichment analysis was conducted by R

package topGO with p < 0.05 as the cut-off. The differential expression analysis for

RNA-seq data between human CD29+/CD56+ myogenic progenitors and mouse MuSCs

was performed according to previous literature[52].

Cell transplantation in TA muscle

A single dose of 18 Grey irradiation was administered to the hind legs of the recipient

NOD/SCID mice. TA muscle was injured by injecting 15μl of 10μM CTX (Sigma), and

50,000 human CD29+/CD56+ myogenic progenitors or 50,000 murine MuSCs suspended

in 10μl PBS were injected intramuscularly to the injury site as described [5].

Tendon injury for lineage tracing

Pax7^{CreERT2}:Rosa26^{tdTomato} mice were obtained and MuSC was labeled with tdTomato

as aforementioned. We used dedicated surgical instruments to imitate a mouse model

similar to previously published human surgery[32]. Firstly, distal gastrocnemius tendon was

woven with a 6-0 polypropylene non-absorbable suture (PROLENE, cat#EH7242H) and

released by microsurgical scissor. After that, a dedicated mini tendon stripper was

introduced over the free end of distal medial gastrocnemius tendon. The medial

gastrocnemius tendon could be totally removed after advancement of tendon stripper. To

further activate MuSCs to guarantee that sufficient amount of activated MuSCs were

available around the tendon injury site, 15μl of 10μM CTX cardiotoxin (CTX) was injected

at the muscle adjacent to the sites where the tendon was removed to induce more muscle

injuries and further activate MuSCs. Then the subcutaneous tissue and skin were sutured.

Tendon injury and cell transplantation

An approximately 1.5mm long and 0.5mm width transverse incision was performed at

5mm from the calcaneus in Achilles tendon for recipient NOD/SCID mice. 50,000 human

CD29+/CD56+ myogenic progenitors or mouse MuSCs resuspended in 20μl hydrogel

were injected to the injury site with 28-gauge needles. PBS mixed with 20μl hydrogel were

injected as control.

Biomechanical analysis

Two months after tendon injury, the tendons were harvested to biomechanical analysis using a universal tester (Instron 3345 load system, USA). The grippers were gradually moved apart with the speed of 5 mm/min until the tested tendon was completely ruptured. Then the max load was obtained and documented. The slope of the stress-strain curve was defined as stiffness. All these data were automatically presented in Instron 3345 load system.

Transmission electron microscope (TEM) examination

After washing the selected target tendon tissues with PBS, tissue fixation was performed for more than 2 hours. Post-fixation was conducted by 1% OsO_4 (Ted Pella Inc) in 0.1M PB for 1-2 hours. Then dehydration and drying were performed and target tendon samples were subsequently attached to metallic stubs for conductive metal coating. Images were obtained by TEM (HITACHI, SU8100) and at least 1000 collagen fibrils were evaluated for each sample. Density of fibrils was evaluated by the percentage of collagen fibril containing area.

Treadmill test

Mouse treadmill test was performed in Exer-3/6 treadmill apparatus with electrical stimulus. Treadmill exercise began with a 5 min warm-up, then each mouse was evaluated with specific protocol. For endurance test, mice ran on the treadmill at the constant speed of 22m/min until exhaustion. For exhaustion test, the treadmill speed was increased (2m/min each 3 min) at the beginning speed at 18m/min until exhaustion. Exhaustion was defined as inability to run on the treadmill for longer than10 seconds.

In vivo muscle force analysis

The 1300A 3-in-1 whole animal system (Aurora Scientific) was used for in vivo muscle force analysis. Mice were first anesthetized and kept warm by heat lamp. The foot was placed on a footplate and kept perpendicular to the tibia. The peroneal branch of the sciatic nerve was stimulated to evaluate the plantarflexion strength. Five repetitive tests were performed for each limb and DMA software (Aurora scientific) was used for results

analysis.

Statistical analysis

At least three biological replicates and technical repeats were performed in each
experimental group. All experiments were analyzed and evaluated by investigators in a
blinded manner. Error bars indicated standard deviation. Two-tailed unpaired Student's
t-tests were used when variances were similar (tested with F-test) for comparison of two
independent groups. One-way ANOVA tests followed by Dunnett's post-test or Tukey's
post-test were used for multiple comparisons. Shapiro-Wilk tests were performed to
determine data normality. Statistical analysis was performed in GraphPad Prism 7
(GraphPad Software, San Diego, USA) or SPSS version 19.0 for Windows (SPSS Inc.,
Chicago, IL, USA). It was considered significant with P value less than 0.05. Data were
presented as mean ± standard unless stated.

Accession numbers

The complete sequencing data have been uploaded on to Sequence Read Archive
database (PRJNA1178160, PRJNA1012476 and PRJNA1012828).

Acknowledgements

We thank Dr. Dangsheng Li for helpful discussions, the National Protein Science
Center (Shanghai) for helps on FACS sorting, the cell biology facility of SIBCB for helps on
imaging and FACS analysis. This work was supported by the Strategic Priority Research
Program of the Chinese Academy of Science (XDA16020400), National Science
Foundation of China (32170804, 81871096, 82372384, 82302657).

References

1.   Fu, X., Wang, H., and Hu, P. (2015). Stem cell activation in skeletal muscle
regeneration. Cell Mol Life Sci *72*, 1663-1677. 10.1007/s00018-014-1819-5.

2.   Relaix, F., Bencze, M., Borok, M.J., Der Vartanian, A., Gattazzo, F., Mademtzoglou, D.,
Perez-Diaz, S., Prola, A., Reyes-Fernandez, P.C., Rotini, A., and Taglietti, t. (2021).
Perspectives on skeletal muscle stem cells. Nat Commun *12*, 692.
10.1038/s41467-020-20760-6.

3.   Seale, P., Asakura, A., and Rudnicki, M.A. (2001). The potential of muscle stem cells.

Dev Cell *1*, 333-342. 10.1016/s1534-5807(01)00049-1.

4.   Collins, C.A., Olsen, I., Zammit, P.S., Heslop, L., Petrie, A., Partridge, T.A., and

Morgan, J.E. (2005). Stem cell function, self-renewal, and behavioral heterogeneity of

cells from the adult muscle satellite cell niche. Cell *122*, 289-301.

10.1016/j.cell.2005.05.010.

5.   Ding, S., Wang, F., Liu, Y., Li, S., Zhou, G., and Hu, P. (2017). Characterization and

isolation of highly purified porcine satellite cells. Cell Death Discov *3*, 17003.

10.1038/cddiscovery.2017.3.

6.   Shao, X., Fu, X., Yang, J., Sui, W., Li, S., Yang, W., Lin, X., Zhang, Y., Jia, M., Liu, H.,

et al. (2023). The asymmetrical ESR1 signaling in muscle progenitor cells determines

the progression of adolescent idiopathic scoliosis. Cell Discov *9*, 44.

10.1038/s41421-023-00531-5.

7.   Barruet, E., Garcia, S.M., Striedinger, K., Wu, J., Lee, S., Byrnes, L., Wong, A.,

Xuefeng, S., Tamaki, S., Brack, A.S., and Pomerantz, J.H. (2020). Functionally

heterogeneous human satellite cells identified by single cell RNA sequencing. Elife *9*.

10.7554/eLife.51576.

8.   Boldrin, L., Muntoni, F., and Morgan, J.E. (2010). Are human and mouse satellite cells

really the same? J Histochem Cytochem *58*, 941-955. 10.1369/jhc.2010.956201.

9.   Bareja, A., Holt, J.A., Luo, G., Chang, C., Lin, J., Hinken, A.C., Freudenberg, J.M.,

Kraus, W.E., Evans, W.J., and Billin, A.N. (2014). Human and mouse skeletal muscle

stem cells: convergent and divergent mechanisms of myogenesis. PloS one *9*,

e90398. 10.1371/journal.pone.0090398.

10.   Spinazzola, J.M., and Gussoni, E. (2017). Isolation of Primary Human Skeletal Muscle

Cells. Bio Protoc *7*. 10.21769/BioProtoc.2591.

11.   Pisani, D.F., Dechesne, C.A., Sacconi, S., Delplace, S., Belmonte, N., Cochet, O.,

Clement, N., Wdziekonski, B., Villageois, A.P., Butori, C., et al. (2010). Isolation of a

highly myogenic CD34-negative subset of human skeletal muscle cells free of

adipogenic potential. Stem cells *28*, 753-764. 10.1002/stem.317.

12.   Xu, X., Wilschut, K.J., Kouklis, G., Tian, H., Hesse, R., Garland, C., Sbitany, H.,

Hansen, S., Seth, R., Knott, P.D., et al. (2015). Human Satellite Cell Transplantation

and Regeneration from Diverse Skeletal Muscles. Stem Cell Rep. *5*, 419-434.

10.1016/j.stemcr.2015.07.016.

13.   Zheng, B., Cao, B., Crisan, M., Sun, B., Li, G., Logar, A., Yap, S., Pollett, J.B., Drowley,

L., Cassino, T., et al. (2007). Prospective identification of myogenic endothelial cells in

human skeletal muscle. Nature biotechnology *25*, 1025-1034. 10.1038/nbt1334.

14.   Castiglioni, A., Hettmer, S., Lynes, M.D., Rao, T.N., Tchessalova, D., Sinha, I., Lee,

B.T., Tseng, Y.H., and Wagers, A.J. (2014). Isolation of progenitors that exhibit

myogenic/osteogenic bipotency in vitro by fluorescence-activated cell sorting from

human fetal muscle. Stem cell reports *2*, 92-106. 10.1016/j.stemcr.2013.12.006.

15.   Pisani, D.F., Clement, N., Loubat, A., Plaisant, M., Sacconi, S., Kurzenne, J.Y.,

Desnuelle, C., Dani, C., and Dechesne, C.A. (2010). Hierarchization of myogenic and

adipogenic progenitors within human skeletal muscle. Stem cells *28*, 2182-2194.

10.1002/stem.537.

16.   Gaida, J.E., Alfredson, H., Kiss, Z.S., Bass, S.L., and Cook, J.L. (2010). Asymptomatic

Achilles tendon pathology is associated with a central fat distribution in men and a

peripheral fat distribution in women: a cross sectional study of 298 individuals. BMC

Musculoskelet Disord *11*, 41. 10.1186/1471-2474-11-41.

17.   Voleti, P.B., Buckley, M.R., and Soslowsky, L.J. (2012). Tendon healing: repair and

regeneration. Annu Rev Biomed Eng *14*, 47-71.

10.1146/annurev-bioeng-071811-150122.

18.   Korcari, A., Nichols, A.E.C., Buckley, M.R., and Loiselle, A.E. (2023). Scleraxis-lineage

cells are required for tendon homeostasis and their depletion induces an accelerated

extracellular matrix aging phenotype. Elife *12*. 10.7554/eLife.84194.

19.   Pennisi, E. (2002). Tending tender tendons. Science *295*, 1011.

10.1126/science.295.5557.1011.

20.   Bi, Y., Ehirchiou, D., Kilts, T.M., Inkson, C.A., Embree, M.C., Sonoyama, W., Li, L.,

Leet, A.I., Seo, B.M., Zhang, L., et al. (2007). Identification of tendon stem/progenitor

cells and the role of the extracellular matrix in their niche. Nat Med *13*, 1219-1227.

10.1038/nm1630.

21.   Kohler, J., Popov, C., Klotz, B., Alberton, P., Prall, W.C., Haasters, F., Muller-Deubert,

S., Ebert, R., Klein-Hitpass, L., Jakob, F., et al. (2013). Uncovering the cellular and
molecular changes in tendon stem/progenitor cells attributed to tendon aging and
degeneration. Aging Cell *12*, 988-999. 10.1111/acel.12124.

22.   Nourissat, G., Berenbaum, F., and Duprez, D. (2015). Tendon injury: from biology to
tendon repair. Nature reviews. Rheumatology *11*, 223-233. 10.1038/nrrheum.2015.26.

23.   Pryce, B.A., Watson, S.S., Murchison, N.D., Staverosky, J.A., Dunker, N., and
Schweitzer, R. (2009). Recruitment and maintenance of tendon progenitors by
TGFbeta signaling are essential for tendon formation. Development *136*, 1351-1361.
10.1242/dev.027342.

24.   Yang, G., Rothrauff, B.B., Lin, H., Yu, S., and Tuan, R.S. (2017). Tendon-Derived
Extracellular Matrix Enhances Transforming Growth Factor-beta3-Induced Tenogenic
Differentiation of Human Adipose-Derived Stem Cells. Tissue engineering. Part A *23*,
166-176. 10.1089/ten.TEA.2015.0498.

25.   Barsby, T., and Guest, D. (2013). Transforming growth factor beta3 promotes tendon
differentiation of equine embryo-derived stem cells. Tissue Eng Part A *19*, 2156-2165.
10.1089/ten.TEA.2012.0372.

26.   Berthet, E., Chen, C., Butcher, K., Schneider, R.A., Alliston, T., and Amirtharajah, M.
(2013). Smad3 binds Scleraxis and Mohawk and regulates tendon matrix organization.
Journal of orthopaedic research : official publication of the Orthopaedic Research
Society *31*, 1475-1483. 10.1002/jor.22382.

27.   Wang, D., Pun, C.C.M., Huang, S., Tang, T.C.M., Ho, K.K.W., Rothrauff, B.B., Yung,
P.S.H., Blocki, A.M., Ker, E.D.F., and Tuan, R.S. (2020). Tendon-derived extracellular
matrix induces mesenchymal stem cell tenogenesis via an integrin/transforming
growth factor-beta crosstalk-mediated mechanism. FASEB J *34*, 8172-8186.
10.1096/fj.201902377RR.

28.   Li, M., Jia, J., Li, S., Cui, B., Huang, J., Guo, Z., Ma, K., Wang, L., and Cui, C. (2021).
Exosomes derived from tendon stem cells promote cell proliferation and migration
through the TGF beta signal pathway. Biochem Biophys Res Commun *536*, 88-94.
10.1016/j.bbrc.2020.12.057.

29.   You, T., Yuan, S., Bai, L., Zhang, X., Chen, P., and Zhang, W. (2020). Benzyl alcohol

accelerates recovery from Achilles tendon injury, potentially via TGF-beta1/Smad2/3

pathway. Injury *51*, 1515-1521. 10.1016/j.injury.2020.03.058.

30.   Jones, E.R., Jones, G.C., Legerlotz, K., and Riley, G.P. (2013). Cyclical strain

modulates metalloprotease and matrix gene expression in human tenocytes via

activation of TGFbeta. Biochim Biophys Acta *1833*, 2596-2607.

10.1016/j.bbamcr.2013.06.019.

31.   Rubenstein, A.B., Smith, G.R., Raue, U., Begue, G., Minchev, K., Ruf-Zamojski, F.,

Nair, V.D., Wang, X., Zhou, L., Zaslavsky, E., et al. (2020). Single-cell transcriptional

profiles in human skeletal muscle. Sci. Rep. *10*, 229. 10.1038/s41598-019-57110-6.

32.   Shao, X., Shi, L.L., Bluman, E.M., Wang, S., Xu, X., Chen, X., and Wang, J. (2020).

Satisfactory functional and MRI outcomes at the foot and ankle following harvesting of

full thickness peroneus longus tendon graft. Bone Joint J. *102-B*, 205-211.

10.1302/0301-620X.102B2.BJJ-2019-0949.R1.

33.   Sassoon, A.A., Ozasa, Y., Chikenji, T., Sun, Y.L., Larson, D.R., Maas, M.L., Zhao, C.,

Jen, J., and Amadio, P.C. (2012). Skeletal muscle and bone marrow derived stromal

cells: a comparison of tenocyte differentiation capabilities. Journal of orthopaedic

research : official publication of the Orthopaedic Research Society *30*, 1710-1718.

10.1002/jor.22135.

34.   Chen, B., Wang, B., Zhang, W.J., Zhou, G., Cao, Y., and Liu, W. (2012). In vivo tendon

engineering with skeletal muscle derived cells in a mouse model. Biomaterials *33*,

6086-6097. 10.1016/j.biomaterials.2012.05.022.

35.   Han, S.H., Kim, H.K., Ahn, J.H., Lee, D.H., Baek, M., Ye, G., Lee, J.M., Min, K., Oh, C.,

and Lee, S. (2018). A Protocol to Acquire the Degenerative Tenocyte from Humans. J

Vis Exp. 10.3791/57634.

36.   Fu, X., Xiao, J., Wei, Y., Li, S., Liu, Y., Yin, J., Sun, K., Sun, H., Wang, H., Zhang, Z., et

al. (2015). Combination of inflammation-related cytokines promotes long-term muscle

stem cell expansion. Cell Res *25*, 655-673. 10.1038/cr.2015.58.

37.   Rossi, L.A., Atala, N., Bertona, A., Tanoira, I., Bongiovanni, S., Maignon, G., and

Ranalletta, M. (2018). Return to Sports After in Situ Arthroscopic Repair of Partial

Rotator Cuff Tears. Arthroscopy : the journal of arthroscopic & related surgery : official

publication of the Arthroscopy Association of North America and the International

Arthroscopy Association. 10.1016/j.arthro.2018.07.037.

38.   Bianco, S.T., Moser, H.L., Galatz, L.M., and Huang, A.H. (2018). Biologics and stem

cell-based therapies for rotator cuff repair. Annals of the New York Academy of

Sciences. 10.1111/nyas.13918.

39.   Lui, P.P. (2015). Stem cell technology for tendon regeneration: current status,

challenges, and future research directions. Stem cells and cloning : advances and

applications *8*, 163-174. 10.2147/SCCAA.S60832.

40.   Liu, D., Black, B.L., and Derynck, R. (2001). TGF-beta inhibits muscle differentiation

through functional repression of myogenic transcription factors by Smad3. Genes Dev

15, 2950-2966. 10.1101/gad.925901.

41.   Brennan, T.J., Edmondson, D.G., Li, L., and Olson, E.N. (1991). Transforming growth

factor beta represses the actions of myogenin through a mechanism independent of

DNA binding. Proc. Natl. Acad. Sci. USA *88*, 3822-3826. 10.1073/pnas.88.9.3822.

42.   Melendez, J., Sieiro, D., Salgado, D., Morin, V., Dejardin, M.J., Zhou, C., Mullen, A.C.,

and Marcelle, C. (2021). TGFbeta signalling acts as a molecular brake of myoblast

fusion. Nat. Commun. *12*, 749. 10.1038/s41467-020-20290-1.

43.   Girardi, F., Taleb, A., Ebrahimi, M., Datye, A., Gamage, D.G., Peccate, C., Giordani, L.,

Millay, D.P., Gilbert, P.M., Cadot, B., and Le Grand, F. (2021). TGFbeta signaling curbs

cell fusion and muscle regeneration. Nat. Commun. *12*, 750.

10.1038/s41467-020-20289-8.

44.   Luo, Y., Fu, C., Wang, Z., Zhang, Z., Wang, H., and Liu, Y. (2015). Mangiferin

attenuates contusive spinal cord injury in rats through the regulation of oxidative stress,

inflammation and the Bcl2 and Bax pathway. Mol Med Rep *12*, 7132-7138.

10.3892/mmr.2015.4274.

45.   Yin, J., Yang, L., Xie, Y., Liu, Y., Li, S., Yang, W., Xu, B., Ji, H., Ding, L., Wang, K., et al.

(2018). Dkk3 dependent transcriptional regulation controls age related skeletal muscle

atrophy. Nat Commun *9*, 1752. 10.1038/s41467-018-04038-6.

46.   Otabe, K., Nakahara, H., Hasegawa, A., Matsukawa, T., Ayabe, F., Onizuka, N., Inui,

M., Takada, S., Ito, Y., Sekiya, I., et al. (2015). Transcription factor Mohawk controls

tenogenic differentiation of bone marrow mesenchymal stem cells in vitro and in vivo.

J Orthop Res *33*, 1-8. 10.1002/jor.22750.

47. Lou, J., Tu, Y., Burns, M., Silva, M.J., and Manske, P. (2001). BMP-12 gene transfer

augmentation of lacerated tendon repair. J Orthop Res *19*, 1199-1202.

10.1016/S0736-0266(01)00042-0.

48. Wang, D., Jiang, X., Lu, A., Tu, M., Huang, W., and Huang, P. (2018). BMP14 induces

tenogenic differentiation of bone marrow mesenchymal stem cells in vitro.

Experimental and therapeutic medicine *16*, 1165-1174. 10.3892/etm.2018.6293.

49. Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).

Reversed graph embedding resolves complex single-cell trajectories. Nat Methods *14*,

979-982. 10.1038/nmeth.4402.

50. Cox, M.P., Peterson, D.A., and Biggs, P.J. (2010). SolexaQA: At-a-glance quality

assessment of Illumina second-generation sequencing data. BMC Bioinformatics *11*,

485. 10.1186/1471-2105-11-485.

51. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with

high-throughput sequencing data. Bioinformatics *31*, 166-169.

10.1093/bioinformatics/btu638.

52. Zhou, Y., Zhu, J., Tong, T., Wang, J., Lin, B., and Zhang, J. (2019). A statistical

normalization method and differential expression analysis for RNA-seq data between

different species. BMC Bioinformatics *20*, 163. 10.1186/s12859-019-2745-1.

Figure legends

792   Shao et al, Figure 1

Figure 1. The CD29+/CD56+ cells isolated from human muscle biopsies display

robust features of myogenic progenitors.

a. UMAP plot of all mononuclear cells isolated from human skeletal muscle which were
colored by cell clusters. Three samples with a total of 57,193 cells were included for
analysis.

b. Dot blot of representative genes in each cell cluster.

c. Bar plot of cell proportion for each cell cluster.

d. Plot of joint expression analysis for CD29+/CD56+ cells in total mononuclear cells
isolated from human skeletal muscle.

e. Proportion of CD29+/CD56+ myogenic progenitors in specific cell population. Error
bars indicated standard deviation (n=3).

f. Immunofluorescence staining of PAX7, MYOD1 and MYF5 in primary human
CD29+/CD56+ myogenic progenitors. Scale bars, 100μm.

g. Statistical analysis of the percentage of PAX7+, MYOD1+ and MYF5+ cells in the
isolated CD29+/CD56+ myogenic progenitors. Error bars indicated standard deviation
(n=5).

h. Analysis of cell proliferation for human CD29+/CD56+ myogenic progenitors in vitro.
10,000 isolated human CD29+/CD56+ myogenic progenitors were plated for
proliferation and counted at each time point. Error bars indicated standard deviation
(n=3).

i. Immunofluorescence staining of MyHC in myotubes differentiated from human
CD29+/CD56+ myogenic progenitors. Primary human CD29+/CD56+ myogenic
progenitors were isolated and differentiated to myotubes for 5 days followed by MyHC
immunofluorescence staining. Scale bars, 100 μm.

j. Statistical analysis of the percentage of nuclei in MyHC+ myotubes after differentiation
of the human CD29+/CD56+ myogenic progenitors. Error bars indicated standard
deviation (n=5).

k. Relative expression level of PAX7, MYF5, and MYOD1 in human CD29+/CD56+
myogenic progenitors (Undifferentiated) and differentiated myotubes (Differentiated).
RT-qPCR assays were performed for human CD29+/CD56+ myogenic progenitors
before and after myogenic differentiation. GAPDH was served as reference gene.
Error bars indicated standard deviation (n=3). *** indicated p<0.001.

I.   Relative expression level of MYOG, MYH1, and MYH3 in human CD29+/CD56+

myogenic progenitors (Undifferentiated) and differentiated myotubes (Differentiated).

RT-qPCR assays were performed for human CD29+/CD56+ myogenic progenitors

before and after myogenic differentiation. GAPDH was served as reference gene.

Error bars indicated standard deviation (n=3). *** indicated $p<0.001$.

Shao et al, Figure 2

Figure 2. Human CD29+/CD56+ myogenic progenitors display tendon differentiation

ability in vitro.

a. Immunofluorescence staining of tendon marker TNC and SCX in human
CD29+/CD56+ myogenic progenitors induced for myogenic and tenogenic
differentiation, respectively. Scale bars, 100 μm.

b. Quantification of TNC and SCX fluorescent intensity in human CD29+/CD56+
myogenic progenitors undergone myogenic and tenogenic differentiation, respectively.
Error bars indicated standard deviation (n=5).

c. Relative expression levels of genes enriched in tendon cells. RT-qPCR assays were
performed with human CD29+/CD56+ myogenic progenitors upon myogenic and
tenogenic differentiation, respectively. GAPDH was served as reference gene. Error
bars indicated standard deviation (n=3). *** indicated $p<0.001$.

d. Scheme of clonal proliferation and differentiation assay.

e. Representative immunofluorescence staining images of MyHC as the marker for
successful myogenic differentiation. Scale bars, 100 μm.

f. Representative immunofluorescence staining images of SCX as the marker for
successful tenogenic differentiation. Scale bars, 100 μm.

g. Statistical analysis of the myogenic and tenogenic differentiation efficiency of human
CD29+/CD56+ myogenic progenitors. Error bars indicated standard deviation (n=3).

Figure 3. Human CD29+/CD56+ myogenic progenitors have tendon differentiation potential.

a. Heat map of gene expression profiles of human CD29+/CD56+ myogenic progenitors, human CD29+/CD56+ myogenic progenitors after myogenic differentiation, human

CD29+/CD56+ myogenic progenitors after tenogenic differentiation, and primary

tenocytes.

b. Volcano plot of genes enriched in myogenic differentiation of human CD29+/CD56+

myogenic progenitors.

c. Volcano plot of genes enriched in tenogenic differentiation of human CD29+/CD56+

myogenic progenitors.

d. Bubble chart of GO analysis of cellular process up-regulated in myogenic

differentiation of human CD29+/CD56+ myogenic progenitors.

e. Bubble chart of GO analysis of cellular process up-regulated in tenogenic

differentiation of human CD29+/CD56+ myogenic progenitors.

Figure 4. Murine MuSCs display poor tenogenic differentiation ability.

a. Immunofluorescence staining of murine MuSCs directed towards myogenic and

tenogenic differentiation, respectively. The undifferentiated murine MuSCs were

stained with Pax7 and DAPI. Murine MuSCs after myogenic differentiation were

stained with MyHC and DAPI. Murine MuSCs after tenogenic differentiation were

stained with Scx and DAPI. Scale bars, 100 μm.

b. Statistical analysis of the percentage of nuclei in MyHC+ myotubes. Error bars

indicated standard deviation (n=5).

c. Relative expression levels of myogenic differentiation specific genes MyoG, Myh1,

and Myh3. Mouse MuSCs were induced to differentiate towards muscle and tendon,

respectively. Mouse MuSCs before and after differentiation together with primary

tenocytes were harvested and subjected for RT-qPCR analysis. Gapdh was served as

reference gene. Error bars indicated standard deviation (n=3). * indicated $p<0.05$, **

indicated $p<0.01$, n.s. indicated $n>0.05$.

d. Relative expression levels of tenogenic differentiation marker genes Scx, Tnc, Col I,

Mkx and Thbs4. Mouse MuSCs were induced to differentiate towards muscle and

tendon, respectively. Mouse MuSCs before and after differentiation together with

primary tenocytes were harvested and subjected for RT-qPCR analysis. Gapdh was

served as reference gene. Error bars indicated standard deviation (n=3). n.s.

indicated $n>0.05$.

e. Scheme of Pax7+ MuSC progeny lineage tracing assay.

f. Immunofluorescence staining of tendon tissue 4 months after injury. tdTomato+ cells

indicated the progeny of Pax7+ MuSCs. TNC+ cells indicated tendon cells. MyHC+

cells indicated muscle fibers. DAPI indicated nuclei staining. Merged indicated the

merged images of tdTomato, TNC, MyHC, and DAPI. The upper panel indicated the

low magnification images. Scale bars, 100 μm. The lower panel indicated the high

magnification images of the region label by white square in the upper panel. Scale

bars, 10 μm.

g. The statistical analysis of the percentage of tdTomato+ TNC+ cell 4 months after

tendon injury. Error bars indicated standard deviation (n=5).

Figure 5. Transplantation of human CD29+/CD56+ myogenic progenitors facilitates

tendon regeneration.

a. Scheme of transplantation of human CD29+/CD56+ myogenic progenitors for

NOD/SCID with tendon injury.

b. Immunofluorescence staining of the regenerated tendon-like tissue after

transplantation of human CD29+/CD56+ myogenic progenitors. Tendon injury was

induced in recipient NOD/SCID mice and 50,000 human CD29+/CD56+ myogenic

progenitors were transplanted to the injured tendon at the injured site. The

regenerated tendon-like tissue, the connected muscle, and the surrounding soft

tissues were harvested to make continuous cryosections. One of the continuous

cryosections were subjected for immunofluorescence staining of tendon marker

human specific TNC and human specific Lamin A/C. DAPI indicated nuclei staining.

Merge indicated the merged images of human TNC, human Lamin A/C, and DAPI.

The white lines indicated the location of regenerated tendon-like tissues from human

CD29+/CD56+ myogenic progenitors based on human TNC staining. The yellow

dashed lines indicated the superimposed location of muscle based on MyHC staining

in panel c. Scale bars, 100 μm.

c. Immunofluorescence staining of MyHC and human Lamin A/C. The regenerated

tendon-like tissue, the connected muscle, and the surrounding soft tissues were

harvested and subjected for continuous cryosection. One of the continuous

cryosections were stained for MyHC which was specifically expressed in skeletal

muscle, and human Lamin A/C to label human cells. DAPI indicated nuclei staining.

Merge indicated the merged images of MyHC, human Lamin A/C, and DAPI. The

white lines indicated the location of regenerated tendon-like tissue from human

CD29+/CD56+ myogenic progenitors based on human TNC staining in panel b. The

yellow dashed lines indicated the location of muscle based on MyHC staining. Scale

bars, 100μm.

d. Statistical analysis of the percentage of human cells expressing skeletal muscle

marker MyHC or tendon marker TNC after being transplanted to the injured tendon.

Error bars indicated standard deviation (n=5).

e. Immunofluorescence staining of Col I, Col III and human Lamin A/C. Two months after

human cell transplantation, continuous cryosections containing the regenerated

tendon-like tissue and native tendon tissue was stained with Col I, Col III and human

Lamin A/C. DAPI indicated the staining of nuclei. T, native tendon tissue; W, wound

tendon tissue. Scale bars, 50 µm.

f. Immunofluorescence staining of tendon tissue after transplantation of tdTomato+

murine MuSCs. tdTomato indicated the progenies of murine MuSCs. TNC indicated

immunofluorescence staining of tendon marker TNC. MyHC indicated

immunofluorescence staining of myofiber marker MyHC. DAPI indicated nuclei

staining. Merge indicated merged images of tdTomato, TNC, MyHC, and DAPI. The

upper panel indicated low magnification images. Scale bars, 100 µm. The lower panel

indicated the amplified images of the region indicated by the white square. Scale bars,

10 µm.

g. Statistical analysis of TNC+ tdTomato+ cells in tendon tissue after transplantation of

murine MuSCs. Error bars indicated standard deviation (n=5).

Figure 6. Transplantation of human CD29+/CD56+ myogenic progenitors improves the locomotor function after tendon injury.

a. TEM images of collagen fibrils in the injured tendon with PBS injection, injured tendon with human CD29+/CD56+ myogenic progenitors transplantation, and uninjured

tendon. Scale bars, 500nm.

b. Statistical analysis of the collagen fibril diameter and percentage of collagen fibril area

for the injured tendon with PBS injection, injured tendon with human CD29+/CD56+

myogenic progenitors transplantation, and uninjured tendon. Error bars indicated

standard deviation (n=5). * indicated $p<0.05$, *** indicated $p<0.001$.

c. Statistical analysis of the max load and stiffness of the injured tendon with PBS

injection, injured tendon with human CD29+/CD56+ myogenic progenitors

transplantation, and uninjured tendon. Error bars indicated standard deviation (n=5). *

indicated $p<0.05$, ** indicated $p<0.01$, *** indicated $p<0.001$.

d. Twitch and tetanus plantarflexion force of the involved limb with PBS injection after

tendon injury, human CD29+/CD56+ myogenic progenitors transplantation after

tendon injury, and uninjured tendon. Error bars indicated standard deviation (n=5). *

indicated $p<0.05$, ** indicated $p<0.01$, *** indicated $p<0.001$.

e. The results of treadmill exercise for tendon injured mice with or without human

CD29+/CD56+ myogenic progenitors transplantation. The endurance time and max

fatigue speed were compared. Error bars indicated standard deviation (n=4). *

indicated $p<0.05$, ** indicated $p<0.01$, ***indicated $p<0.001$ and n.s. indicated $p>0.05$.

Shao et al, Figure 7

Figure 7. TGFβ signaling pathway contributes to tenogenic differentiation of human CD29+/CD56+ myogenic progenitors.

a.   Bubble chart of KEGG enrichment analysis of upregulated genes in human
CD29+/CD56+ myogenic progenitors when compared with mouse muscle stem cells.

b.   Heatmap of detailed upregulated genes in human CD29+/CD56+ myogenic
progenitors which were enriched in TGFβ signaling pathway.

c.   Immunofluorescence staining of tendon marker TNC and SCX in human
CD29+/CD56+ myogenic progenitors induced for tenogenic differentiation with or
without TGFβ signaling pathway inhibitor SB-431542 for 12 days, respectively. Scale
bars, 100 μm.

d.   Protein levels of TNC and SCX. Human CD29+/CD56+ myogenic progenitors were
induced towards tenogenic differentiation with or without TGFβ signaling pathway
inhibitor SB-431542 for 12 days, respectively. Total protein was extracted from cells
before and after differentiation and subjected for TNC and SCX immunoblotting.
GAPDH was served as loading control.

e.   Relative expression levels of tendon related genes. RT-qPCR assays were performed
with human CD29+/CD56+ myogenic progenitors upon tenogenic differentiation with
or without TGFβ signaling pathway inhibitor SB-431542 for 12 days, respectively.
GAPDH was served as reference gene. Error bars indicated standard deviation (n=3).
*** indicated $p<0.001$.

f.   Immunofluorescence staining of myogenic differentiation marker MyHC in human
CD29+/CD56+ myogenic progenitors induced for tenogenic differentiation with or
without TGFβ signaling pathway inhibitor SB-431542 for 12 days, respectively. Scale
bars, 50 μm.

g.   Relative expression levels of muscle related genes. RT-qPCR assays were performed
with human CD29+/CD56+ myogenic progenitors upon tenogenic differentiation with
or without TGFβ signaling pathway inhibitor SB-431542 for 12 days, respectively.
GAPDH was served as reference gene. Error bars indicated standard deviation (n=3).
*** indicated $p<0.001$.

Shao et al, Supplemental Figure 1
Related to Figure2

1000

Supplemental Figure 1. Human CD29+/CD56+ myogenic progenitors display

tenogenic differentiation potential.

a.   Phase contrast images of human CD29+/CD56+ myogenic progenitors before and
after differentiation. Scale bars, 100 μm.

b.   Immunofluorescence staining of myogenic differentiation marker genes MYOG and
MyHC. Human CD29+/CD56+ myogenic progenitors were induced for tenogenic
differentiation. The differentiated cells were stained with MYOG and MyHC antibody.
Red indicated MYOG or MyHC; DAPI indicated nuclei staining; merge indicated the
merged images of red and blue staining. Scale bars, 100μm.

c.   Statistical analysis of the percentage of human cells expressing MYOG or MyHC
under tenogenic differentiation condition. Error bars indicated standard deviation
(n=5).

d.   Expression level of genes marking myogenic differentiation. Human CD29+/CD56+
myogenic progenitors were induced for myogenic and tenogenic differentiation,
respectively. The undifferentiated and differentiated cells were subjected for RT-qPCR
analysis of MYH1, MYH3, DESMIN, and MYL1. GAPDH was served as reference
gene. Error bars indicated standard deviation (n=5). *** indicated $p<0.001$.

e.   Protein levels of TNC and MyHC. Human CD29+/CD56+ myogenic progenitors were
induced towards myogenic and tenogenic differentiation, respectively. Total protein
was extracted from cells before and after differentiation and subjected for TNC and
MyHC immunoblotting. HSP70 and GAPDH were served as loading controls.

1022

Supplemental Figure 2. Murine MuSCs display poor tenogenic differentiation ability.

a. Schematic illustration of the mouse line for murine MuSCs lineage tracing.

b. The surgery in mice mimicking human peroneus longus tendon removal surgery. The

medial gastrocnemius tendon was completely removed. Scale bars, 1mm.

c. Immunofluorescence images of injured muscle adjacent to the removed tendon. The

MuSCs were labeled with tdTomato before injury. The tdTomato positive myofibers

after muscle injury were analyzed to evaluate the tracing system. Scale bars, 100µm.

d. Statistical analysis of the tdTomato positive myofibers adjacent to the removed tendon.

Error bars indicated standard deviation (n=5).

1032

Supplemental Figure 3. Transplantation of human CD29+/CD56+ myogenic

progenitors facilitates tendon regeneration and transplantation of human/murine

myogenic progenitors improve muscle regeneration.

a. Immunofluorescence staining of SCX and human specific Lamin A/C in injured tendon.

Two months after transplantation of human CD29+/CD56+ myogenic progenitors, the

injured tendon was stained with SCX and human specific Lamin A/C. DAPI indicated

nuclei staining. Merge indicated the merged images of SCX, human specific Lamin

A/C, and DAPI. Scale bars, 20 μm.

b. Immunofluorescence staining of TNMD and human specific Lamin A/C in injured

tendon. Two months after transplantation of human CD29+/CD56+ myogenic

progenitors, the injured tendon was stained with TNMD and human specific Lamin A/C.

DAPI indicated nuclei staining. Merge indicated the merged images of TNMD, human

Lamin A/C, and DAPI. Scale bars, 20 μm.

c. Statistical analysis of the percentage of human cells expressing tendon related

markers SCX or TNMD after being transplanted to the injured tendon. Error bars

indicated standard deviation (n=5).

d. Scheme of transplantation of human CD29+/CD56+ myogenic progenitors after

muscle injury. TA muscles were first irradiated to kill the local MuSCs in NOD/SCID

mice. Then transplantation of human CD29+/CD56+ myogenic progenitors to the

irradiated pre-injured recipient mice was performed and TA muscles were harvested

after 28 days.

e. Immunofluorescence staining of MyHC and human specific Lamin A/C after

transplantation of human CD29+/CD56+ myogenic progenitors. Scale bars, 100μm.

f. Statistical analysis of the engraftment efficiency of human CD29+/CD56+ myogenic

progenitors transplantation in muscle. Error bars indicated standard deviation (n=5).

g. Scheme of murine MuSCs transplantation after muscle injury. TA muscles were first

irradiated to kill the local MuSCs in NOD/SCID mice. Then transplantation of tdTomato

positive MuSCs to the irradiated pre-injured recipient mice was performed and TA

muscles were harvested after 28 days.

h. Immunofluorescence staining of Laminin and murine MuSCs constitutively expressing

tdTomato after murine tdTomato positive MuSCs transplantation. Scale bars, 100μm.

i. Statistical analysis of the engraftment efficiency of murine MuSCs transplantation in

muscle. Error bars indicated standard deviation (n=5).

Data availability

The datasets used in this study can be obtained from the corresponding author upon

reasonable request.

1069

1070 **Conflict of interest disclosure**

1071 The authors declared no conflict of interest.

1072

1073 **Funding:**

1074 This work was supported by the Strategic Priority Research Program of the Chinese

1075 Academy of Science (XDA16020400), National Science Foundation of China (32170804,

1076 81871096, 82372384, 82302657).

1077

1078 **Contributors**

1079 XS, XL, HL, HZ, and XF designed the study, collected the experimental data and wrote the

1080 original draft. SL and SZ conducted the image analysis, statistical analysis and edited the

1081 manuscript. WY and MW designed the study, reviewed and edited the manuscript. JW

1082 and PH administrated the project and edited the manuscript.

Shao et al, Figure 1

Shao et al, Figure 2

Shao et al, Figure 3

Shao et al, Figure 4

a

NOD/SCID mouse → Tendon Injury → CD29+/CD56+ myogenic progenitors transplantation → 2 months → Harvest

b

| Human TNC | Hu-Lamin A/C | DAPI | Merge |

PBS

Human Cells

Human Cells

c

| MyHC | Hu-Lamin A/C | DAPI | Merge |

PBS

Human Cells

Human Cells

d

% of human cells

TNC   MyHC

e

| Hu-Lamin A/C | Col I | Col III |

PBS

Human Cells

T W

f

| tdTomato | TNC | MyHC | DAPI | Merge |

g

TNC+/Td tomato+/MyHC− Cells (%)

Shao et al, Figure 5

Shao et al, Figure 7