Human myogenic progenitor cells display tenogenic differentiation potential and facilitate

tendon regeneration

Xiexiang Shao[1], Xingzuan Lin[1], Hao Zhou[1], Lili Han[4], Xin Fu[1], Sheng Li[1], SiyuanZhu[1], Shenao

Zhou[4], Jianhua Wang[1*], Ping Hu[1, 2, 3, 4*]

Affiliations:

1. Xinhua Hospital affiliated to Shanghai Jiao Tong University School of Medicine.

Shanghai,20023, China.

2. Guangzhou Laboratory, Guangzhou, 510005, China.

3. Institute for Stem Cell and Regeneration, Chinese Academy of Sciences, Beijing100101, China.

4. State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology,

Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai 200031,

China.

*Corresponding author: Jianhua Wang, Email: wangjianhua@xinhuamed.com.cn; PingHu, Email:

hup@sibcb.ac.cn;

Abstract

Tendon injury occurs at high frequency and is difficult to repair. Identification of

human stem cells being able to regenerate tendon will greatly facilitate the development of

regenerative medicine for tendon injury. We identified CD29+/CD56+ human muscle

stem/progenitor cells having tendon differentiation potential both in vitro and in vivo.

Transplantation of human myogenic progenitor cells contributes to injured tendon repair

and thus improves locomotor function. Interestingly, the tendon differentiation potential in

mouse muscle stem cells is minimal and the higher TGFβ signaling level in human

myogenic progenitor cells may be the key for the distinct feature of human myogenic

progenitor cells. These findings reveal that CD29+/CD56+ human muscle stem/progenitor

cells are bi-potential adult stem cells and can serve as a new source for tendon

regeneration.

Introduction

Skeletal muscle is a tissue with great regeneration ability due to the existence of muscle stem cells (MuSCs). MuSCs are adult stem cells located at the periphery of myofibers between the basal lamina and the plasmalemma of the myofibers and responsible for muscle regeneration[1,2]. MuSCs have been considered to be unipotent stem cells to have a sole differentiation potential to myofibers[3]. MuSCs undergo expansion and differentiate to multinuclei myofibers after injury in vivo. MuSCs have remarkable abilities to support muscle regeneration. After transplantation, the engrafted mouse MuSCs go through active expansion and regenerate myofibers[4]. Isolation by FACS and expansion of MuSCs have been reported from several species including mouse, pig, and human[5,6]. Single cell sequencing analysis from human skeletal muscles have also revealed the existence of Pax7+ MuSCs[6,7]. Due to the different motion patterns, the regeneration capacity may be different between human and rodents. Current investigations have suggested that human myogenic progenitor cells didn't share same markers as that in mice[8], and the expression pattern of oxidative enzymes and cytokines between these two species are also different[9], suggesting that human muscle progenitor cells may have distinct features from mice.

FACS method to isolate cells with myogenic differentiation potential from human muscle biopsies has been established. CD56 (NCAM)+ and CD56+ CD29+ cells from skeletal muscle display myogenic differentiation potential[9,10,11,12]. The CD56+ CD34+ progenitor cells isolated from human skeletal muscle biopsies have been reported to have chondrogenic, adipogenic, and osteogenic potentials besides myogenic potential[13-15]. CD56+CD34- progenitor cells are free of adipogenic potential[11]. The differentiation potential of human muscle progenitor cells remains to be further explored.

Skeletal muscle directly connects to tendons which is responsible for transmitting forces from skeletal muscle to bone to generate active movement. Tendinopathy affects more than 10% of the population under 45 and compromises the tendon functions[16,17]. Tendon injury healing is slow and incomplete due to the low number of cells in tendon and the hypovascular and anaerobic environment[18,19]. Tendon stem/progenitor cells (TDSCs) are a cell population derived from tendon and considered to be a subgroup of

mesenchymal stem cells that have abilities to improve tendon injury healing[20,21]. However,

the number of TDSCs in tendon is low and retrieving TDSCs is invasive and exacerbates

tendon injury. Morphology and proliferation ability loss in the culture system also hampers

the efforts to obtain sufficient amounts of active TDSCs. To find more cell types supporting

tendon regeneration will facilitate the development of regenerative medicine to treat

tendon injury.

The activity of tenogenesis is tightly regulated by many signaling pathways, while

TGFβ/SMAD2/3 is a crucial molecular axis[22]. TGFβ signaling is indispensable for tendon

development[23], and it could systematically promote the tenogenic differentiation of stem

cells[24,25]. The downstream effectors SMAD2 and SMAD3 are able to activate the

transcription of tendon-specific genes which further facilitates tendon development and

tenogenic differentiation[22,26,27]. After tendon injury, TGFβ/SMAD2/3 promotes the

proliferation, migration and differentiation of TDSCs[28], increases tendon collagen

synthesis[29], and contributes to matrix anabolism for tendon remodeling[30].

Here we found that CD56+ CD29+ human myogenic progenitor cells can be

differentiated to tendon cells both in vitro and in vivo. Transplantation of human myogenic

progenitor cells to injured tendon in mice improved the tendon regeneration, suggesting

that human muscle myogenic progenitor cells are bipotential adult stem cells. Interestingly,

the tendon differentiation potential in mouse muscle stem cells is minimal and the higher

TGFβ signaling level in human myogenic progenitor cells may be the key for the distinct

feature of human myogenic progenitor cells.

Results

Potential tenogenic differentiation potential of myogenic progenitor cells identified

by single cell sequencing

To analyze the cell components of human skeletal muscle, single cell sequencing

analysis was performed using skeletal muscle biopsy. A total of 15,102 cells were included

for analysis. The cells were grouped to 8 cell populations (Fig. 1a and b). Consistent with

the previous single cell sequencing results from human muscles[31], Fibro/Adipogenic

progenitors (FAPs), endothelial cells, muscle cells, and myogenic progenitor cells were

among the identified cell types (Fig. 1a and b).

A population of tenocytes was identified in human muscle biopsies (Fig. 1a and b).

Surprisingly, when we performed unsupervised trajectory analysis using the single cell

sequencing data, two branches were identified by the analysis (Fig. 1c and d), while one

branch linked muscle progenitor cells and tenocytes (Fig. 1c and d). RNA FISH of skeletal

muscle section revealed that SCX, a key transcription factor determining tendon lineage,

could be expressed in muscle stem/progenitor cells (Fig. 1e and f). These analysis results

suggest that muscle stem cells could have tenocyte differentiation potential.

To confirm the single cell analysis results, we first isolated myogenic progenitor cells

from human muscle biopsy using FACS as described previously[6,12]. CD45- CD31-CD56+

CD29+ cells were collected from the single cell suspension dissociated from muscle

tissue. The collected cells were stained with anti-PAX7, anti-MYOD1 and anti-MYF5

antibody to confirm the purity of isolated myogenic progenitor cells (Fig. 1g and h). When

induced to differentiate by 0.4% Ultroser G, human myogenic progenitor cells differentiate

to myotubes robustly. Approximately 90% of nuclei were present in Myosin heavy chain

(MyHC) positive myotubes (Fig. 1i and j). Consistently, the expression of PAX7, MYF5 and

MYOD1 were enriched in the myogenic progenitor cells, their expression decreased after

differentiation as shown by RT-qPCR assays (Fig. 1k). In contrast, the expression of

genes marking differentiation such as Myogenin (MYOG), Myosin heavy chain 1 (MYH1),

and Myosin heavy chain 3 (MYH3) were up-regulated in differentiated myotubes (Fig. 1l).

Taken together, these results suggest that the myogenic progenitor cells isolated from

human muscle biopsies have robust myogenic differentiation potential.

Human myogenic progenitor cells display tenogenic differentiation potential in

vitro

The isolated primary human myogenic progenitor cells were induced to differentiate to

tenocytes by 100ng/ml GDF5, 100ng/ml GDF7 and 0.2mM ascorbic acid for 12 days. After

12 days of tendon differentiation, the morphology of cells display dramatic differences

from those undergoing myogenic differentiation (SFig. 1a). Furthermore, the expression of

tendon markers such as TNC and SCX was significantly increased (Fig. 2a, b, and c).

Moreover, some other tendon related genes, such as COL I, MKX, THBS4, and COMP,

were also enriched after tendon differentiation induction (Fig. 2c). In contrast, the

expression of these genes was not enriched upon myogenic induction (Fig. 2a, b, and c).

The expression of genes marking myogenic differentiation such as MYOG and MyHC

were only detected in a small portion of cells after tenogenic differentiation (SFig. 1b and

c). Compared to the over 90% of differentiation efficiency upon myogenic differentiation

induction, only about 20% of cells showed MYOG and MyHC expression after 12 days of

tenogenic induction (SFig. 1b and c). The genes marking differentiated myotubes such as

MYH1, MYH3, DESMIN, and MYL1 showed moderate elevation after tenogenic

differentiation, while dramatic upregulation of these genes was observed after myogenic

differentiation (SFig. 1d and e). These results combined suggest that human myogenic

progenitor cells are capable of tendon differentiation in vitro.

To further confirm the results, we next performed clonal analysis. The freshly isolated

primary human myogenic progenitor cells were seeded to 96 well plate with the

concentration of 1 cell/well. The wells of single cell were allowed to proliferate for 10 days

followed by tenogenic or myogenic induction. The plates with myogenic induction were

differentiated for 4 days and immunofluorescence staining of MyHC was performed to

determine muscle differentiation. The plates with tenogenic induction were differentiated

for 12 days and immunofluorescence staining of SCX was performed to determine

tenocyte differentiation (Fig. 2d). The number of wells showing positive MyHC staining

was counted and the myogenic differentiation efficiency was calculated (Fig. 2e and g).

MyHC+ myotubes was observed in over 95% of wells with alive cells (Fig. 2g). Similarly,

the number of wells displaying positive SCX staining was counted and the tenogenic

differentiation efficiency was calculated (Fig. 2f and g). Approximately 40% of myogenic

progenitor cells displayed tenogenic differentiation ability (Fig. 2g), suggesting that human

myogenic progenitor cells have tenogenic differentiation potential. Taken together, these

results suggest that human myogenic progenitor cells have dual differentiation potentials

towards muscle or tendon in vitro.

Tenocytes differentiated from human myogenic progenitor cells display similar

expression profile to primary tenocytes

RNA sequencing was then performed to further determine the lineage of the

tenocytes differentiated from myogenic progenitor cells. Myogenic progenitor cells were

induced towards myogenic or tenogenic differentiation respectively. We also isolated

primary human tenocytes as described previously[32]. These cells were subjected for RNA

sequencing analysis. The differentiated myotubes and tenocytes displayed distinct

expression patterns (Fig. 3a). The differentiated tendon cells share high similarity to fresh

isolated tenocytes from human tendons (Fig. 3a), suggesting that human muscle

stem/progenitor cells are capable of tendon differentiation. Consistently, two distinct sets

of genes were up-regulated after myogenic induction and tenogenic induction,

respectively (Fig. 3b and c). These results suggest that human myogenic progenitor cells

are capable of dual direction differentiation

Further GO analysis also displayed the activation of two distinct sets of cell features.

Upon the skeletal muscle differentiation induction, terms related to skeletal muscle

functions such as skeletal muscle thin filament assembly, skeletal muscle contraction,

muscle organ development, and sarcomere organization were enriched (Fig. 3d),

suggesting the muscular identity of the differentiated cells. In contrast, terms related to

tendon formation, tendon development, and tendon cell differentiation were enriched after

tenogenic differentiation (Fig. 3e), suggesting that tendon identity is achieved in the

differentiated cells. Together, these results suggest that human myogenic progenitor cells

are capable of both myogenic and tenogenic differentiation in vitro.

Murine MuSCs display poor tenogenic differentiation ability

We next investigated whether the tenogenic differentiation ability is present in rodent

muscle stem cells (MuSCs). Mouse MuSCs were isolated by positive marker of Vcam1 as

described previously[6] and induced for tenogenic differentiation. In sharp contrast to

human myogenic progenitor cells, murine MuSCs failed to be induced to tendon cells

upon the same induction condition as that for human myogenic progenitor cells, though

the myogenic differentiation is as efficient as the human myogenic progenitor cells. After

myogenic differentiation, 93.9% of nuclei were present in MyHC+ myotubes (Fig. 4a and

b). In sharp contrast, no Scx+ cells were observed after 12 days of induction for tenogenic

differentiation (Fig. 4a). Consistent with the immunofluorescence staining results,

RT-qPCR results revealed that myogenic differentiation marker genes such as MyoG,

Myh1, and Myh3 were up-regulated under both myogenic and tenogenic differentiation

conditions (Fig. 4c), suggesting that murine MuSCs predominantly commit myogenic

differentiation under induction. Different from human myogenic progenitor cells, the

expression of genes indicating tendon cell fate such as Scx, Tnc, Col I, Mkx, and Thbs4

did not increase after tenogenic differentiation (Fig. 4d), suggesting the failure to induce

tenogenic cell fate from murine MuSCs. To further rule out the bias caused by different

FACs strategies between human and mouse myogenic progenitor cells, mouse muscle

CD29+/CD56+ cells were isolated for tenogenic induction. However, very few mouse

muscle CD29+/CD56+ cells expressed myogenic progenitor cell marker Pax7, MyoD1

and Vcam1, indicating the failure strategy to purify mouse myogenic progenitor cells by

gating CD29+/CD56+ (SFig. 2a and 2b). Furthermore, RT-qPCR results of tendon related

markers Scx, Tnc, Mkx, and Thbs4 revealed that there was also no tenogenic or

myogenic differentiation potential of mouse muscle CD29+/CD56+ cells (SFig. 2c).

Together, these results suggest that murine MuSCs display almost no tenogenic

differentiation potential in vitro.

We next performed the lineage tracing experiments in mice to further examine the

tenogenic potential of mouse MuSCs in vivo (Fig. 4e). Pax7CreERT2 mice were crossed

to flox-Stop-flox-tdTomato mice. MuSCs and the descendants of MuSCs will be labeled by

tdTomato (SFig. 2d). Tendon injury in mice was generated by mimicking the peroneus

longus tendon removal surgery in human (SFig. 2e). In human, it has been reported that

the tendon could be regenerated to some extent after the peroneus longus tendon

removal surgery based on MRI imaging[33]. In this surgery, injury of the skeletal muscle

adjacent to the removed tendon was inevitable. The accompanied skeletal muscle injury

could activate MuSCs and make them available for tendon regeneration. To further

activate MuSCs to guarantee that sufficient amount of activated MuSCs were available

around the tendon injury site, we also injected cardiotoxin (CTX) at the muscle adjacent to

the sites where the tendon was removed to induce more muscle injuries and further

activate MuSCs. If MuSCs can participate tendon regeneration, tdTomato+ tendon cells

would be observed after the reparation of tendon injury. As a control, muscle injury was

induced by muscular injection of CTX at the muscles adjacent to the removed tendon.

Expectedly, large amount of tdTomato+ myofibers were observed after muscle injury

(SFig. 2f-g), suggesting that the tracing system works well. Nevertheless, less than 0.2%

tendon cells originated from mouse MuSCs were observed even four months after tendon

removal (Fig. 4f-g). These results suggest that murine MuSCs have poor tendon

differentiation abilities.

Transplantation of human myogenic progenitor cells facilitates tendon

regeneration

We next went on to investigate whether the transplantation of human myogenic

progenitor cells to mouse can improve tendon regeneration. An approximately 1.5mm

long and 0.5mm width transverse incision was performed at 5mm from the calcaneus in

Achilles tendon for NOD/SCID immunodeficient mice. Total 50,000 Pax7+ human

myogenic progenitor cells packed in hydrogel were planted at the incision site (Fig. 5a). As

a control, 50,000 murine MuSCs packed in hydrogel were transplanted in the SCID

recipient mice undergoing the same tendon injury at the cleavage sites. Packing the cells

with hydrogel concentrated the transplanted cells at the local injury sites. Two months

after transplantation, the tendons carrying transplanted human myogenic progenitor cells

or murine MuSCs were harvested, respectively (Fig. 5a).

In mice transplanted with human myogenic progenitor cells, continuous cryosections

containing muscle and tendon tissues were generated. Immunofluorescence staining was

performed to detect tendon and muscle markers with two cryosections adjacent to each

other, respectively. The two sets of images obtained on continuous cryosections were

superimposed on each other to pinpoint the position of tendons. Immunofluorescence

staining of antibody specifically recognizing human TNC indicated the position of

regenerated tendon-like tissue from human cells in the harvested tissues (Fig. 5b). The

presence of human cells was also illustrated by the immunofluorescence staining with the

antibody specifically recognizing human Lamin A/C. In the control PBS injection group,

where the mixture of PBS and hydrogel instead human myogenic progenitor cells was

injected to the incision sites, no human Lamin A/C was detected. These results confirmed

the specificity of human TNC and Lamin A/C antibody. Immunofluorescence staining

revealed that over 75% of the human cells showed TNC expression (Fig. 5b and d),

suggesting that the majority of the transplanted human myogenic progenitor cells

differentiate to tendon cells in vivo. The immunofluorescence staining of MyHC and

human Lamin A/C was performed to detect the muscle cells originated from the

transplanted human myogenic progenitor cells. Only about 12.8% of the human cells

detected expressed MyHC (Fig. 5c and d). Moreover, the human cells were predominantly

enriched at the tendon region (Fig. 5b). Only a few MyHC+ cells originated from the

transplanted human myogenic cells scattered in the muscle region (Fig. 5c). To further

confirm the tendon differentiation potential of the transplanted human myogenic

progenitor cells, immunofluorescence staining of SCX and TNMD was also performed.

The majority (80.0% and 74.6%) of transplanted human myogenic progenitor cells also

express SCX and TNMD (SFig. 3a, b and c). Furthermore, Col I was predominantly

expressed in the regenerated tendon-like tissue rather than Col III after human cells

transplantation (Fig. 5e), indicating human myogenic progenitor cells contributes to

structural repair of injured tendon and facilitates the healing process. Taken together,

these results suggest that human myogenic progenitor cells are capable of tendon

differentiation in vivo and contributing to tendon regeneration.

In sharp contrast to human myogenic progenitor cells, when 50,000 murine MuSCs

constitutively expressing tdTomato were transplanted to pre-injured tendon in NOD/SCID

mice under the same condition, less than 0.3% of tdTomato+ TNC+ cells were detected

(Fig. 5f and g). However, the myogenic differentiation potential of human myogenic

progenitor cells and mouse MuSCs was similar. Muscle injury was induced by muscular

injection of CTX in tibialis anterior (TA) muscle in NOD/SCID mice. TA muscles were

irradiated to kill the local MuSCs as described previously[34]. Transplantation of human

myogenic progenitor cells and mouse MuSCs to the irradiated pre-injured recipient mice

was performed, respectively. TA muscles were harvested after 28 days. Transplantation of

both human myogenic progenitor cells and murine MuSC displayed robust engraftment

efficiency (SFig. 3d-i). These results suggest that human myogenic progenitor cells and

murine MuSCs have similar myogenic differentiation potential.

Combined, the above results suggest that human myogenic progenitor cells have

dual differentiation potentials towards myogenesis or tenogenesis in vivo, while murine

MuSCs predominantly commit myogenesis.

Transplantation of human myogenic progenitor cells improves locomotor function

after tendon injury

We next checked whether the human myogenic progenitor cell transplantation

improves tendon functions. First, transmission electron microscope analysis was used to

evaluate microstructure of injured tendon two months after transplantation. Larger and

denser collagen fibrils of the tendons were identified in transplanted group than control

group with PBS injection, although the maturation level of collagen fibrils could still not

reach uninjured tendon (Fig. 6a and b). We next examined whether transplantation of

human myogenic progenitor cells could improve the biomechanical property of tendon.

The max failure load and stiffness of the tendons from the transplantation group were

significantly better than PBS injection control group (Fig. 6c). These results combined

suggest that transplantation of human myogenic progenitor cells improves the collagen

fibril maturation and biomechanical property of the injured tendon.

Whether the improved tendon regeneration could facilitate the whole organism

locomotor function was next investigated. Since the Achilles tendon transmits the

plantarflexion force from gastrocnemius muscle to calcaneus, the plantarflexion force of

involved leg was also performed two months after tendon injury. Expectedly,

transplantation of human myogenic progenitor cells for injured tendon also contributed to

improving both twitch and tetanus plantarflexion force when compared with PBS injection

control group, although the plantarflexion force could still not reach to level of uninjured

leg (Fig. 6d). Consistent with the improved plantarflexion force, the endurance time and

max fatigue speed of the mice transplanted with human myogenic progenitor cells was

better in treadmill test when compared to no transplantation group (Fig. 6e). Since

immunofluorescence staining with human Lamin A/C revealed that only a small number of

human myogenic progenitor cells engrafted in muscle sporadically (Fig. 5c and d), these

results suggest that the transplanted human myogenic progenitor cells can improve

locomotor function by directly repairing injured tendon.

TGFβ signaling pathway contributes to tenogenic differentiation of human

myogenic progenitor cells

Since the human myogenic progenitor cells and murine MuSCs shared the same

strain of recipient mice and the same tendon injury while being transplanted, they had the

similar microenvironment. Therefore, the distinct differentiation potentials are due to the

cell intrinsic differences between species. We further compared the expression profiles of

human myogenic progenitor cells and murine MuSCs. Interestingly, TGFβ signaling were

identified in KEGG enrichment analysis of upregulated genes in human myogenic

progenitor cells when compared with mouse muscle stem cells (Fig. 7a), indicating that

TGFβ signaling could be the key node for maintaining the tendon differentiation potential.

Furthermore, SMAD2 and SMAD3 were identified in upregulated gene set which was

enriched in TGFβ signaling pathway (Fig. 7b). Since TGFβ/SMAD2/SMAD3 axis plays a

crucial role in tendon development and tenogenic differentiation[22,23,26], we next

investigated the potential function of TGFβ signaling for tenogenic differentiation of human

myogenic progenitor cells. Consistently, increased phosphorylated SMAD2/SMAD3 was

confirmed after tenogenic induction of human myogenic progenitor cells (Fig. 7c). TGFβ

signaling inhibitor SB-431542 was used to significantly impaired TGFβ/SMAD2/SMAD3

signaling during tenogenic induction (Fig. 7c). The immunofluorescent staining, western

blot assay and RT-qPCR results showed SB-431542 significantly suppressed expression

level of tendon related genes SCX, TNC, COL I, MKX, and THBS4 (Fig. 7d, e, f). On the

contrary, myogenic differentiation ability of human myogenic progenitor cells was

increased after treatment of SB-431542 during tenogenic induction (Fig. 7g and h). Taken

together, these data indicated that TGFβ signaling pathway contributes to tenogenic

differentiation of human myogenic progenitor cells.

Discussion

Here we show that human myogenic progenitor cells have dual differentiation

potentials towards muscle and tendon. Transplantation of human myogenic progenitor

cells contributes to injured tendon repair and thus improves locomotor function. Thus, the

human myogenic progenitor cells could be served as a new source for tendon

regeneration.

Tendon disorders widely occur in people of all ages[22]. It disrupts the stability and

mobility of joint, which deeply affects their locomotor function and quality of life. However,

the natural healing of injured tendon is very slow due to hypocellularity and

hypovascularity of tendon. The biomechanical property and structural integrity could be

hardly completely recovered even with surgical treatment[35,36]. It is still a great challenge in

clinical work to treat tendon injury.

The relative inefficient outcome of routine therapy for tendon injury sparked the

exploration of stem cell treatment. Seed cells with the ability to differentiate into tenocytes

and secrete paracrine factors to repair tendon injury are preferred. Thus, tendon derived

stem cells, embryonic stem cells, induced pluripotent stem cells and mensenchymal stem

cells have been introduced as seed cells to treat tendon injury[37]. However, inadequate

sources of tendon derived stem cells, ethical issue and risk of teratoma formation of

embryonic stem cells or induced pluripotent stem cell, and heterogeneity of

mensenchymal stem cells limit the development of these seed cells for tendon injury

treatment. As for muscle progenitor cells, it is high in proliferation and abundant in sources,

and donor site morbidity after muscle harvest was low. Thus, muscle stem cell might be a

promising candidate as seed cells for tendon repair.

Here we also find that the differentiation potential of muscle stem cells is species

dependent. Human muscle myogenic progenitor cells are bipotential adult stem cells. In

contrast, murine muscle stem cells barely have tendon differentiation potential. The

species difference might be due to the higher TGFβ/SMAD2/SMAD3 signaling level in

human myogenic progenitor cells. TGFβ/SMAD2/SMAD3 axis plays a crucial role in

tendon development and tenogenic differentiation[22,23,26]. SMAD2/3-dependent TGFβ

signaling also acts as a crucial molecular brake for myogenesis. It could suppress

myogenic regulatory factors Myod1 and Myogenin[38,39], as well as inhibit myotube fusion

and muscle regeneration[40,41]. The elevated TGFβ/SMAD2/SMAD3 signaling may help

inhibit the myogenic differentiation ability and stimulate the tenogenic differentiation

potential of human myogenic progenitor cells under specific microenvironment.

Studies using mouse models contributed to the majority of our knowledge about

muscle stem cells and laid the foundation for our understanding of mammalian muscle

stem cells. However, human is dramatically different from mouse in many aspects such as

size, life span, and manners of motions. It is more demanding for humans to maintain the

homeostasis of the locomotion system and the whole organism locomotion ability in much

longer life span and bigger body size. Though our knowledge about human muscle

regeneration is limited, the current studies have revealed multiple differentiation potentials

of Pax7+ human progenitor cells in skeletal muscles. CD56+ CD34+ progenitor cells in

human skeletal muscle have been reported to have myogenic, osteogenic, and

adipogenic activity[14,15]. The CD56+ CD34- progenitor cells in human skeletal muscle have

been shown to be free of adipogenic potential[11]. Our results suggest that the progenitor

cells in skeletal muscles also have tenogenic differentiation ability besides their myogenic

differentiation ability. It seems that there are multiple subpopulations of myogenic

progenitor cells in skeletal muscle. They are all capable of muscle regeneration, while with

potentials to regenerate other components of the motion system such as bone, tendon,

and adipocytes. This could be an economic method to maintain the functions of the

motion system for the longer life span and more complicated motion manner in human

beings.

Methods

Animals

Animal care and use were in accordance with the guidelines of the animal facility

hosted by Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of

Sciences, and the operations were approved by the ethical committee of Shanghai

Institute of Biochemistry and Cell Biology. All mice were maintained in specific

pathogen-free (SPF) animal facility in individually ventilated cages (IVC) with controlled

temperature (22±1℃) and light (12h light/dark cycle). NOD/SCID mice were purchased

from Animal Model Research Center of Nanjing University. Pax7CreERT2 and

Rosa26-Flox-Stop-Flox-tdTomato mice were purchased from Jackson Laboratory. All

experiments were conducted on 3-month-old adult male mice.

Human samples

Peroneal longus muscle and remanent tendon were obtained from the wastes of

patients who underwent full-thickness peroneal longus tendon autograft for knee ligament

reconstructive procedures. The study was approved by the ethical committee of Xinhua

Hospital Affiliated to Shanghai Jiao Tong University, School of Medicine (Approval No.

XHEC-D-2019-043) and written informed consents were obtained from all donors.

Isolation of human muscle stem/progenitor cells

Human skeletal muscle from peroneal longus were dissected and digested as

described previously[6]. Briefly, muscle tissues were cut into small pieces and digested by

collagenase II (Worthington biochemical, 700-800U/ml, cat#LS004177) for 60min followed

by 30min of digestion with the mixtures of collagenase II and dispase (Life

Technologies,11U/ml, cat#17105-041). Digested cells were passed 10 times through a

20-gauge needle. Cell suspension was filtered through a 40μm cell strainer (BD Falcon,

cat#352340). Erythrocytes were removed by red blood cell lysis buffer (Thermo Fisher

Scientific, cat#00-433-57). The single cell suspension obtained from human muscle was

stained with a cocktail containing PE-Cy5 anti-human CD45 (BD Pharmingen,

cat#555484), Percp-Cy5.5 anti-human CD31 (BioLegend, cat#303132), AF-488

anti-human CD29 (BioLegend, cat#303016) and PE anti-human CD56 (BioLegend,

cat#304606) for 45 min at 4°C. CD45- CD31- CD29+ CD56+ human muscle

stem/progenitor cells were sorted by BD Influx sorter (BD Biosciences).

Isolation of mouse muscle stem cells

For mouse muscle stem cells isolation, dissected TA muscles were first digested with

10ml muscle digestion buffer (DMEM containing 1% penicillin/streptomycin, 0.125mg/ml

Dispase II (Roche, 04942078001), and 10mg/ml Collagenase D (Roche, 11088866001))

for 90 minutes at 37°C. The digestion was stopped by adding 2ml of FBS. The digested

cells were filtered through 70μm strainers. Red blood cells were lysed by 7ml RBC lysis

buffer (0.802% NH4Cl, 0.084% NaHCO3, 0.037% EDTA in ddH2O, pH7.2-7.4) for 30s,

then filter through 40μm strainers. After staining with antibody cocktails

(AF700-anti-mouse Sca-1, PerCP/Cy5.5-anti-mouse CD11b, PerCP/Cy5.5-anti-mouse

CD31, PerCP/Cy5.5-anti-mouse CD45, FITC anti-mouse CD34, APC-anti-mouse Integrin

α7+), the mononuclear cells were subjected for FACS analysis using Influx (BD

Biosciences). The population of PI-CD45-CD11b-CD31-Sca1- CD34+ Integrin α7+ cells

was collected.

Primary human tenocytes isolation

Tendon tissues were obtained from the discarded materials of tendon autograft

surgery. They were washed with PBS. Epi- and peri-tendon sheath were completely

removed. Tenocytes were isolated as described previously[20]. Briefly, the tendons were

minced to 1mm^3 pieces and digested with 3 mg/ml collagenase I (Worthington

biochemical, cat# LS004194) in DMEM (Gibco, cat#11965118) at 37℃ for 3hrs with

gentle agitation. The digested tissue was filtered through a 40μm cell strainer (BD Falcon,

cat#352340) and the isolated cells were plated for subsequent analysis.

Cell culture and differentiation

Primary human muscle stem/progenitor cells were plated in F10 basal medium

(Gibco, cat#11550043) containing 20% FBS (Gibco, cat#10-013-CV), 2.5ng/ml bFGF

(R&D, cat#233-FB-025) and 1% Penicillin-Streptomycin (Gibco, cat#15140-122) on

collagen-coated dishes. Mouse muscle stem cells were plated in F10 basal medium

(Gibco, cat#11550043) containing 20% FBS (Gibco, cat#10-013-CV), 2.5ng/ml bFGF

(R&D, cat#233-FB-025) and IL-1α, IL-13, IFN-γ and TNF-α as described previously[34].

DMEM (Gibco, cat#11965118) containing 0.4% Ultroser G (Pall Corporation,

cat#15950-17) and 1% Penicillin-Streptomycin were used to differentiate human muscle

stem/progenitor cells [5]. DMEM containing 2% horse serum (HyClone,

cat#HYCLSH30074.03HI) and 1% Penicillin-Streptomycin were used to differentiate

mouse muscle stem cells as described previously[42,43]. DMEM containing 10% FBS,

100ng/ml GDF5 (R&D, cat#8340-G5-050), GDF7 (R&D, cat#8386-G7-050), 0.2mM

ascorbic acid (Sigma-Aldrich, cat#A4403) and 1% Penicillin-Streptomycin were used to

induce tendon differentiation [44-46]. Primary tenocytes were cultured in DMEM containing

20% FBS, 2.5ng/ml bFGF (R&D, cat#233-FB-025) and 1% Penicillin-Streptomycin.

Single-cell RNA sequencing and trajectory analysis

The single cell suspension obtained from human muscle was firstly prepared. PI

(Sangon Biotech, cat#E607328) and Hoechst (Sangon Biotech, cat#A601112) were used

to sort live cells by FACS. Then the sored cells were washed twice with PBS containing

0.04% BSA, followed by library preparasion with Chromium Single Cell 3' Reagent Kits

(10X genomics, cat# 1000121-1000157). The sequencing was performed on Illumina

Novaseq 6000 platform (Illumina).

Single cell RNA-seq data were analyzed by Seurat R (Version 3.2.0) package. Cells

with less than 200 genes, more than 6,000 genes detected, and more than 10%

mitochondrial genes were excluded. A total of 15,102 cells were included for subsequent

analysis. Sequencing reads for each gene were normalized to total UMIs in each cell to

obtain normalized UMI values by "NormalizeData" function. The "ScaleData" function was

used to scale and center expression levels in the data set for dimensional reduction. Total

cell clustering was performed by "FindClusters" function at a resolution of 0.1 and

dimensionality reduction was performed with "RunUMAP" function. Trajectory analysis

was done by R package Monocle2[47].

Immunofluorescence staining

Cryosections were fixed in 4% formaldehyde for 15min, permeabilized in 0.5% Triton

X-100 for 15min, and stained with anti-Pax7 (Developmental Studies Hybridoma Bank),

anti-Laminin (Abcam, cat#ab11575), anti-Lamin A/C (Abcam, cat#ab108595;

cat#ab190380), anti-TNC (Abcam, cat#ab108930; cat#ab3970), anti-Scx (Abcam,

cat#ab58655), anti-Tnmd (Abcam, cat#ab203676), anti-Col I (Abcam, cat#ab260043),

anti-Col III (ABclonal, cat#A0817) or anti-MyHC (Millipore, cat#05-716) at 4°C overnight,

and incubated with Alexa 488-, Alexa 594 or Alexa 647-labeled anti-mouse or anti-rabbit

secondary antibodies (Invitrogen, 1:1000) at room temperature for one hour. The nuclei

were stained with 4,6-diamidino-2-phenylindole (DAPI, Vector Laboratories, cat#H-1200).

All images were acquired by Leica SP8 confocal microscope (Leica).

RNA scope in situ hybridization

Fresh cryosections of human muscle were obtained. Then the target gene-specific

RNAscope probes (RNAscope Probe-Hs-SCX, cat#593041; RNAscope

Probe-Hs-PAX7-C2, cat#546691) were used to investigate the cellular distributions of

Pax7 and SCX. All staining steps strictly followed the manufacturer's protocols by

Advanced Cell Diagnostics. The nuclei were then stained with

4,6-diamidino-2-phenylindole (DAPI, Vector Laboratories, cat#H-1200), and sections were

coverslipped with antifade mounting media (Vector Laboratories, cat#H-100). All images

were acquired by Leica SP8 confocal microscope or Olympus BX53 fluorescence

microscope (Leica).

Gene expression analysis

Total RNA was isolated using TRIzol Reagent (Invitrogen, cat#15596-018) according

to the manufacturer's instruction. GAPDH served as internal control. The primers for

RT-qPCR are listed as below:

Human GAPDH-F	5'-CAAGGCTGAGAACGGGAAGC-3'
Human GAPDH-R	5'-AGGGGGCAGAGATGATGACC-3'
Human SCX-F	5'-AGCGATTCGCAGTTAGGAGG-3'
Human SCX-R	5'-GTCTGTACGTCCGTCTGTCC-3'
Human COL I-F	5'-GGCTCCTGCTCCTCTTAGCG-3'
Human COL I-R	5'-CATGGTACCTGAGGCCGTTC-3'
Human TNC-F	5'-GGTGGATGGATTGTGTTCCTGAGA-3'
Human TNC-R	5'-CTGTGTCCTTGTCAAAGGTGGAGA-3'
Human THBS4-F	5'-TGCTGCCAGTCCTGACAGA-3'
Human THBS4-R	5'-GTTTAAGCGTCCCATCACAGTA-3'
Human MKX-F	5'-TTCAAGGCAATGCTGAACGG-3'
Human MKX-R	5'-CTCCCGCTTTGATGACCGAA-3'
Human COMP-F	5'-GATCACGTTCCTGAAAAACACG-3'
Human COMP-R	5'-GCTCTCCGTCTGGATGCAG-3'
Human PAX7-F	5'-ACCCCTGCCTAACCACATC-3′
Human PAX7-R	5'-GCGGCAAAGAATCTTGGAGAC-3′
Human MYF5-F	5'-AATTTGGGGACGAGTTTGTG-3′
Human MYF5-R	5'-CATGGTGGTGGACTTCCTCT-3′
Human MYOD1-F	5'-GACGGCATGATGGACTACAG-3′
Human MYOD1-R	5'-AGGCAGTCTAGGCTCGACAC-3′
Human MYOG-F	5'-GCTCAGCTCCCTCAACCA-3′
Human MYOG-R	5'-GCTGTGAGAGCTGCATTCG-3′
Human MYH1-F	5'-GGGAGACCTAAAATTGGCTCAA-3′
Human MYH1-R	5'-TTGCAGACCGCTCATTTCAAA-3′
Human MYH3-F	5'-ATTGCTTCGTGGTGGACTCAA-3′
Human MYH3-R	5'-GGCCATGTCTTCGATCCTGTC-3′
Human DESMIN-F	5'-TCGGCTCTAAGGGCTCCTC-3′
Human DESMIN-R	5'-CGTGGTCAGAAACTCCTGGTT-3′
Human MYL1-F	5'-GTTGAGGGTCTGCGTGTCTTT-3′
Human MYL1-R	5'-ACCCAGGGTGGCTAGAACA-3′
Mouse Gapdh-F	5'-ACCCAGAAGACTGTGGATGG-3′
Mouse Gapdh-R	5'-ACACATTGGGGGTAGGAACA-3′
Mouse Col I-F	5'-CCAGCGAAGAACTCATACAGC-3'
Mouse Col I-R	5'-GGACACCCCTTCTACGTTGT-3'
Mouse Scx-F	5'-GAGAACACCCAGCCCAAAC-3'
Mouse Scx-R	5'-TCACCCGCCTGTCCATC-3'
Mouse Tnc-F	5'-GAGCCCCTTTGCCTCAACAA-3'
Mouse Tnc-R	5'-CTTCGCCCGTGAAACCTTCTT-3'

Mouse Mkx-F	5'-TGGTTTCCTGGACAATCCACA-3'
Mouse Mkx-R	5'-CGCTTATGCCTTACCTTCCCTC-3'
Mouse Thbs4-F	5'-GCCACAAGCACAGGAGACTTT-3'
Mouse Thbs4-R	5'-TGACCTGCTGCCTCAGAAGA-3'
Mouse MyoG-F	GAGACATCCCCCTATTTCTACCA
Mouse MyoG-R	GCTCAGTCCGCTCATAGCC
Mouse Myh1-F	5'-GCGAATCGAGGCTCAGAACAA-3'
Mouse Myh1-R	5'-GTAGTTCCGCCTTCGGTCTTG-3'
Mouse Myh3-F	5'-ATGAGTAGCGACACCGAGATG-3'
Mouse Myh3-R	5'-ACAAAGCAGTAGGTTTTGGCAT-3'

Cloning assay

Primary human muscle stem/progenitor cells were first sorted in 96 well plates with

density of single cell per well. Tenogenic induction or myogenic induction was performed

after proliferation for each well. After induction, the immunofluorescence staining of SCX

or MyHC was performed in each well. The differentiation efficiency was determined by

calculating the ratio of total wells with positive fluorescence signal to total wells with alive

cells.

RNA-sequencing

Total RNA was isolated using TRIzol Reagent (Invitrogen, cat#15596-018) according

to the manufacturer's instruction. mRNA was enriched with magnetic oligo (dT) beads

(New England Biolabs, cat#S1419S). The cDNA library was constructed with mean inserts

of 200bp with non-stranded library preparation using NEBNext Ultra RNA Library Prep Kit

for Illumina (New England Biolabs, cat#E7530L). Sequencing was performed by a

paired-end 125 cycles rapid run on the Illumina HiSeq2500. Sequencing data were filtered

by SolexaAQ (Q > 20 and length ≥ 25 bp)[48].The adapter sequences and low quality

segments (Phred Quality Score<20) were trimmed using Cutadapt. Paired-end clean

reads were then mapped to the reference genome GRCh38.98 using HISAT2.

Htseq-count was used to quantify the gene expression value[49]. Read count of each gene

was normalized using FPKM (Fragments Per Kilo bases per Million fragments).

Differential expression (DE) analysis was performed using DESeq and significant DE

genes were defined as those with absolute log2FoldChange>1 and p< 0.05. Heatmap and

volcano plot of DE genes was generated by R package Pheatmap

(https://cran.r-project.org/web/packages/pheatmap/index.html) and EnhancedVolcano

(https://bioconductor.org/packages/release/bioc/vignettes/EnhancedVolcano/inst/doc/Enh

ancedVolcano.html), respectively. Gene enrichment analysis was conducted by R

package topGO with $p < 0.05$ as the cut-off. The differential expression analysis for

RNA-seq data between human and mouse myogenic progenitor cells was performed

according to previous literature[50].

Cell transplantation in TA muscle

A single dose of 18 Grey irradiation was administered to the hind legs of the recipient

NOD/SCID mice. TA muscle was injured by injecting 15μl of 10μM CTX (Sigma), and

50,000 human myogenic progenitor cells or 50,000 murine MuSCs suspended in 10μl

PBS were injected intramuscularly to the injury site as described [5].

Tendon injury for lineage tracing

Pax7^{CreERT2}:Rosa26^{tdTomato} mice were obtained and MuSC was labeled with tdTomato

as aforementioned. We used dedicated surgical instruments to imitate a mouse model

similar to previously published human surgery[33]. Firstly, distal gastrocnemius tendon was

woven with a 6-0 polypropylene non-absorbable suture (PROLENE, cat#EH7242H) and

released by microsurgical scissor. After that, a dedicated mini tendon stripper was

introduced over the free end of distal medial gastrocnemius tendon. The medial

gastrocnemius tendon could be totally removed after advancement of tendon stripper. To

further activate MuSCs to guarantee that sufficient amount of activated MuSCs were

available around the tendon injury site, cardiotoxin (CTX) was injected at the muscle

adjacent to the sites where the tendon was removed to induce more muscle injuries and

further activate MuSCs. Then the subcutaneous tissue and skin were sutured.

Tendon injury and cell transplantation

An approximately 1.5mm long and 0.5mm width transverse incision was performed at

5mm from the calcaneus in Achilles tendon for recipient NOD/SCID mice. 50,000 human

muscle stem/progenitor or mouse muscle stem cells resuspended in 20μl hydrogel were

injected to the injury site with 28-gauge needles. PBS mixed with 20μl hydrogel were

injected as control.

Biomechanical analysis

Two months after tendon injury, the tendons were harvested to biomechanical analysis using a universal tester (Instron 3345 load system, USA). The grippers were gradually moved apart with the speed of 5 mm/min until the tested tendon was completely ruptured. Then the max load was obtained and documented. The slope of the stress-strain curve was defined as stiffness. All these data were automatically presented in Instron 3345 load system.

Transmission electron microscope (TEM) examination

After washing the selected target tendon tissues with PBS, tissue fixation was performed for more than 2 hours. Post-fixation was conducted by 1% OsO_4 (Ted Pella Inc) in 0.1M PB for 1-2 hours. Then dehydration and drying were performed and target tendon samples were subsequently attached to metallic stubs for conductive metal coating. Images were obtained by TEM (HITACHI, SU8100) and at least 1000 collagen fibrils were evaluated for each sample. Density of fibrils was evaluated by the percentage of collagen fibril containing area.

Treadmill test

Mouse treadmill test was performed in Exer-3/6 treadmill apparatus with electrical stimulus. Treadmill exercise began with a 5 min warm-up, then each mouse was evaluated with specific protocol. For endurance test, mice ran on the treadmill at the constant speed of 22m/min until exhaustion. For exhaustion test, the treadmill speed was increased (2m/min each 3 min) at the beginning speed at 18m/min until exhaustion. Exhaustion was defined as inability to run on the treadmill for longer than10 seconds.

In vivo muscle force analysis

The 1300A 3-in-1 whole animal system (Aurora Scientific) was used for in vivo muscle force analysis. Mice were first anesthetized and kept warm by heat lamp. The foot was placed on a footplate and kept perpendicular to the tibia. The peroneal branch of the sciatic nerve was stimulated to evaluate the plantarflexion strength. Five repetitive tests

were performed for each limb and DMA software (Aurora scientific) was used for results analysis.

Statistical analysis

At least three biological replicates and technical repeats were performed in each experimental group. All experiments were analyzed and evaluated by investigators in a blinded manner. Error bars indicated standard deviation. Two-tailed unpaired Student's t-tests were used when variances were similar (tested with F-test) for comparison of two independent groups. One-way ANOVA tests followed by Dunnett's post-test or Tukey's post-test were used for multiple comparisons. Shapiro-Wilk tests were performed to determine data normality. Statistical analysis was performed in GraphPad Prism 7 (GraphPad Software, San Diego, USA) or SPSS version 19.0 for Windows (SPSS Inc., Chicago, IL, USA). It was considered significant with P value less than 0.05. Data were presented as mean ± standard unless stated.

Accession numbers

The complete sequencing data have been uploaded on to Sequence Read Archive database (PRJNA1012476 and PRJNA1012828).

Acknowledgements

We thank Dr. Dangsheng Li for helpful discussions, the National Protein Science Center (Shanghai) for helps on FACS sorting, the cell biology facility of SIBCB for helps on imaging and FACS analysis. This work was supported by the Strategic Priority Research Program of the Chinese Academy of Science (XDA16020400), National Science Foundation of China (32170804, 81871096, 82372384, 82302657).

References

1.   Fu, X., Wang, H., and Hu, P. (2015). Stem cell activation in skeletal muscle regeneration. Cell Mol Life Sci *72*, 1663-1677. 10.1007/s00018-014-1819-5.

2.   Relaix, F., Bencze, M., Borok, M.J., Der Vartanian, A., Gattazzo, F., Mademtzoglou, D., Perez-Diaz, S., Prola, A., Reyes-Fernandez, P.C., Rotini, A., and Taglietti, t. (2021). Perspectives on skeletal muscle stem cells. Nat Commun *12*, 692.

619   10.1038/s41467-020-20760-6.

620 3. Seale, P., Asakura, A., and Rudnicki, M.A. (2001). The potential of muscle stem cells.

621   Dev Cell *1*, 333-342. 10.1016/s1534-5807(01)00049-1.

622 4. Collins, C.A., Olsen, I., Zammit, P.S., Heslop, L., Petrie, A., Partridge, T.A., and

623   Morgan, J.E. (2005). Stem cell function, self-renewal, and behavioral heterogeneity of

624   cells from the adult muscle satellite cell niche. Cell *122*, 289-301.

625   10.1016/j.cell.2005.05.010.

626 5. Ding, S., Wang, F., Liu, Y., Li, S., Zhou, G., and Hu, P. (2017). Characterization and

627   isolation of highly purified porcine satellite cells. Cell death discovery *3*, 17003.

628   10.1038/cddiscovery.2017.3.

629 6. Shao, X., Fu, X., Yang, J., Sui, W., Li, S., Yang, W., Lin, X., Zhang, Y., Jia, M., Liu, H.,

630   et al. (2023). The asymmetrical ESR1 signaling in muscle progenitor cells determines

631   the progression of adolescent idiopathic scoliosis. Cell Discov *9*, 44.

632   10.1038/s41421-023-00531-5.

633 7. Barruet, E., Garcia, S.M., Striedinger, K., Wu, J., Lee, S., Byrnes, L., Wong, A.,

634   Xuefeng, S., Tamaki, S., Brack, A.S., and Pomerantz, J.H. (2020). Functionally

635   heterogeneous human satellite cells identified by single cell RNA sequencing. Elife *9*.

636   10.7554/eLife.51576.

637 8. Boldrin, L., Muntoni, F., and Morgan, J.E. (2010). Are human and mouse satellite cells

638   really the same? J Histochem Cytochem *58*, 941-955. 10.1369/jhc.2010.956201.

639 9. Bareja, A., Holt, J.A., Luo, G., Chang, C., Lin, J., Hinken, A.C., Freudenberg, J.M.,

640   Kraus, W.E., Evans, W.J., and Billin, A.N. (2014). Human and mouse skeletal muscle

stem cells: convergent and divergent mechanisms of myogenesis. PloS one *9*,

e90398. 10.1371/journal.pone.0090398.

10.   Spinazzola, J.M., and Gussoni, E. (2017). Isolation of Primary Human Skeletal Muscle

Cells. Bio Protoc *7*. 10.21769/BioProtoc.2591.

11.   Pisani, D.F., Dechesne, C.A., Sacconi, S., Delplace, S., Belmonte, N., Cochet, O.,

Clement, N., Wdziekonski, B., Villageois, A.P., Butori, C., et al. (2010). Isolation of a

highly myogenic CD34-negative subset of human skeletal muscle cells free of

adipogenic potential. Stem cells *28*, 753-764. 10.1002/stem.317.

12.   Xu, X., Wilschut, K.J., Kouklis, G., Tian, H., Hesse, R., Garland, C., Sbitany, H.,

Hansen, S., Seth, R., Knott, P.D., et al. (2015). Human Satellite Cell Transplantation

and Regeneration from Diverse Skeletal Muscles. Stem Cell Reports *5*, 419-434.

10.1016/j.stemcr.2015.07.016.

13.   Zheng, B., Cao, B., Crisan, M., Sun, B., Li, G., Logar, A., Yap, S., Pollett, J.B., Drowley,

L., Cassino, T., et al. (2007). Prospective identification of myogenic endothelial cells in

human skeletal muscle. Nature biotechnology *25*, 1025-1034. 10.1038/nbt1334.

14.   Castiglioni, A., Hettmer, S., Lynes, M.D., Rao, T.N., Tchessalova, D., Sinha, I., Lee,

B.T., Tseng, Y.H., and Wagers, A.J. (2014). Isolation of progenitors that exhibit

myogenic/osteogenic bipotency in vitro by fluorescence-activated cell sorting from

human fetal muscle. Stem cell reports *2*, 92-106. 10.1016/j.stemcr.2013.12.006.

15.   Pisani, D.F., Clement, N., Loubat, A., Plaisant, M., Sacconi, S., Kurzenne, J.Y.,

Desnuelle, C., Dani, C., and Dechesne, C.A. (2010). Hierarchization of myogenic and

adipogenic progenitors within human skeletal muscle. Stem cells *28*, 2182-2194.

10.1002/stem.537.

16.   Gaida, J.E., Alfredson, H., Kiss, Z.S., Bass, S.L., and Cook, J.L. (2010). Asymptomatic

Achilles tendon pathology is associated with a central fat distribution in men and a

peripheral fat distribution in women: a cross sectional study of 298 individuals. BMC

Musculoskelet Disord *11*, 41. 10.1186/1471-2474-11-41.

17.   Voleti, P.B., Buckley, M.R., and Soslowsky, L.J. (2012). Tendon healing: repair and

regeneration.   Annu   Rev   Biomed   Eng   *14*,   47-71.

10.1146/annurev-bioeng-071811-150122.

18.   Korcari, A., Nichols, A.E.C., Buckley, M.R., and Loiselle, A.E. (2023).

Scleraxis-lineage cells are required for tendon homeostasis and their depletion

induces an accelerated extracellular matrix aging phenotype. Elife *12*.

10.7554/eLife.84194.

19.   Pennisi,   E.   (2002).   Tending   tender   tendons.   Science   *295*,   1011.

10.1126/science.295.5557.1011.

20.   Bi, Y., Ehirchiou, D., Kilts, T.M., Inkson, C.A., Embree, M.C., Sonoyama, W., Li, L.,

Leet, A.I., Seo, B.M., Zhang, L., et al. (2007). Identification of tendon stem/progenitor

cells and the role of the extracellular matrix in their niche. Nat Med *13*, 1219-1227.

10.1038/nm1630.

21.   Kohler, J., Popov, C., Klotz, B., Alberton, P., Prall, W.C., Haasters, F., Muller-Deubert,

S., Ebert, R., Klein-Hitpass, L., Jakob, F., et al. (2013). Uncovering the cellular and

molecular changes in tendon stem/progenitor cells attributed to tendon aging and

degeneration. Aging Cell *12*, 988-999. 10.1111/acel.12124.

22. Nourissat, G., Berenbaum, F., and Duprez, D. (2015). Tendon injury: from biology to

tendon repair. Nature reviews. Rheumatology *11*, 223-233. 10.1038/nrrheum.2015.26.

23. Pryce, B.A., Watson, S.S., Murchison, N.D., Staverosky, J.A., Dunker, N., and

Schweitzer, R. (2009). Recruitment and maintenance of tendon progenitors by

TGFbeta signaling are essential for tendon formation. Development *136*, 1351-1361.

10.1242/dev.027342.

24. Yang, G., Rothrauff, B.B., Lin, H., Yu, S., and Tuan, R.S. (2017). Tendon-Derived

Extracellular Matrix Enhances Transforming Growth Factor-beta3-Induced Tenogenic

Differentiation of Human Adipose-Derived Stem Cells. Tissue engineering. Part A *23*,

166-176. 10.1089/ten.TEA.2015.0498.

25. Barsby, T., and Guest, D. (2013). Transforming growth factor beta3 promotes tendon

differentiation of equine embryo-derived stem cells. Tissue engineering. Part A *19*,

2156-2165. 10.1089/ten.TEA.2012.0372.

26. Berthet, E., Chen, C., Butcher, K., Schneider, R.A., Alliston, T., and Amirtharajah, M.

(2013). Smad3 binds Scleraxis and Mohawk and regulates tendon matrix organization.

Journal of orthopaedic research : official publication of the Orthopaedic Research

Society *31*, 1475-1483. 10.1002/jor.22382.

27. Wang, D., Pun, C.C.M., Huang, S., Tang, T.C.M., Ho, K.K.W., Rothrauff, B.B., Yung,

P.S.H., Blocki, A.M., Ker, E.D.F., and Tuan, R.S. (2020). Tendon-derived extracellular

matrix induces mesenchymal stem cell tenogenesis via an integrin/transforming

growth factor-beta crosstalk-mediated mechanism. FASEB J *34*, 8172-8186.

10.1096/fj.201902377RR.

28.   Li, M., Jia, J., Li, S., Cui, B., Huang, J., Guo, Z., Ma, K., Wang, L., and Cui, C. (2021).

Exosomes derived from tendon stem cells promote cell proliferation and migration

through the TGF beta signal pathway. Biochemical and biophysical research

communications *536*, 88-94. 10.1016/j.bbrc.2020.12.057.

29.   You, T., Yuan, S., Bai, L., Zhang, X., Chen, P., and Zhang, W. (2020). Benzyl alcohol

accelerates recovery from Achilles tendon injury, potentially via TGF-beta1/Smad2/3

pathway. Injury *51*, 1515-1521. 10.1016/j.injury.2020.03.058.

30.   Jones, E.R., Jones, G.C., Legerlotz, K., and Riley, G.P. (2013). Cyclical strain

modulates metalloprotease and matrix gene expression in human tenocytes via

activation of TGFbeta. Biochim Biophys Acta *1833*, 2596-2607.

10.1016/j.bbamcr.2013.06.019.

31.   Rubenstein, A.B., Smith, G.R., Raue, U., Begue, G., Minchev, K., Ruf-Zamojski, F.,

Nair, V.D., Wang, X., Zhou, L., Zaslavsky, E., et al. (2020). Single-cell transcriptional

profiles in human skeletal muscle. Scientific reports *10*, 229.

10.1038/s41598-019-57110-6.

32.   Han, S.H., Kim, H.K., Ahn, J.H., Lee, D.H., Baek, M., Ye, G., Lee, J.M., Min, K., Oh, C.,

and Lee, S. (2018). A Protocol to Acquire the Degenerative Tenocyte from Humans. J

Vis Exp. 10.3791/57634.

33.   Shao, X., Shi, L.L., Bluman, E.M., Wang, S., Xu, X., Chen, X., and Wang, J. (2020).

Satisfactory functional and MRI outcomes at the foot and ankle following harvesting of

full thickness peroneus longus tendon graft. The bone & joint journal *102-B*, 205-211.

10.1302/0301-620X.102B2.BJJ-2019-0949.R1.

34.   Fu, X., Xiao, J., Wei, Y., Li, S., Liu, Y., Yin, J., Sun, K., Sun, H., Wang, H., Zhang, Z.,

730   et al. (2015). Combination of inflammation-related cytokines promotes long-term

731   muscle stem cell expansion. Cell Res *25*, 655-673. 10.1038/cr.2015.58.

732   35.   Rossi, L.A., Atala, N., Bertona, A., Tanoira, I., Bongiovanni, S., Maignon, G., and

733   Ranalletta, M. (2018). Return to Sports After in Situ Arthroscopic Repair of Partial

734   Rotator Cuff Tears. Arthroscopy : the journal of arthroscopic & related surgery : official

735   publication of the Arthroscopy Association of North America and the International

736   Arthroscopy Association. 10.1016/j.arthro.2018.07.037.

737   36.   Bianco, S.T., Moser, H.L., Galatz, L.M., and Huang, A.H. (2018). Biologics and stem

738   cell-based therapies for rotator cuff repair. Annals of the New York Academy of

739   Sciences. 10.1111/nyas.13918.

740   37.   Lui, P.P. (2015). Stem cell technology for tendon regeneration: current status,

741   challenges, and future research directions. Stem cells and cloning : advances and

742   applications *8*, 163-174. 10.2147/SCCAA.S60832.

743   38.   Liu, D., Black, B.L., and Derynck, R. (2001). TGF-beta inhibits muscle differentiation

744   through functional repression of myogenic transcription factors by Smad3. Genes Dev

745   *15*, 2950-2966. 10.1101/gad.925901.

746   39.   Brennan, T.J., Edmondson, D.G., Li, L., and Olson, E.N. (1991). Transforming growth

747   factor beta represses the actions of myogenin through a mechanism independent of

748   DNA binding. Proc Natl Acad Sci U S A *88*, 3822-3826. 10.1073/pnas.88.9.3822.

749   40.   Melendez, J., Sieiro, D., Salgado, D., Morin, V., Dejardin, M.J., Zhou, C., Mullen, A.C.,

750   and Marcelle, C. (2021). TGFbeta signalling acts as a molecular brake of myoblast

fusion. Nat Commun *12*, 749. 10.1038/s41467-020-20290-1.

41.   Girardi, F., Taleb, A., Ebrahimi, M., Datye, A., Gamage, D.G., Peccate, C., Giordani, L.,

Millay, D.P., Gilbert, P.M., Cadot, B., and Le Grand, F. (2021). TGFbeta signaling

curbs cell fusion and muscle regeneration. Nat Commun *12*, 750.

10.1038/s41467-020-20289-8.

42.   Luo, Y., Fu, C., Wang, Z., Zhang, Z., Wang, H., and Liu, Y. (2015). Mangiferin

attenuates contusive spinal cord injury in rats through the regulation of oxidative stress,

inflammation and the Bcl2 and Bax pathway. Mol Med Rep *12*, 7132-7138.

10.3892/mmr.2015.4274.

43.   Yin, J., Yang, L., Xie, Y., Liu, Y., Li, S., Yang, W., Xu, B., Ji, H., Ding, L., Wang, K., et

al. (2018). Dkk3 dependent transcriptional regulation controls age related skeletal

muscle atrophy. Nat Commun *9*, 1752. 10.1038/s41467-018-04038-6.

44.   Otabe, K., Nakahara, H., Hasegawa, A., Matsukawa, T., Ayabe, F., Onizuka, N., Inui,

M., Takada, S., Ito, Y., Sekiya, I., et al. (2015). Transcription factor Mohawk controls

tenogenic differentiation of bone marrow mesenchymal stem cells in vitro and in vivo.

J Orthop Res *33*, 1-8. 10.1002/jor.22750.

45.   Lou, J., Tu, Y., Burns, M., Silva, M.J., and Manske, P. (2001). BMP-12 gene transfer

augmentation of lacerated tendon repair. J Orthop Res *19*, 1199-1202.

10.1016/S0736-0266(01)00042-0.

46.   Wang, D., Jiang, X., Lu, A., Tu, M., Huang, W., and Huang, P. (2018). BMP14 induces

tenogenic differentiation of bone marrow mesenchymal stem cells in vitro.

Experimental and therapeutic medicine *16*, 1165-1174. 10.3892/etm.2018.6293.

47.   Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).

Reversed graph embedding resolves complex single-cell trajectories. Nat Methods *14*,

979-982. 10.1038/nmeth.4402.

48.   Cox, M.P., Peterson, D.A., and Biggs, P.J. (2010). SolexaQA: At-a-glance quality

assessment of Illumina second-generation sequencing data. BMC Bioinformatics *11*,

485. 10.1186/1471-2105-11-485.

49.   Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with

high-throughput   sequencing   data.   Bioinformatics   *31*,   166-169.

10.1093/bioinformatics/btu638.

50.   Zhou, Y., Zhu, J., Tong, T., Wang, J., Lin, B., and Zhang, J. (2019). A statistical

normalization method and differential expression analysis for RNA-seq data between

different species. BMC Bioinformatics *20*, 163. 10.1186/s12859-019-2745-1.

Figure legends

Figure 1. Potential tendon differentiation ability of human myogenic progenitor cells.

a. UMAP plot of all cells colored by cell types.

b. fDot blot of representative genes in each cell type.

c. Pseudotime trajectory inference of muscle progenitor cells, tenocytes and muscle

cells. Different cell types were annotated.

d. Heatmap showing the different expression levels of specific markers during the

trajectory.

e. RNA FISH of PAX7 and SCX in human skeletal muscle section. Scale bars, 10μm.

f. Statistical analysis of the percentage of PAX7+ or SCX+ cells by RNA FISH in human

skeletal muscle section. Error bars indicated standard deviation (n=5).

g. Immunofluorescence staining of PAX7, MYOD1 and MYF5 in primary human

myogenic progenitor cells. Scale bars, 100μm.

h. Statistical analysis of the percentage of PAX7+, MYOD1+ and MYF5+ cells in the

isolated myogenic progenitor cells. Error bars indicated standard deviation (n=5).

i. Immunofluorescence staining of MyHC in myotubes differentiated from human

myogenic progenitor cells. Primary human myogenic progenitor cells were isolated

and differentiated to myotubes for 5 days followed by MyHC immunofluorescence

staining. Scale bars, 100 μm.

j. Statistical analysis of the percentage of the percentage of nuclei in MyHC+ myotubes

after differentiation of the myogenic progenitor cells. Error bars indicated standard

deviation (n=5).

k. Relative expression level of PAX7, MYF5, and MYOD1 in human myogenic progenitor

cells and differentiated myotubes. RT-qPCR assays were performed with human

myogenic progenitor cells. Error bars indicated standard deviation (n=3). *** indicated

p<0.001.

l. Relative expression level of MYOG, MYH1, and MYH3 in human myogenic progenitor

cells and differentiated myotubes. RT-qPCR assays were performed with human

muscle myogenic progenitor cells after myogenic differentiation. Error bars indicated

standard deviation (n=3). *** indicated p<0.001.

Shao et al, Figure 2

Figure 2. Human myogenic progenitor cells display tendon differentiation ability in

vitro.

a. Immunofluorescence staining of tendon marker TNC and SCX in human myogenic

progenitor cells induced for myogenic and tenogenic differentiation, respectively.

Scale bars, 100 µm.

b. Quantification of TNC and SCX fluorescent intensity in human myogenic progenitor

cells undergone myogenic and tenogenic differentiation, respectively. Error bars

indicated standard deviation (n=5).

c. Relative expression levels of genes enriched in tendon cells. RT-qPCR assays were

performed with human myogenic progenitor cells upon myogenic and tenogenic

differentiation, respectively. Error bars indicated standard deviation (n=3). ***

indicated p<0.001.

d. Scheme of clonal proliferation and differentiation assay.

e. Representative immunofluorescence staining images of MyHC as the marker for

successful myogenic differentiation. Scale bars, 100 µm.

f. Representative immunofluorescence staining images of SCX as the marker for

successful tenogenic differentiation. Scale bars, 100 µm.

g. Statistical analysis of the myogenic and tenogenic differentiation efficiency of human

myogenic progenitor cells. Error bars indicated standard deviation (n=5).

Figure 3. Human myogenic progenitor cells have tendon differentiation potential.

a. Heat map of gene expression profiles of human myogenic progenitor cells, human

muscle myogenic progenitor cells after myogenic differentiation, human myogenic

progenitor cells after tenogenic differentiation, and primary tenocytes.

b. Volcano plot of genes enriched in myogenic differentiation of human myogenic

progenitor cells.

c. Volcano plot of genes enriched in tenogenic differentiation of human myogenic

progenitor cells.

d. Bubble chart of GO analysis of cellular process up-regulated in myogenic

differentiation of human myogenic progenitor cells.

e. Bubble chart of GO analysis of cellular process up-regulated in tenogenic

differentiation of human myogenic progenitor cells.

Shao et al, Figure 4

Figure 4. Murine MuSCs display poor tenogenic differentiation ability.

a. Immunofluorescence staining of murine MuSCs directed towards myogenic and

tenogenic differentiation, respectively. The undifferentiated murine MuSCs were

stained with Pax7 and DAPI. Murine MuSCs after myogenic differentiation were

stained with MyHC and DAPI. Murine MuSCs after tenogenic differentiation were

stained with Scx and DAPI. Scale bars, 100 µm.

b. Statistical analysis of the percentage of nuclei in MyHC+ myotubes. Error bars

indicated standard deviation (n=5).

c. Relative expression levels of myogenic differentiation specific genes MyoG, Myh1,

and Myh3. Mouse MuSCs were induced to differentiate towards muscle and tendon,

respectively. Mouse MuSCs before and after differentiation together with primary

tenocytes were harvested and subjected for RT-qPCR analysis. Error bars indicated

standard deviation (n=3). * indicated $p<0.05$, ** indicated $p<0.01$, n.s. indicated

$n>0.05$.

d. Relative expression levels of tenogenic differentiation marker genes Scx, Tnc, Col I,

Mkx and Thbs4. Mouse MuSCs were induced to differentiate towards muscle and

tendon, respectively. Mouse MuSCs before and after differentiation together with

primary tenocytes were harvested and subjected for RT-qPCR analysis. Error bars

indicated standard deviation (n=3). n.s. indicated $n>0.05$.

e. Scheme of Pax7+ MuSC progeny lineage tracing.

f. Immunofluorescence staining of tendon tissue 4 months after injury. tdTomato+ cells

indicated the progeny of Pax7+ MuSCs. TNC+ cells indicated tendon cells. MyHC+

cells indicated muscle fibers. DAPI indicated nuclei staining. Merged indicated the

merged images of tdTomato, TNC, MyHC, and DAPI. The upper panel indicated the

low magnification images. Scale bars, 100 µm. The lower panel indicated the high

magnification images of the region label by white square in the upper panel. Scale

bars, 10 µm.

g. The statistical analysis of the percentage of tdTomato+ TNC+ cell 4 months after

tendon injury. Error bars indicated standard deviation (n=5).

Figure 5. Transplantation of human myogenic progenitor cells facilitates tendon regeneration.

a. Scheme of muscle stem/progenitor cell transplantation.

b. Immunofluorescence staining of the regenerated tendon-like tissue after human

myogenic progenitor cell transplantation. Tendon injury was induced in recipient

NOD/SCID mice and 50,000 human myogenic progenitor cells were transplanted to

the injured tendon at the injured site. The regenerated tendon-like tissue, the

connected muscle, and the surrounding soft tissues were harvested to make

continuous cryosections. One of the continuous cryosections were subjected for

immunofluorescence staining of tendon marker human specific TNC and human

specific Lamin A/C. DAPI indicated nuclei staining. Merge indicated the merged

images of human TNC, human Lamin A/C, and DAPI. The white lines indicated the

location of regenerated tendon-like tissues from human myogenic progenitor cells

based on human TNC staining. The yellow dashed lines indicated the superimposed

location of muscle based on MyHC staining in panel c. Scale bars, 100 µm.

c. Immunofluorescence staining of MyHC and human Lamin A/C. The regenerated

tendon-like tissue, the connected muscle, and the surrounding soft tissues were

harvested and subjected for continuous cryosection. One of the continuous

cryosections were stained for MyHC which was specifically expressed in skeletal

muscle, and human Lamin A/C to label human cells. DAPI indicated nuclei staining.

Merge indicated the merged images of MyHC, human Lamin A/C, and DAPI. The

white lines indicated the location of regenerated tendon-like tissue from human

myogenic progenitor cells based on human TNC staining in panel b. The yellow

dashed lines indicated the location of muscle based on MyHC staining. Scale bars,

100µm.

d. Statistical analysis of the percentage of human cells expressing skeletal muscle

marker MyHC or tendon marker TNC after being transplanted to the injured tendon.

Error bars indicated standard deviation (n=5).

e. Immunofluorescence staining of Col I, Col III and human Lamin A/C. Two months after

human cell transplantation, continuous cryosections containing the regenerated

tendon-like tissue and native tendon tissue was stained with Col I, Col III and human

Lamin A/C. DAPI indicated the staining of nuclei. T, native tendon tissue; W, wound

tendon tissue. Scale bars, 50 µm.

f.   Immunofluorescence staining of tendon tissue after transplantation of tdTomato+

murine MuSCs. tdTomato indicated the progenies of murine MuSCs. TNC indicated

immunofluorescence staining of tendon marker TNC. MyHC indicated

immunofluorescence staining of myofiber marker MyHC. DAPI indicated nuclei

staining. Merge indicated merged images of tdTomato, TNC, MyHC, and DAPI. The

upper panel indicated low magnification images. Scale bars, 100 µm. The lower panel

indicated the amplified images of the region indicated by the white square. Scale bars,

10 µm.

g. Statistical analysis of TNC+ tdTomato+ cells in tendon tissue after transplantation of

murine MuSCs. Error bars indicated standard deviation (n=5).

Figure 6. Transplantation of human myogenic progenitor cells improves the locomotor function after tendon injury.

a.   TEM images of collagen fibrils in the injured tendon with PBS injection, injured tendon

with human myogenic progenitor cell transplantation, and uninjured tendon. Scale

bars, 500nm.

b.   Statistical analysis of the collagen fibril diameter and percentage of collagen fibril area

for the injured tendon with PBS injection, injured tendon with human myogenic

progenitor cell transplantation, and uninjured tendon. Error bars indicated standard

deviation (n=5). * indicated $p<0.05$, *** indicated $p<0.001$.

c.   Statistical analysis of the max load and stiffness of the injured tendon with PBS

injection, injured tendon with human myogenic progenitor cell transplantation, and

uninjured tendon. Error bars indicated standard deviation (n=5). * indicated $p<0.05$, **

indicated $p<0.01$, *** indicated $p<0.001$.

d.   Twitch and tetanus plantarflexion force of the involved limb with PBS injection after

tendon injury, human myogenic progenitor cell transplantation after tendon injury, and

uninjured tendon. Error bars indicated standard deviation (n=5). * indicated $p<0.05$, **

indicated $p<0.01$, *** indicated $p<0.001$.

e.   The treadmill exercise for tendon injured mice with or without human myogenic

progenitor cell transplantation. The endurance time and max fatigue speed were

compared. Error bars indicated standard deviation (n=4). * indicated $p<0.05$, **

indicated $p<0.01$, ***indicated $p<0.001$ and n.s. indicated $p>0.05$.

Figure 7. TGFβ signaling pathway contributes to tenogenic differentiation of human

myogenic progenitor cells.

a. Bubble chart of KEGG enrichment analysis of upregulated genes in human myogenic

progenitor cells when compared with mouse muscle stem cells.

b. Heatmap of detailed upregulated genes in human myogenic progenitor cells which

were enriched in TGFβ signaling pathway.

c. Protein levels of SMAD2/SMAD3 and phosphorylated SMAD2/SMAD3. Human

myogenic progenitor cells were cultured in growth medium, tenogenic induction

medium with or without TGFβ signaling pathway inhibitor SB-431542 for 2 hours. Total

protein was extracted from cells with different treatments and subjected for total

SMAD2/SMAD3 and phosphorylated SMAD2/SMAD3 immunoblotting. GAPDH was

served as loading control.

d. Immunofluorescence staining of tendon marker TNC and SCX in human myogenic

progenitor cells induced for tenogenic differentiation with or without TGFβ signaling

pathway inhibitor SB-431542 for 12 days, respectively. Scale bars, 100 µm.

e. Protein levels of TNC and SCX. Human myogenic progenitor cells were induced

towards tenogenic differentiation with or without TGFβ signaling pathway inhibitor

SB-431542 for 12 days, respectively. Total protein was extracted from cells before and

after differentiation and subjected for TNC and SCX immunoblotting. GAPDH was

served as loading control.

f. Relative expression levels of tendon related genes. RT-qPCR assays were performed

with human myogenic progenitor cells upon tenogenic differentiation with or without

TGFβ signaling pathway inhibitor SB-431542 for 12 days, respectively. Error bars

indicated standard deviation (n=3). *** indicated $p<0.001$.

g. Immunofluorescence staining of myogenic differentiation marker MyHC in human

myogenic progenitor cells induced for tenogenic differentiation with or without TGFβ

signaling pathway inhibitor SB-431542 for 12 days, respectively. Scale bars, 50 µm.

h. Relative expression levels of muscle related genes. RT-qPCR assays were performed

with human myogenic progenitor cells upon tenogenic differentiation with or without

TGFβ signaling pathway inhibitor SB-431542 for 12 days, respectively. Error bars

indicated standard deviation (n=3). *** indicated $p<0.001$.

 Supplemental Figure legend

Shao et al, Supplemental Figure 1
Related to Figure2

Supplemental Figure 1. Human myogenic progenitor cells display tenogenic

differentiation potential.

a. Phase contrast images of human myogenic progenitor cells before and after
differentiation induction. Scale bars, 100 µm.

b. Immunofluorescence staining of myogenic differentiation marker genes MYOG and
MyHC. Human myogenic progenitor cells were induced for tenogenic differentiation.
The differentiated cells were stained with MYOG and MyHC antibody. Red indicated
MYOG or MyHC; DAPI indicated nuclei staining; merge indicated the merged images
of red and blue staining. Scale bars, 100µm.

c. Statistical analysis of the percentage of human cells expression MYOG or MyHC
under tenogenic differentiation condition. Error bars indicated standard deviation
(n=5).

d. Expression level of genes marking myogenic differentiation. Human myogenic
progenitor cells were induced for myogenic and tenogenic differentiation, respectively.
The undifferentiated and differentiated cells were subjected for RT-qPCR analysis of
MYH1, MYH3, DESMIN, and MYL1. Error bars indicated standard deviation (n=5). ***
indicated $p<0.001$.

e. Protein levels of TNC and MyHC. Human myogenic progenitor cells were induced
towards myogenic and tenogenic differentiation, respectively. Total protein was
extracted from cells before and after differentiation and subjected for TNC and MyHC
immunoblotting. HSP70 and GAPDH were served as loading control.

1003

Supplemental Figure 2. Murine MuSCs display poor tenogenic differentiation ability.

a. Immunofluorescence images of Pax7, MyoD1 and Vcam1 for mouse muscle
CD29+/CD56+ cells and Vcam1+ cells isolated by FACs. Scale bars, 50μm.

b. Statistical analysis of percentage of Pax7, MyoD1 and Vcam1 positive cells in mouse
muscle CD29+/CD56+ cells and Vcam1+ cells isolated by FACs. *** indicated
p<0.001.

c. Relative expression levels of tendon related genes. RT-qPCR assays were performed
with murine muscle CD29+/CD56+ cells before and after tenogenic differentiation, and
tenocytes. Error bars indicated standard deviation (n=3). *** indicated p<0.001.

d. Schematic illustration of the mouse line for murine MuSCs lineage tracing.

e. The surgery in mice mimicking human peroneus longus tendon removal surgery. The
medial gastrocnemius tendon was completely removed. Scale bars, 1mm.

f. Immunofluorescence images of injured muscle adjacent to the removed tendon. The
MuSCs were labeled with tdTomato before injury. The tdTomato positive myofibers
after muscle injury were analyzed to evaluate the tracing system. Scale bars, 100μm.

g. Statistical analysis of the tdTomato positive myofibers adjacent to the removed tendon.
Error bars indicated standard deviation (n=5).

1021

Supplemental Figure 3. Transplantation of human myogenic progenitor cells

facilitates tendon regeneration and transplantation of human/murine myogenic

progenitor cells improve muscle regeneration.

a.   Immunofluorescence staining of SCX and human specific Lamin A/C in injured tendon.

Two months after human cells transplantation, the injured tendon was stained with

SCX and human specific Lamin A/C. DAPI indicated nuclei staining. Merge indicated

the merged images of SCX, human specific Lamin A/C, and DAPI. Scale bars, 20 µm.

b.   Immunofluorescence staining of TNMD and human specific Lamin A/C in injured

tendon. Two months after human cell transplantation, the injured tendon was stained

with TNMD and human specific Lamin A/C. DAPI indicated nuclei staining. Merge

indicated the merged images of TNMD, human Lamin A/C, and DAPI. Scale bars, 20

µm.

c.   Statistical analysis of the percentage of human cells expressing tendon related

markers SCX or TNMD after being transplanted to the injured tendon. Error bars

indicated standard deviation (n=5).

d.   Scheme of human myogenic progenitor cell transplantation after muscle injury. TA

muscles were first irradiated to kill the local MuSCs in NOD/SCID mice. Then

transplantation of human myogenic progenitor cells to the irradiated pre-injured

recipient mice was performed and TA muscles were harvested after 28 days.

e.   Immunofluorescence staining of MyHC and human specific Lamin A/C after human

myogenic progenitor cells transplantation. Scale bars, 100µm.

f.   Statistical analysis of the engraftment efficiency of human myogenic progenitor cell

transplantation in muscle. Error bars indicated standard deviation (n=5).

g.   Scheme of murine MuSC transplantation after muscle injury. TA muscles were first

irradiated to kill the local MuSCs in NOD/SCID mice. Then transplantation of tdTomato

positive MuSCs to the irradiated pre-injured recipient mice was performed and TA

muscles were harvested after 28 days.

h.   Immunofluorescence staining of Laminin and murine MuSCs constitutively expressing

tdTomato after murine tdTomato positive MuSCs transplantation. Scale bars, 100µm.

i.   Statistical analysis of the engraftment efficiency of murine MuSCs transplantation in

muscle. Error bars indicated standard deviation (n=5).

1053

1054

Data availability

The datasets used in this study can be obtained from the corresponding author upon reasonable request.

Conflict of interest disclosure

The authors declared no conflict of interest.

Funding:

This work was supported by the Strategic Priority Research Program of the Chinese Academy of Science (XDA16020400), National Science Foundation of China (32170804, 81871096, 82372384, 82302657).

Contributors

XS, XL, HL, HZ, and XF designed the study, collected the experimental data and wrote the original draft. SL and SZ conducted the image analysis, statistical analysis. SZ designed the study, reviewed and edited the manuscript. JW and PH administrated the project and edited the manuscript. Xiexiang Shao, Xingzuan Lin and Hao Zhou contributed equally to this work.