1   Investigating the translational value of Periprosthetic Joint Infection (PJI) models to

2   determine the risk and severity of Staphylococcal biofilms.

Amita Sekar[1,2#], Yingfang Fan[1,2#], Peyton Tierney[1], Madeline McCanne[1], Parker Jones[1], Fawaz

Malick[1], Devika Kannambadi[1], Keith K Wannomae[1], Nicoletta Inverardi[1,2], Orhun Muratoglu[1,2],

Ebru Oral[1,2]*

[1]Harris Orthopaedics laboratory, Massachusetts General Hospital, Boston, USA

[2]Department of Orthopaedic Surgery, Harvard Medical School, Boston USA

*Corresponding author: eoral@mgh.harvard.edu

#These authors contributed equally.

11   Running Title: Translational Value of PJI Models in Assessing Staphylococcal Biofilm Risk

12   Abstract

With the advent of antibiotic-eluting polymeric materials for targeting recalcitrant infections, using

preclinical models to study biofilm is crucial for improving the treatment efficacy in periprosthetic

joint infections. The stratification of risk and severity of infections is needed to develop an

effective clinical dosing framework with better outcomes. Here, using in-vivo and in-vitro

implant-associated infection models, we demonstrate that methicillin-sensitive and resistant

Staphylococcus aureus (MSSA and MRSA) have model-dependent distinct implant and peri-

implant tissue colonization patterns. The maturity of biofilms and the location (implant vs tissue)

were found to influence the antibiotic susceptibility evolution profiles of MSSA and MRSA and

the models could capture the differing host-microbe interactions in vivo. Gene expression studies

revealed the molecular heterogeneity of colonizing bacterial populations. The comparison and

stratification of the risk and severity of infection across different preclinical models provided in

this study can guide clinical dosing to effectively prevent or treat PJI.

Keywords

Implant infections, in vivo models, *Staphylococcus aureus*, antibiotic resistance, drug delivery

devices

Introduction

Periprosthetic joint infections (PJIs) represent a formidable challenge to the success of total joint

replacements. These infections, characterized by microbial colonization on the implant materials

and the surrounding tissue, not only compromise the intended function of the implants but also

significantly reduce the quality of life of the patients [1–3]. In addition, they are increasingly hard

to treat with recurrent infections causing significantly increased morbidity and mortality [4].

Currently, a generalized approach is used in treating suspected PJI, where one of several avenues

of treatment is utilized [5,6]. While systemic antibiotics are the main tool in addressing bacterial

infections, the implants can be retained or replaced in a one- or two-stage revision [7]. Locally,

the elution of aminoglycosides together with vancomycin from antibiotic-eluting bone cement is

used to support the role of systemic antibiotics [8]. It is believed that the high concentrations

achieved by local administration/elution of drugs will lead to higher efficacy in addressing joint

infections while reducing systemic side effects of antibiotics such as nephrotoxicity. However,

there is little conclusive information on the precise effects of local antibiotic administration and

their efficacy in preventing/treating PJI and there is no specific dosing guidance for local

antibiotics [9–13]. Commonly used dosing for prophylaxis may lead to the worsening of outcomes

[14]. Thus, there is a great need to determine the dosing requirements for antibiotics to prevent

and treat PJI locally.

Methicillin-susceptible (MSSA) and methicillin-resistant (MRSA) *S. aureus* have been

established as the most significant causative organisms of periprosthetic joint infections [15,16].

These bacterial strains adhere to inanimate surfaces and the surrounding tissue using specific cell

surface proteins and adhesins such as elastin-binding proteins[17]. Further aggregation triggers

bacterial regulatory pathways promoting the production of extracellular polymeric substances such

as polysaccharide intercellular adhesion (PIA) regulated by *ica* operon) that establishes robust

biofilms[18]. The microbial communities in biofilm status undergo significant molecular,

physiological, and morphological changes which provide bacteria with resilience against host

defenses and conventional antibiotics such as gentamicin[19,20]. These structured microbial

communities, with their cell wall modifications and an intricate architecture comprising live and

dead bacterial aggregates together with host cell components embedded in a complex polymeric

matrix, create diffusion barriers leading to a high tolerance to antibiotics and host factors[21].

They also serve as a reservoir for maintaining the hardy bacterial populations that can persist in

the presence of antibiotics and infiltrate deep tissue spaces[22,23]. Heterogenous subpopulations

of bacteria in the biofilm present varying phenotypic and genotypic profiles within the same

biomass. These subpopulations differentially specialize in pathogenesis, drug resistance, and

evading immune responses[24,25]. Due to the reasons mentioned above, antibiotic-based

treatments are often not effective against these diverse populations and physical removal (radical

debridement) is necessary for definitive treatment. Despite debridement, antibiotic lavage,

replacement of implant components, and local elution from antibiotic-eluting bone cement, failure

rates are high in PJI[26–28].

While the significance of biofilms in periprosthetic joint infections is acknowledged[29,30], there remains a substantial knowledge gap concerning the infection dynamics, phenotypic and genotypic heterogeneity, and the time-dependent properties of antibiotic susceptibility. In-vitro models may not capture the biofilm properties in response to a realistic environment, despite providing a straightforward approach to studying biofilms. More recently, studies have attempted to characterize and correlate biofilm formation and antibiotic susceptibility of clinical isolates to outcomes of PJI[31–33]. However, the characterization of ex vivo-grown bacterial cultures is also limited in capturing the susceptibility dynamics in vivo. There is also little information on emerging heterogenous populations based on colonizing location (implant surface vs tissue), vascularization, and biofilm physiology. There is a need for a better understanding of the infecting organism together with their innate and adaptive behavior within a PJI setting to determine the risk and severity of the infection. This crucial knowledge can aid in the design and development of multi-faceted drug-eluting materials and treatment algorithms.

Our long-term goal is to design antibiotic-eluting polymeric materials that can be more efficiently used locally to prevent and treat PJI. As material advancements and preclinical research enhance our understanding and prevention of periprosthetic joint infections (PJI), notable discrepancies are exposed between promising preclinical findings and clinical testing outcomes. One factor contributing to the lack of predictability of the clinical outcomes associated with PJI treatment is the variability in the bacterial strain and the timing of treatment based on clinical symptoms[34,35]. In this study, we developed an in-vitro implant material infection model, a subcutaneous implantation and infection model, and a periprosthetic joint infection model in the rat, modeling *S. aureus* infections with varying risk and severity. To capture the range of therapeutic dosing dependent on bacterial evolution, we proposed to use two *S. aureus* bacterial

strains, with and without inherent resistance to the aminoglycoside gentamicin. These preclinical

models were designed to monitor bacterial dynamics and bacterial resistance evolution to

gentamicin and to understand the molecular events within biofilms contributing to resistance and

persistence. We hypothesized that we could capture a 'therapeutic window', providing a guideline

for local dosing to prevent or treat PJI for 'low-risk' and 'high-risk' infections clinically. A

secondary goal was to compare the bacterial dynamics and resistance evolution in vitro and in vivo

in preclinical models to enhance the translational value of antibacterial testing of antibiotic-eluting

polymeric materials.

RESULTS

MSSA and MRSA demonstrate distinct colonization patterns in in-vivo and in-vitro models.

The biofilm localization and growth dynamics of MSSA and MRSA were determined on

both the implant material and the peri-implant tissue from the subcutaneous and joint infection

models. In the subcutaneous model, the viability of implant-adherent bacteria recovered was

consistently 10^3 CFU/mL for 21 days. There were≥2log more viable bacteria recovered from peri-

implant tissue samples, with the viable load being highest at POD 1 and 3 (10^8 CFU/mL), which

was subsequently reduced by POD 21 (10^5 CFU/mL) (Figure 1A). SEM observations confirmed

poor bacterial presence/viability on the surface of the implanted plates. MRSA demonstrated more

bacterial aggregates and biofilm matrix structures on SS plates when compared to MSSA (Figure

2A).

High MSSA and MRSA bacterial viability ($>5\times10^5$ CFU/mL) was observed in the joint infection

model on both screw implant and peri-implant tissue (Figure 1B). The MRSA on the screw implant

showed lower viability (~2 log) on POD 3 compared to MSSA but the bacterial load was similar

for both strains at POD 7. The bacterial viability of MSSA recovered from the peri-implant femoral

and tibial tissues did not show any differences. However, the viable MRSA recovered from the

peri-implant (tibia) tissue was consistent over time ($\sim10^7$ CFU/mL) when compared to that of peri-

implant femoral tissue which steadily decreased over the period of 7 days (from $>5 \times10^7$ to 1×10^6

CFU/mL). SEM confirmed increased bacterial adherence to the screw implant surface. Significant

adhesion and biofilm formation comprising varying cell morphologies and dense matrix

components were observed for MSSA on the implanted surface (Figure 2B).

In the in-vitro model, the viable bacteria recovered from MSSA-adhered SS plates and screws

increased from $\sim10^5$ to $> 10^6$ CFU in 24 hours (Figure 1C). SEM observations showed significant

bacterial attachment on implant materials (SS plates and screws) which correlated with the

bacterial viability data (Figure 2B).

Figure 1. Biofilm growth dynamics of MSSA and MRSA across in vivo and in vitro infection models (A) MSSA and MRSA count from stainless-steel implant and peri-implant tissue harvested at POD 1, 3, 7, and 21. (B) MSSA and MRSA count from stainless-steel screw implant and peri-implant femoral and tibial tissue harvested at POD 1, 3, and 7. (C) Adherent MSSA and MRSA bacteria count from in-vitro stainless-steel plate and screw culture at 6 and 24 hours. Bacterial viability data from tissue samples were calculated by normalizing to the respective weight of tissue retrieved. Error bars represent the standard deviation (n=2 for implant, n=4 for tissue culture in (A), n=1 for screw implant, n=3 for tissue culture in (B) and n=3 in (C)). n indicates the number of samples.

Figure 2: *S. aureus* **adhesion on implant materials retrieved from in vivo and in vitro models.** Scanning electron micrographs of A) MSSA and MRSA adhered to implanted stainless-steel materials retrieved on POD3 from in vivo infection models and B) Established (24 hours) biofilms of MSSA and MRSA adhered to stainless-steel materials in vitro. Scale bar = 10, 2, and 1μm for panel A and 10 and 2 μm for panel B respectively. Representative images of n=2 implants from subcutaneous and joint infection model; n=3 for in vitro model

Biofilm formation and dynamics influence bacterial susceptibility to gentamicin.

The susceptibility of tissue-colonized and implant-adherent MSSA and MRSA to the antibiotic gentamicin was determined longitudinally. In the subcutaneous model, the implant-adhered MRSA demonstrated antibiotic resistance as early as POD 1 (200 µg/mL). At POD 3 and until POD 21, the MBEC was found to be above the maximum concentration of gentamicin tested in this study (>500 µg/mL). Implant-adhered MSSA acquired resistance towards gentamicin by POD 7 (100 µg/mL), which remained the same until POD 21 (Figure 3A, Table 1). The tissue-colonized MRSA and MSSA exhibited the highest resistance (>500 and 200 µg/mL, respectively) on POD 1 and 3, which was decreased by POD 21 (200 µg/mL and no growth, respectively) (Figure 3A, Table 1).

In the joint infection model, the implant-adhered MRSA demonstrated very high antibiotic resistance as early as POD 1 (>500 µg/mL) and stayed highly resistant until POD 21. In contrast, the implant adhered MSSA demonstrated a comparatively lower level of acquired resistance (≥10 µg/mL) at POD 1, 3, and 7 (Figure 3B). For the tissue colonized MSSA, the acquired gentamicin resistance was moderate and consistent over 7 days of infection (50 µg/mL). The tissue colonized MRSA showed the highest resistance on POD 1 and 3 and a subsequent reduction of MBEC to 300 µg/mL was observed by POD 7 (Table 2).

For the in-vitro SS plate model, both implant-adhered MSSA and MRSA demonstrated increased resistance due to biofilm formation within 6 hours (100 and >500 µg/mL, respectively). The acquired resistance of MSSA was further increased to >500 µg/mL within 24 hours, whereas MRSA stayed highly resistant throughout the study period (Figure 3C, Table 3). For the in-vitro SS screw model, both 6 hours and 24 hours-grown biofilms of MSSA exposed to gentamicin were eradicated effectively with concentrations close to its planktonic MIC values (5 µg/mL). The 6-

150   hour and 24-hour biofilms of MRSA grown on SS screws exhibited inherent resistance to

151   gentamicin (3log reduction only at 500 µg/mL and no reduction until 500 µg/mL, respectively).

Figure 3: Evolution of gentamicin susceptibility for MSSA and MRSA. Heat maps indicating the %growth frequency observed after 24 hours of indicated gentamicin concentration exposure for A) MSSA and MRSA adhered to the subcutaneously implanted stainless steel material retrieved on POD1,3,7and 21 (POD1,3,7 n=2; POD21 n=3) B) MSSA and MRSA colonizing peri-implant tissue retrieved on POD1,3,7 and 21 (POD1,3,7 n=4; POD21 n=2) C) Nascent (6 hours) and established (24 hours) biofilms of MSSA and MRSA adhered to stainless steel material in vitro (n=3). The susceptibility profiles of gentamicin for implant and tissue samples across the different models are presented in Tables 1-3.

Table 1: Subcutaneous infection model

Infection	Source	MBEC gentamicin (µg/mL)			
		POD1	POD3	POD7	POD21
Low-risk MSSA	Implant	1	100	200	200
	Tissue	50	50	100	No growth
High-risk MRSA	Implant	200	500	>500	>500
	Tissue	>500	>500	>500	200

Table 2: Joint infection model

Infection	Source	MBEC gentamicin (µg/mL)		
		POD1	POD3	POD7
Low-risk MSSA	Implant	>10	>10	10
	Tissue	50	50	50
High-risk MRSA	Implant	>500	>500	>500
	Tissue	>500	>500	300

Table 3: in vitro model

Infection	Source	MBECgentamicin	
		Nascent (6hr)	Established (24hr)
Low-risk MSSA	Implant	100	>500
	Screw	5	5
High-risk MRSA	Implant	>500	>500
	Screw	500	>500

Molecular responses of bacterial biofilms to their environment are strain-dependent.

The gene expression of MSSA and MRSA colonizing the implant materials (in vivo and in vitro) and the peri-implant tissue were performed to reveal their molecular status. In the subcutaneous model, the *vraR* gene expression for tissue colonized MSSA was elevated ($>1.5\log_{10}$) for the length of the study. For MRSA, the *vraR* expression was not altered until POD 7, when the expression was deregulated but the expression was significantly increased on POD 21 ($>1.5\log_{10}$). The *icaA* gene expression of MSSA was found to be largely unaltered until POD 21 where it was somewhat increased ($\sim1\log_{10}$). For MRSA, the *icaA* gene was upregulated significantly at POD 7 ($>1\log_{10}$) and remained elevated on POD 21. The *icaD* gene expression in both MSSA and MRSA was largely unaltered over the entire study period ($< 0.5\log_{10}$). The *ebpS* expression demonstrated a consistent increase for both MSSA and MRSA from POD 1 to POD 21 ($>0.5\log_{10}$) (Figure 4A).

In the joint infection model, the *vraR* expression for MSSA was found to be significantly upregulated ($>2\log_{10}$) and the expression levels were highest on POD7 ($>3.5\log_{10}$). For MRSA, *vraR* expression was slightly altered on POD 3 and 7 ($>0.5\log_{10}$). The *icaA* and *icaD* genes for MSSA were significantly upregulated on POD 3 and 7 ($>1\log_{10}$). For MRSA, the *icaA* expression was somewhat upregulated ($\sim0.5\log_{10}$) and the *icaD* expression was significantly upregulated until POD 7 ($>1\log_{10}$). The *ebpS* gene expression was significantly increased for MSSA with the highest expression demonstrated on POD 3 and 7 ($>2\log_{10}$). For MRSA, the *ebpS* gene expression was increased ($\sim1\log_{10}$) with overall subdued gene expression in comparison to MSSA (Figure 4B).

Figure 4: Regulation of stress responses, adhesion, and biofilm formation in tissue colonizing MSSA and MRSA in vivo. Gene expression analysis of A) MSSA and MRSA colonizing the peri-implant tissue retrieved on POD1,3,7 and 21 from rats subjected to subcutaneous infection and POD1,3,7 from rats subjected to joint infection. B) MSSA and MRSA biofilms adhered to the implant materials retrieved on POD 3 from in vivo models and from in vitro models at time points indicated. The relative gene expression was further normalized to the expression profile of planktonic *S. aureus* suspension. Error bars represent standard deviation (Subcutaneous model; n=3 (POD 1,3,7) n=10 and 11 for POD21 MSSA and MRSA respectively, Joint model; n=3 (POD1,3,7) *Indicates p value <0.05

In the in-vitro model, the implant-adhered MSSA demonstrated downregulation of *vraR* and *ebpS*

genes (-1.5 log$_{10}$) in nascent biofilms (6 hours) and the expression remained unaltered in

established biofilms (24 hours). For MRSA, both genes were significantly downregulated in

nascent biofilms and the expression remained the same in established biofilms (>-1.5 log$_{10}$). The

icaA and *icaD* expression was found to be drastically deregulated in both MSSA and MRSA

harvested from nascent and established biofilms (>-3 log$_{10}$) (Figure 5). In the in-vivo models

(subcutaneous and joint infection) the biofilm RNA extracted from implant materials demonstrated

strain-dependent expression of 16srRNA indicating the presence of bacteria on the surfaces. The

expression profile for the remaining genes was not observed. (Table 5, Supplementary File 1).

Figure 5: Regulation of stress responses, adhesion, and biofilm formation in implant colonizing MSSA and MRSA in vitro. Gene expression analysis of MSSA and MRSA biofilms adhered to the implant materials retrieved from the in vitro stainless-steel plates at time points indicated. Gene expression was normalized to 16srRNA expression. The gene expression was normalized to the planktonic *S. aureus* expression profiles. Error bars represent standard deviation (n=3). * Indicates p-value <0.05

Discussion

Periprosthetic joint infection (PJI) remains a crucial clinical challenge that impacts implant longevity in patients, and results in high morbidity and mortality[37,38]. The persisting populations of bacterial biofilms colonizing the implant surface and surrounding tissue increase the severity of PJI and elevate the risk of recurring infections[39]. A comprehensive understanding of the biofilm establishment timeline, the evolution of drug resistance mechanisms, and the emergence of heterogenous phenotypes and genotypes using suitable in vivo and in vitro PJI models are key to devising effective eradication strategies that can curb bacterial resistance and persistence[18,40,41]. In this study, we investigated and characterized the pathogenesis and biofilm dynamics of a low-risk and high-risk infection in PJI. We aimed to longitudinally evaluate the risk of infection by employing in-vivo and in-vitro models. Our goal in vivo was to characterize the bacteria more extensively by incorporating both longitudinal analysis as well as the determination of resistance evolution. We aimed to elucidate the optimum therapy and therapeutic window to maximize the eradication of the bacteria based on the severity of PJI, simulated by using two strains with varying resistance. Our goal in vitro was to understand the relevance of the bacterial dynamics to in-vivo bacterial behavior and to determine methods to increase this relevance.

Bacterial contamination of the prosthetic components often occurs during their implantation. Despite the wide acceptance of the seminal concept of 'race for the surface' between the bacteria and host, recent findings indicate that the severity of the resulting infection largely depends on the bacterial strain, bacterial load, and the environment[42–47]. In our study, the ATCC 12600 strain was selected to simulate an infection that is susceptible to commonly used antibiotics (low risk) in the clinic. L1101, a clinically isolated Mu50 strain with known resistance

to penicillin, aminoglycosides, and vancomycin, was used to simulate a 'high-risk' infection. To

establish sustainable infections in the rat model in vivo, a high bacterial inoculum (10^8 CFU) was

used[48], and an in-vitro implant infection (10^5 CFU) was established to characterize bacterial

dynamics on surfaces. In the subcutaneous model, preferential colonization of the peri-implant

tissue was observed over the entire study period compared to consistently poor colonization of the

subcutaneous plate implant despite seeding with high bacterial density. Under SEM, the biofilms

of MSSA were sparse compared to MRSA, which validates the advantage of inherent resistance

in evading host immune responses[49]. On the other hand, in the joint infection model, almost

equally high colonization was observed for MSSA and MRSA on the screw implants as well as

the surrounding tibial and femoral tissues until POD 7. These observations indicate a strong

influence of the internal environment within the infected site in driving the biofilm growth

dynamics and in sustaining microbial viability[50]. Moreover, the microbial viability on

contaminated implanted screws was ~3 logs higher than the subcutaneous implant, which was

corroborated by the SEM observations. The increased SA colonization on screw implants further

underlined the flexibility of the facultative anaerobe SA in thriving in hypoxic

environments[51,52]. The biofilm growth dynamics observed on in-vitro screw and plate models

were largely comparable to their respective counterparts in the in-vivo models despite a lower

starting inoculum. The vital role of different factors such as the implant location, site of bacterial

colonization (implant vs tissue), the phenotypic and genotypic differences between bacteria, and

the presence of vascularization in influencing biofilm dynamics in vivo is emphasized. The

observations also facilitated the comparison between in-vivo and in-vitro infection and

underscored the need for more diverse models to capture the biofilm growth dynamics and strain-

specific infection outcomes.

Timely determination of the identity and antibiotic susceptibility profile of contaminating

organisms can be crucial in making clinical dosing decisions to combat PJI[53–55]. Usually, the

minimum inhibitory concentration (MIC) of infecting organisms directs the empirical antibiotic

treatment regimens[13,56]. This approach, although widely accepted, relies on the assumption that

the infecting organisms exist in a 'planktonic state.' In reality, microbes form complex biofilms

during pathogenesis, wherein their physiology, drug susceptibility, and response to the host

environment are continuously evolving, resulting in phenotypic and genotypic

heterogeneity[22,57,58]. Our study highlights the profound impact of biofilm formation on the

antibiotic susceptibility of MSSA and MRSA in periprosthetic joint infection. In the subcutaneous

infection model, the tissue colonizing MSSA rapidly acquired gentamicin resistance as early as

POD 1 ($50 \times$ MIC) and by POD7 required almost $100 \times$ MIC to observe >3log reduction. This

provided a critical insight regarding the minimal antibiotic dose for bacterial eradication, even for

a low-risk susceptible strain of infecting bacteria. In comparison, the inherently resistant MRSA

expectedly remained highly resistant to gentamicin (>$500 \times$ MIC) until POD 7. Notably, the

decrease in the bacterial viability and an increase in the susceptibility of the tissue-colonizing

MRSA on POD21 strongly suggests that the host response can be effective against a high-risk

organism and the timeline of the infection is a vital component in infection characterization. For

the implant-adhered bacteria, the MSSA adhered to the implant, gradually acquired resistance, and

maintained viability over time, whereas the MRSA maintained its resistance profile throughout the

study period. This behavior contrasts with the bacteria colonizing the tissue where there is more

variation in growth dynamics and susceptibility profiles of the bacterial population, presumably

due to an increased interaction with the host.

In the joint infection model, MSSA's susceptibility to gentamicin evolved to $>10 \times$ MIC (implant-

adhered) and $50 \times$ MIC (tissue-colonized) within 7 days of infection which was comparable to

previously published findings[59]. The high-risk MRSA showed consistently high resistance to

the antibiotic over the entire study period. Compared to the subcutaneous infection model, the

antibiotic susceptibility profile of MSSA in the joint infection model was not drastically elevated,

which further emphasized the model-based longitudinal differences in infection risk stratification.

In contrast to the MSSA populations in vivo, implant adhered MSSA acquired rapid and higher

resistance to gentamicin within 24 hours ($500 \times$ MIC) in vitro. The data from plate-adhering

MSSA and MRSA were comparable to the bacterial profiles observed until POD 3 and POD 7,

respectively, in the subcutaneous infection model despite the absence of host factors in the in-vitro

model. The in vitro infection study simulated more advantageous conditions for the bacteria in an

implant-associated infection without host immunity. Based on the antibiotic susceptibility data

from the three different infection models, a high initial dosing of antibiotics ($>100 \times$ MIC) is

required during the early stages of infection that should be sustained for a prolonged period

irrespective of the susceptibility profiles of the bacterial strain. Antibiotic-loaded implant materials

that have been developed to locally release high antibiotic concentrations in a sustained manner

could be best suited for this application[60–62]. For inherently resistant infections, it is advisable

to use stronger tools such as the synergistic use of antibiotic and non-antibiotic compounds to

enhance the antibacterial activity as early as possible to achieve effective eradication[36,63,64].

The study highlights the importance of diagnostics for characterizing the infecting bacteria *in situ*

and developing more nuanced situation-specific and environment-specific guidelines for

antibacterial treatment.

Current PJI diagnosis is largely limited in the identification and determination of antibiotic

susceptibility of the causative organism[34]. The prophylactic and treatment models rely on the

physiological attributes that have been determined using data largely from in-vitro studies[12].

The molecular status of bacterial populations within biofilms undergoes strain and maturity-

dependent changes in their stress responses, cell wall constituents, metabolism, slime, and adhesin

production, resistance to antibiotics, and immune responses[20,65,66]. Besides, the bacterial

populations differ in their physiology depending on the site of colonization[67]. This adds a layer

of complexity in the prevention and treatment approaches required. Taken together, the 'status' of

the infecting bacterial strains (with inherent and acquired antibiotic resistance) and the site of

colonization could directly correlate with the risk and severity of the infection[31,68]. Thus, there

is a lot of uncertainty in predicting the treatment outcomes in specific cases of infections. To

address this knowledge gap, the molecular signatures of the MSSA and MRSA bacterial

populations colonizing implants in vivo, and in vitro were characterized. In the subcutaneous

model, the stress response associated with maintaining bacterial cell wall integrity was triggered

early on for tissue colonized MSSA (POD1) when compared to MRSA (POD21) supporting the

idea that there is a lack of alternative strategies in low-risk strain and the multiplexed mechanisms

present in high-risk strain for evading host-mediated targeting[69–71]. In contrast, both strains

showed consistent upregulation of adhesin expression over time which revealed that bacterial

populations actively increased adhesion on the tissue. The biofilm formation was more pronounced

in MRSA compared to MSSA which is associated with high biofilm-forming properties attributed

to resistant strains of *S. aureus*[72]. This finding significantly correlated to the bacteria counts in

the tissue and to the SEM observations of MRSA on SS plates. For implant-adhered bacteria, due

to the limited RNA availability, we were able to capture only bacterial *16srRNA* expression levels

which were supportive of strain-specific biofilm viability from bacterial numbers and SEM

images.

In the joint infection model, the bacterial stress response to the in-vivo environment was also

found to be strain-dependent with the upregulation of the *vraR* gene for both MSSA and MRSA

within 7 days. Strikingly, the adhesin gene expression was only consistently upregulated for MSSA

and not for MRSA, which indicated the possibility of an alternative site-dependent mechanism in

MRSA mediating tissue colonization[73]. In contrast to the subcutaneous infection model, the

biofilm-associated *ica* genes were all upregulated throughout the study period with strain-specific

longitudinal differences. The biofilm gene expression data validated the bacterial viability and

SEM observations of the SS screw implant indicating the unique site-specific advantage

facilitating increased biofilm survival and immune cell evasion[30,74]. Similar to the

subcutaneous infection model, due to the limitation of biofilm RNA from the screw implants, only

the presence of *16srRNA* could be validated. For the in-vitro infection model, we were able to pool

multiple implant materials colonized by bacteria to assess their molecular status. Due to the

absence of any in-vivo or environmental factors, the implant material-adhered bacterial expression

of *vraR* and *ebpS* was comparable to the planktonic bacteria. The gene expression levels of biofilm-

associated genes were significantly downregulated compared to planktonic bacteria indicating the

limitation of in-vitro models without the integrated response to the immune system to capture and

simulate the biofilm-associated changes in bacteria colonizing the implant[67,75].

Our in-vivo studies were limited by the number of implants. Despite implanting 6 implants per

animal in the subcutaneous model, the determination of MBEC and gene expression of the bacteria

on the implants were still limited by the low bacteria count. In addition, we only had one implant

per animal to work within the joint infection model, further limiting our analysis. Our in-vitro

studies were also limited to growth in one type of medium, which is likely to be a strong

determinant of bacterial dynamics and evolution of resistance[76,77].

Conclusion

The role of inherent and biofilm maturity-associated antibiotic resistance, and site-specific

resistance profiles of *S. aureus* in determining the risk and severity progression of PJI was captured

using three different infection models. This could aid in determining a suitable 'therapeutic

window' for clinical dosing guidelines when encountering a low-resistance or high-resistance

infection. Our study has also provided crucial insights into evaluating the translational value of

in-vivo and in-vitro PJI models. The in-vivo infection studies using two different models revealed

vital information on the strain-specific bacterial colonization patterns, evolution of resistance, and

physiological differences that are governed by the site of infection. Immune response markers

correlating with the infection status could serve as an additional resource to determine the effective

concentrations of antibiotic required. Our results suggest implantation at the desired site is required

for relevant efficacy testing of anti-infective materials. Even though in-vitro implant infection

models are important and practical for studying biofilm dynamics on materials, the absence of in-

vivo factors limits their translational value.

EXPERIMENTAL SECTION

1. Bacteria preparation

Gentamicin-susceptible *S. aureus* ATCC 12600 (MSSA) and Gentamicin-resistant *S. aureus*

L1101 (MRSA) were used in this study (Table 1, supplementary file 1). The bacterial glycerol

stocks at -80°C were grown in tryptic soy agar plates (TSA) for 18-24 hours at 35°C to achieve

optimum growth. *S. aureus* colonies were inoculated in tryptic soy broth (TSB) and cultured

overnight to obtain 10^9 CFU/mL. The bacterial broth cultures were subjected to 10,000 × g

centrifugation and pellets were resuspended in sterile PBS. The bacterial suspension in PBS was

further diluted to 10^8 CFU in sterile PBS and 10^5 CFU/mL in sterile TSB before all animal

infection experiments and in vitro experiments, respectively.

2. **Animal study**

2.1 Ethics statement: The animal study design and protocols were approved by the Institutional

Care and Use Committee of Massachusetts General Hospital (2021N000127).

2.2 Subcutaneous infection model Male Sprague-Dawley rats (Charles River, Wilmington, MA)

weighing 350-400g were randomly divided into two groups. 316L stainless steel plates (10 × 3 ×

1mm; n=6 in each animal) were subcutaneously implanted on each rat dorsum and 10^8 CFU of

gentamicin-sensitive MSSA (n=34 rats), gentamicin-resistant MRSA (n=33) were inoculated into

each of the 6 subcutaneous pockets. Non-infected group (n=14) served as a control for the study.

All rats were given facility chow and water ad libitum. Rats were anesthetized with 1-3%

isoflurane in 1 L of O2/min, and 0.05 mg/kg IP buprenorphine was administered 30 minutes before

surgery. All groups were sacrificed on postoperative days (POD) 1, 3, 7, and 21 (Table 2,

Supplementary File 1)

2.3 Joint infection model:

Male Sprague Dawley Rats were randomly assigned to 'low-risk' gentamicin-sensitive MSSA,

'high-risk' gentamicin-resistant MRSA infection groups (n=3/day) and control group (n=1/day).

10^8 CFU of bacteria were inoculated into the intercondylar canal drilled in the tibia after which, a

stainless-steel screw (1.3 mm diameter, 8 mm in length) was implanted into the contaminated

canal. All rats were given facility chow and water ad libitum. Rats were anesthetized with 1-3%

isoflurane in 1 L of O2/min, and 0.05 mg/kg IP buprenorphine was administered 30 minutes before

surgery. The animals were sacrificed on POD 1, 3, and 7 (Table 3, supplementary file 1).

3. Ex-vivo and in-vitro determination of bacterial colonization

The stainless-steel plates and screws (SS) were retrieved from the animals at specific time points

(POD 1, 3, 7, 21, and POD 1,3,7 respectively) and the explants were washed using sterile 1×

phosphate-buffered saline (PBS) to remove any non-adherent bacteria and debris. The explants

were then transferred to 1.5 mL tubes and subjected to sonication for 40 minutes in 1 mL PBS to

dislodge the adherent bacteria. The sonicate was then diluted and plated on tryptic soy agar plates

and incubated for 18-24 hours at 35°C. The adherent bacteria count was determined the following

day using the colony counting method. The peri-implant tissue surrounding the implant was

harvested at the same time points and homogenized using a Tissue Disruptor (TissueRuptor,

Qiagen). The homogenate was diluted and plated on tryptic soy agar plates and incubated for 18-

24 hours at 35°C. The tissue-colonized bacteria were determined the following day using the

colony count method.

To determine in-vitro biofilm dynamics, staphylococcal bacterial suspension [10^5 CFU/mL] in 1

mL of Luria-Bertani (LB) broth was inoculated on 316 Stainless Steel (SS) plates [$10 \times 3 \times 1$ mm]

or screws placed within 24-well plates. The materials were statically incubated for an indicated

period (6 and 24 hours) at 35°C. At each time point, the spent media was removed, and the

materials were washed thrice using sterile 1× PBS. The surfaces were transferred to 1.5 mL tubes

and subjected to sonication for 40 minutes in 1 mL PBS to dislodge the adherent bacteria, The

sonicate was then plated on tryptic soy agar plates and incubated for 18-24 hours at 35°C. The

adherent bacteria count was determined the following day by the colony counting method.

4. Ex-vivo and in-vitro determination of minimum biofilm eradication concentration

The stainless-steel plates and screws (SS) were retrieved from the animals at specific time points (POD 1, 3, 7, 21 and POD 1, 3, 7 respectively) and the plates were washed using sterile 1× PBS to remove any non-adherent bacteria and debris. The explants were then exposed to a range of gentamicin concentrations in 10% LB [1, 10, 50, 100, 200, and 500 µg/mL for SS plates; 10 (MSSA) and 300 µg/mL (MRSA) for screws)] for 24 hours at 35°C. The explants were then washed using sterile 1× PBS and subjected to sonication for 40 minutes in 1 mL PBS to dislodge the adherent bacteria. The sonicate was then diluted and plated on tryptic soy agar plates and incubated for 18-24 hours at 35°C. The adherent bacteria count was determined the following day using the colony counting method. The retrieved peri-implant tissue was rinsed once with sterile PBS and exposed to a range of gentamicin concentrations in 10% LB [1, 10, 50, 100, 200, and 500 µg/mL (subcutaneous infection study); 10, 50, 100, 300 and 500 µg/mL (joint-infection study)] for 24 hours at 35°C. The tissues were washed with sterile PBS and were sonicated for 40 minutes and plated. The bacterial viability was determined the following day using the colony count method.

To determine MBEC of in vitro-formed biofilms, staphylococcal biofilms were grown for a period of 6 and 24 hours on plates and screws as previously described (section 3). The spent media was removed at each timepoint respectively and the materials were washed thrice with PBS to remove all non-adherent bacteria. The surfaces were then placed in a fresh 24-well plate containing a range of gentamicin concentrations [0.5, 1, 5, 10, 20, 40, 60, 80, 100, 200, 300, 400, 500 µg/mL (SS plates); 1, 5, 10, 50, 100, 300, 500 (SS screws)]. Further to drug exposure for 24 hours at 35°C, the surfaces were gently rinsed thrice using PBS and were transferred to 1.5ml tubes containing 1ml PBS. The materials underwent sonication for 40 minutes and the adherent bacteria count was

determined using the spread plate method. MBEC was determined as the concentration that

achieved >3log10 reduction in adherent bacteria count.

The growth frequency observed from each replicate for each concentration tested was calculated

as %growth frequency and heatmaps were generated using the Complex Heatmaps package in R

studio.

5. Gene expression analysis

The peri-implant tissue retrieved from each time point was subjected to a modified Qiagen RNeasy

extraction protocol to extract bacteria RNA. Briefly, 15-30mg tissue samples were homogenized

and subjected to lysis using RLT buffer. The lysate was transferred to a tube and was further

subjected to enzymatic and mechanical lysis using lysostaphin (200 µg/mL), proteinase K, and

acid-washed beads. RNA extraction was performed using the RNeasy spin column method

according to the manufacturer's instructions. To determine in vitro and in vivo adherent bacteria

gene expression the bacteria were harvested from SS plates (n=10 for in vitro model; n=6 from in

vivo model) and screws (n=4 from in vivo model) for each condition by subjecting the materials

to 40 min sonication. The sonicate fluid was pooled and pelleted by centrifuging at 10,000 ×g for

10 mins. The pellet was then subjected to mechanical and enzymatic lysis and the total RNA was

extracted using RNeasy Power Biofilm RNA extraction kit for gram-positive bacteria. The samples

were subjected to real-time quantitative PCR for *icaA*, *icaD*, *ebpS*, and *vraR* genes for *S. aureus*

using specific primers listed (Table 4, supplementary file 1). The Cq values were normalized to

S. aureus 16srRNA expression. Gene expression analysis of tissue-colonizing and implant-adhered

bacteria relative to planktonic bacterial expression was performed using the $2^{(-\Delta\Delta Ct)}$ method.

6. Scanning Electron Microscopy

Scanning electron microscopy was performed on the implant materials retrieved from rats

on POD3 and from in-vitro experiments[36]. The implant materials with adherent bacteria were

fixed using 2.5% glutaraldehyde in 0.1M PBS for 48 hours. The plates were then washed twice for

10 mins with PBS. The adherent bacteria were then treated with osmium tetroxide (OsO^4) 2% +

Ruthenium red 0.2% 1:1 solution for a period of 1hr. The samples were washed twice thoroughly

with distilled water for 10 mins. Further to this the samples were treated with 1% Tannic acid for

30 mins and then washed twice with distilled water for 10 mins each. The prepared samples were

imaged at 15-20 kV, high vacuum (Zeiss FESEM Ultra Plus).

7. Statistical analysis

The gene expression studies were performed in triplicates and the dataset was analyzed using

Student's T-Test (paired). The p-value was calculated and the lowest significant score of 0.05 was

considered statistically significant.

Author contributions

AS, EO, and YF conceptualized and conducted the experimental design for the study. AS, PT,

MM, PJ, FM, and DK performed the data acquisition and SEM visualization. YF, JY, and SL

performed the animal surgery protocols. KKW and NI performed SEM visualization. AS analyzed

and interpreted the data. AS, EO, and OKM wrote, reviewed, and edited the manuscript. All

authors read and approved the final manuscript.

Acknowledgments

This study was funded by National Institutes of Health Grant R01AR077023. The funder played no role in the study design, data collection, analysis, and interpretation of data, or the writing of this manuscript. The authors thank Jean Yuh and Sashank Lekkala for their technical assistance during animal surgery and Dr. Kerry Laplante at the University of Rhode Island for providing the clinical MRSA strain. This work was performed in part at the Harvard University Center for Nanoscale Systems (CNS); a member of the National Nanotechnology Coordinated Infrastructure Network (NNCI), which is supported by the National Science Foundation under NSF award no. 651 ECCS-2025158.

Competing interests

O.K.M. declares the following disclosures: Royalties - Corin, Mako, Iconacy, Renovis, Arthrex, ConforMIS, Meril Healthcare, Exactech, Cambridge Polymer Group; Stake/Equity-Cambridge Polymer Group, Orthopedic Technology Group, Alchimist. E.O. declares the disclosures: Royalties-Corin, Iconacy, Renovis, Arthrex, ConforMIS, Meril Healthcare, Exactech; Paid consultant – WL Gore & Assoc; Editorial Board – JBMR; Officer/Committee- SFB, ISTA. None of these are in direct conflict with the study.

Data availability

All data generated or analyzed during this study are included in this published article [and its supplementary information files]

References

[1]   Tande AJ, Gomez-Urena EO, Berbari EF, Osmon DR. Management of Prosthetic Joint Infection. Infect Dis Clin North Am 2017;31:237–52. https://doi.org/10.1016/j.idc.2017.01.009.

[2]   Kapadia BH, Berg RA, Daley JA, Fritz J, Bhave A, Mont MA. Periprosthetic joint infection. The Lancet 2016;387:386–94. https://doi.org/10.1016/S0140-6736(14)61798-0.

[3]   Kurtz SM, Lau EC, Son MS, Chang ET, Zimmerli W, Parvizi J. Are We Winning or Losing the Battle With Periprosthetic Joint Infection: Trends in Periprosthetic Joint Infection and Mortality Risk for the Medicare Population. J Arthroplasty 2018;33:3238–45. https://doi.org/10.1016/J.ARTH.2018.05.042.

[4]   Labek G, Thaler M, Janda W, Agreiter M, Stöckl B. Revision rates after total joint replacement. J Bone Joint Surg Br 2011;93-B:293–7. https://doi.org/10.1302/0301-620X.93B3.25467.

[5]   Parvizi J, Tan TL, Goswami K, Higuera C, Della Valle C, Chen AF, et al. The 2018 Definition of Periprosthetic Hip and Knee Infection: An Evidence-Based and Validated Criteria. J Arthroplasty 2018;33:1309-1314.e2. https://doi.org/10.1016/j.arth.2018.02.078.

[6]   Le Vavasseur B, Zeller V. Antibiotic Therapy for Prosthetic Joint Infections: An Overview. Antibiotics 2022;11:486. https://doi.org/10.3390/antibiotics11040486.

[7]   Kandel CE, Jenkinson R, Daneman N, Backstein D, Hansen BE, Muller MP, et al. Predictors of Treatment Failure for Hip and Knee Prosthetic Joint Infections in the Setting of 1- and 2-Stage Exchange Arthroplasty: A Multicenter Retrospective Cohort. Open Forum Infect Dis 2019;6. https://doi.org/10.1093/ofid/ofz452.

[8]   Steadman W, Chapman PR, Schuetz M, Schmutz B, Trampuz A, Tetsworth K. Local Antibiotic Delivery Options in Prosthetic Joint Infection. Antibiotics (Basel) 2023;12. https://doi.org/10.3390/antibiotics12040752.

[9]   Winkler C, Dennison J, Wooldridge A, Larumbe E, Caroom C, Jenkins M, et al. Do local antibiotics reduce periprosthetic joint infections? A retrospective review of 744 cases. J Clin Orthop Trauma 2018;9:S34–9. https://doi.org/10.1016/j.jcot.2017.08.007.

[10]   Bourget-Murray J, Azad M, Gofton W, Abdelbary H, Garceau S, Grammatopoulos G. Is the routine use of local antibiotics in the management of periprosthetic joint infections justified? HIP International 2023;33:4–16. https://doi.org/10.1177/11207000221139467.

[11]   Olearo F, Zanichelli V, Exarchakou A, Both A, Uçkay I, Aepfelbacher M, et al. The Impact of Antimicrobial Therapy Duration in the Treatment of Prosthetic Joint Infections Depending on Surgical Strategies: A Systematic Review and Meta-analysis. Open Forum Infect Dis 2023;10. https://doi.org/10.1093/ofid/ofad246.

[12]   Rottier W, Seidelman J, Wouthuyzen-Bakker M. Antimicrobial treatment of patients with a periprosthetic joint infection: basic principles. Arthroplasty 2023;5:10. https://doi.org/10.1186/s42836-023-00169-4.

[13]   Tillander JAN, Rilby K, Svensson Malchau K, Skovbjerg S, Lindberg E, Rolfson O, et al. Treatment of periprosthetic joint infections guided by minimum biofilm eradication concentration

(MBEC) in addition to minimum inhibitory concentration (MIC): protocol for a prospective randomised clinical trial. BMJ Open 2022;12:e058168. https://doi.org/10.1136/bmjopen-2021-058168.

[14]   Ricciardi BF, Porter KR, Myers TG, Ginnetti JG, Kaplan N, Thirukumaran CP. Demographics and Early Outcomes of Commercial Antibiotic Cement Usage for Infection Prophylaxis During Primary Total Knee Arthroplasty in Patients Older Than 65 Years: An American Joint Replacement Registry Study. Journal of the American Academy of Orthopaedic Surgeons 2024;32:59–67. https://doi.org/10.5435/JAAOS-D-23-00434.

[15]   Scheper H, Gerritsen LM, Pijls BG, Van Asten SA, Visser LG, De Boer MGJ. Outcome of Debridement, Antibiotics, and Implant Retention for Staphylococcal Hip and Knee Prosthetic Joint Infections, Focused on Rifampicin Use: A Systematic Review and Meta-Analysis. Open Forum Infect Dis 2021;8. https://doi.org/10.1093/ofid/ofab298.

[16]   Peng H-M, Zhou Z-K, Wang F, Yan S-G, Xu P, Shang X-F, et al. Microbiology of Periprosthetic Hip and Knee Infections in Surgically Revised Cases from 34 Centers in Mainland China. Infect Drug Resist 2021;14:2411–8. https://doi.org/10.2147/IDR.S305205.

[17]   Kreve S, Reis AC Dos. Bacterial adhesion to biomaterials: What regulates this attachment? A review. Jpn Dent Sci Rev 2021;57:85–96. https://doi.org/10.1016/j.jdsr.2021.05.003.

[18]   Peng Q, Tang X, Dong W, Sun N, Yuan W. A Review of Biofilm Formation of Staphylococcus aureus and Its Regulation Mechanism. Antibiotics 2022;12:12. https://doi.org/10.3390/antibiotics12010012.

[19]   Ricciardi BF, Muthukrishnan G, Masters E, Ninomiya M, Lee CC, Schwarz EM. Staphylococcus aureus Evasion of Host Immunity in the Setting of Prosthetic Joint Infection: Biofilm and Beyond. Curr Rev Musculoskelet Med 2018;11:389–400. https://doi.org/10.1007/s12178-018-9501-4.

[20]   Uruén C, Chopo-Escuin G, Tommassen J, Mainar-Jaime RC, Arenas J. Biofilms as Promoters of Bacterial Antibiotic Resistance and Tolerance. Antibiotics 2020;10:3. https://doi.org/10.3390/antibiotics10010003.

[21]   Tran NN, Morrisette T, Jorgensen SCJ, Orench-Benvenutti JM, Kebriaei R. Current therapies and challenges for the treatment of *Staphylococcus aureus* biofilm-related infections. Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy 2023;43:816–32. https://doi.org/10.1002/phar.2806.

[22]   Karimaei S, Kazem Aghamir SM, Foroushani AR, Pourmand MR. Antibiotic tolerance in biofilm persister cells of Staphylococcus aureus and expression of toxin-antitoxin system genes. Microb Pathog 2021;159:105126. https://doi.org/10.1016/j.micpath.2021.105126.

[23]   Waters EM, Rowe SE, O'Gara JP, Conlon BP. Convergence of Staphylococcus aureus Persister and Biofilm Research: Can Biofilms Be Defined as Communities of Adherent Persister Cells? PLoS Pathog 2016;12:e1006012. https://doi.org/10.1371/journal.ppat.1006012.

[24]   García-Betancur JC, Lopez D. Cell Heterogeneity in Staphylococcal Communities. J Mol Biol 2019;431:4699–711. https://doi.org/10.1016/j.jmb.2019.06.011.

[25]   Klein S, Morath B, Weitz D, Schweizer PA, Sähr A, Heeg K, et al. Comparative Genomic Reveals Clonal Heterogeneity in Persistent Staphylococcus aureus Infection. Front Cell Infect Microbiol 2022;12. https://doi.org/10.3389/fcimb.2022.817841.

[26]   Zhu MF, Kim K, Cavadino A, Coleman B, Munro JT, Young SW. Success Rates of Debridement, Antibiotics, and Implant Retention in 230 Infected Total Knee Arthroplasties: Implications for Classification of Periprosthetic Joint Infection. J Arthroplasty 2021;36:305-310.e1. https://doi.org/10.1016/j.arth.2020.07.081.

[27]   Xu Y, Wang L, Xu W. Risk factors affect success rate of debridement, antibiotics and implant retention (DAIR) in periprosthetic joint infection. Arthroplasty 2020;2:37. https://doi.org/10.1186/s42836-020-00056-2.

[28]   Gonzalez MR, Pretell-Mazzini J, Lozano-Calderon SA. Risk Factors and Management of Prosthetic Joint Infections in Megaprostheses—A Review of the Literature. Antibiotics 2023;13:25. https://doi.org/10.3390/antibiotics13010025.

[29]   Staats A, Li D, Sullivan AC, Stoodley P. Biofilm formation in periprosthetic joint infections. Ann Jt 2021;6:43–43. https://doi.org/10.21037/aoj-20-85.

[30]   Doub JB, Parmiter D, Brantner CA, Moshyedi M, Hughes M, Kolevar M, et al. The Location of Biofilms on Chronic Prosthetic Joint Infections and the Ramifications for Clinical Practice. Arthroplast Today 2024;25:101314. https://doi.org/10.1016/j.artd.2023.101314.

[31]   Svensson Malchau K, Tillander J, Zaborowska M, Hoffman M, Lasa I, Thomsen P, et al. Biofilm properties in relation to treatment outcome in patients with first-time periprosthetic hip or knee joint infection. J Orthop Translat 2021;30:31–40. https://doi.org/10.1016/j.jot.2021.05.008.

[32]   Macias-Valcayo A, Aguilera-Correa J-J, Broncano A, Parron R, Auñon A, Garcia-Cañete J, et al. Comparative *In Vitro* Study of Biofilm Formation and Antimicrobial Susceptibility in Gram-Negative Bacilli Isolated from Prosthetic Joint Infections. Microbiol Spectr 2022;10. https://doi.org/10.1128/spectrum.00851-22.

[33]   Mandell JB, Orr S, Koch J, Nourie B, Ma D, Bonar DD, et al. Large variations in clinical antibiotic activity against *Staphylococcus aureus* biofilms of periprosthetic joint infection isolates. Journal of Orthopaedic Research 2019;37:1604–9. https://doi.org/10.1002/jor.24291.

[34]   Tande AJ, Patel R. Prosthetic Joint Infection. Clin Microbiol Rev 2014;27:302. https://doi.org/10.1128/CMR.00111-13.

[35]   Pannu TS, Villa JM, Higuera CA. Diagnosis and management of infected arthroplasty. SICOT J 2021;7:54. https://doi.org/10.1051/sicotj/2021054.

[36]   Sekar A, Gil D, Tierney P, McCanne M, Daesety V, Trendafilova D, et al. Synergistic use of anti-inflammatory ketorolac and gentamicin to target staphylococcal biofilms. J Transl Med 2024;22:102. https://doi.org/10.1186/s12967-024-04871-y.

[37]   Kurtz SM, Lau EC, Son M-S, Chang ET, Zimmerli W, Parvizi J. Are We Winning or Losing the Battle With Periprosthetic Joint Infection: Trends in Periprosthetic Joint Infection and Mortality Risk for the Medicare Population. J Arthroplasty 2018;33:3238–45. https://doi.org/10.1016/j.arth.2018.05.042.

[38]   Lentino JR. Prosthetic Joint Infections: Bane of Orthopedists, Challenge for Infectious Disease Specialists. Clinical Infectious Diseases 2003;36:1157–61. https://doi.org/10.1086/374554.

[39]   Visperas A, Santana D, Klika AK, Higuera-Rueda CA, Piuzzi NS. Current treatments for biofilm-associated periprosthetic joint infection and new potential strategies. Journal of Orthopaedic Research 2022;40:1477–91. https://doi.org/10.1002/jor.25345.

[40]   Moormeier DE, Bayles KW. *Staphylococcus aureus* biofilm: a complex developmental organism. Mol Microbiol 2017;104:365–76. https://doi.org/10.1111/mmi.13634.

[41]   Irwin S, Mackenzie BW, Matthews BG, Williams DL, Cornish J, Swift S. Improve Integration of In Vitro Biofilm Body of Knowledge to Support Clinical Breakthroughs in Surgical Site Infection. JAAOS: Global Research and Reviews 2021;5. https://doi.org/10.5435/JAAOSGlobal-D-20-00217.

[42]   Shiels S, Mangum L, Wenke J. Revisiting the "race for the surface" in a pre-clinical model of implant infection. Eur Cell Mater 2020;39:77–95. https://doi.org/10.22203/eCM.v039a05.

[43]   Gristina AG, Naylor P, Myrvik Q. Infections from biomaterials and implants: a race for the surface. Med Prog Technol n.d.;14:205–24.

[44]   Ramírez-Granillo A, Bautista-Hernández LA, Bautista-De Lucío VM, Magaña-Guerrero FS, Domínguez-López A, Córdova-Alcántara IM, et al. Microbial Warfare on Three Fronts: Mixed Biofilm of Aspergillus fumigatus and Staphylococcus aureus on Primary Cultures of Human Limbo-Corneal Fibroblasts. Front Cell Infect Microbiol 2021;11. https://doi.org/10.3389/fcimb.2021.646054.

[45]   Martinez-Perez M, Perez-Jorge C, Lozano D, Portal-Nuñez S, Perez-Tanoira R, Conde A, et al. Evaluation of bacterial adherence of clinical isolates of Staphylococcus sp. using a competitive model: An in vitro approach to the "race for the surface" theory. Bone Joint Res 2017;6:315–22. https://doi.org/10.1302/2046-3758.65.BJR-2016-0226.R2.

[46]   Hickok NJ, Li B, Oral E, Zaat SAJ, Armbruster DA, Atkins GJ, et al. The 2023 Orthopedic Research Society's international consensus meeting on musculoskeletal infection: Summary from the in vitro section. Journal of Orthopaedic Research 2024;42:512–7. https://doi.org/10.1002/jor.25774.

[47]   Vidlak D, Kielian T. Infectious Dose Dictates the Host Response during Staphylococcus aureus Orthopedic-Implant Biofilm Infection. Infect Immun 2016;84:1957–65. https://doi.org/10.1128/IAI.00117-16.

[48]   Shiels SM, Bedigrew KM, Wenke JC. Development of a hematogenous implant-related infection in a rat model. BMC Musculoskelet Disord 2015;16:255. https://doi.org/10.1186/s12891-015-0699-7.

[49]   Wimmer MD, Hischebeth GTR, Randau TM, Gathen M, Schildberg FA, Fröschen FS, et al. Difficult-to-treat pathogens significantly reduce infection resolution in periprosthetic joint infections. Diagn Microbiol Infect Dis 2020;98:115114. https://doi.org/10.1016/J.DIAGMICROBIO.2020.115114.

[50] McConoughey SJ, Howlin R, Granger JF, Manring MM, Calhoun JH, Shirtliff M, et al. Biofilms in periprosthetic orthopedic infections. Future Microbiol 2014;9:987–1007. https://doi.org/10.2217/fmb.14.64.

[51] Wilde AD, Snyder DJ, Putnam NE, Valentino MD, Hammer ND, Lonergan ZR, et al. Bacterial Hypoxic Responses Revealed as Critical Determinants of the Host-Pathogen Outcome by TnSeq Analysis of Staphylococcus aureus Invasive Infection. PLoS Pathog 2015;11:e1005341. https://doi.org/10.1371/journal.ppat.1005341.

[52] Balasubramanian D, Harper L, Shopsin B, Torres VJ. *Staphylococcus aureus* pathogenesis in diverse host environments. Pathog Dis 2017:ftx005. https://doi.org/10.1093/femspd/ftx005.

[53] Kim S-J, Cho YJ. Current Guideline for Diagnosis of Periprosthetic Joint Infection: A Review Article. Hip Pelvis 2021;33:11–7. https://doi.org/10.5371/hp.2021.33.1.11.

[54] Lee J, Park H, Bae J, Hyun H, Kim S. Current Diagnostic Methods for Periprosthetic Joint Infection. Biomedical Science Letters 2022;28:1–8. https://doi.org/10.15616/BSL.2022.28.1.1.

[55] Tripathi S, Tarabichi S, Parvizi J, Rajgopal A. Current relevance of biomarkers in diagnosis of periprosthetic joint infection: an update. Arthroplasty 2023;5:41. https://doi.org/10.1186/s42836-023-00192-5.

[56] Mueller M, de la Peña A, Derendorf H. Issues in Pharmacokinetics and Pharmacodynamics of Anti-Infective Agents: Kill Curves versus MIC. Antimicrob Agents Chemother 2004;48:369–77. https://doi.org/10.1128/AAC.48.2.369-377.2004.

[57] Hall-Stoodley L, Costerton JW, Stoodley P. Bacterial biofilms: from the Natural environment to infectious diseases. Nat Rev Microbiol 2004;2:95–108. https://doi.org/10.1038/nrmicro821.

[58] Stewart PS, Franklin MJ. Physiological heterogeneity in biofilms. Nat Rev Microbiol 2008;6:199–210. https://doi.org/10.1038/nrmicro1838.

[59] Okae Y, Nishitani K, Sakamoto A, Kawai T, Tomizawa T, Saito M, et al. Estimation of Minimum Biofilm Eradication Concentration (MBEC) on In Vivo Biofilm on Orthopedic Implants in a Rodent Femoral Infection Model. Front Cell Infect Microbiol 2022;12. https://doi.org/10.3389/fcimb.2022.896978.

[60] Lekkala S, Inverardi N, Yuh J, Wannomae KK, Tierney P, Sekar A, et al. Antibiotic-Loaded Ultrahigh Molecular Weight Polyethylenes. Macromol Biosci 2024;24. https://doi.org/10.1002/mabi.202300389.

[61] Suhardi VJ, Bichara DA, Kwok SJJ, Freiberg AA, Rubash H, Malchau H, et al. A Fully Functional Drug-Eluting Joint Implant. Nat Biomed Eng 2017;1. https://doi.org/10.1038/S41551-017-0080.

[62] Gimeno M, Pinczowski P, Pérez M, Giorello A, Martínez MÁ, Santamaría J, et al. A controlled antibiotic release system to prevent orthopedic-implant associated infections: An in vitro study. European Journal of Pharmaceutics and Biopharmaceutics 2015;96:264–71. https://doi.org/10.1016/j.ejpb.2015.08.007.

[63] Gil D, Daffinee K, Friedman R, Bhushan B, Muratoglu OK, LaPlante K, et al. Synergistic antibacterial effects of analgesics and antibiotics against Staphylococcus aureus. Diagn Microbiol Infect Dis 2020;96:114967. https://doi.org/10.1016/j.diagmicrobio.2019.114967.

[64] Kamble E, Sanghvi P, Pardesi K. Synergistic effect of antibiotic combinations on Staphylococcus aureus biofilms and their persister cell populations. Biofilm 2022;4:100068. https://doi.org/10.1016/j.bioflm.2022.100068.

[65] Coenye T, Bové M, Bjarnsholt T. Biofilm antimicrobial susceptibility through an experimental evolutionary lens. NPJ Biofilms Microbiomes 2022;8:82. https://doi.org/10.1038/s41522-022-00346-4.

[66] Paharik AE, Horswill AR. The Staphylococcal Biofilm: Adhesins, Regulation, and Host Response. Microbiol Spectr 2016;4. https://doi.org/10.1128/microbiolspec.VMBF-0022-2015.

[67] Masters EA, Ricciardi BF, Bentley KL de M, Moriarty TF, Schwarz EM, Muthukrishnan G. Skeletal infections: microbial pathogenesis, immunity and clinical management. Nat Rev Microbiol 2022;20:385–400. https://doi.org/10.1038/s41579-022-00686-0.

[68] Lamret F, Varin-Simon J, Velard F, Terryn C, Mongaret C, Colin M, et al. Staphylococcus aureus Strain-Dependent Biofilm Formation in Bone-Like Environment. Front Microbiol 2021;12. https://doi.org/10.3389/fmicb.2021.714994.

[69] Gerlach D, Guo Y, De Castro C, Kim S-H, Schlatterer K, Xu F-F, et al. Methicillin-resistant Staphylococcus aureus alters cell wall glycosylation to evade immunity. Nature 2018;563:705–9. https://doi.org/10.1038/s41586-018-0730-x.

[70] Hernández-Cuellar E, Tsuchiya K, Valle-Ríos R, Medina-Contreras O. Differences in Biofilm Formation by Methicillin-Resistant and Methicillin-Susceptible Staphylococcus aureus Strains. Diseases 2023;11:160. https://doi.org/10.3390/diseases11040160.

[71] Gao C, Dai Y, Chang W, Fang C, Wang Z, Ma X. VraSR has an important role in immune evasion of Staphylococcus aureus with low level vancomycin resistance. Microbes Infect 2019;21:361–7. https://doi.org/10.1016/j.micinf.2019.04.003.

[72] Jean-Pierre V, Boudet A, Sorlin P, Menetrey Q, Chiron R, Lavigne J-P, et al. Biofilm Formation by Staphylococcus aureus in the Specific Context of Cystic Fibrosis. Int J Mol Sci 2022;24:597. https://doi.org/10.3390/ijms24010597.

[73] Gordon RJ, Lowy FD. Pathogenesis of Methicillin-Resistant *Staphylococcus aureus* Infection. Clinical Infectious Diseases 2008;46:S350–9. https://doi.org/10.1086/533591.

[74] Knabl L, Kuppelwieser B, Mayr A, Posch W, Lackner M, Coraça-Huber D, et al. High percentage of microbial colonization of osteosynthesis material in clinically unremarkable patients. Microbiologyopen 2019;8. https://doi.org/10.1002/mbo3.658.

[75] Lebeaux D, Chauhan A, Rendueles O, Beloin C. From in vitro to in vivo Models of Bacterial Biofilm-Related Infections. Pathogens 2013;2:288–356. https://doi.org/10.3390/pathogens2020288.

[76] Paleczny J, Brożyna M, Dudek-Wicher R, Dydak K, Oleksy-Wawrzyniak M, Madziała M, et al. The Medium Composition Impacts Staphylococcus aureus Biofilm Formation and Susceptibility to Antibiotics Applied in the Treatment of Bone Infections. Int J Mol Sci 2022;23:11564. https://doi.org/10.3390/ijms231911564.

[77] Chen X, Thomsen TR, Winkler H, Xu Y. Influence of biofilm growth age, media, antibiotic concentration and exposure time on Staphylococcus aureus and Pseudomonas aeruginosa biofilm removal in vitro. BMC Microbiol 2020;20:264. https://doi.org/10.1186/s12866-020-01947-9.