A noncanonical polyamine from bacteria antagonizes animal mitochondrial function

Kelsie M. Nauta[1], Darrick R. Gates[1], Matthew Weiland[1], Marco E. Mechan-Llontop[1], Xiao Wang[1], Kim P. Nguyen[1], Christine Isaguirre[2], Megan R. Genjdar[2], Ryan D. Sheldon[2], Connie M. Krawczyk[1], Nicholas O. Burton[1]*

[1]Van Andel Research Institute, Department of Metabolism and Nutritional Programing, Grand Rapids, MI, 49503, USA

[2]Van Andel Research Institute, Mass Spectrometry Core, Grand Rapids, MI, 49503, USA

*Corresponding author: Nick.Burton@vai.org

Keywords

C. elegans, microbiome, insulin, polyamines, N^1-Aminopropylagmatine, *Bacillus subtilis*, mitochondria, *speB, daf-2, hsp-6, catp-5*

Abstract

Canonical polyamines such as agmatine, putrescine, and spermidine are evolutionarily conserved metabolites found in nearly all forms of life ranging from bacteria to humans. Recently, interactions between polyamines produced by gut bacteria and human intestinal cells have been proposed to contribute to both Irritable Bowel Syndrome with Diarrhea (IBS-D) and inflammatory bowel diseases. However, the molecular mechanisms that underlie these effects are often unclear due in part to limitations in the methods used to manipulate and study polyamine functions *in vivo.* Here, we developed a *Caenorhabditis elegans* based screening platform and a modified LC-MS approach for profiling polyamine metabolites. We combined these methods to make the unexpected discovery that dysfunctional polyamine metabolism in both Gram-negative (*E. coli*) and Gram-positive (*B. subtilis*) bacteria can result in the accumulation of a noncanonical polyamine intermediate, N^1-Aminopropylagmatine (N^1-APA). We further find that N^1-APA is produced via spermidine synthase (SpeE) and that it is bioactive when encountered by animals. Specifically, we find that when N^1-APA is produced by bacteria in animal intestines it can be transported into intestinal cells via the polyamine transporter CATP-5 where it antagonizes both animal development and mitochondrial function across diverse animal species. Lastly, we find that N^1-APA functions analogously to the deoxyhypusine synthase inhibitor GC7. For example, like GC7, N^1-APA antagonizes eIF5A hypusination and inhibits the alternative activation of mammalian macrophages. To our knowledge, these findings are the first to demonstrate that N^1-APA is a bioactive metabolite and that bacteria can produce a small molecule that functions similarly to existing deoxyhypusine synthase inhibitors. Furthermore, these results suggest an exciting new mechanistic hypothesis for why the loss of *speB* in gut microbes, including *E. coli,* has been both linked to inflammatory bowel disease (IBD) in humans and found to drive IBD in germ free mice.

Introduction

Canonical polyamine metabolites including agmatine, putrescine, spermidine, and spermine are

found in nearly all domains of life and are generally present at millimolar levels in animal cells

[1,2]. Changes in the metabolism or abundance of polyamines have been linked to numerous

human pathologies. For example, mutations in the polyamine transporter ATP13A2 cause a

heritable form of juvenile onset Parkinson's disease [3–5]. Similarly, changes in the production of

polyamines by gut microbiota have recently been found to both cause irritable bowel syndrome

(IBS-D) and, in other contexts, have been proposed to protect against the development of

inflammatory bowel diseases and colitis [6,7]. Despite their clear importance to biology and

demonstrable relevance to numerous human pathologies, the molecular mechanisms by which

altered polyamine metabolism causes changes in animal physiology are often unclear. For

example, polyamines have been proposed to function as signaling molecules [8–10] or antioxidants

[11,12], to stabilize RNA, protein, and membranes [13,14], and to regulate mRNA translation via the

hypusination of the transcription factor, eIF5A[15]. However, in many cases the exact molecular

mechanism by which changes in polyamine metabolism cause animal disease remain unknown.

In addition, there are several cases of seemingly conflicting models for the mechanisms by

which polyamines regulate animal physiology. For example, using the polyamine analog

inhibitor of deoxyhypusine synthase, GC7, Puleston et al[16] and Nakamura et al.[6] have proposed

that polyamines regulate alternative activation of macrophages (M2-like) through regulating the

hypusination of eIF5A which in turn regulates the translation of a subset of mitochondrial

proteins. By contrast, using a genetic approach, Anderson-Baucum et al.[17] reported no clear

evidence that hypusination of eIF5A regulated mitochondrial protein abundance by proteomics

and suggested that previous findings might be due to alternate target effects of GC7. Instead,

they found that eIF5A hypusination levels decreased the proinflammatory nature of

macrophages by decreasing translation of proinflammatory mRNAs[17], Thus, the precise

mechanisms by which polyamines regulate aspects of animal physiology such as immune cell

activation remain unclear.

Studies describing how the disruption of polyamine metabolism causes or contributes to disease

in animals face multiple challenges. First, manipulating polyamine levels *in vivo* is challenging

because the loss of enzymes required to synthesize putrescine and spermidine are lethal in

many animals. In animal intestines, polyamines can also be transported between intestinal cells

and gut microbes, complicating the interpretation of animal studies due to potential

compensatory effects [2,18]. Despite these challenges, recent studies have found that when

agmatinase (encoded by *speB*) is deleted from bacterial genomes that inhabit animal intestines,

mice develop colitis and *C. elegans* undergo developmental arrest [6,7]. SpeB converts agmatine

to putrescine in the canonically polyamine synthesis pathway[19], and both excess agmatine and

loss of putrescine have been proposed as the underlying molecular mechanism to explain these

changes to animal physiology; however, neither has been shown to be sufficient to cause these

effects [6,7].

The ability to measure polyamine levels *in vivo* also faces technical difficulties. For example,

most kit-based approaches only quantify total polyamines [20]. To overcome this, mass

spectrometry-based approaches have been employed. However, polyamines are poorly

detected in standard metabolomics-based approaches [21]. Additionally, while mass spectrometry

methods have been developed to reliably quantify the core polyamine metabolites (agmatine,

putrescine, and spermidine), most studies are unable to simultaneously identify the many

variants of these molecules. Thus, studies of polyamines are largely restricted to quantifying

canonical polyamines and might overlook alternative polyamine metabolites. As a consequence

of these limitations, defining cause-and-effect relationships between polyamines and mRNA,

protein, or membrane stability, remains difficult. Thus, the precise molecular mechanisms for

how disruptions in polyamine metabolism lead to human diseases remain unclear.

While screening a complete non-essential *Bacillus subtilis* gene knockout library for bacterial

genes that regulate animal development in *C. elegans*; we discovered that the loss of *speB* in

B. subtilis caused complete animal developmental arrest. Two previous studies found the loss of

speB in *E. coli* results in the failure of *C. elegans* to develop in the presence of excess agmatine

or causes mouse lethality in models of inflammatory bowel disease [6,7]. Unexpectedly, however,

we found that the effects of *speB* mutant *B. subtilis* on animals were not due to a requirement

for bacterial putrescine in animals or due to excess agmatine. Instead, we found that diverse

Gram-negative and Gram-positive bacteria produce a non-canonical polyamine (N^1-

Aminopropylagmatine (N^1-APA)) via the activity of the conserved bacterial enzyme SpeE on

agmatine, corroborating recent biochemical findings [22,23]. In addition, we found that N^1-APA

accumulates in *speB* mutant bacteria and is transported into animal intestinal cells via the

polyamine transporter CATP-5. Once inside animal cells, N^1-APA acts analogously to GC7, a

chemical inhibitor of deoxyhypusine synthase. These findings demonstrate that N^1-APA

functions like GC7 in both invertebrates and vertebrates which has numerous downstream

consequences, such as inhibiting *C. elegans* development and immune cell activation in mouse

bone marrow-derived macrophages (BMM$_\Phi$). To our knowledge, these findings are the first to

report that N^1-APA is a bioactive metabolite in any species and the first observation that a

polyamine metabolite derived from bacteria functions analogously to the deoxyhypusine

synthase inhibitor GC7 *in vivo*. These findings demonstrate the importance of investigating

individual non-canonical polyamines when studying how changes in polyamine metabolism

cause disease. They also provide an alternative N^1-APA dependent mechanism of action that

may explain why recent reports link the loss of *speB* in gut microbiome bacteria to inflammatory

bowel disease in both mice and humans [6,24].

Results

Loss of *speB* in bacteria antagonizes animal development

To identify new links between bacterial metabolism and animal physiology, we screened a

complete *B. subtilis* non-essential gene knockout library for changes in animal development,

using *Caenorhabditis elegans* as a model (Fig. 1A – See methods for screen details) [25]. From

our screen of 3,984 mutant *B. subtilis* isolates, we identified only a single *B. subtilis* gene, *speB*,

that was required for *C. elegans* to develop to adulthood (Fig 1B). SpeB encodes an

agmatinase which plays an intermediate role in the synthesis of spermidine (Fig. 1C) [19,26]. We

confirmed that *speB* is required for animals to develop to adulthood by complementing this

mutant with a wild-type copy of *speB* (Fig. 1B). We conclude that *speB* is required in *B. subtilis*

to promote *C. elegans* development to adulthood.

In many bacteria, SpeB (agmatinase) is part of a pathway, together with SpeA and SpeE

(spermidine synthase), that converts arginine to spermidine[19] (Fig. 1C). Previous studies have

proposed that deletion of *speB* causes either the loss of beneficial downstream polyamines

(putrescine, spermidine) or a toxic buildup of agmatine and that these changes in canonical

polyamines results in deleterious effects on animal physiology[6,7]. To determine if *C. elegans* are

unable to develop to adulthood on *B. subtilis* Δ*speB* because they require spermidine from their

diet, we tested if the deletion of *speA* or *speE* in *B. subtilis* also caused animals to arrest their

development (SFig. 1A). In contrast, we found that deletion of *speA* and *speE* did not alter WT

animal development (Fig. 1D, SFig. 1B). Since *speA* converts L-arginine into agmatine[19], we

generated a double Δ*speA* Δ*speB* deletion strain to test if preventing the aberrant accumulation

of agmatine prevented animal developmental arrest. We found deletion of *speA* in the Δ*speB*

background rescued *C. elegans* development, suggesting agmatine causes development arrest

(Fig. 1D, SFig. 1B). We confirmed this phenotype by complementing Δ*speA* Δ*speB* with *speA*

(Fig. 1D, SFig. 1B). Surprisingly, we found that knocking out the *speEB* operon does not

phenocopy Δ*speB* and that Δ*speEB* mutants behave similarly to Δ*speA* Δ*speB* mutant bacteria

(Fig. 1D, SFig. 1B). We confirmed this result by complementing Δ*speEB* with *speE* and found

that the strain no longer promoted animal development (Fig. 1D, SFig. 1B). We conclude that

the deletion of *speB*, but not other genes in the spermidine synthesis pathway, antagonizes

animal development and that this effect is dependent on both *speA* and *speE* activity. These

findings suggest that the simple loss of spermidine or a buildup of agmatine are not the cause of

Δ*speB-mediated* disruption of animal development.

Loss of *speB* causes diverse species of bacteria to accumulate N^1-APA.

We hypothesized that *B. subtilis speB* mutants might produce a noncanonical polyamine that

antagonizes animal development. This would be in line with recent reports that bacteria produce

numerous noncanonical polyamines, sometimes using the same enzymes predicted to be

involved in canonical polyamine metabolism, such as SpeE [22]. To test this, we developed an

untargeted LC-MS method for the profiling of polyamine metabolites in *B. subtilis* Δ*speA*,

ΔspeB, ΔspeEB, ΔspeE, ΔspeB; ΔspeA, and WT cultures. Polyamines are not well retained or

ionized in typical LC-MS metabolomics approaches [21]. So, to do this we generated carbamyl

derivatives of metabolite extracts and performed untargeted metabolomics. Using this approach,

we detected 2641 blank- and peak quality-filtered LC-MS features. Differential abundance

analysis revealed a consistent feature, 388.29207 m/z at retention time of 3.781min, that was

the most significantly increased compound in *ΔspeB* cells versus the other polyamine synthesis

mutants (Fig 2A). Evaluation of extracted ion chromatograms (EIC) indicated that this

compound was only detectable in the *ΔspeB* cells (Fig 2B). Using accurate mass MS1

measurements, we found three possible chemical formulae within 5ppm (Fig 2C). Of these, only

one contained oxygen atoms, of which there are two per carbamyl derivative ($C_5H_9O_2$; Figure

2C). From this we deduced that the unknown compound contained two carbamyl-derivatized

sites, and subtraction of 2x($C_5H_9O_2$) from the predicted chemical formula of the unknown

($C_{18}H_{37}N_5O_4$) yielded a parent compound formula of $C_8H_{21}N_5$. This is consistent with the

non-canonical polyamine, N^1-APA, that was originally reported to accumulate in *speB* mutant

Thermus thermophilus [22,23] (N^1-APA; (Figure 2D). To confirm this putative identification, we

spiked in a synthetic standard to *ΔspeB* media and observed an increase in the 388.2921 EIC

over unspiked *ΔspeB* controls (Figure 2E). MS2 fragmentation spectra of the endogenous

unknown and the synthetic standard further confirmed the identity of the *ΔspeB* specific

polyamine as N^1-APA (Figure 2F).

Previous studies of *C. elegans* found that the loss of *speB* in *E. coli* also caused animals to

arrest their development, similar to our findings for *B. subtilis*[7]. Similar to studies in *C. elegans*,

studies in germ free mice found that *speB* mutant *E. coli* (but not wild-type *E. coli*) alter mouse

physiology and drive inflammatory bowel disease-like pathologies in mammals. Collectively,

these studies suggest that the loss of *speB* in other species of bacteria also affects animal

physiology in taxa ranging from nematodes to mammals. Despite these consistent findings for

speB mutant *E. coli* affecting many important aspects of animal physiology, the mechanistic

explanation remains unclear with different studies proposing the effects might be due to a

buildup of agmatine or a loss of putrescine. Our findings from *B. subtilis* suggest that in addition

to any potential effects on canonical polyamines (agmatine, putrescine), *speB* mutant *E. coli*

might also accumulate the non-canonical polyamine N^1-APA because it disrupts a proposed

alternative spermidine synthesis pathway (Fig. 2G). To test this, we profiled the abundance of

polyamines, including N^1-APA, in *E. coli* with and without *speB*. As in *B. subtilis*, we found that

E. coli lacking *speB* also accumulated N^1-APA (Fig. 2H). We conclude that *speB* mutant *E. coli*

also accumulate N^1-APA. Given that Gram positive (*B. subtilis*) and Gram negative (*E. coli*)

bacteria diverged >2 billion years ago, our findings suggest that most bacteria may produce N^1-

APA in certain contexts. We note that this hypothesis would also be consistent with a recent

biochemical study that found that 17 out of the 18 tested bacterial spermidine synthases

expressed in *E. coli* BL21 (including *speE* from *B. subtilis*) were able to convert agmatine into

N^1-APA, similar to the original report of N^1-APA production in *T. thermophilus* [22,23].

N^1-APA is a bioactive metabolite that drives mitochondrial dysfunction and

developmental arrest in animals when transported via the polyamine transporter CATP-5

Our previous genetics studies of *B. subtilis* indicated that the effects of *speB* mutant *B. subtilis*

on *C. elegans* development were not due to either the loss of putrescine or the accumulation of

agmatine (Fig. 1). Given our mass spectrometry findings, we hypothesized that the

accumulation of N^1-APA in *speB* mutant bacteria might explain why *C. elegans* arrest their

development in response to *speB* mutant *B. subtilis* (Fig. 2). To test this, we synthesized N^1-APA

and grew *C. elegans* on a diet of WT *B. subtilis* on media containing exogenous N^1-APA. We

found exogenous N^1-APA was sufficient to cause developmental arrest in *C. elegans* (Fig. 3A).

This developmental arrest was not observed when spermidine or agmatine were supplemented

into plates (Fig. 3A, SFig. 2). We also found the addition of exogenous spermidine to the media

is sufficient to suppress the effects of exogenous N^1-APA (Fig. 3B) or *B. subtilis* Δ*speB* (Fig. 3C).

We conclude that N^1-APA production by *B. subtilis* antagonizes animal development, possibly by

disrupting the normal function of spermidine.

To determine how N^1-APA antagonizes animal development, we screened 11 *C. elegans* GFP

stress reporters on a diet of either *B. subtilis* WT or *B. subtilis* Δ*speB* (Table 2). We found that

GFP reporters for *hsp-6* and *irg-1* were both substantially increased when animals were fed

Δ*speB* mutant *B. subtilis* when compared to WT *B. subtilis* (Fig. 3D) or compared to other

polyamine synthesis mutants (SFig. 3). *hsp-6* expression is part of the mitochondrial unfolded

protein response and is activated in response to mitochondrial stress. *irg-1* was originally used

as a reporter for bacterial infections, but was later found to be activated in response to

mitochondrial stress[27]. Thus, the activation of both *hsp-6* and *irg-1* is consistent with animals

experiencing mitochondrial stress. This activation is reversed when a wild-type copy of *speB*

was restored to the *B. subtilis* genome (SFig. 3). Importantly, we did not observe a difference in

level of fluorescence for the other nine reporters that were tested: *nlp-29, clec-60, gcs-1, hsp-*

60, hsp-16.2, gst-4, hsp-4, sod-3, bvIs5 (Table 2). The activation of the mitochondrial unfolded

protein response is likely not due to general proteostasis stress because we did not observe

activation of *hsp-4::*GFP which is activated as part of the endoplasmic reticulum unfolded

protein response (SFig.4, Table 2). We conclude that N^1-APA activates the *C. elegans*

mitochondrial unfolded protein response, likely by causing mitochondrial stress.

To determine how a diet of *B. subtilis* Δ*speB* antagonizes animal development, we mutagenized

wild-type *C. elegans* and screened for mutant animals that could develop to adulthood when

feeding on *B. subtilis* Δ*speB.* We isolated 10 lines of mutant animals that developed to

adulthood even when fed *B. subtilis* Δ*speB* (Fig. 3E, 3F, SFig. 5, Table 1). We found the mutant

lines harbored predicted loss-of-function mutations in the *catp-5* polyamine transporter (Fig.

3G). CATP-5 is expressed in intestinal cells and was previously reported to import polyamines

from the intestinal lumen into the intestinal cells [28]. These findings are consistent with a model in

which N^1-APA is produced by *B. subtilis* Δ*speB* and then is imported by CATP-5 into animal

intestinal cells where it antagonizes animal development.

N^1-APA functions analogously to the deoxyhypusine synthase inhibitor GC7 in both

invertebrates and vertebrates

The effects of N^1-APA could be specific to *C. elegans* or could generally antagonize animal

development and mitochondrial function across species. To test this, we exposed the nematode

Pristionchus pacificus to wild-type and *B. subtilis* Δ*speB* (Fig.4A)*. P. pacificus* and *C. elegans*

shared a last common ancestor approximately 100 million years ago [29,30]. We found that *B.*

subtilis Δ*speB* also antagonized the development of *P. pacificus* and led to a >90% reduction

animal fertility (Fig. 4B). We conclude that N^1-Aminiopropylagmatine broadly affects nematode

development and physiology and that this effect is not specific to *C. elegans.*

In the polyamine-eIF5A-hypusine axis, which is conserved in most animals from nematodes to

humans, spermidine is used to hypusinate eIF5A and alter the translation of a subset of

mRNAs. One of the key enzymes in this pathway is deoxyhypusine synthase which is also

conserved from *C. elegans* to humans [31]. Previous chemical inhibition studies of deoxyhypusine

synthase found that GC7, a synthetic polyamine, was a potent inhibitor of deoxyhypusine

synthase[16,32]. Furthermore, this study tested many derivatives of GC7 and found that N^1-APA

could also inhibit deoxyhypusine synthase, but that it was less potent than GC7 [32]. At the time of

this work, it was not known that bacteria synthesize N^1-APA. Given that deoxyhypusine

synthase is evolutionarily conserved, we hypothesized that N^1-APA might cause both

mitochondrial stress and developmental arrest in *C. elegans* by functioning analogously to GC7.

To test this, we first exposed wild-type animals expressing the mitochondrial stress reporter *hsp-*

6::GFP to either N^1-APA or GC7 and found that *hsp-6::GFP* expression is activated by both

molecules (Fig. 4C). We also found that both GC7 and N^1-APA caused animals to arrest their

development (Fig. 4D). Importantly, neither spermidine nor norspermidine activated *hsp-6::GFP*

expression or caused developmental arrest at the relevant concentrations, suggesting these

results are not broadly caused by excess exogenous polyamines (Fig. 4C, 4D, SFig. 6). We

conclude that GC7 and N^1-APA cause similar activation of a mitochondrial stress reporter and

developmental arrest in *C. elegans.*

In mammals, polyamines are important for the regulation of macrophage activation. Activation of

macrophages by IL-4 leads to an anti-inflammatory phenotype of macrophages (M2-like) that

can be inhibited by GC7 and promoted by putrescine [6,16,17]. Therefore, we tested whether N^1-

APA also altered the activation of mammalian macrophages in response to IL-4. Consistent with

N^1-APA being analogous to GC7 and previous studies[16], we found N^1-APA inhibits IL-4 induced

expression of M2-like macrophage markers RELM-α and CD301 (Fig. 4E, 4F, 4G). While GC7

was more effective at inhibiting M2 phenotype than N^1-APA, N^1-APA treatment decreased

hypusination of eIF5A to a greater extent than GC7 (Fig. 4H). These findings demonstrate that

N^1-APA inhibits the hypusination of eIF5A *in vivo* are consistent with a model in which N^1-APA

interferes with a normal function of spermidine in the polyamine-eIF5A-hypusine axis[16].

Discussion

In summary, our findings establish that N^1-APA, produced via an alternative spermidine

synthesis pathway in bacteria, can influence animal development and eIF5a hypusination when

it is taken up by CATP-5 in animal intestinal cells (Fig. 4I). To our knowledge, these findings are

the first to demonstrate that the bacterial polyamine intermediate, N^1-APA, is bioactive in

animals or that bacteria produce an inhibitor that functions similarly to the deoxyhypusine

synthase inhibitor, GC7. Thus, we establish a novel link between a bacterial intermediate

metabolite and the activity of a specific molecular pathway in animals. Our findings also further

establish *C. elegans* as an excellent model species for identifying links between specific

bacterial metabolites and target enzymes in animals.

Importantly, our findings suggest that the role of N^1-APA in biology might be far more profound

that previously appreciated. Specifically, our findings indicate that N^1-APA not only is produced

by both Gram-positive and Gram-negative species of bacteria, but that it can be produced at

significant enough quantitates to alter animal development. Recently, Li *et al.,* 2024 reported the

surprising finding that 17 out of 18 bacterial spermidine synthases, including *speE* from *B.*

subtilis, can produce N^1-APA when expressed in *E. coli*. These unexpected findings led the

authors to conclude that "the N^1-Aminopropylagmatine pathway for spermidine biosynthesis,

which bypasses putrescine, may be far more widespread [in bacteria] than realized and may be

the default pathway for spermidine biosynthesis in species encoding L-Arginine decarboxylase

for agmatine production". Why diverse species of bacteria might use this alternative non-

canonical polyamine synthesis pathway (Fig. 2G) is unknown. However, our findings reported

here indicate that N^1-APA is a bioactive metabolite, that it inhibits nematode development, and

that inhibits the activation of mammalian macrophages. Given these findings, we speculate that

N^1-APA might also function as an immune modulator in nature and/or function to modify animal

physiology in the contentious ecosystem of soil microbiology. For example, non-canonical

polyamine synthesis may act to protect soil microbes from being consumed by predatory

animals such as nematodes. Future studies exploring how bacterial polyamines affect animal

physiology and immune cell activation will further elucidate this relationship.

Our findings also suggest a potentially new mechanistic model for why previous studies found

that disrupting *speB* in intestinal bacteria affects diverse aspect of animal physiology across

both *C. elegans* and mammals. For example, a recent study of the effects of metformin on

animal lifespans found that bacteria-derived agmatine underlies the effects of metformin on host

metabolism and longevity in animals including *C. elegans*[7]. This was in part based on the finding

that supplementing agmatine to *C. elegans* feeding on *E. coli* Δ*speB* resulted in a similar

developmental arrest as we observed for *C. elegans* feeding on *B. subtilis* Δ*speB*. Because

speB converts agmatine to putrescine in the canonical polyamine synthesis pathway, excess

agmatine was hypothesized to accumulate in *speB* mutants and alter animal physiology. Here,

we found there was not a significant difference in agmatine relative abundance between WT and

E. coli Δ*speB*. However, *E. coli* Δ*speB* did accumulate N^1-APA in the presence of exogenous

agmatine. Thus, our findings suggest a potentially alternative mechanistic model for why the

loss of *speB* alters the effects of metformin on animals[7]; the loss of *speB* in *E. coli* leads to a

buildup of N^1-APA which alters mitochondrial function in animals. In this model, the effects of

metformin, a CI inhibitor, may be altered in animals exposed to *speB* mutant bacteria because

N^1-APA accumulates in these bacteria and disrupts animal mitochondrial function[7].

Similar to how loss of bacterial *speB* affects *C. elegans* physiology, studies in mice and humans

have linked loss of *speB* in the gut microbiome to inflammatory bowel disease in mammals [24].

For example, it was recently found that colonizing germ-free mice with *speB* mutant *E. coli* (as

well as deleting a parallel polyamine pathway) promoted the development of severe colitis in

mice[6]. These effects on inflammatory bowel disease and colitis were proposed to be due to a

lack of putrescine. However, our data suggest N^1-APA might also contribute to disease and

could explain why some of the phenotypes caused by *speB* mutant *E. coli* in mice are similar to

those caused by deoxyhypusine synthase inhibition by GC7. Specifically, our findings suggest

that the reason both *speB* mutant *E. coli* and GC7 were found to drive colitis-like changes in

mice or mouse cells might be because *speB* mutant *E. coli* produce N^1-APA which functions

analogously to GC7 and inhibits the development of anti-inflammatory M2-like macrophages

(i.e. "M2-like" macrophages, characterized by the production of RELMa and CD301 (Fig. 4), are

anti-inflammatory and if N^1-APA inhibits their activation it might promote inflammatory bowel

disease like symptoms in the intestine). Along these same lines, our data potentially explain

why the absence of detectable *speB* in certain species of bacteria was significantly associated

with IBD in humans[24] and that the loss of deoxyhypusine synthase in mice causes IBD-like

pathologies in mice [33]; both of these results could be explained by the accumulation of N^1-APA

in *speB* mutant bacteria which inhibits deoxyhypusine synthase function in animals. If true, our

data suggest that existing inhibitors of the bacterial enzymes SpeA or SpeE could potentially be

useful in the treatment of IBD because they would block N^1-APA production. Future studies will

likely be critical in determining if N^1-APA contributes to IBD development in humans or if

inhibiting N^1-APA production in intestinal bacteria might alleviate IBD symptoms.

Acknowledgements

We would like to acknowledge Craig Ellermeier for sharing *Bacillus* strains and plasmids. This research was supported in part by the Van Andel Institute Mass Spectrometry Core (RRID:SCR_024903; Grand Rapids, MI). We thank D. Chandler for helpful comments on the manuscript text, the *Caenorhabditis* Genetics Stock Center (funded by the NIH National Center for Research Resources) and National BioResource Project (NRBP) for *C. elegans* strains, and WormBase as a repository of *C. elegans* data. This work was supported by funds from the VAI MeNu research grant program and NIH grant DP2DK139569.

Diversity and Inclusion Statement

All individuals involved in this study fulfilled the criteria for authorship required by Nature Portfolio journals and thus have been included as authors. Roles and responsibilities of all collaborators were agreed to in advance of research. All efforts were made to ensure no one involved in this study was hindered from participation based on background, ethnicity, or gender. This research also does not result in any known stigmatization, incrimination, discrimination or personal risk to anyone involved in this study.

Declarations of Interest

The authors declare that they have no competing interests.

Author Contributions

Figure 1: Loss of *speB* in bacteria antagonizes animal development. (A) Graphical

representation of the platform used to screen for mutants of *B. subtilis* that did not support

animal development. Germ-free embryos were arrayed on the library of 3,984 *B. subtilis*

mutants and screened for development beyond the L4 stage. Candidate mutants were retested

on 60 mm plates with ~500 embryos. (B) Representative images of WT *C. elegans* on *E. coli*

OP50, *B. subtilis* 168 (*Bs*)(NOBb513), *Bs ΔspeB* (NOBb514), and *Bs ΔspeB; speB*(+)

(NOBb475). The animals were incubated at 20 °C for 3 days on their respective diets, washed

in M9 buffer, and paralyzed with tetramisole (~10 µg/mL). The scale bar is 1 mm. (C)

Canonically, in *B. subtilis*, spermidine is synthesized from L-Arginine by the consecutive

activities of SpeA, SpeB, and SpeE. (D) Percentage of WT *C. elegans* that developed pasts L4+

401   on diets of *Bs ΔspeE* (NOBb517), *Bs ΔspeEB* (NOBb515), *Bs ΔspeEB; speE* (+)(NOBb501), *Bs*

402   *ΔspeA* (NOBb516), *Bs ΔspeB; ΔspeA* (NOBb518), and *Bs ΔspeA; ΔspeB; speA*(+)

403   (NOBb503).The animals were incubated at 26 °C for 5 days on their respective diets and L4+

404   animals were quantified. Experiments were performed in triplicate with ~500 animals per

405   replicate. Error bars = s.d., **** = $p < 0.0001$, ** = $p < 0.0065$, ns = not significant

Figure 2. Identification and validation of N^1-APA accumulated in *ΔspeB* mutant bacteria.

(A) Volcano plots showing differential abundance analysis of untargeted metabolomics on

carbamylated-derivatives between *ΔspeB* cell lysates and the other strains evaluated. Red dots

are LC-MS features with an FDR adjusted P-value <0.05. Note the feature 388.29207 m/z at

retention time 3.781min is the most significantly elevated compound in each comparison. (B)

Representative extracted ion chromatograms of 388.29207 m/z (+/- 5ppm). (C) Top, chemical

formula predictions for m/z 388.29207 using accurate mass and fine isotopic structure in

Compound Discoverer. Bottom, structure of carbamylated-derivatives indicate two oxygens are

present in each derivatized moiety. (D) Structure of N^1-APA with sites of carbamylated-

derivatives denoted with a blue asterisk. Note the chemical formula of N^1-APA (C8H21N5) with

carbamylated-derivatives (C5H9O2 x2) at the indicated sites match the predicted chemical

formula of the unknown (C18H37N5O4). (E) Overlaid extracted ion chromatograms of $\Delta speB$

and $\Delta speB$+N1APA spike-in provides accurate mass and chromatographic profile confirmation

of N1APA as the unknown feature 388.29207@3.7981min. (F) Data dependent MS2

fragmentation mirror plot of parent ion 388.29207@3.7981min in $\Delta speB$ (top) and neat N^1-APA

standard (bottom) provides further confirmation that the unknown is N^1-APA. (G) Canonical

spermidine synthesis pathway (1.) compared to the alternative pathway (2.) found in *Thermus*

thermophilus, which includes N^1-APA as an intermediate metabolite. (H) Volcano plots showing

differential abundance analysis of targeted polyamine carbamylated-derivatives in cell lysates

between $\Delta speB$ and other evaluated lines. The feature identified as N^1-APA is elevated in *E. coli*

OP50 $\Delta speB$ as compared to WT *E. coli* OP50. Both strains were grown in the presence of 50

mM agmatine.

Figure 3: Mutations in the polyamine transporter, *catp-5*, confer resistance to *B. subtilis* Δ*speB*. (A) Percent of WT *C. elegans* (N2) that developed past L4 when grown on media supplemented with a range of concentrations of N^1-APA and seeded with wild type *B. subtilis* after incubating for 3 days at 20 °C. Control plates were supplemented with 1 mM agmatine and spermidine. Experiments were performed in triplicate with ~200 animals per replicate. Error bars = s.d. **** = $p < 0.0001$. (B) Percent of WT *C. elegans* that develop past L4 on modified NGM (see methods) supplemented with 0.25 mM N^1-APA and a concentration gradient of spermidine (0.125 —1 mM). Experiments were performed in triplicate with ~300 animals per replicate. Error bars = s.d. **** = $p < 0.0001$. (C) Percent of WT *C. elegans* that developed past L4 on modified NGM (see methods) seeded with *B. subtilis* (WT) or *B. subtilis* Δ*speB* (NOBb369) and supplemented with spermidine (0.125 and 1 mM). Experiments were performed in triplicate with ~300 animals per replicate. Error bars = s.d., *** = $p < 0.008$, **** = $p < 0.0001$, ns = not significant. (D) Brightfield (BF) and fluorescent (GFP) images of growth stage matched *irg-1p::GFP* and *hsp-6p::GFP* on a diet of *B. subtilis* (NOBb513) or *B. subtilis* Δ*speB* (NOBb514). The *C. elegans* strains were incubated on the respective *B. subtilis* strains for ~30 hours at 20 °C. Representative images were pseudo-colored using ImageJ v1.54f. Scale bars are 0.5 mm. (E) Representative images of WT *C. elegans* and mutant *catp-5*(*bur50*) on *B. subtilis* 168 (*Bs*)(NOBb513) and *B. subtilis* 168 Δ*speB* (NOBb514). The animals were incubated at 20 °C for 3 days on their respective diets, washed in M9 buffer, and paralyzed with tetramisole. The scale bar is 1 mm. (F) Percent of WT *C. elegans* (N2) and *catp-5*(*bur50-59*) that developed past L4 on *B. subtilis* 168 Δ*speB* (NOBb514) after 5 days at 26 °C. WT *C. elegans* on *B. subtilis* 168(*Bs*)(NOBb513) was included as a control. Experiments were performed in triplicate with ~500 animals per replicate. Error bars = s.d., **** = $p < 0.0001$ (G) Graphical representation of mutations in CATP-5 that restored animal development per allele (*bur50-59*). Red = nonsense mutation. Blue = nonsynonymous mutation. Green = insertion mutation. Purple = splice adaptor mutation. Numbers 50-59 correspond to allele *bur50-bur59*.

Figure 4: N^1-APA functions similarly to the deoxyhypusine synthase inhibitor GC7 in di-

verse organisms. (A) Representative images of *Pristionchus pacificus* on a diet of *E.*

coli OP50, *B. subtilis*, and *B. subtilis* Δ*speB* (NOBb369). Scale bars are 1 mm. (B) Quantifica-

tion of brood size in *P. pacificus* on a diet of *E. coli* OP50, *B. subtilis*, and *B. sub-*

tilis Δ*speB* (NOBb369). Error bars = s.d., **** = p < 0.0001. (C) Fluorescent (GFP) images of

growth stage matched *C. elegans* encoding *hsp-6p::GFP* fed a diet of *B. subtilis* (NOBb513) on

NGM agar plates supplemented with 0.125 mM spermidine, N^1-APA, and GC7. Chemical struc-

tures are shown to the right. The *C. elegans* strain were incubated with *B. subtilis* strain for ~30

hours at 20 °C. Representative images were pseudo-colored using ImageJ v1.54f. Scale bars

are 0.5 mm. (D) Percent of WT *C. elegans* that developed past L4 on *B. subtilis* 168 (NOBb513)

after exposure to 0.125 mM spermidine, norspermidine, N^1-APA, and GC7 for 3 days at 20 °C.

Experiments were performed in triplicate with ~500 animals per replicate. Error bars = s.d., ****

= p < 0.0001. ns = not significant. (E) Representative flow plots, gates of RELMα+ BMM$_\Phi$. Pro-

portions of (F) RELMα and (G) CD301 BMM$_\Phi$ (IL-4) following treatment with 200 µM N^1-APA or

GC7 for 24 hours. Experiment performed in triplicate; representative of 2 independent experi-

ments. Error bars = S.E.M., **** = p < 0.0001, *** = p < 0.009, ** = p < 0.01. (H) N^1-APA inhibits

eIF5a hypusination in BMM$_\Phi$. Representative images of 3 independent experiments. (I) N^1-Ani-

mopropylagmatine accumulates in the *C. elegans* intestinal lumen when the animals are grown

on a diet of *B. subtilis* Δ*speB*. CATP-5 transports N^1-APA into intestinal cells where it inhibits

eIF5a hypusination. N^1-APA antagonizes mitochondrial and immune function.

A

B

Supplementary Figure 1: Loss of B. subtilis *speB* causes *C. elegans* developmental ar-

rest. (A) Canonical polyamine pathway. (B) Representative images corresponding to Fig. 1D.

Scale bars = 1 mm.

Supplementary Figure 2: Exogenous spermidine and agmatine do not cause develop-

mental arrest. Representative images corresponding to Fig. 3A of WT *C. elegans* grown on

NGM seeded with *B. subtilis* WT (0) and supplemented with N^1-APA (1 mM), agmatine (1 mM),

or spermidine (1 mM). Scale bar = 1 mm.

Supplementary Figure 3: N^1-APA-induced developmental arrest is mediated by mitochon-

drial stress. A) Brightfield (BF) and fluorescent (GFP) images of growth stage matched *hsp-*

6p::GFP, on *E. coli* OP50, *B. subtilis* (NOBb513), *Bs ΔspeB* (NOBb514), *Bs ΔspeB; speB*(+)

(NOBb475), (B) *Bs ΔspeE* (NOBb517), *Bs ΔspeEB* (NOBb515), *Bs ΔspeEB; speE*(+)

(NOBb501), *Bs ΔspeA* (NOBb516), *Bs ΔspeA; ΔspeB* (NOBb518), and *Bs ΔspeA; ΔspeB;*

speA(+) (NOBb503). The *C. elegans* strains were incubated on the respective *B. subtilis* strains

for ~30 hours at 20 °C. Representative images were pseudo-colored using ImageJ v1.54f. Scale

bars = 0.5 mm.

hsp-4::GFP

Supplementary Figure 4: *hsp-4* is not activated by N[1]-APA. Brightfield (BF) and fluorescent (GFP) images of growth stage matched *hsp-4p::GFP*, on *E. coli* OP50, *B. subtilis* (WT), and *Bs ΔspeB* (NOBb369). Representative images were pseudo-colored using ImageJ v1.54f. Scale bars = 0.5 mm.

| C. elegans: | WT | WT | bur50 | bur51 |
| B. subtilis: | WT | ΔspeB | ΔspeB | ΔspeB |

| C. elegans: | bur52 | bur53 | bur54 | bur55 |
| B. subtilis: | Δ speB | ΔspeB | ΔspeB | ΔspeB |

| C. elegans: | bur56 | bur57 | bur58 | bur59 |
| B. subtilis: | ΔspeB | ΔspeB | ΔspeB | ΔspeB |

Supplementary Figure 5: mutations in *catp-5* promote *C. elegans* development on a diet of *B. subtilis* Δ*speB*. Representative images of WT and *catp-5* (*bur50-59*) on a diet of *B. subtilis* Δ*speB* (NOBb369) after incubation at 20 °C for 3 days. Scale bars = 1 mm.

Supplemental Figure 6: N¹-APA and GC7 cause dose-dependent developmental arrest and expression of *hsp-6p::GFP*. Fluorescent (GFP) images of *C. elegans* encoding *hsp-6p::GFP* fed a diet of *B. subtilis* 168 on NGM agar plates supplemented with 0.0625-0.5 mM spermidine, norspermidine, N¹-APA, and GC7. The *C. elegans hsp6p::GFP* reporter strain was exposed to the various polyamines for ~30 hours at 20 °C on a diet of *B. subtilis* 168. Representative images were pseudo-colored using ImageJ v1.54f. Scale bars are 1 mm.

Supplementary Figure 7: (A) Flow cytometry gating strategy corresponding to Fig. 4E. (B) Percentage of alive BMM$_\Phi$ after 24 hour treatment with 200 μM N^1-APA (APA) or GC7 corresponding to Fig. 4F and 4G. Error bars = S.E.M., **** = p < 0.0001, *** = p < 0.009, ** = p <0.01. (C) Replicate western blots corresponding to Fig. 4H.

	Position	Refer-ence	Alter-nate	Variant Type	CDS Variant Annotation	Protein Variant Annota-tion	Alternate Allele Depth
bur50	8092275	C	T	missense_variant	c.4267G>A	p.Gly855Asp	103
bur51	8094229	C	T	missense_variant	c.2313G>A	p.Gly400Glu	95
bur52	8095771	C	T	stop_gained	c.770G>A	p.Trp59*	106
bur53	8094031	C	T	missense_variant	c.2510G>A	p.Gly466Asp	112
bur54	8091401	C	T	missense_variant	c.5141G>A	p.Gly1101Arg	65
bur55	8095831	C	T	splice_acceptor_variant	c.118-1G>A	intron	78
bur56	8094107	C	T	missense_variant	c.2435G>A	p.Gly441Arg	103
bur57	8093719	A	T	missense_variant	c.2822T>A	p.Ser553Arg	93
bur58	8092772	G	A	missense_variant	c.3770C>T	p.Thr705Met	98
bur59	8094738	G	A	insertion	c.1059+11C>T	p.Lys453-Iso454 in-sArgCys	73

Table 1: List of mutations in *catp-5* that result in *C. elegans* development on *B. subtilis* Δ*speB*

	nlp-29p	clec-60p	gcs-1p	hsp-60	hsp-16.2p	gst-4p	hsp-4	irg-1p	sod-3p	bvIs5	hsp-6p
Difference between WT and *ΔspeB*	-	-	-	-	-	-	-	**+**	-	-	**+**

Table 2: Summary of *C. elegans* GFP reporter strains on a diet of *B. subtilis 168 ΔspeB*.

Strain	Description	Reference
P. pacificus		
PS312	*P. pristioncus* wild isolate	[34]
C. elegans		
N2	*Caenorhabditis elegans* wild isolate	[34]
DR1572	*C. elegans daf-2(e1368) III*	[34]
NOB159	*catp-5(bur50) G855N*	this study
NOB160	*catp-5(bur51) G400E*	this study
NOB161	*catp-5(bur52) W59**	this study
NOB162	*catp-5(bur53) G466N*	this study
NOB163	*catp-5(bur54) G1101R*	this study
NOB164	*catp-5(bur55) intron variant*	this study
NOB165	*catp-5(bur56) G441R*	this study
NOB166	*catp-5(bur57) S553R*	this study
NOB167	*catp-5(bur58) T705M*	this study
NOB168	*catp-5(bur59) intron variant*	this study
IG274	*frIs7 [nlp-29p::GFP + col-12p::DsRed] IV*	[34]
JIN810	*agIs26 [clec-60p::GFP + myo-2p::mCherry]*	[34]
LD1171	*ldIs3 [gcs-1p::GFP + rol-6(su1006)]*	[34]
SJ4058	*zcIs9 [hsp-60::GFP + lin-15(+)]*	[34]
WE5172	*ajIs1 [rCesC05A9.1::GFP + dpy-5(+)] X*	[34]
CL2070	*dvIs70 [hsp-16.2p::GFP + rol-6(su1006)]*	[34]
CL2166	*dvIs19 [(pAF15)gst-4p::GFP::NLS] III*	[34]
SJ4005	*zcIs4 [hsp-4::GFP] V*	[34]
AU133	*agIs17 [myo-2p::mCherry + irg-1p::GFP] IV*	[34]
CF1553	*muIs84 [(pAD76) sod-3p::GFP + rol-6(su1006)]*	[34]
CY573	*bvIs5 [cyp-35B1p::GFP + gcy-7p::GFP]*	[34]
GL347	*zcIs13 [hsp-6p::GFP + lin-15(+)] V*	[34]
B. subtilis		
BGSCID: 1A1	*B. subtilis 168 trpC2*	[35]

NOBb369	*B. subtilis 168 ΔspeB*	this study
NOBb475	*B. subtilis 168 ΔspeB thrC::P$_{speEB}$-speB (pNOB47)*	this study
NOBb501	*B. subtilis 168 ΔspeEB thrC::P$_{speEB}$-speE (pNOB48)*	this study
NOBb503	*B. subtilis 168 ΔspeB speA::kanR thrC::P$_{speA}$-speA (pNOB49)*	this study
NOBb513	*B. subtilis 168 thrC::pDG1664*	this study
NOBb514	*B. subtilis 168 ΔspeB thrC::pDG1664*	this study
NOBb515	*B. subtilis 168 ΔspeEB thrC::pDG1664*	this study
NOBb516	*B. subtilis 168 speA::kanR thrC::pDG1664*	this study
NOBb517	*B. subtilis 168 speE::kanR thrC::pDG1664*	this study
NOBb518	*B. subtilis 168 ΔspeB speA::kanR thrC::pDG1664*	this study
E. coli		
OP50	*ura-, strR, rnc-, (delta)attB::FRT-lacI-lacUV5p-T7)*	
NOBb750	*ura-, strR, rnc-, (delta)attB::FRT-lacI-lacUV5p-T7) ΔspeB*	This study

Table 3: List of nematode and bacterial strains used in this study.

Plasmid	Relevant features	Parent vector	Restriction enzymes to digest parent vector	PCR primers	PCR template	Reference
pDG1664	*thrC, amp, erm, spec*	N/A	N/A	N/A	N/A	(54)
pMAD	*amp, erm, ori-pE194ts*	N/A	N/A	N/A	N/A	(55)
pNOB41	pMAD Δ*speEB*	pMAD	BglII, EcoRI	KN429-KN432	*B. subtilis* 168	this study
pNOB47	*thrC::P$_{speEB}$-speB amp erm spec*	pDG1664	BamHI, EcoRI	KN475-KN478	*B. subtilis* 168	this study
pNOB48	*thrC::P$_{speEB}$-speE amp erm spec*	pDG1664	BamHI, EcoRI	KN492, KN493	*B. subtilis* 168	this study
pNOB49	*thrC::P$_{speA}$-speA amp erm spec*	pDG1664	BamHI, EcoRI	KN490, KN491	*B. subtilis* 168	this study

Table 4: List of plasmids used in this study

Oligo name	Sequence	Relevant features
KN429	ACACATTAACTAGACAGATCACCGTTGTATAGAC-GTTTAA	Clone ~1 kb upstream *speEB* into pMAD digested with BglII and EcoRI FOR
KN430	GACATCGTCTGGATGTTTACTAAAAAC-GATCTTTTTTTCTCATAATAGT	Clone ~1 kb upstream speEB into pMAD digested with BglII and EcoRI REV
KN431	ACTATTATGAGAAAAAAGATCGTTTTTAG-TAAACATCCAGACGATGTC	Clone ~1 kb downstream *speEB* into pMAD digested with BglII and EcoRI FOR
KN432	CTGCAGAAGCTTCTAGAATTCGGA-GATGTAATATATTTATCAGAGT	Clone ~1 kb downstream speEB into pMAD digested with BglII and EcoRI REV
KN475	AAAACTGCTGCCTTCGGATCAACATCG-CAGAACAGGACGC	Clone 144 bp P_{speEB} promoter into pDG1664 digested with BamHI and EcoRI FOR
KN476	TAAGCTTCATCAAATCTcatTT-GTCATCTCCCTTTCGAGG	Clone 144 bp P_{speEB} promoter into pDG1664 digested with BamHI and EcoRI REV
KN477	CCTCGAAAGGGAGATGACAAatgAGATTTGATGAA-GCTTATTCA	Clone *speB* into pDG1664 digested with BamHI and EcoRI FOR
KN478	AACTTTAGGGTTATCGAATTttaTTTCACAAACCCAA-GCAG	Clone *speB* into pDG1664 digested with BamHI and EcoRI REV
KN490	AAAACTGCTGCCTTCGGATCTTTCTTCACTCCTT-GTAGATTTAC	Clone *speA* with 184 bp P_{speA} promoter into pDG1664 digested with BamHI and EcoRI FOR
KN491	AACTTTAGGGTTATCGAATTttaTTGAATTGCTTTTT-GTTCTTTG	Clone *speA* with 184 bp P_{speA} promoter into pDG1664 digested with BamHI and EcoRI REV
KN492	AAAACTGCTGCCTTCGGATCAAC-GGGTTCCCCCTTTTTTT	Clone *speE* with 201 bp P_{speEB} promoter into pDG1664 digested with BamHI and EcoRI FOR

KN493	AACTTTAGGGTTATCGAATTTTATTTAATCAGGTCAC TCACAAA	Clone *speE* with 201 bp P$_{speEB}$ promoter into pDG1664 digested with BamHI and EcoRI REV

Table 5: List of oligonucleotides used in this study

DATA AVAILABILITY

All data reported in this paper are available in the Statistics Source Data.

QUANTIFICATION AND STATISTICAL ANALYSIS

Ordinary one-way ANOVA p-value calculations followed by a Sidak's multiple comparisons test were used to determine statistical significance for figures: 1D, 3A, 3C, 4B, and 4D. 2A and 2H statistics were calculated in Compound Discoverer 3.3.3.200. Ordinary one-way ANOVA p-value calculations followed by a Dunnett's multiple comparisons test were performed to determine statistical significance for figures 3F. A Two-way ANOVA followed by a Tukey multiple comparisons test was performed to determine statistical significance for figure 3B. A Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons correction was performed for figures 4F, 4G, and SFig. 7B. All calculations were performed in GraphPad Prism 10.1.2 (324) [36]. No statistical method was used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.

METHOD DETAILS

C. elegans growth media and conditions

C. elegans strains were maintained at 15 °C or 20 °C on NGM agar plates (3 g/L NaCl, 17 g/L Agar, 2.5 g/L peptone (Bacto), 25 mM KPO_4 (pH 6), 5 mg/L Cholesterol, 1 mM $MgSO_4$, 1 mM $CaCl_2$) seeded with *E. coli* OP50 (CGC, DA837). All *C. elegans* strains used in this study are listed in Table 3. *C. elegans* that were fed a diet of *B. subtilis* or *B. subtilis* mutants for experimentation were grown on modified NGM (3 g/L NaCl, 17 g/L Agar, 5.0 g/L peptone (Bacto), 25 mM KPO_4 (pH 9.2), 5 mg/L Cholesterol, 1 mM $MgSO_4$, 1 mM $CaCl_2$).

Egg Prep Protocol

Adult *C. elegans* were washed from NGM agar plates with M9 buffer (3 g/L KH_2PO_4, 6 g/L

Na_2HPO_4, 5 g/L NaCl, 1 mM $MgSO_4$). *C. elegans* were lysed with Egg Preparation Solution

(30% Sodium hypochlorite, 1 M NaOH) and monitored by dissection microscope for ~[1]-3

minutes. The embryos were washed 2-3 times in M9 buffer.

Bacteria growth media and conditions

All bacterial strains were grown on Lennox LB Agar (5 g/L NaCl, 10 g/L SELECT Peptone 140, 5

g/L SELECT Yeast Extract) (Invitrogen, 12780029) at 30 °C or 37 °C unless specified otherwise

and are listed in Table 3. Liquid cultures were grown at 30 °C and shaken in <5 mL Lennox LB

at 220 rpm in 18 mm glass test tubes. *Bacillus* strains containing pDG1664 and its derivatives

inserted at *thrC* were grown in Lennox LB containing MLS (25 µg/mL erythromycin (Thermo

Fisher Scientific, 50-213-296) and 5 µg/mL lincomycin (Thermo Fisher Scientific, 50-213-395)).

Bacillus non-essential gene knock out library mutants were grown in kanamycin (7.5 µg/mL

(Sigma, K1377)). Plasmid selection in *E. coli* was performed on Lennox LB containing

kanamycin (25 µg/mL, Sigma Aldrich, K1377) or ampicillin (100 µg/mL, Sigma Aldrich, A0166).

Bacteria strains and plasmid construction

All plasmids were constructed by isothermal assembly (ITA) and are listed in Table 4. DNA

regions for plasmid assembly were identified using Biocyc [37] and amplified by touchdown PCR

from the bacterial chromosomes using Q5 DNA polymerase (New England Biolabs, M0492) and

oligonucleotides from Integrated DNA Technologies (Coralville, IA) listed in Table 5. All

oligonucleotides are listed in Table 5. PCR fragments were purified using the Monarch DNA Gel

Extraction Kit (New England Biolabs, T1020) and assemblies were performed using the

NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, E2621). All regions amplified

by PCR were confirmed by DNA sequencing at Genewiz from Azenta Life Sciences (South

Plainfield, NJ). Plasmids were transformed according to the manufacturer's instructions and

maintained in *E. coli* DH5α competent cells (New England Biolabs, C2988J). Plasmids or

purified chromosomal DNA were transformed into *B. subtilis* strains by natural competency as

previously described [25].

All *Bacillus subtilis* strains reported in this study are isogenic derivatives of *B. subtilis* 168

(BGSCID: 1A1) and are reported in Table 3. The operon encoding *speEB* was knocked out by

homologous recombination using the suicide vector, pMAD [38]. Approximately 1 kb upstream and

downstream of *speEB* were amplified and assembled via ITA into pMAD digested with BglII and

EcoRI (New England Biolabs, R0144 and R3101, respectively), resulting in plasmid pNOBb41

(Table 4). The deletion was confirmed by PCR, sanger sequencing, and whole genome

sequencing.

Strain NOBb518 was constructed by transforming purified chromosomal DNA (see methods

section: *Bacillus* genomic DNA extraction) from strain BKK14630 (*speA::kanR*) [25] into NOBb369

(Δ*speB*). Chromosomal DNA was transformed by natural competency as previously described

[25].

Plasmids for complementation studies (pNOB47, pNOB48, and pNOB49) were constructed by

amplifying target genes from the *Bacillus subtilis* 168 chromosomes using the primers listed in

Table 4. These amplicons were assembled in pDG1664 digested with EcoRI and HindIII (New

England Biolabs, R3101 and R3104, respectively) by ITA and were expressed under the control

of their respective native promoters. When necessary *kanR* was removed from the *Bacillus*

subtilis chromosome using pDR244 [25] (Bacillus Genetic Stock Center, BGSCID ECE274).

E. coli OP50 Δ*speB* (NOBb750) was constructed by P1 transduction using *E.coli* K12 Δ*speB*

lysate. The lysate was prepared using *E. coli* P1 bacteriophage (ATCC, 25404-B1). Briefly, *E.*

coli K12 Δ*speB* donor strain was grown overnight at 37 $^{\circ}$C in LB (30 μg/mL kanamycin). The

culture was diluted 1:100 in LB supplemented with 0.2% glucose and 5 mM $CaCl_2$, and incubated at 37 $^{\circ}$C for 45 min, 220 rpm. 100 μL of *E. coli* bacteriophage P1 (ATCC 25404B1^{TM}) was added to the donor bacteria culture and incubated at 37 $^{\circ}$C for 3 hr. 200 μL of chloroform were added and incubated at 37 $^{\circ}$C for 5 min. The mixture was centrifuged at 9,200xg for 5 min at 4 $^{\circ}$C. Lysate was further purified using a 0.45 μm filter before collecting the P1 phage-containing supernatant. Recipient strain *E. coli* OP50 was grown overnight at 37 $^{\circ}$C in 3 mL LB broth. *E. coli* OP50 cells were resuspended in a P1 salt solution (10 mM $MgCl_2$, 5 mM $CaCl_2$). Next, 10 μL of the P1 lysate was mixed with 100 μL of the recipient cells and incubated with shaking (220 rpm) at 37 $^{\circ}$C for 30 min. Then, 1 mL of LB and 200 μL of 1 M sodium citrate was added and incubated at 37 $^{\circ}$C for 1 hr, before being centrifuged at 13,000xg for 2 min. Transformants were selected on kanamycin (30 μg/mL) and sodium citrate (5 mM sodium citrate) LB [39,40].

Screen of wild bacterial isolates for *C. elegans* development

Compost samples from the Grand Rapids, MI area were collected, diluted in M9 buffer, filtered, and plated on LB. Single colonies were grown overnight at 30 °C and 200 µL of culture was seeded on NGMB 12-well plates. The plates were dried in a laminar flow hood. Approximately 100-200 germ-free *C. elegans daf-2* (DR1572) embryos were added to each well. The plates were incubated for 5 days at 26 °C. The abundance of L4+ animals was qualified by eye using a dissection microscope. Strains of bacteria that promoted *C. elegans* development were retested on 90 mm NGM plates to confirm.

Screen of *B. subtilis* 168 knockout library for loss of *C. elegans* development

The *B. subtilis* single gene deletion library--kanamycin was acquired from Addgene (Watertown, MA, 1000000115). The parent strain was acquired from the Bacillus Genetic Stock Center (Columbus, OH, BGSCID: 1A1). The library was replicated into 96-well plates for further use.

Each 96-well plate was replicated and incubated overnight at 30 °C in LB. Each well was seeded on NGMB in a 12-well plate and dried in a laminar flow hood. 50-100 *C. elegans daf-2* (DR1572) embryos were added to each well. The plates were incubated for 5 days at 26 °C. Each *B. subtilis* mutant that did not promote animal development was retested on a larger scale using 90 mm plates.

Bacillus genomic DNA extraction and whole genome sequencing – *B. subtilis*

Single bacteria colonies were grown overnight in 3 mL liquid culture. Bacteria culture was centrifuged at 17,000 x g for 1 min. Genomic DNA was isolated as previously described (58). Briefly, the pelleted cells were resuspended in Chromo Lysis Buffer (50 mM EDTA, 0.1M NaCl, pH 7.5). Lysozyme was added (20 µL of 2.5 mg/mL) and incubated at 37 °C for 10 minutes. Sarkosyl was added (1%) and the cells were incubated for 5 minutes at 37 °C. DNA of each isolate was extracted with phenol/chloroform/isoamyl alcohol 25:24:1 and ethanol precipitation. Pelleted were dried and suspended in sterile DI water. Total genomic DNA was quantified using a qubit fluorometer and visualized on a 1% agarose gel for degradation. Genomic DNA from each isolate was submitted for whole genome sequencing on an Illumina platform PE150 at SeqCenter (Pittsburg, PA). FastQC version 0.12.1 (GPL v3) was used to assess the quality control of the raw sequencing data [41]. Reads were trimmed using Trimmomatic version 0.39 [42] to remove adapter sequences and low-quality reads (Q<30). Unicycler version v0.5.0 with default parameters was used to assemble the bacterial genome [43].

EMS mutagenesis of WT *C. elegans*

L4 staged WT *C. elegans* were washed in M9 buffer (recipe: 3 g/L KH_2PO_4, 6 g/L Na_2HPO_4, 5 g/L NaCl, 1 mM $MgSO_4$) 3 times and resuspended in 4 mL M9 buffer. 20 µL of Ethyl methanesulfonate (EMS) (Sigma Aldrich, M0880) were added and the animals were incubated for 4 hours with a rotary mixer, at 20 °C. The animals were washed three times and plated in 20

pools of ~200 on *E. coli* OP50 seeded NGM. The animals were screened on *B. subtilis ∆speB* for development beyond L4. Approximately one animal from each pool that developed beyond L4 was isolated and whole genome sequenced.

C. elegans genomic DNA extraction and whole genome sequencing

WT *C. elegans* were incubated on NGM for 3 days at 20 °C. The animals were washed in M9 buffer. and suspended 1:1 in Worm Lysis Buffer (50 mM KCl, 10 mM Tris (pH 8.3), 2.5 mM MgCl2, 0.45% Tween 20, 0.01% Gelatin, 0.45% Nonidet P 40 Substitute (Sigma Aldrich, 74385). The animals were incubated at 56 °C for 1 hour with 10 µL of proteinase K (20 mg/mL). After lysis, genomic DNA was purified by phenol chloroform extraction and ethanol precipitation. DNA pellets were resuspended in sterile DI water.

Whole genome Illumina sequencing and variant calling was performed by SeqCenter (Pittsburg, PA). Briefly, Illumina sequencing libraries were prepared using the tagmentation-based and PCR-based Illumina DNA Prep kit and custom IDT 10bp unique dual indices (UDI) with a target insert size of 320 bp. No additional DNA fragmentation or size selection steps were performed. Illumina sequencing was performed on an Illumina NovaSeq 6000 sequencer in one or more multiplexed shared-flow-cell runs, producing 2x151bp paired-end reads. Demultiplexing, quality control and adapter trimming was performed with bcl-convert1 (v4.1.5). The fastq paired end reads were prepared for variant calling using Samtools and then indexed using bwa-mem index. Raw reads were then mapped to the indexed reference using bwamem and subsequently sorted and converted to BAM files using Samtools. GATK's3 MarkDuplicates functionality was used to remove duplicate reads from the alignment file. GATK's Haplotype caller was then used to call variants on the alignment file. BcfTools was used to filter out variants with a QD < 2, or MQ < 40 or MQRankSum < -12.5, or ReadPosRankSum < -8, or FS > 60.0, or SOR >3.

Microscopy

C. elegans strains were washed with M9 buffer prior to imaging to remove excess bacteria. The animals were paralyzed with tetramisole (~10 μg/mL). and incubated for ~2 minutes. Images were taken with a Leica DM6 B upright microscope. Images were pseudo-colored using FIJI [44]. Images were assembled in Adobe Illustrator 2024 28.4.1 64-bit [45].

Metabolomics

Extraction

Metabolites from LB broth were extracted via the addition of ice cold 2:2:1 (v/v) mixture of acetonitrile:methanol:water (#A456, #A955, #W6, respectively; Thermo Fisher Scientific) at a ratio of 30 μl sample per mL of extraction solvent. The samples were vortexed for 10 sec, sonicated for 5 minutes in a water bath sonicator, and incubated on wet ice for 60 min. After incubation, samples were centrifuged at 17,000g and 4 °C to pellet the insoluble fraction. The metabolite-containing supernatant was collected, dried in a vacuum evaporator, and stored at -80°C until derivatization.

Polyamine Derivatization

Carbamylated-derivatives were generated by derivatizing the dried metabolite fraction with isobutylchloroformate**.** Dried metabolite extracts were resuspended in 200 μL of water containing 1 μg/mL of 1,6 Diaminohexane as an internal standard. 5 μL of sodium bicarbonate (1M, pH 9.0) was added to each sample vial followed by the addition 20 μL of isobutyl chloroformate (177989, Sigma). Samples were vortexed for 10 seconds then incubated at 37 °C for 15 minutes. After cooling to room temperature, 1 mL of diethyl ether (309966, Sigma) was added, and samples were vortexed for 10 seconds. Samples were then centrifuged for 1 minute to separate the organic and aqueous layers. 800 μL of the carbamylated metabolite containing organic layer was transferred to an autosampler vial, and subsequently dried under a gentle stream of nitrogen gas. Dried samples were resuspended in 100 μL of 1:1 (v/v)

acetonitrile:water, vortexed, and 90 μL transferred to a fresh autosampler vial for LC-MS

analysis. For LCMS analysis of N^1-APA, chemical standard (EN300-47248193, Enamine) was

prepared at 2 μg/mL in water and derivatized as described above as either a neat standard or

spiked into dried metabolite extracts.

Untargeted LC-MS

LC-MS analysis of samples for Fig. 2A-F was performed on derivatized metabolite extracts

analyzed by high resolution accurate mass spectrometry using an ID-X Orbitrap mass

spectrometer (Thermo Fisher Scientific) coupled to a Thermo Vanquish Horizon liquid

chromatography system. 2 μL of sample volume was injected on column. Chromatographic

separations were accomplished with a Cortecs T3 (120Å, 1.6 μm, 2.1 mm × 150 mm) analytical

column (#186008500, Waters, Eschborn, Germany) fitted with a pre-guard column (120Å, 1.6

μm, 2.1mm × 5 mm; #186008508, Waters, Eschborn, Germany) using an elution gradient with a

binary solvent system. Solvent A consisted of LC/MS grade water (W6–4, Fisher), and Solvent B

was 99% LC-MS grade acetonitrile (A955, Fisher). Both mobile phases contained 0.1% (v/v)

formic acid (A11710X1, Fisher). The 15-min analytical gradient at a flow rate of 400 μL/min was:

0–0.5 min hold at 100% A, 0.5–2.0 min ramp from 0% B to 40% B, 2.0-10.0 min from 40% B to

99% B, then held at 99% B from 10.0-15.0 min. Following the analytical separation, the column

was re-equilibrated for 3.5 min as follows: 0–0.5 min hold at 100% A at 600 μL/min, 0.5–0.6 min

decrease from 600 μL/min to 400 μL/min, 0.6–3.5 min hold at 100% A and 400 μL/min. The

column temperature was maintained at 50°C. The H-ESI source was operated at spray voltage

of 2500 V(negative mode)/3500 V(positive mode), sheath gas: 70 a.u., aux gas: 25 a.u., sweep

gas: 1 a.u., ion transfer tube: 300°C, vaporizer: 250 °C. Sample data were collected via data

dependent MS2 (ddMS2) fragmentation using MS1 resolution at 60,000, MS2 resolution at

7,500, intensity threshold at 2.0×10^4, and dynamic exclusion after two triggers for 10 s.

MS2 fragmentation was completed first with HCD using stepped collision energies at 15, 30,

and 45% and was followed on the next scan by CID fragmentation in assisted collision energy

mode at 15, 30, and 45% with an activation Q of 0.25. MS1 scans used custom AGC target of

25% in positive mode and a maximum injection time of 50 ms. MS2 scans used standard AGC

target and a maximum injection time of 22ms. The total cycle time of the MS1 and ddMS2 scans

was 0.6 s. Untargeted metabolomics data were analyzed in Compound Discoverer (v 3.3,

Thermo). Figures were prepared in GraphPad Prism and FreeStyle™ 1.8 SP2 QF1 Version

1.8.65.0.

Targeted LC-MS

LC-MS analysis of samples for Fig. 2H was performed on derivatized metabolite extracts

analyzed using an Agilent 6470 LC-QQQ mass spectrometer coupled to a Agilent 1290 UHPLC

system as previously reported[21]. Targeted metabolomics data were analyzed in Skyline

23.1.0.455 (Agilent). Statistics were performed using ols and anova_lm from statsmodels

0.14.4, posthoc_ttest from scikit_posthocs 0.7.0 with no corrections, and multipletests, method

= fdr_bh from statsmodels 0.14.4 for global p-value corrections. Figures were prepared in

GraphPad Prism 10.3.0 (507).

Brood size count

Pristionchus pacificus PS312 grew to adulthood on NGM plates seeded with *E. coli* OP50 at 20

°C. Embryos were isolated using egg preparation solution and plated on NGMB plates seeded

with the respective bacterial strains. Bacterial strains were grown overnight at 30 °C before

seeding. Once the animals reached the L4 growth stage, 10 animals were singled onto NGMB

seeded with the respective bacterial strains, prepared from overnight cultures grown at 30 °C.

The plates were incubated at 20 °C for the duration of the experiment. Offspring were counted

each day for approximately 5 days. Animals that died on day 1 or 2 were removed from the data set.

Cell Culture

Bone marrow cells from femur were isolated from C57BL/6 female mice, RBC lysed, pooled (n=3-6 mice), counted, and resuspended at 1x10^7/ml, and plated at 1.5x10^6 per well. To generate Bone Marrow Macrophages (BMMΦ) cells were cultured in RPMI supplemented with 10% FBS (Hyclone), 2mM L-glutamine (Gibco), 100U/mL penicillin/streptomycin (Gibco), with 20ng/ml M-CSF (R&D; 416-ML-050) for 7 days. BMM_Φ(IL4) were generated with 20ng/ml IL-4 (Peprotech 214-14-20ug) overnight from day 7; N^{1}-guanyl-1 7-diaminoheptane (Stock 20mM) (GC7 1:100) 200μM and N^1-APA (Stock 20mM) (APA 1:100) 200uM treatment began on day 7.

BMM Flow

Following treatment on D7, BMM_Φ were harvested, washed and stained for flow cytometry. Samples were incubated with Fc block and eFluor 506 Fixable Viability Dye (ThermoFisher) in PBS, followed by an antibody cocktail prepared in wash buffer (PBS with 1% FBS, 1 mM EDTA, and 0.05% NaN_3). For intracellular staining, cells were fixed with IC fixation buffer (eBioscience/ThermoFisher) for 30 minutes, permeabilized using Permeabilization Buffer (eBioscience/ThermoFisher) and incubated for at least one hour with antibodies targeting intracellular proteins. Samples were acquired on the Cytek Aurora spectral cytometer and data analyzed using FlowJo v10.

BMM_Φ Western Blot for Hypusine and eIF5a

After 24 hours of treatment, BMMΦ were removed from culture by washing with warm 1xPBS followed by a wash with chilled 1xPBS and placed on ice for 5 minutes then gently scraped, resuspended and put into 15ml falcon tubes. Cells were pelleted at 4°C for 5min at 400g. Pellets were lysed with 1x CHAPS lysis buffer (+PI;+PMSF) freeze thawed twice at -80°C, sonicated by 10 seconds in an ice slurry followed by 1 minute rest (repeated 5 times). Supernatant was clarified by centrifuging for 15 minutes 4°C at 17,000xg. Lysed samples were stored at -80°C. Protein quantification was done with the Pierce 660nm Protein Assay Reagent (ThermoFisher; 22660); samples were diluted 1:2 in 1xCHAPS and read on a plate reader along with Pre-diluted Protein Assay Standards: Bovine Serum Albumin Set (ThermoFisher; 23208). For Hypusination and eIF5A blots,1 to 5µg of protein for each sample was removed, brought up to volume with 1xCHAPS and 6xLB. Samples were heated at 70°C for 10 minutes. For Western Blot 4-20% mini-protean gels were used and ran at 70V for 30 minutes and 100V until desired (30min). Proteins were transferred to PVDF membranes, blocked for 1hr 4°C in 5% milk. Membranes were incubated with primary antibodies for anti-hypusine (1:10000) (Millipore ABS1064-I-25ul) and anti-eIF-5a (1:10000) (BD Biosciences 611976) overnight at 4°C in 5% BSA+TBST. Secondary antibodies, 1hr RT 1:10000 in 5% milk+TBST.

References

1. Watanabe, S., Kusama-Eguchi, K., Kobayashi, H. & Igarashi, K. Estimation of polyamine binding to macromolecules and ATP in bovine lymphocytes and rat liver. *J. Biol. Chem.* **266**, 20803–20809 (1991).

2. Matsumoto, M., Kakizoe, K. & Benno, Y. Comparison of fecal microbiota and polyamine concentration in adult patients with intractable atopic dermatitis and healthy adults. *Microbiol. Immunol.* **51**, 37–46 (2007).

3. van Veen, S. *et al.* Cellular function and pathological role of ATP13A2 and related P-type transport ATPases in Parkinson's disease and other neurological disorders. *Front. Mol. Neurosci.* **7**, (2014).

4. Vrijsen, S. *et al.* ATP13A2-mediated endo-lysosomal polyamine export counters mitochondrial oxidative stress. *Proc. Natl. Acad. Sci.* **117**, 31198–31207 (2020).

5. Dang, T. *et al.* ATP13A2 protects dopaminergic neurons in Parkinson's disease: from biology to pathology. *J. Biomed. Res.* **36**, 98–108 (2022).

6. Nakamura, A. *et al.* Symbiotic polyamine metabolism regulates epithelial proliferation and macrophage differentiation in the colon. *Nat. Commun.* **12**, 2105 (2021).

7. Pryor, R. *et al.* Host-Microbe-Drug-Nutrient Screen Identifies Bacterial Effectors of Metformin Therapy. *Cell* **178**, 1299-1312.e29 (2019).

8. Bachrach, U., Wang, Y.-C. & Tabib, A. Polyamines: New Cues in Cellular Signal Transduction. *Physiology* **16**, 106–109 (2001).

9. Alcázar, R. *et al.* Polyamines: molecules with regulatory functions in plant abiotic stress tolerance. *Planta* **231**, 1237–1249 (2010).

10. Polyamines regulate intestinal epithelial restitution through TRPC1-mediated Ca2+ signaling by differentially modulating STIM1 and STIM2 | American Journal of Physiology-Cell Physiology. https://journals.physiology.org/doi/full/10.1152/ajpcell.00120.2012.

11. Toro-Funes, N., Bosch-Fusté, J., Veciana-Nogués, M. T., Izquierdo-Pulido, M. & Vidal-Carou, M. C. *In vitro* antioxidant activity of dietary polyamines. *Food Res. Int.* **51**, 141–147 (2013).

12. Muñoz-Esparza, N. C. *et al.* Polyamines in Food. *Front. Nutr.* **6**, (2019).

13. Acosta-Andrade, C. *et al.* Polyamine-RNA-membrane interactions: From the past to the future in biology. *Colloids Surf. B Biointerfaces* **155**, 173–181 (2017).

14. Dever, T. E. & Ivanov, I. P. Roles of polyamines in translation. *J. Biol. Chem.* **293**, 18719–18729 (2018).

15. Igarashi, K. & Kashiwagi, K. The functional role of polyamines in eukaryotic cells. *Int. J. Biochem. Cell Biol.* **107**, 104–115 (2019).

16.   Puleston, D. J. *et al.* Polyamines and eIF5A Hypusination Modulate Mitochondrial Respiration and Macrophage Activation. *Cell Metab.* **30**, 352-363.e8 (2019).

17.   Anderson-Baucum, E. *et al.* Deoxyhypusine synthase promotes a pro-inflammatory macrophage phenotype. *Cell Metab.* **33**, 1883-1893.e7 (2021).

18.   Kasahara, N. *et al.* Dietary polyamines promote intestinal adaptation in an experimental model of short bowel syndrome. *Sci. Rep.* **14**, 4605 (2024).

19.   Sekowska, A., Bertin, P. & Danchin, A. Characterization of polyamine synthesis pathway in Bacillus subtilis 168. *Mol. Microbiol.* **29**, 851–858 (1998).

20.   Rashid, M. H., Kumar, S. P., Rajan, R. & Mamillapalli, A. Salivary microbiota dysbiosis and elevated polyamine levels contribute to the severity of periodontal disease. *BMC Oral Health* **25**, 2 (2025).

21.   Isaguirre, C. N., Gendjar, M., Nauta, K. M., Burton, N. O. & Sheldon, R. D. Isobutyl chloroformate derivatization of polyamines for LC-MS quantitatio. in *Enzymes of Polyamine Metabolism* (Methods in Enzymology, 2025).

22.   Li, B. *et al.* Functional identification of bacterial spermine, thermospermine, norspermine, norspermidine, spermidine, and N1-aminopropylagmatine synthases. *J. Biol. Chem.* **300**, (2024).

23.   Ohnuma, M. *et al.* N1-Aminopropylagmatine, a New Polyamine Produced as a Key Intermediate in Polyamine Biosynthesis of an Extreme Thermophile, Thermus thermophilus *. *J. Biol. Chem.* **280**, 30073–30082 (2005).

24.   Discovery of disease-adapted bacterial lineages in inflammatory bowel diseases: Cell Host & Microbe. https://www.cell.com/cell-host-microbe/fulltext/S1931-3128(24)00194-X.

25.   Koo, B.-M. *et al.* Construction and Analysis of Two Genome-Scale Deletion Libraries for Bacillus subtilis. *Cell Syst.* **4**, 291-305.e7 (2017).

26.   Sekowska, A., Danchin, A. & Risler, J.-L. Phylogeny of related functions: the case of polyamine biosynthetic enzymes. *Microbiology* **146**, 1815–1828 (2000).

27.   Pellegrino, M. W. *et al.* Mitochondrial UPR-regulated innate immunity provides resistance to pathogen infection. *Nature* **516**, 414–417 (2014).

28.   Heinick, A. *et al.* Caenorhabditis elegans P5B-type ATPase CATP-5 operates in polyamine transport and is crucial for norspermidine-mediated suppression of RNA interference. *FASEB J.* **24**, 206–217 (2010).

29.   Prabh, N. *et al.* Deep taxon sampling reveals the evolutionary dynamics of novel gene families in Pristionchus nematodes. *Genome Res.* **28**, 1664–1674 (2018).

30.   Rödelsperger, C. *et al.* Phylotranscriptomics of Pristionchus Nematodes Reveals Parallel Gene Loss in Six Hermaphroditic Lineages. *Curr. Biol.* **28**, 3123-3127.e5 (2018).

31.   Posttranslational synthesis of hypusine: evolutionary progression and specificity of the hypusine modification | Amino Acids. https://link.springer.com/article/10.1007/s00726-007-0525-0.

32.   Jakus, J., Wolff, E. C., Park, M. H. & Folk, J. E. Features of the spermidine-binding site of deoxyhypusine synthase as derived from inhibition studies. Effective inhibition by bis- and mono-guanylated diamines and polyamines. *J. Biol. Chem.* **268**, 13151–13159 (1993).

33.   Hypusination Maintains Intestinal Homeostasis and Prevents Colitis and Carcinogenesis by Enhancing Aldehyde Detoxification - ScienceDirect. https://www.sciencedirect.com/science/article/abs/pii/S0016508523008144.

34.   Caenorhabditis Genetics Center (CGC) - College of Biological Sciences. https://cgc.umn.edu/.

35.   BGSC - Home. https://bgsc.org/.

36.   Korbie, D. J. & Mattick, J. S. Touchdown PCR for increased specificity and sensitivity in PCR amplification. *Nat. Protoc.* **3**, 1452–1456 (2008).

37.   Karp, P. D. *et al.* The BioCyc collection of microbial genomes and metabolic pathways. *Brief. Bioinform.* **20**, 1085–1093 (2019).

38.   Arnaud, M., Chastanet, A. & Débarbouillé, M. New Vector for Efficient Allelic Replacement in Naturally Nontransformable, Low-GC-Content, Gram-Positive Bacteria. *Appl. Environ. Microbiol.* **70**, 6887–6891 (2004).

39.   Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection | Molecular Systems Biology. https://www.embopress.org/doi/full/10.1038/msb4100050.

40.   Thomason, L. C., Costantino, N. & Court, D. L. Genome Manipulation by P1 Transduction. *Curr. Protoc. Mol. Biol.* **79**, 1.17.1-1.17.8 (2007).

41.   Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

42.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

43.   Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads | PLOS Computational Biology. https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1005595.

44.   Fiji: an open-source platform for biological-image analysis | Nature Methods. https://www.nature.com/articles/nmeth.2019.

45.   Industry-leading vector graphics software | Adobe Illustrator. https://www.adobe.com/products/illustrator/campaign/pricing.html.