Small Extracellular Vesicle Signaling and Mitochondrial Transfer Reprograms T Helper Cell Function in Human Asthma

Kenneth P. Hough[1], Jennifer L. Trevor[1], Balu K. Chacko[2], John G. Strenkowski[1], Yong Wang[1], Kayla Goliwas[1], Nathaniel B. Bone[1], Young-il Kim[1], Renita Holmes[1], Shia Vang[1], Alexandra Pritchard[1], Jay Chin[1], Sandeep Bodduluri[1], Veena B. Antony[1], Sultan Tousif[1], Mohammad Athar[3], Diptiman Chanda[1], Kasturi Mitra[4], Jaroslaw Zmijewski[1], Jianhua Zhang[2], Steven R. Duncan[1], Victor J. Thannickal[1], Susanne Gabrielsson[5,6], Victor M. Darley-Usmar[2], and Jessy S. Deshane[1*]

[1]Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, [2]Mitochondrial Medicine Laboratory, Department of Pathology, [3]Department of Dermatology, [4]Department of Genetics, University of Alabama at Birmingham, AL 35294, [5]Division of Immunology and Allergy, Department of Medicine, Karolinska Institutet and [6]The Karolinska University Hospital, Solna, Stockholm, Sweden

*Correspondence:treena@uab.edu.

Jessy Deshane, Ph.D.
Professor
Department of Medicine
Division of Pulmonary Allergy and Critical Care Medicine
THT 433A, 1900 University Boulevard
University of Alabama at Birmingham, 35294
treena@uab.edu, Tel: 1-205-996-2041

Author's Contributions: KPH and JSD performed experiments, data analysis, interpretations, and wrote the manuscript. JLT performed bronchoscopies, facilitated sample collection, and provided clinical oversight. RH was the clinical co-ordinator. YW processed BAL fluid samples for flow cytometry and did data analysis. KG did co-culture experiments and confocal imaging. JGS, NB, SV, AP, ST, JW and JC helped with co-cultures and ImageStream data analysis. SB helped develop a computation tool to automate co-localization analysis of ImageStream data using MATLAB. YK performed statistical analysis. VJT, VDU, MA, SG, JZ, SR, KM, VBA, JZ, BKC, DC assisted with data interpretation, and feedback on the manuscript.

Grant Support: This work was supported by FAMRI YCSA 2010, Parker B. Francis Fellowship and R01HL128502 awarded to JSD, U54 ES030246-01 awarded to MA, P01HL114470 awarded to VJT, Nathan Shock Center P30 G050886 (JZ, VDU), the Swedish Heart-Lung Foundation Grant # 20200646, 20230528 (SG), The Swedisch Cancer and Allergy Association (SG), NIH P30 AR048311 and NIH P30 AI27667 for service and support provided by the UAB Comprehensive Flow Cytometry Core.

Running Head: Small Extracellular Vesicle mitochondria activates T cells in asthma
Descriptor Number: 1.20 Immunology/Inflammation: Human Studies
Word count of the body: 3499.

Impact: The mechanisms for unremitting airway inflammation in human asthma are unknown. Our studies uncover mitochondrial-small extracellular vesicle-mediated signaling in T cell activation and differentiation that contribute to dysregulated immune responses in asthmatic subjects.

Abstract

Rationale: Asthma is a chronic inflammatory disease of the airways that involves crosstalk between myeloid-derived regulatory cells (MDRCs) and CD4$^+$ T cells. Although small extracellular vesicles (sEVs) are known to mediate cell-cell communication, the role of sEV signaling via mitochondria in perpetuating asthmatic airway inflammation is unknown.

Objectives: We investigated the effects of MDRC-derived exosomes on dysregulated T cell responses in asthmatics.

Methods: Small extracellular vesicles isolated from bronchoalveolar lavage fluid or airway MDRCs of mild to moderate asthmatics or healthy controls were co-cultured with autologous peripheral and airway CD4$^+$ T lymphocytes. sEV internalization, sEV-mediated transfer of mitochondria targeted GFP to T cells, sEV mitochondrial signaling, and subsequent activation, proliferation and polarization of CD4$^+$ T lymphocytes to Th1, Th2 and Th17 subsets were assessed.

Measurements and main results: Airway MDRC-derived sEVs from asthmatics mediated T cell receptor engagement and transfer of mitochondria that induced antigen-specific activation and polarization into Th17 and Th2 cells, drivers of chronic airway inflammation in asthma. CD4$^+$ T cells internalized sEVs containing mitochondria predominantly by membrane fusion, and blocking mitochondrial oxidant signaling in MDRC-derived exosomes mitigated T cell activation. Reactive oxygen species-mediated signaling that elicited T cell activation in asthmatics was sEV-dependent. A Drp1-dependent mitochondrial fission in pro-inflammatory MDRCs promoted mitochondrial packaging within sEVs, which then co-localized with the polarized actin cytoskeleton and mitochondrial networks in the organized immune synapse of recipient T cells.

Conclusions: Our studies indicate a previously unrecognized role for mitochondrial fission and exosomal mitochondrial transfer in dysregulated T cell activation and Th cell differentiation in asthma which could constitute a novel therapeutic target.

Word count: 249

Introduction

CD4[+] T cells orchestrate the immune response in diverse chronic inflammatory diseases, including asthma[1]. Pathogenic T helper type 2 (Th2) and T helper type 17 (Th17) subsets drive airway inflammation, remodeling and smooth muscle hyperplasia in asthmatics[2-11]. Recent advances implicate cellular crosstalk between the innate and adaptive immune systems in the initiation and propagation of the allergic immune response. In allergic asthma, antigen presenting cells (APCs) internalize antigens, present processed peptides bound to class II molecules[12-14] and engage the T cell receptor (TCR), triggering a multi-molecular signaling cascade to activate CD4[+] T cells[15]. HLA-DR expressing APCs, including immature myeloid cells, regulate T cell function through TCR engagement[16-21]. In human asthma, the pro-inflammatory HLA-DR[+] subsets of myeloid-derived regulatory cells (MDRCs) promote proliferation of T cells via redox signaling pathways [21]. Although antigen presentation by MDRCs is not well-defined, TCR engagement and redox mechanisms may synergize to regulate MDRC-mediated T cell activation.

Recently, acellular antigen presentation by exosomes, has been described as an unconventional mechanism of immune activation[22-25]. Small extracellular vesicles (sEVs) that areendosomally-derived class of extracellular vesicles, known for facilitating cellular crosstalk, are decorated with scaffolding proteins for immune receptors called tetraspanins[24, 26-31]. Tetraspanins cluster at the immune synapse, and aid in antigen presentation and immune signaling[30] and, thus, sEVs may modulate the immune response via the immune synapse. While professional APC-derived MHC-peptide bound sEVs induce Th2 cytokines in experimental asthma[22, 24, 32], antigen presentation by

MDRC-derived sEVs as a driver of dysregulated CD4[+] T cell responses has not been

demonstrated.

We reported that MDRC-derived sEVs containing polarized mitochondria are

internalized by CD4[+] T cells[33]. Increasing evidence for mitochondrial components and

respiration in sEVs suggests that functional mitochondria packaged within the sEVs may

elicit diverse physiologic responses[33-39]. This unconventional signaling mechanism may

compensate for the altered bioenergetics of recipient cells and/or provide an alternative

mechanism to outsource autophagy/mitophagy during cellular stress[36, 37, 40, 41].

Here, we show that MDRC-derived sEV mitochondria with intact membrane

potential activate CD4[+] T cell proliferation and polarization into Th2 and Th17 cells via

mitochondrial oxidant signaling; this event is triggered by membrane fusion of sEV-

derived mitochondria and downstream complex formation of polarized cytoskeleton and

mitochondrial network in recipient T cells. We demonstrate that mitochondrial fission,

regulated by Drp1 oligomerization, is required for packaging functional mitochondria

within sEVs and implicates MDRC-derived sEV-dependent mitochondrial signaling in Th

cell polarization and immune regulation of chronic airway inflammation in mice with

asthma and in human asthmatics.

Results

Bronchoalveolar lavage fluid (BALF) sEVs from asthmatics drive CD4[+] T cell

proliferation and Th polarization

Immune regulation by sEVs is attributed to expression of MHC-II (HLA-DR) and

co-stimulatory molecules CD81, CD86 and CD54[25, 42]; the percent of HLA-DR$^+$ exosomes

and HLA-DR expression is increased in airways of asthmatics compared to healthy

controls[24, 43, 44]. In our studies (Table I: Characteristics of study subjects), BALF sEVs

from asthmatics when co-cultured with peripheral autologous CD4$^+$ T cells (10 sEVs:1 T

cell ratio), induced their proliferation (CFSE dilution) (Figure 1A-1B). Furthermore,

proliferation of autologous airway CD4$^+$ T cells of asthmatics was also increased in co-

cultures with BALF sEVs (Figure 1C & Figure S1A). An increase in %CD69$^+$ T cells (an

early activation marker), and an increase in phosphorylated Zap70 (pZap70), a kinase

downstream of the TCR (Figure 1D) was observed, suggesting an MHC Class II-TCR

interaction.

Additionally, airway sEVs from asthmatics modulated polarization of autologous

CD4$^+$ T cells into Th subsets. The increased baseline Th2 responses in asthmatics

compared to healthy controls (Figure 1E) was further enhanced following co-culture with

BALF sEVs (Figure 1E). Th17 polarization was elevated in asthmatics with an increase

in %IL-17$^+$ T cells (Figure S1B) and a robust increase in fold change of both IL-4 and IL-

17 gene expression (Figure 1F).

Modulation of transcription factors, metabolic genes, and epigenetic modifiers

underlie Th polarization[45, 46]. Principal component analyses of gene expression in T cells

from asthmatics co-cultured with or without sEVs indicated unique differences in gene

programs (Figure S1C). Heatmaps with hierarchical clustering revealed differential

modulation of metabolic genes, transcription factors, and cytokines (Figure S1D). The

top 10 differentially expressed genes included sterol pathway genes (SREBP2, HMGCR),

the glutamate transporter (*SLC7A5*), hexokinase 2 (*HK2*), Phosphoglycerate kinase 1

(*PGK1*) and histone deacetylase 1 (*HDAC1)*, in line with Th17 gene programming (Figure

1G and Figure S1D). Thus, airway sEVs from asthmatics enhance T cell proliferation,

activation, and pathogenic Th polarization.

MDRC-derived sEVs containing mitochondria are internalized by T cells via

membrane fusion

We reported that the pro-inflammatory airway MDRC-derived sEVs containing

mitochondria are internalized by peripheral autologous CD4$^+$ T cells[33]. We confirmed that

the %MitoTracker Green$^+$ (MitoT Green$^+$) CD63$^+$ BALF sEVs were not different between

healthy controls and asthmatics (Figure S2A & S2C). Although treatment of MDRCs with

mitochondrial complex I inhibitor (rotenone) and uncoupler (FCCP) did not modulate

the %MitoT Green$^+$ sEVs, evaluation with a mitochondrial membrane potential dye,

tetramethylrhodamine methyl ester (TMRE) showed that TMRE$^+$MitoT Green$^+$ BALF

sEVs were present only in asthmatics (Figure S2B-S2C). Treatment of sEVs with FCCP

or rotenone reduced the %TMRE$^+$MitoT Green$^+$CD63$^+$ sEVs suggesting that the airways

of asthmatics have an increased proportion of sEVs that maintain mitochondrial

membrane potential (Figure S2B). Transduction of purified pro-inflammatory airway

MDRCs with baculovirus constructs CellLight Mitochondria-GFP (Mito-GFP)[33] yielded

MDRC-derived GFP$^+$ sEVs which were also internalized by T cells (Figure S2D).

Since, clathrin and dynamin-dependent endocytosis pathways or micropinocytosis,

facilitate internalization of sEVs by immune and non-immune cells[47, 48], we examined if

their inhibition alters internalization of Mito-GFP$^+$ sEVs by T cells. Although T cells are

non-phagocytic, surface receptor recycling in CD4[+] T cells is clathrin-dependent[49].

Treatment of CD4[+] T cells with PitStop2 or Dynasore (inhibitors of clathrin and dynamin)

decreased membrane- and cytoplasm-associated Mito-GFP[+] sEVs only modestly (Figure

2A-2B), suggesting that endocytosis is not the primary mechanism for sEV internalization.

To assess if sEV membrane fused with recipient T cell membranes, we pre-labeled Mito-

GFP[+] sEV with PKH26; in co-cultures, PKH26 signal remained restricted to the T cell

membrane while the GFP signal was within the CD4[+] T cells (Figure 2E). The percent of

CD4[+] T cells with membrane associated and internalized signals was not different

between study groups (Figure 2C-2D). Thus, membrane fusion may be the primary mode

of internalization of sEVs in lymphocytes.

Mito-GFP[+] MDRC-derived sEVs are sufficient to trigger CD4[+] T cell activation

We transduced MDRCs with baculovirus constructs Mito-GFP and CellLight Early

Endosome-RFP (Endo-RFP) to confirm endosomal derivation of Mito-GFP[+] sEVs. We

observed two distinct populations of either Mito-GFP[+] or Endo-RFP[+] sEVs. The

expression of CD81 and CD63 (Figure S2E) and %CD63[+], CD81, or CD63[+]CD81[+] sEVs

were not different between Mito-GFP[+] and Endo-RFP[+] sEVs. Importantly, GFP and RFP

signals did not overlap within sEVs, or co-localize following internalization in T cells

(Figure 2J). Therefore, the Mito-GFP[+] sEVs may have an alternate pathway of biogenesis

than the Endo-RFP[+] sEVs and/or they follow different paths following internalization by T

cells. We investigated if internalized Mito-GFP[+] sEVs elicit activation of co-cultured

autologous peripheral CD4[+] T cells, by assessing % CD69[+] (early activation marker) and

CD154[+] (marker for antigen-specific activation) T cells.   Only CD4[+] T cells that

internalized Mito-GFP$^+$ or both Endo-RFP$^+$ and Mito-GFP$^+$ sEVs demonstrated a

significant increase in both early and antigen-specific activation signals (Figure 2F-2G &

2J). Furthermore, activation was only seen in T cells of asthmatics and not in the healthy

controls or in T cells that internalized only Endo-RFP$^+$ sEVs (Figure 2H-2I). These data

suggest that internalization of the Mito-GFP$^+$ sEVs were sufficient for promoting T cell

activation in asthmatics.

Inhibition of MHC-II blocks MDRC-derived Mito-GFP$^+$ sEV-mediated activation

We determined if MDRC-derived sEVs activate CD4$^+$ T cells through MHC-II-TCR

interaction (Figure 3A). When MDRC-sEVs were pre-treated with a pan-HLA antibody

(targets HLA-DR, HLA-DP, and HLA-DQ), and then co-cultured with autologous

peripheral CD4$^+$ T cells, the activation of T cells that had internalized Mito-GFP$^+$ (Figure

3B-3D) or both Mito-GFP$^+$ and Endo-RFP$^+$ sEVs (Figure S3A-S3D), was inhibited (Figure

3B-3D) in asthmatics. The internalization of sEVs was not abrogated (Figure 3D),

suggesting that activation was independent of internalization. When MDRC-derived

sEVs were pre-treated with LFA-1 antibody (blocks LFA-1-ICAM1 interaction), differences

in early activation (CD69) were not observed (Figure 3C); however, antigen specific

activation (CD154$^+$) and CD69$^+$CD154$^+$ populations were reduced in asthmatics

compared to sEV treatment alone.

Inhibition of mitochondrial complex III and V abrogates MDRC-derived Mito-GFP$^+$

sEV-mediated CD4$^+$ T cell activation

We then determined if the mitochondria in Mito-GFP$^+$ sEVs have functional effects

on CD4$^+$ T cell activation. Pre-treatment of MDRC-derived Mito-GFP$^+$ sEVs with

antimycin A, a high affinity complex III inhibitor (Figure 4A) before co-culture with T cells

abrogated Mito-GFP$^+$ sEV-mediated CD4$^+$ T cell activation (Figures 4B- 4D, S3C -S3D).

To control for potential transfer of mitochondrial inhibitors from the sEV preparation to

recipient cells, an sEV free isolation media with inhibitors was included which had no

effect (Figures 4B-4D, S3C-S3D). To determine if the inhibition was dependent on ATP

synthesis by sEV- mitochondria, we pre-treated MDRC-derived sEV with oligomycin, a

mitochondrial ATP synthase inhibitor. Oligomycin pre-treatment abrogated Mito-GFP$^+$

sEV-mediated activation of CD4$^+$ T cells (Figure S4A-S4D), consistent with a requirement

of a functional oxidative phosphorylation system within the sEV mitochondria to activate

T cells.

Interesingly, pre-treatment of MDRC-derived sEVs with the complex I inhibitor,

rotenone (10 µM), enhanced Mito-GFP$^+$ sEV-mediated activation of peripheral CD4$^+$ T

cells in asthmatics and importantly, induced activation of healthy T cells (Figure S5A-

S5D). This aberrant activation of CD4$^+$ T cells, short circuiting the conventional MHC-II-

TCR activation pathways, is unlikely to be solely due to effects on oxidative

phosphorylation since oligomycin, antimycin A, and rotenone inhibit mitochondrial ATP

synthesis but differ in their mechanisms. Alternatively, mitochondrial retrograde signaling

mediates these effects and this could include specific sites of mitochondrial ROS

production or accumulation of TCA substrates capable of mediating cell signaling[50].

Oligomycin, rotenone and Antimycin generate ROS through different mechanisms and

sites in the mitochondria[51, 52].

Activation of CD4$^+$ T cells requires ROS generated within MDRC-derived sEVs

Next, we determined if rotenone affects ROS levels within sEVs and if the T cell activation was dependent on sEV ROS. We used MitoTEMPOL, a superoxide dismutase mimetic to scavenge ROS within the MDRC-derived sEVs following rotenone treatment (Figure 5A). To test if blocking complex II resulted in a similar effect to rotenone or antimycin A, we used thenoyltrifluoroacetone (TTFA), a complex II inhibitor. When MDRC-derived sEVs were treated with rotenone and MitoTEMPOL, the rotenone-induced activation of CD4$^+$ T cells was inhibited (Figure 5A-5B) with a reduction in %GFP$^+$CD69$^+$CD4$^+$, GFP$^+$CD154$^+$CD4$^+$ and GFP$^+$CD69$^+$CD154$^+$CD4$^+$ T cells, specifically in CD4$^+$ T cells that have internalized Mito-GFP$^+$ or both Mito-GFP$^+$ and Endo-RFP$^+$ (Figure 5A-5B) but not just the Endo-RFP$^+$ MDRC-derived sEVs (not shown). Furthermore, treatment of MDRC-derived sEVs with TTFA did not induce CD4$^+$ T cell activation suggesting that complex II is not involved.

Since epithelial cells are the first responders to environmental cues that induce oxidative stress and influence sEV production, we investigated if their sEVs modulate CD4$^+$ T cell activation. Although CD45^{neg} cells produced Mito-GFP$^+$ exosomes and T cells internalized these exosomes, T cell activation was not observed (Figure S6A-S7D). Therefore, sEVs derived from MDRCs or other antigen presenting cells (APCs), are important in facilitating sEV-mediated T cell activation.

Mitochondrial fission is required for sEV packaging of mitochondria

We investigated the mechanisms of mitochondrial dynamics involved in packaging

mitochondria within sEVs. We first investigated mitophagy, which is shared with

endosomal sorting complexes required for transport (ESCRT) pathway[53]. We transduced

MDRCs with CellLight Mito-GFP and Lyso-RFP, a baculovirus construct that encodes

lysosomal localization of RFP protein. The Mito-GFP+ and Lyso-RFP+ signals were

mutually exclusive in the purified sEVs, indicating that Mito-GFP+ sEVs were not

generated by mitophagy (Figure 5C-5D).

Mitochondria may be packaged in sEVs by dynamin-dependent mitochondrial

fission, which is regulated by Drp1. We transduced MDRCs with CellLight Mito-GFP,

transfected them with either non-target control siRNA or siRNA targeting *DNM1L*, the

gene encoding Drp1, and confirmed knockdown of *DNM1L* by gene expression analysis

(Figure 5E right panel). The proportion of Mito-GFP+ exosomes was significantly reduced

following *DNM1L* knockdown (Figure 5E left panel & 5F). As dynamin proteins, such as

Drp1, oligomerize to function as facilitators of mitochondrial fission, we compared Drp1

oligomerization in HLA-DR+ MDRCs from healthy individuals and asthmatics. Reduced

levels of Drp1 monomers and oligomers were noted in HLA-DR+ MDRCs from healthy

individuals compared to asthmatics (Figure 5G-5I). Gene expression of DNM1L was not

significantly different comparing healthy, mild to moderate, and allergic asthmatics (Figure

5J). This enhanced mitochondrial fission in MDRCs from asthmatics may alter Drp1

oligomerization (Figure 5G) and promote packaging of mitochondria in sEVs and elicit

sEV-mediated activation in CD4+ T cells.

Mito-GFP+ MDRC-derived sEVs internalize at the site of cytoskeletal polarization and mitochondrial trafficking in CD4+ T cells

We determined if the internalization of Mito-GFP+ sEVs occurs at the CD4+ T cell immune synapse, where cytoskeletal structures polarize with the mitochondria at the site of TCR engagement. When PKH-26 labeled MDRC-derived Mito-GFP+ sEVs were co-cultured with autologous peripheral T cells, the Mito-GFP+ sEVs co-localized with both α-tubulin marking the CD4+ T cell cytoskeleton and the polarized CD4+ T cell mitochondria (Figure 6A). In CD4+ T cells labeled with both α-tubulin and Mitoview (labels mitochondria), the Mio-GFP+ sEVs co-localized with mitochondria and cytoskeleton (Figure 6B). Confocal analysis of phalloidin and 4′,6-diamidino-2-phenylindole (DAPI) labeled CD4+ T cells co-cultured with Mito-GFP+ sEVs, revealed both intracellular and membrane co-localization of the sEVs with actin (Figures 6C & S7). A z-stack slice of the confocal images indicated co-localization (yellow) of the Mito-GFP+ sEVs with cytoplasmic actin in the T cells (Figure S7 & Movie SV1). Live confocal imaging of Tubulin-RFP transduced and Mitoview-labeled T cells, co-cultured with Mito-GFP+ sEVs also showed co-localization of Mito-GFP with the polarized cytoskeleton and mitochondrial network of the recipient CD4+ T cells (Figure 6D & Movie SV2). These data suggest that sEVs are internalized at the site of the immune synapse, which may aid in the organization of the synapse through cytoskeletal and mitochondrial reorganization.

Mito-GFP+ MDRC-derived sEVs drive Th2, Th17, and Th2/17 polarization

As BALF sEVs drive Th2 and Th17 polarization (Figure 1B and S1B), we assessed if airway MDRC-derived Mito-GFP+ sEVs directly promote Th polarization. ImageStream

analyses showed IL-4 producing Th2 and IL-17 producing Th17 responses only in T cells

that have internalized Mito-GFP[+] (Figure 7A,7C, 7D, 7F & 7G), and not the Endo-RFP[+]

sEVs (Figure 7B, 7E & 7G).   Additionally, healthy MDRC-derived Mito-GFP[+] sEVs

induced IFN-γ producing Th1 responses (Figure S6E-S6G).   Th2/Th17 hybrid cells that

produce both IL-4 and IL-17 are also potent inducers of airway inflammation in

asthmatics[11, 54].   The MDRC-derived sEVs from asthmatics induced Th2/17 hybrid

responses only in T cells that internalized Mito-GFP[+] (Figure S6H, S6J & S6K), and not

Endo-RFP[+] sEVs (Figure S6I & S6K).

Mito-GFP[+] MDRC-derived sEVs may signal through mitochondrial DAMPs

Mitochondrial danger-associated molecular patterns (DAMP) signaling[55, 56] via the

sEV mitochondria may trigger T cell responses.   We first determined if T cell activation

was HLA-specific or due to allogeneic responses.   We co-cultured HLA-DR[+] MDRC-

derived sEVs from asthmatics with CD4[+] T cells from healthy subjects, and vice versa.

Modest CD69 activation without unchanged CD154 was seen in CD4[+] T cells that have

internalized Mito-GFP[+] or both Mito-GFP[+] and Endo-RFP[+] MDRC-derived sEVs (Figure

7H-7J) compared to autologous co-cultures (Figures 3, 4, S2-S5).   To investigate

potential mitochondrial DAMP signaling, MDRC-derived sEVs from asthmatics and

healthy subjects were co-cultured with a human myeloid cell line, THP-1 for 24 hours.

Pro-inflammatory cytokines increased in THP-1 cells treated with sEVs of asthmatics

(Figure 7K-7P), while healthy sEVs reduced MCP-1 levels (Figure 7L).   These results

indicate that immune activation by sEVs are partially dependent on mitochondrial DAMP

signaling.

Intranasal transfer of pro-inflammatory lung MDRC-derived sEVs with stable mitochondrial network isolated from Mito-QC donor mice with asthma enhanced overall airway inflammation and pathogenic Th responses *in vivo* in allergic recipients

To ascertain that it is not mitophagy-mediated shuttling of mitochondria in sEVs that elicit pathogenic Th responses in vivo, we performed intranasal transfer of lung MDRC-derived sEVs isolated from Mito-QC asthmatic or OVA sensitized donor mice. Quenched GFP and stable mCherry signal marks mitophagy in Mito-QC mice. We first observed that the intranasal transfer of MDRC-derived sEVs from asthmatic or OVA sensitized Mito-QC donor mice enhanced overall inflammatory cell infiltration in the lungs, airways and lung draining lymph nodes of asthmatic recipient mice compared to OVA sensitized controls (C57BL/6) (Figure S8A-S8C). Followng transfer of MDRC-derived sEVs, OVA-specific IgE in both BAL and serum was increased along with levels of Muc5AC in asthmatic recipient mice compared to OVA sensitized controls (Figure S8D-S8F). MDRC-derived sEVs isolated from OVA sensitized controls elicited inflammatory responses of much lower magnitude when transferred to OVA challenged mice and little to no response in OVA-sensitized mice (Figure S8A-S8F). BALF differential also showed increased airway infiltration of eosinophils, neutrophils, macrophages and lymphocytes in asthmatic recipient mice of MDRC-derived sEVs (Figure S8G- S8J). The absolute numbers and frequencies of lung MDRCs including Ly6C$^+$, Ly6G$^+$ and Ly6C$^+$Ly6G$^+$ subpopulations as well as neutrophils and eosinophils also showed similar mhigher levels in asthmatic recipient mice following sEV transfer (Figures 8A-8E, S9A-S9E). Importantly, the absolute numbers and frequencies of Th2, Th17 cells (Figures 8F-8G, S9F-S9G) as well as Type

2 innate lymphoid cells (Figures 8I, S9I) also increased in asthmatic recipient mice

following exosome transfer compared to relevant control; the absolute numbers and

frequency of Th1 cells declined in mice with asthma (Figures 8H, S9H).

We then confirmed that the activation of CD4$^+$ T cells are elicited by internalization

of Mito-QC$^+$ sEVs *in vivo* during allergic airway inflammation. The oligomerization of Drp1

was enhanced in pro-inflammatory MDRCs sorted from OVA challenged Mito-QC mice

compared to sensitized controls (Figure 9A). The frequency of CD81$^+$MHCII$^+$GFP$^+$ sEVs

produced by pro-inflammatory MDRCs sorted from OVA challenged Mito-QC mice was

higher compared to those from sensitized controls (Figure 9B-9C). Image Stream

analyses confirmed the characterization of GFP$^+$MHC-II$^+$CD81$^+$ sEVs isolated from

proinflammatory Ly6G$^+$ MDRCs (Figure 9C). Further, ImageStream analyses showed that

lung-derived CD45$^+$CD4$^+$CD69$^+$ activated T cells were also GFP$^+$ in transfer recipient

asthmatic mice compared to controls (Figure 9D). Correlation analyses showed that

CD69 MFI correlated with % GFP$^+$ CD4$^+$ T cells as well as % GFP area within T cells of

asthmatic recipient mice (Figure 9E-F).

We then investigated if intranasal transfer of these MDRC-derived sEVs from OVA

sensitized or challenged donor Mito-QC mice to sensitized or challenged recipients would

enhance pathogenic Th responses and exacerbate allergic airway inflammation *in vivo* in

recipients. As stable mitochondrial network without mitophagy is marked by binary signal

of both mCherry and GFP in Mito-QC mice, we evaluated percentages and absolute

numbers of mCherry$^+$GFP$^+$ Th2 and Th17 cells and found that both were increased in

lung tissues, BALF as well as lymph nodes (Figures 10A-10G, S10-S12) of asthmatic

recipient mice compared to control recipients. These data comfirmed that stable

mitochondrial network was transferred *in vivo* via sEVs to CD4[+] T cell subsets to promote

Th polarization. Additionally, increased proportion and absolute numbers of

mCherry[+]GFP[+] ILC2s were also noted in BALF of recipient mice with asthma compared

to sensitized controls (Figure 10G, S12G).

Discussion

Small extracellular vesicles (sEVs) are novel intercellular messengers, both in

homeostasis and disease. Our observations extend the role for sEVs as immune

modulators in human asthma. We report that airway sEVs from asthmatics drive

proliferation of Th cells and their polarization to Th2 and Th17 phenotypes. Accompanying

the changes in epigenetic programming, the airway HLA-DR[+] MDRC-derived

immunogenic sEVs containing mitochondria elicit antigen-specific activation that requires

mitochondrial ROS signaling through the transfer of sEV mitochondria to recipient T cells;

this transfer involves internalization of sEVs by membrane fusion that initiates TCR

engagement/activation in CD4[+] T cells at the site of the immune synapse. We

demonstrate that *in vivo* transfer of stable mitochondrial network carrying sEVs can

enhance *in vivo* pathogenic Th polarization in allergic recipients.

CD4[+] T cells use clathrin-dependent mechanisms to recycle cell surface receptors

[49]; however, transfer of the PKH26 signal from labeled sEVs to CD4[+] T cell membranes,

internalization of the mitochondrial GFP signal and the preserved sEV internalization by

CD4[+] T cells in the presence of inhibitors of clathrin or dynamin, or LFA-1 antibody,

suggest that membrane fusion is the primary mechanism of internalization of

mitochondria-containing sEVs. Although CD4[+] T cell activation correlated with Mito-GFP[+]

sEV internalization, blockade of class II or LFA-1 on sEVs failed to inhibit their

internalization, but attenuated activation of recipient CD4[+] T cells. LFA-1 helps bind T

cells to APCs via ICAM-1 and ICAM-2, which aids in antigen presentation[57] and CD8[+] T

cells utilize LFA-1 to internalize sEVs[58]. Synergistic signaling between class II and LFA-

1 may facilitate MDRC-derived sEV-mediated antigen-specific T cell activation without

APCs.

Our studies indicate that mitochondria within sEVs are bioenergetically functional

and required for sEV-mediated TCR activation. Inhibition of mitochondrial complex II,

complex III, and complex V abrogated Mito-GFP[+] sEV-mediated T cell responses.

Interestingly, pre-treatment of MDRC-derived sEVs with complex I inhibitor, rotenone,

aberrantly activated healthy CD69[+] T cells and promoted antigen specific activation in

asthmatics. Since these mitochondrial inhibitors block oxidative phosphorylation, this

aberrant activation could be ATP-dependent. However, mitochondria-targeted antioxidant,

mitoTEMPOL, scavenged the ROS within MDRC-derived sEVs after rotenone treatment;

implicating ROS in the sEV-dependent activation of healthy T cells. Rotenone-induced

activation may be due to its inhibitory effect on reverse electron transport to suppress T

cell activation, or due to rotenone-mediated ROS production altering normal T cell

function[50, 52, 59, 60]. High rotenone concentrations may cause off target effects;

mechanisms for rotenone-dependent effects needs to be further clarified[61]. Despite these

caveats, our studies implicate mitochondrial ROS as an essential mediator of T cell

activation. T cells produce ROS via complex I after TCR engagement[62], and complex I

is important for T helper differentiation[63]. Retrograde signaling is involved, as Th

transcriptional program is activated in T cells by mitochondrial signaling via MDRC-sEVs.

Taken together, our studies suggest that CD4[+] T cell-activation by sEVs in the context of

asthma or other chronic inflammatory diseases may depend on both TCR-MHC

engagement and transfer of functional mitochondria capable of generating ROS.

We interrogated mechanisms underlying mitochondrial packaging within airway

immune cell-derived sEVs. Mesenchymal stem cells package mitochondria in EVs to

outsource mitophagy [36]. Absence of co-localization of the packaged Mito-GFP[+] signal

with Lyso-RFP[+] associated with mitophagy demonstrates a novel role for MDRC-derived

exosomes in mitochondrial retrograde signaling. Although TLR-mediated activation of

immune cells by mitophagy is possible[64], our results suggest that antigen presentation

and mitochondrial ROS signaling are the primary mechanisms of immune activation. In

addition to ROS, mitochondrial DAMP signaling may account for modest T cell activation,

but not the significant early and antigen-specific activation. Circulating mitochondrial

DAMPs activate neutrophils through TLR- 9[56] and CD4[+] T cells are reactive to

mitochondrial antigens in autoimmune diseases. However, direct activation of CD4[+] T

cells by DAMPs has not been established.

Although transfer of mitochondria by sEVs has been reported, mechanisms of

packaging mitochondria within airway immune cell-derived sEVs are largely unknown.

We provide new insights for mitochondrial fission in the generation of sEVs that transport

functional mitochondria. Our observations that intranasal transfer of Mito-QC mice-

derived sEVs with binary signal of mCherry and GFP results in exacerbation of allergic

airway inflammation and enhancement of pathogenic Th polarization in allergic models

provides *in vivo* evidence for the significance of sEV mitochondria in asthma. Drp1-

dependent mitochondrial fission requires its oligomerization[65]; here, Drp1 knockdown

reduced Mito-GFP[+] sEVs. Mitochondrial fragmentation alters immune function in cancer;

in liver cancers, this limited the function of NK cells[66]. In our studies, mitochondrial fission

generates sEVs with mitochondria that facilitates sEV-mediated immune activation, thus

providing novel insights for mitochondria fission in immune regulation. The extent of sEV

signaling is different between healthy controls and asthmatics. Genetic differences are,

so far not known to contribute to differences in proinflammatory MDRCs reported in

asthmatics. Epigenetic modifications from lung damage/oxidative stress may alter

MDRC-derived sEVs in asthmatics and/or differential presence/presentation of antigens

by sEVs and/or account for the increased mitochondrial fission noted in MDRCs from

asthmatics; this differential response may not be restricted to asthma and may be

important in COPD or sarcoidosis.

Additionally, we demonstrate that Mito-GFP[+] sEV-mediated signaling and

internalization occurs at the site of synapse formation in activated T cells, which provides

an antigen-specific dimension to this cellular crosstalk. Our observation that polarized

cytoskeleton and the mitochondria of the activated CD4[+] T cell co-localizes with sEV

mitochondria from donor MDRCs suggest that upon TCR engagement by the MHC-II on

sEVs, reorganization of the immune synapse occurs at the site of sEV internalization.

Here, we uncover a novel mechanism by which sEVs activate CD4[+] T cells, and

mitochondrial transfer may function as a second signal in CD4[+] T cell activation.

Traditionally, T cell activation/polarization requires antigen presentation by APCs, co-

stimulation and an optimal cytokine milieu[67-69]. Our studies support an acellular form of

T cell activation by sEV mitochondria and a new mechanism of MDRC-T cell crosstalk *in*

vitro and *in vivo* at the intersection of bioenergetics and cellular function. We propose

that these intricate immunological mechanisms facilitated by sEVs participate in both

normal physiological responses and may be dysregulated in pathological processes such

as human asthma.

Methods:

Please see supplementary information

Acknowledgements

We acknowledge Marion Spell at the UAB Flow Cytometry Core Facilities, Shawn

Williams and Robert Grabski at the High Resolution Imaging Facility and Sagar

Hanumanthu at the Shelby Comprehensive Flow Cytometry Core Facility for technical

assistance in exosome flow cytometry, confocal microscopy, and ImageStream flow

cytometry, respectively.

Declaration of Interests

SG holds a patent on B cell derived exosomes in immune therapy, and is part of the

Scientific Advisory Board of Anjarium Biosciences. All other authors declare no conflicts

of interest.

REFERENCES

1. Medoff, B.D., Thomas, S.Y. & Luster, A.D. T cell trafficking in allergic asthma: the
ins and outs. *Annual review of immunology* **26**, 205-232 (2008).
2. Barnes, P.J. Th2 cytokines and asthma: an introduction. *Respir Res* **2**, 64-65
(2001).
3. Newcomb, D.C. & Peebles, R.S., Jr. Th17-mediated inflammation in asthma. *Curr*
Opin Immunol **25**, 755-760 (2013).

4. Djukanovic, R. *et al.* Mucosal inflammation in asthma. *The American review of*
respiratory disease **142**, 434-457 (1990).
5. Randolph, D.A., Stephens, R., Carruthers, C.J. & Chaplin, D.D. Cooperation
between Th1 and Th2 cells in a murine model of eosinophilic airway inflammation.
J Clin Invest **104**, 1021-1029 (1999).
6. Steinke, J.W. & Borish, L. Th2 cytokines and asthma. Interleukin-4: its role in the
pathogenesis of asthma, and targeting it for asthma treatment with interleukin-4
receptor antagonists. *Respir Res* **2**, 66-70 (2001).
7. Wills-Karp, M. The gene encoding interleukin-13: a susceptibility locus for asthma
and related traits. *Respir Res* **1**, 19-23 (2000).
8. Hogan, S.P. *et al.* Eosinophils: biological properties and role in health and disease.
Clin Exp Allergy **38**, 709-750 (2008).
9. Wenzel, S.E. Asthma: defining of the persistent adult phenotypes. *Lancet* **368**,
804-813 (2006).
10. Korn, T., Bettelli, E., Oukka, M. & Kuchroo, V.K. IL-17 and Th17 Cells. *Annu Rev*
Immunol **27**, 485-517 (2009).
11. Liu, W. *et al.* Mechanism of TH2/TH17-predominant and neutrophilic TH2/TH17-
low subtypes of asthma. *J Allergy Clin Immunol* **139**, 1548-1558 e1544 (2017).
12. Cella, M., Sallusto, F. & Lanzavecchia, A. Origin, maturation and antigen
presenting function of dendritic cells. *Curr Opin Immunol* **9**, 10-16 (1997).
13. Steinman, R.M. & Swanson, J. The endocytic activity of dendritic cells. *J Exp Med*
182, 283-288 (1995).
14. Cella, M., Engering, A., Pinet, V., Pieters, J. & Lanzavecchia, A. Inflammatory
stimuli induce accumulation of MHC class II complexes on dendritic cells. *Nature*
388, 782-787 (1997).
15. Wange, R.L. & Samelson, L.E. Complex complexes: signaling at the TCR.
Immunity **5**, 197-205 (1996).
16. Lu, T. & Gabrilovich, D.I. Molecular pathways: tumor-infiltrating myeloid cells and
reactive oxygen species in regulation of tumor microenvironment. *Clin Cancer Res*
18, 4877-4882 (2012).
17. Dumortier, H. *et al.* Antigen presentation by an immature myeloid dendritic cell line
does not cause CTL deletion in vivo, but generates CD8+ central memory-like T
cells that can be rescued for full effector function. *J Immunol* **175**, 855-863 (2005).
18. Santambrogio, L. *et al.* Extracellular antigen processing and presentation by
immature dendritic cells. *Proc Natl Acad Sci U S A* **96**, 15056-15061 (1999).
19. Holgate, S.T. Innate and adaptive immune responses in asthma. *Nat Med* **18**, 673-
683 (2012).
20. Lambrecht, B.N. & Hammad, H. Taking our breath away: dendritic cells in the
pathogenesis of asthma. *Nat Rev Immunol* **3**, 994-1003 (2003).
21. Deshane, J.S. *et al.* Subsets of airway myeloid-derived regulatory cells distinguish
mild asthma from chronic obstructive pulmonary disease. *J Allergy Clin Immunol*
135, 413-424 e415 (2015).
22. Wahlund, C.J.E. *et al.* Exosomes from antigen-pulsed dendritic cells induce
stronger antigen-specific immune responses than microvesicles in vivo. *Sci Rep* **7**,
17095 (2017).

23. Admyre, C., Johansson, S.M., Paulie, S. & Gabrielsson, S. Direct exosome
stimulation of peripheral human T cells detected by ELISPOT. *Eur J Immunol* **36**,
1772-1781 (2006).
24. Admyre, C. *et al.* B cell-derived exosomes can present allergen peptides and
activate allergen-specific T cells to proliferate and produce TH2-like cytokines. *J*
Allergy Clin Immunol **120**, 1418-1424 (2007).
25. Raposo, G. *et al.* B lymphocytes secrete antigen-presenting vesicles. *J Exp Med*
183, 1161-1172 (1996).
26. El Andaloussi, S., Lakhal, S., Mager, I. & Wood, M.J. Exosomes for targeted siRNA
delivery across biological barriers. *Adv Drug Deliv Rev* **65**, 391-397 (2013).
27. Dai, S. *et al.* Phase I clinical trial of autologous ascites-derived exosomes
combined with GM-CSF for colorectal cancer. *Mol Ther* **16**, 782-790 (2008).
28. Andre, F. *et al.* Exosomes as potent cell-free peptide-based vaccine. I. Dendritic
cell-derived exosomes transfer functional MHC class I/peptide complexes to
dendritic cells. *J Immunol* **172**, 2126-2136 (2004).
29. Andreu, Z. & Yanez-Mo, M. Tetraspanins in extracellular vesicle formation and
function. *Front Immunol* **5**, 442 (2014).
30. Rocha-Perugini, V., Sanchez-Madrid, F. & Martinez Del Hoyo, G. Function and
Dynamics of Tetraspanins during Antigen Recognition and Immunological
Synapse Formation. *Front Immunol* **6**, 653 (2015).
31. Petersen, S.H. *et al.* The role of tetraspanin CD63 in antigen presentation via MHC
class II. *Eur J Immunol* **41**, 2556-2561 (2011).
32. Denzer, K. *et al.* Follicular dendritic cells carry MHC class II-expressing
microvesicles at their surface. *J Immunol* **165**, 1259-1265 (2000).
33. Hough, K.P. *et al.* Exosomal transfer of mitochondria from airway myeloid-derived
regulatory cells to T cells. *Redox Biol* **18**, 54-64 (2018).
34. Morrison, T.J. *et al.* Mesenchymal Stromal Cells Modulate Macrophages in
Clinically Relevant Lung Injury Models by Extracellular Vesicle Mitochondrial
Transfer. *Am J Respir Crit Care Med* **196**, 1275-1286 (2017).
35. Hayakawa, K. *et al.* Transfer of mitochondria from astrocytes to neurons after
stroke. *Nature* **535**, 551-555 (2016).
36. Phinney, D.G. *et al.* Mesenchymal stem cells use extracellular vesicles to
outsource mitophagy and shuttle microRNAs. *Nat Commun* **6**, 8472 (2015).
37. Islam, M.N. *et al.* Mitochondrial transfer from bone-marrow-derived stromal cells
to pulmonary alveoli protects against acute lung injury. *Nat Med* **18**, 759-765
(2012).
38. Mahrouf-Yorgov, M. *et al.* Mesenchymal stem cells sense mitochondria released
from damaged cells as danger signals to activate their rescue properties. *Cell*
Death Differ **24**, 1224-1238 (2017).
39. Kiriyama, Y. & Nochi, H. Intra- and Intercellular Quality Control Mechanisms of
Mitochondria. *Cells* **7** (2017).
40. Spees, J.L., Olson, S.D., Whitney, M.J. & Prockop, D.J. Mitochondrial transfer
between cells can rescue aerobic respiration. *Proc Natl Acad Sci U S A* **103**, 1283-
1288 (2006).

41. Hill, B.G., Shiva, S., Ballinger, S., Zhang, J. & Darley-Usmar, V.M. Bioenergetics and translational metabolism: implications for genetics, physiology and precision medicine. *Biol Chem* **401**, 3-29 (2019).

42. Admyre, C. *et al.* Exosomes with major histocompatibility complex class II and co-stimulatory molecules are present in human BAL fluid. *Eur Respir J* **22**, 578-583 (2003).

43. Torregrosa Paredes, P. *et al.* Bronchoalveolar lavage fluid exosomes contribute to cytokine and leukotriene production in allergic asthma. *Allergy* **67**, 911-919 (2012).

44. Hough, K.P. *et al.* Unique Lipid Signatures of Extracellular Vesicles from the Airways of Asthmatics. *Sci Rep* **8**, 10340 (2018).

45. Zhu, J., Yamane, H. & Paul, W.E. Differentiation of effector CD4 T cell populations (*). *Annual review of immunology* **28**, 445-489 (2010).

46. Hirahara, K. *et al.* Helper T-cell differentiation and plasticity: insights from epigenetics. *Immunology* **134**, 235-245 (2011).

47. Tian, T. *et al.* Exosome uptake through clathrin-mediated endocytosis and macropinocytosis and mediating miR-21 delivery. *J Biol Chem* **289**, 22258-22267 (2014).

48. Schneider, D.J. *et al.* Mechanisms and modulation of microvesicle uptake in a model of alveolar cell communication. *J Biol Chem* **292**, 20897-20910 (2017).

49. Compeer, E.B. *et al.* A mobile endocytic network connects clathrin-independent receptor endocytosis to recycling and promotes T cell activation. *Nat Commun* **9**, 1597 (2018).

50. Chouchani, E.T. *et al.* Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* **515**, 431-435 (2014).

51. Lou, P.H. *et al.* Mitochondrial uncouplers with an extraordinary dynamic range. *Biochem J* **407**, 129-140 (2007).

52. Quinlan, C.L., Perevoshchikova, I.V., Hey-Mogensen, M., Orr, A.L. & Brand, M.D. Sites of reactive oxygen species generation by mitochondria oxidizing different substrates. *Redox Biol* **1**, 304-312 (2013).

53. Takahashi, Y. *et al.* An autophagy assay reveals the ESCRT-III component CHMP2A as a regulator of phagophore closure. *Nat Commun* **9**, 2855 (2018).

54. Irvin, C. *et al.* Increased frequency of dual-positive TH2/TH17 cells in bronchoalveolar lavage fluid characterizes a population of patients with severe asthma. *J Allergy Clin Immunol* **134**, 1175-1186 e1177 (2014).

55. Oka, T. *et al.* Mitochondrial DNA that escapes from autophagy causes inflammation and heart failure. *Nature* **485**, 251-255 (2012).

56. Zhang, Q. *et al.* Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature* **464**, 104-107 (2010).

57. Garcon, F. & Okkenhaug, K. PI3Kdelta promotes CD4(+) T-cell interactions with antigen-presenting cells by increasing LFA-1 binding to ICAM-1. *Immunol Cell Biol* **94**, 486-495 (2016).

58. Nolte-'t Hoen, E.N., Buschow, S.I., Anderton, S.M., Stoorvogel, W. & Wauben, M.H. Activated T cells recruit exosomes secreted by dendritic cells via LFA-1. *Blood* **113**, 1977-1981 (2009).

59. Rustin, P. & Lance, C. Succinate-driven reverse electron transport in the
respiratory chain of plant mitochondria. The effects of rotenone and adenylates in
relation to malate and oxaloacetate metabolism. *Biochem J* **274 (Pt 1)**, 249-255
(1991).
60. Scialo, F., Fernandez-Ayala, D.J. & Sanz, A. Role of Mitochondrial Reverse
Electron Transport in ROS Signaling: Potential Roles in Health and Disease. *Front*
Physiol **8**, 428 (2017).
61. Grefte, S., Wagenaars, J.A., Jansen, R., Willems, P.H. & Koopman, W.J.
Rotenone inhibits primary murine myotube formation via Raf-1 and ROCK2.
Biochim Biophys Acta **1853**, 1606-1614 (2015).
62. Akkaya, B. *et al.* Increased Mitochondrial Biogenesis and Reactive Oxygen
Species Production Accompany Prolonged CD4(+) T Cell Activation. *J Immunol*
201, 3294-3306 (2018).
63. Ozay, E.I. *et al.* Rotenone Treatment Reveals a Role for Electron Transport
Complex I in the Subcellular Localization of Key Transcriptional Regulators During
T Helper Cell Differentiation. *Front Immunol* **9**, 1284 (2018).
64. Gkikas, I., Palikaras, K. & Tavernarakis, N. The Role of Mitophagy in Innate
Immunity. *Front Immunol* **9**, 1283 (2018).
65. Cribbs, J.T. & Strack, S. Reversible phosphorylation of Drp1 by cyclic AMP-
dependent protein kinase and calcineurin regulates mitochondrial fission and cell
death. *EMBO Rep* **8**, 939-944 (2007).
66. Zheng, X. *et al.* Mitochondrial fragmentation limits NK cell-based tumor
immunosurveillance. *Nat Immunol* **20**, 1656-1667 (2019).
67. Itano, A.A. & Jenkins, M.K. Antigen presentation to naive CD4 T cells in the lymph
node. *Nat Immunol* **4**, 733-739 (2003).
68. Koulova, L., Clark, E.A., Shu, G. & Dupont, B. The CD28 ligand B7/BB1 provides
costimulatory signal for alloactivation of CD4+ T cells. *J Exp Med* **173**, 759-762
(1991).
69. Curtsinger, J.M. & Mescher, M.F. Inflammatory cytokines as a third signal for T cell
activation. *Curr Opin Immunol* **22**, 333-340 (2010).

Figure Legends

Figure 1 – BALF sEVs from the airways of asthmatics promote proliferation and activation of CD4$^+$ T cells, alters transcriptional signature and enhances T helper cell polarization. (A-B) Percentage of proliferating peripheral CD4$^+$ T cells after co-culture with airway sEVs. Small extracellular vesicles (sEVs) from the BAL fluid (BALF) were co-cultured with CFSE labeled autologous peripheral CD4$^+$ T cells at a ratio of 1:10 (T cells:sEVs) in the presence of rhIL-2 (50 IU/ml) for 7-days. Cells were harvested at day 7 for flow cytometry analysis on the BD LSRII. Mann Whitney T test for comparison between healthy and asthmatics, Wilcoxon matched-pairs signed rank test for intergroup (no sEVs vs sEVs), **<0.01, ***<0.001, ****<0.0001. (C) Percentage of proliferating airway CD4$^+$ T cells after co-culture with airway sEVs. sEVs from the BALF were co-cultured as described above with autologous airway CD4$^+$ T cells. Cells were harvested at day 7 for flow cytometry analysis on the BD LSRII. Mann Whitney T test for comparison between healthy and asthmatics, Wilcoxon matched-pairs signed rank test for intergroup (no sEVs vs sEVs), **<0.01, ***<0.001, ****<0.0001. (D) BALF sEVs were co-cultured with autologous CD4$^+$ T cells (1:10 T cells:sEVs) for 15 minutes and percent CD69 and phospho-Zap70 was assessed by ImageStream imaging flow cytometry (representative image). Mann Whitney T test, *<0.05. (E) sEVs purified from the BALF were co-cultured with autologous peripheral CD4$^+$ T cells in a ratio of 1:10 (T cells: sEVs) in the presence of rhIL-2 (50 IU/ml) for 7 days. Cells were harvested at day 7 for intracellular staining of cytokines and analyzed by flow cytometry using a BD LSRII. Left, percent of Th2 cells (assessed by IL-4 expression), and representative flow plots shown on the right. Mann Whitney T test for comparison between healthy and asthmatics, Wilcoxon matched-pairs signed rank test for intergroup (no sEVs vs sEVs), **<0.01, ****<0.0001. (F) Quantitative real-time PCR analysis of IL-4 and IL-17 gene expression of healthy and asthmatic CD4$^+$ T cells co-cultured with and without sEVs for 7 days. Data are normalized to gene expression in T cells cultured without sEVs (ACTB as reference gene). Co-culture of asthmatic sEVs with T cells significantly increased IL-4 and IL-17 gene expression (**<0.01, ****<0.0001). (E) Clustered heatmap generated using MetaboAnalyst 3.0 from a custom NanoString panel comparing gene expression of CD4$^+$ T cells from asthmatics co-cultured with BALF sEVs (right), and no sEVs control (left). Heatmap and clustering were generated using the nSolver Analysis Software 4.0 from nanoString.

Figure 2 – CD4$^+$ T cells internalize MDRC-derived sEVs predominantly by membrane fusion and not by endocytosis. Activation of CD4$^+$ T cells is associated with internalization of Mito-GFP$^+$ sEVs derived from airway MDRCs. (A-B) Blockade of clathrin dependent and independent mechanism of endocytosis did not completely abrogate MDRC-derived Mito-GFP$^+$ sEV internalization by CD4$^+$ T cells. CD4$^+$ T cells were treated with PitStop2 (50 nM), Dynasore (50 µM), or a combination of the two prior to co-culture with MDRC-derived sEVs for 24 hours and internalization assessed by ImageStream flow cytometry. (A) Quantitation of number of CD4$^+$ T cells with membrane associated Mito-GFP$^+$ MDRC sEVs. (B) Quantitation of number of CD4$^+$ T cells with internalized Mito-GFP$^+$ MDRC sEVs. (A-B) Mann Whitney T, *<0.05, **<0.01, ***<0.001, ****<0.0001. (C-E) MDRCs were transduced with Mito-GFP and membrane labeled with PKH26 red. sEVs derived from MDRCs were purified 48 hours later. Purified MDRC-derived sEVs were co-cultured with autologous peripheral CD4$^+$ T cells for 24 hours in the presence of rhIL-2 (50 IU/ml) in a ration of 1:10 T cells:sEVs. Association of sEVs with the membrane or intracellular localization of sEVs were assessed by ImageStream flow cytometry. (C) Quantitation of membrane associated Mito-GFP$^+$ MDRC sEVs. (D) Quantitation of intracellularly associated Mito-GFP$^+$ MDRC sEVs. (E) Representative image strips from ImageStream flow cytometry analysis illustrating association of PKH26 red with the membrane and intracellular or membrane association of Mito-GFP. (F-J) MDRCs were transduced with CellLight Mito-GFP and CellLight Endo-RFP, and sEVs were purified from the supernatant 48 hours later. Autologous peripheral human T cells were cultured with MDRC sEVs for 24 hours, and internalization and activation assessed by ImageStream imaging flow cytometry. An sEV -generator-cell (MDRCs) to T cell ratio of 2.5x10^5:1x10^6 was maintained. (F-G) Percent of GFP$^+$CD69$^+$CD4$^+$ and GFP$^+$CD154$^+$CD4$^+$ T cells that represent T cells that have internalized GFP$^+$ sEVs and show early activation and antigen specific activation. (H-I) Percent of RFP$^+$CD69$^+$CD4$^+$ and RFP$^+$CD154$^+$CD4$^+$ T cells that represent T cells that have internalized RFP$^+$ sEVs and show early activation and antigen specific activation (F-G) Mann Whitney T, **<0.01. (J) Representative images from ImageStream illustrating MDRC sEV internalization and T cell activation (CD69 and CD154).

Figure 3 – Blockade of class II molecules using a pan-HLA antibody (HLA-DR/DP/DQ) results in loss of MDRC sEV mediated activation of CD4$^+$ T cells. (A) Illustration of experimental approach. (B-D) MDRCs were transduced with CellLight Mito-GFP and CellLight Endo-RFP, and sEVs were purified from the supernatant 48 hours later. sEVs derived from MDRCs transduced with Mito-GFP and Endo-RFP were co-cultured with autologous peripheral CD4$^+$ T cells for 24 hours in the presence of rhIL-2 (50 IU/ml) in a ration of 1:10 T cells:sEVs. MDRC sEVs were pre-treated with pan-HLA-Ab (10 µg/ml), LFA2 antibody (1 µg/ml), IgG control, a combination of pan-HLA-Ab and anti-LFA2 for 30 minutes at room temperature. sEVs were washed and re-purified using the Invitrogen Total Exosome Isolation kit. ImageStream flow cytometry was used to assess early activation (CD69) and antigen-specific activation (CD154) marker expression. (B) Venn diagrams illustrating the percentages for CD154$^+$, CD69$^+$, or CD154$^+$CD69$^+$ T cells that internalized either Mito-GFP$^+$ MDRC sEVs or Endo-RFP$^+$ MDRC sEVs. The top row is untreated MDRC sEVs, and the bottom row is MDRC sEVs pre-treated with pan-HLA-Ab prior to co-culture. (C) Graphs illustrating the percentage of CD69$^+$, CD154$^+$, or CD69$^+$CD154$^+$ T cells that internalized either Mito-GFP$^+$ MDRC sEVs (top row) or Endo-RFP$^+$ MDRC sEVs (bottom row). Mann Whitney T, *<0.05, **<0.01, ***<0.001, ****<0.0001. (D) Representative image strips from ImageStream illustrating that Mito-GFP$^+$ MDRC sEVs activate T cells in asthmatics, while treatment with pan-HLA-Ab abrogates this activation by MDRC sEVs.

Figure 4 – Inhibition of complex III by antimycin A in MDRC sEVs diminishes CD4$^+$ T cell
activation in asthmatics. (A) Illustration of experimental approach. (B-D) MDRCs were
transduced with CellLight Mito-GFP and CellLight Endo-RFP, and sEVs were purified
from the supernatant 48 hours later. sEVs derived from MDRCs transduced with Mito-
GFP and Endo-RFP were pre-treated with antimycin A (10 µM) overnight or untreated
prior to co-culture with autologous peripheral CD4$^+$ T cells for 24 hours in the presence
of rhIL-2 (50 IU/ml) in a ration of 1:10 T cells:sEVs. sEVs were washed and re-purified
using the Invitrogen Total Exosome Isolation kit. ImageStream flow cytometry was used
to assess early activation (CD69) and antigen-specific activation (CD154). (B) Venn
diagrams illustrating the percentages for CD154$^+$, CD69$^+$, or CD154$^+$CD69$^+$ T cells that
internalized either Mito-GFP$^+$ MDRC sEVs or Endo-RFP$^+$ MDRC sEVs. (C) Graphs
illustrating the percentage of CD69$^+$, CD154$^+$, or CD69$^+$CD154$^+$ T cells that internalized
either Mito-GFP$^+$ MDRC sEVs (top row) or Endo-RFP$^+$ MDRC sEVs (bottom row). Mann
Whitney T, **<0.01, ***<0.001, ****<0.0001. (D) Representative image strips from
ImageStream analysis illustrating that Mito-GFP$^+$ MDRC sEVs activate T cells in
asthmatics, and antimycin A blocks activation in asthmatics.

Figure 5 – Reduction of mitochondrial ROS within sEVs by MitoTEMPOL inhibits activation of CD4$^+$ T cells by MDRC-derived Mito-GFP$^+$ sEVs from asthmatics. Packaging of mitochondria inside MDRC-derived sEVs is mediated by mitochondrial fission. (A-B) MDRCs were transduced with CellLight Mito-GFP and CellLight Endo-RFP, and exosomes were purified from the supernatant 48 hours later. sEVs derived from MDRCs transduced with Mito-GFP and Endo-RFP were pre-treated with rotenone (10 nM), MitoTEMPOL (1 µM), rotenone + MitoTEMPOL, thenoyltrifluoroacetone (TTFA, 10 µM) overnight, or untreated prior to co-culture. sEVs were washed and re-purified using the Invitrogen Total Exosome Isolation kit. MDRC-derived sEVs were co-culturedwith autologous peripheral CD4$^+$ T cells for 24 hours in the presence of rhIL-2 (50 IU/ml) in a ration of 1:10 T cells:sEVs. ImageStream flow cytometry was used to assess early activation (CD69) and antigen-specific activation (CD154). (A-B) Graphs illustrating the percentage of CD69$^+$, CD154$^+$, or CD69$^+$CD154$^+$ T cells that internalized (A) Mito-GFP$^+$ MDRC sEVs, (B) both Mito-GFP$^+$ and Endo-RFP$^+$ MDRC sEVs. Mann Whitney T, *<0.05, **<0.01, ***<0.001, ****<0.0001. (C-D) MDRCs were transduced with CellLight Mito-GFP and CellLight Lyso-RFP, and sEVs were purified from the supernatant 48 hours later. Purified sEVs were characterized using ImageStream flow cytometry to assess if the Mito-GFP and Lyso-RFP co-localized. (C) Quantitation of CD63$^+$ MDRC sEVs that are Mito-GFP$^+$Lyso-RFP$^-$, Mito-GFP$^-$Lyso-RFP$^+$, or Mito-GFP$^+$Lyso-RFP$^+$ (D) Representative image strips from ImageStream analysis illustrating Mito-GFP$^+$Lyso-RFP$^-$ MDRC sEVs, and Mito-GFP$^-$Lyso-RFP$^+$ MDRC sEVs. (E-F) MDRCs were transfected with either non-target (NT) control siRNA, or siRNA that targets DNM1L using lipofectamine 3000, and cultured for 48 hours in 10% human serum media that has been depleted of EVs including sEVs. Culture supernatant was isolated for sEV isolation, and cells for gene expression analysis of *DNM1L* (*ACTB* as reference). (E) Left, quantitation of percent CD63$^+$Mito-GFP$^+$ MDRC sEVs. Right, fold-change of *DNM1L* (Drp1) expression in transfected MDRCs to confirm siRNA efficacy. Mann Whitney T test, *<0.05, ***<0.001. (F) Representative flow plots illustrating reduction of CD63$^+$Mito-GFP$^+$ MDRC sEVs after transection with siRNA for *DNM1L*. (G) Protein level and oligomerization of Drp1 was assessed in HLA-DR$^+$ MDRCs sorted from the BAL of healthy and asthmatics by Native-PAGE with β-actin as control. (H) Quantitation of Drp1 monomer ratio in samples from asthmatics to healthy. (I) Quantitation of the Drp1 oligomer ratio in samples from asthmatics to healthy. (J) Quantitation of real-time PCR analysis of *DNM1L* gene expression in purified HLA-enriched peripheral myeloid cells from healthy, mild-to-moderate asthmatics, and allergic asthmatics (house dust mite or fungal).

Figure 6 – Internalization of Mito-GFP⁺ MDRC sEVs occurs at site of tubulin, actin, and
mitochondrial polarization in T cells. (A) Representative images from ImageStream flow
cytometry analysis illustrating polarization of α-tubulin at the site of internalization of Mito-
GFP⁺ sEVs. MDRCs were transduced with CellLight Mito-GFP and sEVs were purified
from the supernatants 48 hours later. Mito-GFP⁺ MDRC sEVs were membrane labeled
with PKH26 red dye. T cells were transduced with CellLight Tubulin-RFP 24 hours prior
to co-culture with sEVs. MDRC sEV co-culture with T cells (1:10 T cell:sEV) was carried
out for 24 hours and analyzed on the ImageStream flow cytometer. (B) Representative
images from ImageStream flow cytometry analysis illustrating polarization of α-tubulin
and host mitochondria to the site of MDRC sEV internalization. T cells were transduced
with CellLight Tubulin-RFP and labeled with MitoView Deep Red dye 24 hours prior to co-
culture with MDRC sEVs. sEVs derived from MDRCs transduced with Mito-GFP were
co-cultured with T cells (1:10 T cell:sEV) for 24-hours and analyzed on the ImageStream
flow cytometer. (C) Confocal image illustrating internalization of MDRC sEVs inside the
cytoplasm of the CD4⁺ T cells. sEVs derived from MDRCs transduced with Mito-GFP
were co-cultured with T cells (1:10 T cell:sEV) for 24-hours in the ibidi μ-dish. Cells were
fixed with 2% PFA and permeabilized with 0.5% triton-X in PBS. The fixed T cells were
labeled with Phalloidine-Rhodamine for actin, and DAPI for the nucleus. Cells were
imaged on the Nikon A1 confocal. (D) Confocal image illustrating polarization of α-tubulin
and host mitochondria to the site of MDRC sEV internalization. T cells were transduced
with CellLight Tubulin-RFP and labeled with MitoView Deep Red dye 24 hours prior to co-
culture with sEVs. sEVs derived from MDRCs transduced with Mito-GFP were co-cultured
with T cells (1:10 T cell:sEV) for 24 hours in the ibidi μ-dish and imaged on the Nikon A1
confocal.
818

Figure 7 – T cells that internalize MDRC-derived Mito-GFP$^+$ sEVs from asthmatics
polarize to Th17 and Th2 subsets. Mito-GFP$^+$ sEVs from asthmatics may activate immune
cells through DAMP signaling. sEVs from asthmatics induced pro-inflammatory cytokine
production by THP-1. (A-F) Airway MDRCs were transduced with CellLight Mito-GFP
and CellLight Endo-RFP, and sEVs were purified from the supernatant 48 hours later.
Purified MDRC sEVs were co-cultured with autologous peripheral CD4$^+$ T cells for 7 days
in the presence of rhIL-2 (50 IU/ml) in a ration of 1:10 T cells:sEV. T helper subsets were
assessed by flow cytometry 7 days later by staining intracellular for IL-4 (Th2) and IL-17
(Th17) cytokines. Cells were gated for CD4$^+$GFP$^+$IFNγ$^-$ cells, and then for either IL-4$^+$IL-
17$^-$ or IL-4$^-$IL-17$^+$. Mann Whitney T test for comparison between healthy and asthmatics,
Wilcoxon matched-pairs signed rank test for intergroup (no exosomes vs exosome),
*<0.05, **<0.01, ***<0.001, ****<0.0001. (G) Representative image strips from
ImageStream analysis. (H-J) MDRCs were transduced as above and sEVs were purified
as above in A-F. Purified MDRC sEVs were co-cultured with their allogenic T cells (i.e.
healthy sEV on asthmatic T cells, and vice versa; at a 1:10 T cell:sEV) for 24 hours.
Activation was assessed by CD69 and CD154 expression using the ImageStream
cytometry (T-test comparison to no-sEV healthy control and no-sEV asthma control;
*<0.05). (K-P) Airway sEVs were purified from BALF using a described ultracentrifugation
method. Purified sEVs were cultured with THP-1 cells (1:10 cell:sEV) for 24 hours.
Supernatants were harvested for BioRad Bioplex analysis of cytokines for (F) TNF-α, (G)
MCP-1, (H) IL1- β, (I) IFNγ, (J) MIP1-α, and (K) MIP1-β. (F-K) One-way ANOVA, *<0.05,
<0.01, *<0.001.
841
842

Figure 8 – Intranasal transfer of pro-inflammatory lung MDRC-derived sEVs from
sensitized and challenged donor Mito-QC mice with asthma exacerbates Th2 and Th17
responses and allergic airway inflammation in sensitized and challenged recipients. Mice
were sensitized by intraperitoneal injection on d0 and d7 with 50 μg of alum-adsorbed
OVA. On d14, d15 & d16 mice were challenged once i.n. with 15 μg OVA in 30 μl PBS or
PBS alone. On d16, i.n. delivery of lung MDRC-derived sEVs (1 x 10^8 particles/mouse in
30 µl PBS) from control or OVA challenged Mito-QC mice were carried out as before.
Lung infiltration of immune cells in lung tissue was determined by FACS analyses. Cell
numbers of Ly6G$^-$Ly6C$^+$ MDRC (A), Ly6G$^+$Ly6C$^-$ MDRC (B), Ly6G$^+$Ly6C$^+$ MDRC (C),
Neutrophils (D), Eosinophils (E), Th2 (F), Th17 (G), Th1 (H), and ILC2 (I) in the lung tissue
were determined by flow cytometry. Statistical significance was evaluated using one-way
ANOVA with Tukey's multiple comparison testing. ** $P < 0.01$, *** $P < 0.005$, **** $P <$
0.001.
856
857

Figure 9 – Activation of CD4$^+$ T cells by internalization *in vivo* of Mito-QC$^+$ sEVs during allergic airway inflammation. (A) Protein level and oligomerization of DRP1 was assessed in Ly6G$^+$ MDRCs sorted from the lungs of control or OVA challenged mice by Native-PAGE. The relative expression of DRP1 oligomer or monomer was normalized with β-actin. (B). sEVs were isolated from conditioned media of proinflammatory Ly6G$^+$ lung MDRCs sorted from control or OVA sensitized and challenged donor Mito-QC mice using the Total Exosome Isolation kit. Frequency of GFP$^+$MHC-II$^+$ sEVs in CD81$^+$ gate was characterized by ImageStream analyses. (C). Image strips showing GFP$^+$MHC-II$^+$CD81$^+$ sEVs isolated from proinflammatory Ly6G$^+$ MDRCs. (D). sEVs isolated from (B) were i.n. delivered (1 x 10^8 particles/mouse in 30 μl PBS) to recipient sensitized and challenged mice as before. Lung tissue was harvested and digested two days after sEV delivery. Image strips showing co-localization of GFP$^+$ signal in CD45$^+$CD4$^+$CD69$^+$ T cells. (E) Pearson correlation analysis between Mito-QC (%) and CD69 expression (MFI) on CD4$^+$ T cells. The data were analyzed by IDEAS 6.2 software. The orange dots represented Con + Con exo group, the blue dots represented Con + Asth exo, the red dots represented OVA + Con exo group, and the dark dots represented OVA + Asth exo group. (F) Pearson correlation analysis between Mito-QC (%) and CD69 expression (MFI) on CD4$^+$ T cells. The percentage area of GFP$^+$ signal or CD69$^+$ mean fluorescence intensity (MFI) within each cell, collected from 400 cells using MATLAB analysis. Each data point represents an individual cell. ** $p < 0.01$, **** $p < 0.0001$.

Figure 10 – Intranasal transfer of pro-inflammatory lung MDRC-derived sEVs from sensitized and challenged donor Mito-QC mice with asthma enhances mCherry$^+$GFP$^+$ Th2 and Th17 cell infiltration in the lung tissue, draining LNs and BALF in sensitized and challenged recipients during allergic airway inflammation. Mice were sensitized by intraperitoneal injection on d0 and d7 with 50 μg of alum-adsorbed OVA. On d14, d15 & d16 mice were challenged once i.n. with 15 μg OVA in 30 μl PBS or PBS alone. On d16, 4 hrs after i.n. challenge with OVA in or PBS, i.n. delivery of lung MDRC-derived sEVs (1 x 10^8 particles/mouse in 30 µl PBS) from control or OVA challenged Mito-QC mice were carried out. Infiltration of immune cells in the lung, draining LNs and BALF were detected by FACS analyses at two days after sEV delivery. Frequencies of mCherry$^+$GFP$^+$ Th2 (A) and Th17 (B) in the lung tissue were determined by flow cytometry. Frequencies of mCherry$^+$GFP$^+$ Th2 (C) and Th17 (D) in the draining LNs were characterized by flow cytometry. Frequencies of mCherry$^+$GFP$^+$ Th2 (E), Th17 (F) and ILC2 (G) in BALF were determined by flow cytometry. Statistical significance was evaluated using one-way ANOVA with Tukey's multiple comparison testing. ** $P < 0.01$, *** $P < 0.005$, **** $P < 0.001$.

Media: Titles and Legends

SV1: MDRC-derived Mito-GFP⁺ sEVs co-localize with actin in activated T cells of subjects with asthma. A z-stack slice of the confocal images indicated co-localization (yellow) of the Mito-GFP⁺ sEVs with cytoplasmic actin (red) in the T cells.

SV2: MDRC-derived Mito-GFP⁺ sEVs co-localize with polarized cytoskeleton and mitochondrial network of recipient CD4⁺ T cells. Confocal images of Tubulin-RFP transduced and Mitoview (labels mitochondria) labeled T cells, co-cultured with MDRC-derived Mito-GFP⁺ sEVs showing co-localization of Mito-GFP⁺ sEVs with the polarized cytoskeleton and mitochondrial network of the recipient CD4⁺ T cells.

909
910

911 **Table 1** – Characteristics of volunteer subjects enrolled in the study.

Characteristics of Enrolled Patients			
Characteristics	Health (N=19)	Asthmatics (N=15)	p-value
Sex			
Male	7	2	
Female	12	13	
Median Age (IQR)	43.5 (22)	36 (24)	0.3654
Median IgE Titer (IQR) – IU/L	23.55 (85.6)	97.4 (598.3)	0.0273
Median Predicted % FEV1 (IQR)	101.5 (17.05)	87 (27)	0.0001
Median Eosinophils (IQR)	113.4 (92.8)	214 (159.55)	0.0038
Median Neutrophils (IQR)	3.46 (1.72)	3.61 (1.96)	0.6959

Figure 1

Figure 2

Figure 4

Figure 5

Figure 6

A

Bright Field	Mito-GFP	PKH26	αTubulin	Merge

B

Bright Field	CD4	Mito-GFP	αTubulin	MitoView	Merge

C

DAPI | Phalloidin | Mito-GFP⁺ sEVs | Composite

D

Mitoview | Tubulin-RFP | Mito-GFP⁺sEVs | Composite

Figure 7

Figure 8

Figure 9

A

CON OVA DRP1

Oligomer

Monomer

β-actin

DRP1 Oligomer Ratio (Asthmatic:Control) — ****
Control, Asthmatic

DRP1 Monomer Ratio (Asthmatic:Control) — ****
Control, Asthmatic

B GFP⁺MHC-II⁺

Percentage of CD81⁺ sEV — **
Con-sEV, Asth-sEV

C

Ch01 GFP CD81 MHC-II

7 µm

D

Ch01 GFP CD45 CD4 CD69 Composite

7 µm

E

Pearson r = 0.9894, *p* < 0.0001

- Con + Con sEV
- Con + Asth sEV
- OVA + Con sEV
- OVA + Asth sEV

CD69 (MFI)

GFP⁺ (% of CD4+ T cells)

F

Pearson r = 0.9872, *p* < 0.0001

CD69 (MFI)

GFP⁺ (% area within the cell)

Figure 10