1 **TITLE**
2 Rapid adaptive evolution of microbial thermal performance curves
3
4 **SHORT TITLE**
5 Evolution of microbial thermal performance curves
6

7 **AUTHORS**
8 Megan H. Liu[1*], Ze-Yi Han[1], Yaning Yuan[1], Katrina DeWitt[1], Daniel J. Wieczynski[1], Kathryn
9 M. Yammine[2], Andrea Yammine[1], Rebecca Zufall[3], Adam Siepielski[4], Douglas Chalker[5],
10 Masayuki Onishi[1], Fabio A. Machado[6], Jean P. Gibert[1*]
11
AFFILIATIONS
[1]Department of Biology, Duke University, Durham, NC, 27708, USA
[2]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, 02139,
USA
[3]Department of Biology and Biochemistry, University of Houston, Houston, TX 77204, USA
[4]Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA
[5]Department of Biology, Washington University, Saint Louis, MI, USA
[6]Department of Integrative Biology, Oklahoma State University, Stillwater, OK, USA
*Corresponding author—email: mhliu58@gmail.com, jean.gibert@duke.edu
21

22 **KEYWORDS:**
23 Thermal Adaptation, Thermal Performance Curve Heritability, Climate Change, Quantitative
24 Genetics
25

STATEMENT OF AUTHORSHIP:
All authors contributed to the conceptual foundation of the study. MHL, ZH, and JPG designed
the study. DC contributed the fluorescent lines and provided cell biology expertise along with
MO. MHL, ZH and AY collected the data. MHL and JPG performed data analyses with support
from DHW, RZ, and AS. JPG did the mathematical modeling and FAM the quantitative genetics.
MHL and JPG wrote the first draft of the manuscript and all authors contributed significantly to
revisions.
33

34 **DATA ACCESSABILITY STATEMENT**
35 All software and data from this study are available on GitHub
36 (https://github.com/JPGibert/TPC_evolution).
37

38 **COUNTS**
39 Abstract words: [148] | Main text words: [4867] | References: [72] | Figures: [4] | Tables: [0]
40 To whom correspondence should be addressed: mhliu58@gmail.com, jean.gibert@duke.edu

ABSTRACT

Microbial respiration is a key biotic driver of climate change. Warming boosts microbial

population growth, which increases biomass and respiration. This feedback might be disrupted

by adaptation in thermal performance curves (TPCs) –whose shape describes how temperature

drives growth. In this study, we uncover substantial genetic variation (G) in microbial intrinsic

population growth rates (r), demonstrate a causal link between G variation in r and G variation in

TPC shape, and show how this variation constrains r-TPC shape evolution along specific

evolutionary paths across temperatures. We also uncover Gene-by-Environment (G × E)

variation in r, which results in specific signatures in TPC shape and predictable temperature-

dependent rapid TPC evolution but also lower G, which could reduce future evolutionary

potential. Overall, we show how temperature-dependent evolution in a linchpin of global

ecosystem function—microbial TPC shape—is determined by a combination of heritable and

non-heritable variation in intrinsic growth rates.

INTRODUCTION

Microbes play a central role in regulating the global carbon (C) cycle that controls

climate change (Falkowski *et al.* 2008). Indeed, soil microbial respiration releases ~94Pg/yr of C

into the atmosphere (Stell *et al.* 2021) while microalgae fix 30-50 Pg of C/yr globally (Falkowski

1994). Global warming is expected to alter these microbial processes (IPCC 2023), but

anticipating these effects requires a deeper understanding of the biotic and abiotic factors

influencing microbial respiration in a warming world (Rocca *et al.* 2022; Wieczynski *et al.*

2023).

One such factor is microbial population growth, which influences total standing biomass,

and hence, total microbial respiration (Savage *et al.* 2004). The thermal performance curve of

intrinsic population growth rate (*r*) describes how microbial population growth changes with

temperature ('*r*-TPCs' henceforth, Fig 1). The shape of r-TPCs is controlled by temperature-

dependent metabolism: metabolism increases with temperature, and so does *r*, until an 'optimal'

temperature (T_{opt}) is reached (Fig 1a), then *r* declines as metabolic costs increase (Fig 1a;

Amarasekare & Savage 2012). While *r*-TPC shape varies across species (Jacob & Legrand

2021;), unimodal shape is the norm (Wieczynski *et al.* 2021), described by shape parameters

linked to thermal ecology: maximum growth rate (r_{peak}), minimum and maximum temperatures

for population growth (CT_{min} and CT_{max} respectively), and rates of *r*-TPC increase/decrease with

temperature (E_a, E_d respectively, Fig 1b). Ultimately, differences in *r*-TPC shape across species

reflect divergent evolutionary trajectories in shape parameters across species and environments

(Angilletta 2009).

While *r*-TPC shape is expected to evolve in novel environments, anticipating this thermal

adaptation is a major open question, as the r-TPC reflects a species' ability to cope with

environmental change. Thermal adaptation hinges on the evolution of intraspecific—heritable—

genetic variation and selection favoring genetic variants suited to novel environments (Frankham

2005), so quantifying heritable intraspecific variation in r-TPC shape parameters is central to

understanding r-TPC shape evolution (Kling *et al.* 2023). From a quantitative genetics

standpoint, the r-TPC represents the reaction norm of the underlying measure of performance, r,

across temperatures among genotypes (Golmulkiewicz *et al.* 2018). r-TPC intraspecific variation

is thus inextricably linked to that of r. Understanding this link is key to quantifying heritable

variation in r-TPC shape parameters and predicting their response to selection in novel

environments.

To understand the link between intraspecific variation in r and r-TPC shape parameters,

we developed a verbal model using classic quantitative genetics of reaction norms (Fig 1, c-e).

First, environmental variation (E) in r, driven by differential expression of r across temperatures,

but not genotypes, classically manifests as completely overlapping genotypic reaction norms (Fig

1C, first row). Correspondingly, genotypic r-TPCs should also overlap (Fig 1D, first row) as they

are reaction norms extended over multiple temperatures. Overlapping r-TPCs have no variation

in shape parameters (Fig 1e, first row), so E in r should not result in r-TPC shape parameter

variation. Second, genetic variation (G) in r, arising from differential expression of r across

genotypes, but not temperatures, results in "additive" shifts in the intercepts of genotypic

reaction norms—but equal slopes (Fig 1C, second row). These additive shifts translate to parallel

r-TPCs (Fig 1D, second row), with additive variation in their x-axis intercepts (CT_{min}, CT_{max})

and their maximum height (r_{peak}), but no variation in T_{opt}, E_a, E_d (Fig 1E, second row). Last,

gene-by-environment (G × E) interactions in r, which manifest as changes in intercept and slope

of reaction norms across genotypes and temperatures (Fig 1C, third row), result in non-parallel r-

TPCs (Fig 1D, third row), with greater variation in all shape parameters (Fig 1E, third row). This

conceptual model therefore predicts covariation between heritable genetic variation in r and

variation in CT_{min}, CT_{max}, and r_{peak}, so r-TPCs' CT_{min}, CT_{max}, and r_{peak} are more likely to be

heritable and respond to selection than T_{opt}, E_a, or E_d, which in turn suggest constraints operating

on possible r-TPC evolutionary trajectories.

To address how r-TPC shape might evolve across temperatures we thus quantify E, G,

and G × E variation in r, and test predictions from our conceptual model. In doing so we reveal

that heritable variation in r results in substantial heritable variation in predicted shape

parameters. We build upon this understanding to quantify the structure of genetic variation and

covariation in shape parameters, and use this information to infer selection and possible r-TPC

evolutionary trajectories of shape parameters across temperatures. Last, we uncover a

mechanism of rapid r-TPC evolution in the presence of G × E variation in r that leads to

temperature-dependent selection on r-TPC shape, but can erase heritable variation, possibly

creating an evolutionary trap that constrains future microbial r-TPC evolution.

METHODS

Study system and genotypes

We used *Tetrahymena thermophila*, a freshwater ciliate protist found across the

Northeastern United States (Zufall *et al.* 2013) and part of a cosmopolitan genus (Lynn &

Doerder 2012). Protists, unicellular Eukaryotes that dominate oceanic biomass and rank third in

terrestrial biomass, comprise twice the biomass of the Animal Kingdom (Bar-On *et al.* 2018) and

underpin global ecosystem functioning (Gao *et al.* 2019). So, while no single species is

representative of the entire group, understanding thermal adaptation in such an ecologically

pivotal group is meaningful.

To do so, we sourced 22 unique *T. thermophila* genotypes: 19 from the Cornell

Tetrahymena Stock Center and 3 from the Chalker lab (Washington University, Appendix 1).

These genotypes, which vary in geographic origin and have well-known genetic differences

(Appendix 1), were chosen to sample existing genetic variation, not for their functional

significance. As most were derived from laboratory cultures, our assemblage likely contains less

variation than natural populations (Zufall *et al.* 2013), making our results a conservative estimate

of how thermal adaptation might proceed in nature.

Quantifying r, r-TPCs, and TPC shape parameters

Culture care and maintenance followed standard practices for ciliate Ecology (Appendix

2). Stock cultures were maintained in Percival AL-22 growth chambers with a 12hr day-night

cycle at 22°C. We quantified *r*-TPCs for all genotypes through growth assays in 3cm diameter

Petri dish microcosms containing 3mL of growth medium at seven temperatures (13, 19, 22, 25,

30, 32, 38°C). Each genotype and temperature combination was replicated six times, totaling 924

microcosms. Experimental temperatures span below and above the average growing season

temperature in *T. thermophila*'s native range (~23ºC ; NOAA 2024). Microcosms were

initialized at densities of 1ind/mL in 3ml Petri dishes, which results in exponential growth for 1-

2 days (Gibert *et al.* 2022, 2023; Singleton *et al.* 2021). After 24hrs, we censused the

microcosms through whole-population counts under a stereomicroscope (Leica M205C) and

calculated *r* across temperatures to characterize the entire *r*-TPC as log(final density/initial

density)/time, with time = 1 day (Wieczynski *et al.* 2021). To obtain *r*-TPC shape parameters, we

fitted a Sharpe-Schoolfield model ("nls.multstart" v1.3.0 package in R, (Padfield 2023)). Shape

parameters E_a, r_{peak}, CT_{min} and T_{opt} (Fig 1b) could be unequivocally estimated from our data, and

so we focused all subsequent analyses on those. These parameters control the rising portion of

the r-TPC (Fig 1b, green) –the "operational temperature range" (DeLong *et al.* 2017), i.e., what

most organisms are likely to experience in their native geographic ranges.

Link between heritable genetic variation in r, and r-TPC shape parameters

We quantified E, G and G × E variation in r using function *gxeVarComps*() in R package

statgenGxE v1.0.5. The function fits a linear model with r as the response variable, and

temperature, genotype, and their interaction as fixed predictors, to calculate effect sizes and

statistical significance. It then fits a second model with all fixed terms treated as random effects

to calculate r variance components (E, G, and G × E). Only G is considered to be heritable, so

the fraction of heritable genetic variation in r, can be estimated as the broad-sense heritability

($H^2 = G/(E+G+G \times E)$). However, this expression does not account for inter-treatment and

replicate variability, which inflates E and underestimates H^2 (Cullis *et al.* 2006; Piepho &

Möhring 2007). In the appendix we present three alternative formulations for H^2 that address this

issue (Appendix 3).

Next, we tested the predictions of our verbal model (Fig 1) linking heritable variation in r

(G) to heritable variation in r-TPC shape parameters. Since each r-TPC yields a single shape

parameter set, only G variation in shape parameters could be quantified. All other components of

variation (e.g., E, G × E, error) were pooled as residual variance. To evaluate the model's

prediction that G variation in r results in greater G variation in shape parameters r_{peak} and CT_{min}

compared to E_a, and T_{opt} (Fig 1e), we tested whether r_{peak} and CT_{min} exhibited stronger genetic

covariance with r than did E_a, and T_{opt} using a multivariate Bayesian Generalized Linear Mixed

Model. The model includes mean-scaled values of r, E_a, r_{peak}, CT_{min} and T_{opt} as jointly modeled

response variables, each with their own fixed mean, and Genotype as a random effect to estimate

variances and covariances implemented in R package MCMCglmm v.2.36 (Hadfield 2010).

Variance-covariance priors were obtained by bootstrapping r-TPC data 100 times within each

genotype and re-fitting the Sharpe-Schoolfield model to each bootstrap replicate, generating a

population of shape parameters for each genotype (Appendix 4). Using inverse Wishart priors

(Murphy 2012) did not alter our results. Last, if G variation in r results in greater G in r_{peak} and

CT_{min}, these shape parameters should also have greater heritability than E_a, and T_{opt}. To confirm

this, we calculated the broad-sense heritability in r-TPC parameters using the standard

expression $H^2 = G/(G + \text{residual variance})$ with G and residual variance estimates from the

MCMCglmm model.

Consequences of heritable variation: selection and evolutionary potential of r-TPC parameters

Heritable r-TPC shape parameters can evolve under selection. To understand how, we

quantified 1) the direction, form, and magnitude of selection on all four shape parameters, 2)

whether and how temperature influenced selection, and, 3) their potential evolutionary responses

across temperatures. We used two approaches: one that neglects genetic correlations between

parameters but allows estimation of non-linear selection (e.g., stabilizing selection), and one that

accounts for genetic correlations and neglects non-linear selection, while enabling predictions of

evolutionary responses across temperatures.

In the first approach, we quantified selection by characterizing the adaptive landscape

(Lande 1976, 1979) as the relationship between shape parameter values and absolute fitness—

classically assumed to a function of r (Lande 1976, 1979) because it reflects average birth and

death rates of individuals, two major fitness components (Lande 1982; Partridge & Harvey

1988). We modeled this relationship for all shape parameter values fitted across genotypes, using

polynomial regression and r as the response, linear and quadratic effects (i.e., non-linear

selection) for each mean-standardized shape parameter, and additive and interactive temperature

effects with the linear and quadratic shape parameter terms (see Appendix 5). We multiplied the

quadratic regression coefficient by two (Stinchcombe *et al.* 2008). A positive relationship

between r and the focal shape parameter would be evidence of positive directional selection, a

negative relationship would indicate negative directional selection, and no relationship suggests

no directional selection. Stabilizing selection would result in a concave-down relationship, and

disruptive selection would manifest as a concave-up relationship where extreme values have

higher fitness (Lande & Arnold 1983).

The second approach predicts r-TPC shape evolution across temperatures while

accounting for genetic correlations among shape parameters. It uses the multivariate breeder's

(Lande 1979) and Price's (Price 1972) equations to estimate direct (i.e., acting on the focal

parameter) and indirect selection (i.e., acting on genetically linked parameters, (Stinchcombe *et*

al. 2014)). This is achieved by estimating the response to selection of all shape parameters, $\mathbf{\Delta z}$,

as their covariance with fitness—i.e., the Price equation aspect of this approach– which requires

estimating a genetic variance-covariance matrix ($\mathbf{G_{zw}}$)—i.e., its multi-variate breeder's equation

aspect. This matrix includes the genetic variance-covariance matrix ($\mathbf{G}$) of shape parameters as

its first four rows/columns, and their genetic covariation with fitness (W) as the last row/column

(Lande 1976, 1979, 1982). The finite rate of increase R=$exp(r)$ was used as our metric of fitness

but using plain r did not alter our results. From the Price equation, this last row/column also

equals the vector of predicted trait change, $\Delta\mathbf{z}$, so $\mathbf{G_{zw}}$ also yields $\Delta\mathbf{z}$. Then, from the multivariate

Breeder's equation, $\Delta\mathbf{z} = \mathbf{G\beta}$, where $\mathbf{\beta}$ is the selection gradient (Lande 1979). Provided that $\mathbf{G}$ is

invertible, then $\mathbf{\beta} = \mathbf{G}^{-1} \Delta\mathbf{z}$, to estimate selection.

Positive/negative Δz_i values indicate an increase/decrease in the i-th shape parameter,

while positive/negative β_i, indicates positive/negative selection operating on the i-th shape

parameter. Alignment between Δz_i and β_i indicates direct responses to selection (Hansen &

Houle 2008), in our case, by temperature, while misalignment suggest indirect selection through

correlated responses with other shape parameters.

To estimate $\mathbf{G_{zw}}$ for each temperature, we used subsets of r for each genotype for each

temperature. Shape parameters do not change across temperatures but r does, so the $\mathbf{G_{zw}}$ matrices

only differed in their last row/column across temperatures, i.e., the covariances between shape

parameters and fitness—or $\Delta\mathbf{z}$. To facilitate this estimation and ensure interpretability, all $\mathbf{G_{zw}}$

variables were rescaled, which was achieved through the expression $\mathbf{G_{zw}}$ as $=\mathbf{SLS}*(1/2F)$ where

$\mathbf{L}$ is a between-genotype covariance matrix, $\mathbf{S}$ is a diagonal matrix containing the inverse of each

trait's standard deviation, and F is the inbreeding coefficient, which equals 1 for clonal lineages

(Falconer 1996). We quantified uncertainty in $\mathbf{G_{zw}}$ entries using a Bayesian posterior distribution

of $\mathbf{L}$ matrices (and hence $\mathbf{G_{zw}}$) and used a multivariate normal likelihood function and non-

informative inverse regularized Wishart priors (nu=traits+1) (Murphy 2012), implemented in

evolqg v3.0 R package (Melo *et al.* 2015). Using MCMCglmm instead of evolqg did not affect

our results (Appendix 6). We took 1000 posterior samples for $\mathbf{G_{zw}}$ to calculate $\Delta\mathbf{z}$, $\mathbf{G}$, and $\mathbf{\beta}$, and

95% maximum density intervals for al estimates.

Consequences of G × E in r for rapid r-TPC evolution

Because G × E variation in r is predicted to result in non-parallel r-TPCs (Fig 1), slight

differences in intrinsic growth rates across temperatures can drive temperature-mediated

differential selection among genotypes (Fig 4a, b). Treating r as proxy for absolute fitness

(Lande 1976), this selection should be strongest at temperatures where r-TPCs differ the most,

i.e., the temperatures with the largest differences in relative fitness.

We empirically tested for temperature-dependent selection among genotypes by setting

up an experimental evolution assay with two fluorescently marked strains (AXS and CU4106,

Fig 4c) with different r-TPCs and relative fitness (Fig 4d, Fig4d inset) across six temperatures

(19, 22, 25, 30, 32, 38ºC), each replicated seven times, and 2 additional single-strain controls per

temperature. The genotypes cannot mate because they belong to the same mating type (mating

type VII, Appendix 1), so only clonal reproduction was possible. We initialized our microcosms

at equal densities (5 ind/mL in mixture treatments, 10ind/mL for single-genotype controls). After

48 hours, we added Cadmium Chloride to induce fluorescence (Appendix 7), confirmed on a

Leica Thunder Cell Culture inverted microscope (Fig 4c and Appendix 8). Thirty minutes later,

we cytometrically censused each sample (Novocyte 2000R). While fluorescently tagged

genotypes can lose their ability to fluoresce over time, they carry a paromomycin resistance gene

(Appendix 7), allowing selection through paromomycin exposure (100μg/mL) prior to counting

and estimating relative frequencies based on fluorescence (Appendices 9-10). Antibiotics can

negatively affect protists and the bacterial communities they feed on, so we replicated this

experiment in paromomycin-free conditions, which did not qualitatively alter our results (see

Appendix Fig S11).

Last, observed changes in genetic frequencies were compared to predicted frequencies by

a classic model of genetic evolution in discrete time parameterized with the r-TPC of both

experimental genotypes (Fig 4f). The model tracks the frequency of each strain f_i, and assumes

that their absolute fitness, W_i, is a function of their average intrinsic growth rate r (Lande 1976,

1979). The frequency of each strain in the population is determined by the classic recursive

equation, $f_i(t + 1) = f_i(t)(W_i/\overline{W})$ [Eq 1.], where $\overline{W}$ is the average fitness of the population

($\sum_{i=1}^{n} f_i W_i = \sum_{i=1}^{n} f_i r_i$), and $W_i/\overline{W}$ is fitness the relative fitness of i-th genotype *relative* to the

entire population. We used each genotype's r-TPC (AXS, CU4106, Fig 4d) to estimate $W_i/\overline{W}$

across temperatures (i.e., replacing r_i). We forward-solved the recursive equation (Eq 1) under

two scenarios: one in which temperature was constant, and one in which temperature fluctuated

randomly across generations, drawn from a normal distribution with known mean and variance.

This minimal model assumes that differential selection across environments is mostly driven by

differences in r-TPCs, and disregards density- and frequency-dependent selection, sexual

reproduction, mutation, and most all ecological processes.

RESULTS

Heritable variation in r leads to heritable variation in r-TPC shape parameters

Intrinsic growth rates showed strong temperature responses within genotypes (F =

3092.70, $D_f = 6$, Generalized Eta-Squared (GES) effect size = 0.964, $p \leq 0.001$), significant

variability across genotypes (F= 163.65, $p \leq 0.001$, $D_f = 21$, GES = 0.832), and significant G ✕

E interactions (F=29.7, $p \leq 0.001$, $D_f = 122$, GES = 0.840), leading to classically unimodal

shapes that varied among genotypes (Fig 2a). Environmental variation (E) accounted for 71.7%

of all observed variation in r, genetic variation (G) explained 6.1% of all variation, and Gene-by-

Environment interactions (G × E) explained 11.7%, with 10.5% residual variation (Fig 2b). This

is well within what is expected for life history traits (Hoffmann & Sgrò 2011). Despite a

relatively modest G compared to E, after accounting for experimental inter-treatment and

replicate variability, r was strongly heritable ($H^2_{standard}$=0.76, H^2_{cullis}=0.95, H^2_{piepho}=0.91,

Appendix 3).

Shape parameters varied widely across genotypes (e.g., genotype AXS was 3.2 times

more thermally sensitive than A1868 and could grow 3.6 times faster at T_{opt}, Fig 2, and

Appendices 12-13). As predicted, CT_{min} and r_{peak} showed strong genetic covariation with r (|

$Gcov_{CTmin}$ | = 0.65, | $Gcov_{rpeak}$ | = 0.63, Appendix 14), but there was no evidence of genetic

covariation between r and T_{opt} or E_a (credible intervals contained 0, Appendix 14). Last, shape

parameter heritabilities matched verbal model predictions (H^2_{CTmin}=0.945, H^2_{rpeak}=0.949,

H^2_{Topt}=0.070, H^2_{Ea}=0.002), confirming that heritable variation in r results in heritable variation in

CT_{min} and r_{peak}, but not T_{opt} or E_a.

Consequences of G in r: selection and evolvability of r-TPC shape parameters

Without accounting for genetic covariances, selection operated differently across shape

parameters and was temperature dependent: r_{peak} was under negative directional selection at low

temperatures (<20°C, Fig 3a, Appendix 15), weakly positive or no directional selection at

intermediate temperatures (between 20 and 30°C, Fig 3a, Appendix 15), and strong positive

directional selection in high temperatures with E_a following a similar pattern (Fig 3b, Appendix

16). CT_{min} was under negative selection at low/intermediate temperatures but no selection at high

temperatures (Fig 3c, Appendix 17). Last, T_{opt} was under no selection at low temperatures but

under weak then strong stabilizing selection at intermediate and high temperatures, respectively

(Fig 3d, and Appendices 18-19).

However, we also found clear positive genetic covariances between r_{peak} and E_a, CT_{min}

and E_a, and CT_{min} and T_{opt} (Fig 3e). Consequently, selection acting on highly heritable

parameters CT_{min} and r_{peak} could still result in correlated evolution in weakly heritable parameters

E_a, and T_{opt}. Accounting for these genetic covariances, we confirmed differences in selection

across shape parameters whose magnitude and direction also shifted with temperature (Fig 3f),

resulting in possible temperature-dependent shifts in parameters (Fig 3g). Specifically, our

multivariate analysis suggested that selection would favor higher r_{peak} (and maybe E_a) at high

temperatures, and low E_a and CT_{min} at low temperatures, but no selection on T_{opt} (Fig 3f),

generally matching univariate predictions (Fig 3g). Overall, the evolutionary responses followed

predicted trajectories from the estimates of selection closely (Fig 3f-g), with a few exceptions

that suggest correlated evolution: positive response of E_a with temperature likely are correlated

responses with r_{peak} (Fig 3g), which in turn shows strong negative response at low temperatures

despite no selection, likely through correlated evolution with E_a whose negative response is

driven by negative selection in CT_{min} (Fig 3g). Despite differences in heritability which impose

constraints in r-TPC evolution, genetic correlations among shape parameters allowed for

temperature-dependent evolutionary responses in all of them except T_{opt}, which our results

suggest is under stabilizing selection (Fig 3d).

Consequences of G × E in r: sorting of standing genetic variation across temperatures

We observed significant temperature-dependent differential selection among genotypes

(Fig 4e), matching theoretical predictions from a simple model of genetic evolution (Fig 4f),

which confirms the hypothesis that G × E in *r* could set the stage for rapid r-TPC shape evolution

through selection on standing genetic variation (Fig 4a, b). Despite quantitative discrepancies

between observed and predicted frequencies—notably at 19ºC where the model predicted a

polymorphic population but the data indicated otherwise (Fig 4e, f)—it still correctly predicted

observed changes in genetic frequencies across most temperatures. These results are also robust

to different experimental conditions (Appendix 11) and fluctuating temperatures (Appendix 20),

the latter showing that temperature fluctuations could play a role in determining r-TPC evolution

if r-TPCs show multiple crossing points in a short temperature span.

The results of this experiment were also consistent with our estimated predicted

responses to selection (cf. Fig 3 and Fig 4): lower temperatures led to higher frequencies of the

CU4106 genotype, whose r-TPC has lower E_a and CT_{min} compared to AXS (Fig 4d), so the

average r-TPC of the population should reflect that and also have lower E_a and CT_{min}, as

predicted by our multivariate model. At higher temperatures, selection favored AXS, which has

higher E_a, so the ensuing population also should have an average r-TPC with higher E_a (Fig 4d).

Last, rising temperatures led to an increase in genetic variance ($\sim p(1-p)$ where p is the frequency

of either genotype and 1-p that of the other) compared to lower temperatures (Fig 4e, f)—which

should facilitate adaptation—then a reduction in genetic variance (as genotype CU4106 becomes

less prevalent, Fig 4e, f), which in turn could impede adaptive evolution in the future.

DISCUSSION

Our study reveals heritable genetic variation in *Tetrahymena thermophila*'s population

intrinsic growth rates that results in heritable variation in some, but not all, r-TPC shape

parameters (Fig 2), allowing for thermal adaptation in new climates (Fig 3). Observed *intra*-

specific variation in r-TPC shape parameters rivals reported *inter*-specific variation across

multiple protist species (Wieczynski *et al.* 2021). We also found that most r-TPC shape

parameters are under different selection regimes across temperatures (Fig 3), with parameters

CT_{min} and r_{peak} being more likely to respond to selection than E_a and T_{opt}, but even weakly-

heritable parameters can still evolve through correlated evolution (Fig 3). Consequently, colder

temperatures select for lower CT_{min}, r_{peak}, and E_a, and warmer temperatures lead to high r_{peak} and

E_a (Fig 3). Last, we showed that G × E in r can lead to rapid, predictable shifts in population

genetic makeup across temperatures and r-TPC evolution (Fig 4), in a "plasticity drives

adaptation" scenario (Ghalambor *et al.* 2007).

While the evolution of microbial TPCs is likely the product of adaptation to local habitats

(Kontopoulos *et al.* 2020), how r-TPCs will adapt to ongoing rising temperatures remains an

open question. Several canonical evolutionary paths have been proposed: 1)'Colder-is-Better'

(CIB), where rising temperatures reduce population growth, leading to lower r_{peak}, T_{opt}, and E_a

(Kingsolver & Huey 2008). 2) Warmer-Is-Better (WIB), where higher growth rates evolve in

warmer temperatures, leading to TPCs with higher r_{peak} and T_{opt} (Pawar *et al.* 2015). And, 3)

Generalist-Specialist-Tradeoff (GST), where species can either evolve towards rapid growth

within a narrow temperature range (i.e., temperature specialists), or slower growth over a broader

temperature range (i.e., temperature generalists), leading to higher r_{peak}, higher CT_{min}, and lower

CT_{max} (Seebacher *et al.* 2015). Tests of these possible evolutionary paths (Kontopoulos *et al.*

2020; Montagnes *et al.* 2022) mostly use inter-species comparisons that often overlook intra-

specific variation and genetic associations between shape parameters, and therefore cannot

readily make predictions about r-TPC evolutionary trajectories for any given species. Indeed,

without accounting for genetic associations, our results would suggest support for WIB with

clear directional selection for higher r_{peak} and E_a under warming climates (Fig 3a, b). Accounting

for genetic covariances, however, suggested more complex evolutionary r-TPC responses than

currently predicted by theory. Specifically: we show support for WIB as warming should favor r-

TPCs with high r_{peak} and high E_a (Fig 3f, g), but no support for GST, as CT_{min} and r_{peak} responded

mostly together (Fig 3f, g). Last, T_{opt} is assumed to evolve under WIB, CIB and GST, but we

found no selection (Fig 3f) –or maybe even weakly stabilizing selection (Fig 3d)– and no

evolutionary response (Fig 3g). This suggesting that neither WIB, CIB and GST can explain

predicted evolutionary responses in *T. thermophila*, which emphasize the importance of

intraspecific variation and genetic associations to fully understand r-TPC evolution.

When G × E in r is prevalent, we showed that thermal adaptation can occur rapidly

through temperature-dependent selection on r-TPC genotypes (Fig 4), resulting in reduced

genetic variation. Loss of G which could slow down, or impede, future adaptation (Pauls *et al.*

2013). In other words, we show that phenotypic plasticity in r could lead to adaptive evolution in

r-TPCs in novel climates. That this form of r-TPC evolution could be anticipated from simple

models of genetic evolution, is striking, considering how little information beyond the r-TPC of

each genotype the model accounts for. Yet, r-TPC adaptation in deep time has been suggested to

occur gradually across six different *Tetrahymena* species (Montagnes *et al.* 2022), directly

countering our claim that r-TPCs could evolve rapidly through temperature-dependent selection.

We argue that this form of rapid r-TPC evolution may only play out during fast-changing

environmental conditions, not always result in long-term r-TPC change, or be masked by

reversals in selection. In fact, selection often "erases its traces" (Haller & Hendry 2014), and is

strongest under novel conditions, weakening as populations adapt (Caruso *et al.* 2017)—i.e., "the

paradox of stasis" (Haller & Hendry 2014).

Resulting temperature-dependent loss of genetic diversity of adaptive r-TPC evolution

could impact the persistence of genetically depauperate species under warming. But whether this

may be the case in nature hinges on quantifying processes that generate diversity (mutation, gene

flow, recombination) in r and r-TPC shape parameters in the wild, and assessing when these

processes could counter loss of genetic diversity after episodes of thermally-induced r-TPC

evolution. *T. thermophila* has the lowest reported base substitution mutation rate of all species

(Zufall et al. 2016), and limited gene flow (Zufall et al. 2013), but even weak dispersal or rare

mutation events could contribute to r and r-TPC variation and rescue genetic diversity in very

large, rapidly reproducing populations. The prevalence of sexual reproduction in nature is

debated –e.g. ~50% of wild individual *T. thermophila* cannot reproduce sexually and some

Tetrahymena lineages have lost this ability entirety (Doerder 2014)– but likely important if

present. In fact, *T. thermophila* possesses two separate genomes, a germinal line (in

micronucleus) and an expressed genome (in macronucleus). Our study focuses on macronuclear

variation, but asexual reproduction following sex results in random distribution of macronuclear

alleles to daughter cells—i.e. phenotypic assortment—potentially generating novel variation in r

and r-TPCs (Tarkington *et al.* 2023). Moreover, seemingly lost genetic variation could persist in

the population in other individual's micronuclei and reemerge following sexual reproduction

(Dimond & Zufall 2016). Fully understanding thermal adaptation in nature thus requires

exploring these processes' interplay, yet, for most organisms, they remain poorly understood.

Caveats

While bodies of water in *T. thermophila*'s native range are warming significantly (US EPA

2021) and experiencing more frequent and extreme heatwaves (Tassone *et al.* 2023), water

temperature typically remains 2-5ºC cooler than air temperature (Stefan & Preud'homme 1993).

Thus, some of the warmer temperatures in our study may be uncommon in *T. thermophila*'s

native range. However, strong temperature effects observed at low temperatures suggest these

patterns can still operate in many *T. thermophila* populations. Also, previous research showed

that most variation in *r*-TPC shape across species could be collapsed into a single dimension of

variation (Rezende & Bozinovic 2019), suggesting that some of the covariation observed in this

study between shape parameters could be spurious. If these *r*-TPCs could actually be collapsed

into a single axis of variation, however, we would have expected strong and multiple covariation

between most if not all shape parameters—which was not the case (Fig 3e, f). Interestingly, most

of the genetic variation in shape parameters was observed, as predicted, in CT_{min} and r_{peak}, but

these two parameters were not correlated to one another. Perhaps our intraspecific *r*-TPC

variation results stand in contrast to those observed across species (Rezende and Bozinovic,

2019), but more research is needed to clarify this point.

Conclusion

Overall, TPCs control the fate of populations (Sinclair *et al.* 2016), ecological

interactions (Enquist *et al.* 2015), food web dynamics (Barbour & Gibert 2021; Gibert *et al.*

2022), and ecosystem processes (Gibert *et al.* 2015). Yet, TPC evolution in a rapidly warming

world remains conspicuously unknown. Here, we show that *r* intraspecific variation can drive

temperature-dependent evolution in microbial *r*-TPCs and have consequences for rapid shifts in

population genetic makeup. Despite studying a single protist species, our findings provide a

foundation for studying thermal adaptation in this diverse and important microbial group and

beyond. Our study emphasizes the importance of temperature in mediating rapid microbial

evolutionary change as we grapple with understanding and predicting organismal responses to an

increasingly warmer world.

ACKNOWLEDGMENTS

We are grateful to Enzo Bruscato and Heather Raslan for their data collection support. We are

indebted to Samraat Pawar, John Stinchcombe, and two anonymous reviewers for their feedback

on earlier versions of our manuscript. AMS was supported by NSF DEB award 2306183. DLC

acknowledges grant support for the Tetrahymena Stock Center by NIH project # 2P40OD010964

and additional support from NSF award # MCB 1947526. JPG acknowledges generous support

from DOE BER DE-SC0020362 (which also supported MHL, AY, and DJW), NSF DEB award

number 2224819, NSF CAREER award number 2337107, and a Simons Foundation Early

Career Fellowship in Microbial Ecology and Evolution LS-ECIAMEE-00001588 (which also

supported MHL and AY).

REFERENCES
Amarasekare, P. & Savage, V. (2012). A Framework for Elucidating the Temperature
Dependence of Fitness. *Am. Nat.*, 179, 178–191.
Angilletta, M.J. (2009). *Thermal Adaptation: A Theoretical and Empirical Synthesis*. OUP
Oxford.
Antiqueira, P.A.P., Petchey, O.L. & Romero, G.Q. (2018). Warming and top predator loss drive
ecosystem multifunctionality. *Ecol. Lett.*, 21, 72–82.
Arrigo, K.R. (2005). Marine microorganisms and global nutrient cycles. *Nature*, 437, 349–355.
Barbour, M.A. & Gibert, J.P. (2021). Genetic and plastic rewiring of food webs under climate
change. *J. Anim. Ecol.*, 90, 1814–1830.
Bar-On, Y.M., Phillips, R. & Milo, R. (2018). The biomass distribution on Earth. *Proc. Natl. Acad.*
Sci., 115, 6506–6511.
Bideault, A., Loreau, M. & Gravel, D. (2019). Temperature Modifies Consumer-Resource
Interaction Strength Through Its Effects on Biological Rates and Body Mass. *Front. Ecol.*
Evol., 7.
Bond-Lamberty, B. (2018). New Techniques and Data for Understanding the Global Soil
Respiration Flux. *Earths Future*, 6, 1176–1180.
Caruso, C.M., Martin, R.A., Sletvold, N., Morrissey, M.B., Wade, M.J., Augustine, K.E., *et al.*
(2017). What Are the Environmental Determinants of Phenotypic Selection? A Meta-
analysis of Experimental Studies. *Am. Nat.*, 190, 363–376.
Cullis, B.R., Smith, A.B. & Coombes, N.E. (2006). On the design of early generation variety
trials with correlated data. *J. Agric. Biol. Environ. Stat.*, 11, 381–393.
Davidson, E.A. & Janssens, I.A. (2006). Temperature sensitivity of soil carbon decomposition
and feedbacks to climate change. *Nature*, 440, 165–173.

490 DeLong, J.P., Gibert, J.P., Luhring, T.M., Bachman, G., Reed, B., Neyer, A., *et al.* (2017). The
491   combined effects of reactant kinetics and enzyme stability explain the temperature
492   dependence of metabolic rates. *Ecol. Evol.*, 7, 3940–3950.
493 Dimond K.L., Zufall R.A (2016). Hidden genetic variation in the germline genome of
494   Tetrahymena thermophila. J Evol Biol 29(6):1284-92
495 Doerder, F.P. (2014). Abandoning sex: multiple origins of asexuality in the ciliate Tetrahymena.
496   *BMC Evol. Biol.*, 14, 112.
497 Enquist, B.J., Norberg, J., Bonser, S.P., Violle, C., Webb, C.T., Henderson, A., *et al.* (2015).
498   Chapter Nine - Scaling from Traits to Ecosystems: Developing a General Trait Driver
499   Theory via Integrating Trait-Based and Metabolic Scaling Theories. In: *Advances in*
500   *Ecological Research*, Trait-Based Ecology - From Structure to Function (eds. Pawar, S.,
501   Woodward, G. & Dell, A.I.). Academic Press, pp. 249–318.
502 Falconer, D.S. (1996). *Introduction To Quantitative Genetics 4th Edition*.
503 Falkowski, P.G. (1994). The role of phytoplankton photosynthesis in global biogeochemical
504   cycles. *Photosynth. Res.*, 39, 235–258.
505 Falkowski, P.G., Fenchel, T. & Delong, E.F. (2008). The Microbial Engines That Drive Earth's
506   Biogeochemical Cycles. *Science*, 320, 1034–1039.
507 Frankham, R. (2005). Stress and adaptation in conservation genetics. *J. Evol. Biol.*, 18, 750–
508   755.
509 Frazier, M.R., Huey, R.B. & Berrigan, D. (2006). Thermodynamics Constrains the Evolution of
510   Insect Population Growth Rates: "Warmer Is Better." *Am. Nat.*, 168, 512–520.
511 Gao, Z., Karlsson, I., Geisen, S., Kowalchuk, G. & Jousset, A. (2019). Protists: Puppet Masters
512   of the Rhizosphere Microbiome. *Trends Plant Sci.*, 24, 165–176.
513 Ghalambor, C.K., McKAY, J.K., Carroll, S.P. & Reznick, D.N. (2007). Adaptive versus non-
514   adaptive phenotypic plasticity and the potential for contemporary adaptation in new
515   environments. *Funct. Ecol.*, 21, 394–407.
516 Gibert, J.P., Dell, A.I., DeLong, J.P. & Pawar, S. (2015). Chapter One - Scaling-up Trait
517   Variation from Individuals to Ecosystems. In: *Advances in Ecological Research*, Trait-
518   Based Ecology - From Structure to Function (eds. Pawar, S., Woodward, G. & Dell, A.I.).
519   Academic Press, pp. 1–17.
520 Gibert, J.P., Grady, J.M. & Dell, A.I. (2022). Food web consequences of thermal asymmetries.
521   *Funct. Ecol.*, 36, 1887–1899.
522 Gibert, J.P., Wieczynski, D.J., Han, Z.-Y. & Yammine, A. (2023). Rapid eco-phenotypic
523   feedback and the temperature response of biomass dynamics. *Ecol. Evol.*, 13, e9685.
524 Gillooly, J.F., Brown, J.H., West, G.B., Savage, V.M. & Charnov, E.L. (2001). Effects of size and
525   temperature on metabolic rate. *Science*, 293, 2248–2251.
526 Golmulkiewicz, R., Kingsolver, J.G., Carter, P.A. & Heckman, N. (2018). Variation and Evolution
527   of Function-Valued Traits. *Annu. Rev. Ecol. Evol. Syst.*, 49, 139–64.
528 Hadfield, J.D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models:
529   The MCMCglmm R Package. *J. Stat. Softw.*, 33, 1–22.
530 Haller, B.C. & Hendry, A.P. (2014). Solving the paradox of stasis: squashed stabilizing selection
531   and the limits of detection. *Evol. Int. J. Org. Evol.*, 68, 483–500.
532 Hansen, T.F. & Houle, D. (2008). Measuring and comparing evolvability and constraint in
533   multivariate characters. *J. Evol. Biol.*, 21, 1201–1219.
534 Hoffmann, A.A. & Sgrò, C.M. (2011). Climate change and evolutionary adaptation. *Nature*, 470,
535   479–485.
536 IPCC. (2023). *Climate Change 2021 – The Physical Science Basis: Working Group I*
537   *Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate*
538   *Change*. Cambridge University Press, Cambridge.

Jacob, S. & Legrand, D. (2021). Phenotypic plasticity can reverse the relative extent of intra-
and interspecific variability across a thermal gradient. *Proc. R. Soc. B Biol. Sci.*, 288,
20210428.
Kingsolver, J. & Huey, R. (2008). Size, temperature, and fitness: Three rules. *Evol. Ecol. Res.*,
10, 251–268.
Kling, J.D., Lee, M.D., Walworth, N.G., Webb, E.A., Coelho, J.T., Wilburn, P., *et al.* (2023). Dual
thermal ecotypes coexist within a nearly genetically identical population of the unicellular
marine cyanobacterium Synechococcus. *Proc. Natl. Acad. Sci.*, 120, e2315701120.
Kontopoulos, D.-G., Smith, T.P., Barraclough, T.G. & Pawar, S. (2020). Adaptive evolution
shapes the present-day distribution of the thermal sensitivity of population growth rate.
PLOS Biol., 18, e3000894.
Lande, R. (1976). Natural Selection and Random Genetic Drift in Phenotypic Evolution.
Evolution, 30, 314–334.
Lande, R. (1979). Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain:body
Size Allometry. *Evolution*, 33, 402–416.
Lande, R. (1982). A Quantitative Genetic Theory of Life History Evolution. *Ecology*, 63, 607–
615.
Lande, R. & Arnold, S.J. (1983). The measurement of selection on correlated characters.
Evolution, 37, 1210–1226.
Lynn, D.H. & Doerder, F.P. (2012). The life and times of Tetrahymena. *Methods Cell Biol.*, 109,
9–27.
MacColl, A.D.C. (2011). The ecological causes of evolution. *Trends Ecol. Evol.*, 26, 514–522.
Malusare, S.P., Zilio, G. & Fronhofer, E.A. (2023). Evolution of thermal performance curves: A
meta-analysis of selection experiments. *J. Evol. Biol.*, 36, 15–28.
Melo, D., Garcia, G., Hubbe, A., Assis, A.P. & Marroig, G. (2015). EvolQG - An R package for
evolutionary quantitative genetics. *F1000Research*, 4, 925.
Montagnes, D.J.S., Wang, Q., Lyu, Z. & Shao, C. (2022). Evaluating thermal performance of
closely related taxa: Support for hotter is not better, but for unexpected reasons. *Ecol.*
Monogr., 92, e1517.
Murphy, K. (2012). *Machine learning: a probabilistic perspective*. MIT Press, Cambridge, MA.
NOAA. (2024). *NOAA NCEI U.S. Climate Normals Quick Access*. Available at:
https://www.ncei.noaa.gov/access/us-climate-normals/#dataset=normals-
monthly&timeframe=30&location=MA&station=USW00014739. Last accessed 23
January 2024.
Padfield, D. (2023). Robust Non-Linear Regression using AIC Scores.
Partridge, L. & Harvey, P.H. (1988). The Ecological Context of Life History Evolution. *Science*,
241, 1449–1455.
Pauls, S.U., Nowak, C., Bálint, M. & Pfenninger, M. (2013). The impact of global climate change
on genetic diversity within populations and species. *Mol. Ecol.*, 22, 925–946.
Pawar, S., Dell, A.I. & Savage, V.M. (2015). Chapter 1 - From Metabolic Constraints
on Individuals to the Dynamics of Ecosystems. In: *Aquatic Functional Biodiversity* (eds.
Belgrano, A., Woodward, G. & Jacob, U.). Academic Press, San Diego, pp. 3–36.
Phillips, B.L., Llewelyn, J., Hatcher, A., Macdonald, S. & Moritz, C. (2014). Do evolutionary
constraints on thermal performance manifest at different organizational scales? *J. Evol.*
Biol., 27, 2687–2694.
Piepho, H.-P. & Möhring, J. (2007). Computing Heritability and Selection Response From
Unbalanced Plant Breeding Trials. *Genetics*, 177, 1881–1888.
Price, G.R. (1972). Extension of covariance selection mathematics. *Ann. Hum. Genet.*, 35, 485–
490.
Rezende, E.L. & Bozinovic, F. (2019). Thermal performance across levels of biological
organization. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 374, 20180549.

Rocca, J.D., Yammine, A., Simonin, M. & Gibert, J.P. (2022). Protist Predation Influences the
Temperature Response of Bacterial Communities. *Front. Microbiol.*, 13.

Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of Body
Size and Temperature on Population Growth. *Am. Nat.*, 163, 429–441.

Seebacher, F., Ducret, V., Little, A.G. & Adriaenssens, B. (2015). Generalist–specialist trade-off
during thermal acclimation. *R. Soc. Open Sci.*, 2, 140251.

Seebacher, F. & Little, A.G. (2021). Plasticity of Performance Curves in Ectotherms: Individual
Variation Modulates Population Responses to Environmental Change. *Front. Physiol.*,
12.

Sinclair, B.J., Marshall, K.E., Sewell, M.A., Levesque, D.L., Willett, C.S., Slotsbo, S., *et al.*
(2016). Can we predict ectotherm responses to climate change using thermal
performance curves and body temperatures? *Ecol. Lett.*, 19, 1372–1385.

Singleton, A.L., Liu, M.H., Votzke, S., Yammine, A. & Gibert, J.P. (2021). Increasing
temperature weakens the positive effect of genetic diversity on population growth. *Ecol.*
Evol., 11, 17810–17816.

Smith, T.P., Clegg, T., Bell, T. & Pawar, S. (2021). Systematic variation in the temperature
dependence of bacterial carbon use efficiency. *Ecol. Lett.*, 24, 2123–2133.

Stefan, H.G. & Preud'homme, E.B. (1993). Stream Temperature Estimation from Air
Temperature. *JAWRA J. Am. Water Resour. Assoc.*, 29, 27–45.

Stell, E., Warner, D., Jian, J., Bond-Lamberty, B. & Vargas, R. (2021). Spatial biases of
information influence global estimates of soil respiration: How can we improve global
predictions? *Glob. Change Biol.*, 27, 3923–3938.

Stinchcombe, J.R., Agrawal, A.F., Hohenlohe, P.A., Arnold, S.J. & Blows, M.W. (2008).
Estimating Nonlinear Selection Gradients Using Quadratic Regression Coefficients:
Double or Nothing? *Evolution*, 62, 2435–2440.

Stinchcombe, J.R., Simonsen, A.K. & Blows, Mark.W. (2014). Estimating uncertainty in
multivariate responses to selection. *Evolution*, 68, 1188–1196.

Tarkington, J., Zhang, H., Azevedo, R.B.R. & Zufall, R.A. (2023). Sex, amitosis, and evolvability
in the ciliate Tetrahymena thermophila. *Evolution*, 77, 36–48.

Tassone, S.J., Besterman, A.F., Buelo, C.D., Ha, D.T., Walter, J.A. & Pace, M.L. (2023).
Increasing heatwave frequency in streams and rivers of the United States. *Limnol.*
Oceanogr. Lett., 8, 295–304.

US EPA, O. (2021). *Climate Change Indicators: Lake Temperature*. Available at:
https://www.epa.gov/climate-indicators/climate-change-indicators-lake-temperature. Last
accessed 13 December 2024.

Voronov, D.A. (2005). Calculating the intrinsic growth rate: comparison of definition and model.
Zh. Obshch. Biol., 66, 425–430.

Wieczynski, D.J., Singla, P., Doan, A., Singleton, A., Han, Z.-Y., Votzke, S., *et al.* (2021).
Linking species traits and demography to explain complex temperature responses
across levels of organization. *Proc. Natl. Acad. Sci.*, 118, e2104863118.

Wieczynski, D.J., Yoshimura, K.M., Denison, E.R., Geisen, S., DeBruyn, J.M., Shaw, A.J., *et al.*
(2023). Viral infections likely mediate microbial controls on ecosystem responses to
global warming. *FEMS Microbiol. Ecol.*, 99, fiad016.

Zufall, R.A., Dimond, K.L. & Doerder, F.P. (2013). Restricted distribution and limited gene flow
in the model ciliate Tetrahymena thermophila. *Mol. Ecol.*, 22, 1081–1091.

640 **FIGURES**
641 **Figure 1**

Figure 2

Figure 3

Figure 4

FIGURE LEGENDS

Figure 1 (a) General shape of the r-TPC. (b) r-TPC shape parameters. Blue-colored shape parameters, measured in this study, include r_{peak}, E_a, CT_{min} and T_{opt}. (c) Top: environmental variation (E) in *r* comes from expressing different *r* across temperatures but not genotypes, and leads to overlapping reaction norms. Middle: genetic variation (G) in *r* results from expressing different *r* across genotypes, but not temperatures, and leads to additive shifts in reaction norm intercepts, but not slopes. Bottom: gene-by-environment interactions (G × E) results from patterns expression of expression of r changing across temperature, leading to shifts in slopes of reaction norms. (d) Because-rTPCs are multi-temperature reaction norms of *r*, classic quantitative genetics can explain how E, G, G × E in *r* influence r-TPC shape: E in *r* produces fully overlapping r-TPCs (top), G in *r* result in non-overlapping r-TPCs with genotype-specific intercepts but not slopes (middle), and G × E leads to r-TPCs that vary in intercepts and slopes (bottom). (e) These r-TPC shape signatures create variation in r-TPC shape parameters: overlapping r-TPCs show no variation in shape parameters (top), additive shifts in r-TPC intercepts lead to additive (heritable) variation in r_{peak}, CT_{min}, and CT_{max} (middle), while slope and intercept variation in r-TPCs creates variation in all shape parameters.

Figure 2 (a) Observed r-TPCs for all 22 genotypes. Dots represent observed *r* values, bold lines represent Sharpe-Schoolfield model fits, and dash lines represent average TPCs across all experimental genotypes. (b) All 22 TPCs are superimposed and dashed lines represent the average r-TPC. Inset: amount of variation due to residual, G × E, G, and E variation in *r*.

Figure 3 (a) Estimated adaptive landscape across temperatures (i.e., change in fitness with a change in the underlying shape parameter) for r_{peak}. Color indicates temperature bins (blue: 10—20ºC, yellow: 20—30ºC, red: 30—40ºC) (b) As in a, but for E_a. (c) As in a, but for CT_{min}. (d) As in a, but for T_{opt}. (e) Observed genetic associations between shape parameters. Each dot is a genotype color-coded as in Fig. 2. In gray, 95% confidence ellipses. ρ represents correlation coefficients. (f) Estimated multivariate selection coefficient (β, 95% maximum density intervals) for all shape parameters across temperatures, or eco-evo landscapes(MacColl 2011). (g) Predicted evolutionary change (Δz, 95% maximum density intervals) for all shape parameters, across temperatures.

Figure 4 (a) G × E variation in r-TPCs leads to differential growth of each genotype across temperatures. (b) Differential growth across temperatures leads to differential selection across environments and rapid shifts in genetic frequencies across temperatures. (c) First column: Differential Interference Contrast (DIC) microscopy for two genotypes of the protist *Tetrahymena thermophila* (CU4106 and AXS). Second column: fluorescence microscopy image overlayed on DIC. Only AXS fluoresces (green) due to the expression of Yellow Fluorescent Protein (YFP). Third column: as in the second column, but for autofluorescence (A. Flu, in pink),

which both genotypes exhibit. Fourth column: Overlayed DIC, YFP and A.Flu images showing
how the different strains fluoresce once all sources of fluorescence are accounted for. (d) r-TPC
for genotypes CU4106 and AXS. Inset: Measures of relative fitness for both CU4106 and AXS.
This predicts an increase in AXS frequency relative to CU4106 at intermediate temperatures
relative to low or high temperatures. (e) Observed genetic frequencies across temperatures. (f)
Predicted genetic frequencies across temperatures.