Waves of light at the bottom of the ocean: insights into the luminous systems of three Pennatuloidea (Anthozoa)

Laurent Duchatelet[1,*,$], Gabriela A. Galeazzo[2,$], Constance Coubris[1], Laure Bridoux[3], René Rezsohazy[3], Marcelo R.S. Melo[4], Martin Marek[5], Sam Dupont[6,7], Anderson G. Oliveira[8], Jérôme Delroisse[9,10]

[1] Marine Biology Laboratory, Earth and Life Institute, University of Louvain – UCLouvain, Belgium

[2] Departamento de Oceanografia Física, Química e Geológica, Instituto Oceanográfico, Universidade de São Paulo, São Paulo, Brazil

[3] Animal Molecular and Cellular Biology, Louvain Institute of Biomolecular Science and Technology, University of Louvain – UCLouvain, Belgium

[4] Departamento de Oceanografia Biológica, Instituto Oceanográfico, Universidade de São Paulo, São Paulo, Brazil

[5] Loschmidt Laboratories, Department of Experimental Biology and RECETOX, Masaryk University, 625 00 Brno, Czech Republic

[6] Department of Biological & Environmental Sciences, University of Gothenburg, Fiskebäckskil, Sweden

[7] International Atomic Energy Agency, Environment Laboratories, Principality of Monaco, Monaco

[8] Department of Chemistry and Biochemistry, Yeshiva University, 245 Lexington Ave, New York, NY 10016, USA

[9] Biology of Marine Organisms and Biomimetics, Research Institute for Biosciences, University of Mons, Belgium

[10] Cellular and Molecular Immunology, University of Liège, Belgium

* corresponding authors: Duchatelet Laurent, laurent.duchatelet@uclouvain.be

$ Both authors contributed equally

Abstract

Bioluminescence is the production of visible light by living organisms. This process occurs biochemically through the primary oxidation of specific luciferin substrates, facilitated by enzymes known as luciferases. Moreover, the bioluminescent reaction often involves additional elements like auxiliary proteins or cofactors. Auxiliary proteins, such as fluorescent proteins and coelenterazine-binding proteins, for example, may serve various roles, such as altering the wavelength of the emitted light–or stabilizing the highly reactive and unstable luciferin, respectively. Calcium has been demonstrated to serve as an essential ion in luminescence processes across various species. Despite the existence of numerous bioluminescent species, only a fraction of bioluminescent systems and their associated molecular actors have been thoroughly studied. Within anthozoans, the bioluminescence system of the *Renilla* genus stands out as the sole extensively characterized system, involving a coelenterazine-dependent luciferase, a calcium-dependent coelenterazine-binding protein, and a green fluorescent protein.

Through a multidisciplinary approach, we investigated the bioluminescent system of three sea pen species: *Pennatula phosphorea, Funiculina quadrangularis*, and *Anthoptilum murrayi* (Pennatuloidea, Anthozoa, Cnidaria). We first confirmed or characterized the emission spectra of these species, elucidated the coelenterazine-based reaction, and generated reference transcriptomes for the two first species. Our transcriptomic analysis revealed transcripts encoding luciferases, fluorescent proteins, and coelenterazine-binding proteins, also shedding light on their expression profiles. Immunodetection confirmed the presence of luciferase in *P.*

phosphorea and *F. quadrangularis*, while autofluorescence patterns attributed to green
fluorescent protein or the coelenterazine were observed in *P. phosphorea*. Furthermore, a
recombinant luciferase was expressed for *A. murrayi*. Our findings also underscore the essential
role of calcium ions in bioluminescence, possibly associated with the mechanism of
coelenterazine binding and substrate release. Finally, we propose a mechanistic model of
bioluminescence in anthozoans, laying the groundwork for further functional characterization
and ecological exploration of these sessile and benthic luminous organisms.

Keywords: Bioluminescence, Luciferase, Coelenterazine, Luciferin-binding protein, Calcium,
Luminous system, Cnidaria, Soft coral

Introduction
Bioluminescence, the direct production of visible light by a living organism, involves
the oxidation of a luciferin catalyzed by a luciferase enzyme. In some cases, luciferase and
luciferin are covalently bound, forming a complex unit called "photoprotein." While the
classical luciferin/luciferase system may not require a cofactor, the photoprotein system
requires ions or cofactor to trigger light emission [1]. Luciferases *sensu lato* (*i.e.,* luciferases
stricto sensu and photoproteins) are classically considered taxon-specific (*i.e.,* each clade being
characterized by its own luciferase) [2]. However, homologous luciferases can sometimes be
used by phylogenetically distant luminous organisms [3]. At least 12 distinct types of luciferase
sensu lato were recently described [3].
When the photogenic reaction takes place, the generated light is defined by a color that
can differ depending on the luciferin but also on the amino acid sequence and structure of the
luciferase [4-7]. In some cases, the light emitted by an organism results from an interaction
between the luciferase and a fluorescent protein [4]. The latter absorbs the photons emitted and
enters an excited state. It then re-emits the light but converts it to longer wavelengths, returning
to its ground state. For example, the Green Fluorescent Protein (GFP) re-emits green light after
absorbing the blue light primarily emitted by the luciferin-luciferase reaction [1,5].
Luminous anthozoans are distributed among four orders (*i.e.,* Actiniria, Zoantharia,
Malacalcyonacea, and Scleralcyonacea). To date, Actiniaria, Zoantharia, and Malacalcyonacea
include four luminous families, while Scleralcyonacea includes 18 luminous families mainly
spread into the Pennatuloidea superfamily (**Table S1**). The bioluminescent system of the
Renilla genus (Pennatuloidea) was characterized, followed by the more recent descriptions of
similar bioluminescent molecular actors in different pennatulaceans [8-16] (**Table S1**). *Renilla*
luminous system involves the most widespread luciferin in marine environments, the
coelenterazine   (6-(4-hydroxyphenyl)-2-[(4-hydroxyphenyl)methyl]-8-(phenylmethyl)-7H-
imidazo[1,2-a]pyrazin-3-one),   and   a   coelenterazine-dependent   luciferase   (*R*Luc-type)
[9,10,15]. This luciferase is homologous to bacterial haloalkane dehalogenases, which were
assumed to be horizontally transferred during evolution [3]. Bessho-Uehara et al. (2020)
demonstrated the widespread occurrence of this bioluminescence system across deep-sea
pennatulacean species such as *Distichoptilum gracile*, *Umbellula* sp., *Pennatula* sp., and
Funiculina sp. [12]. Besides, *Renilla*-type luciferases are also found in phylogenetically distant
organisms such as an echinoderm (*i.e., Amphiura filiformis*; [17]) and a tunicate (*i.e., Pyrosoma*
atlanticum; [18]). Luminescence-associated molecules such as coelenterazine-binding protein

(CBP) and GFP have also been revealed in the sea pansy and other pennatulaceans [15,19-24].
Additionally, the calcium ion cofactor has been demonstrated to be indirectly involved in light
production in *Renilla reniformis* and *Veretillum cynomorium* [25,26]. CBP depends on calcium
ions as cofactors to release the luciferin, allowing light production in *Renilla* [27].
Light production in sea pens occurs within specific endodermal cells (*i.e.,* photocytes)
spread, depending on the species, in the tissue of autozooid and siphonozooid polyps [28-30].
These photocytes often present a green autofluorescence [31,32]. To date, studied
bioluminescent pennatulaceans display a nervous catecholaminergic control of light emission,
with adrenaline and noradrenaline-triggered waves of flashes [33,34]. Although no behavioral
experimental evidence was performed, the ecological function of pennatulacean luminescence
may be an aposematic signal, avoiding predation by misdirection or burglar alarm [2,35].
Current data and gaps in anthozoan bioluminescence knowledge are presented in **Table
S1**.
Through a multidisciplinary study of three different Pennatuloidea species, including
luminometric assays, transcriptome analyses, and immunohistological techniques, results
unveil the bases of the bioluminescent system of the three soft coral sea pens. Cross-reaction
luminometric results underline the involvement of a *Renilla*-like luciferase for both shallow
water species and coelenterazine as the substrate for all three bioluminescent systems. Besides,
P. phosphorea and *A. murrayi de novo* transcriptome analyses demonstrate (*i*) the homology of
both retrieved luciferase sequences with *Renilla* luciferase, (*ii*) the *in silico* presence of CBPs
and a GFP homologous to the one found in other bioluminescent anthozoans. The expression
sites of the luciferase within autozooids and siphonozooids of *P. phosphorea* and autozooids of
F. quadrangularis are described. The presence of the GFP is also pinpointed for *P. phosphorea*
through comparative microscopy. Finally, the involvement of calcium in the light emission
process of *P. phosphorea* and *F. quadrangularis* is described.
117
118   **Materials and methods**
119
Specimen collection
In July 2022 and 2023, common sea pens, *Pennatula phosphorea* (n = 30), were
collected in the Gullmarsfjord, Sweden, using a small 1 m aperture dredge at 35 to 40 m depth.
Similarly, tall sea pens, *Funiculina quadrangularis* (n = 11)*,* were sampled with the same
methodology at 40 m depth in July 2023. Animals were brought back to the Kristineberg Marine
Research Station (Fiskebäckskil, Sweden) and maintained in dark conditions in fresh, running,
deep-sea water pumped from the adjacent fjord.
Specimens of *Anthoptilum murrayi* were collected during the DEEP OCEAN expedition
off the southeast coast of Brazil aboard the R/V Alpha Crucis. The collection was carried out
using a demersal trawl net with doors 19 m long and a mouth opening of 2.5 m at an average
depth of 1,100 m. After hoisting the net, all collected organisms were sorted, and *A. murrayi*
corals were identified and promptly frozen in liquid nitrogen. These samples were maintained
at the Oceanographic Institute of the University of São Paulo (São Paulo, Brazil) in an -80 ºC
ultra-freezer.
134
135   *Dissection*

Specimens of *P. phosphorea* and *F. quadrangularis* were anesthetized by immersion in MgCl$_2$ (3.5%) for 30 minutes [34]. Then, for each tested *Pennatula* specimen, (*i*) pinnules were dissected and weighed, and (*ii*) the rachis was divided into three equivalent portions and weighed. For *Funiculina* specimens, the rachis bearing the polyps was split into 3 cm long segments. Sea pen pinnule/rachis were either directly used for biochemistry assays or rinsed for 3 hours in fresh running deep-sea water for pharmacological calcium assays. Other specimens were directly fixed in PFA 4% phosphate buffer saline (123 mM NaCl, 2.6 mM KCl, 12.6 mM Na$_2$HPO$_4$, 1.7 mM KH$_2$PO$_4$, pH 7.4) for further immunodetection techniques.

Due to the constraints associated with harvesting organisms at depth (*e.g.*, trawling damage), it was impossible to maintain *A. murrayi* specimens in an aquarium. Therefore, all measurements are performed on frozen samples.

Light emission spectrum of Anthoptilum murrayi

The bioluminescence spectrum of *A. murrayi* was obtained using pieces of frozen coral that were thawed and hydrated with deionized water. The spectrum was recorded using a cooled CCD camera (LumiF SpectroCapture AB-1850). The measurements were conducted in triplicate with an exposure time of 2 minutes at room temperature.

Luminescent system biochemistry

For biochemistry analysis, 11 specimens of *P. phosphorea* were used. Tests were done for pinnules (n = 40), rachis (n = 10), and peduncles (n = 10). Pinnule locations along the axis were noticed and classified as upper, middle, and lower depending on the attachment position on the rachis, as in Duchatelet et al., 2023 [34]. Five specimens of *F. quadrangularis* were used just after dissection for these biochemical assays. Measurements were done independently of the polyp-bearing rachis locations among colonies (n = 10). Deep-frozen tissues of two specimens of *A. murrayi* were used for this analysis. For the former two species, the central rigid calcified axis of the rachis was removed before starting experiments.

Measurements of light emission were performed in a dark room using a FB12 tube luminometer (Tirtertek-Berthold, Pforzheim, Germany) calibrated using a standard 470 nm light source (Beta light, Saunders Technology, Hayes, UK). Light responses were recorded using FB12-Sirius PC Software (Tirtertek-Berthold). Light emission was characterized as follows: (*i*) the maximum light intensity (Lmax), expressed in megaquanta per second (10^9 q s^{-1}), and (*ii*) the total amount of light emitted (Ltot) over 3 minutes, expressed in megaquanta. All data were standardized per unit of mass (g).

Luciferase and coelenterazine assays

For the luciferase assay, *P. phosphorea* pinnule and rachis and *F. quadrangularis* rachis were placed in 200 µl of Tris buffer (20 mM Tris, 0.5 mM NaCl; pH 7.4) and crushed with mortar and pestle until a homogenized extract was obtained; 20 and 40 µl of the extract was diluted in 180 and 160 µl Tris buffer, respectively. The diluted *P. phosphorea* and *F. quadrangularis* luciferase solutions were injected into two different tubes filled with 5 µl of 1/200 stock solution of coelenterazine (Prolume Ltd, Pinetop, AZ, USA) in cold methanol (1OD at 430 nm) diluted in 195 µl of Tris buffer. Two measures of Lmax were recorded and averaged

to calculate the maximal light decay rate corresponding to the luciferase activity expressed in
10^9 q s^{-1} g^{-1} [1].

For coelenterazine detection, *P. phosphorea* pinnule, rachis, and *F. quadrangularis*
rachis were put into 200 µl of cold argon-saturated methanol and crushed with mortar and
pestle. Then, 5 µl of the methanolic extract was injected into a tube filled with 195 µl of Tris
buffer and placed in the luminometer. Afterward, 200 µl of *Renilla* luciferase solution
constituted 4 µl of *Renilla* luciferase (Prolume Ltd., working dilution of 0.2 g l^{-1} in a Tris-HCl
buffer 10 mM, NaCl 0.5 M, BSA 1%; pH 7.4) and 196 µl of Tris buffer was injected into the
luminometer tube. The Ltot was recorded and used to calculate the amount of coelenterazine
contained in a gram of pinnule (ng g^{-1}), assuming that 1 ng of pure coelenterazine coupled with
Renilla luciferase emits 2.52×10^{11} photons [1].

A. murrayi tissues were processed using a Potter-Elvehjem homogenizer in 2 mL of Tris
buffer (50 mM Tris-HCl, pH 8.0). Following homogenization, the mixture was centrifuged at
15,000 x g for 10 minutes at 4°C, after which the pellet was discarded. The supernatant was
used for light emission assays. For each assay, the mixture comprised 100 µL of supernatant,
397 µL of Tris buffer (50 mM Tris, pH 7.4), and 3 µL of coelenterazine (Prolume Ltd),
achieving a final volume of 500 µL with a coelenterazine final concentration of 6 µM.

Long-term light production
P. phosphorea specimens (n = 12) were maintained in tanks filled with circulating
artificial seawater (ASW; 400 mM NaCl, 9.6 mM KCl, 52.3 mM MgCl$_2$, 9.9 mM CaCl$_2$, 27.7
mM Na$_2$SO$_4$, 20 mM Tris; pH 8.2) at temperatures following natural temperature variations
encountered in the native fjord (https://www.weather.mi.gu.se/kristineberg/en/data.shtml) with
a 12-12 hours photoperiod. Sea pens were weekly fed with REEF LIFE plancto (Aqua Medic,
Germany), a food devoted to any coelenterazine traces. After six and twelve months, potassium
chloride depolarization, coelenterazine content, and luciferase activity assays were performed
on eight and four specimens, respectively, with two replicates per specimen. Experiments
occurred as described above for the luciferase and coelenterazine assays. Total depolarization
through KCl application allows rapid estimation of the luminous ability of specimens. For KCl
experiments, luminescence inductions were performed on pinnule and rachis portions placed in
a tube luminometer filled with 500 µL of ASW. Then, the light emission was triggered with the
addition of 500 µL of a KCl solution (400 mM KCl, 52.3 mM MgCl$_2$, 9.9 mM CaCl$_2$, 27.7 mM
Na$_2$SO$_4$, 20 mM Tris; pH 8.2), and Ltot was recorded over 3 minutes. Recorded luciferase
activity, luciferin contents, and KCl responses were compared with wild-caught measurements.

De novo transcriptome analyses
Pinnule, rachis, and peduncle tissues of a single *P. phosphorea* specimen were dissected
and directly immersed in a permeabilizing RNAlaterTM-Ice (Life Technologies) solution for
one night at -20°C, following the manufacturer's protocol. Subsequently, samples were stored
at -80°C and processed for RNA extractions. Total RNA was extracted using the trizol-based
method. The quality of RNA extracts (RIN value, fragment length distribution, and 28S/18S
ratio) and their concentrations were assessed using an Agilent 2100 bioanalyzer. The BGI
company (Bejing Genomics Institute, Hong Kong) performed cDNA library preparation and
sequencing using a similar procedure previously described [36-38]. High-throughput

sequencing was conducted via the BGISEQ-500 platform to generate 100 bp paired-end reads.
To exclude low-quality sequences, the raw reads were filtered by removing (*i*) reads with more
than 20% of the qualities of the base lower than 10, (*ii*) reads only containing the adaptor
sequence, and (*iii*) reads containing more than 5% of unknown nucleotide "N." Quality control
of reads was performed using FastQC software [39]. For *P. phosphorea,* a reference *de novo*
transcriptome assembly was then created from the remaining clean reads obtained from pinnule,
rachis, and peduncle tissues using the Trinity software [40] (Trinity-v2.5.1; min_contig_length
150, CPU 8, min_kmer_cov 3, min_glue 3, SS_lib_type RF, bfly_opts'-V 5, edge-thr=0.1,
stderr'). The TGICL software was then used to reduce the transcriptome redundancy by the
further assembly of the contigs into a single set of longer, non-redundant, and more complete
consensus unigenes [41] (Tgicl-v2.5.1; -l 40 -c 10 -v 25 -O '-repeat_stringency 0.95 -minmatch
35 -minscore 35'). Unigenes, then defined as non-redundant assembled sequences obtained
from assembly and/or clustering [42], can form clusters in which the similarity among
overlapping sequences is superior to 70% or singletons that are unique unigenes. For all
transcriptomes, unigene expression was evaluated using the "Fragments per kilobase of the
transcript, per million fragments sequenced" (FPKM) method [36-37,42]. To obtain annotation
for transcriptomes, unigenes were aligned to NCBI Nucleotide (NT), NCBI protein (NR),
EuKariotic Orthologous groups (KOG), Kyoto Encyclopedia of Genes and Genomes (KEGG),
and UniProtKB/Swiss-Prot databases using Blastn, Blastx [43], and Diamond [44]. Blast2GO
[45] with NR annotation results was used to get Gene Ontology annotation according to
molecular function, biological process, and cellular component ontologies. InterPro [46] was
also used to annotate unigenes depending on functional analyses of protein sequences,
clustering them into families and predicting the presence of domains or essential amino acid
residues. The candidate coding area among unigenes was assessed using Transdecoder
(https://transdecoder.github.io). The longest ORFs (open reading frame) were extracted, and
Blast searched the Pfam protein homologous sequences to SwissProt to predict coding regions.
A similar procedure was used for *Anthoptilum*. Whole sea pen tissues (pinnule and
peduncle) were dissected, frozen in liquid nitrogen, and stored at -80°C. The total RNA from
the sea pen was extracted using the RNeasy Plant Mini Kit (QIAGEN), following the extraction
steps according to the manufacturer's instructions. The RNA samples were eluted in DEPC-
treated water in two 16 μL steps. The RNA samples were treated with DNaseI enzyme
(Invitrogen) to remove potential genomic DNA contamination. Subsequently, to clean
impurities and residual DNaseI reaction remnants, they underwent a column cleanup process,
following the instructions of the RNeasy Plant Mini Kit (QIAGEN). The concentrations of the
RNA samples were estimated by fluorescence using the Qubit fluorometer (Thermo Fisher
Scientific). The integrity of the RNA samples was confirmed by agarose gel electrophoresis
(1%) stained with SYBR Safe (Invitrogen), and the RNA content was dried at 30°C for 1 hour
in speed vac mode V-AQ.
The quality of RNA extracts (RIN value, fragment length distribution, and 28S/18S
ratio) and their concentration were assessed by RNA concentrations were assessed using an
Agilent 2100 bioanalyzer. After the QC procedures, mRNA was enriched using oligo(dT)
beads. First, the mRNA was fragmented randomly by adding a fragmentation buffer. Then, the
cDNA was synthesized using an mRNA template and random hexamers primer, after which a
custom second-strand synthesis buffer (Illumina), dNTPs, RNase H, and DNA polymerase I

were added to initiate the second-strand synthesis. Second, after a series of terminal repairs, A
ligation, and sequencing adaptor ligation, the double-stranded cDNA library was completed
through size selection and PCR enrichment. The GenOne Biotechnologies company (Brazil)
performed cDNA library preparation and sequencing (150 bp paired-end reads). To exclude
low-quality sequences, the raw reads were filtered by removing (*i*) reads with more than 50%
of the qualities of the base lower than 5, (*ii*) reads only containing the adaptor sequence, and
(*iii*) reads containing more than 10% of unknown nucleotide "N."

Light emission process-related protein sequence analyses

Potential transcripts of interest were chosen on NCBI online tools according to potential
phylogenetic homologies to identify genes involved in the light production process, such as
luciferase, green fluorescent protein, and coelenterazine-binding protein. These "light emission
process-related genes" were searched within the newly generated *P. phosphorea* transcriptome
using tBlastn analyses (1 hit, E-value < 1e^{-20}). All the retrieved unigenes were individually
reciprocally searched in NCBI NR databases (Reciprocal Blastx; 1hit, E-value < 1e^{-20}). Blast
hits with significant E-values strongly support homologous proteins. *In silico* translation
(ExPASy translate tool, http://expasy.org/tools/dna.html) was performed on the retrieved
sequence from the *P. phosphorea* and *A. murrayi* transcriptomes for all putative candidates.
Multiple alignments were performed for each predicted protein with their respective
homologous "light emission process-related protein" retrieved from the NCBI online tool in
other species via Geneious software [47]. Sequence alignments enabled the identification of
luciferase characteristics features such as the catalytic triads [48].

To validate the *P. phosphorea* luciferase retrieved sequence, primers were designed
based on the *Renilla muelleri* luciferase mRNA (AY015988.1) to amplify the hypothetical
luciferase sequence using Primer Blast software (**Table S2**). A *Pennatula* pinnule was collected
and lysed in 300 µL of 50 mM NaOH for 30 minutes at 95°C with agitation at 800 rpm. The
pH was then adjusted by adding 60 µL of 500 mM Tris-HCl, pH 8. For the PCR amplification,
the reaction mix contained 1U of Expand Long Template (Roche) with the provided buffer, 400
µM of dNTP (#R0191, Life Technologies), and 250 nM of each primer. The amplification was
performed as follows: 95°C for 5 minutes, 35 cycles of denaturation at 95°C for 30 seconds,
hybridization from 52 to 55 °C for 15 seconds, and elongation at 68°C for 45 to 80 seconds.
The final cycle was completed by a final elongation step at 68°C for 7 minutes. Primer pairs
F1-R1, F2-R2, and F8-R2 resulted in amplifications visualized through electrophoresis and
ethidium bromide incorporation. The genomic DNA was purified using the Qiagen PCR
purification kit according to the manufacturer's instructions and subsequently sequenced by
Microsynth.

Phylogenetic analyses

The predicted sequences of the luciferases, the GFP, and the CBPs of *P. phosphorea*
and *A. murrayi* were placed into an anthozoan-focused phylogenetic context using maximum
likelihood phylogenetic reconstructions [49]. *Renilla*-type luciferase sequences from selected
metazoans were collected from public databases based on the literature [12,16,17]. GFP
sequences from cnidarians were collected from public databases using reference literature [50-
53]. CBP sequences from anthozoans were collected from public databases based on the

reference literature [54]. Multiple alignments of all sequences were performed using the
MAFFT algorithm implemented in the Geneious software and trimmed using TrimAL software
[55]. Maximum likelihood phylogenetic analyses were performed using the IQ-tree software
[56]. Before the analyses, ModelTest [57] was used to select the best-fitting evolution model.
Trees were edited using the web tool iTOL.
316
Luciferase and green fluorescent protein presence and localization
To confirm the presence of a *Renilla*-like luciferase, commercial antibodies directed
against *Renilla* luciferase (GTX125851, Genetex). Proteins were extracted from frozen
pinnules, rachis, and peduncle samples. Each sample was homogenized on ice in 1000 µl of 2%
Triton X-100 in phosphate buffer saline (PBS: 10 mM Tris, pH 7,5; 1 mM EDTA, pH 8,0; 100
mM NaCl) supplemented with protease inhibitors (complete–Mini tablets, Roche). The extract
was sonicated and centrifuged at 800 x g for 15 minutes. The supernatant was collected. Protein
concentration in each extract was measured using PierceTM BCA Protein Assay Kit (Thermo
Scientific). Laemmli buffer (Biorad) and β-mercaptoethanol (βMSH, Biorad) were added to
each protein extract, and the proteins were electrophoretically separated at 200 V for 35 minutes
on 12% SDS-PAGE gels. The separated proteins were then electroblotted on a nitrocellulose
membrane. The membrane was incubated overnight with the primary anti-*Renilla* luciferase
antibody and the secondary antibody (ECL HRP conjugated anti-rabbit antibody, Life Sciences,
NA934VS, lot number 4837492) for 1 hour. Antibody detection was performed with the
reagents of the detection kit (HRP Perkin-Elmer, NEL 104) following the manufacturer's
instructions. The dilution for the primary antibody was 1:2000.
The same primary antibody was used to immunolocalize the luciferases within the *P.*
phosphorea pinnules, rachis, peduncle tissues, and the *F. quadrangularis* polyp's tissue. For
in-toto immunohistochemistry, dissected samples were blocked with PBS containing 2% Triton
X-100 and 6% BSA (Amresco). Samples were then incubated for 48 hours with either the anti-
Renilla luciferase antibody diluted 1:200 in PBS containing 1% Triton X-100, 0.01% NaN$_3$,
and 6% BSA. Visualization of luciferase signal was done after 24 hours of incubation of the
samples at RT in the dark with fluorescent dye-labeled secondary antibody (Goat Anti-Rabbit,
Alexa Fluor 594, Life Technologies Limited) diluted 1:500 in PBS containing 1% Triton X-
100, 0.01% NaN$_3$, and 6% BSA. Samples were mounted (Mowiol 4–88, Sigma) on slides and
examined using an epifluorescence microscope (Axio Observer, Zeiss, Oberkochen, Germany)
equipped with Zen microscopy software (Zeiss, Oberkochen, Germany). Control sections were
incubated in PBS containing 1% Triton X-100, 0.01% NaN$_3$, and 6% BSA with no primary
antibody.
346
Calcium assays
Different pharmacological tests were used to investigate the implication of calcium in
the light emission process of *Pennatula* and *Funiculina* (n = 6 for each species). First, the
involvement of calcium concentration was tested using three artificial seawater (ASW)
solutions with different calcium concentrations (0, 10, and 20 mM of CaCl$_2$). To erase any
traces of Ca^{2+} ions in the 0 mM CaCl$_2$ -ASW solution, calcium chelators EGTA (4100, Merck)
and BAPTA (14513, Merck) were added to the solution at 10^{-5} M final concentrations.
Secondly, the effect of a calcium ionophore (A23187; C7522, Merck) was tested to highlight

the potential involvement of calcium storage in the light emission process. Thirdly, the
involvement of calcium was tested in the presence of a previously determined triggering agent
of the light emission in sea pens, the adrenaline at 10^{-5} M. Finally, calcium involvement was
tested on the effect of the potassium chloride depolarization solution usually employed to
trigger the maximum light production in a bioluminescent species.

After 3 hours of rinsing, pinnules of *P. phosphorea* and polyp-bearing rachis of *F.*
quadrangularis were placed in luminometric tubes and subject to various treatments (**Table**
S3). Each sample was pre-incubated for 15 minutes in ASW devoted to calcium (0 mM $CaCl_2$),
and the experiment began. Data was recorded for 15 minutes via the FB12 tube luminometer
(Tirtertek-Berthold, Pforzheim, Germany). Light emission was the total amount of light emitted
(L_{tot}) over 15 min, expressed in megaquanta. All data were standardized per unit of mass (g).
366

Statistical analysis
All statistical analyses were performed with R Studio (version 2023.03.1 + 446, 2022,
Posit Software, USA). Variance normality and equality were tested using the Shapiro–Wilk test
and Levene's test, respectively. When these parametric assumptions were met, a Student t-test
and an ANOVA coupled with Tukey's test were used to perform single or multiple comparisons
between groups. When log transformation did not provide normality and homoscedasticity, the
nonparametric Wilcoxon test and the Kruskal–Wallis test coupled with the Wilcoxon rank-sum
test were used, respectively, to assess whether significant differences were present between two
groups or multiple groups. Each difference was considered to be significant at a minimum *p-*
value < 0.05. Values were graphically illustrated with mean and standard error of the mean
(s.e.m).
378

379 **Results**
380

381 ***Pennatula phosphorea, Funiculina quadrangularis, and Anthoptilum murrayi emit light***
382 ***using a coelenterazine-dependant Renilla-like luciferase.***
383

Biochemistry assays performed on *Pennatula phosphorea* and *Funiculina*
quadrangularis (**Figure 1A, B**) present a similar response pattern. To complete data on the sea
pen light emission spectrum, the *in vivo* light emission spectrum of *A. murrayi* was measured
and revealed a peak wavelength at 513 nm (**Figure 1C**). The cross-reactivity of extracts with
the commercial *Renilla* luciferase highlights the involvement of coelenterazine in the *P.*
phosphorea, *F. quadrangularis*, and *A. murrayi* bioluminescent system (**Figure 1D, E, F**). For
P. phosphorea, measured amounts of coelenterazine present no statistical differences between
areas (ANOVA, *p-value* lower-middle = 0.8122; *p-value* lower-upper = 0.9054; *p-value* upper-
middle = 0.7014). Similarly, no differences were observed between rachis portions (ANOVA,
p-value lower-middle = 0.5905; *p-value* lower-upper = 0.8274; *p-value* upper-middle =
0.4634). Mean coelenterazine content per pinnule and rachis are 110.98 ± 10.45 and $70.00 \pm$
16.79 ng g^{-1}, respectively. Peduncle presents an almost unnoticeable mean value of 0.21 ± 0.01
ng g^{-1}. Comparatively, the mean coelenterazine content of *F. quadrangularis* polyp-bearing
rachis is 3.79 ± 1.20 ng g^{-1}. Cross-reactivity performed on *A. murrayi* also showed that this
system functions as a coelenterazine-based luciferase system, similar to the luminescent

reaction observed in *R. reniformis*. Typical coelenterazine assay curves are presented for each
species in **Figures 1D, E, and F**.

401
402   **Figure 1. Biochemistry assays.** *In situ* pictures of (A) *P. phosphorea* and (B) *F. quadrangularis* in the
403   Swedish Gulmarfjord. (C) The spectrum of the *in vivo* luminescence of *A. murrayi*. Typical luciferin
404   assays for (D) *P. phosphorea* pinnules, (E) *F. quadrangularis* polyp-bearing rachis, and (F) *A. murrayi*
405   whole specimen.
406

Counterpart cross-reaction using synthetic coelenterazine to measure the potential
luciferase activity confirms the implication of a coelenterazine-based luciferase system for *P.*
phosphorea and *F. quadrangularis*. For *P. phosphorea*, luciferase activities of the pinnules do
not present statistical differences considering the pinnule areas (ANOVA, *p-value* lower-
middle = 0.4963; *p-value* lower-upper = 0.2742; *p-value* upper-middle = 0.6251), with a mean
Lmax value of $113.2 \pm 12.7 \ 10^9$ q g^{-1} s^{-1}. Similarly, no statistical differences were observed for

the Lmax values recorded between rachis portions (ANOVA, *p-value* lower-middle = 0.6360;
p-value lower-upper = 0.4438; *p-value* upper-middle = 0.2548), with a mean Lmax value of
$87.9 \pm 16.9 \ 10^9$ q g^{-1} s^{-1}. Finally, tested peduncles present a significantly lower mean Lmax
value ($0.6 \pm 0.4 \ 10^9$ q g^{-1} s^{-1}) than the rachis value (ANOVA, *p-value* = 0.0298). Comparatively,
F. quadrangularis luciferase activity presents a mean Lmax value of $313.8 \pm 99.2 \ 10^9$ q g^{-1} s^{-1}.
418
419 ***Pennatula phosphorea maintains its bioluminescence ability after a captivity period of 1 year***
420 ***without a coelenterazine exogenous supply.***
421
Before the experiment, visual assessments of *P. phosphorea* luminescence were
conducted under dark conditions. These observations confirmed that the species maintained the
ability to produce visible light even after six to twelve months of captivity without any external
sources of coelenterazine-containing food.
Measurements of luminescence parameters underline a general decrease in all three
parameters for both pinnules and the rachis portion (**Figure S1**). For the pinnules, the maximum
light emission presents statistical differences upon KCl application (Kruskal-Wallis test, *p-*
value = 1.8 10^{-5}), with a mean Ltot value after field collection of $361.7 \pm 34.6 \ 10^9$ q g^{-1} s^{-1},
statistically different from the mean value after six months of captivity (Wilcoxon sum-rank
test, *p-value* = 1.5 10^{-6}) but not after 12 months (Wilcoxon sum-rank test, *p-value* = 0.16). The
mean Ltot values after 6 and 12 months are 80.5 ± 21.8 and $223.8 \pm 84.4 \ 10^9$ q g^{-1} s^{-1},
respectively. Coelenterazine content also presents statistical differences (Kruskal-Wallis test,
p-value = 1.3 10^{-5}). The mean coelenterazine content is already drastically reduced after 6
months (30.7 ± 4.5 ng g^{-1}) but remains stable for one year (28.1 ± 4.4 ng g^{-1}) (Wilcoxon sum-
rank test, T0 *VS* T6: *p-value* = 0.0001; T0 *VS* T12: *p-value* = 0.0004), without statistical
differences between the former two (Wilcoxon sum-rank test, *p-value* = 0.91) (**Figure S1**).
Similarly, the luciferase activities decreased over the year and presented statistical differences
(Kruskal-Wallis test, *p-value* = 0.01). Differences occur between the initial Lmax value (113.2
$\pm 12.7 \ 10^9$ q g^{-1} s^{-1}) and the mean Lmax values of 77.0 ± 16.2 and $43.0 \pm 3.7 \ 10^9$ q g^{-1} s^{-1} after
6 and 12 months, respectively (Wilcoxon sum-rank test, T0 *VS* T6: *p-value* = 0.383; T0 *VS* T12:
p-value = 0.013; T6 *VS* T12: *p-value* = 0.383) (**Figure S1**). Similar observations occurred for
the three parameters recorded on the rachis portions of *P. phosphorea* (**Figure S1**).
444
445 ***De novo transcriptomes of Pennatula phosphorea and Anthoptilum murrayi***
446
For *P. phosphorea*, in total, 47.27, 47.27, and 41.96 million raw reads, with a length of
200 bp, were generated from the pinnule, the rachis, and the peduncle libraries, respectively.
Data quality was checked using the FastQC software. The raw reads will be deposited in the
NCBI SRA database and available upon request to the authors. After low-quality reads filtering,
the remaining high-quality reads (*i.e.,* 45.29 for the pinnule transcriptome, 45,53 for the rachis
transcriptome, and 40.29 for the peduncle transcriptome) were used to assemble a reference
transcriptome using the Trinity software. The obtained Trinity-predicted transcripts were
clustered using TGICL to obtain final unigenes.
In total, 49,510 unigenes, *i.e.,* non-redundant unique sequences, were obtained for a
total length of 72,928,901 pb. The average mean length was 1473, and the N50 was 2350.

Within transcriptomes data, 35,439 unigenes for the pinnule dataset, 33,778 unigenes for the
rachis dataset, and 31,001 unigenes for the peduncle dataset were obtained, with a total of
49,510 different unigenes. The length distributions of unigenes are shown in **Figure 2A**, and
numerical data are summarized in **Tables S4** and **S5**.
On a total of 46,258 predicted unigenes, 1,690 were found only in the pinnule
transcriptome, 1,114 only in the rachis transcriptome, and 1,035 only in the peduncle
transcriptome while 36,934 were common to the three transcriptomes. For descriptive purpose,
a comparative gene expression analysis was performed by mapping FPKM values (*e.g.,*
$\log_{10}$(FPKM value pinnule transcriptome) against $\log_{10}$(FPKM value rachis transcriptome))
calculated for all predicted unigenes (**Table S6**). However, it has to be clarified that the
transcriptome data have been generated for new gene discovery, not differential expression
analyses, as no biological or technical replication was performed as a part of the study. The
main represented species within the unigene annotation of the reference transcriptome is the
anthozoan *Dendronephthya gigantea* (79.79%) (**Figure 2B**). On the 49,510 *P. phosphorea*
unigenes present in the filtered reference transcriptome, 34,984 showed significant matches to
molecular databases: 33,896 to NR (68.5%, E-value $> 1e^{-5}$), 8,818 to NT (17.8%), 26,473 to
SwissProt (53.5%), 28,290 to KEGG (57.1%), 24,642 to KOG (49.8%), 26,790 to InterProt
(54.1%), and 13,742 to GO (27.8%) (**Figure 2C**).
For *A. murrayi*, in total, 126,709 unigenes were obtained for a total length of 9,655,825
pb. The average mean length was 635, and the N50 was 682. The length distributions of
unigenes are shown in **Figure S2**, and numerical data are summarized in **Tables S4 and S5**.
For descriptive purposes, a gene expression analysis was performed by mapping FPKM values
(*e.g.,* $\log_{10}$(FPKM value) calculated for all predicted unigenes (**Table S6**). The main
represented species within the unigene annotation of the reference transcriptome is the
anthozoan *Nematostella vectensis* (44.36%) (**Figure S2**). On the 126,709 *A. murrayi* unigenes
present in the filtered reference transcriptome, 35,923 showed significant matches to molecular
databases: 6,993 to NR (62.7%, E-value $> 1e^{-5}$), 2,007 to NT (18.2%), 5,251 to SwissProt
(47.1%), 4,549 to KEGG (40.8%), 5,161 to KOG (46.3%), 5,162 to InterProt (48.8%), and
3,057 to GO (27.4%) (**Figure S2**).
Because of the lack of reference genome in *Pennatula* and *Anthoptilum* and, possibly,
the relatively short length of some unigene sequences, 29.4% and 71,6%, respectively, of the
assembled sequences could not be matched to any known genes.
Based on Benchmarking Universal Single-Copy Orthologs (BUSCO) analysis, 96.23%
metazoa BUSCO genes are predicted to be complete in the merged *P. phosphorea*
transcriptome. In parallel, 1.36% of BUSCO genes are fragmented and 2.41% are missing
(**Figure 2D**). Similar results were obtained for *A. murrayi*, with 94.44% complete BUSCO
genes found while 3.04% BUSCO genes are fragmented and 2.52% are missing (**Figure 2D**).
494

Figure 2. General description of the transcriptomic data. (A) Length distribution of *P. phosphorea* unigenes. (B) Taxonomic annotation of *P. phosphorea* transcriptome. (C) Global annotation of *P. phosphorea* transcriptome. (D) Busco analysis.

Figure 3. Global gene expression and gene ontology of the *Pennatula phosphorea* transcriptomic data. (A) Gene expression in each tissue sample of *P. phosphorea*. (B) gene expression distribution in each sample of *P. phosphorea*. (C) Proportions of unigenes with relative high/low expression in *P. phosphorea* samples. (D) Gene ontology distribution of the 40 most expressed unigenes for each sample of *P. phosphorea*.

Analyses of the unigene expression reveal a similar proportion of FPKM values among the three tissues (**Figure 3A, B**). Comparatively, a large number of unigenes appear to be more expressed in the pinnule and the rachis in comparison to the peduncle tissue (**Figure 3C**). The pinnule and rachis tissues appear more similar in term of unigene expression profile. The "Molecular function" GO functional annotations for the 40 unigenes with the most pronounced expression differences between the tissues, including both those highly expressed in one tissue

511 relative to the other and those with lower expression levels are visualized in **Figure 3D**. We did
512 not perform a formal differential gene expression as no replication was performed. Calcium ion
513 binding function (GO: 0005509) appears predominantly expressed in pinnule and rachis tissues,
514 compared to the peduncle (**Figure 3D**). Similarly, pinnule tissue (containing the feeding polyp)
515 present a higher proportion of gene annotated as digestive enzymes (**Figure 3D**).
516
517 ***Expression of bioluminescence-related genes in Pennatula phosphorea and Anthoptilum***
518 ***murrayi***
519
The *P. phosphorea* and *A. murrayi* transcriptomes contained sequences for several
predicted luciferases, GFPs, and luciferin-binding proteins. The FPKM values retrieved in each
transcriptome are given in **Table S7**. Reciprocal BLAST analyses revealed that the sequences
matched for luciferases, CBPs, and GFPs of Anthozoans.
Several *R*Luc-like enzymes were recovered from both investigated species (*A. murrayi*
and *P. phosphorea*). In addition, the sequence mining allowed us to recover additional *R*Luc-
like from the genomes of *Renilla muelleri* and *R. reniformis*. The phylogenetic analyses
revealed three clades of *R*Luc-like enzymes in Pennatulacea (**Figure 4A**). Clade A contains the
well-known luciferases from *R. muelleri* and *R. reniformis* and probable luciferase sequences
from *A. murrayi*, *P. phosphorea,* and *Cavernularia obesa*. Interestingly, clade A does not
contain any sequence from non-luminous species. The clade B, in comparison, includes several
sequences from non-luminous species (*e.g., Pinnigorgia flava*, *Sinularia cruciata*) in addition
to several sequences from both our model species. The clade C also contains *R*Luc-like enzymes
retrieved from each transcriptome.

Figure 4. Phylogenetic tree of *R*Luc-like enzymes including the *P. phosphorea* and *A. murrayi* amino-acid
sequences. (A) Maximum likelihood tree based on an amino-acid sequence alignment of *R*Luc-like enzymes. The
tree is calculated by IQ-tree software using the LG+I+G4 model of evolution. Numbers at nodes indicate ultrafast
bootstrap values based on 1000 replicates. The scale bar represents the percentage of amino-acid substitutions per
site. Bacterial haloalkane dehalogenase sequences were used to root the tree. (B) Expression level of each retrieved
*R*Luc-like sequence for the distinct portion of *P. phosphorea*. (C) Expression level of each retrieved *R*Luc-like
sequence in *A. murrayi* whole specimen. (D) Superimpose spectrum of the *in vivo* luminescence of *A. murrayi*

(green) and of the *in vitro* luminescent assay of *A. murrayi* luciferase in the presence of synthesized coelenterazine
(fc.: 6 µM) (blue).

The retrieved sequence (CL1333; **Figure 4B**) of *P. phosphorea* luciferase has an
estimated molecular weight of 35.84 kDa. The comparison of the amino-acid sequences of *P.*
phosphorea and anthozoan luciferases demonstrated that the catalytic triad involved in the
luciferase activity was present. These key sites consist of an aspartate residue in position 120,
a glutamate residue in position 144, and a histidine in position 285. The retrieved *P. phosphorea*
luciferase sequence based on RNA-seq data appears highly similar to other known anthozoan
luciferase. It shares 90.26% of identity and 95% of similarity with *Renilla reniformis* luciferase
(*Renilla*-luciferin 2-monooxygenase). This sequence was validated through DNA amplification
and sequencing using *R*Luc primers. FPKM analyses reveal that this sequence is mostly
expressed within pinnule and rachis tissues (**Figure 4B**).

A. murrayi luciferase (Comp24811, **Figure 4C**), with a molecular weight estimated at
34.61 kDa and consisting of 304 amino acids, exhibits 58% sequence identity and 98%
coverage compared to *R*Luc, highlighting significant similarities. Sequence alignment of *A.*
murrayi luciferase with *R*Luc has revealed the conservation of the catalytic triad and active site.
The *R*Luc structure features two distinct domains: a cap domain and an alpha/beta-hydrolase
domain. In the cap domain, key residues essential for enzymatic activity, including the substrate
entry tunnel and catalytic triad (D120, E144, and H285), are located. These elements have also
been identified in *A. murrayi's* luciferase sequence, as reported by Rahnama et al., 2017 [58]
and Khoshnevisan et al. 2018 [59]. Through the expression of recombinant luciferase in
Escherichia coli using degenerate primers derived from *A. murrayi* transcriptome analysis, *A.*
murrayi luciferase was tested for preliminary downstream expression, yielding an active
enzyme capable of producing blue light (λmax = 498 nm) upon coelenterazine addition (**Figure**
4D).

567
568 **Figure 5. Phylogenetic tree of anthozoan fluorescent proteins including the *P. phosphorea* and *A. murrayi***

 (A) Maximum likelihood tree based on an amino-acid sequence alignment of green fluorescent proteins. The tree is calculated by IQ-tree software using the WAG+R4 model of evolution. Numbers at nodes indicate ultrafast bootstrap percentages based on 1000 replicates. The scale bar represents the percentage of amino-acid substitutions per site. Hydrozoan GFP sequences were used to root the tree. (B) Expression level of each retrieved CBP sequence for the distinct portion of *P. phosphorea*. (C) Expression level of each retrieved CBP sequence in *A. murrayi* whole specimen.

Several sequences coding for green fluorescent proteins were retrieved from *P. phosphorea* and *A. murrayi* transcriptomes. *P. phosphorea* GFP (CL380; 20.81 kDa) and *A. murrayi* GFP (Comp24361; 27.24 kDa) appear homologous to other anthozoan sequences, and both sequences clustered with GFP sequences of bioluminescent Scleralcyonacea (**Figure 5A**). The *P. phosphorea* GFP shared 79.42% identity and 86% similarity with the GFP of *R. reniformis*. In comparison, the *A. murrayi* GFP shared 73.06% of identity and 87% similarity with the GFP of *Cavernularia obesa*. Pinnules and rachis were the main sites expressing the *P. phosphorea* GFP (**Figure 5B**). Among the unigene pairs found in the *A. murrayi* transcriptome, Comp24361 appeared to be more expressed (**Figure 5C**).

Figure 6. Phylogenetic tree of coelenterazine-biding proteins including the *P. phosphorea* and *A. murrayi* amino-acid sequences. (A) Maximum likelihood tree based on an amino-acid sequence alignment of coelenterazine-binding proteins. The tree is calculated by IQ-tree software using the LG+R4 model of evolution. Numbers at nodes indicate ultrafast bootstrap percentages based on 1000 replicates. The scale bar represents the percentage of amino-acid substitutions per site. Calmodulin sequences were used to root the tree. (B) Expression level of each retrieved CBP sequence for the distinct portion of *P. phosphorea*. (C) Expression level of each retrieved CBP sequence in *A. murrayi* whole specimen.

As for the luciferase and the GFP, different sequences of CBPs were retrieved from *P.*
phosphorea and *A. murrayi* transcriptomes. Some of these sequences appeared clustered with
the CBPs of luminous Scleralcyonacea, while others were found to be clustered with
photoprotein-like proteins (**Figure 6A**). One sequence appeared to be mainly expressed within
the *P. phosphorea* pinnule and rachis (CL1544; 26.84 kDa) and *A. murrayi* colony (Comp
22336; 21.08 kDa) (**Figure 6B, C**). The most expressed *P. phosphorea* CBP appeared highly
homologous to other luminous anthozoan sequences. *P. phosphorea* CBP shared 85.33%
identity and 94% similarity with the luciferin-binding protein of *R. reniformis*, while *A. murrayi*
CBP shared 50% identity and 74% similarity with the luciferin-binding of *R. reniformis*.
Interestingly, sequences that cluster in the photoprotein-like proteins group appeared to be
mainly expressed within the non-photogenic peduncle tissue in *P. phosphorea*.

Luciferase expression and green autofluorescence in Pennatula phosphorea and Funiculina
quadrangularis

Based on the high sequence similarity of the *P. phosphorea* luciferase with *Renilla*
luciferase, a commercial antibody anti-*Renilla* luciferase was selected for immunodetections.
Immunoblot analyses revealed strong anti-luciferase immunoreactive bands in both extracts of
pinnule and rachis tissues (**Figure S2**). The bands correspond to a protein with an approximate
molecular weight of 35 kDa, matching the molecular weight of the predicted *P. phosphorea*
luciferase and the *Renilla* luciferase molecular weight of 36 kDa. No labeling could be detected
for the peduncle tissue extract (**Figure S3**).
On autozooid polyps of the pinnules (**Figure 7A**), a strong green autofluorescent signal
is observed at the tentacle crown base before (**Figure 7B**) and after (**Figure 7C**)
paraformaldehyde fixation. This green signal is localized at the tentacle junctions as multiple
cells cluster. A strong anti-*Renilla* luciferase immunoreactivity is observed at the same level as
the green fluorescence signal (**Figure 7C, D**). On siphonozooid polyps of the rachis, the
autofluorescent signal is also observable and is located as green dots (from 10 to 25 μm
diameter) spread in the tissue, generally in pairs (**Figure 7E**). Anti-*Renilla* luciferase-positive
cells colocalized with these autofluorescent dots (**Figure 7E, F**). Finally, no green fluorescence
or immunolabeling is detected on the peduncle tissue (data not shown).
For *Funiculina* polyps (**Figure 7G**), a green autofluorescent signal was observed on
freshly dissected specimens before fixation (**Figure 7H**). This green autofluorescence
completely disappeared after fixation (**Figure 7I**). Luciferase localization was different from
P. phosphorea. The luciferase signal was strongly detected within the polyp tentacle tissue and
not at the base of the polyp crown (**Figure 7I, J**).
Negative controls with omission of the primary antibody did not reveal any non-specific
binding of the secondary antibodies (data not shown).

632
Figure 7. Autofluorescence and luciferase immunodetection in *P. phosphorea* and *F. quadrangularis.* (A)
Schematic illustration of *P. phosphorea*. Natural green autofluorescence (B), green fluorescence after fixation
(C), and luciferase immunodetection (D; magenta) of the *P. phosphorea* pinnule autozooids. Green fluorescence
after fixation (E) and luciferase immunodetection (F; magenta) of the *P. phosphorea* rachis siphonozooids. (G)
schematic illustration of *F. quadrangularis*. Natural green autofluorescence (H), observation after fixation with no
autofluorescent signal (I), and luciferase immunodetection (J; magenta) of *F. quadrangularis* autozooids. Scales:
B-F, H - 500 μm; I, J - 250 μm.

Calcium is involved in the bioluminescence of Pennatula phosphorea and Funiculina
quadrangularis

Based on *P. phosphorea* coelenterazine-binding protein retrieval in the *Pennatula*
transcriptome data and the literature mentioning the potential implication of Ca^{2+} in the
releasing of the luciferin from coelenterazine-binding protein, Ca^{2+} involvement in the light
emission process of *P. phosphorea* and *F. quadrangularis* was investigated. Tested specimens
of both species revealed a drastic increase in light production when immersed in ASW with
doubled Ca^{2+} ion concentration (20 mM) (**Figure 8A, B**). At the same time, no statistical
differences are observed between normal ASW (10 mM) and ASW devoted to Ca^{2+} ions (0
mM) (**Figure 8A, B**). Analysis of the A23187 ionophore effects shows no statistical differences

comparatively to each calcium concentration (**Figure 8C, D**). For both specimens, adrenaline triggers light production when calcium is present in ASW (**Figure 8E, F**). While an increase of the mean Ltot is observable between adrenaline-tested samples at 10 mM and at 20 mM of $CaCl_2$ in the medium, this increase is not statistically supported (**Figure 8E, F**). Finally, no significant differences are observed in Ltot after KCl applications at the three Ca^{2+} ion concentrations (**Figure S4**).

Figure 8. Calcium involvement in *P. phosphorea* and *F. quadrangularis* light emission. Experiments performed on *P. phosphorea* (A, C, E) and *F; quadrangularis* (B, D, F). Effect of different concentrations of calcium (0, 10, 20 mM) in the medium on the total light emission (Ltot) (A, B). Effect of different calcium concentrations (0, 10, 20 mM) in the medium in the presence and absence of the calcium ionophore A23187 on the Ltot (C, D). Effect of different concentrations of calcium (0, 10, 20 mM) in the medium in the presence of adrenaline (10^{-5} mol L^{-1}) on the Ltot (E, F). different lettering indicates statistical differences.

Discussion

Through biochemical cross-reaction experiments, this study showed that the bioluminescence of *Pennatula phosphorea*, *Funiculina quadrangularis*, and *Anthoptilum murrayi* is associated with a coelenterazine-dependent luciferase homologous to the luciferase of the sea pansy *Renilla* species. The measured concentration of coelenterazine in *P. phosphorea* pinnules closely aligns with the reported values of 141 and 282.4 ng g^{-1} for the phylogenetically closed species *Renilla muelleri* and *Cavernularia obesa*, respectively [1]. Similarly, the mean luciferase activity of *P. phosphorea* pinnules reaches in the same range as other organisms using *Renilla*-type luciferases such as *R. muelleri* ($\pm$ 130 10^9 q g^{-1} s^{-1}), *C. obesa* ($\pm$ 220 10^9 q g^{-1} s^{-1}), or *Amphiura filiformis* ($\pm$ 69 10^9 q g^{-1} s^{-1}) [1,60]. While the luciferase activity recorded for *F. quadrangularis* is higher than *P. phosphorea*, this species presents smaller coelenterazine content, being closer to recorded values of the sympatric ophiuroid species *A. filiformis* (5.4 ng g^{-1}; [60]) or another echinoderm, *Thalassometra gracilis* (4.5 ng g^{-1}; [61]). The transcriptomic input confirms results for *P. phosphorea* and *A. murrayi* with a high sequence homology of the retrieved luciferase with the *Renilla* luciferase. Consistent with the literature, species such as *Distichoptilum gracile* or undetermined species from *Umbellula*, *Pennatula*, or *Funiculina* genus also present evidence of the use of coelenterazine, an imidazopyrazinone compound as substrate and a *Renilla*-like luciferase as enzyme of their bioluminescent systems [12].

Our phylogenetic analysis underlines three distinctive groups of *Renilla*-like luciferase. One of these groups (Clade A) contains all the known sequences of anthozoan light-productive luciferase. The other two groups (Clades B and C) contain other homologous *R*luc-like sequences, notably from non-luminous species. The related biochemical functionality and bioactivity of these B and C clades sequences are unknown and need further investigations to fully apprehend the evolution of luciferase among anthozoans. Nevertheless, this clustering led to hypothesized duplications of the ancestral gene with either (*i*) a neo functionality as a "real" functional luciferase able to catalyze a bioluminescent reaction or (*ii*) a bi-functionality of these enzymes as light-producing enzyme and another ancestrally conserved enzymatic function. The ancestral functionality of the *R*Luc enzyme has been assumed to originate from bacterial dehaloalkane dehalogenase, an enzyme with a hydrolase activity cleaving bonds in halogenated compounds [62]. A basal gene transfer from bacteria until metazoan has been hypothesized [17,62]. Interestingly, *R*Luc-like luciferases have been demonstrated to be the enzymes involved in the bioluminescence of phylogenetically distinct species such as the brittle star, *A. filiformis* [17] and the tunicate, *P. atlanticum* [18], letting assumed that the sequence was convergently coopted multiple times during the evolution. Consistent with previous research on *R*Luc sequences, our retrieved *P. phosphorea* and *A. murrayi* luciferase (Clade A) present high sequence homologies with the other pennatulacean functional luciferases, also revealing conservation of the catalytic triads essential for the luciferase catalytic activity. The complete characterization of *P. phosphorea* and *A. murrayi* luciferase will allow to better apprehend the functionality and evolution of these luciferase enzymes.

While the retrieved GFP sequences are unique and well clustered with other Scleralcyonacea sequences, a similar observation, as for the luciferase, occurs for the CBPs

with multiple retrieved sequences clustered in different groups. The first group represents the
functional coelenterazine-binding protein retrieved and characterized in luminous species,
while the second group, named photoprotein-like proteins, raised questions on the exact
functionality of these retrieved sequences. These photoprotein-like proteins might also be
involved in calcium binding. Active photoproteins of cnidarians and ctenophores depend on
calcium to trigger light emission [63-66]. Nevertheless, the higher expression within the *P.*
phosphorea peduncle let assumed another function without a relationship with the
bioluminescence.
Kept in captivity without exogenous coelenterazine supply, this sea pen species can still
produce light after one year, even if the coelenterazine content and luciferase activity decrease.
Therefore, these results support a *de novo* synthesis of the coelenterazine substrate in *P.*
phosphorea. Coelenterazine *de novo* synthesis by luminous marine organisms has been
documented for the calanoid copepods *Metridia longa* and *M. pacifica*, the oplophorid shrimp,
Systellaspis debilis, and two ctenophore species, *Mnemiopsis leidyi* and *Bolinopsis*
infundibulum [67-70]. The natural precursors of coelenterazine have been demonstrated to be
the L-tyrosine and L-phenylalanine amino acids in *M. pacifica* [69]. By screening the
transcriptomic data of 24 ctenophores, Francis et al., 2015, assumed the involvement of a non-
heme iron oxidase-like enzyme, similar to isopeniciline-N-synthase, in the biosynthesis
pathway of this luciferin [71]. Future research could be carried out to validate the *de novo*
biosynthesis of the coelenterazine bioluminescent substrate, in particular by maintaining *P.*
phosphorea and their offspring over generations in captivity in the same conditions without
coelenterazine supply [70,72]. This protocol could be applied to other pennatulaceans to
determine whether the coelenterazine genesis is a common trait in this clade. Nevertheless, the
first challenge would be able to maintain and master the life cycle of these species in captivity.
Similarly, it would be interesting to analyze different pennatulacean's transcriptomes searching
for enzymes homologous to the isopenicillin-N-synthase potentially involved in the
coelenterazine biosynthetic pathway, as retrieved in ctenophores.
In *P. phosphorea*, the morphological localization of the luciferase expression matches
to the green fluorescent sites obtained in unfixed and fixed tissues. Green autofluorescence
observed on the unfixed specimen is assumed to be a mix of the native autofluorescence of the
coelenterazine and the autofluorescent reaction occurring through the GFP, while green
autofluorescence observed after tissue fixation corresponds only to the GFP signals.
Comparatively and accordingly, the *in vivo* green fluorescence observed in the *F.*
quadrangularis tissues, disappearing after fixation, is assumed to be only related to the natural
autofluorescence of the coelenterazine and the lack of GFP in this species. Indeed, conversely
to *P. phosphorea,* which emits green waves of light at λmax = 510 nm, *F. quadrangularis* emits
in blue at λmax = 485 nm, supporting the absence of GFP for this species [34,73,74]. This
natural autofluorescence has been recently demonstrated to appear and disappear from the
photogenic site depending on the substrate dietary acquisition of the brittle star *A. filiformis*.
This species depended on a trophic acquisition of the coelenterazine substrate to produce light
[60,75]. When the brittle star is fed with coelenterazine-containing food, green autofluorescent
spots appear at the level of the spine-associated photocytes [75]. As for *F. quadrangularis*, a
similar disparition of the coelenterazine-like green fluorescent signal has been observed on
fixed tissue of the brittle star species [75]. The autofluorescent sites observed along the tentacle

bases of the autozooids of *P. phosphorea* are consistent with the already described location of
the autofluorescent photogenic cell processes in autozooids of *Stylatula elongata*,
Acanthoptilum gracile, and *Renilla koellikeri* made by Satterlie et al., 1980 [31]. For the former
species, it was noticed that the photocytes process followed the same orientation as the
longitudinal muscles, allowing autozooids to retract [31]. On the other hand, the luciferase
expression site in *Funiculina*, in the cellular processes of the apical part of the tentacle, was
never reported before.

Our spectrum measurement performed with the *A. murrayi* recombinant luciferase is
consistent with already observed coelenterazine-dependent systems in other species emitting in
the same range of wavelengths. Recorded spectrum for coelenterazine-dependent luciferase
systems can vary between 475 to 493 nm in *Pyrosoma atlanticum*, 482 nm emitted by *R*Luc,
472 nm emitted by the brittle star *Amphiura filiformis*, 455 nm emitted by *Oplophorus*
gracilirostris and 485 nm emitted by *F. quadrangularis* [17,18,62,76]. According to the natural
spectrum recorded on the whole specimens (λmax = 513 nm), similar to the spectrum measured
for *P. phosphorea* [73], and the transcriptomic presence of a GFP sequence in *A. murrayi*, this
species is strongly assumed to display a GFP-associated coelenterazine-dependent luminous
system. At least six anthozoan species present expression of a GFP and the production of green
light. Some of these species may live in sympatry with anthozoan blue emitters, as *P.*
phosphorea and *F. quadrangularis,* with an overlap of depth repartition and habitat preferences
occurring. Therefore, questions arise concerning luminescence's exact function(s) among
anthozoans. Even if some assumptions are proposed in the literature, no one has ever developed
an ethological protocol to validate them [35]. Therefore, why some anthozoan evolved green
light emission while the primary coelenterazine-luciferase reaction produces blue light remains.
Different function(s) in the same environment may have led to the acquisition or the loss of the
GFP gene by some anthozoan species upon evolutionary constraints.

As demonstrated for other pennatulacean species, CBP seems to be an essential
component of the luminous system [19-23,25]. The retrieved CBPs in *P. phosphorea* and *A.*
murrayi are congruent with this literature. Gene ontology distribution analyses performed on
the different tissues of *P. phosphorea* underline a high expression of calcium ion binding
proteins, including the retrieved CL1333 CBP, in the photogenic tissue (pinnule and rachis) of
this species. The expression of this gene within the photogenic tissues supports it involvement
in the luminous reaction. As demonstrated for *Renilla* by Stepanyuk et al., 2008, CBPs need
calcium as a cofactor to release the coelenterazine [27]. In addition to the retrieved CBPs
sequence in the *Pennatula* transcriptome, our calcium assay results highlight the involvement
of the calcium ion in the light emission process of both *P. phosphorea* and *F. quadrangularis*.
Results obtained for the calcium ionophore A23187 reveal that the ion action does not result
from intracellular calcium storage but rather is provided by external calcium input. Moreover,
calcium is shown to be essential for the physiological luminescent response through adrenaline
application. Pieces of evidence of calcium involvement in anthozoan luminescence were
already described for *R. reniformis* and *V. cynomorium* [25,26].
These results let us assume a conservation of the coelenterazine-dependent *Renilla*-like
luciferase bioluminescent system, involving also a CBP, across luminous Pennatulaceans
(**Table S1**). The involvement of GFP appears species-dependent, with only a few species

emitting blue light (**Table S1**). Nevertheless, deeper investigations are needed to fully
apprehend the conservation of those actors across the diversity of luminous pennatulaceans.

799
800   **Figure 9: Schematic representation of the putative pathway driving the luminescence production in sea**
801   **pens.** Elements of this pathway have been compiled from the present results and the literature [12,13,16,19-23,30-
802   34]. CBP, coelenterazine-binding protein; CTM, coelenteramide; CTZ, coelenterazine; G, G-protein; GFP, green
803   fluorescent protein.
804

A hypothetical scheme of the generalized pathway could be established using our results
and the literature on the pennatulacean luminescence mechanism (**Figure 9**). The first step is
the activation of catecholaminergic receptors through the binding of biogenic amines (mainly
adrenaline and noradrenaline [33,34]), which will release an intracellular-associated G-protein
[77,78]. G-protein could be involved in a large variety of intracellular pathways [79], some of
which involve increasing intracellular calcium (via direct or indirect activation of calcium
channels) [80-83]. This intracellular calcium increase will lead to the release of the
coelenterazine through the binding of this ion on the CBP, leaving this luciferin free to react
with luciferase in the presence of oxygen to produce blue light around 480 nm [9,10,15,19-
23,25]. For pennatulaceans lacking GFP (*e.g., F. quadrangularis*), this scheme ends here with
the direct emission in blue color, while for those displaying GFP expression, the blue light is
captured by this specific fluorescent protein and reemitted in green wavelength [20], such as
for *P. phosphorea*. Future research is needed to validate this hypothetical scheme, and further
investigations will be conducted to establish the functional activities of all these components in
less-studied sea pens. Astonishingly, even if attempts were made on other anthozoans over the
past decades, the *Renilla* bioluminescence system remains the only isolated and cloned system
[13,15,20-23,25]. Despite the demonstrated widespread uses of the sea pansy bioluminescent
system in biotechnology and biomedicine [*e.g.,* 84-86], the *Renilla* luciferase stands as the only
one of the most commercially employed gene reporters in biomolecular sciences. Nonetheless,
in the *Renilla* bioluminescence system, the exact action mode of CBP and calcium is not fully

apprehended. Considering these facts, our introspection into the bioluminescent system of other
luminous pennatulaceans could be of great use for new biotechnological advances. Our results
provided a better understanding of the evolution of the bioluminescence system and associated
molecules from these enigmatic benthic sessile organisms.
829
830 **Supplemental data captions**
831
832 **Table S1. General overview of the bioluminescent knowledge among luminous anthozoans.** All
833 data were extracted from the literature (only scientific data described for species down to the species
834 level have been taken into account to minimize generalizations). CBP, coelenterazine-binding protein;
835 GFP, green fluorescent protein.
836
837 **Table S2. Primers designed for _P. phosphorea_ luciferase validation.**
838
839 **Table S3. Experiment protocol for the calcium assays.**
840
841 **Table S4. Description of the output sequenced data.** Q20 percentage is the proportion of nucleotides
842 with a quality value larger than 20 in reads. GC percentage is the proportion of guanidine and cytosine
843 nucleotides among total nucleotides.
844
845 **Table S5. Summary statistics of assemblies for _Pennatula phosphorea_ pinnules, rachis, and**
846 **peduncle transcriptomes.**
847
848 **Table S6. Annotation file for both transcriptomes.**
849
850 **Table S7. Transcript expression values (FPKM values) and public database sequences used during**
851 **transcriptomic and phylogenetic analyses.**
852
Figure S1: Biochemical assays following luminous parameters activity on _Pennatula phosphorea_
pinnules (A, B, C) and rachis (D, E, F) during 1 year without coelenterazine supply. Mean total
light emission after KCl applications (A, D), mean coelenterazine content (B, E), and mean maximal
light emission during luciferase activity experiments (C, D). Different lettering indicates statistical
differences. The timing corresponds to experiments performed on wild-caught specimens (T0) after 6
months (T6) and twelve months of captivity (T12).
859
860 **Figure S2. General description of the _Anthoptilum murrayi_ transcriptomic data.** (A) Length distribution of
861 _A. murrayi_ unigenes. (B) Taxonomic annotation of _A. murrayi_ transcriptome. (C) Global annotation of _A. murrayi_
862 transcriptome.
863
864 **Figure S3. Luciferase immunoblots performed on _Pennatula phosphorea_ tissues (rachis,**
865 **peduncle, and pinnule).**
866
867 **Figure S4. Ltot after KCl applications with three different calcium concentrations in the medium.**
868

869 **Acknowledgment**
870 The authors acknowledge U. Schwarz, captain of the Alice vessel, and the skillful members of
871 the Kristineberg Center (Goteborg University, Sweden) for their help during the _Pennatula_ and

Funiculina collection. The authors also thank M. Jacquet for contributing to the study during his master's thesis and C. Pels, ELIV laboratory technician who maintained the organisms in the aquaria at the Marine Biology Laboratory - UCLouvain. The authors also want to thank T. Wiegand from the mobile lab (TREC- EMBL) for her help in visualizing coelenterazine autofluorescence and immunolabeling in the wild-caught specimens. The authors also thank J. Mallefet for his helpful advice and help during the first organisms sampling and all along the experiments. LD is a postdoctoral researcher at the Université de Louvain - UCLouvain, GG is a Ph.D. student at Universidade de São Paulo - USP, CC is a Ph.D. student under an FRIA fellowship, LB is a postdoctoral researcher at the Université de Louvain - UCLouvain, RR is an academic professor at UCLouvain, MRSM is an academic professor at Universidade de São Paulo - USP, MM is group leader at the Masaryk University, SD is a Senior Lecturer and Associate professor at the University of Gothenburg, AGO is an assistant professor at Yeshiva University, and JD is a postdoctoral researcher at FNRS. This study is the contribution of BRC#422 of the Biodiversity Research Center (UCLouvain) from the Earth and Life Institute Biodiversity (ELIV) and the "Centre Interuniversitaire de Biologie Marine" (CIBIM).

Competing interests

No competing interests declared

Data availability

The *P. phosphorea* and *A. murrayi* transcriptome raw reads are available upon request to the authors in the meantime. Some supplementary data are not present in this preprint manuscript (Supplementary Table S6 & 7). These datasets are temporarily only accessible from the corresponding author upon reasonable request.

Funding

This work was supported by an F.R.S.-FNRS grant (T.0169.20) awarded to the Université de Louvain – UCLouvain Marine Biology Laboratory and the Université de Mons Biology of Marine Organisms and Biomimetics Laboratory. The research leading to these results also received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 730984, ASSEMBLE Plus project. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 88887.605088/2021-00.

Author contributions

LD, CC, and SD collected the P. phophorea and F. quadrangularis samples, and GG and MRM sampled *A. murrayi*. Data collection was performed by LD, CC, and LB for *P. phosphorea* and *F. quadrangularis*, while data for *A. murrayi* was collected by GG and AO. Transcriptomic analysis was performed by LD, GG, AO, and JD. Phyloproteomic was mainly performed by LD, MM, and JD. Work was supervised by RR, SD, AO, and JD. The original manuscript was written by LD, CC, GG, AO, and JD. All the authors modified and corrected the final version.

Reference

1. Shimomura O, Yampolsky IV. 2019 *Bioluminescence: chemical principles and*
methods. 3rd edition (Singapore: World Scientific Publishing), 1–556.

2. Haddock SHD, Moline MA, Case JF. 2010 Bioluminescence in the sea. *Ann. Rev. Mar.*
Sci. **2**, 443-493. (https://doi.org/10.1146/annurev-marine-120308-081028)

3. Delroisse J, Duchatelet L, Flammang P, Mallefet J. 2021 Leaving the dark side? Insights
into the evolution of luciferases. *Front. Mar. Sci.* **8**, 673620.
(https://doi.org/10.3389/fmars.2021.673620)

4. Fleiss A, Sarkisyan KS. 2019 A brief review of bioluminescent systems. *Curr. Genet.*
65, 877-882. (https://doi.org/10.1007/s00294-019-00951-5)

5. Hastings JW. 1996 Chemistries and colors of bioluminescent reactions: a review. *Gene*
173(1), 5-11. (https://doi.org/10.1016/0378-1119(95)00676-1)

6. Wilson T, Hastings JW. 2012 *Bioluminescence: living lights, lights for living*. Harvard
University Press.

7. Kaskova ZM, Tsarkova AS, Yampolsky IV. 2016 1001 lights: luciferins, luciferases,
their mechanisms of action and applications in chemical analysis, biology and medicine.
Chem. Soc. Rev. **45**, 6048-6077. (https://doi.org/10.1039/C6CS00296J)

8. Cormier MJ. 1960 Studies of the bioluminescence of *Renilla reniformis*: I.
Requirements for luminescence in extracts and characteristics of the system. *Biochimica*
et Biophysica Acta **42**, 333-343. (https://doi.org/10.1016/0006-3002(60)90797-6)

9. Matthews JC, Hori K, Cormier MJ. 1977 Purification and properties of *Renilla*
reniformis luciferase. *Biochemistry* **16(1)**, 85-91.
(https://doi.org/10.1021/bi00620a014)

10. Lorenz WW, McCann RO, Longiaru M, Cormier MJ. 1991 Isolation and expression of
a cDNA encoding *Renilla reniformis* luciferase. *PNAS* **88(10)**, 4438-4442.
(https://doi.org/10.1073/pnas.88.10.4438)

11. Loening AM, Fenn TD, Gambhir SS. 2007 Crystal structures of the luciferase and green
fluorescent protein from *Renilla reniformis*. *J. Mol. Biol.* **374(4)**, 1017-1028.
(https://doi.org/10.1016/j.jmb.2007.09.078)

12. Bessho-Uehara M, Francis WR, Haddock SHD. 2020 Biochemical characterization of
diverse deep-sea anthozoan bioluminescence systems. *Mar. Biol.* **167**, 114.
(https://doi.org/10.1007/s00227-020-03706-w)

13. Schenkmayerova A, Toul M, Pluskal D, Baatallah R, Gagnot G, Pinto GP, Santana VT,
et al. 2023 Catalytic mechanism for *Renilla*-type luciferases. *Nat. Catal.* **6**, 23-38.
(https://doi.org/10.1038/s41929-022-00895-z)

14. DeLeo D, Bessho-Uehara M, Haddock SHD, McFadden CS, Quattrini AM. 2024
Evolution of bioluminescence in Anthozoa with emphasis on Octocorallia. *Proc. R. Soc.*
B. **291**, 20232626. (https://doi.org/10.1098/rspb.2023.2626)

15. Inoue S, Kakoi H, Murata M, Goto T, Shimomura O. 1977 Complete structure of *Renilla*
luciferin and luciferyl sulfate. *Tetrahedron Lett.* **18(31)**, 2685-2688.
(https://doi.org/10.1016/S0040-4039(01)83046-X)

16. Cormier MJ, Hori K, Karkhanis YD, Anderson JM, Wampler JE, Morin JG, Hastings
JW. 1973 Evidence for similar biochemical requirements for bioluminescence among
coelenterates. *J. Cell. Physiol.* **81(2)**, 291-297.
(https://doi.org/10.1002/jcp.1040810218)

17. Delroisse J, Ullrich-Lüter E, Blaue S, Ortega-Martinez O, Eeckhaut I, Flammang P, Mallefet J. 2017 A puzzling homology: a brittle star using a putative cnidarian-type luciferase for bioluminescence. *Open Biol.* **7(4)**, 160300. (https://doi.org/10.1098/rsob.160300)

18. Tessler M, Gaffney JP, Oliveira AG, Guarnaccia A, Dobi KC, Gujarati NA, et al. 2020 A putative chordate luciferase from a cosmopolitan tunicate indicates convergent bioluminescence evolution across phyla. *Sci. Rep.* **10**, 17724. (https://doi.org/10.1038/s41598-020-54373446-w)

19. Anderson J, Charbonneau H, Cormier MJ. 1974 Mechanism of calcium induction of *Renilla* bioluminescence. Involvement of a calcium-triggered luciferin binding protein. *Biochemistry* **13(6)**, 1195-1200. (https://doi.org/10.1021/bi00703a602)

20. Ward WW, Cormier MJ. 1979 Energy transfer protein in coelenterate bioluminescence. Characterization of the *Renilla* green-fluorescent protein. *J. Biol. Chem.* **254(3)**, 781-788.

21. Kumar S, Harrylock M, Walsh KA, Cormier MJ, Charbonneau H. 1990 Amino acid sequence of the Ca^{2+}-triggered luciferin binding protein of *Renilla reniformis*. *FEBS Lett.* **268(1)**, 287-290. (https://doi.org/10.1016/0014-5793(90)81029-N)

22. Inouye S. 2007 Expression, purification and characterization of calcium-triggered luciferin-binding protein of *Renilla reniformis*. *Protein Expr. Purif.* **52(1)**, 66-73. (https://doi.org/10.1016/j.pep.2006.07.028)

23. Titushin MS, Markova SV, Frank LA, Malikova NP, Stepanyuk GA, Lee J, Vysotski ES. 2008 Coelenterazine-binding protein of *Renilla muelleri*: cDNA cloning, overexpression, and characterization as a substrate of luciferase. *Photochem. Photobiol. Sci.* **7**, 189-196. (https://doi.org/10.1039/b713109g)

24. Ogoh K, Kinebuchi T, Murai M, Takahashi T, Ohmiya Y, Suzuki H. 2013 Dual-color-emitting green fluorescent protein from the sea cactus *Cavernularia obesa* and its use as a pH indicator for fluorescent microscopy. *Luminescence* **28(4)**, 582-591. (https://doi.org/10.1002/bio.2497)

25. Henry J-P, Ninio M. 1978 Control of the Ca^{2+}-triggered bioluminescence of *Veretillum cynomorium* lumisomes. *Biochimica et Biophysica acta* **504(1)**, 40-59. (https://doi.org/10.1016/0005-2728(78)90005-1)

26. Charbonneau H, Cormier MJ. 1979 Ca^{2+}-induced bioluminescence in *Renilla reniformis*. Purification and characterization of a calcium-triggered luciferin-binding protein. *J. Biol. Chem.* **254(3)**, 769-780. (https://doi.org/10.1016/S0021-9258(17)37872-9)

27. Stepanyuk GA, Liu Z-J, Vysotski ES, Lee J, Rose JP, Wang B-C. 2008 Structure based mechanism of the Ca^{2+}-induced release of coelenterazine from the *Renilla* binding protein. *Proteins* **74(3)**, 583-593. (https://doi.org/10.1002/prot.22173)

28. Nicol JAC. 1955 Observations on luminescence in *Renilla* (Pennatulacea). *J. Exp. Biol.* **51232(2)**, 299-320.

29. Davenport D, Nicol JAC. 1956 Observations on luminescence in sea pens (Pennatulacea). *Proc. R. Soc. B* **144(917)**, 480-496. (https://doi.org/10.1098/rspb.1956.0005)

30. Wampler JE, Karkhanis YD, Morin JG, Cormier MJ. 1973 Similarities in the bioluminescence from the Pennatulacea. *Biochimica et Biophysica Acta* **314**, 104-109. (https://doi.org/10.1016/0005-2728(73)90068-6)

31. Satterlie RA, Anderson PAV, Case JF. 1980 Colonial coordination on anthozoans: Pennatulacea. *Mar. Behav. Physiol.* **7**, 25-46. (https://doi.org/10.1080/10236248009386969)

32. Germain G, Anctil M. 1988 Luminescent activity and ultrastructural characterization of photocytes dissociated from the coelenterate *Renilla köllikeri*. *Tissue and Cell* **20(5)**, 701-720. (https://doi.org/10.1016/0040-8166(88)90017-1)

33. Anctil M, Boulay D, Larivière L. 1982 Monoaminergic mechanisms associated with control of luminescence and contractile activities in the coelenterate, *Renilla köllikeri*. *J. Exp. Zool.* **223(1)**, 11-24. (https://doi.org/10.1002/jez.1402230103)

34. Duchatelet L, Coubris C, Pels C, Dupont S, Mallefet J. 2023 Catecholamine involvement in the bioluminescence control of two species of Anthozoans. *Life* **13(9)**, 1798. (https://doi.org/10.3390/life13091798)

35. Morin JG. 1976 Probable functions of bioluminescence in the Pennatulacea (Cnidaria, Anthozoa). In: Mackie GO.(Ed) *Coelenterate Ecology and Behavior*. Springer, Boston, MA, USA. (https://doi.org/10.1007/978-1-4757-9724-4_65)

36. Delroisse J, Mallefet J, Flammang P. 2016 *De novo* adult transcriptomes of two European brittle stars: spotlight on opsin-based photoreception. *PLoS ONE* **11(4)**, e0152988. (https://doi.org/10.1371/journal.pone.0152988)

37. Delroisse J, Duchatelet L, Flammang P, Mallefet J. 2018 *De novo* transcriptome analyses provide insights into opsin-based photoreception in the lanternshark *Etmopterus spinax*. *PLoS ONE* **13(12)**, e0209767. (https://doi.org/10.1371/journal.pone.0209767)

38. Delroisse J, Duchatelet L, Flammang P, Mallefet J. 2021 Photophore distribution and enzymatic diversity within the photogenic integument of the cookie-cutter shark *Isistius brasiliensis* (Chondrichthyes: Dalatiidae). *Front. Mar. Sci.* **8**, 627045. (https://doi.org/10.3389/fmars.2021.627045)

39. Cock PJA, Fields CJ, Goto N, Heuer ML, Rice PM. 2010 The Sanger FASTQ file format for sequences with quality scores and the Solexa/Illumina FASTQ variants. *Nucleic Acids Res.* **38(6)**, 1767-1771. (https://doi.org/10.1093/nar/gkp1137)

40. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. 2011 Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **29**, 644-652. (https://doi.org/10.1038/nbt.1883)

41. Pertea G, Huang X, Liang F, Antonescu V, Sultana R, Karamycheva S, et al. 2002 TIGR gene indices clustering tools (TGICL): a software system for fast clustering of large EST datasets. *Bioinformatics* **19(5)**, 651-652. (https://doi.org/10.1093/bioinformatics/btg034)

42. Das S, Shyamal S, Durica DS. 2016 Analysis of annotation and differential expression methods used in RNA-seq studies in crustacean systems. *Integr. Comp. Biol.* **56(6)**, 1067-1079. (https://doi.org/10.1093/icb/icw117)

43. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990 Basic local alignment search tool. *J. Mol. Biol.* **215(3)**, 403-410. (https://doi.org/10.1016/S0022-2836(05)80360-2)

44. Buchfink B, Xie C, Huson DH. 2014 Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* **12**, 59-60. (https://doi.org/10.1038/nmeth.3176)

45. Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. 2005 Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* **21(18)**, 3674-3676. (https://doi.org/10.1093/bioinformatics/bti610)

46. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. 2005 InterProScan: protein domains identifier. *Nucleic Acids Res.* **33**, W116-W120. (https://doi.org/10.1093/nar/gki442)

47. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. 2012 Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28(12)**, 1647-1649. (https://doi.org/10.1093/bioinformatics/bts199)

48. Woo J, Howell MH, von Arnim AG. 2009 Structure-function studies on the active site of the coelenterazine-dependent luciferase from *Renilla*. *Protein Sci.* **17(4)**, 725-735. (https://doi.org/10.1110/ps.073355508)

49. McFadden CS, van Ofwegen LP, Quattrini AM. 2022 Revisionary systematics of Octocorallia (Cnidaria: Anthozoa) guided by phylogenomics. *Bulletin of the Society of Systematics Biologists* **1(3)**. (https://doi.org/10.18061/bssb.v1i3.8735)

50. Labas YA, Gurskaya NG, Yanushevich YG, Fradkov AF, Lukyanov KA, Lukyanov SA, Matz MV. 2002 Diversity and evolution of the green fluorescent protein family. *Proc. Natl Acad. Sci.* **99(7)**, 4257-4261. (https://doi.org/10.1073/pnas.062552299)

51. Alieva NO, Konzen KA, Field SF, Meleshkevitch EA, Hunt ME, Beltran-Ramirez V, Miller DJ, Wiedenmann J, Salih A, Matz MV. 2008 Diversity and evolution of coral fluorescent proteins. *PLoS ONE* **3(7)**, e2680. (https://doi.org/10.1371/journal.pone.0002680)

52. Li G, Zhang QJ, Zhong J, Wang YQ. 2009 Evolutionary and functional diversity of green fluorescent proteins in cephalochordates. *Gene* **446(1)**, 41-49. (https://doi.org/10.1016/j.gene.2009.07.003)

53. Shagin DA, Barsova EV, Yanushevich YG, Fradkov AF, Lukyanov KA, Labas YA, Semenova TN, Ugalde JA, Meyers A, Nunez JM, Widder EA, Lukyanov SA, Matz MV. 2004 GFP-like proteins as ubiquitous metazoan superfamily: evolution of functional features and structural complexity. *Mol. Biol. Evol.* **21(5)**, 841-850. (https://doi.org/10.1093/molbev/msh079)

54. Schnitzler CE, Pang K, Powers ML, Reitzel AM, Ryan JF, Simmons D, Tada T, Park M, Gupta J, Brooks SY, Blakesley RW, Yokoyama S, Haddock SHD, Martindale MQ, Baxevanis AD. 2012 Genomic organization, evolution, and expression of photoprotein and opsin genes in *Mnemiopsis leidyi*: a new view of ctenophore photocytes. *BMC Biol.* **10**, 1-26. (https://doi.org/10.1186/1741-7007-10-107)

55. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009 trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973. (https://doi.org/10.1093/bioinformatics/btp348)

56. Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015 IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol.*
Biol. Evol. **32(1)**, 268-274. (https://doi.org/10.1093/molbev/msu300)

57. Posada D, Crandall KA. 1998 MODELTEST: testing the model of DNA substitution.
Bioinformatics (Oxford, England) **14(9)**, 817-818.
(https://doi.org/10.1093/bioinformatics/14.9.817)

58. Rahnama S, Saffar B, Kahrani Z.F, Nazari M, Emamzadeh R. 2017 Super RLuc8: a
novel engineered *Renilla* luciferase with a red-shifted spectrum and stable light
emission. *Enzym. Microb. Technol.* **96**, 60-66.
(https://doi.org/10.1016/j.enzmictec.2016.09.009)

59. Khoshnevisan G, Emamzadeh R, Nazari M, Rasa SMM, Sariri R, Hassani L. 2018
Kinetics, structure, and dynamics of *Renilla* luciferase solvated in binary mixtures of
glycerol and water and the mechanism by which glycerol obstructs the enzyme emitter
site. *Int. J. Biol. Macromol.* **117**, 617-624.
(https://doi.org/10.1016/j.ijbiomac.2018.05.160)

60. Mallefet J, Duchatelet L, Coubris C. 2020 Bioluminescence induction in the ophiuroid
Amphiura filiformis (Echinodermata). *J. Exp. Biol.* **223(4)**, jeb218719.
(https://doi.org/10.1242/jeb.218719)

61. Mallefet J,Martinez-Soares P, Eléaume M, O'Hara T, Duchatelet L. 2023 New insights
on crinoid (Echinodermata; Crinoidea) bioluminescence. *Front. Mar. Sci.* **10**, 1136138.
(https://doi.org/10.3389/fmars.2023.1136138)

62. Loening AM, Fenn TD, Wu AM, Gambhir SS. 2006 Consensus guided mutagenesis of
Renilla luciferase yields enhanced stability and light output. *Protein Eng. Des. Sel.*
19(9), 391–400. (https://doi.org/10.1093/protein/gzl023)

63. Tsuji FI, Ohmiya Y, Fagan TF, Toh H, Inouye S. 1995 Molecular evolution of the Ca^{2+}-
binding photoproteins of the hydrozoa. *Photochem. Photobiol.* **62(4)**, 657-661.
(https://doi.org/10.1111/j.1751-1097.1995.tb08713.x)

64. Powers ML, McDermott AG, Shaner NC, Haddock SHD. 2013 Expression and
characterization of the calcium-activated photoprotein from the ctenophore *Bathocyroe*
fosteri: Insights into light-sensitive photoproteins. *Biochem. Biophys. Res. Commun.*
431(2), 360-366. (https://doi.org/10.1016/j.bbrc.2012.12.026)

65. Burakova LP, Stepanyuk GA, Eremeeva EV, Vysotski ES. 2016 Role of certain amino
acid residues of the coelenterazine-binding cavity in bioluminescence of light-sensitive
Ca^{2+}-regulated photoprotein berovin. *Photochem. Photobiol. Sci.* **15**, 691-704.
(https://doi.org/10.1039/c6pp00050a)

66. Burakova LP, Vysotski ES. 2019 Recombinant Ca2+-regulated photoproteins of
ctenophores: Current knowledge and application prospects. *Appl. Microbiol.*
Biotechnol. **103**, 5929-5946. (https://doi.org/10.1007/s00253-019-09939-0)

67. Buskey EJ, Stearns DE. 1991 The effects of starvation on bioluminescence potential
and egg release of the copepod *Metridia longa*. *J. Plankton Res.* **13(4)**, 885-893.
(https://doi.org/10.1093/plankt/13.4.885)

68. Thomson CM, Herring PJ, Campbell AK. 1995 Evidence for *de novo* biosynthesis of
coelenterazine in the bioluminescent midwater shrimp, *Systellaspis debilis* C. *J. Mar.*
Biol. Assoc. U.K. **75(1)**, 165-171. (https://doi.org/10.1017/S0025315400015277)

69. Oba Y, Kato S-I, Ojika M, Inouye S. 2009 Biosynthesis of coelenterazine in the deep-sea copepod, *Metridia pacifica*. *Biochem. Biophys. Res. Commun.* **390(3)**, 684-688. (https://doi.org/10.1016/j.bbrc.2009.10.028)

70. Bessho-Uehara M, Huang W, Patry WL, Browne WE, Weng J-K, Haddock SHD. 2020 Evidence for *de novo* biosynthesis of the luminous substrate coelenterazine in ctenophores. *iScience* **23(12)**, 101859. (https://doi.org/10.1016/j.isci.2020.101859)

71. Francis WR, Shaner NC, Christianson LM, Powers ML, Haddock SHD. 2015 Occurrence of isopenicillin-N-synthase homologs in bioluminescent ctenophores and implications for coelenterazine biosynthesis. *PLoS ONE* **10(6)**, e0128742. (https://doi.org/10.1371/journal.pone.0128742)

72. Haddock SHD, Rivers TJ, Robison BH. 2001 Can coelenterates make coelenterazine? Dietary requirement for luciferin in cnidarian bioluminescence. *Proc. Ntl Acad. Sci.* **98(20)**, 11148-11151.(https://doi.org/10.1073/pnas.201329798)

73. Nicol JAC. 1958 Observations on the luminescence of *Pennatula phosphorea*, with a note on the luminescence of *Virgularia mirabilis*. *J. Mar. Biol. Assoc. UK* **37**, 551–563. (https://doi.org/10.1017/S0025315400005610)

74. Herring PJ. 1991 Observations on bioluminescence in some deep-water anthozoans. In *Coelenterate Biology: Recent Research on Cnidaria and Ctenophora, Proceedings of the Fifth International Conference on Coelenterate Biology*, Southampton, UK, 10–14 July 1989; Springer: Dordrecht, The Netherlands; pp. 573–579.

75. Coubris C, Duchatelet L, Delroisse J, Bayaert WS, Parise L, Eloy MC, Pels C, Mallefet J. 2024 Maintain the light, long-term seasonal monitoring of luminous capabilities in the brittle star *Amphiura filiformis*. *Sci. Rep. Submitted.*

76. Inouye S, Watanabe K, Nakamura H, Shimomura O. 2000 Secretional luciferase of the luminous shrimp *Oplophorus gracilirostris*: cDNA cloning of a novel imidazopyrazinone luciferase. *FEBS Lett.* **481(1)**, 19-25. (https://doi.org/10.1016/S0014-5793(00)01963-3)

77. Kobilka B. 1992 Adrenergic receptors as models for G protein-coupledreceptors. *Ann. Rev. Neurosci.* **15(1)**, 87-114. (https://doi.org/10.1146/annurev.ne.15.030192.000511)

78. Caron MG, Lefkowitz RJ. 1993 Catecholamine receptors: structure, function, and regulation. *Recent Prog. Horm. Res.* **1993**, 277-290. (https://doi.org/10.1016/B978-0-12-571148-7.50014-2)

79. Neves SR, Ram PT, Iyengar R. 2002 G protein pathways. *Science* **296(5573)**, 1636-1639. (https://doi.org/10.1126/science.1071550)

80. Brown AM, Birnbaumer L. 1988 Direct G protein gating of ion channels. *Am. J. Physiol. Heart Circ. Physiol.* **254(3)**, H401-H410. (https://doi.org/10.1152/ajpheart.1988.254.3.H401)

81. Hille B. 1994 Modulation of ion-channel function by G-protein-coupled receptors. *Trends Neurosci.* **17(12)**, 531-536. (https://doi.org/10.1016/0166-2236(94)90157-0)

82. Dolphin AC. 2003 G protein modulation of voltage-gated calcium channels. *Pharmacol. Rev.* **55(4)**, 607-627. (https://doi.org/10.1124/pr.55.4.3)

83. Tedford HW, Zamponi GW. 2006 Direct G protein modulation of Cav2 calcium channels. *Pharmacol. Rev.* **58(4)**, 837-862. (https://doi.org/10.1124/pr.58.4.11)

84. Roda A, Pasini P, Mirasoli M, Michelini E, Guardigli M. 2004 Biotechnological
applications of bioluminescence and chemiluminescence. *Trends Biotechnol.* **22(6)**,
295-303. (https://doi.org/10.1016/j.tibtech.2004.03.011)
85. Kirkpatrick A, Xu T, Ripp S, Sayler G, Close D. 2019 Biotechnological advances in
luciferase enzymes. *Bioluminescence-Analytical Applications and Basic Biology*, 1-23.
(https://doi.org/10.5772/intechopen.85313)
86. Welsh DK, Kay SA. 2005 Bioluminescence imaging in living organisms. *Curr. Opin.*
Biotechnol. **16(1)**, 73-78. (https://doi.org/10.1016/j.copbio.2004.12.006)