1 FRONT MATTER
2

3 Title: **Mechanical regulation of cellular energy metabolism in cancer microenvironments**
4 **Short title:** Mechano-metabolism of metastatic cells.
5

6 Joshua M. Toth[1,2]†, Anuja Jaganathan[1,2]†, Ramin Basir[1,2], Laurent Pieuchot[3], Yihui Shen[4],
7 Cynthia A. Reinhart-King[5], Vivek B. Shenoy[1,2,4,6]*
8

[1]Center for Engineering Mechanobiology, University of Pennsylvania, PA, US
[2]Department of Materials Science and Engineering, University of Pennsylvania, PA, US
[3]Université de Haute-Alsace, CNRS, IS2M UMR 7361, 61800, Mulhouse, France
[4]Department of Bioengineering, University of Pennsylvania, PA, US
[5]Department of Bioengineering, Rice University, Houston, TX, US
[6]Department of Mechanical Engineering and Applied Mathematics, University of Pennsylvania,
PA, US
16

17 †Equal Contribution
18

*Correspondence to Vivek B. Shenoy
Email: vshenoy@seas.upenn.edu

Abstract
Cells dynamically regulate their morphology, contractility, and metabolism in response to the
mechano-chemical properties of their microenvironment. Here, we show matrix stiffness and ligand
density jointly govern the bioenergetics of contractile cells through a nonequilibrium active chemo-
mechanical model built around a newly introduced cellular metabolic potential. This concept links
ATP hydrolysis to mechanosensitive signaling, quantifies the energetic cost of stress fiber
assembly, and determines mechanically stable states. The metabolic potential enables quantitative
prediction of cell contractility, morphology, and ATP consumption in different stiffness 2D and 3D
environments, and we find quantitative agreement with experimental measurements in MDA-MB-
231 breast cancer cells. The model further predicts activation of AMPK accompanies increased
energetic demands in stiffer microenvironments which we experimentally validate and correlate
with increased mitochondrial membrane potential, glucose uptake, and intracellular ATP levels.
Together, these findings establish a predictive quantitative framework unifying mechanosensitive
control of cell shape and contractility with the metabolic pathways sustaining cellular function
across diverse mechanical environments.

Teaser
Matrix stiffness reshapes the cellular energy budget, driving metabolic adaptation to mechanical
demand.
41

42 **MAIN TEXT**
43

44 **INTRODUCTION**
45

46 Cells sense and respond to mechanical cues through cytoskeletal reorganization and regulation of
   actomyosin contractility (*1*). Changes in contractility alter cell stiffness (*2*), shape (*3*), and
47 differentiation (*4,5*) via mechanosensitive biochemical pathways. Mechanical inputs also regulate
48 cellular metabolism through diverse mechanisms, including cytoskeletal sequestration of glycolytic
49 enzymes (*6–9*), modulation of lipid synthesis (*10,11*), alterations in mitochondrial morphology and

function (*12,13*), and YAP/TAZ signaling (*14,15*). A rapid consequence of increasing matrix stiffness is AMP-activated protein kinase (AMPK)-dependent upregulation of ATP production (*16–18*). As a master energy sensor, AMPK monitors intracellular ATP levels and coordinates glucose uptake (*19,20*), glycolysis (*21,22*), oxidative phosphorylation (*23*), protein synthesis (*24*), and autophagy (*25*), thereby maintaining energy homeostasis. Since generating and sustaining contractile forces is energetically costly, mechanotransduction and metabolic regulation are necessarily coupled. This coupling is particularly relevant in cancer progression and metastasis (*26,27*), where tumor cells reprogram metabolism in response to mechanical cues from their microenvironment (*28*). Despite extensive evidence linking matrix mechanics to metabolic adaptation, this interplay remains largely qualitative. A predictive quantitative framework that integrates mechanosensitive signaling with metabolic regulation is lacking.

Here, we develop a nonequilibrium chemo-mechanical model of contractility and adhesion centered on a newly introduced cellular "metabolic potential." This nonequilibrium thermodynamic quantity explicitly links ATP hydrolysis by myosin-II motors to mechanosensitive stress-fiber assembly, accounting for the partitioning of chemical energy into contractile work and dissipation. Within this framework, cells adopt contractile states and morphologies that minimize their metabolic potential in a given mechanical microenvironment. The model predicts cell shape, contractility, and ATP consumption across substrates of varying stiffness and reveals that differences between 2D and 3D morphologies emerge solely from geometry-dependent stress distributions that reshape downstream chemo-mechanical signaling. We further incorporate AMPK-mediated energy replenishment and show that homeostatic regulation can sustain elevated intracellular ATP levels in highly contractile cells. These predictions are validated experimentally through measurements of cell morphology, ATP:ADP ratios, phosphorylated AMPK levels, glucose uptake, and mitochondrial membrane potential in MDA-MB-231 cancer cells cultured across stiffness gradients. Together, this work establishes a predictive quantitative framework linking cellular mechanics, mechanosignaling, and metabolic adaptation.

RESULTS

The Metabolic Potential Links ATP Hydrolysis to Cellular Tension

To elucidate the key principles underlying mechanosensitive metabolism, we begin with a 1D formulation, which applies to mesenchymal cells adhered between deformable microposts (Figure 1C). In contractile cells, myosin II motors bind to F-actin filaments and generate contractile forces by forming stress fibers, thereby converting the chemical energy released by ATP hydrolysis into mechanical work (*29,30*). We model individual myosin motors as force dipoles (Figure S1). The contractile stress generated by motors engaged in stress fibers is represented by ρ, while the cytoskeletal stress and strain are denoted by σ and ε, respectively. In a quiescent, non-adherent cell, we assume a baseline level of contractility, $\rho = \rho_0$, determined by the balance between the decrease in enthalpy upon motor binding to actin and the associated loss of entropy relative to unbound motors in the cytosol. Upon establishing adhesions, mechanosensitive signaling increases both myosin motor activity and stress fiber assembly. Specifically, cell-generated tension activates signaling pathways including Rho–ROCK (*31,32*) and Ca^{2+}/calmodulin (*33*) (Figure 1A) which increase motor recruitment and contraction via ATP hydrolysis. A portion of the energy released by ATP hydrolysis is converted into (i) chemical energy associated with actin–myosin binding ($U_{binding}(\rho, \varepsilon)$), (ii) mechanical work done by active motors ($U_{motor}(\rho, \varepsilon)$), and (iii) elastic strain energy stored in the cytoskeleton ($U_{strain}^{cell}(\varepsilon)$) and matrix ($U_{strain}^{ECM}(\varepsilon)$) (Figure 1B).

To describe the energy available from ATP hydrolysis, we incorporate the mechanosensitive kinetics of stress fiber assembly and disassembly through stress-dependent rate constants $k_{on}(\sigma)$ and $k_{off}(\sigma)$ (Figure 1A). These rates represent the timescale of stress fiber assembly and the lifetime of existing fibers, respectively (Figure S2). A lower disassembly rate corresponds to longer-lived stress fibers, such that $k_{off} \propto 1/\tau_{off}$, with τ_{off} the fiber lifetime. The fraction of time that myosin motors remain engaged with actin and hydrolyze ATP is then $\tilde{\tau} = k_{on}(\sigma)/(k_{on}(\sigma)+k_{off}(\sigma))$. When assembly is fast and disassembly is slow, motor engagement—and thus ATP consumption—is maximized. Therefore, the time-averaged energy released by ATP hydrolysis at steady state is $U_{ATP} = \tilde{\tau}\rho\Delta G/p_0$, where $\tilde{\tau}\rho$ is the time-averaged density of ATP-hydrolyzing stress fibers, ΔG is the free energy released per ATP hydrolyzed, and p_0 is the dipole strength of a single myosin dipole.

ATP hydrolysis supplies the energy needed to drive otherwise unfavorable processes, including increases in contractility and cytoskeletal and ECM deformation (Figure 1B). Because mechanosensitive signaling modulates myosin ATPase activity and stress fiber organization, it also determines how much ATP-derived energy is accessible to the cell. To capture this relationship, we introduce a metabolic potential, R, defined as the sum of conservative energies and the energy released by ATP hydrolysis (Equation S1.11). As shown in SI Section 1.1.4, minimizing R with respect to contractility and strain is equivalent to predicting the nonequilibrium steady state arising from stress fiber assembly and disassembly kinetics. Thus, the cell selects the configuration (ρ, σ, ε) that minimizes the metabolic potential—corresponding to maximal ATP hydrolysis.

To illustrate this concept, we examine a cell adhered between two microposts (Figure 1C). The cell consists of a passive elastic element of bulk modulus K and an active contractile element representing stress fibers, while each micropost is modeled as an elastic element of modulus K_p in series with the cell (parameter values listed in Table S1). Minimizing the metabolic potential (SI Section 1.2) predicts that steady-state contractility increases with micropost stiffness (Figure 1D). The predicted cellular traction force (SI Section 1.3) also increases with post stiffness and agrees well with experimental measurements of myoblasts contracting against deformable substrates (34,35). Stiffer microposts elevate mechanosensitive signaling, activate more myosin motors, and increase ATP hydrolysis, which in turn drives stronger cell contraction. The detailed dependence of mechanical energies on K, K_p, and other parameters is provided in SI Section 1.2 and Figure S3.

Stress regulates the coupling between signaling and contractility such that $k_{on}(\sigma)$ increases and $k_{off}(\sigma)$ decreases with stress. This formulation is consistent with experimentally measured kinetics of myosin II isoforms. Myosin IIB exhibits strong catch-bond behavior, with tension decreasing its detachment rate (lower k_{off} at higher σ) and thus extending stress fiber lifetime (36). However, its force-dependent recruitment (increase in k_{on}) is modest and cell-type specific, depending on co-assembly with myosin IIA (37,38). Conversely, myosin IIA is strongly recruited under tension (higher k_{on}), although its catch-bond behavior is weaker (38). The stress-dependent forms of k_{on} and k_{off} (Equation S1.22) therefore capture both mechano-signaling–dependent motor activation and isoform-specific recruitment and dissociation kinetics.

Figure 1: A model for the establishment of stress-regulated contractility driven by ATP hydrolysis. (A) Schematic illustrating the stress-dependent pathways that regulate cell contractility. Tensile forces transmitted from the micro-environment act through adhesion complexes at the cell-matrix interface and trigger: (I) Stress-assisted activation of signaling molecules (like Src Family Kinases (SFKs)), which act on Rho-GTPases and promote ROCK-mediated phosphorylation of myosin light chain (MLC) and consequently, stress fiber assembly. ROCK activation also leads to inhibition of myosin phosphatase (MYPT) which downregulates myosin unbinding and stress fiber disassembly. (II) Generation of cell membrane tension leading to the opening of stress-controlled ion-channels, an influx of Ca^{2+} into the cytoplasm, and upregulation of motor activation by myosin light chain kinase (MLCK) activation. **(B)** Schematic showing how the energy released from ATP

hydrolysis by active motors in stress fibers depends on the kinetics of stress fiber assembly and
comprises the dissipative part of the energy, some of which is converted into other conservative
forms while the rest is dissipated as heat. We define the sum of all energies as the metabolic
potential. (**C**) A cell adhered between microposts is modeled with active contractile element
representing myosin motors in parallel with a linear elastic spring that describes passive
cytoskeleton components. (**D**) Contributions to the metabolic potential for cells adhered to
microposts as a function of contractility. Plots indicate how the various energies evolve as
microposts become stiffer. The metabolic potential shows an optimum at a higher contractility on
stiff (ρ_{stiff}) than on soft microposts (ρ_{soft}).

A 3D Chemo-mechanical formulation of the metabolic potential

While the 1D model captures stiffness sensing, real cells exhibit spatially heterogeneous and
anisotropic stress distributions. We therefore generalize the framework to three dimensions, where
contractility, stress, and strain become tensors, ($\rho_{ij}(\mathbf{x})$, $\sigma_{ij}(\mathbf{x})$, and $\epsilon_{ij}(\mathbf{x})$) that can vary with spatial
coordinate, $\mathbf{x}$, in the cell. In elongated cells, stress concentrates near elongated edges (Figure S4A),
triggering mechanosensitive signaling and promoting preferential stress fiber formation along the
long axes, producing a polarized contractility field ($\rho_{11}(\mathbf{x}) \neq \rho_{22}(\mathbf{x}) \neq \rho_{33}(\mathbf{x})$) (*39*). To capture this
anisotropy, the assembly and disassembly rates, $k_{on,ij}$ and $k_{off,ij}$, are also treated as spatially
dependent tensor quantities. Their volumetric components ($k^v_{on}(\sigma_{kk}(\mathbf{x}))$, $k^v_{off}(\sigma_{kk}(\mathbf{x}))$) represent
mean-stress–dependent changes in assembly/disassembly, while their deviatoric components
($\tilde{k}_{on,ij}(\tilde{\sigma}_{ij}(\mathbf{x}))$ and $\tilde{k}_{off,ij}(\tilde{\sigma}_{ij}(\mathbf{x}))$) capture anisotropic stress-driven polarization of stress fibers. This
tensorial description is necessary to reproduce experimentally observed spatial polarization of
myosin densities in cells with anisotropic morphologies. Because anisotropic stress also upregulates
contractility in elongated cells (*39*), the 3D model includes an additional anisotropy-dependent term
in the assembly rate (Equation S2.14). We further introduce a cell–matrix interfacial energy, $\Gamma(\Theta)$,
which accounts for membrane/cortical tension and adhesion energy, and depends on cell shape,
described by Θ (Figure 2A).

Performing a spatial integration of all contributions over their respective domains yields the
generalized metabolic potential:

$$R = \int_{V_c} R_l\left(\rho_{ij}(\mathbf{x}), \varepsilon_{ij}(\mathbf{x}), \Theta\right) dV_c$$

$$+ \int_{V_{ECM}} F_{nuc}\left(\varepsilon_{ij}^{nuc}(\mathbf{x})\right) dV_{nuc} + \int_{V_{ECM}} F_{ECM}\left(\varepsilon_{ij}^{ECM}(\mathbf{x})\right) dV_{ECM} + \Gamma(\Theta) \tag{1}$$

171 where R_l is the local (non-integrated) metabolic potential at every coordinate, $\mathbf{x}$ (Equation 3 in
172 Materials and Methods), and F_{ECM} and F_{nuc} are strain energy densities of the ECM and nucleus. As
173 in the 1D formulation, minimizing R corresponds to the steady state of the full chemo-mechanical
174 system governed by stress fiber kinetics (SI Section 2.2.3). Thus, both the metabolic potential and
175 the ATP hydrolysis energy follow directly from nonequilibrium thermodynamic arguments.
176

Metabolic potential predicts biphasic changes in cell morphology with increasing collagen ECM density

We implemented the generalized 3D model within a finite element framework (Figure S4C) to
investigate the physical mechanisms that govern the establishment of cell morphology, cytoskeletal
contractility, and metabolic activity in MDA-MB-231 breast cancer cells embedded in three-

dimensional collagen matrices. The cell body is represented as an ellipsoid characterized by two
semiaxes of equal radius and a third distinct radius ($a = c \neq b$), and the cell aspect ratio is defined
as AR = a/b (Figure 2B). Thus, a spherical cell has AR = 1, whereas AR > 1 denotes an elongated
cell. To consistently parameterize changes in cellular geometry, we normalized the aspect ratio by
the cell circularity, circ=4π(area)/(perimeter)^2, resulting in an elongation factor (EF) that captures
shape variations across conditions (Figure 2B). Although cellular volume can vary with ECM
stiffness, we assume that shape change is a volume-conserving process and set the volume of each
cell in the model to 1500 μm^3 (*40*). This assumption is appropriate because changes in cell volume
are regulated by water transport, which requires osmotic pressures on the order of ~1 MPa (*41*) -
approximately three orders of magnitude larger than typical cytoskeletal tension (~1–10 kPa) -
making volume changes negligible relative to mechanically driven shape changes within collagen
matrices.

Cell morphology, cytoskeletal contractility, and energy expenditure in 3D collagen matrices depend
strongly on collagen density (*16,42*). To capture this mechano-regulation, we modeled the collagen
network using a nonlinear fiber formulation (*43*) (see SI Section 2.2.5 for details; model parameters
are listed in Table S2), which accounts for tension-induced fiber alignment and the resulting
anisotropic matrix stiffening (Figure S4B). For each collagen density, we determined the steady-
state cell shape, strain field, and contractility by solving the constitutive equations (Equations
S2.20–S2.21) obtained from minimizing the metabolic potential, R (Equation 1), with respect to
these variables. We repeated this analysis across a range of matrix densities and iteratively
compared model predictions of cell elongation and contractility with experimental measurements
for MDA-MB-231 cells to calibrate the mechanical and biochemical model parameters (Table S3).
204
205 *Actomyosin contractility and ATP hydrolysis rates are higher in elongated cells*
206
Having established the computational framework, we examined how cell shape and collagen
density jointly influence emergent contractility and energetics. As predicted in the 1D micropost
setting, the steady-state volume-averaged contractility, $1/V_c \int \rho_{kk}(\mathbf{x})dV_c$, increases with collagen
density because initial cell contraction generates larger volumetric stresses, $1/V_c \int \sigma_{kk}(\mathbf{x})dV_c$ in
denser matrices, which in turn lead to stronger stress-dependent actomyosin upregulation.
Similarly, for any fixed matrix density, elongated cells exhibit higher steady-state contractility than
spherical cells (Figure S5). This enhancement arises because elongated geometries generate larger
stresses at the cell–matrix interface (Figure 2C), thereby increasing the activation of
mechanosensitive signaling pathways and promoting the recruitment and activation of actomyosin
machinery (Figure 1A), analogous to the effects observed in stiffer matrices.

Interestingly, in high-density collagen matrices ($\geq$ 4 mg/ml), a different mode of stress-dependent
regulation emerges. While the volume-averaged contractility increases monotonically with
elongation, the volumetric stress exhibits a biphasic trend (Figure S5). This occurs because, at
constant cell volume, further elongation eventually produces diminishing increases in the stress
along the elongated axis (σ_{33}) that are outweighed by reductions in the stresses along the transverse
axes (σ_{22} and σ_{11}). In soft collagen matrices, this behavior only appears at unrealistically large
elongation (EF > 50), but in stiff collagen it occurs within physiologically relevant elongation levels
(0 < EF < 10). However, our 3D model also includes stress-dependent actomyosin regulation arising
from stress anisotropy (Equation S2.14), and anisotropy increases continuously with elongation at
all collagen densities. Consequently, in high-density collagen, anisotropy-dependent signaling

dominates over mean-stress–dependent signaling, causing the volume-averaged contractility to
increase monotonically with elongation even when the axial stress, σ_{33} begins to decrease.

Figure 2: Minimizing the metabolic potential predicts a biphasic variation in cell shape with increasing collagen density. (A) Interfacial energy comprised of membrane/cortical tension, γ_0, which resists increases in cell surface area, and adhesion energy of integrin-collagen binding, γ_1, which favors cell elongation as increased surface area allows for more binding interactions. (B) Schematic description of cell body aspect ratio (AR), circularity (circ) and their variation with cell shape. (C) Optimum cell shape shows a biphasic transition from spherical in low density collagen to elongated (spindle-shaped) in medium density collagen, back to nearly spherical in high density collagen resulting from the competition between the volumetric part of the metabolic potential (red block arrows) and the interfacial energy (yellow block arrows). (D) Metabolic potential as a function of cell shape for low (0.4 mg/mL), medium (1 mg/mL), and high density (5 mg/mL) collagen, where the minimum in metabolic potential corresponds to the steady state optimum cell

shape. Simulation predictions as compared to reported experimental measurements of MDA-MB-231 cells in 3D collagen for (**E**) cell elongation (*16*) and (**G**) normalized ATP consumption rate (*42*). The experimentally measured optical retardance signal (*45*) indicating contractile strain on f-actin is displayed in (**F**) along with model predictions of steady state contractility vs collagen density. Bar plots show mean ±SE, and box plots show median, $25^{th}/75^{th}$ percentiles, and $5^{th}/95^{th}$ percentiles *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.001$ with Kruskal-Wallis one-way ANOVA (**E**) or Tukey's HSD post hoc test (**G**). Scale bar = 25 μm.

Consistent with this interpretation, our model captures the emergence of stress polarization in anisotropic cell shapes. Examination of the deviatoric stress components reveals that the component aligned with the elongated axis ($\tilde{\sigma}_{33}$) is significantly greater than those along the other axes. This polarization drives the preferential alignment of stress fibers along the first principal stress direction (Figure S6). These trends in stress and contractility determine the observed behavior of the metabolic potential and its volumetric portion, v^{cell} (all terms in Equation 1 except $\Gamma(\Theta)$), reflecting the central role of stress-dependent signaling in modulating actomyosin activity.

The optimum cell aspect ratio minimizing metabolic potential exhibits biphasic variation with collagen density

Metabolic potential dictates the optimum cell morphology with changing cell shape and collagen density through competition between its volumetric component, v^{cell}, and the interfacial energy, $\Gamma(\Theta)$. Although all energetic terms in the volumetric metabolic potential closely track the changes in contractility across matrix densities and cell shapes, the decrease in ATP hydrolysis energy shows the largest variation with cell elongation (Figure S7). Since elongated cells develop higher contractility and thus higher ATP consumption, they exhibit lower v^{cell}. However, elongation also increases cell surface area thereby increasing interfacial energy, which incorporates both membrane tension and adhesion-related contributions (Figure 2A). As detailed in SI Section 2.2.5, membrane tension is approximately two orders of magnitude larger than receptor–ligand adhesion energy and increases with both cell surface area and matrix density. Moreover, dense collagen matrices present fewer accessible adhesion sites because matrix degradation is limited (*44*). To capture these effects, the interfacial energy, $\Gamma(\Theta)$, is implemented as an increasing function of both cell surface area and collagen density and introduces an energetic penalty for elongation (Figure 2C). The specific functional form was determined through sensitivity analysis (Figure S8 and Table S3).

The competition between the decrease in the volumetric part of the metabolic potential, v^{cell}, and the increase in the interfacial energy, $\Gamma(\Theta)$, generates a biphasic trend in the optimum cell aspect ratio as collagen density increases:
Low collagen density (< 0.5 mg/ml): In soft collagen matrices, cells cannot generate substantial stresses even when elongated (Figure 2C). The resulting stress-regulated enhancement of stress fiber assembly, and thus the ATP hydrolysis energy, is small for both spherical and elongated shapes (Figure S7). Consequently, the small decrease in the volumetric part of the metabolic potential, v^{cell}, associated with elongation is insufficient to overcome the increase in the interfacial energy, $\Gamma(\Theta)$. The metabolic potential therefore predicts a spherical shape (Figure 2D), consistent with experimental observations (Figure 2E).

Intermediate collagen density (0.5–3 mg/ml):   At moderate collagen densities, the matrix can support the generation of larger stresses and higher contractility when the cell elongates (Figure 2C). As a result, the reduction in the volumetric part of the metabolic potential due to elongation becomes large enough to overcome the interfacial energy penalty, and the model predicts elongated cell morphologies (Figure 2D), again matching experimental data (Figure 2E).

High collagen density (> 3 mg/ml): In dense matrices, the decrease in the volumetric part of the metabolic potential, v^{cell}, that accompanies elongation is insufficient to compensate for the increase in interfacial energy beyond an aspect ratio of approximately 3, and the cell therefore adopts a moderately elongated shape (Figure 2D). This prediction is in close agreement with experimental measurements (Figure 2E). Notably, the difference in v^{cell} between spherical and elongated cells is smaller in dense matrices, since high matrix stiffness already enables cells of nearly any shape to generate substantial contractility (Figure 2C). Thus, the metabolic potential does not provide a strong driving force for elongation. For a clear visualization of how the variation in the volumetric part of the metabolic potential and interfacial energy produces the trends described here, we plot and compare their exact values in Figure S9.

In summary, only at intermediate collagen densities is the stress- and contractility-dependent reduction in the volumetric portion of the metabolic potential large enough to offset the interfacial energy penalty associated with elongation. This competition produces an energy landscape that leads to a biphasic dependence of the optimum cell aspect ratio on collagen density (Figure 2D). Indeed, we find that without interfacial energy, the shape that minimizes the volumetric portion of the metabolic potential at any density is always very highly elongated (Figure S10).

Metabolic and mechanical consequences of morphology selection

Despite the nonmonotonic dependence of cell shape on collagen density, both the steady-state volume-averaged contractility and the ECM strain energy of the optimal cell shape increase monotonically with collagen density because the amount of stress generated by the cell (and thus the amount of stress-related signaling the cell experiences) increases with ECM stiffness (Figures 2F, S11A). These predictions agree with quantitative polarization microscopy (QPOL) measurements, which report optical retardance due to F-actin strain and increase with collagen density (Figure 2F). The model also predicts a monotonic increase in ATP consumption rate at steady state (Equation S2.27), consistent with PercevalHR measurements of ATP depletion following inhibition of oxidative phosphorylation and glycolysis (*40*) (Figure 2G).

Finally, to further validate our framework, we simulated myosin inhibition by reducing the stress fiber assembly rate $k_{on,ij}$. This perturbation significantly reduced steady-state cytoskeletal stress, mirroring the effects of Y-27632 treatment in 3D collagen (Figure S12A,B) (*45*). Myosin inhibition also prevented cells from adopting elongated shapes in intermediate-density collagen (1.5 mg/ml), reflecting the fact that reduced contractility weakens the reduction in the volumetric portion of the metabolic potential, v^{cell}, normally associated with elongation.

Metabolic potential predicts cell flattening and elongation on stiffer 2D substrates

We next asked whether the metabolic potential framework could accurately predict single-cell morphologies in environments beyond 3D collagen, such as on two-dimensional polyacrylamide (PA) substrates commonly used to study mechanosensitivity as a function of matrix stiffness (*46*). To address this, we predicted MDA-MB-231 cell shape and contractility on 2D substrates of varying stiffness (Young's modulus) using *the same biophysical parameters* determined for cells in 3D collagen (Table S3), without introducing any additional fitting parameters. The only

modification is the interfacial energy term, $\Gamma(\Theta)$, which must account for the fact that cells on 2D substrates interact with the ECM only along their basal surface rather than being fully encapsulated (SI Section 2.2.5). In addition, because cells form larger and stronger adhesions on stiffer substrates (47), the receptor–ligand adhesion energy is scaled with substrate stiffness, rather than implemented as a function decreasing with matrix density as in 3D collagen.

Figure 3: Minimizing the metabolic potential predicts increasing cell elongation and flattening with 2D matrix stiffness. (**A**) Schematic depiction of the geometry of a cell on a 2D substrate. (**B**) Optimum cell shape shows a transition from hemispherical on soft gels to increasingly elongated and flattened on intermediately and very stiff PA gels as determined by the interaction between the volumetric part of the metabolic potential (red block arrows) and the interfacial energy (yellow block arrows). (**C**) Contour plots of the metabolic potential as a function of the cell aspect ratios a/b and c/b with the optimum cell shape indicated. Comparison of (**D**) cell spread area, (**E**) cell circularity, and (**F**) cell contractility at steady state predicted by the model with the corresponding data from experiments with MDA-MB-231 cells seeded on 2D substrates reported in (45,48). Bar plots show mean ±SD. Scale bar = 25 μm.

Cells on 2D substrates are modeled as half-ellipsoids with semiaxes a, b, and c (Figure 3A), where in general a ≠ b ≠ c. When all three semiaxes are equal, the cell adopts a hemispherical shape. To parameterize cell morphology, we define two aspect ratios: a/b, which captures elongation in the plane of the cell–matrix interface, and c/b, which characterizes the cell height relative to its in-plane

width (Figure 3A). Cell volume is held constant, so the spreading area ($A=\pi ab$) increases as the vertical axis, c, shortens. Thus, a decrease in c/b indicates cell flattening, while an increase in a/b indicates cell elongation. Although a continuous family of shapes was explored, we found that three representative morphologies capture the dominant energetic effects: (i) hemispherical (a/b=1, c/b=1), (ii) moderately elongated spindle-like (a/b>1,c/b<1), and (iii) highly elongated and flattened spindle-like (a/b≫1,c/b≪1) (Figure 3B). The PA substrate is treated as a linear elastic material (*46*) with stiffness varying between 1 and 200 kPa.

Anisotropic cell shapes on 2D substrates develop elevated and polarized cytoskeletal stresses

Cells on 2D substrates exhibit mechanosensitive trends similar to those observed in 3D collagen. As substrate stiffness increases, they develop larger volume-averaged stresses, which lead to higher steady-state contractility and increased ATP hydrolysis by active motors (Figure S13). Likewise, as the cell elongates within the 2D plane, both the stress field and the contractility field become increasingly polarized due to preferential stress fiber assembly along the direction of elongation (Figure S14A,B). The optimum cell shape on 2D substrates is still determined by the competition between the volumetric portion of the metabolic potential, v^{cell}, and the interfacial energy, $\Gamma(\Theta)$. Substrate stiffness modulates this competition by regulating the magnitude of stress the cell can generate at steady state, and thus the degree of mechanosensitive upregulation of stress fiber assembly.

However, the predicted dependence of optimal cell morphology on matrix stiffness differs substantially from the biphasic trend observed in 3D collagen. In 2D, cells *monotonically elongate and flatten* with increasing substrate stiffness (Figure 3B). This monotonic trend arises because the cell can only adhere to and transmit traction forces through its basal surface, meaning the region of highest stress is always located near the cell–matrix interface regardless of shape (Figure 3B). Flattening the cell increases the fraction of the cytoskeletal network that lies close to this high-stress region and prevents stress relaxation in regions farther from the interface. Consequently, flattened morphologies support larger stresses throughout the cell body. These larger stresses increase ATP hydrolysis and reduce the volumetric portion of the metabolic potential, v^{cell}, providing a strong energetic incentive for both flattening and elongation as stiffness increases.

Interfacial energy opposes both flattening and elongation because each increases cell surface area. However, as substrate stiffness rises, the cell's ability to develop larger contractility more effectively reduces v^{cell}, enabling it to overcome the increasing interfacial energy penalty. As a result, the competition between v^{cell} and $\Gamma(\Theta)$ produces a minimum in the metabolic potential that shifts systematically toward shapes with higher a/b (more elongated) and lower c/b (more flattened) as stiffness increases (Figure 3C). The resulting morphology ranges from a nearly hemispherical cell on soft substrates (< 1 kPa) to a highly elongated, spindle-like cell on stiff substrates (> 100 kPa). This predicted monotonic elongation and flattening is in excellent agreement with experimental measurements of MDA-MB-231 spread area and cell circularity on PA substrates (*48*) (Figure 3D,E). Consistent with the predictions from 3D collagen, the volume-averaged stress, contractility, and ATP hydrolysis all monotonically increase with substrate stiffness (Figure S14C–E). These predictions also match experimental assessments of cell contractility via optical retardance measurements (*45*) (Figure 3F).

In summary, we find cells do not exhibit a biphasic morphology trend in 2D because, unlike in 3D matrices, a flattened configuration always provides the most favorable means of maintaining high stresses on stiff substrates. Therefore, there is no energetic advantage to returning to a less flattened shape once stiffness becomes large. Interestingly, this driving force for a cell to adopt the shape

393 which maximizes contractility and minimizes metabolic potential can accurately predict the array
394 of morphologies observed when cells are placed in environments with different stiffness and
395 different geometry.

A Predictive Framework for Mechanosensitive ATP Replenishment in Microenvironments with Variable Matrix Stiffness

Thus far, we have shown that despite differences in optimum morphologies, cells in stiffer
environments in both 2D and 3D develop higher contractility and correspondingly exhibit greater
ATP consumption. We now extend our model to develop a theoretical framework describing how
cells regulate their energetic budget in response to the increased bioenergetic demands imposed by
stiff microenvironments. Our objective is to quantitatively predict key metabolic indicators—
including ATP levels, ATP:ADP ratios, glucose uptake, and oxidative phosphorylation (OXPHOS)
rates—as functions of ECM stiffness. We first provide a quantitative description of ATP
consumption in adherent, contractile cells and then outline the proposed mechanosensitive ATP
replenishment pathways, along with the model parameters used to quantify them. Full derivations
and parameter determination are provided in SI Section 3.

Living cells maintain a high, nonequilibrium ATP:ADP ratio, typically in the range of 1–10 (*49*).
To model ATP consumption, we consider the ATP hydrolysis reaction (reaction 1 in Table S4), in
which the rate constants k_1^+ and k_1^- govern ATP consumption and ATP synthesis, respectively.
Although cells may adjust many energy-intensive processes in different microenvironments, here
we focus on ATP consumption associated with changes in motor activity driven by stress fiber
assembly. We therefore take the ATP consumption rate constant k_1^+ to depend on the rates of stress
fiber assembly and disassembly, which themselves depend on cytoskeletal stress. Accordingly, k_1^+
increases with the total volumetric stress (see SI Section 3 for the functional form).

Once the finite element formulation is solved for a given cell shape on a 2D substrate or in 3D
collagen, we compute the total stress by integrating $\sigma_{kk}(\mathbf{x})$ over the cell volume. We also incorporate
the conversion of ADP to AMP (reaction 3 in Table S4) by introducing the ratio γ' of the reverse
and forward rate constants (k_2^+ and k_2^-). Increased cytoskeletal tension and ATP consumption on
stiff substrates lead to a greater reduction in ATP:ADP ratio and a corresponding rise in the
AMP:ATP ratio compared with soft substrates.

Mechanosensitive, calcium-dependent activation of AMPK drives ATP replenishment

A central regulator of cellular metabolism is AMP-activated protein kinase (AMPK), which
integrates signals related to cellular energy levels and controls major catabolic and anabolic
pathways, including OXPHOS and glycolysis (*17*). AMPK is a heterotrimeric complex composed
of the catalytic AMPKα subunit and regulatory AMPKβ and AMPKγ subunits. The activation of
AMPK is initiated by the allosteric binding of AMP to AMPKγ. As cellular energy levels fall, AMP
concentration increases, promoting higher levels of AMP-bound AMPK. This effect is amplified in
stiffer microenvironments, where greater contractility and ATP consumption elevate AMP levels
(Figure 4A).

AMP binding induces a conformational change that enhances phosphorylation of AMPKα by
upstream kinases. Two kinases—LKB1 (liver kinase B1) and CaMKKβ (Ca²⁺/calmodulin-
dependent protein kinase kinase β)—promote phosphorylation of AMPKα at Thr172. Although
LKB1 expression does not appear to depend on ECM stiffness, cytoplasmic calcium levels increase
on stiffer substrates (*50–52*), especially in mesenchymal cells where elevated RhoA signaling and

increased membrane tension both enhance calcium influx (*52,53*). Higher intracellular calcium leads to higher CaMKKβ activation and thus higher levels of phosphorylated (activated) AMPK, denoted here as AMPK*(Figure 4B).

Figure 4: Model predicts ATP replenishment based on the mechano-sensitive calcium-mediated activation of AMPK. (**A**) Schematic representation of the changing ATP, ADP and AMP levels immediately after a cell is seeded, before reaching steady state, and at steady state on soft and stiff ECMs. (**B**) Schematic depiction of AMPK activation in response to AMP and calcium levels. ECM stiffness regulates AMPK activation by affecting both cell energy consumption and calcium signaling as the cell evolves towards steady state. (**C**) Predicted ATP:ADP ratio vs. collagen density and (**D**) PA gel stiffness compared to experimental measurements with PercevalHR from (*16*). (**E**) Confocal max intensity projections of phosphorylated AMPK in MDA-MB-231 cells cultured on 10kPa 2D PA gel and glass. (**F**) Quantification of (**E**) compared with

model predictions of [AMPK*]. Model predictions of [AMPK*] are compared with change in rate
of glucose uptake for MDA MB-231 cells in (**G**) 3D collagen and (**H**) hMSCs on 2D gels from
references (*18,42*). (**I**) Predicted ATP levels compared with experimental measures in different cell
types from (*18*). (**J**) Predicted ATP:ADP levels in cells with normal and lowered stress fiber
assembly rates compared against experimental measurements for MDA MB-231 cells in 3D
collagen treated with contractility inhibitor ML7 from (*42*). Bar and line plots show mean ±SE, and
box plots show median, $25^{th}/75^{th}$ percentiles, and $5^{th}/95^{th}$ percentiles *p<0.05, **p<0.01,
***p<0.001, ****p<0.001 with Kruskal-Wallis one-way ANOVA (**C,D,F**), Tukey's HSD post-hoc
test (**G,H**) or Wilcoxon rank test (**J**). Scale bar = 50 μm.

AMPK activation is represented by reaction 4 in Table S4, where k_3^+ denotes the activation rate
constant and k_3^- the deactivation rate constant. To capture the mechanosensitive nature of AMPK
activation, we take k_3^+ to be a function of cytoskeletal stress (and thus intracellular calcium), i.e.,
$k_3^+ = k_3^+(\sigma_{kk})$. The ATP synthesis rate constant k_1^- is an increasing function of AMPK*, reflecting
AMPK-mediated upregulation of ATP production (reaction 2 in Table S4).

Mathematical expressions for steady-state concentrations of ATP, ADP, AMP, and AMPK* are
obtained by setting time derivatives to zero and enforcing conservation of total nucleotide
concentration and total AMPK. This yields a system of five nonlinear equations that we solve
simultaneously.

Cells sustain higher ATP:ADP ratios and AMPK activation on stiffer matrices

467
468
469   Simplifying our system of equations results in a predicted relationship between the ATP:ADP
470   ratio and total volumetric stress:
471

$$\frac{[\text{ATP}]}{[\text{ADP}]} = \frac{1.3\left(\dfrac{c}{\sigma_{kk}^n} + 1 - \sqrt{\left(\dfrac{c}{\sigma_{kk}^n} + 1\right)^2 - 1}\right)}{\sqrt{\left(\dfrac{c}{\sigma_{kk}^n} + 1\right)^2 - 1} - \dfrac{c}{\sigma_{kk}^n}} \qquad (2)$$

472
where c and n represent combinations of other model parameters that arise while solving (see SI
Section 3.1). Equation 2 predicts that the steady state ATP:ADP ratio will increase with
increasing stress (or increasing ECM stiffness) if both c and n are positive. Parameter c is always
positive since all its factors such as k_2^+, k_2^-, and the total AMPK concentration, [AMPKtot], are
all positive by definition. However, n describes the difference between the change in the rate
parameters k_1^+ and k_3^+ in response to a change in stress and will only be positive if k_3^+ (the rate of
calcium-dependent AMPK activation) shows a larger increase in response to stress than k_1^+ (the
rate of ATP consumption by motors). In other words, Equation 2 reaffirms that the ATP:ADP
ratio will increase with ECM stiffness only when cells experience sufficient AMPK activation and
subsequent ATP production to overcome the rate at which ATP is consumed by motors. We fitted
Equation 2 to experimentally measured ATP:ADP ratios for collagen densities ranging from 0.5
to 5 mg/mL (Figure S15A) to determine the value of parameters c and n (Table S5). Our
framework captures the trend that ATP:ADP ratios increase with collagen density (Figure 4C) and
accurately predicts that cells maintain steady-state nucleotide concentrations with
[ATP]>[ADP]>[AMP] (Figure S15B,C).

To validate the generality of the model, we predicted ATP:ADP ratios for cells cultured on 2D PA substrates using the same parameters. The predicted ATP:ADP ratio increases monotonically with substrate stiffness, in agreement with experimental observations (Figure 4D). Although cell morphology responds differently to stiffness in 3D collagen and 2D PA gels, the ATP:ADP ratio increases in both environments because the optimum configuration adopted by cells on stiffer substrates supports higher stress and contractility (Figures 2F and 3F), leading to greater ATP consumption and a corresponding increase in ATP replenishment.

Our framework also predicts that AMPK activation levels increase as a function of matrix stiffness (Equation S3.18). Measurements of phosphorylated AMPK (Thr172) from immunostained cells cultured on soft versus stiff substrates show significant increases in AMPK activation on stiffer substrates, in good agreement with our predictions (Figure 4F). These findings are consistent with reports showing that AMPK activation increases with stiffness on both short and long timescales (*18*). We find that activated AMPK localizes strongly to the nucleus (Figure 4E), consistent with other reports of nuclear accumulation of AMPK under conditions of metabolic stress (*54,55*).

Fig. 5. Model predicts decreases in intracellular energy levels and mitochondria membrane potential observed in cells on confining micropatterns. (**A**) Normalized [ATP]:[ADP] ratio as measured with PercevalHR and (**B**) mitochondria membrane potential as measured with tetramethylrhodamine methyl ester (TMRM) in MDA-MB-231 cells cultured on collagen micropatterns of fixed area fabricated on PA gels of 10 kPa or 30 kPa along with model predictions. (**C**) Normalized ATP:ADP ratio measured in cells on micropatterns of fixed elongation (elongation ~1) compared to model predictions. Plots show mean ±SD. Scale bar = 50 μm. Experimental data and images obtained from (*16*).

Model predictions of increased AMPK activation in stiff microenvironments correlate with experimentally observed increases in glucose uptake (Figure 4G,H), elevated mitochondrial membrane potential (a proxy for OXPHOS rate) (Figure S16), and increased ATP levels across multiple cell types (Figure 4I) (*16,18,42*). Increased ATP levels with stiffness are a general feature of mesenchymal cells, and the model captures this trend well. To assess whether changes in ATP:ADP ratio indeed reflect changes in motor activity, we simulated myosin inhibition by reducing the stress fiber assembly rate $k_{on,ij}$ over 200-fold. Motor inhibition reduced steady-state cytoskeletal stress (Figure S12B), AMPK activation, and ATP replenishment, producing significantly lower ATP:ADP ratios consistent with experimental results using myosin light chain

kinase inhibitors such as ML7 (*42*) (Figure 4J). These findings support the idea that cytoskeletal
contractility is a major determinant of energy budget regulation across stiffness environments.

In summary, we find a mathematical analysis accounting for the mechanosensitive chemical
reactions governing the generation and interconversion of ATP, ADP, and AMP is sufficient to
explain experimental measurements of intracellular ATP:ADP ratio and AMPK activation in
contractile cells in different stiffness environments. On stiffer substrates, motor ATP consumption
and calcium mechanosignaling work in parallel to upregulate AMPK activation and ensure cellular
energy production meets energy demand.

Restricted spreading reduces cytoskeletal stress and metabolic activity

Previous work has shown that constraining cells to micropatterned substrates that limit their ability
to spread or elongate alters their metabolic state, including the ATP:ADP ratio (*16*). Using our
model, we examined how cytoskeletal stress and ATP consumption are modulated in cells confined
to restricted geometries. For cells adhered to PA substrates of 10 or 30 kPa, we considered two
cases: 1) fixed spread area with varying elongation, and 2) fixed elongation with varying spread
area. For each shape, we solved for the steady-state stress field and computed the volume-averaged
cytoskeletal stress ($1/V_c \int \sigma_{kk} dV_c$) followed by the predicted ATP:ADP ratio using Equation 2 and
the biophysical parameters in Table S5.

Our model predicts that ATP:ADP ratios increase with cell elongation when spread area is held
constant, consistent with experimental measurements (Figure 5A). We obtain similar results when
modeling cells on patterns of increasing spread area at a constant elongation (Figure 5C). Confining
patterns that restrict elongation or spreading force the cell into more hemispherical shapes, which
exhibit greater stress relaxation and lower volume-averaged cytoskeletal stress, ultimately reducing
contractility. Cells with lower contractility consume less ATP and consequently exhibit reduced
AMPK activation and ATP replenishment at steady state. For a given elongation, ATP:ADP ratios
are higher on stiffer substrates, reflecting the enhanced stress-supported contractility (Figure 5A).
Model predictions of AMPK activation also agree closely with observed mitochondrial membrane
potential changes across micropattern geometries that limit elongation (Figure 5B). When spread
area increases at fixed elongation, ATP:ADP ratios again increase, matching experimental findings,
as does mitochondria membrane potential (Figure S17).

Together, these results further validate the proposed mechanosensitive ATP replenishment model
in contexts where cell spreading is externally controlled and where long-term cytoskeletal
remodeling dominates cellular energy expenditure.

Figure 6. A model to predict how cell shape, contractility, and metabolism depend on mechanosensitive stress fiber assembly in adherent cells. Cell shape and ECM stiffness determine the distribution and magnitude of stresses generated in an adherent cell which, through mechanosensitive pathways, regulate the establishment of stress fibers. The decrease in energy achieved by coupling the ATP hydrolysis with stress fiber assembly drives the cell towards a more contractile state and to adopt morphologies that maximize ATP consumption and minimize interfacial energy. Simultaneously, the same mechanosensitive pathways upregulate cytosolic calcium levels which leads to the activation of AMPK through a calmodulin mediated pathway, replenishing cellular ATP stores.

DISCUSSION

In this study, we developed a nonequilibrium chemo-mechanical model for adherent cells that quantitatively connects mechanosensitive signaling to the regulation of cellular energy budgets and morphological adaptations in response to microenvironment stiffness (Figure 6). Central to this framework is the concept of a metabolic potential, defined as the sum of energies associated with establishing contractility, including myosin binding energy, motor work, cytoskeletal and matrix strain energies, energy released from ATP hydrolysis, and the cell–matrix interfacial energy. We demonstrate that minimizing this metabolic potential yields contractility and strain fields identical to those derived from the kinetics of mechanosensitive stress fiber assembly, consistent with nonequilibrium thermodynamics.

Using this formulation, we predicted the steady-state cell shape and contractility of MDA-MB-231 cells in two distinct contexts: embedded within 3D collagen matrices and adhered to 2D polyacrylamide substrates. Across both environments, elongated cells exhibit increased contractility and stress-fiber polarization, which lower the metabolic potential, while elongation simultaneously increases interfacial energy, creating a competition that determines the cell's optimal morphology. The model predicts a biphasic dependence of cell elongation on collagen density in 3D matrices and a monotonic elongation and flattening with increasing stiffness on 2D substrates, matching experimental observations. These results highlight the central role of geometry and stress distribution in determining cell morphology under different mechanical constraints.

Our predictions agree with experimentally observed cell shapes in different collagen densities and PA stiffnesses (16), and they are consistent with behaviors reported for fibroblasts and other mesenchymal cells (56). Previous reports of increased collagen fiber alignment with cell contraction in dense matrices support our finding that cells generate larger contractile forces in stiffer microenvironments (57,58). Very high-density matrices can suppress traction when they restrict matrix degradation, as observed for hMSCs in covalently crosslinked hydrogels (59), but this effect is unlikely to influence MDA-MB-231 cells at the range of densities examined here. Our model also predicts matrix strain energies that are consistent with reported experimental measurements (60,61). Differences in strain energy between nonlinear collagen gels and linear PA gels are captured by the model's incorporation of tension-driven fiber alignment, which enables long-range force transmission (43).

We further introduced a theoretical description of ATP replenishment mediated by mechanosensitive activation of AMPK. Because ATP hydrolysis fuels contractility, cells increase metabolic output to meet the demands imposed by stiffer microenvironments. The model predicts increases in ATP:ADP ratio, glucose uptake, mitochondrial potential, and AMPK activation with ECM stiffness, consistent with experimental data in both 2D and 3D environments (16,42). The reduction in ATP:ADP ratio observed upon myosin inhibition provides additional support for the direct coupling between cytoskeletal tension, AMPK activation, and cellular energy homeostasis. Predictions for cells on micropatterns further demonstrate how reduced spreading or elongation lowers cytoskeletal stress, metabolic activity, and AMPK signaling.

An important aspect of our framework is that all ATP consumed at steady state is dissipated as heat, since there is no net addition of stress fibers and no net change in conservative energy. The heat dissipation rate therefore represents the maximum mechanical power output of the cell. Using experimentally constrained parameters (Table S6), we estimate that a cell in 0.5 mg/mL collagen generates approximately 0.81 fW of power. Since 1 fW corresponds to the hydrolysis of roughly 20,000 ATP molecules per second (62), this power output implies that such a cell consumes approximately 16,000 ATP molecules per second to maintain its steady-state configuration. On soft 1 kPa PA gels, the predicted power output is around 0.38 fW. In dense collagen (5 mg/mL), this value increases to approximately 6.5 fW, corresponding to more than 100,000 ATP molecules per second. These predictions are comparable to reported single-cell power measurements and illustrate the substantial energetic investment required to sustain high contractility in stiff environments (35).

The results of this study suggest several natural extensions of the framework. The metabolic potential formulation can be generalized to describe dynamic behaviors such as cell spreading, migration, durotaxis, and mechanically induced transitions in contractile state. Such processes involve time-dependent changes in stress fiber organization, signaling, and AMPK activation, and incorporating these dynamics could provide deeper insight into transient and adaptive mechanosensitive responses. The model could also be adapted to account for cell-type-specific

mechanosensitive programs, enabling comparisons across different cancer cell lines, stem cells, or
immune cells that experience distinct mechanical and metabolic constraints. Differences in motor
kinetics, adhesion dynamics, or metabolic capacity could be incorporated to explore how distinct
populations balance mechanical demand with energetic supply.

More broadly, the framework provides a foundation for linking single-cell energetics to tissue-level
mechanical environments. Mechanical heterogeneity within tumors, fibrotic tissues, or regenerating
matrices may impose spatial patterns of energetic demand that govern collective behavior,
metabolic adaptation, and invasiveness. Understanding these relationships may help identify
mechanically regulated vulnerabilities in cancer cells or inform the design of engineered
microenvironments that direct stem cell function or modulate immune activation.

Altogether, by establishing a quantitative link between mechanical signaling, cytoskeletal
organization, and energetic regulation, this model offers a predictive and extensible approach for
understanding how cells sense, respond to, and adapt to the mechanical properties of their
surroundings. These insights provide a basis for future explorations of mechanosensitive metabolic
regulation across diverse physiological and pathological settings

METHODS

A 3D chemo-mechanical formulation of the metabolic potential

Here, we briefly outline (refer to SI Section 2 for details) a general 3D model for the metabolic
potential of a contractile cell to describe the effect of stress-activated signaling pathways on the
metabolic budget. The key mechanical components of this model are a) stress fibers composed of
active contractile elements in parallel with passive components representing the cell cytoskeleton,
b) a poroelastic nucleus, and c) the extracellular matrix or substrate.

A) Stress fibers: In the 3D model, contractility, stress, and strain are treated as tensor fields,
reflecting the cell's ability to assume nonspherical shapes with spatially nonuniform anisotropic
stresses and locally varying motor orientations (Figure S4A). In this case, the contractility field
(motor density associated with stress fibers) is anisotropic, i.e., $\rho_{11} \neq \rho_{22} \neq \rho_{33}$, as is the stress field,
and both can vary spatially across the cell body necessitating a dependence on spatial coordinates
represented by vector $\mathbf{x}$. The volumetric part of the contractility tensor $\rho_{kk}(\mathbf{x}) =$
$(\rho_{11}(\mathbf{x})+\rho_{22}(\mathbf{x})+\rho_{33}(\mathbf{x}))/3$ represents contraction averaged over all directions and hence is a measure
of the density of motors per unit volume in the cell. The deviatoric part, $\tilde{\rho}_{ij}(\mathbf{x}) = \rho_{ij}(\mathbf{x}) - 1/3\rho_{kk}(\mathbf{x})\delta_{ij}$,
is a measure of the deviation of each component of ρ_{ij} from the mean and represents the degree of
motor polarization and, by extension, stress fiber polarization.

The cell experiences activation of mechanosensitive pathways dependent on the strength of the
stress generated, σ_{ij}. These pathways regulate myosin motor activity and associated ATP
consumption by affecting the stress-dependent rates of stress fiber assembly ($k_{on,ij}(\sigma_{ij})$) and
disassembly ($k_{off,ij}(\sigma_{ij})$). $k_{on,ij}$ and $k_{off,ij}$ are tensorial quantities which allow for anisotropy in stress
fiber assembly and disassembly governed by anisotropy in the stress field. The exact functional
dependence of the components of $k_{on,ij}$ and $k_{off,ij}$ on the components of σ_{ij} is informed by the
biophysical mechanisms that underlie motor activation and stress fiber formation. For example, we
assume that stress fiber kinetics are linearly proportional to the average stress at every point such
that the volumetric parts of the on and off rates (k^v_{on} and k^v_{off}) are proportional to the volumetric
stress (i.e., $k^v_{on}(\sigma_{kk}(\mathbf{x})) = (k_{on,11}(\sigma_{kk}(\mathbf{x})) + k_{on,22}(\sigma_{kk}(\mathbf{x})) + k_{on,33}(\sigma_{kk}(\mathbf{x})))/3$) while the deviatoric parts
$(\tilde{k}_{on,ij}(\tilde{\sigma}_{ij}(\mathbf{x}))$ and $\tilde{k}_{off,ij}(\tilde{\sigma}_{ij}(\mathbf{x})))$ allow for upregulated stress fiber formation rates in the direction of

larger stresses. This captures the fact that mechanosignaling and an increase in the total number of active motors may arise from an increase in stress along any direction, but the cell will experience a greater rate of stress fiber formation in the direction of higher stress. However, we have previously shown that contractility regulated solely by volumetric (average) stress is insufficient to explain experimentally observed increases in cell contractility in elongated cell morphologies characterized by anisotropic stress distributions (*39*). This discrepancy arises from the formation of anisotropic adhesions aligned with polarized stress fibers, which establish bidirectional mechano-chemical feedback between cell shape and signaling (*63*). To model this process, we introduce a non-linear term to the rate of stress fiber assembly, $k^a_{on,ij}$, that accounts for increases in cell contractility but only when the stress field is anisotropic. This is readily accomplished by expressing tensor quantities in terms of their principal components to identify stress anisotropy independent of any coordinate system. The principal components of stress, $\sigma_i(\mathbf{x})$ where $\sigma_1(\mathbf{x}) \geq \sigma_2(\mathbf{x}) \geq 0$, reveal stress anisotropy exists whenever the first principal component, $\sigma_1(\mathbf{x})$, is larger than the second principal component, $\sigma_2(\mathbf{x})$. We introduce this mathematically by assigning $k^a_{on,i}$ a functional dependence on the term $(\sigma_1(\mathbf{x})/\sigma_2(\mathbf{x}) -1)$ such that $k^a_{on,i}$ is nonzero and leads to increased stress fiber assembly only when $\sigma_1(\mathbf{x}) > \sigma_2(\mathbf{x})$, or when the stress fields are sufficiently anisotropic (see SI Section 2 for a full description).

Using these quantities, we define the conservative energy densities at every point in the cell, including the myosin binding energy (Equation S2.8), motor work (Equation S2.9), strain energy of the passive components (Equation S2.10), and the energy from ATP hydrolysis (Equation S2.14) (see SI Section 2.2 for a full tensorial description of these energies). Physically, this formulation captures the fact that ATP hydrolysis is energetically downhill and can therefore drive otherwise unfavorable processes, such as myosin binding and the straining of the cytoskeleton and extracellular matrix. However, because the release of this energy is directly coupled to the mechanosensitive activity of myosin motors within stress fibers, the kinetics of stress-dependent and stress-anisotropy-dependent signaling pathways regulate the rate at which ATP energy can be utilized. We incorporate this effect by defining the stress fiber assembly rates, $k_{on,i}$ and $k^a_{on,i}$, as increasing functions of the local principal stress, σ_i, and the disassembly rate, $k_{off,i}$, as a decreasing function of stress. Consequently, when the cell experiences high levels of cytoskeletal tension - such as on stiff extracellular matrices - the energy available from ATP consumption increases due to both an elevated rate of myosin motor engagement and an increased motor binding lifetime.

B) Adhesions at the cell-matrix interface: Transmembrane proteins at the cell-ECM interface, such as integrins, form a bond with ligands on the ECM and play an important role in transmitting forces to the cell cytoskeleton. To capture the cellular morphology in 2D and 3D microenvironments, we also introduce an interfacial energy, Γ, which is a function of the cell shape. There are two contributions to the interfacial energy: the cell surface tension, γ_0 which resists any increase in the cell area or elongation, and an adhesive contribution from membrane protein–ligand binding interactions, γ_1, which favors an increase in the cell surface area along the cell-matrix interface (Figure 2A). Cell membranes consist of lipid bilayers tethered to an underlying rigid actin–myosin cortex. Hence, any change in the cell membrane area is accompanied by unfavorable stretching of membrane-engaged proteins that resist expansion in the cell area as well as by sustained contraction of the underlying actomyosin cortex. We consider these effects to comprise the energetic cost of stretching the cell membrane. The adhesive element represents the reduction in energy achieved due to bond formation between receptors such as integrin at the cell-matrix interface and ligands present in the matrix. The relative strengths of these two quantities determine whether the total interfacial energy $\Gamma = (\gamma_0 - \gamma_1)S$ will increase or decrease with cell surface area, S. Additionally, we account for changes in the adhesive energy with changing ECM ligand density and changing membrane protein expression with ECM stiffness, as explained further in SI Section 2.2.5.

C) The nucleus: We assume that the viscous relaxation of the nucleus is negligible (see SI Section 2.1.2) and model the nucleus as a linear elastic ellipsoidal inclusion encapsulated within the cell with elastic properties K_{nuc} and μ_{nuc} denoting its bulk and shear modulus, respectively. We assign the nucleus a volume 1/5 of the cell volume and vary its aspect ratios in the same manner as cell aspect ratios when considering different shape cells.

D) Mechanical response of the microenvironment: We consider two types of cell microenvironments in this work: 1) cells encapsulated in 3D collagen and 2) cells seeded on 2D PA gels. Following our previous work, the mechanical behavior of collagen is described using a fibrous nonlinear constitutive model that captures the strain-stiffening effect of fiber alignment under tension (*43*) (see SI Section 2.2.6). The bulk and shear moduli of collagen as a function of its density were determined from experimental measurements of the equilibrium modulus of collagen gels (*16*). 2D PA gel substrates are modeled as linear elastic, as AFM measurements have shown that gels with stiffness values of physio-pathological relevance exhibit ideal linear elastic behavior even at the nanometer scale (*46*).

Generalized 3D expression for the metabolic potential of an adherent cell

We define the local metabolic potential, R_l, at every point $\mathbf{x}$ in the cell in terms of principal components by summing all energetic contributions:

$$R_l\big(\rho_i(\mathbf{x}), \varepsilon_i(\mathbf{x}), \Theta\big)$$

$$= \underbrace{F_c\big(\rho_i(\mathbf{x}), \varepsilon_i(\mathbf{x}), \Theta\big)}_{\substack{\text{Conservative contributions including motor work,}\\ \text{myosin binding energy, and cytoskeleton strain energy}}}$$

$$+ -\sum_i \underbrace{\frac{k_{on,i}\big(\sigma_i(\mathbf{x})\big) + k_{on,i}^a\left(\frac{\sigma_1(\mathbf{x})}{\sigma_2(\mathbf{x})} - 1\right)}{k_{off,i}\big(\sigma_i(\mathbf{x})\big) + k_{on,i}\big(\sigma_i(\mathbf{x})\big) + k_{on,i}^a\left(\frac{\sigma_1(\mathbf{x})}{\sigma_2(\mathbf{x})} - 1\right)} \frac{\Delta G}{p_0}\rho_i(\mathbf{x})}_{\text{ATP hydrolysis energy due to motor recruitment}} \quad (3)$$

The total metabolic potential, R, is then calculated by integrating R_l over the cell volume and adding the strain energy from the nucleus and ECM as well as the interfacial energy, $\Gamma(\Theta)$ (Equation 1). The metabolic potential represents the potential of a cell to enhance its contractility and alter its morphology/shape in response to the mechanical properties of its microenvironment by considering mechanosensitive cytoskeletal kinetics. Minimization of the metabolic potential with respect to the contractility, strain, and shape yields the rate of ATP consumption in the cell at steady state (Equation S2.25).

To determine the optimum cell shape and contractility as a function of matrix stiffness, we employed a finite element procedure, as described in SI Section 2.3 (Figure S4C). Briefly, we construct cell geometries with varying cell body aspect ratios and discretize the geometry within a finite element framework. At every spatial coordinate $\mathbf{x}$, we determine the steady-state cellular contractility and strain by minimizing the local metabolic potential R_l with respect to $\rho_{ij}(\mathbf{x})$ and $\epsilon_{ij}(\mathbf{x})$ through calculation of functional derivatives of R with respect to these variables. This yields a set of equations that we solve using simulations for a given cell body aspect ratio (Equations S2.20-S2.21). Then, knowing the spatial variation of the contractility and strain, we determine the cell metabolic potential, R (Equation 1), corresponding to a particular cell body aspect ratio by integrating the local metabolic potential, R_l, over the cell and matrix volume and adding the

748 interfacial energy corresponding to that shape. This process is repeated for different cell body aspect
749 ratios, and the optimum cell shape is identified as the cell body aspect ratio that has minimum
750 metabolic potential. We repeated this procedure for different matrix stiffness values to generate the
751 observed variations in the optimum cell shape and contractility with stiffness.

Experimental measurement of AMPK activation

To compare with model results, we performed immunostaining and confocal imaging of cells
cultured on 2D polyacrylamide substrates to determine activation of AMPK at the single-cell level
in response to changing microenvironment stiffness. MDA-MB-231 cells (ATCC HTB-26) were
cultured in growth media consisting of DMEM supplemented with glutamax, sodium pyruvate,
Anti-Anti (Gibco), and 10% FBS (Cytiva) before trypsinization and resuspension for growth on
substrates. Polyacrylamide substrates were fabricated on pre-treated glass coverslips by mixing
different volumes of acrylamide, bis-acrylamide, tetramethylethylenediamine (TEMED), and
ammonium persulfate (APS) as previously described (*56,64*). Final percentages by volume were
7.7%:0.26%:0.27%:0.9% (10kPa) in dH_2O. Coverslips were pretreated to promote polyacrylamide
binding by incubation in 0.1M NaOH for 3 minutes, 0.5% (3-Aminopropyl)trimethoxysilane (3-
APTMS, Sigma) for 30 minutes, and 0.5% glutaraldehyde (Sigma) overnight. All incubations were
performed at room temperature and preceded by multiple washes with dH_2O. Coverslips were dried,
and the acrylamide mixture was polymerized for 30 minutes at room temperature while sandwiched
between the coverslips and siliconized glass, the latter of which could be easily removed with gentle
force afterward. Resulting PA gels were surface coated with collagen to promote cell adherence.
This was accomplished by incubation in 0.05% sulfo-sanpah (Sigma) in PBS during treatment with
UV light for 10 minutes. Gels were then washed multiple times with PBS and incubated in 0.1
mg/ml rat tail type I collagen (Corning) for 2 hours at room temperature. Gels were again washed
multiple times with PBS and sterilized under UV for 1 hour. After at least 24 hours, cells were
seeded in growth media at 25% confluency. For the glass stiffness condition, coverslips simply
underwent the final incubation step in 0.1 mg/ml collagen before sterilization and cell seeding.

Samples were cultured for 24 hours prior to 15 minute fixation with 4% paraformaldehyde,
permeabilization with 0.5% triton X-100 (Sigma) for 20 minutes, and blocking with 3% bovine
serum albumin (Sigma) for 1 hour with repeated washing between each step. Samples were then
stained overnight at 4°C with phosphorylated AMPK primary antibody (Cell Signaling Technology
#2531) and 1 hour at room temperature with secondary antibodies (Abcam 150077) prior to
confocal imaging. For each cell, a bright field image as well as a confocal z-stack at 235 x 235 μm
(512 x 512 pixels) and 0.5 μm vertical spacing was obtained using a Zeiss LSM980 point scanning
confocal microscope equipped with a 40x water immersion lens and a Prime BSI Express sCMOS
camera. Samples were imaged by inverting the gels into a No. 0 35 mm glass-bottom dish (MatTek)
containing Fluoromount-G mounting media (Invitrogen) to allow imaging of the cells through the
glass bottom of the dish. Relative levels of phosphorylated AMPK were quantified for each cell by
finding the total fluorescence signal using ImageJ software (*65*). Confocal images were z-projected
for max intensity and the extent of each cell body was manually determined by using reference
bright field images. Total expression for each cell (Figure 4F) was then calculated by normalizing
the sum of all pixel values to cell area and the mean expression of all cells on glass. Full details of
how model predictions were compared with previously published experimental data are available
in SI Section 3.2.

Statistical Analysis

796 All box and whisker plots show median (center line) $\pm$ interquartile ranges (box edges) and 5^{th} and
797 95^{th} percentiles (whiskers). Bar plots show mean and standard error or mean and standard deviation

where indicated. Statistical analysis was performed with MATLAB version 2023b (*66*) and significance was found using Kruskal-Wallis ANOVA (*67*) with Dunn's post hoc analysis (*68*). Experiments were reproduced independently at least three times. Nonlinear least squares regression was performed using MATLAB's lsqcurvefit algorithm.

REFERENCES

1. D.E. Discher, P. Janmey, Y. Wang. Tissue cells feel and respond to the stiffness of their Ssbstrate. *Science* **310**, 1139–1143 (2005).
2. M. Nagayama, H. Haga, M. Takahashi, T. Saitoh, K. Kawabata, Contribution of cellular contractility to spatial and temporal variations in cellular stiffness. *Exp Cell Res* **300**, 396–405 (2004).
3. N. Jain, K.V. Iyer, A. Kumar, G.V. Shivashankar. Cell geometric constraints induce modular gene-expression patterns via redistribution of HDAC3 regulated by actomyosin contractility. *Proc Natl Acad Sci U.S.A*. **110**, 11349–11354 (2013).
4. S. Sonam, S.R. Sathe, E.K.F. Yim, M.P. Sheetz, C.T. Lim. Cell contractility arising from topography and shear flow determines human mesenchymal stem cell fate. *Sci Rep* **6**, 1–12 (2016).
5. M.A. Griffin, S. Sen, H.L. Sweeney, D.E. Discher. Adhesion-contractile balance in myocyte differentiation. *J Cell Sci* **117**, 5855–5863 (2004).
6. J.S. Park, C.J. Burckhardt, R. Lazcano, L.M. Solis, T. Isogai, L. Li, C.S. Chen, B. Gao, J.D. Minna, R. Bachoo, R.J. DeBerardinis, G. Danuser. Mechanical regulation of glycolysis via cytoskeleton architecture. *Nature* **578**, 621–626 (2020).
7. T. Kusakabe, K. Motoki, K. Hori. Mode of Interactions of Human Aldolase Isozymes with Cytoskeletons. *Arch Biochem Biophys* **344**, 184–193 (1997).
8. J. Wang, A.J. Morris, D.R. Tolan, L. Pagliaro. The molecular nature of the F-actin binding activity of aldolase revealed with site-directed mutants. *J Biol Chem* **271**, 6861–6865 (1996).
9. H. Hu, A. Juvekar, C.A. Lyssiotis, E.C. Lien, J.G. Albeck, D. Oh, G. Varma, Y.P. Hung, S. Ullas, J. Lauring, P. Seth. M.R. Lundquist, D.R. Tolan, A.K. Grant, D.J. Needleman, J.M. Asara, L.C. Cantley, G.M. Wulf. Phosphoinositide 3-Kinase Regulates Glycolysis through Mobilization of Aldolase from the Actin Cytoskeleton. *Cell* **164**, 433–446 (2016).
10. P. Romani, I. Brian, G. Santinon, A. Pocaterra, M. Audano, S. Pedretti, S. Mathieu, M. Forcato, S. Bicciato, J.B. Manneville, N. Mitro, S. Dupont. Extracellular matrix mechanical cues regulate lipid metabolism through Lipin-1 and SREBP. *Nat Cell Biol* **21**, 338–347 (2019).
11. R. Bertolio, F. Napoletano, M. Mano, S. Maurer-Stroh, M. Fantuz, A. Zannini, S. Bicciato, G. Sorrentino, G, Del Sal. Sterol regulatory element binding protein 1 couples mechanical cues and lipid metabolism. *Nat Commun* **10**, 1–11 (2019).
12. S.C.J. Helle, Q. Feng, M.J. Aebersold, L. Hirt, R.R. Grüter, A. Vahid, A. Sirianni, S. Mostowy, J.G. Snedeker, A. Šarić, T. Idema, T. Zambelli, B. Kornmann. Mechanical force induces mitochondrial fission. *Elife* **6**, 1–26 (2017).
13. K.M. Tharp, R. Higuchi-Sanabria, G.A. Timblin, B. Ford, C. Garzon-Coral, C. Schneider, J.M. Muncie, C. Stashko, J.R. Daniele, A.S. Moore, P.A. Frankino. S. Homentcovschi, S.S. Manoli, H. Shao, A.L. Richards, K. Chen, J. Hoeve, G.M. Ku, M. Hellerstein, D.K. Nomura, K. Saijo, J. Gestwicki, A.R. Dunn, N.J. Krogan, D.L. Swaney, A. Dillin, V.M. Weaver. Adhesion-mediated mechanosignaling forces mitohormesis. *Cell Metab* **33**, 1322-1341.e13 (2021).
14. J.H. Koo, K.L. Guan. Interplay between YAP/TAZ and Metabolism. *Cell Metab* **28**, 196–206 (2018).

15. Q. Liu, Q. Luo, B. Deng, Y. Ju, G. Song. Stiffer matrix accelerates migration of
hepatocellular carcinoma cells through enhanced aerobic glycolysis via the MAPK-YAP
signaling. *Cancers* **12**, 490–505 (2020).

16. Y. Wu, M.R. Zanotelli, J. Zhang, C.A. Reinhart-King. Matrix-driven changes in metabolism
support cytoskeletal activity to promote cell migration. *Biophys J* **120**, 1705–1717 (2021).

17. D.G. Hardie. AMP-activated protein kinase-an energy sensor that regulates all aspects of
cell function. *Genes and Development* **25**, 1895–1908 (2011).

18. J. Xie, M. Bao, X. Hu, W.J.H. Koopman, W.T.S. Huck. Energy expenditure during cell
spreading influences the cellular response to matrix stiffness. *Biomaterials* **267**, 120494
(2021).

19. M.J. Tan, J.M. Ye, N. Turner, C. Hohnen-Behrens, C.Q. Ke, C.P. Tang, T. Chen, H.C.
Weiss, E.R. Gesing, A. Rowland, D.E. James. Y. Ye. Antidiabetic activities of triterpenoids
isolated from bitter melon associated with activation of the AMPK pathway. *Chem Biol* **15**,
263–273 (2008).

20. R.B. Ceddia, R. Somwar, A. Maida, X. Fang, G. Bikopoulos, G. Sweeney. Globular
adiponectin increases GLUT4 translocation and glucose uptake but reduces glycogen
synthesis in rat skeletal muscle cells. *Diabetologia* **48**, 132–139 (2005).

21. A.S. Marsin, L. Bertrand, M.H. Rider, J. Deprez, C. Beauloye, M.F. Vincent, G. Van den
Berghe, D. Carling, L. Hue. Phosphorylation and activation of heart PFK-2 by AMPK has
a role in the stimulation of glycolysis during ischaemia. *Curr Biol* **10**, 1247–1255 (2000).

22. A.S. Marsin, C. Bouzin, L. Bertrand, L. Hue. The stimulation of glycolysis by hypoxia in
activated monocytes is mediated by AMP-activated protein kinase and inducible 6-
phosphofructo-2-kinase. *J Biol Chem* **277**, 30778–30783 (2002).

23. H.M. O'Neill, G.P. Holloway, G.R. Steinberg. AMPK regulation of fatty acid metabolism
and mitochondrial biogenesis: Implications for obesity. *Mol Cell Endocrinol* **366**, 135–151
(2013).

24. S. Horman, G.J. Browne, U. Krause, J.V. Patel, D. Vertommen, L. Bertrand, A. Lavoinne,
L. Hue, C.G. Proud, M.H. Rider. Activation of AMP-activated protein kinase leads to the
phosphorylation of elongation factor 2 and an inhibition of protein synthesis. *Curr Biol* **12**,
1419–1423 (2002).

25. J. Kim, M. Kundu, B. Viollet, K.L. Guan. AMPK and mTOR regulate autophagy through
direct phosphorylation of Ulk1. *Nat Cell Biol* **13**, 132–141 (2011).

26. J.A. Eble, S. Niland. The extracellular matrix in tumor progression and metastasis. *Clin Exp
Metastasis* **36**, 171–198 (2019).

27. F. Spill, D.S. Reynolds, R.D. Kamm, M.H. Zaman. Impact of the physical
microenvironment on tumor progression and metastasis. *Curr Opin Biotechnol* **40**, 41–48
(2016).

28. D. Hanahan, R.A. Weinberg. Hallmarks of Cancer: The Next Generation. *Cell* **144**, 646–
674 (2011).

29. M.A. Conti, R.S. Adelstein. Nonmuscle myosin II moves in new directions. *J Cell Sci* **121**,
11–18 (2008).

30. M. Vicente-Manzanares, X. Ma, R.S. Adelstein, A.R. Horwitz. Non-muscle myosin II takes
centre stage in cell adhesion and migration. *Nat Rev Mol Cell Biol* **10**, 778–790 (2009).

31. S.T. Sit, E. Manser. Rho GTPases and their role in organizing the actin cytoskeleton. *J Cell
Sci* **124**, 679–683 (2011).

32. K. Katoh, Y. Kano, Y. Noda. Rho-associated kinase-dependent contraction of stress fibres
and the organization of focal adhesions. *Journal of the Royal Society Interface* **8**, 305–311
(2011).

33. B.D. Matthews, C.K. Thodeti, J.D. Tytell, A. Mammoto, D.R. Overby, D.E. Ingber. Ultra-rapid activation of TRPV4 ion channels by mechanical forces applied to cell surface β1 integrins. *Integr Biol* **2**, 435–442 (2010).

34. V.B. Shenoy, H. Wang, X.A. Wang. A chemo-mechanical free-energy-based approach to model durotaxis and extracellular stiffness-dependent contraction and polarization of cells. *Interface Focus* **6**, 20150067 (2016).

35. D. Mitrossilis, J. Fouchard, A. Guiroy, N. Desprat, N. Rodriguez, B. Fabry, A. Asnacios. Single-cell response to stiffness exhibits muscle-like behavior. *Proc Natl Acad Sci U.S.A.* **106**, 18243–18248 (2009).

36. M. Kovács, K. Thirumurugan, P.J. Knight, J.R. Sellers. Load-dependent mechanism of nonmuscle myosin 2. *Proc Natl Acad Sci U.S.A.* **104**, 9994–9999 (2007).

37. E.S. Schiffhauer, T. Luo, K. Mohan, V. Srivastava, X. Qian, E.R. Griffis, P.A. Iglesias, D.N. Robinson. Mechanoaccumulative Elements of the Mammalian Actin Cytoskeleton. *Curr Biol* **26**, 1473–1479 (2016).

38. E.S. Schiffhauer, Y. Ren, V.A. Iglesias, P. Kothari, P.A. Iglesias, D.N. Robinson. Myosin IIB assembly state determines its mechanosensitive dynamics. *J Cell Biol* **218**, 895–908 (2019).

39. F. Alisafaei, D.S. Jokhun, G.V. Shivashankar, V.B. Shenoy. Regulation of nuclear architecture, mechanics, and nucleocytoplasmic shuttling of epigenetic factors by cell geometric constraints. *Proc Natl Acad Sci U.S.A.* **116**, 13200–13209 (2019).

40. M. Wang, Y. Yang, L. Han, S. Han, N. Liu, F. Xu, F. Li. Effect of three-dimensional ECM stiffness on cancer cell migration through regulating cell volume homeostasis. *Biochemical and Biophysical Research Communications* **528**, 459-465 (2020).

41. M. Guo, A.F. Pegoraro, A. Mao, E.H. Zhou, P.R. Arany, Y. Han, D.T. Burnette, M.H. Jensen, K.E. Kasza, J.R. Moore, F.C. Mackintosh. J.J. Fredberg, D.J. Mooney, J. Lippincott-Schwartz, D.A. Weitz. Cell volume change through water efflux impacts cell stiffness and stem cell fate. *Proc Natl Acad Sci U.S.A.* **114**, E8618–E8627 (2017).

42. M.R. Zanotelli, Z.E. Goldblatt, J.P. Miller, F. Bordeleau, J. Li, J.A. VanderBurgh, M.C. Lampi, M.R. King, C.A. Reinhart-King. Regulation of ATP utilization during metastatic cell migration by collagen architecture. *Mol Biol Cell* **29**, 1–9 (2018).

43. H. Wang, A.S. Abhilash, C.S. Chen, R.G. Wells, V.B. Shenoy. Long Range Force Transmission in Fibrous Matrices Enabled by Tension-Driven Alignment of Fibers. *Biophys J* **107**, 2592-2603 (2014).

44. D.O. Velez, S.K. Ranamukhaarachchi, A. Kumar, R.N. Modi, E.W. Lim, A.J. Engler, C.M. Metallo, S.I. Fraley. 3D collagen architecture regulates cell adhesion through degradability, thereby controlling metabolic and oxidative stress. *Integr Biol* **11**, 221-234 (2019).

45. W. Wang, J.P. Miller, S.C. Pannullo, C.A. Reinhart-King, F. Bordeleau. Quantitative assessment of cell contractility using polarized light microscopy. *J Biophotonics* **11**, e201800008 (2018).

46. S. Hassan, J. Kim. Polyacrylamide hydrogels. IV. Near-perfect elasticity and rate-dependent toughness. *J Mech Phys Solids* **158**, 104675 (2022).

47. F. Lin, H. Zhang, J. Huang, C. Xiong. Substrate stiffness coupling TGF-β1 modulates migration and traction force of MDA-MB-231 human breast cancer cells in vitro. *ACS Biomater Sci Eng* **4**, 1337-1345 (2018).

48. S. Syed, J. Schober, A. Blanco, S.P. Zustiak. Morphological adaptations in breast cancer cells as a function of prolonged passaging on compliant substrates. *Plos one* **12,** e0187853 (2017).

49. J.J. Zimmerman, A. von Saint André-von Arnim, J. McLaughlin. "Cellular Respiration." in *Pediatric Critical Care (Fourth Edition)*, B.P. Fuhrman, J.J. Zimmerman, Eds. (Mosby, 2011), chap. 76.

50. K.E. Derricks, V. Trinkaus-Randall, M.A. Nugent. Extracellular matrix stiffness modulates VEGF calcium signaling in endothelial cells: individual cell and population analysis. *Integr Biol* **7**, 1011–1025 (2015).

51. S.A. Wicher, B.B. Roos, J.J. Teske, Y.H. Fang, C. Pabelick, Y.S. Prakash. Aging increases senescence, calcium signaling, and extracellular matrix deposition in human airway smooth muscle. *PLoS One* **16**, e0254710 (2021).

52. T.J. Kim, J. Seong, M. Ouyang, J. Sun, S. Lu, J.P. Hong, N. Wang, Y. Wang. Substrate rigidity regulates Ca2+ oscillation via RhoA pathway in stem cells. *J Cell Physiol* **218**, 285–293 (2009).

53. T.J. Kim, J. Sun, S. Lu, Y.X. Qi, Y. Wang. Prolonged mechanical stretch initiates intracellular calcium oscillations in human mesenchymal stem cells. *PLoS One* **9**, (2014).

54. S.L. McGee, K.F. Howlett, R.L. Starkie, D. Cameron-Smith, B.E. Kemp, M. Hargreaves. Exercise increases nuclear AMPK α2 in human skeletal muscle. *Diabetes* **52**, 926-928 (2003).

55. M. Kodiha, J.G. Rassi, C.M. Brown, U. Stochaj. Localization of AMP kinase is regulated by stress, cell density, and signaling through the MEK→ ERK1/2 pathway. *American Journal of Physiology-Cell Physiology* **293**, C1427-C1436 (2007).

56. T. Yeung, P.C. Georges, L.A. Flanagan, B. Marg, M. Ortiz, M. Funaki, N. Zahir, W. Ming, V. Weaver, P.A. Janmey. Effects of substrate stiffness on cell morphology, cytoskeletal structure, and adhesion. *Cell Motil Cytoskeleton* **60**, 24–34 (2005).

57. Y.L. Han, P. Ronceray, G. Xu, A. Malandrino, R.D. Kamm, M. Lenz, C.P. Broedersz, M. Guo. Cell contraction induces long-ranged stress stiffening in the extracellular matrix. *Proc Natl Acad Sci U.S.A.* **115**, 4075–4080 (2018).

58. P. Pakshir, M. Alizadehgiashi, B. Wong, N.M. Coelho, X. Chen, Z. Gong, V.B. Shenoy, C.A. McCulloch, B. Hinz. Dynamic fibroblast contractions attract remote macrophages in fibrillar collagen matrix. *Nat Commun* **10**, 1850 (2019).

59. S. Khetan, M. Guvendiren, W.R. Legant, D.M. Cohen, C.S. Chen, J.A. Burdick. Degradation-mediated cellular traction directs stem cell fate in covalently crosslinked three-dimensional hydrogels. *Nat Mater* **12**, 458-465 (2013).

60. M. Cóndor, C. Mark, R.C. Gerum, N.C. Grummel, A. Bauer, J.M. García-Aznar, B. Fabry. Breast Cancer Cells Adapt Contractile Forces to Overcome Steric Hindrance. *Biophys J* **116**, 1305–1312 (2019).

61. D.M. Graham, T. Andersen, L. Sharek, G. Uzer, K. Rothenberg, B.D. Hoffman, J. Rubin, M. Balland, J.E. Bear, K. Burridge. Enucleated cells reveal differential roles of the nucleus in cell migration, polarity, and mechanotransduction. *J Cell Biol* **217**, 895–914 (2018).

62. R. Milo, R. Phillips, *Cell Biology by the Numbers*. (Garland Science, New York, 2015).

63. F. Alisafaei, D. Shakiba, Y. Hong, G. Ramahdita, Y. Huang, L. E. Iannucci, M. D. Davidson, M. Jafari, J. Qian, C. Qu, D. Ju, D. R. Flory, Y. Huang, P. Gupta, S. Jiang, A. Mujahid, S. Singamaneni, K. M. Pryse, P. G. Chao, J. A. Burdick, S. P. Lake, E. L. Elson, N. Huebsch, V. B. Shenoy, G. M. Genin. Tension anisotropy drives fibroblast phenotypic transition by self-reinforcing cell–extracellular matrix mechanical feedback. *Nat Materials* **24**, 955-965 (2025).

64. A.E. Loneker, F. Alisafaei, A. Kant, D. Li, P.A. Janmey, V.B. Shenoy, R.G. Wells. Lipid droplets are intracellular mechanical stressors that impair hepatocyte function. *Proc Natl Acad Sci U.S.A.* **120**, e2216811120 (2023).

65. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez. D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, A. Cardona. Fiji: An open-source platform for biological-image analysis. *Nat Methods* **9,** 676–682 (2012).

66. The MathWorks Inc. MATLAB version: 9.13.0 (R2023b), https://www.mathworks.com. Natick, Massachusetts: The MathWorks Inc (2024).

67. W.H. Kruskal, W.A. Wallis. Use of Ranks in One-Criterion Variance Analysis. *J Am Stat Assoc* **47**, 583–621 (1952).

68. O.J. Dunn. Multiple Comparisons Using Rank Sums. *Technometrics* **6**, 241–252 (1964).

69. J. Fu, Y.K. Wang, M.T. Yang, R.A. Desai, X. Yu, Z. Liu, C.S. Chen. Mechanical regulation of cell function with geometrically modulated elastomeric substrates. *Nat Methods* **7**, 733–736 (2010).

70. H. Wang, A.A. Svoronos, T. Boudou, M.S. Sakar, J.Y. Schell, J.R. Morgan, C.S. Chen, V.B. Shenoy. Necking and failure of constrained 3D microtissues induced by cellular tension. *Proc Natl Acad Sci U.S.A.* **110**, 20923–20928 (2013).

71. F. Guilak, J.R. Tedrow, R. Burgkart. Viscoelastic Properties of the Cell Nucleus. *Biochem Biophys Res Commun* **269**, 781–786 (2000).

72. R. Vishavkarma, S. Raghavan, C. Kuyyamudi, A. Majumder, J. Dhawan, P.A. Pullarkat. Role of actin filaments in correlating nuclear shape and cell spreading. *PLoS One* **9**, (2014).

73. P. Hotulainen, P. Lappalainen. Stress fibers are generated by two distinct actin assembly mechanisms in motile cells. *J Cell Biol* **173**, 383–394 (2006).

74. M. Kovács, F. Wang, A. Hu, Y. Zhang, J.R. Sellers. Functional divergence of human cytoplasmic myosin II. Kinetic characterization of the non-muscle IIA isoform. *J Biol Chem* **278**, 38132–38140 (2003).

75. K. Burridge, E.S. Wittchen. The tension mounts: Stress fibers as force-generating mechanotransducers. *J Cell Biol* **200**, 9–19 (2013).

76. M.M. Kozlov, L.V. Chernomordik, Membrane tension and membrane fusion. *Curr Opin in Struct Biol* **33**, 61–67 010 (2015).

77. Z. Shi, Z.T. Graber, T. Baumgart, H.A. Stone, A.E. Cohen. Cell Membranes Resist Flow. *Cell* **175**, 1769-1779 (2018).

78. Y. Chen, A.C. Munteanu, Y.F. Huang, J. Phillips, Z. Zhu, M. Mavros, W. Tan. Mapping Receptor Density on Live Cells by Using Fluorescence Correlation Spectroscopy. *Chemistry – A European Journal* **15**, 5327–5336 (2009).

79. D.J. Irvine, K.A. Hue, A.M. Mayes, L.G. Griffith. Simulations of cell-surface integrin binding to nanoscale-clustered adhesion ligands. *Biophys J* **82**, 120–132 (2002).

80. COMSOL Multiphysics® v. 6.0. www.comsol.com. COMSOL AB, Stockholm, Sweden (2024).

81. M. Menegollo, I. Tessari, L. Bubacco, G. Szabadkai. "Determination of ATP, ADP, and AMP Levels by Reversed-Phase High-Performance Liquid Chromatography in Cultured Cells." in *Calcium Signalling: Methods and Protocols*. A. Raffaello, R.D. Vecellio, Eds. (Springer New York, 2019) pp. 223–232.

Acknowledgments

We express our gratitude to X. Chen for assistance with development of early versions of our models, E. Sorokina for assistance in maintaining cell cultures, and C. Madl for allowing generous use of his imaging system.

Funding: This work was supported by:
National Cancer Institute award U54CA261694 (V.B.S.)
National Institute of Biomedical Imaging and Bioengineering awards R01EB017753 and R01EB030876 (V.B.S.)
National Institute of General Medical Sciences award R01GM155943 (V.B.S.)
NSF Center for Engineering Mechanobiology Grant CMMI-154857 (V.B.S.)

1045 NSF Grant DMS-2347834 (V.B.S.)
1046

Author contributions:
Conceptualization: VBS, CRK, JT, AJ
Methodology: VBS, JT, AJ
Investigation: JT, AJ
Visualization: JT, AJ
Supervision: VBS
Writing—original draft: JT, AJ
Writing—review & editing: VBS, CRK, LP, YS
1055

1056 **Competing interests:**
1057 Authors declare that they have no competing interests.
1058

Data and materials availability:
All data are available in the main text or the supplementary materials. We provide open access to
all files used to generate and run our models as well as custom written code used to analyze model
results in a public GitHub repository: https://github.com/ShenoyLab/Metabolism. Within, readers
will find instructions to download, install, and reproduce model data. Unprocessed images from
novel experiments are also available within this repository.
1065