Article

Nuclear Translocation of SIRT4 Mediates Deacetylation of U2AF2 to Modulate Renal

Fibrosis Through Alternative Splicing-mediated Upregulation of CCN2

Guangyan Yang[1,2,4], Jiaqing Xiang[1,2,4], Xiaoxiao Yang[3,4], Xiaomai Liu[1,2], Yanchun

Li[1,2], Lixing Li[1,2], Lin Kang[1,2,*], Zhen Liang[1,2,*], Shu Yang[1,2,5,*]

[1]Department of Geriatrics, The First Affiliated Hospital of Southern University of

Science and Technology, Shenzhen, Guangdong 518020, China

[2]Guangdong Provincial Clinical Research Center for Geriatrics, Shenzhen Clinical

Research Center for Geriatrics, The First Affiliated Hospital of Southern University of

Science and Technology, Shenzhen, Guangdong 518020, China

[3]Key Laboratory of Metabolism and Regulation for Major Diseases of Anhui Higher

Education Institutes, College of Food and Biological Engineering, Hefei University of

Technology, Hefei, China

[4]These authors contributed equally

[5]Lead contact

Running title: SIRT4 exacerbates renal fibrosis

Corresponding author:

Shu Yang

Email: yang.shu@szhospital.com

Zhen Liang

Email: liang.zhen@szhospital.com

Lin Kang

Email: kang.lin@szhospital.com

Abstract

TGF-β stimulates CCN2 expression which in turn amplifies TGF-β signaling, thereby promoting extracellular matrix production and accelerating the pathological progression of fibrotic diseases. Alternative splicing plays an important role in multiple disease development, while U2 small nuclear RNA auxiliary factor 2 (U2AF2) is an essential pre-mRNA splicing factor in an early step of splicing. However, the molecular mechanism underlying abnormal *CCN2* expression upon TGF-β stimulation remains unclear. This study elucidates that SIRT4 acts as a master regulator for CCN2 expression in response to TGF-β by modulating U2AF2-mediated alternative splicing. Analyses of renal biopsy specimens from patients with CKD and mouse fibrotic kidney tissues revealed marked nuclear accumulation of SIRT4. The tubulointerstitial fibrosis was alleviated by global deletion or tubular epithelial cell (TEC)-specific knockout of *Sirt4*, and aggravated by adeno-associated virus-mediated SIRT4 overexpression in TECs. Furthermore, SIRT4 was found to translocate from the mitochondria to the cytoplasm through the BAX/BAK pore under TGF-β stimulation. In the cytoplasm, TGF-β activated the ERK pathway and induced the phosphorylation of SIRT4 at Ser36, further promoting its interaction with importin α1 and subsequent nuclear translocation. In the nucleus, SIRT4 was found to deacetylate U2AF2 at K413, facilitating the splicing of CCN2 pre-mRNA to promote CCN2 protein expression. Importantly, exosomes containing anti-SIRT4 antibodies were found to effectively mitigate the UUO-induced kidney fibrosis in mice. Collectively, these findings indicated that SIRT4 plays a role in kidney fibrosis by regulating CCN2 expression via the pre-mRNA splicing.

Key Words: Kidney fibrosis; pre-mRNA splicing; U2 snRNP; SIRT4; U2AF2; CCN2

Graphical abstract

1. Introduction

In the kidney and other organ systems, the overexpression (OE) of cellular communication network 2 (CCN2), also known as connective tissue growth factor, is widely recognized as a marker of fibrotic activity[1]. Transforming growth factor-β (TGF-β) is a master regulator of tissue growth, regeneration, remodeling, and fibrosis, and most TGF-β responses involve CCN2 stimulation at some level, such as the stimulation of extracellular matrix (ECM) components and fibrosis[2,3]. Numerous signaling molecules are involved in the crosstalk and integration of TGF-β and CCN2 effects and vary depending on the cell type and the physiological or pathological process involved. For instance, TGF-β-stimulated SMADs are necessary for the induction of CCN2 expression in normal fibroblasts and basal CCN2 induction in scleroderma fibroblasts, while maintenance of CCN2 expression is independent of SMADs[4]. However, it remains unknown whether TGF-β1 regulates CCN2 expression via non-transcriptional pathways.

Sirtuins (Sirts), mammalian Sir2 orthologs, are a highly conserved family of nicotinamide adenine dinucleotide-dependent protein deacetylases and act as

important regulators of the aging process, inflammation, cancer, and metabolic diseases[5-7]. Among the seven known mammalian Sirts, SIRT4 possesses ADP-ribosyltransferase, lipoamidase, and deacylase activities[8,9]. The inhibition of SIRT4 expression has been shown to increase the fat-oxidation capacity of the liver and mitochondrial function in the muscle[10]. As loss of fatty acid β-oxidation in the proximal tubule is a critical mediator of acute kidney injury and eventual fibrosis[11,12], we hypothesize that SIRT4 may act as a pro-fibrotic factor. However, other studies have demonstrated that SIRT4 OE inhibits glutamine metabolism[13], which is necessary for collagen protein synthesis[14], suggesting a potential protective role of SIRT4 in the context of fibrosis. However, the specific mechanism by which SIRT4 regulates renal fibrosis remains unclear. Therefore, research is needed to further explore the role and mechanisms of SIRT4 in renal fibrosis as well as potential therapeutic strategies.

An estimated 60% of all human genes undergo alternative splicing, a highly regulated process that produces splice variants with different functions[15]. There are five small nuclear ribonucleoproteins (snRNPs)—U1, U2, U5, and U4/U6. The status of these proteins helps in maintaining sufficient proteomic diversity for the functional requirements of cell fates and body homeostasis[16]. U2 small nuclear RNA auxiliary factor 1 (U2AF1), together with U2AF2, forms the U2AF complex that recognizes and binds to the 3′ splice site of pre-mRNA and recruits U2 snRNPs, thereby facilitating the assembly of the spliceosome, a large RNA-protein complex responsible for the splicing of pre-mRNA[17-20]. The SF3B complex, the core of the U2 snRNP, comprises SF3B1, SF3B2, SF3B3, SF3B4, SF3B5, SF3B6, and PHF5A/ SF3B14b[21]. Dysregulation of U2AF2 can lead to the disruption of splicing events, which may contribute to the development and progression of kidney fibrosis. Accordingly, understanding the role of U2AF2 in kidney fibrosis may provide valuable insights into its underlying molecular mechanisms.

Here, we demonstrated that TGF-β1 induced SIRT4 nuclear translocation, resulting in the deacetylation of U2AF2 and recruitment of U2 snRNP, subsequently contributing to pre-mRNA splicing and enhanced protein expression of CCN2.

Overall, our study revealed that SIRT4 inhibition can alleviate the progression of
renal fibrosis by suppressing CCN2 expression.

2. Methods

2.1 Studies in animals

All Animal care and experimental protocols for in vivo studies conformed to the

Guide for the Care and Use of Laboratory Animals published by the National

Institutes of Health (NIH; NIH publication no.:85–23, revised 1996). The sample size

for the animal studies was calculated based on a survey of data from published

research or preliminary studies. $Sirt4^{flox/flox}$ (C57BL/6J-$Sirt4^{em1flox}$/Cya; Strain ID:

CKOCMP-75387-$Sirt4$-B6J-VA), Col1a2-Cre/ERT2 mice (Cat#: C001248) and

Cdh16-Cre mice (Cat#: C001452) obtained from Cyagen Biosciences (Guangzhou)

Inc (Guangzhou, Guangdong, China). C57BL/6J mice, $Sirt4^{-/-}$

(C57BL/6JGpt-$Sirt4^{em13Cd1976}$/Gpt; Strain ID: T011568) and $U2af2^{-/-}$

(C57BL/6JGpt-$U2af2^{em2Cd5259}$/Gpt; Strain ID: T029898) mice based on C57BL/6J

background were purchased from Gempharmatech Co. Ltd (Jiangsu, Nanjing, China).

These mice were maintained in SPF units of the Animal Center of Shenzhen People's

Hospital with a 12 h light cycle from 8 a.m. to 8 p.m., 23 ± 1 °C, $60 - 70$ % humidity.

Mice were allowed to acclimatize to their housing environment for 7 days before the

experiments. At the end of the experiments, all mice were anesthetized and euthanized

in a CO_2 chamber, followed by the collection of kidney tissues. All animals were

randomized before treatment. Mice were treated in a blinded fashion as the drugs used

for treating animals were prepared by researchers who did not carry out the treatments.

No mice were excluded from the statistical analysis. Studies were performed in

accordance with the German Animal Welfare Act and reporting follows the ARRIVE

guidelines.

2.2 Generation of cell type–specific *Sirt4* conditional knockout mice

To generate fibroblast-specific conditional *Sirt4* knockout mouse line, $Sirt4^{flox/flox}$ mice

were bred with Col1a2-Cre/ERT2 transgenic mice. To activate the Cre-ERT system,

tamoxifen (80 mg/kg/day, dissolved in olive oil) was injected intraperitoneally for 4 consecutive days 2 weeks before the induction of renal fibrosis in control and S4FKO mice. After UUO, uIRI, or FA, tamoxifen diet was administered until sacrifice in order to ensure the deletion of Sirt4 in newly generated myofibroblasts. To generate TECs-specific conditional Sirt4 knockout mouse line (S4TKO), $Sirt^{flox/flox}$ mice were bred with Cdh16-Cre transgenic mice.

2.3 Mouse kidneys were transfected with adeno-associated virus vector (AAV)

8-week-old mice received in situ renal injection with AAV9- empty vector (AAV9-*Ctrl*; control group), AAV9-*Ksp-Sirt4* (*Sirt4^{OE}* group; *Ksp*, tubule specific promoter), AAV9-*Ksp*-wild type U2af2 (*wtU2af2*), and AAV9-*Ksp*-mutant U2af2 (*mU2af2*) at three independent points (10 - 15 μl virus per poin; virus injected dose: 2.5 E + 11 v.g.) in the kidneys of mice (n = 6). Adeno-associated virus type 9 constructs, including GV501 empty vector, *Sirt4*, *wtU2af2*, and *mU2af2* were provided by GeneChem Company (Shanghai, China).

2.4 Mice kidney fibrotic models

Male C57BL/6, $Sirt4^{-/-}$, $Sirt^{flox/flox}$, and cell type–specific conditional knockout mice (~8 to 10 weeks old) were subjected to various kidney injury models to induce renal fibrosis. UUO was performed by permanent ligation of the right ureter with 6-0 silk. Ureter-ligated kidneys and contralateral kidneys (CLs), used as nonfibrotic controls, were collected 10 days after surgery. To establish uIRI, left renal pedicles were clamped with microaneurysm clips for 30 minutes followed by reperfusion. During uIRI surgery, mice were placed on a heating pad to maintain body temperature at 37°C. Injured and contralateral kidneys were collected 1 day or 28 days after surgery for analyses. Folic acid (FA)–induced renal fibrosis was conducted by single intraperitoneal injection of 250 mg/kg folic acid (Sigma-Aldrich, 7876) dissolved in 0.3M sodium bicarbonate, and mice were sacrificed 14 days after FA treatment. Mice injected with sodium bicarbonate served as vehicle control.

2.5 Study approval

All animal care and experimental protocols for in vivo studies conformed to the Guide

for the Care and Use of Laboratory Animals, published by the National Institutes of

Health (NIH; NIH publication no.: 85–23, revised 1996), was approved by the Animal

Care Committees of the First Affiliated Hospital of Southern University of Science

and Technology (No. AUP-230809-LZ-0426-01), and were performed in compliance

with the ARRIVE guidelines. Studies with human participants were conducted in line

with the Declaration of Helsinki. The studies were approved by Ethics Committee of

the First Affiliated Hospital of Southern University of Science and Technology. The

written consent obtained from patients was informed.

2.6 Quantification and Statistical Analysis

All data were generated from at least three independent experiments. Each value was

presented as the mean ± SD. All raw data were initially subjected to a normal

distribution and analysis by one-sample Kolmogorov-Smirnov (K-S) nonparametric

test using SPSS 22.0 software. For animal and cellular experiments, a two-tailed

unpaired student's t-test was performed to compare the two groups. One-way ANOVA

followed by the Bonferroni's post-hoc test was used to compare more than two groups.

To avoid bias, all statistical analyses were performed blindly. Statistical significance

was indicated at $^*P < 0.05$, $^{**}P < 0.01$, and $^{***}P < 0.001$.

3. Results

3.1. Nuclear localization of SIRT4 increases in fibrotic kidney

The application of mitochondrial SIRT4 tripartite abundance reporter, a tripartite

probe for visualizing the distribution of SIRT4 between mitochondria and the nucleus

in single cells[22], proved the importation of SIRT4 into the mitochondrial matrix and

demonstrated its localization in the nucleus under mitochondrial stress conditions.

Nuclear accumulation of SIRT4 was observed in the kidneys following unilateral

ureteral occlusion (UUO) or unilateral renal ischemia-reperfusion injury (uIRI)

surgery **(Fig. 1A–D)**. Consistently, elevated nuclear accumulation of SIRT4 was

observed in kidney sections from patients with chronic kidney disease (CKD) with

severe collagen deposition **(Fig. 1E, F)**.

3.2 Global deletion of *Sirt4* protects against kidney fibrosis

To determine the role of SIRT4 in kidney fibrosis development in vivo, wild-type

(WT) and *Sirt4* global knockout (S4KO) mice were subjected to UUO, uIRI, and folic

acid (FA) treatment. Sirius red and Masson's trichrome staining of the kidney sections

revealed extensive renal fibrosis in WT mice following UUO **(Fig. 2A)**, uIRI **(Fig.**

2E), and FA administration **(Fig. S1B)**. In contrast, the global deletion of *Sirt4*

resulted in a remarkably reduction in the extent of renal fibrosis in all three kidney

injury models **(Fig. 2A, E and S1B)**. Immunoblots of whole-kidney tissue lysates

showed that UUO, uIRI, and FA treatment led to induced levels of the markers of

kidney fibrosis (CCN2, FN1, COL1A1, COL3A1, and α-SMA), while the expression

of E-cadherin, a hallmark of epithelial-mesenchymal transition, was decreased, with

the changes being more pronounced in WT mice than in S4KO mice **(Fig. 2B, F and**

S1A). Consistent with the attenuated post-injury fibrotic response in S4KO mouse

kidneys observed via imaging studies, deletion of *Sirt4* also mitigated the

upregulation of the profibrotic genes *Col1a1*, *Fn1*, *Eln*, *Ccn2*, *Acta2*, and *Col3a1* **(Fig.**

2C, G and S1C). In addition, the mRNA levels of both neutrophil

gelatinase-associated lipocalin (*Ngal*) and kidney injury molecule 1 (*Kim-1*), which

are markers of acute kidney injury, were significantly increased in the kidney tissues

from WT and S4KO mice following injury **(Fig. 2D, H and S1D)**. Notably, the levels

of *Ngal* and *Kim-1* were significantly reduced in S4KO mice compared to those in

WT mice following injury **(Fig. 2D, H and S1D)**. Collectively, these results suggest

that *Sirt4* deletion protects mice against renal fibrosis.

3.3. Deletion of *Sirt4* in renal tubule epithelial cells markedly attenuates the

extent of kidney fibrosis following injury

Using conventional agarose gel-based RT-PCR and Western blot analysis, we

analyzed the expression of SIRT4 in renal parenchymal cells, including mouse

podocytes (MPCs), glomerular endothelial cells (GECs), kidney fibroblasts (KFs),

and tubule epithelial cells (TECs). Compared to their levels in TECs, the basal levels

of SIRT4 in MPCs and GECs were relatively low, while in KFs is moderate (**Fig.**

S1E). To determine the *in vivo* contribution of SIRT4 in renal TECs and fibroblasts to

the development of kidney fibrosis, we generated TEC-specific (S4TKO;

Cdh16-cre/ERT2×*Sirt4$^{flox/flox}$*)   and   fibroblast-specific   (S4FKO;

Col1a2-Cre/ERT2×*Sirt4$^{flox/flox}$*) *Sirt4* knockout mice; in these experiments, Cadh16-cre

mice and tamoxifen-treated Col1a2-Cre mice were used as controls, respectively. The

extent of kidney fibrosis induced by UUO was markedly reduced in S4TKO mice

compared to that in control mice (**Fig. 2I, J**). In addition, the expression levels of

kidney fibrosis markers (CCN2, FN1, COL1A1, COL3A1, and α-SMA) were also

significantly reduced in S4TKO compared to those in control mice following UUO

surgery (**Fig. 2J**). Consistently, the profibrotic genes (*Col1a1, Fn1, Eln, Ccn2, Acta2*,

and *Col3a1*) and acute kidney injury markers *Nagl* and *Kim-1* were downregulated in

S4TKO mice compared to those in control mice (**Fig. 2K, L**). Nevertheless, the

targeted deletion of SIRT4 in fibroblasts did not affect the extent of UUO-induced

kidney fibrosis (**Fig. S1F–H**). These results suggest that SIRT4 expressed in TECs but

not in fibroblasts primarily contributes to the pathogenesis of kidney fibrosis.

3.4 SIRT4 OE aggravates kidney fibrosis

Since the deletion of *Sirt4* attenuated the post-injury fibrotic response, we next

determined whether *Sirt4* OE can aggravate the response. We introduced SIRT4 into

WT mouse kidneys using the adeno-associated virus serotype 9 vector with a

tubule-specific *Ksp-cadherin* promoter, and AAV9-*Ksp-null* transfection was used as

the control treatment. Targeting *Sirt4* OE in kidney TECs markedly aggravated the

extent of kidney fibrosis induced by UUO, uIRI, and FA treatment compared to that in

the control mice (**Fig. 3A, D, G**). The expression levels of kidney fibrosis markers

(CCN2, FN1, COL1A1, COL3A1, and α-SMA) and the decreased E-cadherin were

also significantly enhanced in *Sirt4* OE mice compared to those in the control mice

following kidney injury (**Fig. 3B, E, H**). Consistently, *Sirt4* OE upregulated the

transcription of profibrotic genes, namely *Col1a1*, *Fn1*, *Eln*, *Ccn2*, *Acta2*, and *Col3a1*,

in mouse kidneys following injury, compared to that in the control group **(Fig. 3C, F,**

I). Further analysis revealed an evident upregulation of *Ngal* and *Kim-1* transcripts in

the kidneys of *Sirt4* OE mice compared to that in the kidneys of control mice in

response to UUO **(Fig. 3C)**. Notably, targeting WT *Sirt4* OE in kidney TECs

markedly reversed the decline in collagen deposition and the reduction in

ECM-related protein expression in S4KO mice **(Fig. S2A–C)**. These results

demonstrate that interventional SIRT4 OE in renal TECs exacerbates kidney fibrosis

progression.

3.5. SIRT4 interacts with U2AF2 under TGF-β stimulation

To explore the action mechanism of SIRT4 in kidney fibrosis, we first identified the

proteins interacting with SIRT4 in TECs. We employed rapid immunoprecipitation

and mass spectrometry of endogenous proteins (RIME), which is an efficient and

unbiased proteomic approach for identifying interacting proteins. Human TECs were

treated with TGF-β1 or the control (DMSO) for 24 h, followed by protein-DNA

crosslinking in 1% formaldehyde. Cells were sonicated, followed by

immunoprecipitation with an SIRT4 antibody **(Fig. 4A)**. A total of 1081 unique

proteins copurified with SIRT4 were identified using mass spectrometry. We only

considered SIRT4-associated proteins that occurred in three of three independent

replicates and excluded any proteins that appeared in any one of the IgG control

RIMEs. As a result, we selected of 22 SIRT4-associated proteins. Notably, the SIRT4

interaction involved numerous specific nuclear localization proteins, such as PUF60,

U2AF2, RPS2, TSR1, ZC3H15, and SART1. Of these, U2AF2 and PUF60 function

cooperatively in pre-mRNA splicing[23]. The levels (normalized) of PUF60 and U2AF2

are shown in the **Figure 4B**. We next determined whether PUF60 or U2AF2 was

recruited to SIRT4 in TECs stimulated with TGF-β1. Immunoblotting of

immunoprecipitated SIRT4 with anti-PUF60 and -U2AF2 antibodies showed that

TGF-β1 treatment resulted in an interaction among SIRT4, PUF60, and U2AF2 **(Fig.**

4C, left panel). Whole-kidney tissue lysates consistently showed an interaction

among SIRT4, PUF60, and U2AF2 **(Fig. 4C, middle panel)**. As U2AF2 has been

shown to interact with PUF60[23], we tested whether SIRT4 interacts with either of the

two proteins via the U2AF2-PUF60 interaction. The interaction between SIRT4 and

PUF60 was abolished in U2af2-knockout cells **(Fig. 4D)**. However, the interaction of

SIRT4 and U2AF2 was not affected in *Puf60* knockdown cells (data not shown).

Furthermore, exogenous co-IP revealed an interaction between SIRT4 and U2AF2

(Fig. 4C, right panel). Together, these results suggest that SIRT4 interacts with

U2AF2 under TGF-β1 stimulation or kidney injury.

3.6. U2AF2 acetylation is decreased under cellular stress

Accumulating   evidence   suggests   that   SIRT4   can   show   weak

ADP-ribosyltransferase[8,24,25] as well as substrate-specific deacetylase activity[26-28],

akin to that observed for SIRT6 and SIRT7[29,30]. Hence, we tested whether SIRT4

regulates the ADP-ribosylation or acetylation of U2AF2. Our results showed that *Sirt4*

OE reduced the levels of acetylated U2AF2 **(Fig. 4E)**. However, the ADP-ribosylation

of U2AF2 was extremely low in both SIRT4 OE and control cells **(Fig. 4E)**.

According to the database Compendium of Protein Lysine Modifications 4.0, lysine

(K) 70, 276, 292, 413, 453, and 462 on U2AF2 can be acetylated. Hence, we mutated

all six K residues to arginine (R), which mimics the deacetylated state of the protein.

As shown in **Figure 4F**, the U2AF2-K413R mutant showed reduced acetylation

compared to U2AF2 WT. Conservation analysis of U2AF2 indicated that K413 is a

highly conserved site spanning from *Schizosaccharomyces pombe* to *Homo sapiens*

(Fig. 4G). Next, we examined whether U2AF2 K413 is the key acetylation site in

response to TGF-β1 stimulation. TGF-β1 treatment reduced the levels of acetylated

U2AF2 in WT but had little effect in the K413R mutant U2AF2 **(Fig. 4H)**.

Consistently, U2AF2 acetylation decreased in the kidney following injury **(Fig. 4I)**.

Interestingly, U2AF2 acetylation was also remarkably reduced in the cells subjected

to serum starvation, endoplasmic reticulum (ER) stress (induced by tunicamycin),

viral infection [poly (I:C)], or DNA damage stress (induced by cisplatin) **(Fig. 4J)**.

These results collectively demonstrate that U2AF2 deacetylation occurs universally in

response to cellular stress.

3.7. U2AF2 acetylation affects the interaction of U2AF and SF3B complexes

Next, we investigate the effect of U2AF2 acetylation at K413 on the formation of the
U2AF complex and recruitment of the SF3B complex **(Fig. 4K)**. The expression of
K413R did not regulate the stability of the SF3B and U2AF components. Next, we
performed an immunoprecipitation assay to detect the interaction between the U2AF
and SF3B complexes using U2AF2-WT- and U2AF2-K413R-expressing cells.
Interestingly, U2AF2 K413R interacted strongly with components of the SF3B
complex and with U2AF1 compared to that with U2AF2 WT **(Fig. 4K)**. Notably, the
interaction of U2AF2 and U2AF1 was not obviously regulated after RNAase
treatment, while the interaction of U2AF2 and SF3B complex was reduced following
RNase treatment **(Fig. 4K)**. These findings implied the interactions between U2AF1
and U2AF2 were independent of RNA binding, while the interaction of U2AF2 and
SF3B complex were dependent on RNA binding. As a component of the U2AF
complex, U2AF2 interacts with U2AF1 to form the U2AF complex, which recognizes
the 3′ splice site (3' SS) of U2 introns and recruits U2 snRNP[17] **(Fig. 4L)**. Taken
together, these data suggest that U2AF2 acetylation regulates the formation of the
U2AF complex and recruitment of the SF3B complex.

3.8. U2AF2 acetylation under cellular stress is not HAT1-dependent

To identify the histone acetyltransferase (HAT) responsible for U2AF2 acetylation
under cellular stress, we co-transfected U2AF2 with different HATs, namely CBP,
p300, HAT1, P/CAF, KAT5, and KAT8. Of these, HAT1 mainly promoted U2AF2
acetylation **(Fig. 4M)**. Both endogenous and exogenous co-IP results showed an
interaction between HAT1 and U2AF2 **(Fig. 4N)**. Further experiments were
performed to determine whether U2AF2 acetylation in response to stress is dependent
on HAT1. We found that knockdown of *Hat1* reduced U2AF2 acetylation, while *Hat1*
OE induced the acetylation **(Fig. 4O)**. Furthermore, U2AF2 acetylation was not
affected by enzymatically defective HAT1 **(Fig. 4P)**. Although U2AF2 interacted with

HAT1, the interaction was altered weakly under TGF-β1 stimulation (**Fig. 4Q, left**

panel). Consistently, under *Hat1* knockdown conditions as well, TGF-β1 stimulation

further reduced U2AF2 acetylation (**Fig. 4Q, right panel**). Thus, although HAT1

acetylated U2AF2, the decreased acetylation level of U2AF2 by TGF-β1 stimulation

was independent of HAT1.

3.9. SIRT4 deacetylates U2AF2 at Lys413

As SIRT4 interacted with U2AF2, we intend to figure out the directly role of SIRT4 in

U2AF2 acetylation. We found that SIRT4 OE reduced U2AF2 acetylation, whereas

SIRT4 knockdown increased its acetylation (**Fig. 5A**). Additionally, TGF-β1

stimulated the interaction between SIRT4 and U2AF2 in a dose dependent manner

(**Fig. 5B**). Consistently, the TGF-β1-induced decrease in U2AF2 acetylation was

blocked by the knockdown of *Sirt4* (**Fig. 5C**). In contrast, SIRT4 OE further enhanced

the TGF-β1-induced U2AF2 deacetylation (**Fig. 5C**). Compared to U2AF2 WT,

U2AF2 K413R blocked the SIRT4 OE-induced deacetylation of U2AF2 as well as the

interaction of U2AF2 with U2AF1 and the SF3B complex under TGF-β1 stimulation

(**Fig. 5D**). Accordingly, these phenotypes induced by SIRT4 OE were restricted when

an enzyme-defective SIRT4 (SIRT4 H161Y) was used (**Fig. 5E**). Molecular docking

simulations showed that U2AF2 Lys413 was present at the contact surface between

SIRT4 and U2AF2 (**Fig. 5F**), further demonstrating that SIRT4 can interact with

U2AF2 to promote its deacetylation.

To determine whether S4KO alleviates renal fibrosis by upregulating the

acetylation of U2AF2 in mouse kidney, we crossed $Sirt4^{+/-}$ and $U2af2^{+/-}$ mice to

generate $Sirt4^{-/-}U2af2^{-/-}$ (DKO) mice and then performed UUO surgery. No body

weight differences were noted due to genetic manipulation (data not shown).

Compared to that in WT mice, the acetylation level of U2AF2 was increased, but

renal fibrosis was reduced in S4KO mice after UUO surgery (**Fig. 5G, H, K**). The

UUO-induced kidney injury was largely reduced in DKO and $U2af2^{-/-}$ mice, which

showed comparable collagen deposition and expression of ECM-related proteins and

mRNA (**Fig. 5G, H, K**). We next treated $U2af2^{-/-}$ mice with AAV-wt*U2af2* or

AAV-m*U2af2* (*U2af2* K413R) to re-express the two types of U2AF2 proteins through

in situ renal injection. Mice were followed by administered of AAV-*Sirt4* or AAV-Ctrl

and then subjected to UUO surgery. The U2AF2 K413R OE mice showed a greater

extent of renal fibrosis than did the wtU2AF2 OE mice, as evidenced by collagen

deposition and increased ECM protein levels **(Fig. 5I, J, L).** Importantly, the extent of

renal fibrosis was remarkably augmented in mice injected with wt*U2af2* OE and *Sirt4*

OE compared to that in wtU2AF2 OE mice **(Fig. 5I, J, L)**. However, *Sirt4* OE had

little synergistic effect on mice injected with AAV-m*U2af2* **(Fig. 5I, J, L)**. Together,

these data support that SIRT4-mediated deacetylation of U2AF2 at K413 is an

important step in SIRT4-induced renal fibrosis.

3.10. SIRT4 deacetylates U2AF2 at K413 to regulate *CCN2* expression and

pre-mRNA splicing

U2AF1 S34F and Q157R mutations have been reported to compromise U2AF2–RNA

interactions, resulting predominantly in intron retention and exon exclusion[31]. Hence,

we hypothesized that the SIRT4-mediated deacetylation of U2AF2 participates in the

regulation of these processes. RNA-sequencing was performed to identify the changes

in gene expression and alternative splicing in adenovirus-mediated

SIRT4[OE]-transfected cells or adenovirus-Ctrl-transfected cells after TGF-β1

stimulation **(Fig. 6A)**. Surprisingly, SIRT4 stimulation upregulated several genes ($P <$

0.01; Table EV1) that are involved in mRNA processing and the TGF-β signaling

pathway. A total of 142 differentially expressed genes ($P < 0.0005$) were identified

(Fig. 6A). Meanwhile, 248 genes showed differential intron retention (FDR < 0.05;

Table EV2) **(Fig. 6A)**. This analysis revealed four genes that were common between

the differentially expressed genes and the differentially spliced genes **(Fig. 6A)**. *Ccn2*,

one of these four genes, ranked third among the all upregulated genes ($P < 0.0005$)

after SIRT4 overexpression. Given that intron retention due to abnormal splicing can

trigger nonsense-mediated mRNA decay, gene expression could be affected by a

change in splicing pattern. The mRNA levels of *CCN2* were remarkably increased in

SIRT4[OE] cells but reduced in *Sirt4*-knockdown cells **(Fig. 6B)**. In addition, *CCN2*

mRNA levels were significantly increased in U2AF2 K413R-transfected cells but

remarkably reduced in U2AF2 K413Q [lysine to glutamine (Q) mutant for protein

hyperacetylation mimic]-transfected cells **(Fig. 6B)**. In line with the mRNA levels,

SIRT4 OE significantly elevated the protein levels of CCN2 in U2AF2 WT or U2AF2

K453Q-transfected cells but not in U2AF2-K413Q or K413R-transfected cells **(Fig.**

6C, D), suggesting that only the U2AF2 acetylation at K413 is efficient to regulate

CCN2 expression.

CCN2 is transcriptionally regulated by Sp1 in response to TGF-β1 stimulation[32].

Thus, we examined whether SIRT4 induced CCN2 expression through transcription

co-activator function. An analysis of the promoter region of *Ccn2* (including Sp1

binding site) showed that there was no increase in the promoter activity following

overexpression of U2AF2 WT, U2AF2-K413R or U2AF2-K413Q upon TGF-β1

stimulation **(Fig. 6E)**, implying that the SIRT4-induced CCN2 expression is not

transcription dependent. According to the RNA sequencing data, SIRT4 stimulation

increased the efficiency of *Ccn2* pre-mRNA exon2/intron2 and exon3/intron3 splicing

(Fig. 6F). Upon analyzing the gene sequence of *Ccn2*, we found that both *Ccn2*

introns 2 and 3 contain a TGA stop codon, implying that they could trigger the

degradation of abnormal mRNAs. Furthermore, we confirmed that SIRT4 knockdown

or U2AF2 K413Q increased the retention index of *Ccn2* introns 2 and 3, whereas

SIRT4 OE or U2AF2 K413 reduced the retention index **(Fig. 6G, H)**. U2AF2 K413R

or K413Q impeded the changes in intron 2 and 3 retention indices of *Ccn2* caused by

SIRT4 OE or *Sirt4* knockdown, respectively **(Fig. 6G, H)**. Next, we assessed whether

U2AF2 regulates CCN2 expression by regulating mRNA stability. We treated SIRT4

OE or *Sirt4* knockdown cells with Actinomycin D to block transcription and

measured *Ccn2* mRNA stability over time. The mRNA stability of *Ccn2* was

significantly enhanced in SIRT4 OE cells, whereas it was reduced in *Sirt4* knockdown

cells compared to that in control cells **(Fig. 6I)**. As expected, compared to that in

U2AF2 WT-transfected cells, U2AF2 K413R increased the mRNA stability of *Ccn2*,

while U2AF2 K413Q reduced its stability **(Fig. 6I)**. In addition, K413R and K413Q

repressed the changes in the stability of *Ccn2* mRNA caused by SIRT4 OE or *Sirt4*

knockdown, respectively **(Fig. 6I)**. Next, a RiboIP experiment was performed wherein

U2AF2 was immunoprecipitated to determine whether it can bind endogenous *Ccn2*

transcripts **(Fig. 6J)**. SIRT4 OE enhanced the interaction between U2AF2 and *Ccn2*

pre-mRNA (indicated by the PCR product containing intron 2 and exon 3 or intron 3

and exon 4) in U2AF2 WT-transfected cells but had little effect on their interaction in

U2AF2 K413R-transfected cells **(Fig. 6J)**. This indicates that the SIRT4-induced

deacetylation of U2AF2 at K413 increases CCN2 expression by regulating the

alternative splicing of *Ccn2*.

3.11. SIRT4 translocates from mitochondria to the cytoplasm through the

BAX/BAK pore after TGF-β stimulation

SIRT4 is located in the mitochondrial matrix under normal conditions[33]. A previous

study showed that SIRT4 translocates from the mitochondria into the cytoplasm upon

Wnt stimulation[34]. Hence, we conducted an immunofluorescence assay and organelle

separation experiment to compare the localization of SIRT4 before and after TGF-β1

treatment. Interestingly, SIRT4 localization changed from mitochondria to the

cytoplasm or even nucleus at after 12 h of TGF-β1 treatment **(Fig. 7A–C)**. Moreover,

TGF-β1 stimulation did not induce U2AF2 release from the nucleus into the

cytoplasm, even after 24 h (data not shown). Other stimulations such as serum

starvation, tunicamycin, poly (I: C), and cisplatin also induced this translocation of

SIRT4 **(Fig. 7D)**. It is well known that BAX and BAK are two key molecules of the

mitochondrial permeability transition pore. They form polymers on the mitochondrial

outer membrane and mediate the release of mitochondrial contents such as mtDNA,

mitochondrial dsRNA, and cytochrome c[35,36]. Therefore, we tested whether the

TGF-β1-induced release of SIRT4 is dependent on the BAX/BAK oligomeric pore.

Notably, BAX or BAK deficiency or MSN-125 (an effective oligomeric inhibitor of

Bax and Bak) treatment almost abolished the release of SIRT4 from mitochondria to

the cytoplasm as well as the upregulation of *Ccn2* under TGF-β1 stimulation **(Fig. 7E,**

F). As expected, the *in vivo* results showed that MSN-125 inhibited the translocation

of SIRT4 from mitochondria to the cytoplasm in the kidneys of mice following UUO

surgery **(Fig. 7G)**. Together, these results suggest that TGF-β1 induces the

translocation of SIRT4 from mitochondria to the cytoplasm in a BAX/BAK-dependent

manner.

3.12. ERK phosphorylates SIRT4 at Ser36 to promote the binding of SIRT4 to

importin α1 and nuclear translocation

As SIRT4 showed nuclear localization **(Fig. 7C)** and interacted with the nuclear

protein U2AF2 **(Fig. 4C)** under TGF-β1 stimulation, we investigated the mechanisms

underlying SIRT4 accumulation in the nucleus. Pretreatment of TECs with LY290042

(phosphoinositide 3-kinase inhibitor), SU6656 (Src inhibitor), SP600125 (JNK

inhibitor), and U0126 (MEK/ERK inhibitor) blocked TGF-β1-induced

phosphorylation of AKT, c-Src, c-Jun, and ERK1/2, respectively. Immunoblotting

analyses showed that only U0126 treatment abrogated the TGF-β1-induced nuclear

translocation of SIRT4 **(Fig. 8A, upper panel)**. Compared to the amount of cytosolic

SIRT4, nuclear SIRT4 is present in a small portion **(Fig. 8A, bottom panel)**. These

results were further supported by the immunofluorescence analyses **(Fig. 8B)**.

Additionally, expression of the Flag-ERK2 K52R kinase-dead mutant blocked the

TGF-β1-induced nuclear accumulation of SIRT4 and resulted in the accumulation of

phosphorylated SIRT4 in the cytosol **(Fig. 8C, left panel)**. Co-expression of a

constitutively active MEK1 Q56P mutant (expression of constitutively active MEK1

Q56P with WT ERK2) with Flag-tagged ERK2 WT or ERK2 K52R in TECs **(Fig. 8C,**

right panel) showed that expression of WT ERK2, but not ERK2 K52R, induced the

nuclear translocation of SIRT4. These results indicate that ERK activation is required

for the TGF-β1-induced nuclear translocation of SIRT4.

To further determine the relationship between ERK1/2 and SIRT4, we performed a

Co-IP assay and found that TGF-β1 treatment resulted in the binding of ERK1/2 to

SIRT4 **(Fig. 8D)**. MAP kinases bind to their substrates through a docking groove

comprising an acidic common docking (CD) domain and glutamic acid-aspartic acid

(ED) pockets[37]. Our results showed that mutation of either the ERK2 CD domain

(D316/319N) or the ED pocket (T157/158E) reduced the binding to endogenous

SIRT4 compared to that in the WT ERK2 control **(Fig. 8E)**. Combined mutations at both the CD domain and ED pocket (T/E-D/N) abrogated the binding of ERK2 to SIRT4 entirely **(Fig. 8E)**, indicating that ERK2 binds SIRT4 through its docking groove. Furthermore, the server provided docking information regarding SIRT4‑ERK2, indicating SIRT4 and ERK2 CD domain and ED pocket interactions **(Fig. 8F)**. ERK substrates often have a docking domain characterized by a cluster of basic residues, followed by an LXL motif (L represents Leu, but can also be Ile or Val; X represents any amino acid)[37]. An analysis of the docking information of SIRT4 - ERK2 and the amino acid sequence of SIRT4 revealed the putative ERK-binding sequence 77-EKVGLYARTDRR-88 and 203-GDVFLSE-209, which contain LXL motifs at V79/L81 and V205/L207, respectively. Immunoblotting of the immunoprecipitated Flag-SIRT4 proteins with an anti-ERK1/2 antibody showed that a SIRT4 V205/L207 mutant, but not a SIRT4 V79/L81 mutant, markedly reduced its binding to ERK1/2 **(Fig. 8G)**. These results indicate that the ERK2 docking groove binds to a CD domain in SIRT4 at V205/L207. Sequence analysis of SIRT4 revealed that it contains an ERK consensus phosphorylation motif (Ser-Pro) at the S36/P37 residues. Notably, S36A mutation completely abrogated the TGF-β1-dependent phosphorylation of SIRT4 **(Fig. 8H)**. Consistently, pretreatment with U0126 blocked the TGF-β1-induced S36 phosphorylation and nuclear translocation of SIRT4 **(Fig. 8I)**. In addition, expression of constitutively active MEK1 Q56P with WT ERK2, but not of ERK2 K52R, induced SIRT4 phosphorylation **(Fig. 8J)**. These results indicate that ERK2 specifically phosphorylates SIRT4.

Notably, we found that SIRT4 contains a potential nuclear localization signal (NLSs), a single type containing 3–5 basic amino acids with the weak consensus Lys-Arg/Lys-X-Arg/Lys[38], 248-KRVK-251. We mutated the K248/251 and R249 residues in the putative NLS sequences of SIRT4 to alanine (named as mNLS). Cell fractionation and immunofluorescence analyses showed that Flag-SIRT4-mNLS, unlike the WT SIRT4, was unable to translocate into the nucleus upon TGF-β1 treatment **(Fig. 8K, L)**. These results indicate that the NLS in SIRT4 is essential for TGF-β1-induced nuclear translocation of SIRT4. Importin α functions as an adaptor

and links NLS-containing proteins to importin β, which then docks the ternary complex at the nuclear-pore complex to facilitate the translocation of these proteins across the nuclear envelope[39,40]. Six importin α family members (α1, α3, α4, α5, α6 and α7) have been identified in humans[40]. We found that the endogenous SIRT4 only binds importin α1 **(Fig. 8M)**. Importin a1/Rch1 is barely detectable in the glomeruli of normal SD rat kidneys but is highly expressed in tubular cells, and the importin α1/Rch1 staining is significantly enhanced in the kidneys of diabetic rats[41]. Depletion of importin α1 with *Rch1* (coding for importin α1) shRNA **(Fig. 8N)** largely blocked the TGF-β1-induced nuclear translocation of SIRT4 and resulted in the accumulation of phosphorylated SIRT4 in the cytosol **(Fig. 8N)**. *In vivo*, U0126 restrained the *Sirt4*^{OE}-induced nuclear translocation of SIRT4 **(Fig. S3A, B)**. Furthermore, SIRT4 S36A or SIRT4 mNLS overexpression failed to accumulate SIRT4 in the nucleus and aggravated renal fibrosis in S4KO UUO mice **(Fig. 8O–R)**. Taken together, these results strongly suggest that TGF-β1-induced SIRT4 nuclear translocation is mediated by SIRT4 phosphorylation at S36, which is regulated by the ERK1/2 signaling pathway.

3.13. Exosomes containing anti-SIRT4 antibodies alleviate renal fibrosis in UUO mice

SIRT4 is necessary for maintaining mitochondrial function[8,42]. Since knockout or complete inhibition of SIRT4 may not be an advisable therapeutic strategy, we constructed exosomes containing anti-SIRT4 antibodies (αSRIT4) to treat UUO mice. Exosomes containing αSRIT4 effectively inhibited renal fibrosis in UUO mice, which was accompanied by the inhibition of SIRT4 expression in the nucleus, with little effect on SIRT4 content in mitochondria **(Fig. S4A−D)**.

Reportedly, the initial stage of the canonical Wnt signaling pathway in which SIRT4 translocates from mitochondria into the cytoplasm leads to β-catenin protein accumulation[34]. To determine whether the β-catenin accumulation is involved in the SIRT4-mediated kidney fibrosis *in vivo*, we generated mice expressing SIRT4^{OE} TECs and subjected them to UUO surgery with or without MSAB treatment (MSAB binds

to β-catenin and promotes its degradation). Our results showed that SIRT4 OE remarkably aggravated renal fibrosis under MSAB treatment **(Fig. S5A–C)**. Moreover, downregulation of β-catenin accumulation by MSAB can inhibit renal fibrosis in WT mice, but not the SIRT4^{OE} mice **(Fig. S5A–C).** These findings suggest that SIRT4-mediated the pathogenesis of renal fibrosis is independent of β-catenin accumulation.

4. Discussion

Renal fibrosis, especially tubulointerstitial fibrosis, is an inevitable common pathway of progressive chronic kidney disease[43,44]. However, there is a lack information regarding the pathogenesis of renal fibrosis, which hampers the development of effective therapeutics[45]. Here, we demonstrate that the nuclear translocation of SIRT4 is a prime initiator of kidney fibrosis. SIRT4 significantly accumulates in the nucleus during fibrosis following obstructed nephropathy and renal ischemia reperfusion injury. Global knockout or target deletion of *Sirt4* in TECs attenuated UUO-induced kidney fibrosis, whereas TEC-specific SIRT4^{OE} aggravated the fibrosis. Mechanistically, we found that TGF-β1 promoted SIRT4 release from mitochondria through the BAX/BAK pore. Furthermore, TGF-β1 activation resulted in the nuclear translocation of SIRT4, which was mediated by the ERK1/2-dependent phosphorylation of SIRT4 at S36, and consequently the binding of SIRT4 to importin α1. Nuclear SIRT4 deacetylates U2AF2 and promotes U2 snRNP formation, which promoting the *Ccn2* pre-mRNA splicing, ultimately leading to the increased CCN2 expression. In vivo, SIRT4 S36 or NLS mutants blocked SIRT4^{OE}-aggravated kidney fibrosis in UUO mice, which implied that SIRT4 nuclear translocation plays a significant role in the progression of kidney fibrosis.

Acetyl-CoA is an important energy-rich metabolite for homeostasis. In normal states, abundant acetyl-CoA in the cytosol shuttles freely in the nucleus or mitochondria to modulate the acetylation of histone or non-histone proteins. A rapid reduction in total acetyl-CoA levels in renal cells is observed after TGF-β1 stimulation[46]. Non-enzymatic acetylation levels are strongly reduced by acetyl-CoA.

Under such circumstances, few exceptional hyperacetylated proteins can function as

key regulators of kidney fibrosis development. In accordance with this hypothesis, we

found that U2AF2 acetylation decreased and U2 snRNP formation increased after

deacetylation of U2AF2 (**Fig. 4H–K**). Strikingly, U2AF2-K413, the deacetylation

form, promotes the pre-mRNA splicing and expression of *Ccn2* (**Fig. 6C, D**),

supporting the idea that acetylation of U2AF2 is responsible for fibrotic reaction

under TGF-β1. Additionally, a recent study showed that U2AF2 can directly bind and

stabilize circNCAPG, which participates in the nuclear translocation of ras responsive

element binding protein 1, thereby activating the TGF-β pathway and promoting

glioma progression[47]. Taking this into consideration, we speculate that U2AF2 may be

a positive feedback regulator of TGF-β1, although further investigations are required

to ascertain this.

SIRT4 is mainly located in the mitochondria and participates in various

mitochondrial metabolic processes[13]. Some studies have revealed a potential role of

SIRT4 in fibrosis[48,49]. In heart, loss of SIRT4 has been found to result in the

development of fibrosis. These studies indicated the protective role of SIRT4 in

mitochondria. In recent research, the authors found that SIRT4 abolish can ameliorate

CCl4-induced hepatic encephalopathy phenotypes, which was mediated by

downregulating detoxify ammonia through the urea cycle[50]. In our study, we indicated

that SIRT4 translocates from the mitochondria to the cytoplasm, which was caused by

TGF-β induced mitochondria damage (**Fig. 7**). Furthermore, the cytoplasm SIRT4

was phosphorylated by ERK2, which is a downstream of TGF-β signaling (**Fig.**

8A−J). The phosphorylated SIRT4 further translocated to the nuclear for promoting

CCN2 expression by regulating the alternative splicing, then accelerated the kidney

fibrosis (**Fig. 5**). Some studies have shown that SIRT4 exerts a protective effect on

podocytes. SIRT4 OE prevents glucose-induced podocyte apoptosis and ROS

production, thereby alleviating diabetic kidney disease (DKD)[51]. Furthermore,

FOXM1 transcriptionally activates SIRT4 and inhibits NF-κB signaling and the

expression of the NLRP3 inflammasome to alleviate kidney injury and podocyte

pyroptosis in DKD[52]. In the present study, we suggest a profibrotic role of SIRT4 in

TECs, which was contributed by upregulated expression of CCN2. Although these are complicatedly related to one another in CKD pathophysiology, injury to TECs is considered a core element that initiates progressive fibrosis[53]. Therefore, we indicated that it is SRIT4 may perform different role in different kinds of cells or subcellular organelles. Moreover, further studies on the role of SIRT4 in DKD are needed to evaluate the safety of anti-SIRT4 therapy.

Overall, our study reveals that TGF-β1 activation resulted in the nuclear translocation of SIRT4, mediated by the ERK1/2-dependent phosphorylation of SIRT4 at S36, and consequently the binding of SIRT4 to importin α1. In nucleus, SIRT4-mediated U2AF2 deacetylation at K413, a key protein for the spliceosome, acts as a responder under TGF-β1 stimulation. SIRT4 promotes CCN2 expression through alternative pre-mRNA splicing by deacetylation U2AF2, which contributes to the progression of kidney fibrosis. These findings expand the field of epigenetic regulation of fibrogenic gene expression and provide a potential therapeutic target for kidney fibrosis.

Non-standard abbreviations:

Cellular Communication Network Factor 2, CCN2; Tubular epithelial cells, TECs; Sirtuins, Sirts; Diabetes kidney disease, DKD; U2 Small Nuclear RNA Auxiliary Factor 1 or 2, U2AF1/U2AF2; Mitochondrial sirtuin 4 tripartite abundance reporters, mito-STAR; *Sirt4* knockout, S4KO; Unilateral ureteral occlusion, UUO; Unilateral renal ischemia-reperfusion injury, uIRI; Folic acid, FA; Neutrophil gelatinase-associated lipocalin, Ngal; Kidney injury molecule 1, Kim-1; Mouse podocytes, MPC; Mouse glomerular endothelial cells, GEC; Mouse fibroblast , MF; Tubule epithelial cells, TEC; Overexpression, OE; Rapid immunoprecipitation and mass spectrometry of endogenous proteins, RIME; Wild type, WT; endoplasmic reticulum, ER; Small nuclear ribonucleoproteins, snRNPs; Histone Acetyltransferase, HAT; Common docking, CD

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China (82370876 to **Shu Yang**, 82170842 and 82371572 to **Zhen Liang**, 82171556 to **Lin Kang**), Shenzhen Sustainable Development Science and Technology Special Project, China (No. KCXFZ20201221173600001 to **Zhen Liang**), Key Program Topics of Shenzhen Basic Research, China (No. JCYJ20220818102605013 to **Lin Kang**). Sequencing service was provided by Bioyi Biotechnology Co., Ltd. Wuhan, China.

Contribution statement

Guangyan Yang, **Jiaqing Xiang, Lixing Li,** and **Yanchun Li** conducted experiments. **Xiaoxiao Yang** contributed to the acquisition of data, analysis and interpretation of data. **Shu Yang** and **Zhen Liang** drafted the work or revised it critically for important intellectual content. **Lin Kang and Xiaoxiao Yang** analysed the data and revised the article critically for important intellectual content. **Shu Yang** contributed to the conception and the study design. All authors gave their approval of the version to be published.

Conflict of interest

The authors have declared that no conflict of interest exists.

Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

Reference

1   Gupta, S., Clarkson, M., Duggan, J. & Brady, H. Connective tissue growth factor: potential role in glomerulosclerosis and tubulointerstitial fibrosis. *Kidney international* **58**, 1389-1399, doi:10.1046/j.1523-1755.2000.00301.x (2000).

2   Kinashi, H. *et al.* Connective tissue growth factor regulates fibrosis-associated renal lymphangiogenesis. *Kidney international* **92**, 850-863, doi:10.1016/j.kint.2017.03.029 (2017).

3   Tampe, D. & Zeisberg, M. Potential approaches to reverse or repair renal fibrosis. *Nature reviews. Nephrology* **10**, 226-237, doi:10.1038/nrneph.2014.14 (2014).

4   Holmes, A. *et al.* CTGF and SMADs, maintenance of scleroderma phenotype is independent of SMAD signaling. *The Journal of biological chemistry* **276**, 10594-10601,

676   doi:10.1074/jbc.M010149200 (2001).

677 5  Morigi, M., Perico, L. & Benigni, A. Sirtuins in Renal Health and Disease. *Journal of the*
678   *American Society of Nephrology : JASN* **29**, 1799-1809, doi:10.1681/asn.2017111218 (2018).

679 6  Chalkiadaki, A. & Guarente, L. Sirtuins mediate mammalian metabolic responses to nutrient
680   availability. *Nature reviews. Endocrinology* **8**, 287-296, doi:10.1038/nrendo.2011.225 (2012).

681 7  Baur, J., Ungvari, Z., Minor, R., Le Couteur, D. & de Cabo, R. Are sirtuins viable targets for
682   improving healthspan and lifespan? *Nature reviews. Drug discovery* **11**, 443-461,
683   doi:10.1038/nrd3738 (2012).

684 8  Haigis, M. *et al.* SIRT4 inhibits glutamate dehydrogenase and opposes the effects of calorie
685   restriction in pancreatic beta cells. *Cell* **126**, 941-954, doi:10.1016/j.cell.2006.06.057 (2006).

686 9  Mathias, R. *et al.* Sirtuin 4 is a lipoamidase regulating pyruvate dehydrogenase complex
687   activity. *Cell* **159**, 1615-1625, doi:10.1016/j.cell.2014.11.046 (2014).

688 10  Nasrin, N. *et al.* SIRT4 regulates fatty acid oxidation and mitochondrial gene expression in
689   liver and muscle cells. *The Journal of biological chemistry* **285**, 31995-32002,
690   doi:10.1074/jbc.M110.124164 (2010).

691 11  Kang, H. *et al.* Defective fatty acid oxidation in renal tubular epithelial cells has a key role in
692   kidney fibrosis development. *Nature medicine* **21**, 37-46, doi:10.1038/nm.3762 (2015).

693 12  Gu, M. *et al.* Protein phosphatase 2Acα modulates fatty acid oxidation and glycolysis to
694   determine tubular cell fate and kidney injury. *Kidney international* **102**, 321-336,
695   doi:10.1016/j.kint.2022.03.024 (2022).

696 13  Csibi, A. *et al.* The mTORC1 pathway stimulates glutamine metabolism and cell proliferation
697   by repressing SIRT4. *Cell* **153**, 840-854, doi:10.1016/j.cell.2013.04.023 (2013).

698 14  Stegen, S. *et al.* HIF-1α metabolically controls collagen synthesis and modification in
699   chondrocytes. *Nature* **565**, 511-515, doi:10.1038/s41586-019-0874-3 (2019).

700 15  Sazani, P. & Kole, R. Therapeutic potential of antisense oligonucleotides as modulators of
701   alternative splicing. *The Journal of clinical investigation* **112**, 481-486, doi:10.1172/jci19547
702   (2003).

703 16  Maniatis, T. & Tasic, B. Alternative pre-mRNA splicing and proteome expansion in metazoans.
704   *Nature* **418**, 236-243, doi:10.1038/418236a (2002).

705 17  Zorio, D. & Blumenthal, T. Both subunits of U2AF recognize the 3' splice site in Caenorhabditis
706   elegans. *Nature* **402**, 835-838, doi:10.1038/45597 (1999).

707 18  Wu, S., Romfo, C., Nilsen, T. & Green, M. Functional recognition of the 3' splice site AG by the
708   splicing factor U2AF35. *Nature* **402**, 832-835, doi:10.1038/45590 (1999).

709 19  Merendino, L., Guth, S., Bilbao, D., Martínez, C. & Valcárcel, J. Inhibition of msl-2 splicing by
710   Sex-lethal reveals interaction between U2AF35 and the 3' splice site AG. *Nature* **402**, 838-841,
711   doi:10.1038/45602 (1999).

712 20  Agrawal, A. *et al.* An extended U2AF(65)-RNA-binding domain recognizes the 3' splice site
713   signal. *Nature communications* **7**, 10950, doi:10.1038/ncomms10950 (2016).

714 21  Will, C. *et al.* Characterization of novel SF3b and 17S U2 snRNP proteins, including a human
715   Prp5p homologue and an SF3b DEAD-box protein. *The EMBO journal* **21**, 4978-4988,
716   doi:10.1093/emboj/cdf480 (2002).

717 22  Ramadani-Muja, J. *et al.* Visualization of Sirtuin 4 Distribution between Mitochondria and the
718   Nucleus, Based on Bimolecular Fluorescence Self-Complementation. *Cells* **8**,
719   doi:10.3390/cells8121583 (2019).

23   Hastings, M., Allemand, E., Duelli, D., Myers, M. & Krainer, A. Control of pre-mRNA splicing by
the general splicing factors PUF60 and U2AF(65). *PloS one* **2**, e538,
doi:10.1371/journal.pone.0000538 (2007).

24   Schwer, B., North, B., Frye, R., Ott, M. & Verdin, E. The human silent information regulator
(Sir)2 homologue hSIRT3 is a mitochondrial nicotinamide adenine dinucleotide-dependent
deacetylase. *The Journal of cell biology* **158**, 647-657, doi:10.1083/jcb.200205057 (2002).

25   Ahuja, N. *et al.* Regulation of insulin secretion by SIRT4, a mitochondrial
ADP-ribosyltransferase. *The Journal of biological chemistry* **282**, 33583-33592,
doi:10.1074/jbc.M705488200 (2007).

26   Laurent, G. *et al.* SIRT4 coordinates the balance between lipid synthesis and catabolism by
repressing malonyl CoA decarboxylase. *Molecular cell* **50**, 686-698,
doi:10.1016/j.molcel.2013.05.012 (2013).

27   Huang, H. *et al.* Acetylation of SCFD1 regulates SNARE complex formation and
autophagosome-lysosome fusion. *Autophagy* **19**, 189-203,
doi:10.1080/15548627.2022.2064624 (2023).

28   Gu, L. *et al.* Stabilization of FASN by ACAT1-mediated GNPAT acetylation promotes lipid
metabolism and hepatocarcinogenesis. *Oncogene* **39**, 2437-2449,
doi:10.1038/s41388-020-1156-0 (2020).

29   Barber, M. *et al.* SIRT7 links H3K18 deacetylation to maintenance of oncogenic
transformation. *Nature* **487**, 114-118, doi:10.1038/nature11043 (2012).

30   Zhong, L. *et al.* The histone deacetylase Sirt6 regulates glucose homeostasis via Hif1alpha.
Cell **140**, 280-293, doi:10.1016/j.cell.2009.12.041 (2010).

31   Biancon, G. *et al.* Precision analysis of mutant U2AF1 activity reveals deployment of stress
granules in myeloid malignancies. *Molecular cell* **82**, 1107-1122.e1107,
doi:10.1016/j.molcel.2022.02.025 (2022).

32   Holmes, A. *et al.* Constitutive connective tissue growth factor expression in scleroderma
fibroblasts is dependent on Sp1. *The Journal of biological chemistry* **278**, 41728-41733,
doi:10.1074/jbc.M305019200 (2003).

33   Argmann, C. & Auwerx, J. Insulin secretion: SIRT4 gets in on the act. *Cell* **126**, 837-839,
doi:10.1016/j.cell.2006.08.031 (2006).

34   Wang, Y., Yue, J., Xiao, M., Lu, X. & Chin, Y. SIRT4-Catalyzed Deacetylation of Axin1 Modulates
the Wnt/β-Catenin Signaling Pathway. *Frontiers in oncology* **12**, 872444,
doi:10.3389/fonc.2022.872444 (2022).

35   Ma, K. *et al.* Dynamic PGAM5 multimers dephosphorylate BCL-xL or FUNDC1 to regulate
mitochondrial and cellular fate. *Cell death and differentiation* **27**, 1036-1051,
doi:10.1038/s41418-019-0396-4 (2020).

36   McArthur, K. *et al.* BAK/BAX macropores facilitate mitochondrial herniation and mtDNA efflux
during apoptosis. *Science (New York, N.Y.)* **359**, doi:10.1126/science.aao6047 (2018).

37   Lu, Z. & Xu, S. ERK1/2 MAP kinases in cell survival and apoptosis. *IUBMB life* **58**, 621-631,
doi:10.1080/15216540600957438 (2006).

38   Miyamoto, Y. *et al.* Differential modes of nuclear localization signal (NLS) recognition by three
distinct classes of NLS receptors. *The Journal of biological chemistry* **272**, 26375-26381,
doi:10.1074/jbc.272.42.26375 (1997).

39   Cingolani, G., Bednenko, J., Gillespie, M. & Gerace, L. Molecular basis for the recognition of a

764   nonclassical nuclear localization signal by importin beta. *Molecular cell* **10**, 1345-1353,
765   doi:10.1016/s1097-2765(02)00727-x (2002).

766   40   Goldfarb, D., Corbett, A., Mason, D., Harreman, M. & Adam, S. Importin alpha: a
767   multipurpose nuclear-transport receptor. *Trends in cell biology* **14**, 505-514,
768   doi:10.1016/j.tcb.2004.07.016 (2004).

769   41   Köhler, M. *et al.* Increased importin alpha protein expression in diabetic nephropathy. *Kidney*
770   *international* **60**, 2263-2273, doi:10.1046/j.1523-1755.2001.00069.x (2001).

771   42   Csibi, A. *et al.* The mTORC1 pathway stimulates glutamine metabolism and cell proliferation
772   by repressing SIRT4. *Cell* **184**, 2256, doi:10.1016/j.cell.2021.03.059 (2021).

773   43   Humphreys, B. Mechanisms of Renal Fibrosis. *Annual review of physiology* **80**, 309-326,
774   doi:10.1146/annurev-physiol-022516-034227 (2018).

775   44   Venkatachalam, M., Weinberg, J., Kriz, W. & Bidani, A. Failed Tubule Recovery, AKI-CKD
776   Transition, and Kidney Disease Progression. *Journal of the American Society of Nephrology :*
777   *JASN* **26**, 1765-1776, doi:10.1681/asn.2015010006 (2015).

778   45   Liu, Y. Renal fibrosis: new insights into the pathogenesis and therapeutics. *Kidney*
779   *international* **69**, 213-217, doi:10.1038/sj.ki.5000054 (2006).

780   46   Smith, E., Wigg, B., Holt, S. & Hewitson, T. TGF-β1 modifies histone acetylation and
781   acetyl-coenzyme A metabolism in renal myofibroblasts. *American journal of physiology. Renal*
782   *physiology*, doi:10.1152/ajprenal.00513.2018 (2019).

783   47   Li, H. *et al.* The U2AF65/circNCAPG/RREB1 feedback loop promotes malignant phenotypes of
784   glioma stem cells through activating the TGF-β pathway. *Cell death & disease* **14**, 23,
785   doi:10.1038/s41419-023-05556-y (2023).

786   48   Yin, X. *et al.* Targeting glutamine metabolism in hepatic stellate cells alleviates liver fibrosis.
787   *Cell death & disease* **13**, 955, doi:10.1038/s41419-022-05409-0 (2022).

788   49   Luo, Y. *et al.* SIRT4 accelerates Ang II-induced pathological cardiac hypertrophy by inhibiting
789   manganese superoxide dismutase activity. *European heart journal* **38**, 1389-1398,
790   doi:10.1093/eurheartj/ehw138 (2017).

791   50   Hu, S. *et al.* Amino acids downregulate SIRT4 to detoxify ammonia through the urea cycle.
792   *Nature metabolism* **5**, 626-641, doi:10.1038/s42255-023-00784-0 (2023).

793   51   Shi, J., Wang, Q., Li, H. & Huang, Q. SIRT4 overexpression protects against diabetic
794   nephropathy by inhibiting podocyte apoptosis. *Experimental and therapeutic medicine* **13**,
795   342-348, doi:10.3892/etm.2016.3938 (2017).

796   52   Xu, X. *et al.* FOXM1-activated SIRT4 inhibits NF-κB signaling and NLRP3 inflammasome to
797   alleviate kidney injury and podocyte pyroptosis in diabetic nephropathy. *Experimental cell*
798   *research* **408**, 112863, doi:10.1016/j.yexcr.2021.112863 (2021).

799   53   Friedman, S., Sheppard, D., Duffield, J. & Violette, S. Therapy for fibrotic diseases: nearing the
800   starting line. *Science translational medicine* **5**, 167sr161, doi:10.1126/scitranslmed.3004700
801   (2013).

808   **Figure and Figure legend**

809
Figure 1. Nucleus accumulation of SIRT4 was increases in renal tubules after
injury. (**A**) Western blots analysis of SIRT4, FN1, COL1A1, and Tubulin in the
kidney of UUO, uIRI, and sham mice (control group). (**B**) Nuclear fractions were
prepared from the kidney of UUO, uIRI, and sham mice. Nuclear PCNA and
cytoplasmic tubulin were used as controls. (**C, D**) Representative images of
immunohistochemical staining of SIRT4 (scale bar, 50 μm) in the kidneys from mice
that underwent sham surgery, UUO surgery on day 10 post surgery or uIRI surgery on
day 28 post surgery. (**E, F**) Representative images of Masson's trichrome staining (the
upper panel; scale bar = 50 μm) and SIRT4 immumohistochemical staining (the
bottom panel; scale bar = 20 μm) in the kidney sections from patients with CKD (n =
8) and minimal change disease (control group, n = 1).

Figure 2

Figure 2. Knockout of *Sirt4* or targeted deletion of *Sirt4* in renal TECs alleviates renal fibrosis induced by UUO or uIRI. (A to D) WT or S4KO mice were randomly assigned to sham or UUO surgery according to an established protocol. Kidney samples were obtained from mice on day 10 post UUO or sham surgery (n = 6 per group). A: Representative images of Masson's trichrome staining and Sirius red in kidneys from mice (scale bar = 100 μm). B: Western blot analysis of the expression of

SIRT4, CCN2, FN1, COL1A1, COL3A1, E-cadherin, α-SMA, and Tubulin in the
kidney of mice. C, D: The mRNA level of *Col1a1, Fn1, Eln, Ccn2, Acta2, Col3a1,*
Ngal, and *Kim-1* in the kidney of mice. **(E to H)** WT or S4KO mice received uIRI
surgery, and the sham surgery kidneys were used as control. Kidney samples collected
from mice after surgery at 28 d (n = 6 per group). E: Representative images of
Masson's trichrome staining and Sirius red in kidneys from mice (scale bar = 100 μm).
F: Western blot analysis of the expression of SIRT4, CCN2, FN1, COL1A1, COL3A1,
E-cadherin, α-SMA, and Tubulin in the kidney of mice. G, H: The mRNA level of
Col1a1, Fn1, Eln, Ccn2, Acta2, Col3a1, Ngal, and *Kim-1* in the kidney of mice. **(I to**
L) WT or S4TKO mice were randomly assigned to sham or UUO surgery according
to an established protocol. Kidney samples were obtained from mice on day 10 post
UUO or sham surgery (n = 6 per group). **I:** Representative images of Masson's
trichrome staining and Sirius red in kidneys from mice (scale bar = 100 μm). **J:**
Western blot analysis of the expression of SIRT4, CCN2, FN1, COL1A1, COL3A1,
α-SMA, and Tubulin in the kidney of mice. **K, L:** The mRNA level of *Col1a1, Fn1,*
Eln, Ccn2, Acta2, Col3a1, Ngal, and *Kim-1* in the kidney of mice. For all panels, data
are presented as mean ± SD. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ by one-way ANOVA
with Bonferroni correction test.
847

Figure 3

Figure 3. TECs specific SIRT4^OE aggravated kidney fibrosis. (A to I) AAV9-Ctrl
or AAV9-*Ksp-Sirt4* was injected into the kidneys of mice in situ at three independent
points. After 2-week transfection, the mice received UUO surgery, uIRI surgery, or FA
treatment (vehicle treatment as control) (n = 6 per group). A, D, G: Representative
images of Masson's trichrome staining and Sirius red in kidneys from mice (scale bar
= 100 μm). B, E, H: Western blot analysis of the expression of SIRT4, CCN2, FN1,
COL1A1, COL3A1, E-cadherin, α-SMA, and Tubulin in the kidney of mice. C, F, I:
The mRNA level of *Col3a1, Fn1, Eln, Ccn2, Acta2, Col1a1, Ngal*, and *Kim-1* in the
kidney of mice. For all panels, data are presented as mean ± SD. *$P < 0.05$, **$P < 0.01$,
***$P < 0.001$ by one-way ANOVA with Bonferroni correction test.

Figure 4

Figure 4. U2AF2 acetylation at K413 is increased under TGF-β1 stimulation. (A) Proteins interacted with SIRT4 in chromatin were identified by RIME. **(B)** The abundances (Normalized) of PUF60 and U2AF2 were shown. **(C)** Protein lysates from TECs were exposed to TGF-β1(left panel), kidney from the sham, uIRI, and UUO mice (middle panels), and WT TECs transfected with HA-*Sirt4* and/or Flag-*U2af2* and then treated with TGF-β1 (right panel), the protein lysates were subjected to Co-IP with anti-SIRT4 antibody, and determined protein expression using Western blotting by the indicated antibodies. **(D)** TECs isolated from WT or *U2af2⁻/⁻* mice and then treated with TGF-β1, cells lysates subjected to Co-IP with anti-SIRT4 antibody, and Western blotting using indicated antibodies. **(E)** WT TECs transfected with Ad-*Sirt4* or Ad-null as indicated, and cells lysates subjected to Co-IP with anti-U2AF2 antibody, and western blotting using indicated antibodies. **(F)** HK2 cells

 (human renal TECs) transfected with Flag- U2AF2 WT, K70R, K276R, K272R,
 K292R, K413R, K453R, or K462R under Ac-CoA treatment, cells lysates subjected to
 Co-IP with anti-Flag antibody, and Western blotting using indicated antibodies. **(G)**
 The conservation of U2AF2 lysine 413 in different species. **(H)** TECs transfected with
 Flag-*U2af2* WT under TGF-β1 stimulation, cells lysates subjected to Co-IP with
 anti-Flag antibody, and western blotting using indicated antibodies (left panel). HK2
 cells transfected with Flag-*U2AF2* WT or K413R with or without TGF-β1, cells
 lysates subjected to Co-IP with anti-Flag antibody, and western blotting using
 indicated antibodies (right panel). **(I)** Kidney lysates subjected to Co-IP with
 anti-U2AF2 antibody in uIRI, and UUO mice, and western blotting using indicated
 antibodies (middle and right panels). **(J)** WT TECs transfected with Flag-*U2af2* WT
 and then under serum starvation, Tunicamycin, Poly (I:C), and Cisplatin stimulation.
 Cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using
 indicated antibodies. **(K)** HK2 cells transfected with Flag-*U2AF2* WT or K413R and
 cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using
 indicated antibodies (left and middle panels). HK2 cells transfected with Flag-*U2AF2*
 WT or K413R with or without RNase stimulation, cells lysates subjected to Co-IP
 with anti-Flag antibody, and western blotting using indicated antibodies (right panel).
 (L) Schematic representation of components of SF3B complex performing pre-mRNA
 BPS recognition. **(M)** WT TECs transfected with Flag-*U2af2* WT with CBP, p300,
 HAT1, P/CAF, KAT5, or KAT8, cells lysates subjected to Co-IP with anti-Flag
 antibody, and Western blotting using indicated antibodies. **(N)** Whole-kidney lysates
 were immunoprecipitated with anti-U2AF2 antibody, and precipitated proteins were
 detected by indicated antibodies (the upper panel). WT TECs was transfected with
 HA-*Hat1* and/or Flag-*U2af2*. Cells lysates subjected to Co-IP with anti-Flag antibody,
 and Western blotting using indicated antibodies. **(O)** WT TECs was transfected with
 Flag-*U2af2* WT with Ad-sh*Hat1* or Ad-*Hat1*. Cells lysates subjected to Co-IP with
 anti-Flag antibody, and Western blotting using indicated antibodies. **(P)** HK2 cells
 were transfected with Flag-*U2AF2* WT, Ad-*HAT1*, or Ad-*HAT1* E276Q as indicated in
 figure. Cells lysates subjected to Co-IP with anti-Flag antibody, and western blotting
 using indicated antibodies. **(Q)** WT TECs was transfected with Flag-*U2af2*, Ad-*Hat1*
 or Ad-sh*Hat1* under TGF-β1 stimulation. Cells lysates subjected to Co-IP with
 anti-Flag antibody, and western blotting using indicated antibodies.

Figure 5

Figure 5. U2AF2 deacetylation under TGF-β1 is SIRT4 dependent. (A) WT TECs were transfected with Flag-*U2af2* WT, Ad-*Sirt4* or Ad-sh*Sirt4* under Ac-CoA treatment. Cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using indicated antibodies. **(B)** WT TECs were transfected with Flag-*U2af2* WT and HA-*Sirt4* under TGF-β1 stimulation. Cells lysates subjected to Co-IP with anti-HA antibody and Western blotting using indicated antibodies. **(C)** WT TECs were transfected with Ad-sh*Sirt4* or Ad-*Sirt4* with or without TGF-β1 stimulation. Cells lysates subjected to Co-IP with anti-U2AF2 antibody, and Western blotting using indicated antibodies. **(D)** HK2 cells were transfected with Flag-*U2AF2* WT or K413R, Ad-sh*SIRT4* under TGF-β1 stimulation. Cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using indicated antibodies. **(E)** HK2 cells

were transfected with HA-*SIRT4* WT or H161Y and Flag-*U2AF2* under TGF-β1 treatment. Cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using indicated antibodies. **(F)** SIRT4-U2AF2 docking with the HDOCK server. High magnification of boxed areas is presented on the left. Arrow indicates K413 of U2AF2 protein. **(G, H, K)** WT, S4KO, U2AF2$^{-/-}$ or DKO mice were randomly assigned to UUO surgery according to an established protocol. Kidney samples were obtained from mice on day 10 post surgery or sham surgery (n = 6 per group). G: Western blot analysis of the expression of SIRT4, U2AF2, CCN2, FN1, COL1A1, COL3A1, and Tubulin in the kidney from mice. Kidney lysates subjected to Co-IP with anti-U2AF2 antibody, and Western blotting using indicated antibodies. H: Representative images of Masson's trichrome staining and Sirius red in kidneys from mice (scale bar, 100 μm). K: The mRNA level of *Col1a1, Fn1, Eln, Ccn2, Acta2* and *Col3a1* in the kidney of mice. **(I, J, L)** *U2af2*$^{-/-}$ mice received *in situ* renal injection of AAV9-*Ksp-U2af2 WT* (*wtU2af2*), *wtU2af2* plus *Sirt4,* AAV9-*Ksp-U2af2 K413R* (*mutU2af2*), *K413R* plus *Sirt4* at 6 weeks of age. After 2 weeks, the mice were randomly assigned to sham surgery or UUO surgery according to an established protocol. Kidney samples were obtained from mice on day 10 post surgery or sham surgery (n = 6 per group). I: Western blot analysis of the expression of SIRT4, U2AF2, CCN2, FN1, COL1A1, COL3A1, and Tubulin in the kidney from mice. Kidney lysates subjected to Co-IP with anti-U2AF2 antibody, and Western blotting using indicated antibodies. J: Representative images of Masson's trichrome staining and Sirius red in kidneys from mice (scale bar, 100 μm). L: The mRNA level of *Col1a1, Fn1, Eln, Ccn2, Acta2 and Col3a1* in the kidney of mice. For all panels, data are presented as mean ± SD. $^*P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$ by one-way ANOVA with Bonferroni correction test.

Figure 6

Figure 6. Acetylation of U2AF2 at K413 regulates CCN2 gene expression and pre-mRNA splicing. (**A**) The crosstalk of differences of the gene expression profiling analysis and splicing analysis. (**B**) The *Ccn2* mRNA level in HK2 cells (n = 5). (**C, D**) HK2 cells were transfected with Flag-*U2AF2* WT, K413R, K413Q or K453Q, and Ad-*SIRT4* or *SIRT4* siRNA (si*SIRT4*) as indicated in figure. Cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using indicated antibodies. (**E**) Luciferase activity of *Ccn2* promoter in HK2 cells transfected with Ad-*SIRT4*, Ad-sh*SIRT4*, WT *U2AF2, U2AF2* K413R, or *U2AF2* K413Q under TGF-β1 stimulation was measured (n = 5). (**F**) Schematics of *Ccn2* alternative splicing pattern regulated by hyperacetylated U2AF2. Full line and dotted line represent normal splicing and abnormal intron retention, respectively. Grey and yellow boxes represent untranslated region (UTR) and protein coding region (CDS), respectively. The first stop codon in intron 2 or 3 was indicated with red arrow. (**G, H**) Quantitative real-time PCR analysis of normal or abnormal splicing isoforms of *CCN2* (n = 5). (**I**) Actinomycin D (Act D, 2 mg/mL) treatment and quantitative real-time PCR were performed to measure *CCN2* mRNA levels (n = 5). (**J**) HK2 cells were transfected with Flag-*U2AF2* WT, Flag-*U2AF2* WT, or Flag-*U2AF2* K413R with or without Ad-*SIRT4* under TGF-β1 stimulation. Cells lysates subjected to RNA Binding Protein Immunoprecipitation (RIP) with anti-U2AF2 antibody (left panel). Quantitative real-time PCR results of RIP assays (right panel). For all panels, data are presented as mean ± SD. **$P < 0.01$, ***$P < 0.001$, by Student's t-test for B. **$P < 0.01$, ***$P <$

967   0.001 by one-way ANOVA with Bonferroni correction test for E, G, H, I and J.

Figure 7. SIRT4 is released through BAX/BAK pore in a TGF-β1-dependent
manner. (A) Organelle separation experiment and immunoblot analysis detected the
expression/localization of SIRT4, Tubulin, and COX IV. **(B)** TECs were treated with
TGF-β1 for indicated time. Organelle separation experiment and immunoblot analysis
detected the localization of SIRT4, Tubulin, and COX IV. **(C)** Representative image
of immunofluorescent staining of SIRT4 in TECs treated with TGF-β1 or not (scale
bar, 50 μm). **(D)** TECs were treated with Tunicamycin, Cisplatin, Poly(I:C), or under
serum starvation. Organelle separation experiment and immunoblot analysis detected
the localization of SIRT4, Tubulin, and COX IV. **(E)** WT TECs incubated with *Bax*
siRNA or *Bak* siRNA under TGF-β1 stimulation. Organelle separation experiment and
immunoblot analysis detected the localization of SIRT4, Tubulin, and COX IV (left
panels). Western blot analysis of the expression of BAX, BAK and Tubulin in TECs
(middle panels). The mRNA level of *Ccn2* was detected by qPCR (n = 5) (right
panels). **(F)** WT TECs were treated with MSN-125 (10 μM) or vehicle. Organelle
separation experiment and immunoblot analysis detected the localization of SIRT4,
Tubulin, and COX IV (left panel). The mRNA level of *Ccn2* was detected by qPCR (n
= 5) (right panel). **(G)** Kidney tissues from WT or S4KO mice were treated with
MSN-125 or vehicle. The mRNA level of *Ccn2* was detected by qPCR in the kidney
of mice (n = 5). For all panels, data are presented as mean ± SD. ****P < 0.01, ***P <**
0.001 by one-way ANOVA with Bonferroni correction test.

Figure 8. ERK2 phosphorylates SIRT4 at S36 and promotes it nucleus translocation. (A) (The upper panel) Nuclear fractions were prepared from TECs pretreated with LY290042 (30 μM), SU6656 (4 μM), SP600125 (25 μM), and U0126 (20 μM) for 30 min before TGF-β1 (2 ng/ml) for 12 h. Nuclear PCNA and cytoplasmic Tubulin were used as controls. (The bottom panel) TECs were treated with TGF-β1 (2 ng/ml) for 12 h. Organelle separation experiment and immunoblot analysis detected the localization of SIRT4, PCNA, and Tubulin. **(B)** TECs were pretreated with or without U0126 (20 μM) for 30 min, then treated with TGF-β1 (2 ng/ml) for 12 h. Immunofluorescence analyses were performed with the indicated antibodies. **(C)** HK2 cells were stably transfected with Flag-ERK2, Flag-ERK2 K52R, or siRNA ERK2 (left panel) or transiently transfected with HA-MEK1 Q56P and indicated Flag-tagged ERK2 proteins (right panel). The cells were treated with or without TGF-β1 (2 ng/ml) for 12 h, and the total cell lysates, cytosol and nuclear fractions were prepared for determination of indicated proteins by Western blot. **(D)** TECs were treated with TGF-β1, and cells lysates subjected to Co-IP with anti-SIRT4

antibody, and Western blotting using indicated antibodies. **(E)** HK2 cells transfected with vectors expressing the indicated Flag-tagged ERK proteins were treated with or without TGF-β1 (2 ng/ml) for 12 h. **(F)** SIRT4-ERK2 docking with the HDOCK server. **(G)** HK2 cells expressing the indicated Flag-*SIRT4* proteins were treated with or without TGF-β1 (2 ng/ml) for 12 h. **(H, I)** HK2 cells transfected with Flag-Sirt4 WT or S36A and then incubated with TGF-β1 and/or U0126 as indicated in figure, the total cell lysates and nuclear fractions were prepared. Total cells lysates subjected to Co-IP with anti-Flag antibody, and Western blotting using indicated antibodies. **(J)** HK2 cells were transfected with MEK1 Q56P, Flag-ERK2 WT or K52R, and ERK2 siRNA as indicated in figure, and the total cell lysates and nuclear fractions were prepared. Total cells lysates subjected to Co-IP with SIRT4 antibody and Western blotting using indicated antibodies. **(K, L)** HK2 cells transfected with Flag-*Sirt4* WT or mNLS with or without TGF-β1 treatment, and the total cell lysates, cytosol, and nuclear fractions were prepared. Western blotting using indicated antibodies (K). Immunofluorescence analyses were performed with the indicated antibodies (L). **(M)** The indicated Flag-importin α proteins were transfected in HK2 cells, then treated with TGF-β1 (2 ng/ml) for 12 h. Flag-importin α proteins were immunoprecipitated with an anti-Flag antibody. **(N)** HK2 cells transfected with importin α1 were treated with or without TGF-β1 (2 ng/ml) for 12 h. The total cell lysates, cytosolic and nuclear fractions were prepared and Western blotting using indicated antibodies. **(O to R)** S4KO mice received *in situ* renal injection of AAV9-*Ksp-Sirt4 S36A* (S36A) or AAV9-*Ksp-Sirt4 mNLS* (mNLS) at 6 weeks of age. After 2 weeks, the mice were randomly assigned to sham surgery or UUO surgery according to an established protocol. O: Western blot analysis of the expression of SIRT4, CCN2, FN1, COL3A1, E-cadherin, α-SMA and Tubulin in mouse kidney. P: Representative images of Masson's trichrome staining and Sirius red staining in kidney sections from mice (scale bar, 100 μm). Q: The mRNA level of *Col1a1, Fn1, Eln, Ccn2 and Acta2* in the kidney of mice. R: The total cell lysates and nuclear fractions were prepared from kidney and Western blotting using indicated antibodies (left panel). (Right panel) Representative images of immumohistochemical staining of SIRT4 in kidneys from mice (scale bar, 100 μm). For all panels, data are presented as mean ± SD. ***$P < 0.001$ by one-way ANOVA with Bonferroni correction test.

        Electronic supplementary material

        1. ESM Figure and Figure legend

1053

Figure S1. Knockout of SIRT4 aggravated FA-induced kidney fibrosis. (A to D)
WT or S4KO mice were randomly assigned to FA or vehicle according to an
established protocol. Kidney samples were obtained from mice on day 14 post FA or
vehicle treatment (n = 6 per group). A: Western blot analysis of the expression of
SIRT4, CCN2, FN1, COL1A1, COL3A1, E-cadherin, α-SMA, and Tubulin in the
kidney of mice. B: Representative images of Masson's trichrome staining and Sirius
red in kidneys from mice (scale bar, 100 μm). C, D: The mRNA level of *Col1a1, Fn1,*
Eln, Ccn2, Acta2, Col3a1, Ngal, and *Kim-1* in the kidney of mice. **(E)** RT-PCR (n = 5)
and western blot analysis (n = 3) of the expression of SIRT4 in selected mouse renal
cells including mouse podocytes (MPC), glomerular endothelial cells (GEC), mouse
TECs and mouse fibroblast (MF). **(F to H)** Control or S4FKO mice were randomly
assigned to sham or UUO surgery according to an established protocol. F: Western
blot analysis of the expression of SIRT4, CCN2, FN1, COL1A1, COL3A1, α-SMA,
and Tubulin in the kidney of mice. G: The mRNA level of *Col1a1, Fn1, Eln, Ccn2,*

Acta2, *Col3a1, Ngal*, and *Kim-1* in the kidney of mice. H: Representative images of
Masson's trichrome staining and Sirius red in kidney sections of mice (scale bar, 100
μm). For all panels, data are presented as mean ± SD. *P < 0.05, **P < 0.01, ***P <
0.001 by one-way ANOVA with Bonferroni correction test. For all panels, data are
presented as mean ± SD. **P < 0.01, ***P < 0.001 by one-way ANOVA with
Bonferroni correction test.

Figure S2

1074

Figure S2. TECs targeted overexpression of SIRT4 reverses S4KO reduced
kidney fibrosis in UUO mice. (A to C) AAV9-*Ksp-Sirt4* was injected into kidneys of
mice *in situ* at three independent points in situ (AAV9-*Ksp*-null used as control). After
2-week transfection, the mice received UUO surgery (WT mice treated with
AAV9-Ctrl followed by UUO surgery as control) (n = 6 per group). A: Western blot
analysis of the expression of SIRT4, CCN2, FN1, COL1A1, COL3A1, E-cadherin,
α-SMA, and Tubulin in the kidney of mice. B: Representative images of Masson's
trichrome staining and Sirius red in kidney sections of mice (scale bar, 100 μm). C:
The mRNA level of *Col1a1, Fn1, Eln, Ccn2,* and *Acta2*. For all panels, data are
presented as mean ± SD. *P < 0.05, **P < 0.01, ***P < 0.001 by one-way ANOVA
with Bonferroni correction test.

1086

1087

Figure S3. U0126 prevents SIRT4 overexpression induced kidney fibrosis in
UUO mice. (A, B) AAV9-*Ksp-Sirt4* was injected into kidneys of mice *in situ* at three
independent points *in situ*. After 2-week transfection, the mice received UUO surgery,
and then mice were treated with U0126 (10 mg/kg body weight) or vehicle for 10
days (n = 6 per group). A: Organelle separation experiment and immunoblot analysis
detected the localization of SIRT4 and Tubulin in the kidneys of mice. B:
Representative images of immunohistochemical staining of SIRT4 (scale bar, 50 μm)
in the kidney sections of mice.

1096

Figure S4. Exosomes contain anti-SIRT4 antibody alleviated UUO-induced
kidney fibrosis. (A to D) After sham or UUO surgery, WT mice were treated with
exosomes null or contain anti-SIRT4 for 10 days (n = 6 per group). A: Western blot
analysis of the expression of CCN2, FN1, COL1A1, COL3A1, E-cadherin. α-SMA
and Tubulin in the kidney from mice. B: Representative images of Masson's
trichrome staining and Sirius red staining in kidneys sections of mice (scale bar, 100
μm). C: The mRNA level of *Col1a1, Fn1, Eln, Ccn2, Acta2*, and *Col3a1* in the kidney
of mice. D: Organelle separation experiment and immunoblot analysis detected the
localization of SIRT4 in the kidneys of mice. For all panels, data are presented as
mean ± SD. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ by one-way ANOVA with Bonferroni
correction test.

1108

Figure S5

1109

Figure S5. Inhibition of nucleus accumulation of β-catenin failed to suppresses
kidney fibrosis induced by SIRT4 overexpression in UUO mice. (A to C)
AAV9-*Ksp-Sirt4* or AAV-Ctrl was injected into kidneys of mice in situ at three
independent points in situ. After 2-week transfection, the mice received UUO surgery.
After UUO surgery, mice were treated with vehicle or MSAB for 10 days (n = 6 per
group). A: Western blot analysis of the expression of SIRT4, CCN2, FN1, COL3A1,
α-SMA, and Tubulin in the kidneys from mice. B: Representative images of Masson's
trichrome staining and Sirius red staining in kidney sections of mice (scale bar, 100
μm). C: The mRNA level of *Col1a1, Fn1, Eln, Ccn2, Acta2,* and *Col3a1* in the kidney
of mice. For all panels, data are presented as mean ± SD. ns: not significant difference,
*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ by one-way ANOVA with Bonferroni correction
test.

1122

2. ESM Methods

2.1 Protein extraction and Western blot analysis

Cytoplasmic or nuclear extracts were prepared from cells or kidneys using a

cytoplasmic and nuclear protein isolation kit (Cat# 78833, Thermo Fisher Scientific).

Briefly, the tube was vortexed vigorously in the highest setting for 15 s to fully

suspend the cell pellet (or tissues were homogenized in PBS). The tubes were

incubated on ice for 10 min. Next, ice-cold cytoplasmic extraction reagent II solution

was added to the tubes. The tube was vortexed and incubated on ice. The tube was

centrifuged for 5 min at maximum speed in a microcentrifuge (~16,000 × g), and the

supernatant (cytoplasmic extract) was transferred to a tube. The insoluble (pellet)

fraction, which contains nuclei, was suspended in ice-cold nuclear extraction reagent

and then vortexed at the highest setting for 15 s. The sample was placed on ice and

vortexed for 15 s every 10 min for 40 min. The tube was centrifuged at maximum

speed (~16,000 × g) in a microcentrifuge for 10 min, and the supernatant (nuclear

extract) fraction was transferred to a tube.

Mitochondria were extracted using a Mitochondria Isolation Kit (Sigma) following

the manufacturer's protocol. In brief, 2×10^7 cells were pelleted by centrifuging the

harvested cell suspension, and then mitochondria isolation reagent was added to the

cell pellets. The cell resuspension was centrifuged at 700 × g for 10 min at 4°C, and

then the supernatant was transferred to a new 1.5 ml tube and centrifuged at 12,000 ×

g for 15 min at 4°C. The supernatant (cytosolic fraction) was transferred to a new tube,

and the pellet contained the isolated mitochondria. The pellet was further lysed to

yield the final mitochondrial lysate. The extracted proteins were prepared for

subsequent western blotting analysis.

For western blot analysis, 50 µg of lysate was loaded onto sodium dodecyl

sulfate-polyacrylamide gel electrophoresis gels, and transferred onto polyvinylidene

difluoride membranes (Millipore). Proteins were analyzed with their corresponding

specific antibodies. Densitometry analysis was performed using Quantity One®

Software and quantified relative to the loading control, Tubulin.

1152

1153 **2.2 Antibodies used in Western blottings**

ANTIBODIES	SOURCE	IDENTIFIER
SIRT4 (1：1000)	Thermo Fisher Scientific	Cat# PA5-114377; RRID: AB_2787525
SF3B1 (1:1000)	Thermo Fisher Scientific	Cat# MA5-42938; RRID:AB_2912079
KAT5 (1:1000)	Thermo Fisher Scientific	Cat# PA5-34548; RRID:AB_2551900
SF3B5 (1:1000)	Thermo Fisher Scientific	Cat# A305-586A-T; RRID:AB_2782745
SF3B3 (1:1000)	Abcam	Cat# ab209402; RRID:AB_2910146
CCN2 (1:1000)	Abcam	Cat# ab6992; RRID:AB_305688
FN1 (1:1000)	Abcam	Cat# ab2413; RRID:AB_2262874
α-SMA (1:1000)	Abcam	Cat#ab7817; RRID:AB_262054
U2AF2 (1:50/1: 250)	Abcam	Cat# ab37530; RRID:AB_883336
Flag (1:50/1:1000)	Abcam	Cat# ab205606; RRID:AB_2916341
Anti-acetyl Lysine (1:1000)	Abcam	Cat# ab21623; RRID:AB_446436

PCAF (1:1000)	Abcam	Cat# ab96510; RRID:AB_10679933
KAT8 (1:1000)	Abcam	Cat# ab200660; RRID:AB_2891127
COX IV (1:1000)	Abcam	Cat# ab202554; RRID:AB_2861351
KPNA2 (1:1000)	Abcam	Cat# ab84440; RRID:AB_1860701
p300 (1:1000)	Abcam	Cat# ab275378; RRID:AB_2935873
HAT1 (1:1000)	Abcam	Cat# ab194296; RRID:AB_2801641
BAK	Abcam	Cat# ab104124; RRID:AB_10712355
COL1A1 (1:1000)	Proteintech	Cat# 67288-1-Ig; RRID:AB_2882554
HA-Tag (1:50/1:1000)	Cell signaling Technology	Cat# 3724; RRID:AB_1549585
Poly/Mono-ADPRibose (1:1000)	Cell signaling Technology	Cat# 83732; RRID:AB_2749858
CBP (1:1000)	Cell Signaling Technology	Cat# 7389; RRID:AB_2616020
E-cadherin (1:1000)	Proteintech	Cat# 20874-1-AP; AB_10697811
Tubulin (1:1000)	Proteintech	Cat# 11224-1-AP; RRID:AB_2210206
PCNA (1:2000)	Novus	Cat# NB500-106; RRID:AB_2252058
PUF60 (1:1000)	Novus	Cat# NBP1-49906; RRID:AB_10012096
SF3B4 (1:1000)	Novus	Cat# NBP1-92692; RRID:AB_11025842
COL3A1 (1:1000)	Novus	Cat# NB600-594; RRID:AB_10001330
SF3B2 (1:1000)	Novus	Cat# NBP1-92380; RRID:AB_11031649
BAX (1:1000)	Novus	Cat# NBP1-28566; RRID:AB_1852819
ERK1/ERK2 (1:1000)	Novus	Cat# AF1576; RRID:AB_354872
p-ERK1/2 (1:1000)	Novus	Cat# AF1018; RRID:AB_354539
U2AF1（1:100/1:1000）	Novus	Cat# NBP1-32515; RRID:AB_2288123

1154

2.3 Histological analysis

1155

1156 Kidney tissue was embedded in paraffin and sliced into 5 μm thick serial sections

1157 using a paraffin slicer. For kidney histology, the paraffin sections were stained using

1158 Masson's trichrome and Sirius red staining kit (Solarbio & Technology, Beijing,

1159 China).

1160

2.4 Quantitative real-time PCR

1161

Total RNAs were extracted by Trizol (Invitrogen) and then dissolved in an appropriate

amount of RNase water. cDNA was obtained by a reverse transcription kit purchased

from TransGen Biotech (Beijing, China). qPCR was performed using the ABI

StepOnePlusTM Real-time PCR system (Applied Biosystem) with specific primers

(Table S2). The relative mRNA levels of target genes were analyzed using the $2^{-\Delta\Delta CT}$

method. Tubulin was used as a housekeeping gene for analysis.

1168

2.5 RIME

The procedure was performed as previously described. The hTECs (2×10^7) were

treated with DMSO or DMOG (1mM) for 8 h and cross-linked with 1% formaldehyde

for 8min and quenched by 0.125M glycine. Cells were harvested and the nuclear

fraction was extracted. After sonication, the cell lysate was immunoprecipitated with

bead-prebound SIRT4 antibody (NB100-479, Novus Biologicals). Precipitated

proteins in 30 μl of 100mM ammonium hydrogen carbonate were reduced in 2.5mM

dithiothreitol at 60 °C for 1 h, then alkylated with 5mM iodoacetamide in the dark at

room temperature for 15min. Proteins were digested with 20 ng/μl Trypsin/LysC

(Promega) at 37 °C overnight. Peptides were desalted on Oasis HLB μ-elution plates

(Waters), eluted with 65 % acetonitrile/0.1 % trifluoroacetic acid, dried by vacuum

centrifugation, and analyzed by liquid chromatography/tandem mass spectrometry

using a nano-Easy LC 1000 interfaced with an Orbitrap Fusion Lumos Tribrid Mass

Spectrometer (Thermo Fisher Scientific). Tandem mass spectra were extracted by

Proteome Discoverer version 2.3 (Thermo Fisher Scientific) and searched against the

SwissProt_Full_Synchronized_2018_08 database using Mascot version 2.6.2 (Matrix

Science). Mascot ".dat" files were compiled in Scaffold version 3 (Proteome Software)

to validate MS/MS-based peptide and protein identifications. Peptide identifications

were accepted if false discovery rate (FDR) was less than 1%, based on a

concatenated decoy database search by the Peptide Prophet algorithm with Scaffold

delta-mass correction.

1190

1191 **2.6 Co-immunoprecipitation (co-IP)**

1192 After treatment, the cells were lysed in an ice-cold co-IP buffer containing 20 mM

1193 Tris-HCl (pH 8.0), 100 mM NaCl, 1 mM EDTA, and 0.5% NP-40, supplemented with

1194 a protease inhibitor cocktail (Roche, 04693132001), for 30 min. The cell homogenates

were then centrifuged at 13,000g for 15 min, and the resulting supernatant was incubated overnight at 4 °C on a shaker with primary antibodies (anti-Flag, anti-SIRT4, anti-HA, anti-U2AF2 or anti-IgG). To ensure complete saturation of the primary antibodies, sufficient cell lysates were cultured and collected for IP. The mixture of antibodies and proteins was subsequently incubated with protein A/G-agarose beads (Thermo Fisher Scientific, Cat#: 78610) at 4 °C for 3 h. The beads were washed 5-6 times with cold IP buffer and resuspended in loading buffer. The cell lysates and immunoprecipitates were denatured in loading buffer at 95 °C for 5 min, and western blotting analysis was performed.

2.7 Protein-protein docking simulation

Visual protein-protein docking of SIRT4 and U2AF2 or ERK2 was performed online using HDOCK (http://hdock.phys.hust.edu.cn/).

2.8 Analysis of the conservation of lysine 413 in U2AF2

A gene panel of the NCBI database (https://www.ncbi.nlm.nih.gov/gene/) was used to download amino acid sequences of proteins from multiple species and subsequently compare sequence conservation between species using the UGENE software (version 39; the software can be downloaded from the following website: http://ugene.net/).

2.9 Bioinformatics methods predict acetylase of Lys413 in U2AF2

A Group-based Prediction System (GPS) online tool (http://pail.biocuckoo.org) was used to predict Lys413 in U2AF2 modified by histone acetyltransferases, with high score values suggesting better results. The amino acid sequence of the protein in FASTA format was downloaded from UniProt and uploaded to the GPS website.

2.10 Luciferase report assay for TGF-β1

Specific primers for the 5'-ACTCTCGAGCAGTGTTCCCACCCTGACAC-3' and for 5'-ACTAAGCTTGGTTGGCACTGCGGGCGGAG-3' were designed to amplify the DNA (-835/+214) in the CTGF promoter region of the human genome through the polymerase chain reaction (PCR) experiment. The sequence of the human CTGF

promoter was cloned into the pGL3.0-Basic vector (LMAI Bio; Cat# LM-1554) through molecular cloning. Positive plasmids were screened and identified by methods such as monoclonal colony PCR, plasmid double enzyme digestion identification, and sequencing. Then, using Lipofectamine 3000 transfection reagent, pGL3.0-Basic-CTGF and Renilla luciferase were transfected into HEK293T. Firefly and Renilla luciferase activities were then measured by a dual luciferase reporter gene system (Promega, Madison, WI, USA)

2.11 Cell culture, transfection

The human proximal tubular epithelial cell line (HK2 cells) (Cat# CRL-2190) were purchased from the American Type Culture Collection (Manassas, VA, USA). Cells were cultured in Minimum Essential Medium (Thermo Fisher Scientific, Shanghai, China; Cat# 10373017).   Mouse podocytes (MPC) (Cat# BNCC342021) and mouse renal glomerular endothelial cells (GEC) (Cat# BNCC360313) were purchased from BeNa Culture Collection (Beijing, China), cultured in DMEM-H complete medium (BeNa Culture Collection, Cat# BNCC338068). Mouse renal fibroblasts (MF) (Procell Life Science&Technology, Wuhan, china; Cat# CP-M069) were cultured in mouse kidney fibroblast complete culture medium (Procell, Cat#CM-M069). All kinds of media were supplemented with 10% FBS (Gibco, Grand Island, NY), 100 U·mL−1 of penicillin (Gibco), and 100 mg•mL−1 of streptomycin (Gibco). All cells were cultured in a 37 °C incubator containing 5% CO_2 and 95% air.Human embryonic kidney 293 cells (HEK293) were purchased from ATCC and maintained in HyClone Dulbecco's Modified Eagle Medium (SH30022, Cytiva) with 10 % FBS, 1 % glutamine, and 1 % penicillin/streptomycin solution in a humidified incubator supplemented with 95 % air/5 % CO_2 at 37 °C. Cells were regularly checked for mycoplasma in a standardized manner, by a qPCR test, performed under ISO17025 accreditation to ensure work was conducted in mycoplasma-negative cells.

Adenovirus particles used in the article were provided by GeneChem Company (Shanghai, China). Adenovirus was used to transfect HK2 cells or TECs for 6 h (adenovirus particles containing empty vectors were used as controls), and then

1256 replace the fresh medium and continue to culture. The commercial siRNA Ctrl (Cat#

1257 sc-37007, Santa Cruz), siRNA *Hat1* (Cat# sc-145898, Santa Cruz), siRNA *Sirt4* (Cat#

1258 sc-63025, Santa Cruz), siRNA *Bax* (Cat# sc-29213, Santa Cruz), and siRNA *Bak*

1259 (Cat# sc-29785, Santa Cruz) were transfected using Lipofectamine 3000 (Invitrogen,

1260 Carlsbad, California, USA) according to the manufacturer's instructions.

1261

2.12 Isolation of primary renal tubular epithelial cells (TECs)

Primary mouse renal tubular epithelial cells were isolated using an established

protocol we previously published[1]. Briefly, the cortex of the kidneys was carefully

dissected and chopped into small pieces. Then, 1 mg/ml of collagenase solution was

applied and incubated at 37 °C for 30 min with gentle agitation. The digestion was

terminated by FBS and then filtered sequentially. Fragmented tubules were collected

and maintained in a renal epithelial cell basal medium using a growth kit. The

medium was changed for the first time, after 72 h. The purity of the primary mouse

renal tubular epithelial cells was confirmed by immunofluorescence staining of

SGLT1 (Novus, NBP2-20338). Cells at passages 2-5 were used for the experiments.

1272

2.13 Immunofluorescent staining of cells

After treatment, cells was fixed for 15 min using fresh, methanol-free 4%

formaldehyde, and then, rinsed thrice with PBS for 5 min each. After blocking with

goat serum for 30 min, the cells were incubated with primary antibody against SIRT4

(Thermo Fisher Scientific, Cat# PA5-114377) at 4 °C overnight. Alexa Fluor®488

goat antibodies against murine IgG (Invitrogen, Shanghai, China; Cat# A-11078,

1:400) were included as secondary antibodies. As negative controls, the primary

antibodies were exchanged for nonimmune serum from the same species. The samples

were counterstained with DAPI for 15 min. The sections were sealed with a cover

glass, and the specimens were examined using the appropriate excitation wavelength.

Images were captured and processed with a Laica microscope (Wetzlar, Germany).

1284

2.14 Immunohistochemical (IHC) staining

Paraffin-embedded mouse kidney samples were sliced into 4 μm-thick sections and
subjected to IHC staining using a Rabbit two-step detection kit (Rabbit enhanced
polymer detection system, ZSGB Bio, Beijing, China). Briefly, mouse kidney sections
were deparaffinized, followed by antigen retrieval, treatment with peroxidase block,
and incubation with rabbit anti-SIRT4 (1:100, Thermo Fisher Scientific, Cat#
PA5-114377) primary antibody overnight at 4°C. Tissue sections were then washed
and incubated with the reaction-enhancing solution at room temperature for 20 min.
After washing three times with phosphate buffer saline (PBS), tissue sections were
treated with enhanced enzyme-labeled goat anti-rabbit IgG polymerase for 20 min and
then developed with 3,3' Diaminobenzidine solution, counterstained with hematoxylin,
and mounted with mounting medium. The kidney frozen slices from patients with
minimal change disease and DKD were fixed with 4 % paraformaldehyde for 20 min,
incubated with anti-SIRT4 primary antibody overnight at 4 °C, and subjected to
immunohistochemistry staining as described above. The stained area was measured
using ImageJ software.
1301
2.15 RNA immunoprecipitation (RIP)
RIP is a powerful method to investigate the interactions between RNA and RNA
binding proteins in vivo by immunoprecipitating RNA-protein complexes using
specific antibodies against RNA binding proteins. An example of the RIP protocol is
as follows: Covalently conjugate protein-specific antibody to the beads. Wash
extensively with lysis buffer. Collect cell lysate and incubate with antibody
conjugated beads allowing RNA-protein complexes formation. Use normal IgG as
negative control. Incubate the lysate for 2-4 hours at 4°C with gentle rotation. Wash
beads stringently with ice-cold lysis buffer containing RNase inhibitors, protease
inhibitors and nonionic detergents. Perform 4-6 quick washes. Elute RNA-protein
complexes from beads, generally with 1% SDS buffer heated to 95°C for 5min.
Extract RNA with phenol/chloroform. Isolate RNA for downstream applications like
RT-qPCR or RNA sequencing.
1315

2.16 The stability and intron retention index of CCN2 mRNA

To detect the CCN2 mRNA stability, the cells were treated with Actinomycin D (2 mg/mL) for 0 h, 1 h, 2 h and 3 h. The cDNA from different time point was diluted in nuclease-free water for the same times, and mRNA was quantified by qPCR. The fold of changing CCN2 level was normalized to the initial expression level of CCN2 (0 h). Intron Retention index = (mRNA level of normal splicing isoform/total CCN2 mRNA level)/ (mRNA level of abnormal splicing isoform/total CCN2 mRNA level). The CCN2 mRNA stability qPCR primers are as listed in Table S3.

2.17 Loading of Exosomes

The approaches for αSIRT4 incorporation into exosomes were executed as mentioned before[3]. Naive exosomes released from HEK293T were diluted in PBS to a concentration 0.15 mg/mL of total protein, then αSIRT4 solution in PBS (0.5 mg/mL) was added to 250 μl of exosomes to the final concentration 0.1 mg/mL total protein, and incubated at RT for 18 hours. In case of a saponin treatment, a mixture of αSIRT4 and exosomes was supplemented with 0.2% saponin and placed on shaker for 20 min at room temperature.

3. Table

Table S1. Clinical characteristics of CKD in biopsy samples, Related to Figure 1

Parameters	Mild CKD (n=4)	Severe CKD (n=4)	*P* value
Sex	Male:Female=3:1	Male:Female=2:2	NS
Age	36.3±7.8	47.5±17.5	NS
FBG (mmol/L)	6.57±1.69	5.82±0.81	NS
Proteinuria (g/24h)	2.49±2.01	10.96±7.95	0.042
eGFR (ml/min/1.73m^2)	84.42±27.49	38.875±12.94	0.012
Duration of disease (y)	4±2.3	5±4.8	NS

Data are calculated by One-way ANOVA and presented as mean ± SEM. The number of patients in each group was as indicated. NS: not significant difference, y: year.

Table S2. Primers used in qPCR

Gene	Forward Primer (5'→3')	Reverse Primer (5'→3')
m*Col1a1* (ID: 12842)	*GCTCCTCTTAGGGGCCACT*	*CCACGTCTCACCATTGGGG*
m*Col3a1* (ID: 12825)	*CTGTAACATGGAAACTGGGGAAA*	*CCATAGCTGAACTGAAAACCACC*

m*Fn1* (ID: 14268)	ATGTGGACCCCTCCTGATAGT	GCCCAGTGATTTCAGCAAAGG
m*Eln* (ID: 13717)	TTGCTGATCCTCTTGCTCAAC	GCCCCTGGATAATAGACTCCAC
m*Ccn2* (ID: 14219)	GGGCCTCTTCTGCGATTTC	ATCCAGGCAAGTGCATTGGTA
m*Acta2* (ID: 11475)	CCCAACTGGGACCACATGG	TACATGCGGGGGACATTGAAG
m*Ngal* (ID: 16819)	TGGCCCTGAGTGTCATGTG	CTCTTGTAGCTCATAGATGGTGC
m*Kim-1* (ID: 171283)	GTTAAACCAGAGATTCCCACACG	TCTCATGGGGACAAAATGTAGTG
m*β-actin* (ID:11461)	GGCTGTATTCCCCTCCATCG	CCAGTTGGTAACAATGCCATGT

1340

Table S3. Primers used in IR Index

CCN2 pre-mRNA	Forward Primer (5'→3')	Reverse Primer (5'→3')
In 2+ex 3	GCCTATTCTGTCACTTCGGC	CACTCCTCGCAGCATTTCC
In 3+ex 4	TCAGGGTCGTGATTCTCTCC	TCTCTTCCAGGTCAGCTTCG

1342
1343

Reference

1. Yang S, Xiang J, Ma C, Yang G, Wang X, Liu H, Fan G, Kang L and Liang Z. Sp1-like protein KLF13 acts as a negative feedback regulator of TGF-β signaling and fibrosis. *Cell reports*. 2023;42:112367.

2. Glowacki F, Savary G, Gnemmi V, Buob D, Van der Hauwaert C, Lo-Guidice J, Bouyé S, Hazzan M, Pottier N, Perrais M, Aubert S and Cauffiez C. Increased circulating miR-21 levels are associated with kidney fibrosis. *PloS one*. 2013;8:e58014.

3. Haney M, Klyachko N, Zhao Y, Gupta R, Plotnikova E, He Z, Patel T, Piroyan A, Sokolsky M, Kabanov A and Batrakova E. Exosomes as drug delivery vehicles for Parkinson's disease therapy. *Journal of controlled release : official journal of the Controlled Release Society*. 2015;207:18-30.

1353
1354