1

Engineering fluorescent reporters in human pluripotent cells

and strategies for live imaging human neurogenesis

Alwyn Dady*[1], Lindsay Davidson, Nicolas Loyer, Timothy Sanders[2],

Jens Januschke & Kate G. Storey*

Division of Molecular, Cell & Developmental Biology, School of Life Sciences,

University of Dundee, Dundee DD1 5EH, UK

* co-corresponding authors

[1] current position: Université Paris Est Créteil (UPEC), Glycobiology, Cell Growth and Tissue Repair Research Unit (Gly-CRRET), F-94010 Créteil, France

[2]University of California, San Francisco, Benioff Children's Hospital Oakland, USA

Running title: human PSC reporters for live imaging

Key words: human pluripotent cells, human spinal cord development, human cortical development, live cell imaging, piggyBac-mediated fluorescent reporters

Abstract

Investigation of cell behaviour and cell biological processes in human embryonic tissues is facilitated by creation of fluorescent reporters in human pluripotent stem cell lines, which can be differentiated into cell types of choice. Here we report use of a piggyBac transposon-mediated stable integration strategy to engineer human pluripotent stem cell reporter lines. These express a plasma membrane (pm) localised protein tagged with the fluorescent protein eGFP or mKate2, the photoconvertible nuclear marker H2B-mEos3.2, with or without pm-mKate2, and the cytoskeletal protein F-tractin tagged with mKate2. Focussing on neural development some of these lines were used to live image and quantify cell behaviours, including cell cycle progression and cell division orientation in spinal cord rosettes. Further, lipofection-mediated introduction of piggyBac constructs into human neural progenitors labelled single cells and small cell groups within rosettes, allowed monitoring of individual cell behaviours including neuronal delamination. Finally, using the F-tractin-mKate2 hiPSC line, actin dynamics were captured during proliferation in cortical neural rosettes. This study presents new tools and techniques with which to interrogate human cell behaviour and cell biology using live imaging approaches.

Introduction

Live imaging is used to monitor the behaviour of cells within tissues and to reveal sub-cellular proteins dynamics. This includes important cellular processes such as cell cycle progression and changes in cell shape and adhesion as individual cells differentiate and cells group together to form structures during tissue morphogenesis. Such approaches provide insight into developmental and tissue homeostasis mechanisms and elucidate changes in disease states. The ability to genetically engineer human pluripotent stem cells to express fluorescent proteins facilitates such live imaging approaches and allows high resolution investigation of cell behaviour and cell biological processes in human tissue models.

DNA constructs can be introduced into human pluripotent cells in numerous ways. The piggyBac transposon system has advantages compared to other stable integration approaches, particularly for live cell imaging studies. In comparison with viral vectors, such as adenovirus or lentivirus (Lee, et al., 2017), the piggyBac system is more secure and convenient to generate (Ding, et al., 2005, Wang, et al., 2008, Woltjen, et al., 2009). Viral integration sites are non-random but diverse and there is no control over copy number and gene expression levels, (Cereseto and Giacca, 2004, Engelman, 2003, Sultana, et al., 2017): indeed viral integrations can alter endogenous gene expression, including vital regulatory genes and/or inactivation of tumour-suppressor genes, impacting cell behaviour and differentiation (e.g. (Hacein-Bey-Abina, et al., 2010, Hacein-Bey-Abina, et al., 2003). In contrast, piggyBac constructs flanked by a 13bp perfect terminal invert repeat sequence insert specifically into P-elements (TTAA segments) in the genome (Elick, et al., 1996, Lee, et al., 2011, Liu, et al., 2013). Other transposon systems are also used including hAT-Tol2, (Kawakami, 2007, Kawakami and Shima, 1999) or Tcl-like transposons, *Sleeping Beauty* and *Frog Prince*, (Ivics, et al., 1997, Miskey, et al., 2003, Narayanavari, et al., 2017). However, while *Sleeping Beauty* and piggyBac have the highest transposition activity in mammalian cells (Kim and Pyykko, 2011), *Sleeping Beauty* is less reliable, with a tendency for local hopping and leaving a CAG footprint following excision (Wang, et al., 2008, Wu, et al., 2006).

In comparison with other stable integration approaches, Transcription Activator-like Effector Nucleases (TALENs) (Hockemeyer, et al., 2011, Joung and Sander, 2013, Miller, et al., 2011, Wood, et al., 2011) and CRISPR/Cas9 (Charpentier and Doudna, 2013, Doudna and Charpentier, 2014, Jinek, et al., 2012, Pacesa, et al., 2024), which achieve precision gene engineering of single copy insertions at a defined locus, the piggyBac transposon system will insert multiple copies of fluorescently tagged proteins within the many P-elements/ TTAA chromosome sites (Elick, et al., 1996, Lee, et al., 2011) (Liu, et al., 2013, Woodard and Wilson,

2015). This feature of the piggyBac system has a particular advantage for long term live-imaging

by increasing transgene copy number and expression, and so detection sensitivity for

fluorescently tagged proteins. This in turn reduces exposure times and thereby potential

phototoxicity.

Nevertheless, gene editing using CRISPR/Cas9 and other nucleases has been

successfully used to express fluorescently tagged endogenous proteins for live cell imaging

(Dambournet, et al., 2014) and these include hiPSC lines expressing, for example, mEGFP-

nuclear lamin B1 (Haupt, et al., 2018, Roberts, et al., 2017). This has the advantage of

endogenous regulatory control of fusion protein expression but does rely on this having

sufficient strength for live detection and for longer term live cell imaging. Fusion of an

endogenous protein with large exogenous sequences such as GFP may also compromise

protein function. Indeed, such approaches typically report low recovery of biallelic edits, which

may reflect lack of tolerance for two tagged copies of an indispensable protein, but this is likely

to be protein specific (Roberts, et al., 2017). Inclusion of an amino acid linker sequence

between the native and fluorescent protein can mitigate this risk. Such engineered hPSC lines

require quality controls, including confirmation of pluripotency, karyotype and growth rate as

well as comparison with endogenous protein localisation and cell behaviour in well-defined

differentiation assays.

Here we describe use of a piggyBac mediated engineering strategy to create a series of

human pluripotent stem cell reporter lines by introducing fluorescent proteins which localise

to the plasma-membrane, chromatin, or actin cytoskeleton. We characterise these lines and

deploy them, or lipofected piggyBac constructs introduced during differentiation, in live

imaging protocols to uncover fundamental characteristics of human neural progenitor cell

behaviour in *in vitro* assays for spinal cord and cerebral cortex development.

Results and Discussion

Generation of human pluripotent cell fluorescent reporter lines using the piggyBac-
transposon stable integration system

To facilitate analysis of cell behaviour in live tissue derived from human pluripotent stem cells

we generated a series of fluorescent reporter hESC or iPSC lines (Dady, et al., 2022)(Figures

1A,B). To monitor cell shape and dynamics we used a plasma membrane (pm) localised protein

tagged with eGFP or mKate2 (pm-eGFP or pm-mKate2) (Sanders, et al., 2013). To allow

monitoring of the cell nucleus we engineered a line expressing H2B-mEos3.2, the latter

localising in the nucleus and mEos3.2 having the potential for laser-mediated photo-conversion

(Zhang, et al., 2012) and so the labelling and tracking of individually selected cells. This provides

the potential for clonal analysis in combination with immunofluorescence for cell type specific

marker proteins. We also co-transfected H2B-mEos3.2 and pm-mKate2 expressing plasmids to

allow monitoring of labelled nuclei and changes in cell shape. Finally, to monitor actin

cytoskeletal dynamics we created an F-tractin-mKate2 cell line.

To generate these reporters, we used the hESC line H9 (for pm-eGFP, pm-mKate2,

H2B-mEos3.2), or hiPSC line ChiPS4 for F-Tractin-mKate2 and a further pm-eGFP line (Dady, et

al., 2022) and we deployed the piggyBac transposon-mediated stable integration strategy

(Sanders, et al., 2013, Sarkar, et al., 2003, Yusa, et al., 2009, Yusa, et al., 2011). The piggyBac

vectors are composed of the specific fusion protein of interest whose expression is driven by a

CAG promoter (Figures 1A,B). This cargo is then flanked by two Inverted Terminal Repeat (ITR)

sequences which allow a hyperactive piggyBac transposase (HyPBase, characterised by highly

stable enzymatic activity), to cut and paste the cargo within the specific TTAA-chromosomal

sites (Yusa, et al., 2011). Critical to efficient stable integration was transfection of 5 times more

piggyBac vector than transposase.

The cloning strategy for generating these constructs is described in the Methods. Once

complete these were introduced to a suspension of 1×10^6 hPSCs by electroporation with a final

concentration of 1.2 µg/µl of piggyBac constructs and HyPBase (Yusa, et al., 2011) (Figures 1C

Methods). Transfected hPSCs were then FAC-sorted and selected using their specific

fluorescence properties: this optimized protocol delivered a high rate of transfection (Figure

S1). Using this approach, polyclonal cell lines were generated for each fusion protein of

interest: these were composed of cells with this construct randomly inserted within each

genome. Polyclonal lines were sorted with easily detectable, but not high, fluorescence for live

imaging studies. We also made a monoclonal line for pm-eGFP in ChiPS4 for future use as this

approach would obviate the remote possibility of cell-to-cell variation impacting cell behaviours

captured in live imaging experiments (Figure 1D).

Using human pluripotent cell fluorescent reporter lines to monitor and quantify cell behaviour

in human spinal cord rosettes

These fluorescent reporter hPSC lines allowed the development of *in vitro* assays for monitoring

and quantifying human cell behaviour. Here we first focussed on capturing fundamental

parameters of human neurogenesis using our established spinal cord rosette assay (Dady, et

al., 2022, Verrier, et al., 2018)(Figure 2A). This involves differentiating hPSCs into Neuro-

Mesodermal Progenitor-like cells (NMP-like cells) and their subsequent differentiation into

SOX2 expressing neural progenitors, which self-organise into neural rosettes exhibiting

localised expression of apico-basal polarity markers such as NCAD/CDH2 by Day 6 (D6) following

NMP-L cell plating (Dady, et al., 2022). This differentiation assay generates neural crest

progenitors and dorsal neural progenitors, with the first neurons apparent from D8, peak

neurogenesis at D14 and gliogenesis onset by ~ D20. This sequence of events is in line with

those documented in the human embryonic spinal cord with dorsal neurogenesis at forelimb

level commencing by CS12/13 and gliogenesis by CS18 (Dady, et al., 2022, Rayon, et al., 2020).

The cell lines expressing pm-eGFP and pm-mKate with H2B-mEos3.2 lend themselves

to analysis of cell shape change. We wished to monitor the dynamics of neuroepithelial cell

behaviour as these cells progressed through the cell cycle during inter-kinetic nuclear migration

(IKNM)(Sauer, 1935, Spear and Erickson, 2012). This behaviour is characteristic of pseudo-

stratified epithelia and involves movement of the nucleus from the basal to the apical side of a

cell in G2 phase, subsequent mitosis at the apical (ventricular) surface and return to the basal

surface in G1: here a neuroepithelial cell may then either re-enter S-phase or exit the cell cycle

and commence neuronal differentiation. To monitor this behaviour in spinal cord rosettes we

first optimised our assay to address a major challenge of longer-term live imaging, increasing

thickness of the developing tissue over time (D6 rosettes are typically ~ 100μm). This reduces

the ability to maintain cells of interest in focus, even when deploying a long working-distance

lens and deconvolution algorithms (see Methods). To reduce the thickness of rosettes we

developed a re-plating regime in which D6 rosettes were manually micro-dissected and

replated on glass-bottom dishes suitable for microscopy (Figure 2B). This resulted in formation

of small satellite micro-rosettes (SmR) surrounding the replated primary rosette (PrR) (Figure

2C). Satellite micro-rosettes were polarised as indicated by apical actin localisation. Most

importantly, SmR were flatter (~ 50µm thick) allowing longer term high-resolution live cell

imaging.

Human spinal cord satellite rosette cells derived from hESCs expressing pm-eGFP or

pm-mKate and H2B-mEos3.2 were then imaged at intervals for 24h at D13-D14 and at D18-D19

(Figs. 2D-E, movie 1). Apical visiting time (defined as the period during which a basally located

nucleus transits to the apical surface, divides and returns to a basal position) and used as an

indication of changing cell cycle duration, increased from D13/D14 to D18/D19 and

approximately doubled over this 6-day period (D13, 11h ± 3h; D19, 25h ± 7h (Fig. 2F). This real

time measurement confirmed that cell cycle time extends as human ESC derived spinal cord

progenitors age (Rayon, et al., 2020). These findings further correlate with studies in mouse

(Calegari, et al., 2005, Calegari and Huttner, 2003, Takahashi, et al., 1995), and chick (Olivera-

Martinez, et al., 2014) embryos showing that neuron-generating divisions, which appear later

in development, are characterized by longer IKNM and cell cycle duration.

Labelling and live-imaging individual human spinal cord rosette cells using temporally defined

lipofection of piggyBac constructs

While it is possible to follow individual cell profiles as pm-FP expressing cells approach the

apical surface, divide and return to a basal position we also wished to monitor sub-cellular

behaviour within individual cells, including mitotic spindle orientation and final cleavage plane

as well as cell delamination events within this human neuroepithelium. To achieve this, we

designed a new strategy taking advantage of the precise timing of neurogenesis (Fig. 2A) (Dady,

et al., 2022) and a less efficient mode of cell transfection (Fig.3A): a mixture of piggyBac

constructs and the HyPBase transposase was transfected into D8/D9 human spinal cord neural

progenitors using lipofection (Felgner, et al., 1987)(Figure 3A, see Methods). Lipofected

neurogenic neural rosettes were then allowed to differentiate to later stages (Fig.3A). This

approach led to labelling of single or small groups of human spinal cord progenitors within each

rosette (Figs.3B,C).

Lipofection of cells with the pm-mKate and H2B-mEos3.2 plasmids allowed us to

monitor cell shape and chromatin configuration and so facilitated analysis of the orientation of

the mitotic spindle and final cleavage plane. Rosettes containing such cells were cultured until

D13 or D18 and imaged at 10 min intervals for 24-30h. Most divisions within each rosette

between D13 and D14 were symmetrical, with vertical cleavage planes and resulting horizontal

divisions, while between D18 and D19 most divisions were asymmetric, with horizontal

cleavage planes and resulting vertically orientated divisions (Figs 3D-F, movie 3). These findings

demonstrate that division orientation alters as development proceeds in human spinal cord neuroepithelium. This change correlates with the longer apical visiting times and the accumulation of neurons in later stage rosettes (Dady, et al., 2022).

To monitor delamination of new-born neurons within rosettes we followed individual cells with a basally located nucleus and a long cellular process inserted into the apical surface of the rosette. In a subset of these cells, we were able to observe delamination from the apical surface, a step characteristic of new-born neurons (Das and Storey, 2014). Moreover, this involved the abscission of an apical membrane particle in a manner characteristic of the apical abscission mechanism first found in the chicken spinal cord (Das and Storey, 2014, Kasioulis, et al., 2017)(Fig. 3G, 3H and movie 5). Overall, these analyses of human neuroepithelial cell behaviour during differentiation revealed the expansion of the early human neural progenitor cell pool in developing spinal cord, followed by a later phase characterized by slower cell cycle progression, increased asymmetric cell division and the generation of neurons.

Live imaging F-tractin in human cerebral cortex progenitors uncovers known and novel actin dynamics

Filamentous actin is a fundamental cytoskeletal component involved in the mechanics of cell division, cell motility and force transduction within tissues, and recent work has further uncovered roles around (Davidson and Cadot, 2021) and within the nucleus (Gunasekaran, et al., 2022, Hyrskyluoto and Vartiainen, 2020) where actin regulates gene expression as part of the nucleoskeleton in cell type and cell state specific ways, e.g. (Okuno, et al., 2020). Within the developing vertebrate neuroepithelium dynamic actin mediates aspects of IKNM (Spear and Erickson, 2012, Yanakieva, et al., 2019), apical cell adhesion and constriction leading to neuronal delamination, as well as intercellular filopodial contacts between non-neighbouring cells, e.g. (Das and Storey, 2014, Kasioulis, et al., 2022, Kasioulis, et al., 2017). In humans, actin cytoskeleton dysfunction in the cerebral cortex leads to periventricular heterotopia characterised by failure of neuronal delamination, altered cell adhesion and reduced brain size/cell proliferation, and is associated with seizures, dyslexia and psychiatric disorders (Lian and Sheen, 2015). Periventricular heterotopia is linked to mutation of actin binding proteins such as Filamin A (Sheen, et al., 2005). Moreover, the uniquely human fusion gene *CHRFAM7A* acts at least in part by activating actin in neuronal progenitors, impacting cell shape, growth cone extension and dendritic spine formation and so synaptic clustering (Szigeti, et al., 2023). The mechanisms by which actin mediates such critical processes in the developing human brain have yet to be elucidated.

The F-tractin-mKate2 hiPSC line was used to capture detailed actin dynamics in

cerebral cortex (cortical) neural rosettes. A dual-SMAD forebrain differentiation protocol was

followed, after (Chambers, et al., 2009) and neuroepithelial rosette structures were generated

by day 28. The appearance of Tbr2/EOMES positive cells (Fig. S2A) indicated that some basal

progenitors were present and confirmed forebrain tissue identity, while Beta-Tubulin labelling

(Fig. S2B) established neuron generation. Live imaging analyses focussed on actin dynamics in

cell behaviours characteristic of early cortical development. F-Tractin localised as expected to

the neuroepithelial cell cortex and was enriched at the apical cell junctions around apical end

feet and in rounding mitotic cells (Fig. 4A and movie 4): a similar distribution was revealed by

Phalloidin staining of fixed rosettes (Fig S2C). Apical enrichment of F-Tractin was also observed

during cell delamination from the apical surface (Fig 4B, movie 5). F-Tractin was detected at

low levels in the cytoplasm, distinguishing the nucleus (where F-Tractin was not detected) and

nuclear envelope breakdown (NEB) (Figs 4C-E). Most divisions (97%) took place at the apical

surface; 92% were symmetrical with a vertical cleavage plane (Fig. 4C, movie 6), while the

remaining divisions were asymmetric with a horizontal cleavage (Fig. 4D, movie 7). A few basal

divisions within the plane of rosettes (Fig. S2D, movie 8) correlated with low numbers of Tbr2

expressing cells at this time (Fig. S2A).

Focussing on entry into prophase, a transient F-Tractin enrichment localised to the

basal nuclear membrane prior to NEB: this was never observed on the apical side of the nuclear

membrane and disappeared as soon as NEB occurred (7.8±2.8' later, Fig 4C, and Fig 4E, 5'). This

pattern of actin localisation is a novel observation, which to the best of our knowledge has not

been described before. Actin polymerises around the entire nuclear envelope of mitotic U2OS

human cells just prior to NEB and associates with nuclear envelope remnants to facilitate

chromosome congression and correct segregation (Booth, et al., 2019). We speculate that, in

cortical progenitor cells, polarised actin polymerisation at the basal nuclear surface may

similarly prevent basal chromosome displacement and enable apically localised mitosis.

Following NEB, a strong enrichment of F-Tractin was further observed in the bulk of the basal

cortex as it moved toward the apical surface (Figs 4C-E). This movement ended 15.6±5.2' after

NEB (Fig 4C) with the cell rounding at the apical surface. At this stage, a thin basal process

persisted (Fig 4C, 26') and cytokinesis occurred 12.4±6.1' later. A role for actin in apical nuclear

movement during IKNM has been reported in the vertebrate neuroepithelium (Norden, et al.,

2009, Spear and Erickson, 2012, Yanakieva, et al., 2019) and in *Drosophila* larval wing disk cells,

where perturbation of actin polymerisation in the basal process disrupted apical movement

during IKNM (Meyer, et al., 2011). Overall, this analysis of F-tractin localisation in human

cortical rosettes revealed novel (accumulation at the basal nuclear membrane) as well as
expected actin localisations and dynamics. Next steps might include identification of actin
binding proteins that regulate specific actin functions and development of this tool to include
a nuclear localisation signal to investigate the roles of nuclear actin.

The human pluripotent fluorescent protein reporter cell lines generated and
characterised here and the piggyBac strategy used to generate such reporters, will be valuable
for the investigation of the cell biology of human neurogenesis. This could be extended from
neural rosettes to brain organoids to elucidate mechanisms underlying human cortical
expansion and indeed to any cell or tissue type that can be differentiated from these
pluripotent cell lines. Such tissue assays could be used to screen for the impact of
environmental factors (e.g. extreme heat, toxins, and other stressors) on human cell behaviour
and differentiation. The ability to visualise the actin cytoskeleton and other key cell biological
elements in pluripotent cells and their derivatives may also pave the way for biomechanical
studies in human tissues. These reporter cell lines could also be used in combination with
pharmacological studies, chemical genetics approaches, or in functional assays involving acute
and targeted degradation of proteins using proteolysis targeting chimeras (PROTACs) to
elucidate molecular mechanisms underlying cellular and developmental processes.

Materials and Methods
329
Human pluripotent cell authentication and quality control

Human ESC line H9 was obtained from WiCell and human iPSC line ChiPS4 from Cellartis AB,

now Takara Bio Europe. These suppliers provided documentation confirming cell identity. STR

DNA profiling for ChiPS4 and ChiPS4-pmGFP (polyclonal), H9 (WA09) confirmed identity of

these lines in our laboratory. All cell lines were expanded and banked on arrival. Prior to

experiments representative lots of each cell bank were thawed and tested for post-thaw

viability and to ensure sterility and absence of mycoplasma contamination. Sterility testing was

performed by direct inoculation of conditioned medium into tryptic soya broth and soya bean

casein broth and no contamination was observed. Mycoplasma testing was carried out by DAPI

staining of fixed cultures and using the mycoalert mycoplasma detection kit (Lonza). We

confirm that the cell lines used in this study were free of mycoplasma and aerobic bacteria and

fungi.

342   H9 (WA09) hES cells were supplied at passage 24, the cells were thawed, and cell banks

prepared at passage 29: for experiments the cells were used between passage 31 and 41.

ChiPS4 hiPS cells were supplied at passage 9 and cell banks prepared at passage 13: for

experiments the cells were used between passage 15 and 25. Two passages after thawing cells

were checked for uniform expression of pluripotency markers (OCT4, SOX2, NANOG, SSEA-3,

SSEA-4 TRA-1-60 and TRA-1-81), absence of differentiation markers (SSEA-1, HNF3-beta, beta-

III-tubulin and smooth muscle alpha actinin) by immunofluorescence.

Human Pluripotent cell culture

Human pluripotent stem cells were maintained as feeder-free cultures in TESR medium

(Ludwig, et al., 2006) supplemented with bFGF (30ng/mL, Peprotech) and Noggin (10ng/ml,

Peprotech) on Geltrex matrix coated plates (10 μg/cm^2, Life Technologies) at 37^{o}C in a

humidified atmosphere of 5% CO_2 in air. Cells were enzymatically dispersed twice a week using

TrypLE select (Life Technologies), counted using a hemacytometer and seeded on Geltrex

coated plates at a density of 3e4-5e4 cells/cm^2. For single-cell passaging, the medium was

supplemented by addition of the Rho kinase inhibitor Y-27632 (10 μM, Tocris). All work with

hESCs was undertaken in accordance with the UK Stem Cell Bank steering committee (license

number SCSC14-29).

Plasmid expression constructs

The APX1 (**A**dvanced **P**iggyBac E**x**pression) plasmid was synthetically engineered to harbor improved PiggyBac inverted terminal repeats (ITRs), the cytomegalovirus (CMV) enhancer/chicken Beta-actin (CAG) promoter, woodchuck post regulatory element (WPRE), and the Life Technologies Gateway recombination cassette allowing for efficient phiC31-mediated recombination (Sanders, et al., 2013)(Figure 1A). Subcellularly localized fluorescent protein gene cassettes were synthesized by overlap extension PCR and cloned into the Gateway Entry Vector, pENTRDTOPO, sequence verified. High efficiency transposase-mediated insertion of the PiggyBac transposon cassette was accomplished through transient co-expression of the mammalian optimized PiggyBac transposase (CMV HyPBase, (Yusa, et al., 2011)).

For multicolour labelling of subcellular structures, we utilized monomeric enhanced green fluorescent protein (mEGFP) (gift from Karel Svoboda, Addgene plasmid #18696) (Harvey, et al., 2008) and far-red mKate2 (Evrogen). These proteins were selected for their photostability, low toxicity, high brightness and distinct emission spectra for dual-colour assays making them suitable for live-cell imaging. To label the outer leaflet of the cell membrane, membrane-associated palmitoylated proteins were generated by appending 20-amino acid sequence of ratGAP-43 (MLCCMRRTKQVEKNDEDQKI) to the N-terminus (pm-XFP). To label the cell nucleus, mEos3.2 and green-to-red photoconvertible protein (a gift from Michael Davidson & Tao Xu (Addgene plasmid 54525) was coupled to Histone 2B (H2B) (Zhang, et al., 2012). For labeling of filamentous actin, F-Tractin which contains the inositol triphosphate 3-kinase actin binding domain was coupled to mKate2. Cell lines expressing both H2B-mEos3.2 and pm-mKate2 were then created by co-transfection of each construct and selection for dual expressing cells using FACS.

hPSCs transfection

H9 hESCs or hIPSCs were transfected using a Neon electroporator using 10 µl tips (Life Technologies). Briefly, cells were dispersed, resuspended in TESR medium and counted as described above. For each transfection 1e6 cells were collected by centrifugation at 300 xg for 2 minutes, washed with 1ml of PBS, then resuspended with 10µl of Neon electroporation buffer R. 1 µg of piggyBac vector and 0.2 µg of CMV HyPBase vector was then added to the cells in a volume of 2 µl. The cells were then electroporated (1150v, 30 mSec, 1 pulse), resuspended in TESR medium supplemented with bFGF, Noggin and Y27632 and plated on Geltrex coated dishes. Cells were maintained as described above then FAC-sorted 10 days after transfection.

Flow cytometry and cell sorting

Transfected hESCs or hiPSCs were dispersed and re-suspended as described above, then sorted

by FACS using a SH800 (Sony) cell sorter as a bulk pool of cells with a narrow gate selected for

robustly expressing cells. Quadrant gates used to estimate the percentage of positive cells and

level of fluorescence needed were designed based on fluorescence levels detected in the non-

transfected control samples. Sorted cells expressing medium levels of fluorescence were

expanded and frozen then representative lots of each cell line were thawed and tested for

mycoplasma contamination using the MycoAlert assay (Lonza), and fungi and aerobic bacteria

sterility by inoculating conditioned medium into trypticase soya broth (Sigma Aldrich). No

contamination was observed. In some instances (pm-eGFP in hIPSC4) monoclonal cell lines

were made from the polyclonal FACS sorted pools. To make monoclonal cell lines, 400 cells

were seeded into 60 mm dishes and cultured until the colonies were 2 – 3 mm in diameter.

They were then picked using 3.2 mm cloning discs (Sigma Aldrich) soaked in TrypLE select and

seeded into 96 well plates. Individual clones were then expanded in TESR medium. Controls

for pluripotency were undertaken as described above.

Neural differentiation assays

Spinal cord rosette differentiation: This was carried out following (Dady, et al., 2022, Verrier,

et al., 2018). Briefly, hESC were plated on Geltrex matrix (20 μg/cm2, Life Technologies) at a

density of 4e4 cells/cm^2 in TSER medium as described and allowed to attach for 24h. Cells were

differentiated for 48 hours in Neurobasal medium plus 1X B27, 1xN2, 1x non-essential amino

acids and 1x glutamax (all Thermo Fisher) (N2B27 medium) supplemented with 3 μM

Chiron99021 (Tocris) and 20 ng/ml bFGF (PeproTech) followed by a further 24 hours in N2B27

medium supplemented with 3 μM Chiron99021, 20 ng/ml bFGF, 10 μM SB431542 (Tocris) and

50 μg/ml Noggin (Peprotech) to obtain NeuroMesodermal progenitors (NMPs). For neural

differentiation of NMPs, these cells were then passaged using PBS-EDTA 0.5mM and seeded

back at controlled density (1e5 cells/cm2) on Geltrex (20μg/cm2, Life Technologies) in N2B27

medium supplemented with 10 μM SB431542, 50 μg/ml Noggin, 100 nM all-trans retinoic acid

(Sigma Aldrich) and 10 μM Y27632 (Tocris). After 24 hours the medium was removed and

replaced with N2B27 medium containing 100 nM all-trans retinoic acid for the remainder of

the differentiation. As described previously, dorsal spinal cord neural rosettes formed from day

6 (Dady, et al., 2022) and medium was refreshed every 3 days.

Forebrain/Cortical rosette differentiation: hPSCs were dispersed as described above, counted,

and re-suspended in TESR medium supplemented with bFGF (30 ng/ml), Noggin (10 ng/ml) and

Y27632 (10 μM) at a concentration of 1e5 cells/ml. 100 μl of cell suspension (1e4 cells) was

then added to the required number of wells of 96 well v-bottom plates and the cells pelleted

by centrifugation at 300 xg for 5 minutes to form embryoid bodies (EBs). EBs were cultured for

48 hours then collected from the 96 well plates using an 8-channel pipette, pooled together,

and washed twice with DMEM/F12 medium (Gibco). EBs were then resuspended in DMEM/F12

containing 20% knockout serum, 1x glutamax, 1x NEAA, 0.1 mM beta-mercaptoethanol (all

Gibco), SB431542 (10 μM, Tocris) and LDN193189 (0.5 μM Sigma Aldrich) and 50 EBs

transferred to each well of a low attachment 6-well plate in 2ml of medium to begin

differentiation. Every 48 hours 1ml of medium was removed from each well and replaced with

Neurobasal (Gibco) containing 1x glutamax, 1x NEAA, 1x N2 (Gibco), 1x B27 (Gibco), SB431542

(10 μM) and LDN193189 (0.5 μM). On day 9 of differentiation the EBs were recovered from the

6 well plates and replated on Geltrex coated (10 μg/cm2) 8 well chamber slides (Ibidi) using 10

– 15 EBs per well in Neurobasal containing 1x glutamax, 1x NEAA, 1x N2, 1x B27, SB431542 (10

μM) and LDN193189 (0.5 μM). The medium was refreshed every 48 hours until day 15 of

differentiation when it was changed to Neurobasal containing 1x glutamax, 1x NEAA, 1x N2, 1x

B27, BDNF (20 ng/ml, Peprotech) and GDNF (20 ng/ml, Peprotech). The medium was then

refreshed every 3 days, neural rosettes were evident from day 18 of differentiation.

Labeling individual cells in human spinal cord rosettes

For single cell or small cell group transfection, NMP-L cells were replated at end of D3 and

cultured for 8 to 9 days and transfected using Lipofectamine 3000 kit (Thermo Fisher) following

manufacturer's instructions with a combination of piggyBac plasmids (pm-eGFP + HypBase or

pm-mKate +H2B-mEos3.2 + HypBase) for a final concentration of 2 μg/μl of DNA mix. These

lipofected human neural progenitors were then cultured and imaged for the desired period.

Immunofluorescence

Commercially available primary antibodies were used to characterize cell types:

OCT4 Cell Signaling Technology (C30A3) rabbit mAb (1:400); SOX2 Cell Signaling Technology

(D6D9) rabbit mAb (1:400); NANOG Cell Signaling Technology (D73G4) rabbit mAb (1:400);

SSEA-4 Developmental Studies Hybridoma Bank (MC-813-70) mouse mAb (2 micro g per ml)

TRA-1-60 Cell Signaling Technology mouse mAb (1:200);TRA-1-81 Cell Signaling Technology

mouse mAb (1:200);SSEA-1 Developmental Studies Hybridoma Bank (MC-480) mouse mAb (2

micro g per ml);HNF-3 beta Cell Signaling Technology (D56D6) rabbit mAb (1:400) beta-III-

tubulin Sigma Aldrich (SDL.3D10) mouse mAb (1:800); Smooth muscle alpha actinin

Developmental Studies Hybridoma Bank (1E12) mouse mAb (2 micro g per ml); NCad, SIGMA,

1:500, PKC-ζ, Santa Cruz, SC-216; TBR2/EOMES 1:200 Abcam 23345, TUJ1, Covance Cat# MMS-

435P, mouse monoclonal 1:1000. hPSCs and neural rosettes were fixed in 1%

paraformaldehyde solution overnight at 4°C. Samples were then permeabilized with 0.1%

Triton X-100 in PBS and subjected to immunofluorescence labeling using appropriate

secondary antibodies conjugated to Alexa-fluor 488, 568 or 594 (Life Technologies), and

processed for DAPI staining to visualize cell nuclei. F-Actin cytoskeleton was labeled using

Phalloidin coupled with FITC fluorophore (1:1000; ThermoFisher). Fluorescence images were

taken on an EVOS Cell Imaging System (Thermo Fisher Scientific) for hPSCs, on a Leica SP8

confocal (using a 20X air or 63X NA1.2 APO water immersion objective) for spinal cord rosettes,

while cortical rosettes were imaged using LEICA SP8 Stellaris confocal microscope or a Zeiss

spinning disc confocal microscope equipped with a ×63 NA 1.4 oil immersion objective.

Generating satellite human spinal cord rosettes

From NMP cell passaging, transfected cells were cultured in N2B27 containing 100nM RA

(SIGMA) until day 6 when neural rosettes were easily observable. Using a very sharp glass

needle (0.29-0.31 mm) (Vitrolife), neural rosettes were individually micro-dissected and gently

deposited on a 35 mm glass bottom petri dish (WPI) previously coated with Geltrex ($20\mu g/cm2$,

Life Technologies). Micro-dissected rosettes were then cultured in N2B27 containing 100nM

RA (SIGMA) until the desired stage of observation.

Live imaging of human neural rosettes

Cells in satellite rosettes formed from primary micro-dissected spinal cord rosettes or individual

lipofected neural progenitors were imaged using a DeltaVision Core microscope system,

allowing detection of fine cellular detail at low fluorescence levels with minimal exposure time,

in a WeatherStation environmental chamber maintained at 37°C. The spinal cord rosette

culture medium was buffered with a 5% CO_2/95% air mix and maintained in a humid chamber.

Images were acquired using an Olympus 40x 1.30 NA objective using a Xenon light source and

a CoolSnap HQ2 cooled CCD camera (Photometrics). 30-40 optical sections (exposure time, 10-

50 milliseconds for each channel, 1024x1024 pixels, 2x2 binning) spaced 1.5µm apart were

imaged for each micro-satellite neural rosette at 10-minute intervals for periods up to 36h.

Images were deconvolved and maximum intensity projections of Z-stacks were made using

SoftWorx imaging software (Applied Precision) (Das, et al., 2012).

Confocal live imaging of cortical rosettes expressing F-Tractin::mKate2 was undertaken at 37°C

with a 5% CO_2/95% air mix using a Leica SP8 Stellaris confocal microscope with a x63 NA1.4 oil

objective. Stacks of 18 slices optical z-section separated by 0.8 μm, covering a 154×154×14.4

μm region were acquired over multiple positions every 310 seconds for up to 24 hours. We

applied a 3D gaussian blur with a sigma of 0.8x 0.8x 0.8 μm.

Quantification of cell behaviour

In spinal cord rosettes, **apical visiting time** was assessed by measuring time taken for a single

labelled cell located in the basal region of one spinal cord rosette (rosette periphery) to move

to the apical region and divide (rosette centre). To be consistent, observed rosettes were

selected by their similar diameter ~100-150 μm. At least 5 different cells per spinal cord rosette

and at least 5 different rosettes were analyzed for each time point (D13, D14, D18, 1D9).

Cell division symmetry was assessed by measuring the angle of the metaphase plate relative

to the apical surface at the lumen wall of the spinal rosette during cell division using Fiji. Cell

divisions were considered symmetric when the metaphase plate was perpendicular to the axis

of cell division. On the contrary, cell divisions were considered asymmetric when the

metaphase plate formed an angle of 40-45° to the axis of cell division (with one daughter cell

remaining longer at the apical surface. All cells were analysed in 3D. At least 5 different cells

per spinal cord rosette and at least 5 different rosettes were analyzed for each time point.

In cortical rosettes, we measured the time between cell cycle stages in rosettes expressing F-

Tractin-mKate2. Nuclear Envelope Breakdown was identified as the moment when cytoplasmic

F-Tractin entered the position held by the nucleus during the previous timepoint. Completion

of apical-to-basal migration as the moment when the cell appeared rounded at the apical

surface. Cytokinesis onset was defined as the timepoint when the first membrane cortical

contraction was detected. Cells in 12 different rosettes imaged for 22 hours from D29-D30

were analysed.

Statistical analysis

Statistical analysis for spinal cord rosettes was performed using non-parametric Mann-

Whitney U test for non-normally distributed data using Prism V8. Results are represented as

means ± SEM or SD as indicated in figures and p-values are indicated with *$p<0.05$, **$p<0.01$,

*** $p<0.001$, and ****$p<0.0001$.

Acknowledgements
532
The authors declare that they have no competing interests. We are grateful to Professor
Timothy Sanders for gifts of APX-1 plasmid and FP-constructs. AD conducted also cloning work
for some expression constructs in the Sanders Laboratory. This work was supported by an
enhancement to a Wellcome Trust Investigator award to KGS (WT102817AIA). Research in the
J.J. lab is supported by a BBSRC grant (BB/V001353/1). The Leica Stellaris microscope used for
confocal live cell imaging was supported by BBSRC grant BB/T017546/1. The Leica SP8 was
funded by a Wellcome Trust Multi-User Equipment grant (WT101468).
540
541
542
543

Figure Legends

Figure 1 Creation of fluorescent reporters in human pluripotent cells

A) Schematic of the APX1 plasmid used for high level expression of fluorescent protein fusion constructs highlighting the relevant elements of the system; **B)** Schematic of fluorescent-protein reporter constructs for introduction into human pluripotent cells using piggyBac transposase; **C)** transfection regime and FACS for selection of cells expressing constructs; **D)** Expression of reporters in human pluripotent cell lines, pm-eGFP, pm-mKate2, H2B-mEos3.2, pm-mKate2::H2B-mEos or F-Tractin-pm-mKate2. Scale bars 50μm.

Figure 2 Spinal cord differentiation and rosette generation for live imaging

A) Neural differentiation regime, after (Verrier, et al., 2018) and primary neural rosette formation and expansion from NMPs at (Day 4 to D14) apical/ luminal surface identified by N-Cadherin (NCAD) expression, and nuclei by DAPI staining; **B)** manual microdissection of day 6 (D6) primary neural rosettes (**PrR**), replating and culture to 14/15 and beyond generates; **C)** small satellite micro-rosettes (**SmR**) around the PrR, suitable for live imaging; **D)** Satellite micro-rosette made using pm-eGFP hESC line (movie 1); **E)** selected frames from live imaging of this SmR following IKNM including mitosis; **F)** Apical visiting time quantifications, revealing increase as development progresses (5 independent experiments for each day, measurement of 25 cells for each of D13 and D14, 23 cells for each of D18 and 19), means ± SD, p-values *$p<0.05$, **$p<0.01$, *** $p<0.001$, and ****$p<0.0001$. Scale bars 50 μm

Figure 3 Mosaic transfection of spinal cord rosettes using lipofection

A) Timing of lipofection-mediated introduction of piggyBac constructs into spinal cord rosette cells and live-cell imaging; **B,C)** pm-mKate2 construct expression revealing individual neuroepithelial cell morphology across apico-basal axis; **D,E)** Mosaic pm-mKate2:H2B-mEOS3.2 expression allowing live-cell imaging of individual cells and facilitating measurement of division symmetry (movie 2); and **F)** Percentage of asymmetric and symmetric divisions at apical surface in individual rosettes during early and late D18-19 neurogenesis (5 independent experiments for each stage: D13-14 (181 divisions in 26 rosettes), D18-19 (163 divisions in 36 rosettes) means ± S.E.M; **G,H)** Live cell imaging of apical abscission in individual cell expressing the pm-mKate:H2B-mEOS construct (movie 3). Scale bars 50 μm.

Figure 4 Live-cell imaging of F-Tractin in Cortical rosettes

Live imaging of cortical rosettes expressing F-Tractin showing, **A)** whole rosette monitored over 24h, note F-Tractin enrichment in apical end feet in rosette centre and prominent apical mitoses (movie 4); **B)** Enrichment of F-Tractin::mKate2 at the apical tip of a delaminating cell (9 cases across 30 rosettes) (movie 5); **C)** Vertical cleavage plane (symmetric; n=101/110 divisions occurring at the apical surface) division of a neuroepithelial cell in a cortical rosette. Values above black arrows: average time between specific stages: respectively, n=83, n=98 and n=110 (movie 6); **D)** Horizontal cleavage plane (asymmetric; n=101/110 divisions occurring at the apical surface) division of a neuroepithelial cell in a cortical rosette (movie 7); **E)** Neuroepithelial cell in a cortical rosette before and during Nuclear Envelope Breakdown (NEB). Arrowheads: transient F-Tractin enrichment at basal nuclear surface; arrows: F-Tractin-rich basal process (except in B, where indicates apical tip); asterisks: daughter cells following cytokinesis. Scale bars: A), B), 20 µm, C), E), 10 µm, D) 5 µm, t0: NEB, all cortical rosette imaging from D28 to 29.

Supplementary Figures

Figure S1 - FASC data for human pluripotent cells expressing fluorescent reporters

(related to Figure 1)

FASC data for selection of human pluripotent cells expressing fluorescent reporters, showing expression of eGFP in 82.82% of cells, mKate2 in 92.74% of cells, and H2B-mEos3 only in 22.75% of cells, while 13.79% of cells co-expressed pm-Kate and H2B-mEos3.2.

Figure S2 - Characterisation of cortical rosettes

(related to Figure 4)

A) Fixed cortical rosette stained for DAPI (magenta) and TBR2 (green); **B)** Fixed cortical rosette stained for DAPI (blue), apically localised PKC-ζ (magenta) and neuronal Beta-Tubulin (green). Lower panel: orthogonal re-slice. Scale bar: 50 µm; **C)** Fixed cortical rosette stained for Phalloidin displaying similar localisation to F-tractin::mKate in Cortical rosette; **D)** Basal Cortical progenitor cell division within the plane of the rosette (movie 8). Scale bars: A), B), C), 50 µm; D) 20 µm (top) and 10 µm (bottom).

Movie legends

Live imaging of cell behaviour in human spinal cord rosettes

Movie 1 - related to Figs 2D ,E: example of nuclear apical-basal visiting in spinal cord rosette

expressing pm-eGFP

Movie 2 - related to Figs 3D ,E: examples of symmetric and asymmetric (final division) in spinal

cord rosette following lipofection-mediated mosaic expression pm-mKate2::H2B-mEos3.2

Movie 3 - related to Figs 3G, H: Apical delamination of individual neuroepithelial cell following

lipofection-mediated mosaic expression pm-mKate2::H2B-mEos3.2

Live imaging of cell behaviour in human Cortical rosettes

Movie 4 - related to Fig 4A: Live cortical rosette expressing F-Tractin-mKate2.

Scale bar: 20 µm.

Movie 5 - related to Fig 4B: Apical delamination of cortical progenitor cell. Scale bar: 10 µm.

Movie 6 - related to Fig 4C: Apical symmetrical division of cortical progenitor cell. Scale bar:

10 µm.

Movie 7 - related to Fig 4D: Apical asymmetric division of cortical progenitor cell.

Scale bar: 5 µm.

Movie 8 - related to Fig 4SD: Basal cortical progenitor cell division. Scale bars: 20 µm (top)

and 10 µm (bottom).

References

Booth, A. J. R., Yue, Z., Eykelenboom, J. K., Stiff, T., Luxton, G. W. G., Hochegger, H. and Tanaka, T. U.
(2019). Contractile acto-myosin network on nuclear envelope remnants positions human
chromosomes for mitosis. Elife 8.

Calegari, F., Haubensak, W., Haffner, C. and Huttner, W. B. (2005). Selective lengthening of the cell
cycle in the neurogenic subpopulation of neural progenitor cells during mouse brain development. J
Neurosci 25, 6533-6538.

Calegari, F. and Huttner, W. B. (2003). An inhibition of cyclin-dependent kinases that lengthens, but
does not arrest, neuroepithelial cell cycle induces premature neurogenesis. J Cell Sci 116, 4947-4955.

Cereseto, A. and Giacca, M. (2004). Integration site selection by retroviruses. AIDS Rev 6, 13-21.

Chambers, S. M., Fasano, C. A., Papapetrou, E. P., Tomishima, M., Sadelain, M. and Studer, L. (2009).
Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. Nat
Biotechnol 27, 275-280.

Charpentier, E. and Doudna, J. A. (2013). Biotechnology: Rewriting a genome. Nature 495, 50-51.

Dady, A., Davidson, L., Halley, P. A. and Storey, K. G. (2022). Human spinal cord in vitro differentiation
pace is initially maintained in heterologous embryonic environments. Elife 11.

Dambournet, D., Hong, S. H., Grassart, A. and Drubin, D. G. (2014). Tagging endogenous loci for live-cell
fluorescence imaging and molecule counting using ZFNs, TALENs, and Cas9. Methods Enzymol 546,
139-160.

Das, R. M. and Storey, K. G. (2014). Apical abscission alters cell polarity and dismantles the primary
cilium during neurogenesis. Science 343, 200-204.

Das, R. M., Wilcock, A. C., Swedlow, J. R. and Storey, K. G. (2012). High-resolution live imaging of cell
behavior in the developing neuroepithelium. Journal of visualized experiments : JoVE.

Davidson, P. M. and Cadot, B. (2021). Actin on and around the Nucleus. Trends Cell Biol 31, 211-223.

Ding, S., Wu, X., Li, G., Han, M., Zhuang, Y. and Xu, T. (2005). Efficient transposition of the piggyBac (PB)
transposon in mammalian cells and mice. Cell 122, 473-483.

Doudna, J. A. and Charpentier, E. (2014). Genome editing. The new frontier of genome engineering
with CRISPR-Cas9. Science 346, 1258096.

Elick, T. A., Bauser, C. A. and Fraser, M. J. (1996). Excision of the piggyBac transposable element in vitro
is a precise event that is enhanced by the expression of its encoded transposase. Genetica 98, 33-41.

Engelman, A. (2003). The roles of cellular factors in retroviral integration. Curr Top Microbiol Immunol
281, 209-238.

Felgner, P. L., Gadek, T. R., Holm, M., Roman, R., Chan, H. W., Wenz, M., Northrop, J. P., Ringold, G. M.
and Danielsen, M. (1987). Lipofection: a highly efficient, lipid-mediated DNA-transfection procedure.
Proc Natl Acad Sci U S A 84, 7413-7417.

Gunasekaran, S., Miyagawa, Y. and Miyamoto, K. (2022). Actin nucleoskeleton in embryonic
development and cellular differentiation. Curr Opin Cell Biol 76, 102100.

Hacein-Bey-Abina, S., Hauer, J., Lim, A., Picard, C., Wang, G. P., Berry, C. C., Martinache, C., Rieux-
Laucat, F., Latour, S., Belohradsky, B. H., et al. (2010). Efficacy of gene therapy for X-linked severe
combined immunodeficiency. N Engl J Med 363, 355-364.

Hacein-Bey-Abina, S., Von Kalle, C., Schmidt, M., McCormack, M. P., Wulffraat, N., Leboulch, P., Lim, A.,
Osborne, C. S., Pawliuk, R., Morillon, E., et al. (2003). LMO2-associated clonal T cell proliferation in two
patients after gene therapy for SCID-X1. Science 302, 415-419.

Harvey, C. D., Yasuda, R., Zhong, H. and Svoboda, K. (2008). The spread of Ras activity triggered by
activation of a single dendritic spine. Science 321, 136-140.

Haupt, A., Grancharova, T., Arakaki, J., Fuqua, M. A., Roberts, B. and Gunawardane, R. N. (2018).
Endogenous Protein Tagging in Human Induced Pluripotent Stem Cells Using CRISPR/Cas9. J Vis Exp.

Hockemeyer, D., Wang, H., Kiani, S., Lai, C. S., Gao, Q., Cassady, J. P., Cost, G. J., Zhang, L., Santiago, Y.,
Miller, J. C., et al. (2011). Genetic engineering of human pluripotent cells using TALE nucleases. Nat
Biotechnol 29, 731-734.

Hyrskyluoto, A. and Vartiainen, M. K. (2020). Regulation of nuclear actin dynamics in development and
disease. Curr Opin Cell Biol 64, 18-24.

Ivics, Z., Hackett, P. B., Plasterk, R. H. and Izsvak, Z. (1997). Molecular reconstruction of Sleeping
Beauty, a Tc1-like transposon from fish, and its transposition in human cells. Cell 91, 501-510.

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A. and Charpentier, E. (2012). A programmable
dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. Science 337, 816-821.

Joung, J. K. and Sander, J. D. (2013). TALENs: a widely applicable technology for targeted genome
editing. Nat Rev Mol Cell Biol 14, 49-55.
Kasioulis, I., Dady, A., James, J., Prescott, A., Halley, P. A. and Storey, K. G. (2022). A lateral protrusion
latticework connects neuroepithelial cells and is regulated during neurogenesis. J Cell Sci 135.
Kasioulis, I., Das, R. M. and Storey, K. G. (2017). Inter-dependent apical microtubule and actin dynamics
orchestrate centrosome retention and neuronal delamination. Elife 6.
Kawakami, K. (2007). Tol2: a versatile gene transfer vector in vertebrates. Genome Biol 8 Suppl 1, S7.
Kawakami, K. and Shima, A. (1999). Identification of the Tol2 transposase of the medaka fish Oryzias
latipes that catalyzes excision of a nonautonomous Tol2 element in zebrafish Danio rerio. Gene 240,
239-244.
Kim, A. and Pyykko, I. (2011). Size matters: versatile use of PiggyBac transposons as a genetic
manipulation tool. Mol Cell Biochem 354, 301-309.
Lee, C. S., Bishop, E. S., Zhang, R., Yu, X., Farina, E. M., Yan, S., Zhao, C., Zheng, Z., Shu, Y., Wu, X., et al.
(2017). Adenovirus-Mediated Gene Delivery: Potential Applications for Gene and Cell-Based Therapies
in the New Era of Personalized Medicine. Genes Dis 4, 43-63.
Lee, C. Y., Li, J. F., Liou, J. S., Charng, Y. C., Huang, Y. W. and Lee, H. J. (2011). A gene delivery system for
human cells mediated by both a cell-penetrating peptide and a piggyBac transposase. Biomaterials 32,
6264-6276.
Lian, G. and Sheen, V. L. (2015). Cytoskeletal proteins in cortical development and disease: actin
associated proteins in periventricular heterotopia. Front Cell Neurosci 9, 99.
Liu, X., Li, N., Hu, X., Zhang, R., Li, Q., Cao, D., Liu, T., Zhang, Y. and Liu, X. (2013). Efficient production of
transgenic chickens based on piggyBac. Transgenic Res 22, 417-423.
Ludwig, T. E., Levenstein, M. E., Jones, J. M., Berggren, W. T., Mitchen, E. R., Frane, J. L., Crandall, L. J.,
Daigh, C. A., Conard, K. R., Piekarczyk, M. S., et al. (2006). Derivation of human embryonic stem cells in
defined conditions. Nat Biotechnol 24, 185-187.
Meyer, E. J., Ikmi, A. and Gibson, M. C. (2011). Interkinetic nuclear migration is a broadly conserved
feature of cell division in pseudostratified epithelia. Curr Biol 21, 485-491.
Miller, J. C., Tan, S., Qiao, G., Barlow, K. A., Wang, J., Xia, D. F., Meng, X., Paschon, D. E., Leung, E.,
Hinkley, S. J., et al. (2011). A TALE nuclease architecture for efficient genome editing. Nat Biotechnol
29, 143-148.
Miskey, C., Izsvak, Z., Plasterk, R. H. and Ivics, Z. (2003). The Frog Prince: a reconstructed transposon
from Rana pipiens with high transpositional activity in vertebrate cells. Nucleic Acids Res 31, 6873-
6881.
Narayanavari, S. A., Chilkunda, S. S., Ivics, Z. and Izsvak, Z. (2017). Sleeping Beauty transposition: from
biology to applications. Crit Rev Biochem Mol Biol 52, 18-44.
Norden, C., Young, S., Link, B. A. and Harris, W. A. (2009). Actomyosin is the main driver of interkinetic
nuclear migration in the retina. Cell 138, 1195-1208.
Okuno, T., Li, W. Y., Hatano, Y., Takasu, A., Sakamoto, Y., Yamamoto, M., Ikeda, Z., Shindo, T., Plessner,
M., Morita, K., et al. (2020). Zygotic Nuclear F-Actin Safeguards Embryonic Development. Cell Rep 31,
107824.
Olivera-Martinez, I., Schurch, N., Li, R. A., Song, J., Halley, P. A., Das, R. M., Burt, D. W., Barton, G. J. and
Storey, K. G. (2014). Major transcriptome re-organisation and abrupt changes in signalling, cell cycle
and chromatin regulation at neural differentiation in vivo. Development 141, 3266-3276.
Pacesa, M., Pelea, O. and Jinek, M. (2024). Past, present, and future of CRISPR genome editing
technologies. Cell 187, 1076-1100.
Rayon, T., Stamataki, D., Perez-Carrasco, R., Garcia-Perez, L., Barrington, C., Melchionda, M., Exelby, K.,
Lazaro, J., Tybulewicz, V. L. J., Fisher, E. M. C., et al. (2020). Species-specific pace of development is
associated with differences in protein stability. Science 369.
Roberts, B., Haupt, A., Tucker, A., Grancharova, T., Arakaki, J., Fuqua, M. A., Nelson, A., Hookway, C.,
Ludmann, S. A., Mueller, I. A., et al. (2017). Systematic gene tagging using CRISPR/Cas9 in human stem
cells to illuminate cell organization. Mol Biol Cell 28, 2854-2874.
Sanders, T. A., Llagostera, E. and Barna, M. (2013). Specialized filopodia direct long-range transport of
SHH during vertebrate tissue patterning. Nature 497, 628-632.
Sarkar, A., Sim, C., Hong, Y. S., Hogan, J. R., Fraser, M. J., Robertson, H. M. and Collins, F. H. (2003).
Molecular evolutionary analysis of the widespread piggyBac transposon family and related
"domesticated" sequences. Mol Genet Genomics 270, 173-180.
Sauer, F. C. (1935). Mitosis in the neural tube. J. Comp. Neurol. 62, 377-405.

Sheen, V. L., Jansen, A., Chen, M. H., Parrini, E., Morgan, T., Ravenscroft, R., Ganesh, V., Underwood, T.,
Wiley, J., Leventer, R., et al. (2005). Filamin A mutations cause periventricular heterotopia with Ehlers-
Danlos syndrome. Neurology 64, 254-262.
Spear, P. C. and Erickson, C. A. (2012). Interkinetic nuclear migration: a mysterious process in search of
a function. Dev Growth Differ 54, 306-316.
Sultana, T., Zamborlini, A., Cristofari, G. and Lesage, P. (2017). Integration site selection by retroviruses
and transposable elements in eukaryotes. Nat Rev Genet 18, 292-308.
Szigeti, K., Ihnatovych, I., Rosas, N., Dorn, R. P., Notari, E., Cortes Gomez, E., He, M., Maly, I., Prasad, S.,
Nimmer, E., et al. (2023). Neuronal actin cytoskeleton gain of function in the human brain.
EBioMedicine 95, 104725.
Takahashi, T., Nowakowski, R. S. and Caviness, V. S., Jr. (1995). The cell cycle of the pseudostratified
ventricular epithelium of the embryonic murine cerebral wall. J Neurosci 15, 6046-6057.
Verrier, L., Davidson, L., Gierlinski, M., Dady, A. and Storey, K. G. (2018). Neural differentiation,
selection and transcriptomic profiling of human neuromesodermal progenitor-like cells in vitro.
Development 145.
Wang, W., Lin, C., Lu, D., Ning, Z., Cox, T., Melvin, D., Wang, X., Bradley, A. and Liu, P. (2008).
Chromosomal transposition of PiggyBac in mouse embryonic stem cells. Proc Natl Acad Sci U S A 105,
9290-9295.
Woltjen, K., Michael, I. P., Mohseni, P., Desai, R., Mileikovsky, M., Hamalainen, R., Cowling, R., Wang,
W., Liu, P., Gertsenstein, M., et al. (2009). piggyBac transposition reprograms fibroblasts to induced
pluripotent stem cells. Nature 458, 766-770.
Wood, A. J., Lo, T. W., Zeitler, B., Pickle, C. S., Ralston, E. J., Lee, A. H., Amora, R., Miller, J. C., Leung, E.,
Meng, X., et al. (2011). Targeted genome editing across species using ZFNs and TALENs. Science 333,
307.
Woodard, L. E. and Wilson, M. H. (2015). piggyBac-ing models and new therapeutic strategies. Trends
Biotechnol 33, 525-533.
Wu, S. C., Meir, Y. J., Coates, C. J., Handler, A. M., Pelczar, P., Moisyadi, S. and Kaminski, J. M. (2006).
piggyBac is a flexible and highly active transposon as compared to sleeping beauty, Tol2, and Mos1 in
mammalian cells. Proc Natl Acad Sci U S A 103, 15008-15013.
Yanakieva, I., Erzberger, A., Matejcic, M., Modes, C. D. and Norden, C. (2019). Cell and tissue
morphology determine actin-dependent nuclear migration mechanisms in neuroepithelia. J Cell Biol
218, 3272-3289.
Yusa, K., Rad, R., Takeda, J. and Bradley, A. (2009). Generation of transgene-free induced pluripotent
mouse stem cells by the piggyBac transposon. Nat Methods 6, 363-369.
Yusa, K., Zhou, L., Li, M. A., Bradley, A. and Craig, N. L. (2011). A hyperactive piggyBac transposase for
mammalian applications. Proc Natl Acad Sci U S A 108, 1531-1536.
Zhang, M., Chang, H., Zhang, Y., Yu, J., Wu, L., Ji, W., Chen, J., Liu, B., Lu, J., Liu, Y., et al. (2012). Rational
design of true monomeric and bright photoactivatable fluorescent proteins. Nat Methods 9, 727-729.
784

A

APX1
7345

5' ITR(e)
CAG
3' ITR
bghpA
WPRE
Gateway Recombination Cassette
AmpR Cassette
E. Coli Backbone

B

ITR CAGp pm-eGFP ITR

or

pm-mKate2

H2B-mEos

pm-mKate2

F-Tractin-mKate2

C

hESCs
or hiPSCs

Electroporation
1x1150 V/30ms
10⁷ cells
2μg/μl DNA

cell
sorting
FACS

D

pm-eGFP
H9 hESC

pm-mKate2
H9 hESC

pm-eGFP
ChiPSC4 (clonal)

H2B-mEos3.2
H9 hESC

H2B-mEos3.2
+ pm-mKate2
H9 hESC

F-Tractin-mKate2
ChiPSC4

Figure 1 Dady et al

Figure 2 Dady et al

Figure 3 Dady et al

Figure 4 Dady et al

Figure S1 Dady et al