Long title

PAN deadenylase ensures proper mitosis under conditions of microtubule stress by regulating

spindle integrity and promoting cell survival

Short title

PAN deadenylase is required for mitosis in response to microtubule stress

Authors

Jigyasa Verma[1], Zhengcheng He[2], Joshua A.R. Brown[3], Pamela Dean[1], Barry P. Young[1],

Stephane Flibotte[4], LeAnn J. Howe[3], Christopher D. Maxwell[2], Calvin D. Roskelley[1], and

Christopher J.R. Loewen[1*]

Author Affiliations

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, University of

British Columbia, Vancouver, Canada, V6T1Z3

[2] Department of Pediatrics, British Columbia Children's Hospital, University of British

Columbia, Vancouver, Canada, V6H 0B3

[3] Department of Biochemistry and Molecular Biology, Life Sciences Institute, University of

British Columbia, Vancouver, Canada, V6T1Z3

[4] Bioinformatics Facility, Life Sciences Institute, University of British Columbia, Vancouver,

Canada, V6T1Z3

[*] Corresponding author: Department of Cellular and Physiological Sciences, Life Sciences

Institute, University of British Columbia, Vancouver, Canada, V6T1Z3 cloewen@mail.ubc.ca

Abstract

The Poly(A) Tail Length (PATL) of mRNAs of certain cell-cycle regulatory genes undergo significant trimming during M-phase, however the functional importance is unknown. The Ccr4-Not and PAN complexes account for the majority of cytoplasmic poly(A) deadenylation, however a role in M phase has not been described. We find that under conditions of microtubule stress in yeast, loss of PAN deadenylase activity leads to arrest in M phase, defective spindles, and increased cell death. PAN consists of the catalytic subunit Pan2 and the RNA binding subunit Pan3. Consistent with a role in mitosis, *PAN2* interacts genetically with tubulin genes, prefoldin complex genes and the mitotic cyclin *CLB1*. *PAN2* knockdown in human cultured cells disrupts mitosis and results in spindle fragmentation leading to abnormal cell division, while expression of human *PAN2* in yeast rescues *pan2Δ* cell-cycle phenotypes. Hence, we reveal an important highly conserved role for PAN in ensuring proper mitosis when cells are under microtubule stress. We propose PAN regulates PATLs of mRNAs of key cell-cycle/mitotic proteins in response to defective spindles.

Author Summary

Proper cell division is essential for the growth and survival of all living organisms. Our study investigates the role of PAN deadenylase complex in yeast and human cultured cells under microtubule stress, induced by microtubule inhibitors used in cancer treatment. The PAN complex, consisting of Pan2 and Pan3, trims the poly(A) tails of mRNAs. We found that loss of PAN activity leads to cell cycle arrest in M-phase, spindle defects and increased cell death in yeast. Similarly, *PAN2* knockdown in human cultured cells disrupts mitosis and causes abnormal cell division, indicating a conserved function across species. We propose that PAN regulates mRNA poly(A) tail lengths of key mitotic proteins to ensure proper mitosis under stress. This regulation likely prevents faulty spindle formation by repressing translation of these mRNAs. Interestingly, PAN's role is specific to stress conditions, as cells without PAN function normally otherwise. Our findings highlight PAN's critical role in maintaining genomic stability and proper cell division during microtubule stress, providing insights into the post-transcriptional regulation of cell cycle and potential targets for cancer therapy.

Introduction

In eukaryotes, poly(A) tails are present on the 3' end of almost every mRNA and regulate

mRNA stability and translation (1–6). Poly(A) tail length (PATL) varies greatly from as short as

~20 nucleotides to as long as ~250 nucleotides and is governed by poly(A) deadenylases, which

trim or remove the tails completely (7–10). Two protein complexes, PAN (PolyA Nuclease) and

Ccr4-Not, carry out the majority of deadenylation, where PAN initiates trimming of longer tails

while Ccr4-Not removes the remainder (11–14). Poly(A)-Binding Proteins (PABs) bind poly(A)

tails promoting deadenylation by recruiting PAN complex to poly(A) tails. Genes such as *CDK1*,

TOP2A and *FBXO5*, which orchestrate timing and coordination of mitotic events in somatic

cells, exhibit reduced PATLs specifically in M-phase (15), suggesting deadenylation of specific

mRNAs is important in cell-cycle regulation. However, roles for specific deadenylases in the

cell-cycle are poorly defined.

The structure of PAN complex is well understood. It comprises the catalytic subunit Pan2 and an

asymmetric homodimer of the RNA-binding regulatory subunit Pan3 (16–18). Pan2 contains a

C-terminal DEDD-family exoribonuclease domain that is responsible for its poly-A deadenylase

activity. Adjacent to the exoribonuclease domain is a ubiquitin-specific protease (USP) domain,

which in human cells is functional (19,20). *Saccharomyces cerevisiae* Pan2 has this domain also,

but it lacks critical residues in the catalytic triad indicating yeast Pan2 lacks deubiquitinase

activity (16,17). Pan3 contains a CCCH-type zinc finger domain and a Poly(A) interacting motif

2 (PAM2) motif that interacts with PABs facilitating RNA binding. Pan3 also contains a

pseudokinase domain (21). Pan2 is a substrate of Cdk1 phosphorylation (22). Pan3

phosphorylation regulates binding to PABs which regulates PAN activity (23), and Pan3 is a

substrate of Cdk5 phosphorylation (24,25).

The biological relevance of PANs deadenylase activity is poorly understood. In human cells,

knockdown of *PAN2* leads to an increase in mRNAs with long poly(A) tails (> ~150 nt), but

does not alter mRNA half lives (14). Deletion of either *PAN2* or *PAN3* in yeast leads to longer

mRNA poly(A) tails, but does not completely abolish deadenylation (11,12). This suggests that

PAN's role may be more specialized than regulating stability of bulk mRNAs, which has been

shown for *HIF1A* regulation (26). In yeast, a protein serine/threonine kinase involved in

replication fork protection and the DNA damage checkpoint, Dun1, interacts with Pan2 and Pan3

and regulates the composition and activity of post-replication repair complexes. The

dun1Δpan2Δ strain is hypersensitive to replication stress and has elevated levels of an important

regulator of error-free post-replication repair, Rad5 (27). This suggests PAN plays a role in post-

transcriptional regulation of DNA repair and replication stress although the mechanism and

mRNA targets are unknown.

We have identified a previously unknown role of PAN complex in cell-cycle regulation. High

throughput data indicate that both yeast *PAN2* and *PAN3* exhibit strong aggravating genetic

interactions with genes that encode tubulin and tubulin folding proteins (28), and Δ*pan2* and

Δ*pan3* yeast are among the most sensitive in the yeast deletion collection to various microtubule

biogenesis inhibitors, including nocodazole and benomyl (29). This suggested to us an important,

yet uncharacterized function for PAN in mitosis in yeast in response to microtubule stress. We

found that nocodazole caused PAN-deficient cells to arrest in M phase, likely due to defective

spindle assembly, resulting in increased cell death. Transcriptional profiling of cells expressing

deadenylase-dead Pan2 in the presence of nocodazole revealed changes in key cell-cycle

regulators, further supporting a function for PAN in mitosis. We extended our findings from

yeast to mammalian cells by knocking down *PAN2* in human cells, which caused spindle

fragmentation and defective mitosis upon treatment with nocodazole and colchicine, indicating

this function of PAN is conserved in metazoans.

Results

Yeast deletion mutants *pan2Δ* and *pan3Δ* are sensitive to nocodazole

On querying for information on the effects of small molecules on the growth of *pan2Δ* and

pan3Δ yeast deletion mutants, we noticed that homozygous deletion of *PAN2* or *PAN3* sensitizes

yeast to multiple drugs that disrupt tubulin biogenesis (29). These include nocodazole, benomyl,

and a number of other small molecules that belong to the response signature 'tubulin folding and

SWR complex'. Importantly, both *pan2Δ* and *pan3Δ* are ranked in the top 20 out of over ~4,800

yeast deletion mutants for nocodazole and benomyl sensitivity (29) (**SFig 1A**). Other

homozygous deletion strains which occupied top ranks are *pac2Δ, cin1Δ, cin2Δ,* and *tub3Δ*.

TUB3 encodes α-tubulin, which associates with β-tubulin (*TUB2*) to form tubulin dimers, which

polymerize to form microtubules. Pac2 is an alpha tubulin-binding protein, Cin1 is a beta-

tubulin-binding protein, and Cin2 is a GTPase activator involved in tubulin complex assembly

(30–33). Hence, Pac2, Cin1, and Cin2 are involved in protein folding and assembly of the

tubulin complex. This suggested a role for PAN complex in microtubule related functions.

 Figure S1

S1A

Fig S1A: Sensitivity of yeast homozygous deletion strains to nocodazole and benomyl. Ratio of

fitness defect scores for over ~4,800 mutants (x-axis: nocodazole; y-axis: benomyl). Blue circles:

microtubule complex mutants; green circles: PAN complex mutants. Data from A. Y. Lee et al.

2014. **Fig S1B**: Exponential growth constants for WT, *pan2Δ*, and *pan3Δ* with DMSO. Error

bars: +/- SD (n=6, three experiments). Paired Student's *t*-test for strain differences.

Fig S1C: GFP-Pan2 on plasmid assesses Pan2 expression in D1020A. Upper band: Pan-GFP;

lowermost band: free GFP. PGK1 - loading control. CEN, centromeric; GFP, green fluorescent

protein. **Fig S1D**: Growth curves of yeast strains with DMSO (blue) or 6 µM nocodazole

(yellow) at 30°C for 24 hours. Absorbance plotted with SD (n=6). min, minutes; SC, Synthetic

Complete media; SD, standard deviation; EV, empty vector; D1020A, deadenylase dead Pan2.

The nocodazole sensitivity screen described above was performed at 6 µM, while the typical

concentration used for inducing G2/M arrest is 40 µM (34). Since the screen produced a

phenotype at 6µM (29), we treated log phase cells of *pan2Δ*, *pan3Δ,* and their isogenic wild-type

strain with 6 µM nocodazole to mimic the screen and measured yeast growth. While no

difference in growth rate was observed for WT, *pan2Δ* and *pan3Δ* grew substantially slower in

the presence of nocodazole (**Fig 1A**), but not in the absence of nocodazole (**SFig 1B**). Since

growth dynamics were monitored immediately upon the addition of nocodazole, we observed a

lag period of approximately 100 minutes (~ 1 1/2 hours) before the effect of the drug was

noticeable. For this reason, exponential growth rates were not determined for these experiments.

Figure 1

Fig 1: Yeast deletion mutants *pan2Δ* and *pan3Δ* are sensitive to nocodazole. (A-D) Growth

curves of yeast strains in SC media with DMSO (blue) or 6 μM nocodazole (yellow) at 30°C for

24 hours. Absorbance plotted with SD (n=6). Normalized exponential growth constants have

been represented in **Fig 1C**. Error bars are +/- SD from three separate experiments (n=6).

Significant differences between different strains were tested using paired Student's *t*-test.WT,

wild type; min, minutes; SC, Synthetic Complete media; SD, standard deviation; EV, empty

vector; D1020A, deadenylase dead Pan2; Hs, *Homo sapiens.*

Expressing WT *PAN2* from a CEN plasmid eliminated the nocodazole sensitivity of *pan2Δ* cells

(**Fig 1B**). In contrast, expression of a catalytic-dead mutant of *PAN2* with a mutation within the

active site DEDD motif (D1020A) that is unable to perform in-vitro deadenylation on model-

polyA RNAs (35) was unable to eliminate the nocodazole sensitivity (**Fig 1B**). To verify that the

D1020A mutant was stably expressed, we tagged *PAN2* and the D1020A mutant with GFP and

examined protein expression and function. We found no difference in the level of GFP-Pan2 and

GFP-Pan2(D1020A) protein (**SFig 1C**). GFP-Pan2 largely eliminated nocodazole sensitivity of

pan2Δ cells whereas GFP-Pan2(D1020A) did not, confirming that the nocodazole sensitivity is

dependent upon Pan2 catalytic activity (**SFig 1D**).

To determine if the role of Pan2 in nocodazole sensitivity is conserved in mammalian cells, we

codon-optimized the human *PAN2* gene for expression in yeast (**SFile 1**) and expressed the gene

in the *pan2Δ* strain from a high copy 2-micron plasmid (2μ-HsPan2; **Fig 1C**). As was the case

with the WT yeast Pan2, expression of HsPAN2 eliminated sensitivity to nocodazole (there was

no difference in growth with or without nocodazole in cells expressing HsPan2). Interestingly,

overexpression of HsPAN2 in the *pan2Δ* strain caused a decrease in growth rate (**Fig 1C**). We

hypothesized that this reduction in growth would be Pan3 dependent if this phenotype was a

result of PAN activity. Indeed, expression of HsPan2 in the *pan3Δ* strain did not not slow growth

and did not prevent nocodazole sensitivity (**Fig 1D**). We found a similar slow-growth phenotype

with overexpression of yeast Pan2 (see **Fig 5D**). Thus, human Pan2 complemented the function

of yeast *pan2Δ* in response to nocodazole, indicating this function of PAN was highly conserved.

***PAN2* and *PAN3* genetically interact with tubulin and genes governing tubulin folding**

To better understand how PAN function was related to nocodazole sensitivity, we examined

known common genetic interactions of *PAN2* and *PAN3 (28)* (**SFig 2A**). Gene-ontology (GO)

analysis indicated roles for PAN in "chromosome segregation" (*APC1, BUB3, CBF1, MPS1,*

SLI15, SMC3, TUB2, and *TUB3*), "cytoskeletal protein binding" (*ARP2, CIN1, GIM3, GIM5,*

PAC10, RBL2, and *YKE2*), "secretion" (*ACT1, SEC1, SSO2,* and *VPS27*), and "RNA

polymerase-II-specific transcription factor binding" (*CBF1, GCR2,* and *SWI1*) (**SFig 2A**). Within

the "chromosome segregation" network, multiple functions relating to microtubules were

identified including "spindle", "microtubule cytoskeleton", and "mitotic spindle assembly

checkpoint signaling", consistent with a role for PAN in spindle function during mitosis.

S2A

S2B

S2C

Fig S2A: Heatmap and Cytoscape analysis of genetic interactions of PAN2 and PAN3 by

Costanzo et al., 2016. In the heatmap, darker shades of blue represent strong aggravating genetic

interactions and darker shades of yellow represent strong alleviating genetic interactions. In the

functional enrichment, node size indicates statistical significance of enriched GO terms and

edges indicate statistically significant associations between terms. **Fig S2B-C**: Growth curves of

yeast strains with DMSO or 1.5 μM nocodazole (B) and single or double knockout strains in

DMSO (C). Absorbance plotted with SD (n=6). **Fig S2D**: Gene expression changes in

pan2Δ+WT-PAN2 or *pan2Δ*+D1020A-PAN2 post 6 hour treatment with DMSO or 6 μM

nocodazole assessed by qRT-PCRs. The ΔCt values are normalized to actin. Data are mean +/-

S.D. n.s., not significant. Significant differences between all pairwise combinations of groups

was detected using two-way ANOVA comparison. Ct, cycle threshold; μM, micromolar; qRT,

quantitative reverse transcriptase; ANOVA, Analysis of Variance. min, minutes; SC, Synthetic

Complete media; SD, standard deviation; WT, wild type.

To validate the genetic interactions described above and expand the interaction landscape to

include microtubule stress from nocodazole treatment, we performed synthetic genetic array

(SGA) double mutant screens with a *pan2Δ* query and the non-essential deletion collection of

~4800 strains in the absence and presence of nocodazole (1.5µM nocodazole as this does not

lead to differential growth between WT and *pan2Δ* in liquid growth assays, **SFig 2B**). In the

absence of nocodazole using a cut-off ratio of 0.7 for aggravating genetic interactions (three

standard deviations, 99.7% confidence interval) and p-value of less than 0.05 in 3/3 replicates,

we found a total of 52 genetic interactions. For deletion mutants in the array whose growth was

rescued upon deleting *PAN2* in them (referred to as alleviating genetic interactions from here

onwards), we chose a cut-off ratio of 1.3 and found 27 genetic interactors (**SFile 2**). The 4 genes

in common with the previously identified genetic interactors (28) were *TUB3, BUB3, RPL12B,*

and *RPL14A*. Surprisingly, all the genes with functions related to chromosome segregation were

not identified in the absence of nocodazole, but instead required addition of low levels of

nocodazole (see below).

For SGA screens in the presence of nocodazole, we used a low nocodazole concentration (1.5

µM) anticipating that the genetic interactors themselves might be nocodazole sensitive, and that

this dose would not produce a growth phenotype in the *pan2Δ* query strain (**SFig 2B**). Using the

same cut-offs as the no-nocodazole screens, we identified a total of 30 aggravating and 21

alleviating genetic interactions (**SFile 2**). Of these, 7 genetic interactors were in common with

the PAN interactions from the CellMap (**Fig 2A**). These genes were *CIN1, GIM3, MDM32,*

PAC2, TUB3, VPS68 and *YKE2. CIN1, GIM3, PAC2,* and *YKE2* are involved in tubulin folding

and tubulin complex assembly. *CIN1* encodes for tubulin folding factor D involved in beta-

tubulin (Tub2) folding (36). *GIM3*, *PAC2,* and *YKE2* encode for subunits of the prefoldin co-

chaperone complex involved in tubulin folding (37). *TUB3* encodes alpha-tubulin which

associates with beta-tubulin (Tub2) to form the tubulin dimer. It is unclear why nocodazole was

required to detect genetic interactions with microtubule related components in our screens, but

not those performed by the Boone lab (28) however, screen conditions (query strain background,

media composition, temperature, array density) varied between our labs, suggesting perhaps that

our screens had reduced microtubule stress, which prevented identification of the interactions, or

that their screens had increased stress mimicking nocodazole addition.

    Figure 2

2A

2B

Fig 2: PAN2 and PAN3 genetically interact with tubulin and genes governing tubulin folding. (**A**) Venn diagram shows overlap between aggravating genetic interactions for PAN2 according to TheCellMap (yellow), PAN2 SGA screen in either absence (blue) or presence of nocodazole (light brown). (**B**) Growth curves of single knockout (blue) or double knockout (yellow) yeast strains in SC media with 1.5 µM nocodazole for 24 hours at 30°C. Absorbance plotted with SD (n=6). min, minutes; SC, Synthetic Complete media; SD, standard deviation.

To confirm genetic interactions of *PAN2* with genes involved in microtubule function, we assessed the sensitivities of single and double knock-out strains in liquid growth assays using the same low dose of nocodazole used in our screens (1.5 µM), which did not affect the growth of *pan2Δ* single mutant cells (**SFig 2B**). We confirmed slow growth phenotypes in double mutants with *CIN1, GIM3, PAC2,* and *TUB3* in the presence of 1.5 µM nocodazole (**Fig 2B**), indicating aggravating genetic interactions with these genes. There was no difference in growth of the single and double knockouts in the absence of drug (**SFig 2C**), consistent with our screen results. We also expanded our test set to include other genes involved in tubulin folding including *CIN2* (GTPase activating protein, tubulin folding cofactor C), *CIN4* (GTPase involved in beta-tubulin folding, regulated by *CIN2*), and *GIM4* (prefoldin co-chaperone complex). These all showed aggravating genetic interactions with *PAN2* in the presence of 1.5 µM nocodazole but not in the absence. Since PAN regulates mRNA stability, we were intrigued to test the mRNA levels of tubulin genes *TUB1, TUB2,* and *TUB3*, in the presence and absence of nocodazole, but found no significant changes in transcript abundance (**SFig 2D**). Thus, the genetic interactions between *PAN2* and *PAN3* and genes governing tubulin folding provided strong genetic evidence for a role for PAN in microtubule function.

PAN2 **deletion exacerbates nocodazole-induced G2/M arrest**

Upon establishing the nocodazole sensitivity of *pan2Δ* and *pan3Δ* and genetic interactions between

PAN2 and *PAN3* and genes governing tubulin folding in yeast, we were interested in understanding

this phenotype further. We performed flow cytometry to determine the distribution of cells in G1/S

and G2/M phases. Log phase cells of *pan2Δ* and WT were treated with either DMSO or 6 μM

nocodazole and collected at different time points after treatment (3, 6 and 10 hours). They were

fixed in ethanol and stained to analyze DNA content by flow cytometer. In the absence of drug

treatment, we saw that both *pan2Δ* and WT cells have typical distributions for asynchronous log

phase yeast cultures of 1N and 2N cells at all the time points, suggesting normal cell-cycle

progression (**Fig 3A**). However, upon treatment with nocodazole, we found that there was a shift

from 1N to 2N and a broadening of the 2N peak in *pan2Δ* as compared to WT cells at 3 hours,

indicating a G2/M delay (**Fig 3A**). We did not observe a discrete 1N peak and the DNA content

extended beyond 2N, suggesting genomic instability or an extensive G2/M arrest in *pan2Δ* cells

after 6 hours of treatment with nocodazole (and also at 10 hours) (**Fig 3A**).

Figure 3

Fig 3: G2/M delay in *pan2Δ* cells treated with nocodazole. (A) FACS profiles post treatment

with DMSO or 6 μM nocodazole. 1N and 2N respectively represent DNA content. Green lines

represent the position of 1N and 2N DNA peaks in untreated cells. FACS, fluorescence activated

cell sorting; hr, hours; WT, wild type; Noc, nocodazole. **(B-C)** Live imaging of yeast cells

treated for 6 hours. Mother cell size quantified (n > 100). Scale bars: 5 μm. Paired Student's t-

test for significant differences. WT, wild type; ns, not significant, ****, p<0.0001.

Analysis of the gates during flow cytometry revealed that *pan2Δ* cells upon 3 hours of treatment

with nocodazole, had a different forward scatter than WT cells, suggesting that *pan2Δ* cells upon

treatment with nocodazole for 3 hours were larger than WT cells (**SFig 3A**), while there was no

difference in cell-size upon DMSO treatment (**SFig 3A**). This trend was consistent for the next

two time-points as well (6 and 10 hours post treatment with nocodazole), so we determined the

cell morphology of *pan2Δ* upon treatment with nocodazole. On treatment of log phase cells with

6 μM nocodazole for 6 hours, *pan2Δ* exhibited a significant increase in cell diameter (~1.75

times), relative to the WT cells, while there was no significant difference in size upon treatment

with DMSO (**Fig 3B**). *pan2Δ* cells treated with nocodazole were predominantly large-budded,

and together with their increased cell size, were consistent with our FACS data indicating a

G2/M delay. Upon expression of WT *PAN2* in *pan2Δ* cells, we were able to eliminate the large

cell phenotype, whereas expression of deadenylase-dead *PAN2* did not, indicating the

deadenylase activity of Pan2 was required to prevent the G2/M delay (**Fig 3C**).

Figure S3

S3A

Fig S3A: FACS profiles post DMSO or 6 µM nocodazole treatment. FACS, fluorescence

activated cell sorting; hr, hours; WT, wild type.

Nocodazole treatment leads to increased cell-death in *pan2Δ* cells due to impaired spindle

formation

We were intrigued to determine if there was a difference in spindle morphology in *pan2Δ* cells

upon nocodazole treatment. Thus, we used (Tub1+3'UTR)-yomWasabi cassette (38) to first tag

tubulin. Upon testing the sensitivity of *pan2Δ*-(Tub1+3'UTR)-yomWasabi yeast and WT-

(Tub1+3'UTR)-yomWasabi yeast, similar to *pan2Δ* yeast, we found that *pan2Δ*-(Tub1+3'UTR)-

yomWasabi yeast was sensitive to nocodazole, and that WT-(Tub1+3'UTR)-yomWasabi yeast

exhibited a mild slow-growth phenotype compared to WT (**SFig 4A**). We set up an assay to

observe spindle morphology as well as measure the total number of cells, dead cells and cell area

using live-cell imaging of yeast and analyzing the data using a machine-learning assisted

pipeline (see Methods). In untreated cells, cell growth appeared to be normal and we did not

observe significant differences in cell number or number of dead cells in WT vs *pan2Δ* cells over

8 h of imaging (**Fig 4A, left panels; also see movies in SFiles 4 and 6**). Cell area was largely

unchanged in both WT and *pan2Δ* cells over the time-course (**Fig. 4B**). Treatment with 6μM

nocodazole did not result in a significant difference in total cells or dead cells between WT and

Δ*pan2* cells, and cell area remained largely unchanged (**SFig 4B-4C**). This was different from

what we observed in liquid growth assay and we suspect was due to the different growth

conditions during live-cell imaging (no shaking or aeration), which altered the level of effective

drug exposure. Hence, we tested increased drug concentrations. At 12 μM nocodazole we

observed a general slowing of growth and a significant difference in total cells ($p<0.0001$) and

dead cells ($p<0.0001$) between WT and *pan2Δ*. For WT treated with nocodazole, the mean total

cell number was 120 at t=0 and 344.5 at t=480 minutes, while the mean number of dead cells

increased from 0.75 to 19.75 (**Fig 4A, top right panel; also see movie SFile 5**), whereas for

pan2Δ treated with nocodazole the mean total cell number was 125 at t=0 and 199 at t=480 min

and the mean number of dead cells increased from 0.25 to 27.75 (**Fig 4A, bottom right panel;**

also see movie SFile 7). We also observed a steady increase in cell area of both WT and *pan2Δ*

until ~360 min, which then leveled off (**Fig 4B**). This increase in cell area was much greater for

pan2Δ (~13 μm^2 to ~23 μm^2) than for WT (~12 μm^2 to ~17 μm^2) (**Fig 4B**). Taken together the

decreased cell growth, increased cell death and increased cell area of *pan2Δ* cells treated with

nocodazole were consistent with these cells having a defective cell-cycle.

Figure S4

340
Fig S4A: *pan2Δ* Tub1-yW yeast is sensitive to nocodazole and WT-Tub1-yW exhibits a mild

slow-growth phenotype compared to WT. Growth curves of yeast strains grown in liquid SC

media with or without 6 µM nocodazole and monitored for 24 hours. An average of 6 replicates

has been plotted with SD. Absorbance adjusted for baseline correction. µM, micromolar; SC,

Synthetic Complete media; SD, standard deviation. **Fig S4B:** Number of total cells (left y-axis)

and number of dead cells (right y-axis) upon treatment of WT (top) or *pan2Δ* (below) with 6µM

nocodazole for every 5-minutes over a total time interval of 480 minutes. **Fig S4C:** Cell area of

individual cells of WT (brown) or *pan2Δ* (yellow) is represented with 95% confidence interval

upon treatment with 6µM nocodazole over a total duration of 480 minutes. Each time point

represents the cell area of 500-2500 cells recognised by cell segmentation.

 Figure 4

4A

4B

Fig 4: **Spindle defects in *pan2Δ* upon treatment with nocodazole. (A)** Number of total cells

(left y-axis) and number of dead cells (right y-axis) upon treatment of WT or *pan2Δ* with DMSO

(left) or 12µM nocodazole (right) for every 5-minutes over a total time interval of 480 minutes

were calculated using machine-learning assisted pipelines. **(B)** Cell area of individual cells of

WT (brown) or *pan2Δ* (yellow) is represented with 95% confidence interval upon treatment with

DMSO (left) or 12µM nocodazole (right) over a total duration of 480 minutes. Each time point

represents the cell area of 500-2500 cells recognised by cell segmentation. **(C)** Live imaging of

log phase yeast cells expressing (Tub1+3'UTR)-yom-Wasabi WT or *pan2Δ*. The cells are treated

with DMSO or 12 µM nocodazole for ~75 minutes prior to imaging. The time-stamps are

indicated on top left of each snapshot. Arrow indicates new bud, arrowhead indicates mitotic

spindle and asterisk indicates dead cell. WT, wild-type; DMSO, dimethyl sulfoxide; µM,

microMolar; Tub1: alpha-tubulin; UTR, untranslated region.

Now next we examined cell morphology and tubulin-tagged spindles in detail. In untreated WT

and *pan2Δ* cells, budding and cell division appeared normal with a cell-cycle duration of ~75

min (**Fig 4C**). Spindles appeared normal, with the spindle pole body being clearly visible at the

time of bud emergence (15 min, S phase) and then migrating to the bud neck (30 min, G2 phase)

before elongating through the neck and into the bud (55 min, M phase). For nocodazole treated

cells, we chose to examine cells after 75 min of treatment with 12µM nocodazole because this

time corresponded to approximately one round of cell division and there was little cell death at

this time point. The WT cell shown had an elongated cell-cycle of ~90 min (compared to ~75

min for untreated cells), however cell morphology and spindles appeared normal (**Fig 4C**; times

indicated are relative to the start of the cell-cycle for the given cell). This was consistent with our

quantification, which indicated that over this time period cell number increased, there were few

dead cells, and there was only a marginal increase in cell size. In contrast, the *pan2Δ* cell shown

did not have clearly detectable spindle pole bodies or spindles, and although bud growth

progressed normally for ~90 min, cell division failed to complete and there was continued

mother and bud growth leading to very large cells (see 270 min). For this cell, at 300 min the

mother died and the now orphan "daughter" failed to grow or divide (as late as 380 min). This

was consistent with our quantification, which indicated that cell number did not increase

substantially after nocodazole addition, that the number of dead cells increased and there was a

substantial increase in cell size. Thus, in *pan2Δ* cells treated with nocodazole, the spindle was

compromised which led to defective mitosis and a failure to complete the cell-cycle.

pan2Δ cells exhibit dysregulated cell-cycle gene expression in response to nocodazole

Because deadenylase dead PAN2 did not eliminate the nocodazole sensitivity of *pan2Δ* cells, we

were interested in the transcriptional profile of these cells and if there were altered expression of

cell-cycle genes. Therefore, we performed an RNA-sequencing experiment on nocodazole-

treated populations of *pan2Δ* + WT Pan2 and *pan2Δ* + D1020A Pan2 cells to identify differences

in the transcriptome. Log phase cultures of the above genotypes in duplicate were treated with 6

µM of nocodazole for 6 hours and mRNA was collected and sequenced. Spearman correlations

for expression of all the genes were plotted for the two replicates of the two samples (**SFig 5A**).

S5A

S5B

S5C

Fig S5A: Spearman correlations for expression of all the genes were plotted for the two

replicates per sample of nocodazole treated Control and D1020A samples. **Fig S5B**: Growth

curves of yeast strains in SC media with DMSO (left) or 6 μM nocodazole (right) at 30°C for 24

hours. Absorbance plotted with SD (n=6). min, minutes; SC, Synthetic Complete media; SD,

standard deviation; WT, wild type. **Fig S5C**: Normalized exponential growth constants (data

from **Fig 5D**). Error bars are +/- SD from three separate experiments (n=6). Significant

differences between different strains were tested using paired Student's *t*-test. OE,

overexpression.

Out of the 6,031 ORFs analyzed, we chose an absolute z-score cutoff of 2 (equivalent to a log2

fold change cutoff of 1.182 and -1.0554). For a p value < 0.01 and log2 CPM > 1 (counts per

million), we identified 129 upregulated and 39 downregulated genes (**Fig 5A**) (**SFile 3**). These

genes were enriched in GO terms related to replication fork protection complex, arginine

biosynthetic process, carboxylic acid metabolic process, ascospore wall assembly, thiamine

biosynthetic process, hexitol metabolic process and glyoxalase III activity (**Fig 5B**). These

processes are known to be cell-cycle dependent and regulated at the transcriptional level, with

histone mRNAs (replication fork complex genes) being synthesized exclusively in S phase (39).

Hence, the change we see in the expression of these genes was consistent transcriptionally with

nocodazole-treated PAN-deficient cells having a G2/M arrest. The transcriptional upregulation

of hexitol metabolic process and glyoxalase III activity in *pan2Δ* + D1020A Pan2 compared to

pan2Δ + WT Pan2, indicates that the cells are likely undergoing a metabolic shift and producing

energy differently (hexitol metabolism) and are responding to the oxidative stress in the

environment (glyoxalase III activity). While ascospore wall assembly and thiamine biosynthetic

processes were also found to be upregulated, since these are haploid cells, it is likely that

upregulation of ascospore wall assembly mRNAs is consistent with increased demand for cell

wall synthesis during bud growth in G2/M and the increased cell-size of PAN-deficient cells.

Some of the genes involved in sporulation might have alternative functions in conditions of

nutrient limitation, environmental stress or even mutations that disrupt normal cell-cycle.

Fig 5: **mRNA levels of *CLB1* and *CLN3* are altered in deadenylase dead mutant of *PAN2* in**

nocodazole. (A) Fold change and p-value of ~6000 ORFs analyzed in nocodazole treated

Control and D1020A cells by RNA-sequencing. Each closed circle represents a single ORF,

categorized into upregulated (green), downregulated (blue) or not-significantly different (gray)

(**B**) Functional enrichment of the upregulated and downregulated genes identified in (**A**). Node

size indicates statistical significance of enriched GO terms and edges indicate statistically

significant associations between terms. Node color indicates the % of aggravating or alleviating

interactions in each term. Unspecific terms in gray. (**C**) Gene expression changes in *pan2Δ+WT*-

PAN2 or *pan2Δ+D1020A-PAN2* post 6 hour treatment with DMSO or 6 μM nocodazole

assessed by qRT-PCRs. The ΔCt values are normalized to actin. Data mean +/- S.D. 0.4X and

2X indicate fold-change in *CLB1* and *CLN3* mRNA levels respectively. n.s., not significant;

**** p < 0.0001; *** p < 0.001; **, p < 0.01. Two-way ANOVA for significant differences. Ct,

cycle threshold; μM, micromolar; qRT, quantitative reverse transcriptase; ANOVA, Analysis of

Variance. (**D**) Growth curves of yeast strains grown in liquid SC media and monitored for 24

hours at 30°C. Absorbance adjusted for baseline correction is plotted on the y-axis and time (in

minutes) is plotted on the x-axis. An average of 6 replicates have been plotted with SD. min,

minutes; SC, Synthetic Complete media; SD, standard deviation; WT, wild type; EV, empty

vector; OE, overexpression. Base 2 exponential growth constants calculated from growth of

different yeast strains. Error bars are +/- SD from three separate experiments. Each experiment

has 6 replicates. Significant differences between different strains were tested using two-way

ANOVA comparison.

Two genes, *CLN3* and *CLB1*, with functions related to regulation of cyclin-dependent protein

serine/threonine kinase (CDK) activity, were significantly different between WT and D1020A

Pan2 treated with nocodazole. *CLN3* encodes a CDK regulatory subunit that regulates cell-cycle

passage through START by regulating transcription at the G1/S transition. *CLB1* encodes a B-

type cyclin involved in the regulation of the G2/M transition of both the mitotic cell-cycle and

first meiotic division (40). *CLB1* is also involved in mitotic spindle assembly and spindle pole

body separation through its interaction with CDK1 (41). By qPCR we found that *CLB1* mRNA

levels decreased ~2.5 fold in cells with WT Pan2 in response to nocodazole, whereas in cells

with D1020A Pan2, *CLB1* decreased ~5.6 fold (**Fig 5C**). There was no difference in *CLB1*

expression between cells with WT or D1020A Pan2 in the absence of nocodazole. Similarly, for

CLN3 there was no difference between cells with WT or D1020A Pan2 in the absence of

nocodazole, however in the presence of nocodazole *CLN3* mRNA levels increased ~1.7 fold in

cells with WT Pan2 and ~3.5 fold in cells with D1020A Pan2 (**Fig 5C**). Thus, PAN was required

for regulation of *CLB1* and *CLN3* mRNA levels in response to nocodazole.

Given the changes we observed in *CLB1* and *CLN3* expression in cells with D1020A Pan2, we

tested for functional associations between PAN and *CLB1* and *CLN3*. We constructed double

mutants between *PAN2* and *CLB1* and *CLN3* and tested for genetic interactions in the absence

and presence of nocodazole. Growth of the double mutants was the same as the *PAN2* single

mutant both in the absence and presence of nocodazole, indicating deletion of either *CLB1* or

CLN3 did not modify the Δ*pan2* nocodazole sensitivity (**Fig S5B**). Next we tested

overexpression of *PAN2* in the absence of *CLB1* or *CLN3*. Similar to HsPAN2, overexpression

of yeast *PAN2* caused a slow growth phenotype in WT yeast (**Fig 5D**). This phenotype was

enhanced in cells lacking *CLB1* and *CLN3* (**Fig 5D**), indicating a functional association between

PAN and both *CLB1* and *CLN3*. Thus, these data supported that the role for PAN in response to

microtubule stress involved regulation of Clb1 and Cln3 function.

PAN2 **depletion in mammalian cells increases sensitivity to microtubule destabilizing drugs**

To determine if the function of Pan2 in regulating the cell-cycle upon microtubule stress is

conserved from yeast to mammalian cells, we implemented a human cell culture system to

examine sensitivity to microtubule destabilizing drugs and spindle integrity. We generated a

population of HEK293 cells where Pan2 was stably knocked down using an shRNA targeting the

ORF of *PAN2* (**see Methods**). Specifically, we observed ~ 75% reduction in protein levels of

Pan2 by western blot in this population (**Fig 6A**).

Fig 6: HEK293 PAN2KD cells are sensitive to nocodazole and colchicine. (**A**) Cell lysates

from HEK293 Control and Pan2KD cells were analyzed on an SDS gel and probed for Pan2 and

Actin levels, quantification for which is shown on right, n=3. Significant differences between the

2 strains were tested using paired Student's *t*-test. **, p < 0.01. (**B-C**) Absorbance at 562 nm (y-

axis) and days post treatment (x-axis) with nocodazole **(B)** or colchicine **(C)** was plotted for

control (blue) and Pan2KD (gray) at various concentrations of drug. Absorbance of each treated

condition was normalized to the absorbance of the untreated condition for each cell line on the

day 3 post-treatment to plot cell viability (% of untreated). Average of 6 replicates has been

plotted. Mean +/- S.D. Significant differences between Control and Pan2KD cells at each

concentration was tested using multiple Student's *t*-test. nM, nanomolar; n.s., not significant; *,

$p<0.05$.

To determine if this reduction in Pan2 reduced viability and/or slowed the growth of cells treated

with the microtubule inhibitors nocodazole or colchicine, we used a colorimetric MTT (3-(4, 5-

dimethylthiazol-2)-2, 5-diphenyltetrazolium bromide) assay for live, metabolically active cells

that reads out as absorbance at 562 nm (42). We seeded wells at 1,000 cells per well, added drug

16 h later and at a further 0, 24, 48 and 72 h we performed MTT assay (**Fig S6A**). We observed a

dose-dependent decrease in cell viability as measured by A562 in both control and Pan2KD cells

treated with nocodazole or colchicine over the time course (**Fig 6B &C, left panels**). We

compared the viability of Pan2KD and control cells at the endpoint of the assay on day 3. For

nocodazole treatment, we observed a significant decrease in viability of Pan2KD compared to

control cells at 40, 60 and 80 nM nocodazole (**Fig 6B, right panel**). For colchicine we observed

significant decreases at all concentrations (**Fig 6C, right panel**). This suggested that, as was the

case in yeast, depletion of Pan2 increased the vulnerability of mammalian cells to microtubule

depolymerization.

Figure S6

S6A

520
521
522
523
524   **Fig S6A**: Protocol for recording MTT activity of cells over multiple days.
525

PAN2KD cells exhibit extended metaphase and multipolar spindle formation upon

treatment with colchicine

We were curious to determine the cause of increased sensitivity of HEK293 Pan2KD cells to

microtubular depolarization and how this related to the cell-cycle. Thus, we live-imaged

microtubules in the control and Pan2KD cells using a tubulin staining dye, CellLight™ Tubulin-

RFP BacMam 2.0. Upon treatment with a low dose of 10 nM of colchicine, we imaged cells for a

period of 24 hours to capture the spindle morphology, as well as to determine the duration of

different stages of mitosis (**Fig S7A**). In the absence of drug treatment, progression through the

cell-cycle appeared normal for both control and Pan2KD cells (**Fig 7A; also see movie SFile 8**).

We observed no differences in metaphase duration or spindle morphology. In contrast, upon

treatment with 10 nM colchicine, as compared to control cells, Pan2KD showed abnormal

progression through the cell-cycle (**Fig 7A; also see movie SFile 9**). Pan2KD cells spent longer

time in metaphase (**Fig 7A**) and had a higher propensity to form multipolar spindles (~75% in

Pan2KD vs ~40% in Ctrl) (**Fig 7A**; also see 100 min image in **Fig 7A**). This indicated that Pan2

played a role in spindle integrity during microtubule stress and was consistent with the decreased

viability we observed in Pan2KD cells treated with nocodazole and colchicine.

Figure S7

S7A

546   **Fig S7A**: Protocol for live-cell imaging of colchicine treated cells using tubulin staining dye.

 Figure 7

7A

Fig 7: **HEK293 PAN2KD cells exhibit defective spindle phenotypes upon treatment with colchicine** (**A**) Representative images of HEK293 Control and Pan2KD cells with or without treatment with 10 nM colchicine. These cells are shown undergoing mitosis at the transition from G2 to M and different phases of M-phase (prophase, metaphase 1, metaphase 2, anaphase and telophase). The time taken for each stage of mitosis is indicated on top left of each snapshot. Arrows indicate cells with multipolar spindles. Metaphase duration (in minutes) and percentage of cells having 2, 3 or 4 spindles is plotted for colchicine treatment of HEK293 Control and Pan2KD cells. Each closed circle represents one cell. 50 cells were counted for each condition. Mean +/- S.D. has been plotted. nM, nanomolar; min, minutes; ns, not significant; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Discussion

Since the role of PAN in response to microtubule stress was dependent on its deadenylase activity, we propose that PAN regulates the PATLs of mRNAs required for cellular survival of microtubule stress (**Fig 8**). PAN activity during microtubule stress would lead to translational repression of key proteins involved in either sensing microtubule stress and/or responding to/recovering from formation of defective spindles. Some obvious targets are proteins involved in the spindle assembly checkpoint, however we did not find expression of any of these to be altered in our mRNA seq screen. We also did not detect altered expression of genes with functions directly related to microtubule biogenesis, including the tubulin genes themselves, making it unlikely that PAN regulates microtubules directly. However, it is possible that our screen missed these targets because it did not identify mRNAs with altered poly-A tail lengths.

Altered poly-A tail length could lead to altered translation, independent of mRNA stability,

hence screening for altered poly-A tail length would likely produce novel insights into PAN's

mechanism of action (15,43). It is interesting that cells lacking PAN appear to have a normal

cell-cycle, normal spindles, and show no decrease in fitness under regular growth conditions.

This further supports a specific role for PAN in response to microtubule/spindle stress and

perhaps explains its elusive role in the cell-cycle until now.

Figure 8

Fig 8: Model schematic of PAN regulating the mRNA polyA tails lengths (PATLs) of mRNAs

required for cellular survival of microtubule stress.

We did identify a functional connection between PAN and two important cell-cycle regulators,

Clb1 and Cln3, although the significance remains obscure. Decreased expression of *CLB1* in

response to nocodazole is consistent with failure of Δ*pan2* cells to complete mitosis, however

because *CLB1* is decreased in Δ*pan2* cells it is unlikely to be a direct target. The fact that *CLB1*

expression increases in response to nocodazole in WT yeast (which complete mitosis apparently

normally) supports that *CLB1* is part of the microtubule stress response. This is also supported by

the genetic interaction we observed between PAN and *CLB1*. It is possible that the low

expression of *CLB1* in Δ*pan2* cells treated with nocodazole prevents completion of mitosis,

however we might have expected to see loss of *CLB1* enhance the nocodazole sensitivity of

Δ*pan2* cells, which wasn't the case. It is possible that *CLB1* expression is already limiting, in

which case overexpression of *CLB1* should reduce the nocodazole sensitivity of Δ*pan2* cells if

Clb1 is playing an important role in the response. *CLN3* expression was found to be upregulated

in response to nocodazole which was enhanced in Δ*pan2* cells. Since Cln3 is not known to play

roles in mitosis it is unclear why Cln3 is being impacted. If *CLN3* was a direct target of PAN and

its elevated expression was preventing completion of mitosis, we would have expected loss of

CLN3 to suppress the nocodazole sensitivity of Δ*pan2* cells, which wasn't the case. Because we

identify a genetic interaction between PAN and *CLN3* it is likely Cln3 is functionally relevant,

perhaps indirectly by affecting cell-cycle dynamics or by regulating PAN activity outside of M

phase.

There is evidence for regulation of cell-cycle progression through regulation of PATLs. For

example, genes such as *CDK1, TOP2A, KPNA2, UBE2C* and *FBXO5*, which orchestrate timing

and coordination of mitotic events in somatic cells, have been found to have reduced PATLs in

M-phase (15). Transcription of these genes peaks at G2/M phase and is repressed at G1 phase

(44). In contrast, other well-known mitotic regulators which are also transcriptionally

upregulated in G2/M phase, such as *CCNA, CCNB*, and *CENP-E*, do not show poly(A) length

changes (44). This suggests that a subset of genes are selectively deadenylated to shut down once

cells enter M phase. Specifically, *CDK1* displayed the most extreme change, whose median

poly(A) tail length changed from 77 nt in S phase to 2 nt in M phase and displayed translational

repression (15), unlike the previous report showing translational upregulation of *CDK1* in M

phase (45). Analysis of gene ontology for genes exhibiting shortened poly(A) tails during the M

phase identified enriched terms such as "regulation of mitotic cell-cycle", "microtubule

cytoskeleton", and "regulation of ubiquitin-protein ligase activity". This implies that

deadenylation during the M phase may be linked to regulatory functions within the cell cycle. It

is currently unknown which deadenylases are responsible for which modifications and our results

suggest that PAN may play an important role. Ccr4-Not may also play a role, however, the fact

that loss of Ccr4-Not activity in yeast does not lead to sensitivity to microtubule destabilizing

drugs implies a prominent function for PAN (29).

Post-transcriptional regulation of mRNAs plays a critical role during the cell cycle and can be

achieved through their localization to specific subcellular compartments (46–48). A high-

throughput smFISH (single molecule Fluorescence *In Situ* Hybridization) screen of 602 proteins

with centrosomal functions identified 8 human mRNAs (*NIN, CEP350, PCNT, BICD2,*

CCDC88C, ASPM, NUMA1, and *HMMR)* that are locally translated at the centrosome (49).

Seven of these are translated during mitosis (*CCDC88C* is translated only during interphase).

CEP350, ASPM, NUMA1, and *HMMR* are microtubule binding proteins with functions in

centrosome/spindle maturation and *NIN, PCNT* and *BICD* contribute to microtubule anchoring at

the centrosome (50–56). Hence, these proteins are potential PAN targets in response to

microtubule stress. However, we did not find altered levels of yeast spindle pole body mRNAs in

our RNA-seq screen, suggesting either we missed detecting such changes or that PAN acts via

other targets. Given that the yeast spindle pole body shares no structural similarity to the

centrosome and that RNAs have yet to be localized to spindle pole bodies further suggests that

PAN targets components other than the centrosome/spindle pole body in yeast and humans.

The spindle phenotypes we observed in Pan2KD cells treated with colchicine coupled with their

decreased cell viability suggested these cells could have undergone abnormal chromosome

segregation. Cells with mis-segregated chromosomes often have abnormal nuclei and have

increased cell death (57–59). An imaging based whole-genome siRNA screen in HeLa cells

found that the absence of *PAN2* led to irregular nuclear shape and chromosome mis-segregation

(60). Upon chromatin extraction during G2, Pan2 and Pan3 were found to be part of a soluble

complex containing the spindle checkpoint proteins Bub1 and Mad1l1, the kinetochore proteins

Ndc80 and Nuf2, and 265 other proteins enriched for the molecular functions poly(A)-specific

ribonuclease activity, poly(A) binding, translation initiation factor activity, mRNA 3'UTR

binding, RNA-helicase activity, ribosome binding, microtubule motor activity and gamma-

tubulin binding (60). Hence, this study provides evidence that PAN's role in responding to

microtubule stress may involve direct associations between PAN complex and key spindle

regulatory proteins, which is required for proper chromosome segregation.

Previous studies have shown that the PAN localizes to P-bodies in yeast cells. P-bodies serve as

sites of mRNA storage, decay, and translational repression. Sweet et al. find that treating yeast

with the microtubule depolymerizing drug benomyl induces formation of P-bodies containing

tubulin (61). However, according to the HIPHOP chemical genetics database (29), none of the P-

body protein mutants are sensitive to nocodazole or benomyl, suggesting the role for PAN in

microtubule stress is independent of P bodies. Additionally, many groups have reported an

increase in P-body numbers during S phase, the disappearance of PB structures before mitosis,

and their reassembly during cytokinesis in mammalian cells, further supporting a non-P body

role for PAN during mitosis (62,63).

Finally, if the role of PAN is restricted to responding to microtubule stress in M-phase, this raises

the question of whether PAN activity is regulated in a cell-cycle dependent manner and/or in

response to stress. Pan3 has previously been identified as a substrate of the CDK Pho85 (24).

Different groups have identified 9 phosphorylation sites on Pan3, one in the pseudokinase

domain and the rest in unstructured regions. Holt et al. identified residue 415 in the linker

domain of Pan2 as a substrate of CDK1 phosphorylation (22). Phosphorylation of Pan3 in

mammalian cells has previously been shown to decrease its interaction with PABPC1 and reduce

PAN activity (23). Together this suggests PAN activity is regulated by phosphorylation by CDKs

in a cell-cycle dependent manner. Interestingly, our SGA screen identified an alleviating genetic

interaction between *PAN2* and *PHO85*, supporting a functional connection between PAN and

this important cellular stress response regulator. The need for cell-cycle dependent regulation of

PAN might also explain our finding that having too much Pan2 in the cell leads to reduced cell

growth, although the cause of this fitness defect has yet to be elucidated.

Materials and Methods

Yeast strains, growth conditions, and manipulations

Deletion and tagged strains were constructed in JHY716 or BY4741 (derivatives of S288C)

backgrounds using homologous recombination of PCR-generated linear fragments amplified

from pKT209, pHIS3p:yomWasabi-Tub1+3'UTR::HPH, and pFA6a-kanMX6 (38,64,65), with

successful recombination assessed by colony PCR. All yeast strains were maintained on standard

yeast growth medium of either yeast extract, peptone, and dextrose (YPD) media or synthetic

complete (SC) media consisting of Yeast Nitrogen Base (YNB) supplemented with appropriate

amino acids and 2% dextrose. Yeast cultures were grown at 30°C. A complete list of all the

strains used in this study is provided in **Table 1**.

Table 1: List of strains

Strain	Genotype	Source	Parental strain	Notes
BY4741	MAT**a**; *his3Δ1*; *leu2Δ0*; *met15Δ0*; *ura3Δ0*; S288C background	Lab collection		Wild type
BY4742	MATα; *his3Δ1*; *leu2Δ0*; *ura3Δ0*; S288C background	Lab collection		Wild type

Y7092	MATα; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0; LYS2+; can1Δ::STE2pr-Sp_his5; lypΔ1; S288C background	Kind gift from Dr. Charlie Boone		SGA query strain background
JV1	pan2Δ::uraMX	This study	Y7092	PCR knock-in
JV2	pan3Δ::uraMX	This study	Y7092	PCR knock-in
JV3	HIS3pr:yomWasabi-Tub1+3'UTR::HPH	This study	Y7092	PCR knock-in
JV4	pan2Δ::uraMX HIS3pr:yomWasabi-Tub1+3'UTR::HPH	This study	Y7092	PCR knock-in
JV5	pan2Δ::uraMX cin1Δ::kanMX6	This study	SGA background	SGA
JV6	pan2Δ::uraMX cin2Δ::kanMX6	This study	SGA background	SGA
JV7	pan2Δ::uraMX cin4Δ::kanMX6	This study	SGA background	SGA
JV8	pan2Δ::uraMX gim3Δ::kanMX6	This study	SGA background	SGA
JV9	pan2Δ::uraMX gim4Δ::kanMX6	This study	SGA background	SGA
JV10	pan2Δ::uraMX pac2Δ::kanMX6	This study	SGA background	SGA

JV11	*pan2Δ::uraMX tub3Δ::* kanMX6	This study	SGA background	SGA

Plasmids and constructs

CEN plasmids were taken from the MoBY-ORF collection (66). Galactose-driven

overexpression plasmids were obtained from the FLEX overexpression library (67) generously

provided by Tejomayee Singh from the laboratory of Dr. Phil Hieter. All plasmids were

sequenced to confirm their identity (Plasmidsaurus). The coding sequence of *Saccharomyces*

cerevisiae PAN2 gene, along with its native promoter and 3'UTR sequence, and the codon

optimized version of *Homo sapiens PAN2* (sequence is provided in **SFile 1**) were purchased

from Twist Biosciences to insert into various plasmids for tagging or overexpression using the

Gateway cloning kit (ThermoFisher Scientific). These plasmids were pAG426*GAL*-ccdB,

pAG426-*GAL*-GFP-ccdB, pAG425-*GAL*-ccdB, pAG415-*GPD*-ccdB, and pAG425-*GPD*-ccdB.

Details of these plasmids are provided in **Table 2**.

Table 2: List of plasmids

Plasmid name	Description	Source
pKT0209/pKT209	pFA6a–link–yEGFP–Ca*URA3*; used for URA amplification for making gene deletion cassettes	Addgene (64)
pHIS3pr:yomWasabi -Tub1+3'UTR::HPH	*pHIS3pr*:yomWasabi-Tub1+3'UTR::HPH: used for tagging alpha-tubulin	Addgene (38)

pFA6a-kanMX6	for one-step PCR deletion of gene of interest	Addgene (65)
pAG426-*GAL*-Pan2	URA3 marker, multi-copy, GAL promoter	This study
pAG426-*GAL*-Pan2-D1020A	URA3 marker, multi-copy, GAL promoter	This study
pAG426-*GAL*-GFP-Pan2	URA3 marker, multi-copy, GAL promoter, N-terminal GFP-tagging	This study
pAG426-*GAL*-GFP-Pan2-D1020A	URA3 marker, multi-copy, GAL promoter, N-terminal GFP-tagging	This study
pAG425-*GAL*-Pan2	LEU2 marker, multi-copy, GAL promoter	This study
pAG425-*GAL*-Pan2-D1020A	LEU2 marker, multi-copy, GAL promoter	This study
pAG425-*GAL*-HsPan2	LEU2 marker, multi-copy, GAL promoter	This study
pAG415-*GPD*-Pan2	LEU2 marker, low copy, GPD promoter	This study
pAG415-*GPD*-Pan2-D1020A	LEU2 marker, low copy, GPD promoter	This study
pAG425-*GPD*-Pan2	LEU2 marker, multi-copy, GPD promoter	This study
pAG425-*GPD*-Pan2-D1020A	LEU2 marker, multi-copy, GPD promoter	This study
pFLEX-Pan2	*GAL1-10pr, URA3*	A gift from Dr. Phil Hieter (67)

The deadenylase dead mutation (D1020A) in *PAN2*'s coding sequence was generated by site-

directed mutagenesis (Q5 Site Directed Mutagenesis Kit from NEB E0554) with primer pairs 5'

GGTTTGAATAATGCCTTCAAACACATTAATATTAATGTC 3' and 5'

ATGACCAACAAATACATTATTCAAACCATGACCAAC 3'. These PCR products were

recombined in DHFpir bacteria which was a generous gift from Dr. J.T. Beatty.

For tagging Tub1 with yomWasabi-Tub1+3'UTR, the strategy used has been described in

Markus et al., 2015 (38). Briefly, the yomWasabi-Tub1 tagging plasmid was altered by adding

an additional 618 nucleotides corresponding to the region downstream of the *TUB1* locus. This

sequence was chosen because it extends well beyond the last potential polyadenylation sequence

(AATAAA, 145 nt downstream of the stop codon; ATTAAA, 540 nt downstream of the stop

codon). Following transformation with the BsaBI-digested plasmid (*pHIS3p*:yomWasabi-

TUB1+3'UTR), both the endogenous, untagged *TUB1* and the tagged *TUB1* genes are

immediately followed by the 3'UTR sequence.

Quantitative liquid growth assays

Yeast were grown overnight in liquid SC media supplemented with appropriate amino acids,

diluted into fresh medium, then grown until reaching logarithmic growth phase (OD_{600} 0.2-0.5).

Yeast cells were then transferred to liquid media supplemented with an appropriate concentration

of nocodazole (Sigma) or equivalent percentage (<0.5%) of DMSO (ThermoFisher). Strains

were normalized to the same OD_{600} then diluted to an OD_{600} value of 0.005-0.010 in 96-well

plates in triplicate wells to a volume of 200 µL. Plates were then incubated at 30°C with orbital

shaking at 600 RPM in a Biotek Instruments Epoch 2 Microplate Spectrophotometer. Data was

$\log_2$ transformed and appropriate time points were fitted to a linear equation, with exponential

growth constants derived from the slopes of the linear equation (except in drug treatments where

yeast showed a non-exponential growth pattern). Statistical analysis was performed in GraphPad

Prism 10, and significant differences between pairs were detected using Student's *t*-test or

between pairwise combinations of groups using a one-way ANOVA with *post-hoc* multiple

comparisons corrections using Tukey's method.

Western blotting

Yeast cell lysates were prepared using the TCA (Trichloroacetic acid) precipitation method (68).

The following primary antibodies were used: chicken anti-GFP (1:5000 of Abcam ab13970) and

mouse anti-PGK1 (1:5000 of Abcam ab113687). The following secondary antibodies were used:

donkey anti-chicken Alexafluor 488 (1:10,000 of Jackson ImmunoReserach Lab 703-545-155)

and goat anti-mouse DyLight 800 (1:15,000 of Invitrogen SA5-10176). The experiments were

performed in triplicates, and the results were quantified using the Image Lab software (BioRad).

Mammalian cells were lysed using RIPA (Radio Immuno Precipitation Assay) buffer (50 mM

Tris–HCl (pH 7.4), 150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS,

and 5 mM EDTA) supplemented with cOmplete protease inhibitors (Roche). The following

primary antibodies were used: rabbit anti-PAN2 (1:1000 of Proteintech 16427-1-AP) and mouse

anti-actin (1:5000 of Abcam ab8226). The following secondary antibodies were used: goat anti-

mouse HRP (Jackson 115-035-003) and goat anti-rabbit HRP (Jackson 111-035-003).

Synthetic Genetic Array Analysis

Creation of the genome-wide array of double mutants with the *pan2Δ* deletion strain was

undertaken as previously described (69,70). Briefly, *PAN2* was knocked-out in the SGA query

strain background Y7092 by genomic integration of the *URA3* cassette at the *PAN2* genomic

locus amplified from pKT209. This *pan2Δ* query strain was mated to an array of non-essential

deletion mutants marked with kanMX6 using a Singer RoToR HDA robot. Diploids were

selected for and induced to undergo sporulation. From spores, *MAT***a** cells were obtained by

pinning onto haploid selection media (SC-His/Arg/Lys supplemented with 100 μg/mL

canavanine and 100 μg/mL thialysine). Control plates were obtained by re-pinning to 5-FOA

media to counter select for the *PAN2* deletion, while experimental plates were generated by

plating on media lacking uracil. These were then separately pinned onto plates containing 1.5μM

of nocodazole. Plates were imaged using a CanoScan flatbed scanner and were analyzed using

Balony software (69). Gene Ontology (GO) term enrichment and protein-protein interaction

network analysis was performed using the GeneMANIA web server, with GO term enrichment

presented as Benjamini-Hochberg FDR-adjusted p values (q values) from hypergeometric tests

(71,72).

FACS analysis of DNA content

Yeast cells were cultured under specified conditions, fixed in 70% ethanol, and stored at 4°C.

Fixed cells were rehydrated in phosphate-buffered saline, incubated with 100 μg/mL RNase A

(ThermoFisher) for 3 hrs, and stained with 5 μg/mL propidium iodide (Invitrogen) for 1 hour.

Cells were analyzed using a Cytoflex LX Analyser, and yeast cells were separated from debris

by gating with forward scatter and side scatter. The experiments were repeated three times using

freshly grown cultures.

Confocal microscopy

Log phase (OD_{600} 0.2-0.5) yeast cells were centrifuged at 2,000 g and imaged in sterile filtered

SC liquid medium (yeast nitrogen base without amino acids and with ammonium sulphate,

amino acid mix, dextrose). Squashes of yeast were made and imaged on the Leica SP5 system

with DIC imaging. Images were analyzed using the Fiji distribution (73) of ImageJ software

(74).

Sample preparation for live-cell imaging of yeast and cell-segmentation

Live-cell imaging of yeast was conducted in Concanavalin A (ConA) (Sigma L7647) coated 96-

well microscope dishes using the Cell Discoverer 7 imaging system. To prepare the ConA-coated

dishes, a stock solution of ConA at a concentration of 10 mg/mL was initially prepared and

stored in aliquots at -20°C for a maximum of 6 months. A working solution was then prepared

by diluting the stock to a final concentration of 1 mg/mL and stored at 4°C for several weeks.

The ConA solution was evenly applied to each well of the 96-well dishes (Cellvis, P96-1.5H-N).

Following a 5-minute incubation on the benchtop, excess solution was aspirated, and the dishes

were allowed to dry for 5 minutes. For sample preparation, yeast cultures were grown overnight

in SC medium (sterile filtered). The following day, yeast cultures were diluted and allowed to

grow to log phase. Cells were adhered to the ConA-coated plates by applying 20 μL of the cell

suspension to each well. After 10 minutes of incubation to ensure proper adhesion, the media

was removed, and 40 μL of fresh media without cells was added to eliminate potential optical

artifacts. The CellDiscoverer7 (Carl Zeiss) microscope was set up prior to starting the live-cell

imaging to reach 30°C and allow the objectives to acclimate to that temperature. An imaging

time course was set up using ZEN Blue software (Carl Zeiss) to take images at 2 evenly

dispersed, non-overlapping locations per well. For each condition, 2 biological replicates were

imaged in separate wells. Images were taken in brightfield (TL LED lamp) and GFP (LED

module 470nm) channels. The Plan Apochromat water immersion 50x/1.2 objective and 1x

Tubelens were used for a final magnification of 50x, with an effective NA of 1.2. Images were

taken every 5 minutes for 480 minutes using definite focus. The individual images were

deconvolved using the constrained iterative algorithm and max intensity images were generated

using extended depth of focus with an effective depth of ~1μm (9 z-stacks of ~0.125μm). Images

were analyzed using the ZEN Blue software (Carl Zeiss). GFP segmentation was performed to

collect data for individual cells after Gaussian smoothing, sigma=2, using iso data thresholding

for light regions. For each GFP segmented cell, area and GFP mean intensity were measured.

Data for the cells in each of the 4 images per well were compiled to generate the total cell

number, number of dead cells (brighter, GFP mean intensity > 8000) and cell area.

RNA isolation and qRT-PCR analysis

Total RNA was isolated from cells using the RNeasy kit from Qiagen (74104) following the

manufacturer's instructions. Quantitative reverse-transcriptase PCR was performed using Power

SYBR® Green RNA-to-CT™ 1-Step Kit from ThermoFisher (4389986). All data were

normalized to *ACT1* transcript levels and are presented as ΔCt values. Where a statistically

significant change was observed, fold-increase against control was indicated in the plots. The

primers used in this study are listed in **Table 3**.

Table 3: RT-PCR primers

Primer	Sequence
Sc_Act1_qFwd	TTTGCCGGTGACGACGCTCC
Sc_Act1_qRev	ACAGGGTGTTCTTCTGGGGCA
Sc_Clb1_qFwd	CAGTCTAGGACGTTAGC
Sc_Clb1_qRev	GTCGTGAATAGTAGATCC
Sc_Cln3_qFwd	TCAGCGCTGCCTCATGTC
Sc_ Cln3_qRev	ATGGCCCGCCTCTTTTG
Sc_ Tub1_qFwd	GGCCACTGTGGATAGGGCCG
Sc_ Tub1_qRev	TCAGCGTATGAGTCGGCACCC
Sc_ Tub2_qFwd	CGTCGATCTAGAACCTGGGA
Sc_ Tub2_qRev	GACGCTGTCTACAAGCTCAG
Sc_ Tub3_qFwd	TGGCCGACCAATGTGACGGT
Sc_ Tub3_qRev	AGGCGCAGGATAAACGGCGA

RNA sequencing and analysis

Sample quality control was performed using the Agilent 2100 Bioanalyzer. Qualifying samples

were then prepped following the standard protocol for the Illumina Stranded mRNA prep

(Illumina). Sequencing was performed on the Illumina NextSeq2000 with Paired End 59bp ×

59bp reads by BRC-sequencing facility. As previously described (75), multiple differential

expression analysis pipelines were applied, and the results combined. The pipelines used the

reference genome and transcriptome R64-1-1 downloaded from Ensembl (www.ensembl.org,

Release 107).

Generation of stable PAN2 knockdown mammalian HEK293 cell populations

HEK293A cells were cultured in DMEM/F12 medium (Sigma) supplemented with 10% FBS and

grown in an incubator at 5% CO_2 and 37°C and transfected with Dharmacon pGIPZ shRNA set

(RHS4531-EG9924; shRNAs fused to GFP) targeting PAN2 ORF/3'UTR using Lipofectamine

2000 according to the manufacturer's protocol (ThermoFisher Scientific 11668027). Puromycin

selection was initiated 48 hours post-transfection at a concentration of 1 µg/mL to generate

stably integrated shRNA-mediated knockdowns. After 14 days of selection, the puromycin-

resistant cells were subjected to FACS to isolate a stably transfected population of GFP-positive

cells. Pan2 knockdown efficiency was then determined by Western blotting (1:1000 of

Proteintech 16427-1-AP against human Pan2).

Mammalian cell viability assessment using MTT

HEK293 cells were plated in 96-well plates at 10,000 cells/well. The number of viable cells was

quantified using a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT; M2128,

Sigma) assay at 0, 24, 48, and 72 h. After the designated treatment period, the culture medium

was removed, MTT was added to each well at 1 mg/mL in regular growth media for 2 h to allow

the formation of formazan crystals (purple precipitate). Following incubation, the MTT solution

was aspirated, and the formazan crystals were solubilized using 100 μL of DMSO per well. The

plate was gently agitated to ensure complete dissolution of the formazan crystals. The

absorbance was measured at 560 nm using a spectrophotometer plate, with DMSO serving as the

blank control. Cell viability was calculated as the percentage of viable cells relative to the

untreated control group, and the results were expressed as mean ± standard deviation (SD) of at

least three independent experiments. Statistical analysis was performed in GraphPad.

Live-cell imaging of mammalian cell spindle formation and mitosis

For spindle analysis of living cells, cells were grown in a 96-well plate (Corning) and a tubulin

staining dye, CellLight™ Tubulin-RFP BacMam 2.0 was added prior to imaging the cells in a

37°C and 5% CO_2 environmental chamber (ImageXpress Micro XL). Images were taken using a

40X 0.75 NA dry objective with the MetaXpress 5.0.2.0 software on the ImageXpress Micro XL

epifluorescence microscope (Molecular Devices Inc.). Images were taken every 15 minutes using

100-ms exposures, 2x2 binned resolution, with 25% of full lamp intensity per channel. Movies of

the fluorescence channels were made in the MetaXpress 5.0.2.0 software (Molecular Devices

Inc.).

Acknowledgements

We thank members of the Loewen and Roskelley labs for critical discussions. We thank Dr. Phil

Hieter and Dr. J. Thomas Beatty for sharing reagents. We thank the following core facilities at

UBC: LSI imaging facility, ubcFLOW, and School of Biomedical Engineering (SBME)

Sequencing Core (formerly BRC-Seq). Research was funded by the Canadian Institutes of

Health Research (PJT-152967) and CFI Innovation Fund (2020) #39914 awarded to CJRL. This

study was partially funded by the Canadian Institutes of Health Research (CIHR 169111)

awarded to CDM and Canadian Institutes of Health Research (PJT-162253) awarded to LJH.

Author contributions

JV, CDR and CJRL designed the experiments. JV performed the majority of the experiments.

ZH collected live-cell imaging data for HEK293 cells, under the supervision of CDM. JARB

collected the flow-cytometry data for cell-cycle analysis in yeast cells, under the supervision of

LJH. PD confirmed HEK293 Pan2KD GFP positive cells by flow cytometry. BPY performed the

initial testing of sensitivity of *pan2Δ* to nocodazole. SF performed the bioinformatics analysis on

RNA-seq data. JV, ZH, JARB, PD, BPY, CDM, CDR and CJRL analyzed the data. JV and

CJRL wrote the manuscript. CDM and CDR edited the manuscript.

Competing interests

The authors declare no competing financial interests.

References

1. Jackson RJ, Standart N. Do the poly(A) tail and 3′ untranslated region control mRNA

translation? Cell. 1990;62(1):15–24.

2. Sachs AB, Kornberg RD. Purification and characterization of polyadenylate-binding protein.

Methods Enzym. 1990;181:332–52.

3. Zlotorynski E. The short tail that wags the mRNA. Nat Rev Mol Cell Biol. 2018;19(1):2–3.

4. Nicholson AL, Pasquinelli AE. Tales of Detailed Poly(A) Tails. Trends Cell Biol.

2019;29(3):191–200.

5. Tang TTL, Passmore LA. Recognition of Poly(A) RNA through Its Intrinsic Helical Structure.

Cold Spring Harb Symp Quant Biol. 2020;84:039818.

6. Passmore LA, Coller J. Roles of mRNA poly(A) tails in regulation of eukaryotic gene

expression. Nat Rev Mol Cell Biol. 2022;23(2):93–106.

7. Edmonds M, Vaughan MH, Nakazato H. Polyadenylic Acid Sequences in the Heterogeneous

Nuclear RNA and Rapidly-Labeled Polyribosomal RNA of HeLa Cells: Possible Evidence for a

Precursor Relationship. Proc Natl Acad Sci. 1971;68(6):1336–40.

8. McLaughlin CS, Warner JR, Edmonds M, Nakazato H, Vaughan MH. Polyadenylic Acid

Sequences in Yeast Messenger Ribonucleic Acid. J Biol Chem. 1973;248(4):1466–71.

9. Weill L, Belloc E, Bava FA, Méndez R. Translational control by changes in poly(A) tail

length: recycling mRNAs. Nat Struct Mol Biol. 2012;19(6):577–85.

10. Jalkanen AL, Coleman SJ, Wilusz J. Determinants and implications of mRNA poly(A) tail

size – Does this protein make my tail look big? Semin Cell Dev Biol. 2014;34:24–32.

11. Boeck R, Tarun S, Rieger M, Deardorff JA, Müller-Auer S, Sachs AB. The Yeast Pan2

Protein Is Required for Poly(A)-binding Protein-stimulated Poly(A)-nuclease Activity ∗. J Biol

Chem. 1996;271(1):432–8.

12. Brown CE, Tarun SZ, Boeck R, Sachs AB. PAN3 Encodes a Subunit of the Pab1p-

Dependent Poly(A) Nuclease in Saccharomyces cerevisiae. Mol Cell Biol. 1996;16(10):5744–53.

13. Yamashita A, Chang TC, Yamashita Y, Zhu W, Zhong Z, Chen CYA, et al. Concerted action

of poly(A) nucleases and decapping enzyme in mammalian mRNA turnover. Nat Struct Mol

Biol. 2005;12(12):1054–63.

14. Yi H, Park J, Ha M, Lim J, Chang H, Kim VN. PABP Cooperates with the CCR4-NOT

Complex to Promote mRNA Deadenylation and Block Precocious Decay. Mol Cell.

2018;70(6):1081-1088.e5.

15. Park JE, Yi H, Kim Y, Chang H, Kim VN. Regulation of Poly(A) Tail and Translation

during the Somatic Cell Cycle. Mol Cell. 2016;62(3):462–71.

16. Jonas S, Christie M, Peter D, Bhandari D, Loh B, Huntzinger E, et al. An asymmetric PAN3

dimer recruits a single PAN2 exonuclease to mediate mRNA deadenylation and decay. Nat

Struct Mol Biol. 2014;21(7):599–608.

17. Schäfer IB, Rode M, Bonneau F, Schüssler S, Conti E. The structure of the Pan2–Pan3 core

complex reveals cross-talk between deadenylase and pseudokinase. Nat Struct Mol Biol.

2014;21(7):591–8.

18. Wolf J, Valkov E, Allen MD, Meineke B, Gordiyenko Y, McLaughlin SH, et al. Structural

basis for Pan3 binding to Pan2 and its function in mRNA recruitment and deadenylation. EMBO

J. 2014;33(14):1514–26.

19. Yang S, Liu L, Cao C, Song N, Wang Y, Ma S, et al. USP52 acts as a deubiquitinase and

promotes histone chaperone ASF1A stabilization. Nat Commun. 2018;9(1):1285.

20. Gao M, Guo G, Huang J, Kloeber JA, Zhao F, Deng M, et al. USP52 regulates DNA end

resection and chemosensitivity through removing inhibitory ubiquitination from CtIP. Nat

Commun. 2020;11(1):5362.

21. Christie M, Boland A, Huntzinger E, Weichenrieder O, Izaurralde E. Structure of the PAN3

Pseudokinase Reveals the Basis for Interactions with the PAN2 Deadenylase and the GW182

Proteins. Mol Cell. 2013;51(3):360–73.

22. Holt LJ, Tuch BB, Villén J, Johnson AD, Gygi SP, Morgan DO. Global Analysis of Cdk1

Substrate Phosphorylation Sites Provides Insights into Evolution. Science.

2009;325(5948):1682–6.

23. Huang KL, Chadee AB, Chen CYA, Zhang Y, Shyu AB. Phosphorylation at intrinsically

disordered regions of PAM2 motif-containing proteins modulates their interactions with

PABPC1 and influences mRNA fate. RNA. 2013;19(3):295–305.

24. Dephoure N, Howson RW, Blethrow JD, Shokat KM, O'Shea EK. Combining chemical

genetics and proteomics to identify protein kinase substrates. Proc Natl Acad Sci.

2005;102(50):17940–5.

25. Huang D, Patrick G, Moffat J, Tsai LH, Andrews B. Mammalian Cdk5 is a functional

homologue of the budding yeast Pho85 cyclin-dependent protein kinase. Proc National Acad Sci.

1999;96(25):14445–50.

26. Bett JS, Ibrahim AFM, Garg AK, Kelly V, Pedrioli P, Rocha S, et al. The P-body component

USP52/PAN2 is a novel regulator of HIF1A mRNA stability. Biochem J. 2013;451(Pt 2):185–

94.

27. Hammet A, Pike BL, Heierhorst J. Posttranscriptional regulation of the RAD5 DNA repair

gene by the Dun1 kinase and the Pan2-Pan3 poly(A)-nuclease complex contributes to survival of

replication blocks. J Biol Chem. 2002;277(25):22469–74.

28. Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, et al. A global

genetic interaction network maps a wiring diagram of cellular function. Science.

2016;353(6306):aaf1420–aaf1420.

29. Lee AY, St.Onge RP, Proctor MJ, Wallace IM, Nile AH, Spagnuolo PA, et al. Mapping the

Cellular Response to Small Molecules Using Chemogenomic Fitness Signatures. Science.

2014;344(6180):208–11.

30. Hoyt MA, Stearns T, Botstein D. Chromosome Instability Mutants of Saccharomyces

cerevisiae That Are Defective in Microtubule-Mediated Processes. Mol Cell Biol.

1990;10(1):223–34.

31. Hoyt MA, Macke JP, Roberts BT, Geiser JR. Saccharomyces cerevisiae PAC2 Functions

With CIN1, 2 and 4 in a Pathway Leading to Normal Microtubule Stability. Genetics.

1997;146(3):849–57.

32. Bhamidipati A, Lewis SA, Cowan NJ. Adp Ribosylation Factor-like Protein 2 (Arl2)

Regulates the Interaction of Tubulin-Folding Cofactor D with Native Tubulin. J Cell Biol.

2000;149(5):1087–96.

33. Geiser JR, Schott EJ, Kingsbury TJ, Cole NB, Totis LJ, Bhattacharyya G, et al.

Saccharomyces cerevisiae genes required in the absence of the CIN8-encoded spindle motor act

in functionally diverse mitotic pathways. Mol Biol Cell. 1997;8(6):1035–50.

34. Amberg DC, Burke DJ, Strathern JN. Inducing yeast cell synchrony: nocodazole arrest. CSH

Protoc. 2006;2006(1):pdb.prot4175.

35. Schäfer IB, Yamashita M, Schuller JM, Schüssler S, Reichelt P, Strauss M, et al. Molecular

Basis for poly(A) RNP Architecture and Recognition by the Pan2-Pan3 Deadenylase. Cell.

2019;177(6):1619-1631.e21.

36. Tian G, Huang Y, Rommelaere H, Vandekerckhove J, Ampe C, Cowan NJ. Pathway Leading

to Correctly Folded β-Tubulin. Cell. 1996;86(2):287–96.

37. Siegers K, Waldmann T, Leroux MR, Grein K, Shevchenko A, Schiebel E, et al.

Compartmentation of protein folding in vivo: sequestration of non-native polypeptide by the

chaperonin–GimC system. EMBO J. 1999;18(1):75–84.

38. Markus SM, Omer S, Baranowski K, Lee W. Improved Plasmids for Fluorescent Protein

Tagging of Microtubules in Saccharomyces cerevisiae. Traffic. 2015;16(7):773–86.

39. Hereford LM, Osley MA, Ludwig JR, McLaughlin CS. Cell-cycle regulation of yeast histone

mRNA. Cell. 1981;24(2):367–75.

40. Fitch I, Dahmann C, Surana U, Amon A, Nasmyth K, Goetsch L, et al. Characterization of

four B-type cyclin genes of the budding yeast Saccharomyces cerevisiae. Mol Biol Cell.

1992;3(7):805–18.

41. Richardson H, Lew DJ, Henze M, Sugimoto K, Reed SI. Cyclin-B homologs in

Saccharomyces cerevisiae function in S phase and in G2. Genes Dev. 1992;6(11):2021–34.

42. Kamiloglu S, Sari G, Ozdal T, Capanoglu E. Guidelines for cell viability assays. Food Front.

2020;1(3):332–49.

43. Xiang K, Bartel DP. The molecular basis of coupling between poly(A)-tail length and

translational efficiency. eLife. 2021;10:e66493.

44. Müller GA, Wintsche A, Stangner K, Prohaska SJ, Stadler PF, Engeland K. The CHR site:

definition and genome-wide identification of a cell cycle transcriptional element. Nucleic Acids

Res. 2014;42(16):10331–50.

45. Stumpf CR, Moreno MV, Olshen AB, Taylor BS, Ruggero D. The Translational Landscape

of the Mammalian Cell Cycle. Mol Cell. 2013;52(4):574–82.

46. Czaplinski K, Singer RH. Pathways for mRNA localization in the cytoplasm. Trends

Biochem Sci. 2006;31(12):687–93.

47. Engel KL, Arora A, Goering R, Lo HG, Taliaferro JM. Mechanisms and consequences of

subcellular RNA localization across diverse cell types. Traffic. 2020;21(6):404–18.

48. Loedige I, Baranovskii A, Mendonsa S, Dantsuji S, Popitsch N, Breimann L, et al. mRNA

stability and m6A are major determinants of subcellular mRNA localization in neurons. Mol

Cell. 2023;83(15):2709-2725.e10.

49. Safieddine A, Coleno E, Salloum S, Imbert A, Traboulsi AM, Kwon OS, et al. A

choreography of centrosomal mRNAs reveals a conserved localization mechanism involving

active polysome transport. Nat Commun. 2021;12(1):1352.

50. Karasu OR, Neuner A, Atorino ES, Pereira G, Schiebel E. The central scaffold protein

CEP350 coordinates centriole length, stability, and maturation. J Cell Biol.

2022;221(12):e202203081.

51. Jiang K, Rezabkova L, Hua S, Liu Q, Capitani G, Altelaar AFM, et al. Microtubule minus-

end regulation at spindle poles by an ASPM–katanin complex. Nat Cell Biol. 2017;19(5):480–

92.

52. Sun M, Jia M, Ren H, Yang B, Chi W, Xin G, et al. NuMA regulates mitotic spindle

assembly, structural dynamics and function via phase separation. Nat Commun.

2021;12(1):7157.

53. He Z, Mei L, Connell M, Maxwell CA. Hyaluronan Mediated Motility Receptor (HMMR)

Encodes an Evolutionarily Conserved Homeostasis, Mitosis, and Meiosis Regulator Rather than

a Hyaluronan Receptor. Cells. 2020;9(4):819.

54. Vineethakumari C, Lüders J. Microtubule Anchoring: Attaching Dynamic Polymers to

Cellular Structures. Front Cell Dev Biol. 2022;10:867870.

55. Delgehyr N, Sillibourne J, Bornens M. Microtubule nucleation and anchoring at the

centrosome are independent processes linked by ninein function. J Cell Sci. 2005;118(8):1565–

75.

56. Panic M, Hata S, Neuner A, Schiebel E. The Centrosomal Linker and Microtubules Provide

Dual Levels of Spatial Coordination of Centrosomes. PLoS Genet. 2015;11(5):e1005243.

57. He Z, Wilson A, Rich F, Kenwright D, Stevens A, Low YS, et al. Chromosomal instability

and its effect on cell lines. Cancer Rep. 2023;6(6):e1822.

58. Nicholson JM, Macedo JC, Mattingly AJ, Wangsa D, Camps J, Lima V, et al. Chromosome

mis-segregation and cytokinesis failure in trisomic human cells. eLife. 2015;4:e05068.

59. Bona MD, Bakhoum SF. Micronuclei and Cancer. Cancer Discov. 2024;14(2):214–26.

60. Cai Y, Hossain MJ, Hériché JK, Politi AZ, Walther N, Koch B, et al. Experimental and

computational framework for a dynamic protein atlas of human cell division. Nature.

2018;561(7723):411–5.

61. Sweet TJ, Boyer B, Hu W, Baker KE, Coller J. Microtubule disruption stimulates P-body

formation. RNA. 2007;13(4):493–502.

62. Aizer A, Kafri P, Kalo A, Shav-Tal Y. The P Body Protein Dcp1a Is Hyper-phosphorylated

during Mitosis. Plos One. 2013;8(1):e49783.

63. Luo Y, Na Z, Slavoff SA. P-Bodies: Composition, Properties, and Functions. Biochemistry.

2018;57(17):2424–31.

64. Sheff MA, Thorn KS. Optimized cassettes for fluorescent protein tagging in Saccharomyces

cerevisiae. Yeast. 2004;21(8):661–70.

65. Bähler J, Wu J, Longtine MS, Shah NG, III AM, Steever AB, et al. Heterologous modules

for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe. Yeast.

1998;14(10):943–51.

66. Ho CH, Magtanong L, Barker SL, Gresham D, Nishimura S, Natarajan P, et al. A molecular

barcoded yeast ORF library enables mode-of-action analysis of bioactive compounds. Nat

Biotechnol. 2009;27(4):369–77.

67. Douglas AC, Smith AM, Sharifpoor S, Yan Z, Durbic T, Heisler LE, et al. Functional

Analysis With a Barcoder Yeast Gene Overexpression System. G3: GenesGenomesGenet.

2012;2(10):1279–89.

68. Cox JS, Chapman RE, Walter P. The unfolded protein response coordinates the production of

endoplasmic reticulum protein and endoplasmic reticulum membrane. Mol Biol Cell.

1997;8(9):1805–14.

69. Young BP, Loewen CJR. Balony: a software package for analysis of data generated by

synthetic genetic array experiments. BMC Bioinform. 2013;14:354.

70. Tong AHY, Boone C. Yeast Protocol. Methods Mol Biol (Clifton, NJ). 2005;313:171–91.

71. Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al. The

GeneMANIA prediction server: biological network integration for gene prioritization and

predicting gene function. Nucleic Acids Res. 2010;38(suppl_2):W214–20.

72. Zuberi K, Franz M, Rodriguez H, Montojo J, Lopes CT, Bader GD, et al. GeneMANIA

Prediction Server 2013 Update. Nucleic Acids Res. 2013;41(W1):W115–22.

73. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676–82.

74. Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, et al. ImageJ2:

ImageJ for the next generation of scientific image data. BMC Bioinform. 2017;18(1):529.

75. Vuilleumier R, Lian T, Flibotte S, Khan ZN, Fuchs A, Pyrowolakis G, et al. Retrograde BMP

signaling activates neuronal gene expression through widespread deployment of a conserved

BMP-responsive cis-regulatory activation element. Nucleic Acids Res. 2019;47(2):679–99.

1090

Supporting information captions

SFile 1.docx: Sequence of human PAN2 codon optimized for yeast

SFile 2.xlsx: Synthetic Genetic Analysis (SGA) data for *PAN2* in presence or absence of nocodazole

SFile 3.xlsx: RNA-seq data for *pan2Δ* + WT Pan2 and *pan2Δ* + D1020A Pan2 in presence of nocodazole

SFile 4.avi: Yeast WT with DMSO

SFile 5.avi: Yeast WT with nocodazole

SFile 6.avi: Yeast *pan2Δ* with DMSO

SFile 7.avi: Yeast *pan2Δ* with nocodazole

SFile 8.avi: HEK293 cells - bipolar spindle formation

SFile 9.avi: HEK293 cells - tetrapolar spindle formation