Title: Cell type and cell signaling innovations underlying mammalian pregnancy

Authors:
Daniel J. Stadtmauer*†[1,2,3], Silvia Basanta Martínez*†[3], Jamie D. Maziarz[1], Alison G. Cole[4], Gülay Dagdas[3], Gilbecca Rae Smith[5], Frank van Breukelen[5], Mihaela Pavličev*[3,6], Günter P. Wagner* [1,3,7]

Affiliations:
1. Department of Ecology & Evolutionary Biology, Yale University, New Haven, CT, USA
2. Konrad Lorenz Institute for Evolution and Cognition, Klosterneuburg, Austria
3. Department of Evolutionary Biology, University of Vienna, Vienna, Austria
4. Department of Neuroscience and Developmental Biology, University of Vienna, Vienna, Austria
5. School of Life Sciences, University of Nevada, Las Vegas, Las Vegas, NV, USA
6. Complexity Science Hub Vienna, Vienna, Austria
7. Department of Animal Science, Texas A&M University, College Station, TX, USA

* Corresponding authors:

daniel.stadtmauer@yale.edu

silvia.basanta@univie.ac.at

mihaela.pavlicev@univie.ac.at

gunter.wagner@yale.edu

† These authors contributed equally

Abstract

How fetal and maternal cell types have co-evolved to enable mammalian placentation poses a unique evolutionary puzzle. Here, we present a multi-species atlas integrating single-cell transcriptomes from six species bracketing therian mammal diversity. We find that invasive trophoblasts share a gene-expression signature across eutherians, and evidence that endocrine decidual cells evolved stepwise from an immunomodulatory cell type retained in *Tenrec* with affinity to human decidua of menstruation. We recover evolutionary patterns in ligand-receptor signaling: fetal and maternal cells show a pronounced tendency towards disambiguation, but a predicted arms race dynamic between them is limited. We reconstruct cell communication networks of extinct mammalian ancestors, finding strong integration of fetal trophoblast into maternal networks. Together, our results reveal a dynamic history of cell type and signaling evolution.

250-Word Synopsis

The fetal-maternal interface is one of the most intense loci of cell-cell signaling in the human body. Invasion of cells from the fetal placenta into the uterus, and the corresponding transformation of maternal tissues called decidualization, first evolved in the stem lineage of eutherian mammals[1, 2]. Single-cell studies of the human fetal-maternal interface have provided new insight into the cell type diversity and cell-cell interactions governing this chimeric organ[3–5]. However, the fetal-maternal interface is also one of the most rapidly evolving, and hence most diverse, characters among mammals[6], and an evolutionary analysis is missing. Here, we present and compare single-cell data from the fetal-maternal interface of species bracketing key events in mammal phylogeny: a marsupial (opossum, *Monodelphis domestica*), the afrotherian *Tenrec ecaudatus,* and four Euarchontoglires - guinea pig and mouse (Rodentia) together with recent macaque and human data (primates) [*4*, *5*, *7*]. We infer cell type homologies, identify a gene-expression signature of eutherian invasive trophoblast conserved over 99 million years, and discover a predecidual cell in the tenrec which suggests stepwise evolution of the decidual stromal cell. We reconstruct ancestral cell signaling networks, revealing the integration of fetal cell types into the interface. Finally, we test two long-standing theoretical predictions, the disambiguation hypothesis[8] and escalation hypothesis[9], at transcriptome-wide scale, finding divergence between fetal and maternal signaling repertoires but arms race dynamics restricted to a small subset of ligand-receptor pairs. In so doing, we trace the co-evolutionary history of cell types and their signaling across mammalian viviparity.

Main Text

The origin of novel tissues and organs is essential for the evolution of complex multicellular organisms. However, understanding how evolution at higher levels of organization relates to evolution of cellular composition and signaling is currently limited. One of the most intense loci of cell signaling in the body is the fetal-maternal interface, where cells from the fetal placenta invade and establish communication with the maternal endometrium. Placental invasion, and the corresponding transformation of maternal tissues called decidualization, first evolved in the stem lineage of placental (eutherian) mammals[1, 2]. In

this paper, we trace the fetal-maternal interface to its evolutionary origin, and ask a
fundamental question: how does a complex novelty, composed of highly interdependent
constituent parts, come to be assembled in evolution?
Answering this question demands a comparative perspective. Single-cell
transcriptomic atlases of the human fetal-maternal interface have yielded insight into the cell-
cell interaction networks governing this chimeric organ(*3–5*). However, the placenta is also
one of the most rapidly evolving, and hence most diverse, characters among mammals. Here,
we present and compare single-cell data from the mid-gestation fetal-maternal interface of
five species bracketing key events in mammal phylogeny (**Figure 1a**). These include the gray
short-tailed opossum, *Monodelphis domestica*, a non-deciduate marsupial, and four eutherian
species: the Malagasy common tenrec *Tenrec ecaudatus*, an afrotherian mammal considered
to have a "primitive" form of hemochorial (invasive) placentation (*10*, *11*), and four
Euarchontoglires - the guinea pig *Cavia porcellus*, the mouse *Mus musculus*, and recently
published data from two primates *Macaca fascicularis*(*7*) and *Homo sapiens*(*4*, *5*).
We establish putative cell type homologies, identifying trophoblast populations of
guinea pig and tenrec which share a gene expression signature with primate extravillous
trophoblast, and discover a primitive "predecidual" cell type in the tenrec which suggests a
stepwise evolution of the decidual stromal cell. We reconstruct ancestral cell-cell signaling
networks, revealing the integration of novel cell types into the fetal-maternal interface. Finally,
we test two long-standing theoretical predictions, the disambiguation hypothesis(*8*) and
escalation hypothesis(*9*), at transcriptome-wide scale. In so doing, we trace the co-
evolutionary history of cell types and cell signaling across the origin and diversification of
mammalian placentation.
90

Results

Therian mammals differ in placental and uterine cell type inventories

Single-cell transcriptomic data were collected from the uteroplacental interface of
mammals at informative phylogenetic positions for the evolution of placentation (**Figure 1b**).
Samples were collected after placental development had completed but before the
physiological changes leading to the onset of parturition. Datasets were analyzed following a
standardized approach (**Figure S1**). The resulting curated cell atlas (**Figure 1c**) consists of
404,118 cells and 145,153 nuclei totaling 549,271 libraries, with an average of 10,253 unique
transcripts per library. Cell clusters were identified using Leiden clustering and annotated
using marker gene expression and histology. To reduce over- or under-clustering, we used
non-negative matrix factorization (NMF)(*12*) to identify coregulated gene modules (**Figure
S2**). Cell clusters lacking uniquely distinguishable NMF gene modules or markers were
merged. Detailed cell type descriptions are provided as Supplementary Material.
Tissue organization (**Figure 1b**) and cell type composition (**Figure 1d**) vary across the
mammals in our study. In eutherian mammals, both a vascular interface where maternal
circulation is made available to the fetus and an interface where invading trophoblast contacts
maternal decidua are present. The latter was the focus of analysis due to greater potential for
co-evolved cell-cell signaling. In the opossum, however, fetal-maternal contact and nutrient
exchange occur in the same location, a lamellar matrix of trophoblast which coats maternal
epithelium (**Figure S3a**). Uterine glands, which provide histotrophic nutrition, are expansive
(**Figure 1b**): more than 50% of the cells captured in the opossum were epithelial, of which

85% were glandular (**Figure 1d**). Stromal cells such as fibroblasts are more scarce (8%) in
opossum than in other species (**Figure 1b,d**). In the tenrec, guinea pig, and mouse, the
vascular interface is a trophoblast labyrinth surrounding maternal blood (**Figure 1b**), whereas
the human and macaque vascular interface consists of fetal villi reaching into a maternal
blood sinus. In the tenrec, we identified glands containing mucus and leukocytes, suggesting
continued histotrophic nutrition (**Figure S3b**), whereas in the other eutherians histotrophy has
ceased by the time placentation is complete. Lymphoid cells, primarily NK-like innate cells,
constituted 58% of the total cells recovered in this species (**Figure 1d**), which contrasts with
the opossum, where no uterine NK cells could be identified. The guinea pig, mouse, macaque
and human have similar cell type compositions (**Figure 1d**), but one notable structural
difference: the guinea pig's interface is organized into a folded cytotrophoblast structure
called the subplacenta, which anchors the placenta to the maternal decidua (**Figure 1b**) and
gives rise to invasive trophoblast(*13*).
Assessment of cross-species homology between cell populations was assisted by self-
assembling manifold mapping (SAMap)(*14*), which integrates cells from all species into a
unified manifold (**Figure 1c**) while accounting for complex gene homology. Pairwise SAMap
mapping scores between cell clusters (**Table S1**) formed reciprocally linked groups (**Figure
S4**). Conserved maternal cell types including smooth muscle, pericytes, leukocytes,
endothelial, epithelial, and mesothelial cells showed consistent mapping across species. In
contrast, trophoblast and decidual cell types displayed lower transcriptomic conservation
across species, suggesting rapid cell type and gene expression evolution.
To obtain a global picture of cell type relationships preserving within-species
hierarchical structure, we calculated a neighbor-joining tree of cell type transcriptomes. This
yielded monophyletic or near-monophyletic groupings corresponding to major cell type
families (**Figure 1e**; see **Methods**). The tree contains a hematopoietic cell clade with myeloid
and lymphoid sub-groups, a clade of non-immune mesenchymal cells divided into one group
consisting of fibroblasts, smooth muscle, and pericytes and the other of endothelial cells, and
finally three mixed epithelial/trophoblast clades. These cell type family groupings were largely
maintained when the genes used to generate the phylogeny were restricted to transcription
factors (**Figure S5**), suggesting that the pattern derives from regulatory identity and not solely
phenotypic convergence. Rapid innovation of trophoblast and decidual cell types made one-
to-one homology assessment complicated, so we explored their evolutionary history with
more detailed analyses.

Placental cell types

Trophoblast cells are traditionally classified by topological location (e.g., extravillous),
phenotype (e.g., glycogen trophoblast, syncytiotrophoblast), or ploidy (giant trophoblast), but
whether these divisions correspond to cell type identities conserved across species is
unknown.
Human trophoblast has been well documented by single-cell methods (**Figure 2a**)(*5*):
cytotrophoblast of the placental villi (Hs_VCT) can undergo fusion, becoming
syncytiotrophoblast (Hs_SCT), or invade, becoming extravillous trophoblast (EVT). EVTs can
migrate interstitially (Hs_iEVT) into the decidua, or endovascularly (Hs_eEVT) to replace
maternal spiral artery endothelium. Macaque trophoblast divide into the same three major cell
type groupings of cytotrophoblast of placental villi (Mf_iCTB), fused/syncytial trophoblast
(Mf_fcCTB, Mf_SCT), and extravillous trophoblast (Mf_EVT)(*7*). As not all mammals develop

villi, we refer to other species' migratory trophoblast cells as "extraplacental trophoblast"
(EPT).
In the opossum, two trophoblast populations were identified (**Figure 2a**).
Cytotrophoblast cells (Md_CTB) express transcription factors *WFDC2* and *ZBTB7C*,
desmosome components *DSC2* and *DSG2,* as well as *HAND1*, detected via in situ
hybridization in mononuclear cells of the trophectoderm (**Figure 2b**). These cells also express
the t-SNARE protein *SNAP25* involved in vesicle fusion(*15*), possibly related to fusion with
maternal apocrine secretory bodies (**Figure S3a**). The second population consists of syncytial
trophoblast, which includes giant cells of the trophectoderm, termed syncytial knots(*16*)
(Md_SCT) (**Figure 2b**). Syncytial knots express *GCM1* and *DLX3,* cathepsins *CTSV* and
CTSH, proteases *PRSS22* and *PRSS36*, high levels of the growth factor *VEGFA* and
inflammatory mediators *IL1A, IL6, IL17A* and *PTGS2* as well as *PTGES*.
In the tenrec we identified four trophoblast populations (**Figure 2a**). One cluster
(Te_CTB-CYP11A1) expresses *CYP11A1*, the enzyme catalyzing the first step in steroid
hormone production(*17*), which we localized to the superficial-most trophoblast coating the
endometrium (**Figure 2b**). These cells express heme oxidase *HMOX1*, consistent with reports
in other tenrec species of catabolic breakdown of maternal hemoglobin by
cytotrophoblast(*18*). Steroid hormone-producing cytotrophoblast (Te_CTB-HSD3B1)
expresses the progesterone synthase *HSD3B1* and the cortisol/cortisone converting enzyme
HSD11B1. Non-steroidogenic cytotrophoblast cells are enriched for the somatotropin *IGF2*
(Te_CTB-IGF2) as well as *WFDC2*. Finally, invasive extraplacental trophoblast (Te_EPT)
expresses *ADAMTS18*, *PAPPA2*, *QSOX1*, and cell invasion regulators *SCAI*(*19*) and
ZEB1(*20*)*.* Cells immunoreactive for the proteoglycan GPC1, a marker of invasive
trophoblast(*21*) and angiogenesis(*22*), surround the deepest maternal vasculature (**Figure
2b**), suggesting deep interstitial invasion. Unlike other eutherians, no expression of *prolactin*
or other growth hormone paralogs was present in any *Tenrec* placental cell type.
In the guinea pig, seven trophoblast cell clusters were distinguished. Invasion is
evident as KRT7$^+$ trophoblast cells are present in the decidua and surrounding maternal
vasculature (**Figure S3c**). Extraplacental trophoblast (Cp_iEPT) expresses *ADAM19*, *CNFN,
FLT1, QSOX1,* and *GPC1* (**Figure 2a**). GPC1$^+$ cells surround maternal vasculature and
reside in the myometrium (**Figure 2b**). A population of trophoblast progenitors (Cp_CTB-
CCC) share expression of proliferation regulators *DLX5*(*23*) and *HMGA1* with human cell
column cytotrophoblast (Hs_VCT-CCC); indeed, cell aggregations in the guinea pig
subplacenta have been identified as homologs to the human cell column(*24*). Non-invasive
cytotrophoblasts share the expression of the genes *CTSE, GJB2,* and *TMEM213* (**Figure 2a**).
Syncytiotrophoblast (Cp_SCT) expresses three paralogs of *prolactin*, transcription factors
TFAP2A and TFAP2C, and *LCN2*, a gene linked to invasion in human EVT(*25*).
Syncytiotrophoblast "tongues", identified by elongate cell morphology combined with nuclear
TFAP2C expression (**Figure 2b; Figure S3d**), derive from subplacental cytotrophoblast and
invade deeply into the decidua(*13*). Coarse-meshed syncytiotrophoblast between the
subplacenta and the labyrinth and giant cells between the subplacenta and labyrinth and
invading the decidua, are present histologically (**Figure S3e**) but were not captured
transcriptomically due to their size.
Mouse trophoblast cells were grouped into eight populations: spongiotrophoblast,
syncytiotrophoblast, labyrinth cytotrophoblast, glycogen trophoblast, and three populations of
giant cells (**Figure 2a**). A subset of trophoblast are invasive, united by the expression of
Cdx2, *Pcdh12*(*26*), *Plac1*, and *Tfap2c*. These include interstitially invasive glycogen
trophoblast (Mm_GlyT), enriched in *Arid3a*, and their progenitors (Mm_GlyT_Prog), enriched

for *Ascl2* and *Prdm1*(*27*). Endovascularly invasive spiral artery-remodeling giant cells
(Mm_TGC-SpA) express the prolactins *Prl2a1* and *Prl7b1*. Noninvasive cells of the labyrinth,
canalicular (TGC-C) and sinusoidal (TGC-S) giant cells, express *Cited1*(*28*) and *Ctsj*(*29*)
(**Figure 2a**). Syncytiotrophoblast (Mm_SCT), in close proximity to maternal blood in the
labyrinth, share some of these noninvasive cell markers and are also marked by *Slc16a1*,
Hsd11b2, and the syncytin *Syna*(*30*).

Placental cell type homology inference identifies a eutherian radiation of invasive trophoblasts

211
212

We analyzed fine-grained relationships among trophoblasts by building a network of
SAMap scores and identifying reciprocally linked cell clusters using Leiden community
detection (**Figure 2c**). The Leiden algorithm grouped the 35 total trophoblast cell populations
into five communities which we considered putative homology groups. The resulting
communities divided by degree of invasiveness, with one consisting of invasive cell types and
three of noninvasive cell types (**Figure 2d**).
The grouping of invasive (extraplacental) trophoblast was driven by genes
demonstrated to regulate cell invasion, including proteases *ADAM19*, *MMP15*(*31*)*,* and
PAPPA2(*32*), *ANXA4*(*33*)*, GPC1*(*22*)*,* and *PLAC8*(*34*) (**Figure 2d**; **Table S1**). This group
united extravillous trophoblast from human (Hs_EVT, Hs_eEVT/iEVT) and macaque
(Mf_EVT) with rodent invasive cells (Mm_GlyT, Mm_TGC-SpA, Cp_iEPT) Opossum syncytial
knot cells (Md_SCT) and extraplacental trophoblast from tenrec (Te_EPT) also fell into this
group, suggesting deep conservation of this invasive placental gene signature across
therians*.* Guinea pig SCT, the only syncytial cell type of the 6 species studied to have
invasive properties, was also included in this homology group.
Three communities of noninvasive - primarily cytotrophoblast - cell types were also
identified (**Figure 2d**). Opossum cytotrophoblast (Md_CTB) exhibited linkage to
cytotrophoblast-like cells of all other species (**Figure 2d**): tenrec cytotrophoblast (Te_CTB-
HSD3B1), guinea pig subplacental progenitors (Cp_CTB-CCC), and human villous cell
column progenitors (Hs_VCT-CCC). The similarity of the earlier developmental stages of
human and guinea pig cytotrophoblast to marsupial cytotrophoblast suggests that eutherian-
specific placental cell types may have evolved from a conserved cytotrophoblast cell type.
In contrast to invasiveness, other phenotypic traits of cells, such as fusion and
endoreduplication, did not consistently sort into SAMap communities (**Figure 2d**). Human
giant cells (Hs_TGC), which develop from deeply migratory extravillous cells(*5*), mapped to
extraplacental cells of other species rather than to giant cells of the mouse. Furthermore,
placental cell types which undergo cell fusion, "syncytiotrophoblast", did not show cross-
species conservation except between human and macaque, the two species in our study with
the most recent evolutionary divergence (**Figure 2c**). These formed a homology community
with mouse spongiotrophoblast, driven by expression of large numbers of pregnancy-specific
glycoproteins, although independent evolution of *PSG* and *CEACAM* gene families in rodents
and primates(*35*) suggests that cell type homology based on expression of these genes is
unlikely. While giant opossum knot cells (Md_SCT) are histologically syncytial, their gene
expression differs from other therian trophoblasts by inflammatory and growth factor
production. Likewise, guinea pig SCT clustered with other invasive cell types rather than with
noninvasive mouse or human SCT*.* We conclude that, with the exception of macaque and

human SCT, syncytial trophoblasts and giant trophoblast cell types most likely arose
independently in major mammalian lineages.
Overall, the patterns we observe are consistent with eutherian invasive trophoblasts
having descended from a single cell type with invasive potential in the therian common
ancestor. This cell type was evidently retained in *Tenrec* and *Monodelphis,* which both have
only one cell population belonging to invasive homology groups. This ancestral
extraplacentally migratory cell likely radiated into interstitially migratory and artery-remodeling
subtypes in Boreoeutherian mammals, and acquired distinct morphological phenotypes in
different lineages such as endoreduplication. Conservation of a suite of invasion-enhancing
genes, including matrix proteases and metastasis drivers (**Figure 2d**), suggest that an ancient
potential for cellular invasion was elaborated upon during this process.

Cross-species decidual cell diversity and a predecidual cell type in Tenrec

Decidualization is a coordinated transformation of the endometrium during pregnancy,
including arterial remodeling, recruitment of tissue-specific leukocytes (NK cells), and
differentiation of endometrial stromal fibroblasts into epithelioid decidual stromal cells.
Decidualization is histologically observed only in mammals with hemochorial placentation(*1*).
Molecular mechanisms of decidual development are poorly known outside of humans and
mice, but regulatory interactions essential for decidual cell development, such as protein-
protein interactions between HOXA11 and FOXO1, are shared among eutherian mammals
and lacking in marsupials(*36*, *37*). Using our transcriptomic atlas, we investigated whether the
decidual stromal cell is indeed a unitary cell type novel to eutherians and whether the gene-
regulatory signature of decidual stromal cells is conserved across eutherian mammals.
Recent single-cell studies(*4*, *5*) suggest greater human decidual stromal cell diversity
than the single prolactin-producing cell type canonically recognized in the field(*38*). We used
matrix factorization to identify co-regulated gene sets within the human endometrial stroma:
this identified gene expression modules corresponding to the three decidual cell populations
annotated by Vento-Tormo and colleagues(*4*) (**Figure S2; Table S2**). The gene expression
module of "Type I" stromal cells (Hs_dS1) includes contractility- and myofibroblast-related
genes *ACTA2* and *TAGLN* and myosin light chain kinase *MYLK*. The "Type II" (Hs_dS2) gene
expression module includes *IL15*, *HAND2, MEIS1, FOXO1*, and *LEFTY2.* These genes are
upregulated during the initial wave of human decidualization(*39*) and commonly called
"predecidual"(*40*). The "Type III" (Hs_dS3) gene expression module includes *IGFBP1* and
PRL(*38*)*,* as well as *WNT4* and the decidual proteoglycans *DCN* and *LUM*; we refer to these
as "endocrine decidual cells". Comprehensive sequencing of stromal cells across the human
menstrual cycle found that predecidual cells generated during the secretory phase in the
absence of an embryo, or spontaneous decidual cells, are $IL15^+ PRL^-$ (*41*), similar to dS2 of
pregnancy, not dS3. Thus, human $IL15^+ PRL^+$ endocrine decidual cells are pregnancy-
specific, whereas $IL15^+ PRL^-$ predecidual cells also develop spontaneously during the
proliferative phase of the menstrual cycle. The decidua of the pregnant macaque, an Old-
World Monkey with similar reproductive characteristics including menstruation, shows the
same three stromal cell types: $ACTA2^+$ Mf_dS1, $IL15^+ PRL^-$ Mf_dS2, and $PRL^+ IGFBP1^+$
Mf_dS3.
In the opossum, two stromal cell populations are present. Both express *FOXO1*,
HOXA11, estrogen receptor *ESR1* and progesterone receptor *PGR* (**Figure 3a**). Endometrial
fibroblasts (Md_eS1) are enriched for *SMOC2*, *WNT5A,* and the relaxin receptor *RXFP1*, and
reside in the stroma surrounding endometrial glands(*36*). The other population (Md_TF) are

enriched for *FBLN1*, *CLEC3B*, *MFAP4*, and are present in the myometrium (**Figure 3b**) and
we thus consider them tissue fibroblasts. 29% of Md_eS1 cells express *HAND2* and *IL15*,
markers of human predecidual cells thought to be eutherian-specific(*42*), but these cells did
not group together consistently with sub-clustering. This suggests that *HAND2 and IL15*
expression is a developmentally accessible gene-regulatory state even in stromal cells of the
opossum, but became stabilized into a persistent cell type identity only during eutherian
evolution.
In the tenrec, two major endometrial stromal cell populations are present, united by
expression of *HOXA11* and *HOXA9* (**Figure 3a**). HOXA11 is localized to peri-glandular
endometrial stroma (**Figure 3b**), like the opossum's Md_eS1 cells. RNA velocity(*43*) analysis
inferred a developmental trajectory from fibroblast-like (Te_dS1) to predecidual-like (Te_dS2)
cells (**Figure 3c-d**). Tenrec dS1 are enriched in *CLEC3B***,** *COL14A1, C7*, and *SMOC2,*
whereas predecidual cells express hormone receptors *ERBB4*, *RXFP1, ESR1* and *PGR*,
decidual transcription factor *KLF7*, and predecidual markers *HAND2* and *IL15*. IL15 is a
chemokine for uterine NK cells[4], with the potential to act via binding a receptor
(*IL2RB+IL2RG*) expressed in tenrec uterine NK cells. Treatment of uterine stromal cells
isolated from the pregnant uterus of *T. ecaudatus* with deciduogenic stimuli cAMP and MPA
for 6 days elicited upregulation of these predecidual markers (**Figure 3e**), consistent with the
predecidual cells observed *in vivo* representing this species' full decidual response. The
discovery of a cell population in *Tenrec* resembling *IL15*+ *HAND2*+ human predecidual cells
and lack of a clear endocrine decidual cell suggests that in this member of the basally
divergent mammalian clade Afrotheria, uterine stroma persists in a state similar to the human
predecidua of menstruation, rather than developing endocrine decidual cells.
The guinea pig decidual stroma contains one ECM-remodeling and two
endocrinologically active populations. All three express *HOXA11*, *HAND2, ESR1, PGR,* and
the rodent-specific decidual marker *HOPX*(*45*) **(Figure 3a)**. Endocrine decidual cells include a
cluster enriched in oxytocin *OXT* (Cp_dS3-OXT) and another expressing a *prolactin* gene
(Cp_dS3-PRL) surrounding maternal spiral arteries (**Figure 3b**). The majority (84%) of
stromal cells, however, are marked by expression of *SMOC1*, *SMOC2*, and *WNT5A*. Based
on their size (**Figure 3b**), expression of the cell cycle gene *E2F8* implicated in
endoreduplication(*46*), and previous descriptions of giant decidual cells in necrotic areas of
guinea pig decidua(*13*), we identify these as giant decidual cells, and based on SAMap
affinity and lack of an endocrine profile, classify them as a Type II cell type (Cp_dS2).
The mouse day 15.5 decidua consists of a single endocrine cell type (Mm_dS3)
marked by the prolactin paralog *Prl8a2*, *Pgr*, *Esr1*, and *Hand2*. This cell type also expresses 6
genes recently discovered to have maternally-biased parent-of-origin imprinting(*47*) –
Adamts5, *Erv3*, *Mfap5*, *Tfpi2*, *Tnfrsf11b*, and *Tnfrsf23* (**Figure 3a**): these genes are tumor
suppressors(*48*, *49*) and may represent mechanisms to regulate trophoblast invasion. A
subpopulation of decidual cells show reduced expression of *Pgr, Esr1, Hand2,* and *Hoxa11*
but maintain expression of stress-related genes, including the chaperone *Cryab*, the
inflammatory enzyme *Ctsk*, and the oxidative stress gene *Gpx3*. These we term
progesterone-resistant decidual cells (Mm_dS3-Pgr⁻). They have a matching gene expression
profile with a *Pgr⁻* oxidative stress-enriched endocrine decidual cell population which a
previous study identified as postmature decidua(*50*), and share similarities with so-called
senescent *PGR⁻* endocrine decidual cells of the human which exist in balance with normal
endocrine decidual cells(*51*). Cells corresponding to the predecidual cells of the tenrec and
human were absent in the mouse.

Stepwise evolution of the uterine decidua in Eutheria

While our cell type phylogeny (**Figure 1e**) identified high-level family groupings, it could not resolve one-to-one decidual cell type homologies. We investigated fine-scale decidual cell relationships using SAMap community detection and comparison of gene co-regulatory modules across species.

Two stromal cell type communities in the SAMap network were highly distinct from other mesenchymal cell types, smooth muscle cells and placental fibroblasts (**Figure 3f**). One community consists predominantly of endocrine decidual cells and the other of contractile fibroblasts and predecidual stromal cells. The endocrine or "Type III" community united the two endocrine decidual cell populations of the guinea pig (OXT^+ and PRL^+), mouse Pgr^+ and Pgr^- decidual cells, and human and macaque PRL^+ $IGFBP1^+$ cells. The predecidual and non-decidual stromal cell community consisted of the opossum's $SMOC2^+$ eS1 and $FBLN1^+$ tissue fibroblasts, tenrec dS1 and predecidual cells, guinea pig dS2, mouse tissue fibroblasts, endometrial stromal fibroblasts from macaque and human, and macaque predecidual cells. Human predecidual Hs_dS2 had strong mapping affinities to both the predecidual cells of other mammals and to endocrine decidua and were formally assigned to the same group as endocrine decidual cells. This result was robust across all iterations of community detection attempted, although we suspect that this more reflects the fuzziness of drawing hard boundaries in cells undergoing a continuous developmental process than it does an evolutionary difference between the predecidua of humans and other mammals.

We next looked for conservation of co-regulated gene modules in decidual cells of our 6-species comparison (**Figure S2**). A hierarchically-clustered correlation matrix of gene expression programs identified by NMF analysis grouped them into two families (**Figure 3g; Table S2**): Predecidual-like gene-regulatory modules from all species tended to weigh highly *MEIS1, WT1, ESR1, PGR,* and *EGFR*, and were active in human, macaque, tenrec, and guinea pig. Endocrine gene-regulatory modules, which had high weightings of *PRL, LUM, DCN, VIM, PPIB*, and *DUSP1*, were active in endocrine decidua of all eutherians (mouse, guinea pig, human and macaque), but no counterpart was identified in the opossum. Predecidual and endocrine gene expression modules showed low or negative correlation with one another (**Figure 3g**), meaning that genes expressed in the predecidual cells tended not to be expressed in mature decidual cells (**Figure S2**). We interpret this as evidence that predecidual cells and endocrine decidual cells are robustly distinct alternative gene-regulatory states.

Integrating these lines of evidence leads to the following model for decidual cell evolution (**Figure 3h**). Un-decidualized, contractile endometrial fibroblasts, or "Type I" cells, have homologs across therian mammals. These cells predominate in the non-pregnant uterus. Eutherians display decidual stromal cells falling into two states – predecidual or "Type II" decidual cells, which secrete immunomodulatory peptides, and "Type III" decidual cells, which secrete endocrine and growth-regulatory signals. Type II decidual cells likely represent a novel eutherian stromal cell type, and are evolutionarily modified from mesenchymal cells of the ancestral therian endometrium by upregulation of *HAND2* and *IL15*(52). In the two rodents sampled, no stromal cell type produces *IL15*; instead, decidual *IL15* is solely produced by macrophages. However, the deep phylogenetic age of the tenrec-human divergence suggests that the $IL15^+$ predecidual cell type was lost in rodents, or is restricted to early stages of pregnancy. These findings suggest that decidual stromal cell diversity expanded within crown Placentalia in two steps – establishment of the decidual stromal cell type in its initial

390 predecidual *IL15+ PRL-* state, followed by later evolution of the endocrine (*PRL+*) decidual cells
391 observed in rodents and primates.

Co-evolutionary dynamics of cell communication

393   We next asked to what degree the cell-cell communication between fetal and maternal
394 cell types shows evidence of co-evolution. We inferred signaling between cell types from
395 complementary secreted ligand-receptor gene expression and traced the evolutionary history
396 of these interactions within Theria.
397

Invasive trophoblast cells are integrated into the endometrial signaling network

The evolution of placentation is predicted to result in increased functional integration of
fetal and maternal cells(*53*). As an approximation of functional integration, we compared the
number of non-autocrine ("allocrine") signaling interactions each cell type engages in (see
Methods). Cells producing more allocrine ligands tend to also receive more allocrine signals
themselves (Pearson's *p* < 0.05 in all species except *Monodelphis*), and degrees of allocrine
ligand production varied across major cell type families (one-way ANOVA *p* < 0.05 in all
species except *Cavia*; **Figure 4a**). Stromal cells, particularly decidual stromal cells, display
the greatest degrees of allocrine signaling. Invasive trophoblast (EVT/EPT, Mm_GlyT in
mouse and Md_SCT in opossum) also show high degrees of allocrine signaling (**Figure 4a**).
Macrophages and endothelial cells rank intermediately, whereas eutherian
syncytiotrophoblast and immune effector cells (BC, TC, NKC, PMN) show the fewest allocrine
interactions (**Figure 4a**). *Monodelphis GCM1+* trophoblast and the human Type III decidual
cell have the highest Kleinberg(*54*) hub scores in their respective signaling networks, whereas
lymphoid cells from all species rank among the lowest (**Figure S6a**). We interpret these
findings to mean that fetal and maternal cell types that make direct contact have evolved to
function as signaling hubs, highly integrated into the signaling network of the fetal-maternal
interface, whereas peripheral immune cells without a stable niche do not evolve integration to
the tissue microenvironment(*55*).
418

Reconstruction of common ancestral fetal-maternal signaling

We inferred the fetal-maternal signaling of common ancestors in our tree using
maximum-parsimony ancestral state reconstruction on ligand and receptor expression of
pooled fetal and maternal cell types (**Table S4**). We identify a total of 83 signals which the
placenta of the first viviparous mammal likely sent to its mother, including relaxin *RLN2*,
platelet-derived growth factors *PDGFA* and *PDGFC*, prostaglandin E2 via *PTGES3*, and
TGFB1 and *TGFB2* (**Figure 4b**). We traced gains and losses of signaling interactions
throughout mammal evolution by comparing ancestral states. At the Placentalia node, an
additional 44 signals were inferred to have been gained, including vascular growth factors
VEGFB and *VEGFC* and the insulin-like growth factor *IGF2*. To gain more precise insight, we
reconstructed full ancestral signaling networks between major cell type classes (**Table S4**).
The reconstructed network of the Placental mammal common ancestor revealed that early
eutherian trophoblast had already acquired a signaling potential towards maternal stromal,
vascular, epithelial, and immune cells (**Figure 4c**).
Human, macaque, mouse, and guinea pig belong to the clade Euarchontoglires. Their
stem lineage evolved 4 fetal-maternal signals absent in the basally branching *Tenrec,*

including *CCL2* and *CCL4*, and also lost fetal expression of 11 ligands, including cytokines
IL1A and *TNF* and neutrophil chemoattractant *CXCL2*, suggesting that inflammatory signaling
by the placenta was suppressed after the divergence of tenrec and opossum. The two
rodents in our sample, *Cavia porcellus* and *Mus musculus,* share an additional 10 ligands,
including *FGF7*, *IL34*, *OSM* (related to *LIF*), and *prolactin*. Many fetal-maternal ligand-
receptor interactions are species-specific. *Mus musculus* saw an expansion of novel
CEACAM genes, and the midgestation guinea pig placenta is unique in *OXT* production,
produced only at parturition in other species. The human lineage saw the acquisition of novel
growth hormone paralogs, and the *Monodelphis domestica* placenta is inferred to have
gained expression of additional pro-inflammatory cytokines.
We calculated rates of evolutionary change in fetal-to-maternal signaling using a time-
calibrated mammalian tree(*56*) (**Figure 4b**). The greatest rates of evolutionary change were in
the stem lineage leading to Euarchontoglires, with as many as 1.25 changes/Myr, many of
these being losses of inflammatory signaling (e.g. *IL1A, PTGIS, CXCL2, TNF*) by fetal cell
types. High rates of change were also observed in both primate lineages. In contrast, the
lineage leading to *Monodelphis domestica* and the lineage leading to *Tenrec ecaudatus* show
slower turnover. These findings suggest that the diversification of Euarchontoglires entailed
an accelerated evolution of fetal-maternal communication compared to basally branching
placentals and marsupial lineages.
454

Fetal and maternal signaling show disambiguation

A model of placental hormone evolution based on evolutionary signaling theory (*8*)
predicts that fetal-maternal signaling systems will evolve towards maternal-only or fetal-only
expression to reduce fetal manipulation of maternal physiology. Selection against fetal and
maternal co-expression of ligands would manifest as a pattern of "disambiguation". To
quantify disambiguation, we compared the observed number of fetal-only and maternal-only
expressed ligands from different signaling families (e.g. WNT, NOTCH) against a null
statistical model based on random distribution of ligands to cell types regardless of origin
(**Figure 5a-b;** see **Methods**). In all species, fetal and maternal co-expression of ligands was
lower than expected by our null model (**Figure 5b**). The most consistently disambiguated
ligand families across species included WNTs, steroids, FGFs, and various families of
immunomodulatory factors (**Figure 5b; Table S3**).
Gene duplication can increase disambiguation by the creation of novel placenta- or
decidua-expressed paralogs. In the murine lineage, the *prolactin* gene family includes 26
members and a strongly disambiguated expression pattern with 10 fetal-specific paralogs
(**Figure 5c; Table S3**). In human and macaque, duplication of the *GH1* locus has led to the
origin of various placenta-specific paralogs, whereas the somatoropin *PRL* is maternal-only
(**Figure 5c**). Cell type-specific growth factors involved in core tissue homeostasis, such as
VEGF, *PDGF,* and *CSF1*, showed overwhelmingly bilateral expression (**Table S3**).
Extreme production of the angiogenic factor *VEGFA* by the opossum trophoblast can
be interpreted as a nutrient-soliciting signal. The disambiguation model(*8*) predicts that in an
evolutionary stable state, the mother should silence her own copy or utilize a different VEGF
paralog. However, all 3 VEGF paralogs in the opossum were co-expressed. This may be
explained by developmental constraints countervailing the selection for disambiguation, due
to the importance of VEGF in tissue homeostasis. Alternatively, disambiguation may be
achieved by maternal utilization of a splice variant of VEGFA with differential signaling

properties, as documented with the *VEGF[111]* isoform in the marsupial *Sminthopsis*
crassicaudata, the rat, and viviparous skinks(*57*).
483

484 *Escalation of fetal-maternal signaling*

485
Parent-offspring conflict over nutrient allocation during pregnancy is predicted to lead
to an asymmetric evolutionary arms race(*58*) in which gains to fetal signaling strength are
matched by reduced maternal responsiveness(*9*). We tested for correlated fetal ligand-
maternal receptor co-evolution at the transcriptomic level by comparing reconstructed
evolutionary changes in ligand and receptor expression across internal branches of the tree.
We tested for a quantitative escalation pattern without complete suppression by modeling
changes in expression levels as continuous traits and testing for phylogenetically-
independent(*59*) associations between ligands and cognate receptor expression (**Figure 5d;**
Table S5). Inferred evolutionary changes along internal branches of the tree did not show a
global prevalence towards escalation regardless of whether all secreted signaling peptides
are considered or just growth factors (gene ontology group GO:0008083), (**Figure S6b-c;**
Table S5). Across the tree, negative ligand-receptor correlations were observed for the
inhibin-activin pair INHA–ACVR1B+ACVR2A ($p = 1.1 \times 10^{-3}$) and the well-studied gene pair
IGF2–IGF1R(*60*) ($p = 2.7 \times 10^{-2}$) (**Figure 5d**). Other pairs such as VEGFA–FLT1 showed
positive correlation ($p = 5.6 \times 10^{-3}$), i.e. gains to both fetal ligand and maternal receptor
expression along the same branches, possibly reflecting a cooperative trend towards greater
placental vascular development in multiple eutherian lineages rather than a manipulation-
desensitization dynamic.
In the branch leading from the human-macaque ancestor to humans, for which the
ancestral state confidence was high due to the sampling of two rodents and two outgroups,
only the growth factors *PGF* and *TGFB3* showed asymmetric escalation (**Figure 5e**), whereas
PRL, *KITLG*, and *IGF2* were inferred to be undergoing de-escalation relative to the human-
macaque ancestor. When modeling ligand and receptor expression as binary on/off traits
using expression thresholding, lineages in which more fetal ligands were gained saw more
gains in maternal receptors, rather than losses (**Figure 4b**). These findings suggest that only
a select subset of placental-maternal signaling is subject to antagonistic gene expression
evolution. That said, our analysis cannot exclude antagonistic dynamics at the post-
transcriptional level, as in the case of decoy receptors(*60*) or the chemical properties of signal
peptides such as their half-life(*61*).
515
516

Discussion

The eutherian fetal-maternal interface is the product of three major evolutionary events
– the origin of endometrial decidualization, the evolution of a distinct placental organ (in
contrast to simple apposition of fetal and maternal tissue), and the origin of trophoblast
invasion(*2*). The latter is extreme in species with deep interstitial invasion such as humans
and guinea pigs. In this paper, we have compared the cell type composition and cell-cell
signaling networks of species spanning these transitions revealing a complex history of cell
type and cell signaling evolution.

Cross-species mapping suggests that eutherian mammals underwent an expansion in trophoblast cell type diversity. We infer that the therian ancestor, like the extant opossum, had two trophoblast cell types, one homologous to eutherian cytotrophoblast and the other specialized in signaling to the mother with weak invasive potential. Extraplacental trophoblast of tenrec, guinea pig, macaque and human and spiral artery-remodeling trophoblast in the mouse shared a common gene signature. The grouping of phylogenetically distant trophoblast transcriptomes by invasive potential suggests that the invasive/noninvasive phenotypic divide may reflect an ancient divide in trophoblast cell type identity, rather than convergent evolution.

Endometrial stromal cells fall into homology groups of different phylogenetic ages, suggesting a stepwise accrual of decidual cell diversity in eutherians. The discovery of a primitive $IL15^+$ predecidual cell in *Tenrec*, with gene expression similarity to human Type II decidual cells as well as endometrial stromal cells of other species, suggests that the predecidual cell is conserved over the 99 million year *Tenrec-Homo* divergence. Endocrine (PRL^+) decidual cells likely arose later in eutherian evolution, restricted to humans and rodents in our sampling (Euarchontoglires) and lacking in *Tenrec*; investigation of decidua in more mammals from the basally-branching Afrotheria and Xenarthra clades will allow this evolution to be more thoroughly untangled.

Our findings are consistent with a model of cell type evolution playing out at two distinct levels: broad transcriptomic "cell type identities" and functional gene expression modules or "apomeres"(*62*), a division established in neuronal(*63*) and photoreceptor(*64*) evolution. In addition to novel cell types of decidual stroma and extraplacental trophoblast, fetal and maternal cells both show evidence of acquisition of functional apomeres. On the fetal side, syncytiotrophoblasts are transcriptomically dissimilar across species, and perform diverse functions - inflammatory and growth factor signaling in the opossum, invasion in the guinea pig, vascular interface in the mouse, and gonadotropin production in the human. Molecular genetic evidence suggests that trophoblast cell fusion is associated with five independent insertions of retroviral envelope genes in the opossum, tenrec, guinea pig, muroid, and primate lineages(*65–67*), a pattern congruent with our inferred lack of homology between syncytiotrophoblast beyond the human-macaque pair. Likewise, prolactin expression in endocrine decidual cells is associated with independent transposable element insertions upstream of the *prolactin* promoters in rodents and primates(*68*). These findings suggest that both trophoblast and decidual evolution have involved parallel evolution of new phenotypes, like cell fusion and hormone production, by recruitment of retroviral and transposable elements into gene regulatory networks of these cell types. Identifying which of the cell populations we identify here correspond to developmentally robust cell type identities, and the factors determining their differentiation, will require further targeted investigation.

Compelling theoretical models have long been advanced for the co-evolutionary signaling dynamics between the fetus and mother(*9*). Our test of the disambiguation hypothesis(*8*) revealed consistent and strong overrepresentation of both fetal-specific and maternal-specific ligand repertoires. This pattern was driven in part by recent gene duplication into fetal-specific paralogs(*35*). A logical next step would be to investigate genetic mechanisms, such as capture of ligand genes by fetal-specific enhancers(*69*). The escalation hypothesis predicts that increased fetal ligand production should be complemented by reduction in corresponding receptor expression by the mother. Only a select few ligand-receptor pairs in our sample consistently followed this dynamic, suggesting it does not hold for placental-uterine signaling universally; however, some of the outlier ligand-recetor pairs we identified are plausibly involved in escalation such as *IGF2*. Finally, we used ancestral

state reconstruction to trace the co-evolutionary history of ligand-receptor signaling, reconstructed the cell signaling interactions of the Placental common ancestor, and found that Boreoeutherians underwent a more rapid divergence in fetal-maternal signaling than the opossum and tenrec lineages. This implies that fetal-maternal co-evolution was not solely concentrated at the origin of eutherian placentation; rather, the pace of gene-regulatory innovation in mammals has even accelerated in eutherian evolution, including substantial changes in recent primate and rodent evolution.

We acknowledge limitations to inferring ligand-receptor signaling from RNA expression. The potential for false negatives due to sequencing drop-out, incomplete genome annotation in non-model species, and protein-protein interaction divergence likely inflate the inferred number of species-specific signaling innovations. Differences in gestation length and placental physiology between our six species may limit the comparability of the chosen gestational stages for trophoblast and decidua, which are developmentally dynamic. Nevertheless, the outlines of a multistage history of cell type and cell phenotypic evolution can clearly be seen.

The fetal-maternal interface has attracted attention in recent years as a model system for single-cell genomics and ligand-mediated cell signaling at scale[3-5]. The atlas we present here demonstrates that the field has reached the point where long-standing evolutionary questions about parent-offspring co-evolution can be addressed at a resolution heretofore impossible. The molecular genetic mechanisms behind signaling innovation, integration, and disambiguation will require future investigation. As high-throughput genomics of non-traditional model species becomes more technologically accessible, expanded taxon sampling will allow the kinds of phylogenetic reconstructions we introduce here to expand in precision and scope.

The evolutionary approach we introduce for cell-cell communication holds potential for broader application. Comparative analysis may help uncover the co-evolution of cell types within other tissues, and in other inter-organismal symbioses, an avenue opened by recent advances in single-cell transcriptomics of multicellular parasites(*70*) and commensals(*71*).

References and Notes

1. A. Mess, A. M. Carter, Evolutionary transformations of fetal membrane characters in Eutheria with special reference to Afrotheria. *J. Exp. Zoolog. B Mol. Dev. Evol.* **306B**, 140–163 (2006).
2. D. E. Wildman, C. Chen, O. Erez, L. I. Grossman, M. Goodman, R. Romero, Evolution of the mammalian placenta revealed by phylogenetic analysis. *Proc. Natl. Acad. Sci.* **103**, 3203–3208 (2006).
3. M. Pavličev, G. P. Wagner, A. R. Chavan, K. Owens, J. Maziarz, C. Dunn-Fletcher, S. G. Kallapur, L. Muglia, H. Jones, Single-cell transcriptomics of the human placenta: inferring the cell communication network of the maternal-fetal interface. *Genome Res.* **27**, 349–361 (2017).
4. R. Vento-Tormo, M. Efremova, R. A. Botting, M. Y. Turco, M. Vento-Tormo, K. B. Meyer, J.-E. Park, E. Stephenson, K. Polański, A. Goncalves, L. Gardner, S. Holmqvist, J. Henriksson, A. Zou, A. M. Sharkey, B. Millar, B. Innes, L. Wood, A. Wilbrey-Clark, R. P. Payne, M. A. Ivarsson, S. Lisgo, A. Filby, D. H. Rowitch, J. N. Bulmer, G. J. Wright, M. J. T. Stubbington, M. Haniffa, A. Moffett, S. A. Teichmann, Single-cell reconstruction of the early maternal–fetal interface in humans. *Nature* **563**, 347–353 (2018).
5. A. Arutyunyan, K. Roberts, K. Troulé, F. C. K. Wong, M. A. Sheridan, I. Kats, L. Garcia-Alonso, B. Velten, R. Hoo, E. R. Ruiz-Morales, C. Sancho-Serra, J. Shilts, L.-F. Handfield, L. Marconato,

E. Tuck, L. Gardner, C. I. Mazzeo, Q. Li, I. Kelava, G. J. Wright, E. Prigmore, S. A. Teichmann,
O. A. Bayraktar, A. Moffett, O. Stegle, M. Y. Turco, R. Vento-Tormo, Spatial multiomics map of
trophoblast development in early pregnancy. *Nature* **616**, 143–151 (2023).

6.   M. Garratt, J.-M. Gaillard, R. C. Brooks, J.-F. Lemaître, Diversification of the eutherian placenta
is associated with changes in the pace of life. *Proc. Natl. Acad. Sci.* **110**, 7760–7765 (2013).

7.   X. Jiang, J. Zhai, Z. Xiao, X. Wu, D. Zhang, H. Wan, Y. Xu, L. Qi, M. Wang, D. Yu, Y. Liu, H.
Wu, R. Sun, S. Xia, K. Yu, J. Guo, H. Wang, Identifying a dynamic transcriptomic landscape of
the cynomolgus macaque placenta during pregnancy at single-cell resolution. *Dev. Cell* **58**, 806-
821.e7 (2023).

8.   D. Haig, Placental hormones, genomic imprinting, and maternal—fetal communication. *J. Evol.*
Biol. **9**, 357–380 (1996).

9.   D. Haig, Genetic conflicts in human pregnancy. *Q. Rev. Biol.* **68**, 495–532 (1993).

10. O. Grosser, Die Placenta von Centetes und ihre Lehren betreffs der Stoffaufnahme in den
Placenten. *Z. Für Anat. Entwicklungsgeschichte* **88**, 509–521 (1928).

11. H. Strahl, Beiräge zur vergleichenden Anatomie der Placenta. *Abhandl Senckenberg Naturf*
Gesell **27**, 263–319 (1905).

12. D. Kotliar, A. Veres, M. A. Nagy, S. Tabrizi, E. Hodis, D. A. Melton, P. C. Sabeti, Identifying
gene expression programs of cell-type identity and cellular activity with single-cell RNA-Seq.
eLife **8**, e43803 (2019).

13. J. Davies, E. W. Dempsey, E. C. Amoroso, The subplacenta of the guinea-pig: development,
histology and histochemistry. *J. Anat.* **95**, 457–473 (1961).

14. A. J. Tarashansky, J. M. Musser, M. Khariton, P. Li, D. Arendt, S. R. Quake, B. Wang, Mapping
single-cell atlases throughout Metazoa unravels cell type evolution. *eLife* **10**, e66747 (2021).

15. Y. Zhao, Q. Fang, A. D. Herbst, K. N. Berberian, W. Almers, M. Lindau, Rapid structural change
in synaptosomal-associated protein 25 (SNAP25) precedes the fusion of single vesicles with the
plasma membrane in live chromaffin cells. *Proc. Natl. Acad. Sci.* **110**, 14249–14254 (2013).

16. Zeller, Freyer, Early ontogeny and placentation of the grey short-tailed opossum, Monodelphis
domestica (Didelphidae: Marsupialia): contribution to the reconstruction of the marsupial
morphotype. *J. Zool. Syst. Evol. Res.* **39**, 137–158 (2001).

17. Y.-N. Zhu, F. Pan, X.-W. Gan, Y. Liu, W.-S. Wang, K. Sun, The Role of DNMT1 and C/EBPα in
the Regulation of CYP11A1 Expression During Syncytialization of Human Placental
Trophoblasts. *Endocrinology* **165**, bqad195 (2024).

18. A. M. Carter, Evolution of Placental Function in Mammals: The Molecular Basis of Gas and
Nutrient Transfer, Hormone Secretion, and Immune Responses. *Physiol. Rev.* **92**, 1543–1576
(2012).

19. D. T. Brandt, C. Baarlink, T. M. Kitzing, E. Kremmer, J. Ivaska, P. Nollau, R. Grosse, SCAI acts
as a suppressor of cancer cell invasion through the transcriptional control of β1-integrin. *Nat. Cell*
Biol. **11**, 557–568 (2009).

20. X. Yang, T. Meng, MicroRNA-431 affects trophoblast migration and invasion by targeting ZEB1
in preeclampsia. *Gene* **683**, 225–232 (2019).

21. R. L. Scott, H. T. H. Vu, A. Jain, K. Iqbal, G. Tuteja, M. J. Soares, Conservation at the uterine–
placental interface. *Proc. Natl. Acad. Sci.* **119**, e2210633119 (2022).

22. T. Aikawa, C. A. Whipple, M. E. Lopez, J. Gunn, A. Young, A. D. Lander, M. Korc, Glypican-1
modulates the angiogenic and metastatic potential of human and mouse cancer cells. *J. Clin.*
Invest. **118**, 89–99 (2008).

23. J. Zadora, M. Singh, F. Herse, L. Przybyl, N. Haase, M. Golic, H. W. Yung, B. Huppertz, J. E.
Cartwright, G. Whitley, G. M. Johnsen, G. Levi, A. Isbruch, H. Schulz, F. C. Luft, D. N. Müller,
A. C. Staff, L. D. Hurst, R. Dechend, Z. Izsvák, Disturbed Placental Imprinting in Preeclampsia

Leads to Altered Expression of DLX5, a Human-Specific Early Trophoblast Marker. *Circulation*
136, 1824–1839 (2017).

24. A. Mess, N. Zaki, M. Kadyrov, H. Korr, P. Kaufmann, Caviomorph Placentation as a Model for
Trophoblast Invasion. *Placenta* **28**, 1234–1238 (2007).

25. H. Kobara, T. Miyamoto, A. Suzuki, R. Asaka, Y. Yamada, K. Ishikawa, N. Kikuchi, S. Ohira, T.
Shiozawa, Lipocalin2 enhances the matrix metalloproteinase-9 activity and invasion of
extravillous trophoblasts under hypoxia. *Placenta* **34**, 1036–1043 (2013).

26. B. Marsh, R. Blelloch, Single nuclei RNA-seq of mouse placental labyrinth development. *eLife* **9**,
e60266 (2020).

27. S. J. Tunster, E. D. Watson, A. L. Fowden, G. J. Burton, Placental glycogen stores and fetal
growth: insights from genetic mouse models. *Reproduction* **159**, R213–R235 (2020).

28. T. A. Rodriguez, D. B. Sparrow, A. N. Scott, S. L. Withington, J. I. Preis, J. Michalicek, M.
Clements, T. E. Tsang, T. Shioda, R. S. P. Beddington, S. L. Dunwoodie, Cited1 Is Required in
Trophoblasts for Placental Development and for Embryo Growth and Survival. *Mol. Cell. Biol.*
24, 228–244 (2004).

29. A. Nakajima, K. Kataoka, Y. Takata, N. Huh, Cathepsin-6, a novel cysteine proteinase showing
homology with and co-localized expression with cathepsin J/P in the labyrinthine layer of mouse
placenta. *Biochem. J.* **349**, 689–692 (2000).

30. D. G. Simmons, D. R. C. Natale, V. Begay, M. Hughes, A. Leutz, J. C. Cross, Early patterning of
the chorion leads to the trilaminar trophoblast cell structure in the placental labyrinth.
Development **135**, 2083–2091 (2008).

31. A. Majali-Martinez, D. Hoch, C. Tam-Amersdorfer, J. Pollheimer, A. Glasner, N. Ghaffari-
Tabrizi-Wizsy, A. G. Beristain, U. Hiden, M. Dieber-Rotheneder, G. Desoye, Matrix
metalloproteinase 15 plays a pivotal role in human first trimester cytotrophoblast invasion and is
not altered by maternal obesity. *FASEB J.* **34**, 10720–10730 (2020).

32. X. Chen, K. Chen, Y. Feng, C. Ren, W. Li, J. Xiao, L. Fan, R. Beejadhursing, L. Xi, S. Chen, The
potential role of pregnancy-associated plasma protein-A2 in angiogenesis and development of
preeclampsia. *Hypertens. Res.* **42**, 970–980 (2019).

33. Y. Xu, L. Sui, B. Qiu, X. Yin, J. Liu, X. Zhang, ANXA4 promotes trophoblast invasion via the
PI3K/Akt/eNOS pathway in preeclampsia. *Am. J. Physiol.-Cell Physiol.* **316**, C481–C491 (2019).

34. W.-L. Chang, Y.-W. Liu, Y.-L. Dang, X.-X. Jiang, H. Xu, X. Huang, Y.-L. Wang, H. Wang, C.
Zhu, L.-Q. Xue, H.-Y. Lin, W. Meng, H. Wang, PLAC8, a new marker for human interstitial
extravillous trophoblast cells, promotes their invasion and migration. *Development* **145**,
dev148932 (2018).

35. R. Kammerer, W. Zimmermann, Two waves of evolution in the rodent pregnancy-specific
glycoprotein (Psg) gene family lead to structurally diverse PSGs. *BMC Genomics* **24**, 468 (2023).

36. K. Kin, J. Maziarz, G. P. Wagner, Immunohistological Study of the Endometrial Stromal
Fibroblasts in the Opossum, Monodelphis domestica: Evidence for Homology with Eutherian
Stromal Fibroblasts1. *Biol. Reprod.* **90** (2014).

37. K. J. Brayer, V. J. Lynch, G. P. Wagner, Evolution of a derived protein–protein interaction
between HoxA11 and Foxo1a in mammals caused by changes in intramolecular regulation. *Proc.*
Natl. Acad. Sci. **108**, E414–E420 (2011).

38. B. Gellersen, J. J. Brosens, Cyclic Decidualization of the Human Endometrium in Reproductive
Health and Failure. *Endocr. Rev.* **35**, 851–905 (2014).

39. K. T. Rytkönen, E. M. Erkenbrack, M. Poutanen, L. L. Elo, M. Pavlicev, G. P. Wagner,
Decidualization of Human Endometrial Stromal Fibroblasts is a Multiphasic Process Involving
Distinct Transcriptional Programs. *Reprod. Sci.* **26**, 323–336 (2019).

40. T. M. Rawlings, K. Makwana, D. M. Taylor, M. A. Molè, K. J. Fishwick, M. Tryfonos, J.

Odendaal, A. Hawkes, M. Zernicka-Goetz, G. M. Hartshorne, J. J. Brosens, E. S. Lucas,
Modelling the impact of decidual senescence on embryo implantation in human endometrial
assembloids. *eLife* **10**, e69603 (2021).
41. M. Marečková, L. Garcia-Alonso, M. Moullet, V. Lorenzi, R. Petryszak, C. Sancho-Serra, A.
Oszlanczi, C. I. Mazzeo, S. Hoffman, M. Krassowski, K. Garbutt, I. Kelava, K. Gaitskell, S.
Yancheva, E. V. Woon, V. Male, I. Granne, K. Hellner, K. T. Mahbubani, K. Saeb-Parsy, M.
Lotfollahi, E. Prigmore, J. Southcombe, R. A. Dragovic, C. M. Becker, K. T. Zondervan, R.
Vento-Tormo, An integrated single-cell reference atlas of the human endometrium. bioRxiv
[Preprint] (2023). https://doi.org/10.1101/2023.11.03.564728.
42. K. Mika, C. M. Whittington, B. M. McAllan, V. J. Lynch, Gene expression phylogenies and
ancestral transcriptome reconstruction resolves major transitions in the origins of pregnancy. *eLife*
11, e74297 (2022).
43. V. Bergen, M. Lange, S. Peidli, F. A. Wolf, F. J. Theis, Generalizing RNA velocity to transient
cell states through dynamical modeling. *Nat. Biotechnol.* **38**, 1408–1414 (2020).
44. A. Erlebacher, Immunology of the maternal-fetal interface. *Annu. Rev. Immunol.* **31**, 387–411
(2013).
45. R. Li, T. Wang, X. Xu, O. M. Emery, M. Yi, S.-P. Wu, F. J. DeMayo, Spatial transcriptomic
profiles of mouse uterine microenvironments at pregnancy day 7.5. *Biol. Reprod.* **107**, 529–545
(2022).
46. Q.-R. Qi, X.-Y. Zhao, R.-J. Zuo, T.-S. Wang, X.-W. Gu, J.-L. Liu, Z.-M. Yang, Involvement of
atypical transcription factor E2F8 in the polyploidization during mouse and human
decidualization. *Cell Cycle* **14**, 1842–1858 (2015).
47. F. Rodriguez-Caro, E. C. Moore, J. M. Good, Evolution of parent-of-origin effects on placental
gene expression in house mice. bioRxiv [Preprint] (2023).
https://doi.org/10.1101/2023.08.24.554674.
48. S. Kumar, S. Sharghi-Namini, N. Rao, R. Ge, ADAMTS5 Functions as an Anti-Angiogenic and
Anti-Tumorigenic Protein Independent of Its Proteoglycanase Activity. *Am. J. Pathol.* **181**, 1056–
1068 (2012).
49. X. Xiao, X. Tao, Y. Wang, L. Zhu, Y. Ye, H. Liu, Q. Zhou, X. Li, Y. Xiong, Hypomethylation of
tissue factor pathway inhibitor 2 in human placenta of preeclampsia. *Thromb. Res.* **152**, 7–13
(2017).
50. M. Yang, J. Ong, F. Meng, F. Zhang, H. Shen, K. Kitt, T. Liu, W. Tao, P. Du, Spatiotemporal
insight into early pregnancy governed by immune-featured stromal cells. *Cell* **186**, 4271-4288.e24
(2023).
51. E. S. Lucas, P. Vrljicak, J. Muter, M. M. Diniz-da-Costa, P. J. Brighton, C.-S. Kong, J. Lipecki,
K. J. Fishwick, J. Odendaal, L. J. Ewington, S. Quenby, S. Ott, J. J. Brosens, Recurrent pregnancy
loss is associated with a pro-senescent decidual response during the peri-implantation window.
Commun. Biol. **3**, 1–14 (2020).
52. K. Mika, M. Marinić, M. Singh, J. Muter, J. J. Brosens, V. J. Lynch, Evolutionary transcriptomics
implicates new genes and pathways in human pregnancy and adverse pregnancy outcomes. *eLife*
10, e69584 (2021).
53. O. W. Griffith, G. P. Wagner, The placenta as a model for understanding the origin and evolution
of vertebrate organs. *Nat. Ecol. Evol.* **1**, 1–10 (2017).
54. J. M. Kleinberg, Authoritative sources in a hyperlinked environment. *J. ACM* **46**, 604–632 (1999).
55. M. Pavlicev, G. Wagner, Reading the Palimpsest of Cell Interactions: What Questions May We
Ask of the Data? *iScience,* doi: 10.20944/preprints202310.1685.v1 (2024).
56. S. Kumar, M. Suleski, J. M. Craig, A. E. Kasprowicz, M. Sanderford, M. Li, G. Stecher, S. B.
Hedges, TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol. Biol. Evol.* **39**,

msac174 (2022).

57.   C. M. Whittington, K. Danastas, G. E. Grau, C. R. Murphy, M. B. Thompson, Expression of
VEGF111and other VEGF-A variants in the rat uterus is correlated with stage of pregnancy. *J.*
Comp. Physiol. B **187**, 353–360 (2017).

58.   R. Dawkins, J. R. Krebs, Arms Races between and within Species. *Proc. R. Soc. Lond. B Biol.*
Sci. **205**, 489–511 (1979).

59.   J. Felsenstein, Phylogenies and the comparative method. *Am. Nat.* **125**, 1–15 (1985).

60.   C. Williams, H.-J. Hoppe, D. Rezgui, M. Strickland, B. E. Forbes, F. Grutzner, S. Frago, R. Z.
Ellis, P. Wattana-Amorn, S. N. Prince, O. J. Zaccheo, C. M. Nolan, A. J. Mungall, E. Y. Jones, M.
P. Crump, A. B. Hassan, An Exon Splice Enhancer Primes IGF2:IGF2R Binding Site Structure
and Function Evolution. *Science* **338**, 1209–1213 (2012).

61.   D. E. McCoy, D. Haig, Embryo Selection and Mate Choice: Can 'Honest Signals' Be Trusted?
Trends Ecol. Evol. **35**, 308–318 (2020).

62.   D. Arendt, J. M. Musser, C. V. H. Baker, A. Bergman, C. Cepko, D. H. Erwin, M. Pavlicev, G.
Schlosser, S. Widder, M. D. Laubichler, G. P. Wagner, The origin and evolution of cell types.
Nat. Rev. Genet. **17**, 744–757 (2016).

63.   D. Arendt, P. Y. Bertucci, K. Achim, J. M. Musser, Evolution of neuronal types and families.
Curr. Opin. Neurobiol. **56**, 144–152 (2019).

64.   E. S. Lau, T. H. Oakley, Multi-level convergence of complex traits and the evolution of
bioluminescence. *Biol. Rev.* **96**, 673–691 (2021).

65.   G. Cornelis, C. Vernochet, Q. Carradec, S. Souquere, B. Mulot, F. Catzeflis, M. A. Nilsson, B. R.
Menzies, M. B. Renfree, G. Pierron, U. Zeller, O. Heidmann, A. Dupressoir, T. Heidmann,
Retroviral envelope gene captures and syncytin exaptation for placentation in marsupials. *Proc.*
Natl. Acad. Sci. **112**, E487–E496 (2015).

66.   G. Cornelis, C. Vernochet, S. Malicorne, S. Souquere, A. C. Tzika, S. M. Goodman, F. Catzeflis,
T. J. Robinson, M. C. Milinkovitch, G. Pierron, O. Heidmann, A. Dupressoir, T. Heidmann,
Retroviral envelope syncytin capture in an ancestrally diverged mammalian clade for placentation
in the primitive Afrotherian tenrecs. *Proc. Natl. Acad. Sci.* **111**, E4332–E4341 (2014).

67.   C. Vernochet, O. Heidmann, A. Dupressoir, G. Cornelis, P. Dessen, F. Catzeflis, T. Heidmann, A
syncytin-like endogenous retrovirus envelope gene of the guinea pig specifically expressed in the
placenta junctional zone and conserved in Caviomorpha. *Placenta* **32**, 885–892 (2011).

68.   D. Emera, C. Casola, V. J. Lynch, D. E. Wildman, D. Agnew, G. P. Wagner, Convergent
Evolution of Endometrial Prolactin Expression in Primates, Mice, and Elephants Through the
Independent Recruitment of Transposable Elements. *Mol. Biol. Evol.* **29**, 239–247 (2012).

69.   V. J. Lynch, R. D. Leclerc, G. May, G. P. Wagner, Transposon-mediated rewiring of gene
regulatory networks contributed to the evolution of pregnancy in mammals. *Nat. Genet.* **43**, 1154–
1159 (2011).

70.   C. R. Henthorn, P. M. Airs, E. K. Neumann, M. Zamanian, Resolving the origins of secretory
products and anthelmintic responses in a human parasitic nematode at single-cell resolution. *eLife*
12, e83100 (2023).

71.   M. Hu, Y. Bai, X. Zheng, Y. Zheng, Coral–algal endosymbiosis characterized using RNAi and
single-cell RNA-seq. *Nat. Microbiol.* **8**, 1240–1251 (2023).

72.   A. L. Keyte, K. K. Smith, Basic Maintenance and Breeding of the Opossum Monodelphis
domestica. *Cold Spring Harb. Protoc.* **2008**, pdb.prot5073 (2008).

73.   N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

74.   J. Kiernan, *Histological and Histochemical Methods* (Scion Publishing Ltd, 5th Edition., 2015).

75.   B. M. Kirilenko, C. Munegowda, E. Osipova, D. Jebb, V. Sharma, M. Blumer, A. E. Morales, A.-

W. Ahmed, D.-G. Kontopoulos, L. Hilgers, K. Lindblad-Toh, E. K. Karlsson, Zoonomia
Consortium, M. Hiller, Integrating gene annotation with orthology inference at scale. *Science* **380**,
eabn3107 (2023).

76. S. L. Wolock, R. Lopez, A. M. Klein, Scrublet: Computational Identification of Cell Doublets in
Single-Cell Transcriptomic Data. *Cell Syst.* **8**, 281-291.e9 (2019).

77. A. Gayoso, J. Shor, JonathanShor/DoubletDetection: doubletdetection v4.2, version v4.2, Zenodo
(2022); https://doi.org/10.5281/zenodo.6349517.

78. F. A. Wolf, P. Angerer, F. J. Theis, SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol.* **19**, 15 (2018).

79. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk, C.
Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T.
Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P.
Smibert, R. Satija, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
(2021).

80. I. Korsunsky, N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M. Brenner,
P. Loh, S. Raychaudhuri, Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

81. V. A. Traag, L. Waltman, N. J. van Eck, From Louvain to Leiden: guaranteeing well-connected
communities. *Sci. Rep.* **9**, 5233 (2019).

82. H. Heaton, A. M. Talman, A. Knights, M. Imaz, D. J. Gaffney, R. Durbin, M. Hemberg, M. K. N.
Lawniczak, Souporcell: robust clustering of single-cell RNA-seq data by genotype without
reference genotypes. *Nat. Methods* **17**, 615–620 (2020).

83. G. La Manno, R. Soldatov, A. Zeisel, E. Braun, H. Hochgerner, V. Petukhov, K. Lidschreiber, M.
E. Kastriti, P. Lönnerberg, A. Furlan, J. Fan, L. E. Borm, Z. Liu, D. van Bruggen, J. Guo, X. He,
R. Barker, E. Sundström, G. Castelo-Branco, P. Cramer, I. Adameyko, S. Linnarsson, P. V.
Kharchenko, RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

84. J. M. Musser, G. P. Wagner, Character trees from transcriptome data: Origin and individuation of
morphological characters and the so-called "species signal." *J. Exp. Zoolog. B Mol. Dev. Evol.*
324, 588–604 (2015).

85. S. A. Lambert, A. Jolma, L. F. Campitelli, P. K. Das, Y. Yin, M. Albu, X. Chen, J. Taipale, T. R.
Hughes, M. T. Weirauch, The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

86. E. M. Erkenbrack, J. D. Maziarz, O. W. Griffith, C. Liang, A. R. Chavan, M. C. Nnamani, G. P.
Wagner, The mammalian decidual cell evolved from a cellular stress response. *PLOS Biol.* **16**,
e2005594 (2018).

87. J. DiFrisco, A. C. Love, G. P. Wagner, Character identity mechanisms: a conceptual model for
comparative-mechanistic biology. *Biol. Philos.* **35**, 44 (2020).

88. C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T. L. Madden,
BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

89. A. A. Hagberg, D. A. Schult, P. J. Swart, "Exploring Network Structure, Dynamics, and Function
using NetworkX" in *Proceedings of the 7th Python in Science Conference*, G. Varoquaux, T.
Vaught, J. Millman, Eds. (Pasadena, CA USA, 2008), pp. 11–15.

90. A. Brückner, J. M. Badroos, R. W. Learsch, M. Yousefelahiyeh, S. A. Kitchen, J. Parker,
Evolutionary assembly of cooperating cell types in an animal chemical defense system. *Cell* **184**,
6138-6156.e28 (2021).

91. B. Muzellec, M. Teleńczuk, V. Cabeli, M. Andreux, PyDESeq2: a python package for bulk RNA-
seq differential expression analysis. *Bioinformatics* **39**, btad547 (2023).

92. L. Garcia-Alonso, L.-F. Handfield, K. Roberts, K. Nikolakopoulou, R. C. Fernando, L. Gardner,
B. Woodhams, A. Arutyunyan, K. Polanski, R. Hoo, C. Sancho-Serra, T. Li, K. Kwakwa, E.

Tuck, V. Kleshchevnikov, A. Tarkowska, T. Porter, C. I. Mazzeo, S. van Dongen, M. Dabrowska, V. Vaskivskyi, K. T. Mahbubani, J. Park, M. Jimenez-Linan, L. Campos, V. Kiselev, C. Lindskog, P. Ayuk, E. Prigmore, M. R. Stratton, K. Saeb-Parsy, A. Moffett, L. Moore, O. A. Bayraktar, S. A. Teichmann, M. Y. Turco, R. Vento-Tormo, "Mapping the temporal and spatial dynamics of the human endometrium in vivo and in vitro" (2021); https://doi.org/10.1101/2021.01.02.425073.

93. D. Dimitrov, P. S. L. Schäfer, E. Farr, P. R. Mier, S. Lobentanzer, A. Dugourd, J. Tanevski, R. O. R. Flores, J. Saez-Rodriguez, LIANA+: an all-in-one cell-cell communication framework. bioRxiv [Preprint] (2023). https://doi.org/10.1101/2023.08.19.553863.

94. S. Louca, M. Doebeli, Efficient comparative phylogenetics on large trees. *Bioinformatics* **34**, 1053–1055 (2018).

95. E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2019).

96. E. Armingol, A. Officer, O. Harismendy, N. E. Lewis, Deciphering cell–cell interactions and communication from gene expression. *Nat. Rev. Genet.* **22**, 71–88 (2021).

97. M. Efremova, M. Vento-Tormo, S. A. Teichmann, R. Vento-Tormo, CellPhoneDB: inferring cell–cell communication from combined expression of multi-subunit ligand–receptor complexes. *Nat. Protoc.* **15**, 1484–1506 (2020).

98. S. Jin, C. F. Guerrero-Juarez, L. Zhang, I. Chang, R. Ramos, C.-H. Kuan, P. Myung, M. V. Plikus, Q. Nie, Inference and analysis of cell-cell communication using CellChat. *Nat. Commun.* **12**, 1088 (2021).

99. M. Büttner, J. Ostner, C. L. Müller, F. J. Theis, B. Schubert, scCODA is a Bayesian model for compositional single-cell data analysis. *Nat. Commun.* **12**, 6876 (2021).

Acknowledgments

The authors thank Katie Grabek of Fauna Bio, Michael Hiller and Evgeny Leushkin for sharing the draft *Tenrec ecaudatus* genome. The study benefited from input from Caitlin McDonough-Goldstein and Jan Engelhardt. Steve Stearns and Arun Chavan provided critical comments on the manuscript. The Yale Center for Genomic Analysis (NIH 1S10OD030363-01A1) and Yale West Campus Imaging Core supported this work.

Funding

The research presented was supported by the Yale Institute for Biospheric Studies, the John Templeton Foundation (#61329), the National Science Foundation (NSF IOS 1655091) and the Austrian Science Fund (FWF 33540). D.J.S. was supported by the NIH (T32 GM 007499) and D.J.S. and S.B. were supported by the NSF g2p2pop Research Coordination Network (RCN 1656063). G.P.W. receives research support from the Hagler Institute of Advanced Studies at Texas A&M University.

Author Contributions

Conceptualization: DJS, SBM, GPW, MP
Funding acquisition: GPW, MP, FvB, DJS, SBM
Experimentation: DJS, SBM, JDM, AGC, GPW
Animal Colony Maintenance: GRS, GD, JDM, DJS, SBM
Formal analysis: DJS, SBM
Software: DJS

903 Supervision: GPW, MP
904 Writing - original draft: DJS, SBM
905 Writing - review & editing: DJS, SBM, GPW, MP
906

Competing Interests

908   Authors declare that they have no competing interests.

Data and Materials Availability

Sequencing data have been uploaded to the NCBI Gene Expression Omnibus at
accession GSE274701. Human gene expression data used in this analysis were retrieved
from https://www.reproductivecellatlas.org/ and macaque data from the NCBI accession
GSE180637.
Data to reproduce figures and output from analyses are available in the Supplementary
Materials. Analytical code for this study is available at https://gitlab.com/dnjst/fmi2024. Additional
processing scripts for cell-cell communication data are archived in a python package at
https://gitlab.com/dnjst/chinpy.
918

Supplementary Materials

920 Figs. S1 to S6
921 Tables S1 to S8

Materials and Methods

Animal Husbandry
Pregnancy samples were taken at gestation points after the establishment of the
definite placenta and, in the case of guinea pigs, before the luteal-placental shift in
progesterone production. Time points chosen were in opossum day 13.5 of 14.5 total
gestation, in tenrec day 28-29 of 56, in guinea pig day 30.5 of 62, and in mouse day 15.5 of
19.5 days gestation. These were integrated with the human atlas generated from 4-13 weeks,
roughly approximate days 28-91 out of a 280-day gestation, and macaque samples generated
from stages ranging from 20-140 of a 162-day gestation.
M. domestica were raised in a breeding colony at Yale University according to
established technical protocols(*72*) and ethical protocols approved by the Yale University
Institutional Animal Care and Use Committee (#2020-11313). Male and female animals were
housed separately after 3 months of age, at which point female opossums were introduced to
the male room for sexual preconditioning, and breeding was attempted after 6 months. During
breeding, non-cycling female opossums were introduced to the male room for 1 day, and then
subsequently swapped into the used cage of a prospective male partner for 5 days. After this
period, both individuals were placed into a breeding cage and video recorded to assess the
time of copulation. If multiple copulations were observed, the first was always used to
calibrate 0 dpc; samples were taken one day before parturition at 13.5 dpc (n=2).
T. ecaudatus were maintained in a breeding colony at the University of Nevada, Las
Vegas according to approved University of Nevada, Las Vegas Institutional Animal Care
protocols. Individuals in the colony descend from a population of 40 wild-caught animals
imported from Mauritius in June 2014. Animals were mated and mid-gestation was sampled

for scRNA-seq between 28 and 29 days (n=3) following the first exposure of females to
males.
947   *C. porcellus* (Charles River) were maintained at the University of Vienna according to
948   Institutional Animal Care protocols, on standard chow and water ad libitum. The estrus cycle
949   was monitored by examination of vaginal membrane opening and the animals were mated in
950   estrus at 3-4 months of age, and video recorded to detect copulation. Midgestation samples
951   (n=2) were collected 30.5 days post copulation.
952   *M. musculus* (C57BL/6J) were maintained at the University of Vienna on standard
chow and water ad libitum according to University Institutional Animal Care protocols. Animals
of age 2.5-4 months were used for the experiments. Estrus cycle was monitored by vaginal
swabbing, and the females were mated when approaching estrus. Copulation was determined
by the presence of a copulatory plug. The start of gestation is counted from the midnight of
the night preceding the detection of copulatory plug. Midgestation samples (n=2) were
collected 15.5 days post copulation.

Single-Cell RNA Sequencing

961   Uteri were dissected into phosphate-buffered saline (PBS), separated from the cervix
and fallopian tube, and opened longitudinally. Embryos and directly attached extraembryonic
membranes were removed via severing of the umbilical cords. In the guinea pig, the labyrinth
was removed prior to sample preparation. Tissue samples were kept at 4°C for approximately
30 minutes until they could be dissociated into a single-cell suspension. Due to necessity,
Tenrec samples were shipped in RPMI medium on ice overnight and dissociated the next
day.
968   All tissue was processed as follows, with small species-specific modifications as noted.
Tissue was minced with a scalpel into ~1 mm^3 cubes in 2 mL of digestive solution containing
0.2 mg/mL Liberase TL (05401020001, Sigma). Tissue suspensions were heated at 37°C for
15 minutes and then passed 10 times through a 16-gauge needle attached to a 3-mL syringe.
This incubation and needle passage process was repeated two more times, with the
substitution of an 18-gauge needle. Finally, 2 mL of charcoal-stripped fetal bovine serum
(100-199, Gemini) was added and the suspension was immediately passed through a 70-µm
cell strainer then a 40-µm cell strainer. The flowthrough was pelleted by centrifugation at 500
g for 5 minutes and cells were resuspended in 1x ACK red blood cell lysis buffer (A1049201,
Thermo-Fisher), incubated at room temperature for 5 minutes, centrifuged again and
resuspended in PBS containing 0.04% bovine serum albumen (A9647, Sigma). At this point,
cells were examined on a hemocytometer to gauge concentration and check for debris and
cell death using trypan blue stain (15250061, ThermoFisher).
981   Mouse tissue was processed as above with the addition of Accumax (Stem Cell
Technologies) to the final 0.04% bovine serum albumen solution. Guinea pig tissue was
processed as with the mouse with the following modifications: a wide-bore 1 mL pipette tip
was substituted for the 16-gauge needle, and 0.2 mg/mL collagenase I (17018029, Thermo-
Fisher) was added to the digestion solution. As tenrec and guinea pig tissue showed higher
cell death, a fractionated protocol was adopted to reduce exposure of cells to digestive
enzymes. For these species, cells which had dislodged following each passage step were
separated from intact tissue and immediately passed through a 70-µm cell strainer into 2 mL
of charcoal-stripped fetal bovine serum (100-199, Gemini), while remaining intact tissue was
allowed to continue with further digestion. Each fraction was centrifuged at 300 rpm for 5
minutes and resuspended in 500 µL ACK red blood cell lysis buffer (A1049201, Thermo-

Fisher) for 5 minutes, then centrifuged at 500 rpm for 2 minutes and resuspended in 0.04%
bovine serum albumen (A9647, Sigma) and kept on ice until dissociation was complete. Cells
were counted on a hemocytometer and recombined in a proportional manner before
sequencing.
Cells were captured using the 10X Chromium platform (3' chemistry, version 3),
libraries were generated according to manufacturer protocols (CG000315). Libraries were
sequenced using an Illumina NovaSeq by the Yale Center for Genomic Analysis (*M.*
domestica and T. ecaudatus) and at the Next Generation Sequencing Facility of Vienna
Biocenter (*C. porcellus and M. musculus*).

Histology and Histochemistry
Histological samples were fixed for 24 hours in 10% neutral buffered formalin, followed
by dehydration to 70% Ethanol and kept at 4°C (days-weeks) or -20°C (months) before
paraffin embedding. Staining with Gill 2 hematoxylin and Y alcoholic eosin was performed
according to(*74*). Bright-field images for Figure 1 were taken with a Leica Thunder Imager
(*Tenrec*), a Nikon Eclipse E600 (*Monodelphis*), and an EVOS M7000 (*Cavia* and *Mus*).
Immunohistochemistry was performed using antibodies recorded in **Table S6**.
For chromogenic immunohistochemistry, slides were incubated for 1 hour at 65°C,
dewaxed with xylene, washed in 100% ethanol, and rehydrated in running tap water. Antigen
retrieval was performed for 1 hour in a 95°C vegetable steamer containing 10 uM sodium
citrate (pH 6.0). Slides were washed in phosphate-buffered saline and blocked with 0.1%
mass/volume bovine serum albumen. Peroxidases were blocked by incubation in 0.03%
hydrogen peroxide containing sodium azide (DAKO) in a humidification chamber for 30
minutes. Slides were incubated overnight at 4°C in primary antibody solutions, blocked,
treated with horseradish peroxidase-conjugated secondary antibody for 1 hour at room
temperature. Finally, slides were washed in blocking solution once more and treated with 3,3'-
diaminobenzidine (DAKO K401011-2) for 5 minutes and counterstained with Gill 2
hematoxylin before brightfield imaging.
For fluorescent immunohistochemistry, the procedure had the following modifications.
Peroxidase blocking and 3,3'-diaminobenzidine was not used in favor of secondary
antibodies, conjugated with the fluorophore. Images were acquired using the EVOS M7000
Imaging System (Invitrogen). Fluorescent images were also captured in the red channel and
superimposed to leverage the autofluorescent nature of erythrocytes to highlight the location
of vasculature.
1025

In Situ Hybridization
Single-molecule fluorescence *in situ* hybridization was conducted on formalin-fixed
paraffin-embedded sections using the RNAscope platform following the manufacturer's
protocols (ACD Bio 323100-USM for fluorescent and 322310-USM for chromogenic). Probes
were designed against the equivalent transcript sequences used for RNA-seq alignment and
deposited into the RNAscope database as standardized probes. Probes used are listed in
Table S7.
Fluorescent RNAscope was performed using the MultiPlex platform (ACD Bio 323100)
with Opal 520, Opal 570, Opal 620, and Opal 690 fluorophores (Akoya Biosciences).
Fluorescent images were captured using a laser scanning confocal microscope (Leica
Stellaris 8 Falcon). For Figure 3b, Fluorescent images were also captured at 690 nm and
superimposed to leverage the autofluorescent nature of the opossum glands to highlight the

location of glandular epithelia. Chromogenic RNAscope was performed using the RNAscope
2.5 HD Brown Assay (ACD Bio 322300) and imaged on an EVOS M7000 microscope.

Single-Cell Data Analysis

Reads were obtained in FASTQ format from the respective sequencing cores. *M.*
fascicularis reads were downloaded from the NCBI Gene Expression Omnibus (GSE180637),
and *H. sapiens* aligned counts were downloaded from https://www.reproductivecellatlas.org.

Reads were aligned to reference genomes using the 10X Genomics CellRanger
software (≥v7.0.0). *Monodelphis domestica*, *Cavia porcellus*, *Mus musculus*, and *Macaca*
fascicularis were mapped to their respective Ensembl genome annotations (ASM229v1 v104,
cavPor3 v104, GRCm39 v104, and Macaca_fascicularis_6.0 v112, respectively). For *Tenrec*
ecaudatus a novel unpublished genome was provided by the laboratory of Michael Hiller and
Fauna Bio, annotated using the TOGA pipeline(*75*).

Quality control included filtering cells with low count (less than 700 unique features for
maternal cells and less than 500 unique features for fetal cells) or high mitochondrial gene
expression (more than 25%), followed by doublet detection by scrublet (v0.2.3)(*76*) and
doubletdetection (v4.2)(*77*), to remove clusters consisting of majority doublets. Library
size normalization, log1p normalization, feature selection, and dimensionality reduction
were performed using scanpy ≥ v1.9.1(*78*), with the exception of the mouse, which was
annotated using equivalent functions in Seurat v4(*79*) and exported as raw counts and then
reprocessed in scanpy. Principal components were adjusted to correct for batch effect across
biological replicates using harmony (harmonypy, v0.0.9, r-harmony, v0.1)(*80*). UMAP
embeddings were calculated and Leiden(*81*) clustering was performed to partition cells into
putative cell types. Marker genes were calculated based upon differential expression (logistic
regression, Wilcoxon rank-sum test, and t-test as implemented by
scanpy.pp.rank_genes_groups) and used to annotate groups. Cells were annotated using a
combination of these markers and active genes belonging to NMF gene expression modules
(see below), and cell clusters lacking uniquely distinguishable NMF gene modules or markers
were merged. Fetal cells were identified by a combination of placenta-specific marker gene
expression, and non-overlap with non-pregnant uterine samples. In the case of the tenrec
which was sufficiently outbred, single nucleotide polymorphism-based inference of the
genome of origin using souporcell (v2.5)(*82*) (**Figure S1**). Clusters of cells which souporcell
assigned to the genome of lesser abundance (putative fetal) overlapped with those annotated
as fetal by differential gene expression. Spliced and unspliced transcript calling was
performed using the velocyto package (v0.17.17)(*83*) and RNA velocity was modeled using
scVelo (v0.30)(*43*).

Because the ground-truth database was curated from the human literature and
insufficient data exist to curate ligand-receptor pairs for each species, the non-human
transcriptomes were converted into "human-equivalent" transcriptomes by mapping of genes
to their top human ortholog as detected by BLAST of translated peptide sequence(*14*). To
maximize coverage, in cases of many:one human orthology, counts of all detected paralogs
were pooled together. This transformed expression matrix was used only for cell
communication analysis and cell phylogeny inference. Dimensionality reduction, cluster
identification, differential expression testing, and marker gene identification and plotting were
conducted on unadulterated species-relevant gene sets.
1082

Cell Type Phylogeny

A phylogenetic tree of cell type transcriptomes was calculated using cell type-aggregated transcriptomes from all species, with orthology resolved using the human BLAST method described above. Clusters with high annotation uncertainty, extremely low capture rate, or possible artifactual nature ('Md_LE-OAT', 'Te_T-gd', 'Md_PMN-NE', 'Mm_TC-NKT', 'Mm_NEU', 'Te_NEU-CHL1', 'Hs_Granulocytes', 'Mm_fMES-Myoblast', 'Mm_fFB-ribo'), with high erythroid signature ('Mf_Mega_progenitor', 'Mf_Mye_progenitor', 'Mf_e&ysEry2', 'Mf_e&ysEry3', 'Mf_EMP', 'Mf_BP'), and those originating from the embryo proper ('Mf_LP._Meso', 'Mf_AI', 'Mf_PS', 'Mf_PGC', 'Mf_Node', 'Mf_EPI') were excluded from analysis. Gene expression matrices were corrected for "species signal"(*84*) by linear regression to remove the species of origin covariate from cell transcriptomes with the scanpy.pp.regress_out function. A euclidean distance matrix between cell type transcriptomes was then calculated and a neighbor-joining tree generated using the TreeMethods python package (v1.0.3; https://github.com/BradBalderson/TreeMethods). An alternative tree was also generated using the same approach with the additional subsetting of genes to only those annotated as transcription factors in Lambert et al.(*85*). Cartoons of cell types were modified from BioRender, Erkenbrack et al.(*86*), DiFrisco et al.(*87*), and original art.

Homology Inference by SAMap

SAMap (v.1.3.4)(*14*) was used to calculate gene homology-aware similarity scores between cells of different species. BLAST (v2.14.1)(*88*) graphs were generated from the ENSEMBL proteomes of respective species, with the exception of *Tenrec ecaudatus* for which the transcriptome was used with tblastx mode. SAMap mapping tables were exported and analyzed using hierarchical clustering (seaborn, v0.13.1) as well as network analysis (networkx, v3.2.1)(*89*). For network analysis, SAMap mapping scores were used as edge weights and cell types as nodes. Communities were detected using the Leiden algorithm (leidenalg, v0.10.2)(*81*). Stability of the resulting communities was assessed by repeating community detection with a range of 10000 random initial states and constructing a co-occurrence matrix between cell types: cluster stability was defined as the mean of all co-occurrence values between members of the final community.

Gene Expression Program Identification and Cross-Species Comparison

cNMF v1.4.1(*12*) was used to infer gene expression programs from count matrices of each species. Analysis was performed following a modified protocol from(*90*). Optimal numbers of factors (K) were chosen based upon manual examination of stability-error curves over a range of possible K values from 10-50, selecting a value immediately preceding a sudden drop in stability score (**Figure S2**).

For cross-species comparison, spectra scores from cNMF (i.e., gene loadings in NMF factors) were used. Non-human genes were mapped to their closest human orthologs as determined by BLAST score (see above), and in cases of many-to-one mapping, the highest loading score was used. Pearson correlation was calculated between programs and hierarchical clustering conducted using the clustermap function in seaborn (v0.13.1).

In Vitro Decidualization of Primary Tenrec Stromal Cells

Mesenchymal stromal cells were isolated from the midgestation *Tenrec ecaudatus* uterus by differential attachment. Cells were expanded in T25 flasks in a growth medium consisting of 15.5 g/L Dulbecco's modified eagle medium (30-2002, ATCC), 1.2 g/L sodium bicarbonate, 10 mL/L sodium pyruvate (11360, Thermo Fisher), and 1 mL/L ITS supplement (354350, VWR) in 10% charcoal-stripped fetal bovine serum (100-199, Gemini), and then seeded into 12-well plates (3.9 cm^2) for 1 day. Samples were treated with either base medium (Base), 1 µM medroxyprogesterone acetate (MPA; M1629, Sigma-Aldrich), or 1 µM MPA and 0.5 mM 8-bromoadenosine 3',5'-cyclic adenosine monophosphate (cAMP; B7880, Sigma-Aldrich) for 6 days, with replenishment on day 3 (n=4 for each condition, total n=12). Bulk RNA was isolated using a Qiagen RNeasy Micro Kit (74004), libraries were prepared by the Yale Center for Genomic Analysis and sequenced using an Illumina NovaSeq. Reads were aligned to the draft genome using kallisto (v0.45.0). Significance of differentially-expressed genes was calculated using the Wald test function of pyDESeq2 (v0.4.10)(*91*) with multiple test adjustment using the Benjamini-Hochberg method.

Cell-Cell Communication Analysis

Putative cell interactions were inferred using expression thresholding, followed by statistical testing for significantly cell type-enriched interactions. A ground truth ligand-receptor database was built as a manually extended fork of CellPhoneDB v5.0.0(*92*) with additional curation and metadata. This modified list is archived at https://gitlab.com/dnjst/ViennaCPDB/. From these transcriptomes and the ground-truth list, interaction scores for all cell type pairs were generated using the LIANA+ (v1.2.0)(*93*) and chinpy (https://gitlab.com/dnjst/chinpy; v0.0.55) frameworks. Single-cell transcriptomes from multiple biological replicates (sample sizes reported above) were pooled before inference of cell communication was performed, following common practice(*96*). Instead of biological replicate-based testing, other statistical tests, such as CellPhoneDB(*97*) and CellChat(*98*) permutation tests, are standard; output from these tests (calculated as described above using LIANA+) are available for ligand-receptor interactions of all species as **Supplementary Data**. Likewise, the sample size of scRNA-seq studies complicates statistical testing of differential cell type abundance(*99*), so the relative abundance measures in **Figure 1d** are presented as pooled values.

For co-evolutionary integration analysis, all ligand-receptor interactions in the ground truth database were classified into "off" (ligand and receptor off), "allocrine ligand" (ligand on, receptor off), "allocrine receptor" (receptor on, ligand off), or "autocrine" (ligand and receptor on) states, based upon an expression threshold of 20% of cells in a cluster. Classification of ligands was done in this manner to measure intrinsic signaling potential for each individual cell type in a way which is not inflated by the number of clusters of other cell types identified in the tissue, in contrast to network analysis, where the splitting of a given cell type into multiple clusters inflates the number of inferred outgoing interactions for other cell types. Statistical associations between cell class and number of allocrine ligands were assessed using a one-way analysis of variance. Kleinberg hub and authority scores(*54*) were calculated for all cell types within their respective species signaling networks using the HITS algorithm as implemented in the hub_score() and authority_score() functions in igraph (v0.11.6), with edge weights set to the LIANA+ "expression products" of each inferred interaction (product of ligand and receptor log-normalized transcripts per 10,000).

For disambiguation analysis, a manually-corrected version of CellPhoneDB "classification" metadata was used to group ligands into ligand families. To preserve gene

duplication events, in cases of many-to-one homology, all possible homologs to the human
ground truth ligand list were included in the target species' ligand family using Ensembl
(mouse, guinea pig, opossum) or BLAST (tenrec) orthology. Expected values of
disambiguated ligands were calculated as described in Statistical Analysis.
For discrete ancestral state reconstruction, interactions were scored in a boolean on/off
manner based on a nonzero expression threshold of 10% of cells in a cluster for all subunits
of both ligand and receptor. Maximum parsimony was calculated using the castor package
(v1.8.0)(94). Edge changes were calculated for all internal branches of the 5-species tree by
identifying genes considered "on" in descendant nodes but inferred "off" in the immediate
common ancestor node, or vice versa.
For continuous ancestral state reconstruction, restricted maximum likelihood via the
ape package (v5.8)(95) was used to estimate ancestral states. Expression values for
continuous analysis were in natural log-transformed transcripts per 10,000 scale
(scanpy.pp.log1p). To reduce artifacts, only genes expressed in at least 4 of 5 species were
subjected to continuous ancestral state reconstruction (i.e., one loss/absence of a gene was
tolerated). Changes in continuous expression values for ligands and receptors along internal
branches of the 5-species tree were calculated by subtracting descendant node expression
values from inferred immediate ancestral node values.
For escalation analysis, phylogenetic independent contrasts(59) (PIC) were calculated
for ligand and corresponding receptor expression values (scaled as above) using the pic()
function in ape (v5.8)(95). Linear regression was performed on the PIC values using the
model ligand PIC ~ receptor PIC + 0. That is, with ligand PIC as the independent variable and
receptor PIC as the dependent variable, with the intercept forced to zero (assuming that when
there is no change in ligand expression, there is no change expected in receptor expression).
P-values reported as "PIC P-value" result from a two-sided t-test on the estimated slope
coefficient of linear regression of ligand versus receptor PICs. Linear regression was also
performed on observed expression values to return a regression slope.

Statistical Analysis of "Disambiguation"
Disambiguation of ligands was compared to a null model in which the probability of
randomly redistribution of ligands to cell types leading to exclusively maternal or exclusively
fetal expression is assessed. First, all secreted peptide and small molecule-mediated ligand-
receptor interactions were classified into "fetal-only", "maternal-only", or "both" expression
groups based on the threshold value of 20% or more of the cells sequenced expressing the
ligand in. Ligands were grouped into ligand families using an additionally curated version of
the CellPhoneDB (v5.0.0) "classification" metadata. For each ligand family, the per-cell type
probability of a given ligand being expressed was calculated according to the equation:

$$P_{on} = \frac{L_{expressed}}{S_{pathway} \times (N_{fetal} + N_{maternal})}$$

1209
1210
1211 where $L_{expressed}$ is the observed number of "on" calls within the ligand family across fetal and
1212 maternal cell types, N_{fetal} is the number of fetal cell types, $N_{maternal}$ the number of maternal cell
1213 types, and $S_{pathway}$ the number of ligands in the family.
1214   The null probabilities of fetal-only and maternal-only expression for a ligand,
1215 respectively, were given by:

$$P_{fetalonly} = \left(1 - (1 - P_{on})^{N_{fetal}}\right) \cdot (1 - P_{on})^{N_{maternal}}$$

1216
1217

1218
$$P_{maternalonly} = \left(1 - \left(1 - P_{on}\right)^{N_{maternal}}\right) \cdot \left(1 - P_{on}\right)^{N_{fetal}}$$

1219

The expected fetal-only and maternal-only ligands in each family were obtained by
multiplying $P_{fetalonly}$ and $P_{maternalonly}$ by the number of ligands in the family.
Finally, a one-way chi-squared goodness of fit test for identity of observed and
expected counts of the 3 scoring categories (coexpressed, fetal-only, and maternal-only)
was used to obtain chi-squared and p-values, with individual observations
corresponding to ligands within a family. The Benjamini-Hochberg correction was used
(statsmodels.multitest.multipletests v0.14.1) with an alpha of 0.05 to obtain false
discovery rate q-values. For plotting, ligand families were considered significant if p ≤
0.05 and q ≤ 0.05. Bonferroni-corrected p-values, obtained by multiplying *p* by the
number of ligands in a family, are also reported in **Table S3**.

Fig. 1. Single-cell transcriptomic atlases of five mammalian species spanning the diversification of viviparity – from left to right: *Monodelphis domestica*, *Tenrec ecaudatus, Cavia porcellus, Mus musculus, Homo sapiens*. (**A**) UMAP embeddings of single-cell transcriptomes from the species listed. (**B**) Histology of the fetal-maternal interface of the species in (a). Color bars reflect tissue regions (legend on right) from fetal (F) to maternal (M). (**C**) Unified cross-species UMAP (SAMap), colored by cell type, with prominent divisions annotated in gray. (**D**) Proportional abundances of cell types belonging to major cell type families in each sample. (**E**) A neighbor-joining phylogenetic tree showing hierarchical relationships of cell clusters recovered, colored by broad families. For a full list of cell type descriptions, see Supplementary Material. CTB: cytotrophoblast; DC: dendritic cell; dS1/dS2/dS3: decidual stromal cell; eS1: endometrial stromal fibroblast; EPT: extraplacental trophoblast; iEPT: invasive extraplacental trophoblast; EVT: extravillous trophoblast; eEVT: endovascular extravillous trophoblast; iEVT: interstitial extravillous trophoblast; fFB/fMES: fetal mesenchymal cell; GE: glandular epithelial cell; HB: Hofbauer cell; ILC: innate lymphoid cell; LE: luminal epithelial cell; LEC: lymphatic endothelial cell; MET: mesothelial cell; MO: monocyte; MPH: macrophage; NK/NKC: natural killer cell; PC/PV: pericyte; PMN: polymorphonuclear cell; SCT: syncytiotrophoblast; SMC: smooth muscle cell; SpT: spongiotrophoblast; TC: T cell; T-reg: regulatory T cell; TF: tissue fibroblast; TGC: trophoblast giant cell; TGC-SpA: spiral artery-remodeling trophoblast giant cell; GlyT: glycogen trophoblast cell; GlyT_Prog: glycogen trophoblast progenitor; TGC_S: sinusoidal trophoblast giant cell; TGC_C: canal trophoblast giant cell; VEC: vascular endothelial cell; fVEC: fetal vascular endothelial cell; YSC: yolk sac cell; YSC-P: parietal yolk sac cell; YSC-V: visceral yolk sac cell.

Fig. 2. Trophoblast cell types and homology inference. (**A**) Expression of trophoblast marker genes. SCT: syncytiotrophoblast; CTB: cytotrophoblast; EPT/EVT: extraplacental/extravillous trophoblast; YSC: yolk sac cell; TGC: giant cell; CCC: cell column. (**B**) Histological localization of select cell type markers from (a). epi.: epithelium; myom.: myometrium; ery.: erythrocytes. Scale bars = 100 μm. (**C**) Network of SAMap linkages between trophoblast cells, with edge thicknesses are proportional to mapping scores. Edges with scores less than 0.1 are not shown. Nodes are colored by inferred Leiden community. (**D**) Expression of select "pan-invasive" genes in trophoblast of all species, grouped by SAMap homology group. Phenotypic covariates (endoreduplication, invasion, fusion) are annotated below cell type labels. Invasiveness groups by community, whereas fusion and endoreduplication are widespread. Md: *Monodelphis domestica*; Te: *Tenrec ecaudatus*; Cp: *Cavia porcellus*; Mm: *Mus musculus*; Mf: *Macaca fascicularis;* Hs: *Homo sapiens.*

Fig. 3. Cross-species comparison of decidual stroma. (**A**) Marker gene expression of endometrial stromal and decidual cell types. (**B**) Comparison of human midgestation decidual cells to those generated spontaneously during the secretory phase of the ovarian cycle shows a similarity of predecidua (dS2) to late luteal decidual cells. (**C**) *In situ* (italics) and immunohistochemical (upright) localization of select markers genes from (**A**); cells of interest are marked by an asterisk (*). (**D**) Force atlas embedding of tenrec endometrial stromal cells with RNA velocity vectors overlaid. Expression of select velocity-associated genes in the dS1→dS2 transition in all cells from the left plot, ordered by increasing velocity-informed pseudotime. (**E**) Expression of predecidual markers in primary tenrec uterine stromal cells cultured in vitro with deciduogenic stimuli (MPA: medroxyprogesterone acetate; cAMP) (**F**) Network of SAMap mapping scores between stromal cell types, with edge weight proportional to mapping scores, colored by community as identified by leiden clustering of the graph. Edges with scores less than 0.1 are not shown. (**G**) Correlation heatmap of NMF gene expression programs expressed in decidual stromal cells, with hierarchically-clustered dendrogram on top showing division into two families. (**H**) Tree showing inferred evolutionary relationships of decidual cell types.

Fig. 4. Integration of cell types into cell-cell communication networks and reconstruction of evolutionary changes. (**A**) Numbers of allocrine ligands (receptor unexpressed) and allocrine receptors (ligand unexpressed) for secreted peptide signals. Degrees of integration range from low to high and stratify by cell type class. P-values of analysis of variance relating allocrine ligand degrees to cell type class in each species are plotted below. (**B**) Co-evolution of binary expression of signals by placental cells (left tree) and cognate maternal receptors (right tree) inferred from maximum-parsimony reconstruction. Branches are colored by rate (changes/Myr) using divergence dates from (56). (**C**) Reconstructed secreted signaling interactions in the Placentalia common ancestor, colored by phylogenetic age of gene expression (eutherian novelty: P; ancestral therian: T), partitioned by cell type class and subset to only those derived from the trophoblast.

Fig. 5. Tests of two evolutionary hypotheses for evolutionary dynamics of fetal-maternal communication, "disambiguation" (8) (**A** to **C**) and "escalation" (9) (**D** to **E**). (**A**) Design for disambiguation test. (**B**) Observed vs. expected numbers of coexpressed, fetal-only, and maternal-only ligands, by ligand family. Points with solid color have both $p < 0.05$ and Benjamini-Hochberg false discovery rate $q < 0.05$. (**C**) Disambiguation status of select ligands. Blue: maternal-only; Red: fetal-only; faint: coexpressed. (**D**) Volcano plot of ligand-receptor pairs showing significant phylogenetically-independent correlation in expression (59) on the vertical axis and correlation slope on the horizontal axis. (**E**) Inferred evolutionary changes to ligand and corresponding receptor expression in humans with respect to the human-macaque common ancestor (Catarrhini).

A

.mtx .csv Raw Count Matrices

Empty cell calling

Filtered Count Matrices

Quality control
Normalization
Dimensional reduction
Clustering
Cell type identification

.h5ad Count Matrix with
Cell Type Annotations

cNMF Analysis +
Cluster Refinement

Analysis Platform

python
R

Gene orthology mapping
(BLAST)

.h5ad Standardized Gene +
Cell Type Naming

SAMap (Cell Type
Orthology Mapping)

.csv Interactome
Calculation

Interaction
Phylogenetics

Interactome
Analysis/Plotting

scRNA-seq data
collection

B Souporcell Minority Genome

C Cell Type Annotation

UMAP2

UMAP1

UMAP2

UMAP1

Fig. S1. Single cell RNA-seq analysis. (**A**) Flow chart of analytical design. (**B** and **C**) Souporcell calls for *Tenrec ecaudatus* (k=2 genomes). UMAP embeddings of all cells are shown, colored by cell subtype on the left, and by whether or not the cell was assigned to the less frequent genome (assumed fetal) by souporcell on the right.

Fig. S2. NMF analysis. (**A** to **F**) Optimal numbers of gene expression programs (NMF factors) were determined by identifying local stability values preceding a sudden drop. *Monodelphis domestica*: K=26; *Tenrec ecaudatus*: K=28; *Cavia porcellus*: K=31; *Mus musculus*: K=25. *Homo sapiens:* K=29. *Macaca fascicularis:* K=39. (**G** to **L**) Confusion matrices showing the mean activity score (ranging from 0 to 1) of cells in each cluster, taken from all cells with scores greater than 0.25. (**M** to **P**) Representative "Type II" and "Type III" decidual gene expression program activity scores across individual stromal cells of human (**M**), macaque (**N**), and guinea pig (**O**) show a diversity from exclusively Type II or Type III to a mixture of both programs. An equivalent comparison of Type I and II in the tenrec is depicted in (**P**).

Fig. S3. Additional histological identification of placental cell types. (**A** and **B**) The opossum fetal-maternal interface, showing 3 layers of yolk sac endoderm (YSC), cytotrophoblast (CTB), and syncytial knots (SK), in apposition with maternal luminal epithelium (LE) which produces secretory bodies (*); maternal glandular epithelium (GE) lies closely underneath the lumen. (**C** and **D**) The tenrec invasion front, where trophoblast invades into mucus-filled glands, eroding the glandular epithelium (GE) immunoreactive for pan-cytokeratin stain (CK AE1/AE3). (**E** and **F**) KRT7$^+$ trophoblast in the guinea pig surrounding decidual stroma (left) and maternal arteries (MA; right). (**G** and **H**) Invasive syncytiotrophoblast (SCT) in the guinea pig emanates from subplacental cytotrophoblast (left) and extends into stroma where necrotic patches develop (right). (**I**) Trophoblast giant cell observed in the subplacenta of the guinea pig (proximate to subplacental CTB), but these cells were not captured by droplet-based sequencing. Scale bars: 200 μm.

Fig. S4. Cell type cross-species mapping scores (SAMap). (**A**) Clustered heatmap of pairwise cell type mapping scores (0-1), with selected 1:1 orthogroups annotated in green. (**B** to **E**) Networks subset from the full SAMap mapping table of (**A**) for specific cell type families, colored by Leiden community after community detection was run on the subnetwork. Nodes are cell clusters, and edge weights are equal to score magnitude, with scores less than 0.10 not shown.

Fig. S5. Cell type dendrogram of the cell types studied. (**A**) Neighbor-joining tree of transcription factor-only transcriptomes from all species studied.

Fig. S6. Extended cell-cell communication analysis. (**A**) Graph statistics of all cell types within their respective species' signaling networks. Kleinberg hub scores (a measure of outgoing signaling) are shown on the left and Kleinberg authority scores (a measure of incoming signaling) on the right. Graphs are weighted based on the product of expression values of ligand and receptor. (**B**) Inferred changes in inferred expression magnitude of placental ligands and cognate maternal receptors of secreted signals along internal branches of the phylogeny. Asymmetric escalation and de-escalation events fall into the lower-right and upper-left quadrants, respectively, and uncorrelated changes cluster around the center. (**C**) Phylogenetic tree of species sampled with numbering of nodes and branches labeled as in (**A**).