Flexible reproductive seasonality in Africa-dwelling papionins is associated with low environmental productivity and high climatic unpredictability

Jules Dezeure[1*], Julie Dagorrette[2], Lugdiwine Burtschell[2,3], Shahrina Chowdhury[4,7,8], Dieter Lukas[5], Larissa Swedell[6,7,8,9,10], Elise Huchard[2]

[1]Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France

[2]Institute of Evolutionary Sciences of Montpellier (ISEM), Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France.

[3]AgroParisTech, Paris, France

[4]Department of Anthropology, Brooklyn College, CUNY, 2900 Bedford Ave, Brooklyn, NY 112

[5]Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

[6]Department of Anthropology, Queens College, City University of New York, Flushing, NY 11367-1597, USA

[7]New York Consortium in Evolutionary Primatology, New York, NY, USA

[8]Anthropology Program, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016, USA

[9]Biology and Psychology Programs, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016, USA

[10]Department of Archaeology, University of Cape Town, Rondebosch 7701, Cape Town, South Africa

* corresponding author: dezeurejules@gmail.com

ABSTRACT

At a time when seasonal cycles are increasingly disrupted, the ecology and evolution of reproductive seasonality in tropical vertebrates remains poorly understood. In order to predict how changes in seasonality might affect these animals, it is important to understand which aspects of their diverse patterns of reproductive phenology are linked to either the equally diverse patterns of rainfall seasonality (within-year variations) or instead the marked climatic unpredictability (year-to-year variations) occurring across the intertropical belt. Here, we gather birth and climatic seasonality data from 21 populations of 11 Africa-dwelling primate species from the papionin tribe, occupying a wide range of environments, including equatorial, tropical, temperate and arid climates. We investigate (1) the environmental variations that influence the intensity of reproductive seasonality, and (2) the reproductive stage that is synchronized with increased resource availability. Our results demonstrate wide variation in the intensity of birth seasonality between and within species. Across multiple measures of climatic variation, we found rainfall unpredictability to be the only clear predictor of the intensity of reproductive seasonality across populations, i.e., greater year-to-year variation in the amount of rainfall was associated with lower to no reproductive seasonality. Finally, we identified diverse patterns of reproductive phenology, with the most seasonal breeders generally aligning lactation with the peak in resource availability while other populations show more diverse patterns, where conception, lactation or weaning can all be synchronized with maximal food availability. This study sheds new light on the extent and

ecological drivers of flexible reproductive phenology in long-lived tropical mammals, and

may even contribute to our understanding of why humans give birth year-round.

INTRODUCTION

Most animals face variation in their environment across the year (Boyce, 1979) in the form of

seasonal fluctuations in rainfall, temperature and resource availability that affect their energy

balance. Reproductive seasonality, the temporal clustering of reproductive events in the

annual cycle, is thought to be beneficial because it synchronizes the most energetically costly

reproductive stage with the seasonal food peak, thereby enhancing the condition and survival

probability of mothers and offspring (Bronson, 2009; Bronson & Heideman, 1994). Variation

in birth frequencies across the annual cycle is a continuous trait, ranging from a complete

absence of reproductive seasonality (i.e., random distribution of births throughout the year),

as in mountain gorillas (Campos et al., 2017), to cases in which all births occur within a few

weeks each year, as in many lemurs (Wright, 1999).

Comparative studies investigating determinants of variation in reproductive

seasonality across mammals have often been conducted at the level of the order (Rodents:

Heldstab, 2021, Lagomorphs: Heldstab, 2021, ruminants: Rutberg, 1987; Zerbe et al., 2012,

Carnivora: Heldstab et al., 2018, Primates: Di Bitetti & Janson, 2000; Heldstab et al., 2020;

Janson & Verdolin, 2005) and thus focus on broad-scale macro-evolutionary patterns. These

studies have typically detected a relationship between geographic latitude and birth

seasonality, suggesting that at higher latitudes, birth seasonality is more pronounced, with a

more intense birth peak (a birth peak being the temporal period in the annual cycle during

which most birth occur). However, important gaps remain in our understanding of the

determinants of reproductive seasonality. Few studies have attempted to quantify the extent of

variation in reproductive seasonality across multiple populations of the same species (but see

in African wild dogs (*Lycaon pictus*): McNutt, Groom, & Woodroffe, 2019, and in red-tailed

monkeys (*Cercopithecus ascanius*): Struhsaker, 1997), or across closely related species

sharing relatively similar diets, body sizes and life histories (but see in several ungulate

species: Pereira, Dos Santos Zanetti, & Furlan Polegato, 2010; Spinage, 1973; Brogi et al.,

2022; macaque species: Trébouet, Malaivijitnond, & Reichard, 2021). Thus, studies that

control for major sources of variation in life history or broad dietary categories should be

particularly useful for identifying the climatic drivers of variation in reproductive phenology.

The well-known association between latitude and reproductive seasonality fails to

explain the diversity of reproductive seasonality patterns observed within restricted latitudinal

ranges, such as in the tropics (Heldstab et al., 2020; Janson & Verdolin, 2005). In addition,

latitude encapsulates multiple components of climatic variation, which need to be

disentangled in order to identify the main climatic factors at play (Burtschell, Dezeure,

Huchard, & Godelle, 2023). First, latitude correlates positively with the degree of

environmental seasonality, measured as the magnitude of within-year variation (such as the

difference between maximal and minimal monthly rainfall in the annual cycle) (Botero, Dor,

McCain, & Safran, 2014). Further, latitude covaries negatively with environmental

productivity, i.e., overall food availability in a given environment. Variation in productivity

may alter the benefits of seasonal breeding, as populations living in more productive habitats

may face less pressure to breed seasonally (Burtschell et al., 2023).

Finally, environmental predictability, independently of latitude and seasonality

(Tonkin, Bogan, Bonada, Rios-Touma, & Lytle, 2017), could also influence breeding

schedules. In locations with intense year-to-year environmental variation, a flexible

reproductive phenology (i.e., individual ability to start a reproductive cycle at different

timings of the year, in response to internal or external factors) may be more advantageous

than a strictly seasonal reproduction (Brockman & van Schaik, 2005a; van Schaik & van

Noordwijk, 1985). Indeed, regular delays or decreases in the food peak may lead to

reproductive failures in strict seasonal breeders, thus reducing the fitness benefits of breeding

seasonally. However, few studies have investigated the effects of environmental

unpredictability on the intensity of reproductive seasonality, with mixed results so far. While

English, Chauvenet, Safi, & Pettorelli (2012) found that higher inter-annual variation in food

availability decreased the intensity of birth synchrony across 38 ungulate species, two studies

of red deer, *Cervus elaphus* L. (Loe *et al.*, 2005) and chacma baboons, *Papio ursinus*

(Dezeure *et al.*, 2023) found no effect of environmental unpredictability on reproductive

seasonality. A recent modelling study similarly detected limited effects of environmental

unpredictability on evolutionary transitions to nonseasonal breeding (Burtschell et al., 2023).

Aside from the selective pressures favouring a flexible reproductive phenology,

relatively little is known about how birth timing varies in relation to the annual resource peak

in long-lived species. In short-lived species, the full reproductive cycle, from conception to

offspring nutritional independence (such as weaning in mammals or fledging in birds), can

take place within a single productive season (Bronson, 2009). However, this is not the case

for long-lived species, in which multiple stages of a female's reproductive cycle can be

aligned with the annual food peak, with varying fitness consequences (Dezeure et al., 2021).

For example, females of some species may have to reach a certain threshold of body condition

for the onset of reproduction and conception to take place (Brockman & van Schaik, 2005a),

meaning that most conceptions are expected to follow a peak of food availability (Brockman

& van Schaik, 2005a). In some other species, females may instead synchronize the costliest

part of their reproductive cycle with the most productive season so as to enhance maternal

condition and survival (Bronson, 2009; Bronson & Heideman, 1994), such that early- or mid-

lactation occurs during the annual food peak, as in many primates (J. Altmann, 1980;

Brockman & van Schaik, 2005a; Janson & Verdolin, 2005). Lastly, weaning is a critically

vulnerable life stage, where juveniles must begin to forage for themselves (J. Altmann, 1980;

Lee, 1996). Accordingly, several species have been shown to time their births so as to align

weaning with the seasonal food peak (Brockman & van Schaik, 2005a; Janson & Verdolin,

2005), as occurs in most lemurs (Wright, 1999). Overall, the reasons underlying the observed

variation in alignment of reproductive stages with the food peak across species and

populations remain largely unknown.

In this study, we attempt to address the above gap in our understanding by

investigating the evolutionary determinants of the intensity and timing of reproductive

seasonality in Africa-dwelling papionin monkeys. We focus on papionins for several reasons.

First, they exhibit relatively similar body sizes (large-bodied), life history traits (slow) and

diet (mostly omnivorous) (Kingdon et al., 2012; Swedell, 2011), allowing us to investigate

environmental effects on reproductive seasonality while controlling for these – potentially

confounding – factors. Second, this taxonomic group displays a wide diversity of patterns of

reproductive seasonality. Indeed, most baboon (*Papio* spp.) species are non-seasonal breeders

(Bercovitch & Harding, 1993; Swedell, 2011) despite exhibiting variation in monthly birth

frequencies (Cheney et al., 2004; Lycett, Weingrill, & Henzi, 1999), while mandrills

(*Mandrillus sphinx*) (Setchell, Lee, Wickings, & Dixson, 2002), Kinda baboons (*Papio*

kindae) (Petersdorf, Weyher, Kamilar, Dubuc, & Higham, 2019) and most mangabey species

(i.e., *Cercocebus* and *Lophocebus* spp.) (Swedell, 2011) are seasonal breeders. Third, this

species constellation exhibits great ecological flexibility, inhabiting arid areas, woodland

savannahs, equatorial forests, and high altitude grasslands (J. Fischer et al., 2019; Kingdon et

al., 2012; Swedell, 2011) (see also Figure 1). Fourth, within the African members of this tribe,

baboons are one of the most well studied primate taxon, with data available from multiple

populations of some species within the genus *Papio* (J. Fischer et al., 2019). Lastly, species

from this taxonomic group, and in particular from the genus *Papio*, possess a variety of

features shared with early hominins (Alberts et al., 2005; Brockman, 2005; Jolly, 2001): they

are large, terrestrial and eclectic omnivorous primates (Alberts et al., 2005; Rhine, Norton,

Wynn, & Wynn, 1989) that, unlike the great apes, have colonized African savannahs (Bobe,

Martínez, & Carvalho, 2020) and give birth to a single offspring every one to three years (J.

Altmann & Alberts, 2005; Swedell, 2011). Investigating the environmental determinants of

their reproductive seasonality may thus shed new light on the evolution and maintenance of

non-seasonal breeding in early hominins (King, 2022).

Here we ask three main questions regarding reproductive seasonality in the papionins

in our sample:

(i) What is the extent of inter- and intra-specific variation in patterns of reproductive

seasonality, specifically regarding the height and width of the birth peak, as well as its timing

in the annual cycle?

(ii) What are the main environmental factors responsible for variation in the intensity of

reproductive seasonality? We isolated eight components of environmental variation: latitude,

environmental productivity, magnitude of seasonal variation in rainfall, number of rainy

seasons, breadth of the rainy season, amount of between-year (unpredictable) variation in

rainfall, between-year variation in the timing of the rainfall season, and the type of habitat.

We tested the eight corresponding hypotheses (H1.1-1.8) and their associated predictions,

which are listed in Table 1.

(iii) In seasonally breeding populations, which stage of the reproductive cycle is synchronized

with the food peak? We tested whether females match the seasonal food peak with

conceptions (H2.1 - the 'conception hypothesis'), lactation (H2.2 - the 'lactation hypothesis'),

or weaning (H2.3 - the 'weaning hypothesis').

Table 1: Hypotheses and predictions proposed on the effects of various environmental components on the intensity of reproductive seasonality

Predictor		Description	Hypothesis	Prediction	Results
Latitude		Latitude, in degrees	H1.1	When latitude increases, r_{birth} increases	No
Environmental productivity		Mean annual rainfall ($12 \times K_{rain}$[1])	H1.2	When environmental productivity increases, r_{birth} decreases	No (opposite effect)
Environmental seasonality	Magnitude of rainfall peak	Relative standard deviation of the seasonal component of rainfall (Rainfall S)	H1.3	When the amount of within-year rainfall variation increases, r_{birth} increases	No
	Modality of annual rainfall distribution	Number of rainy seasons (1 or 2)	H1.4	When there is more than one rainy season, r_{birth} decreases	No
	Breadth of the annual rainy season (broadness)	Rainfall peak breadth (RPB): minimum number of consecutive months with >80% of annual rainfall	H1.5	When RPB increases, r_{birth} decreases	No
Environmental unpredictability	Magnitude	Relative standard deviation of the non-seasonal component of rainfall (Rainfall NS)	H1.6	When the amount of rainfall unpredictability (between-year variation) increases, r_{birth} decreases	Yes
	Timing	Standard deviation of the yearly mean rainfall dates	H1.7	When the level of rainfall unpredictability in terms of timing increases, r_{birth} decreases	No
Habitat		Type of habitat: tropical forest, open savannah, or mosaic forest-savannah	H1.8	Populations living in open savannahs have lower r_{birth} than those living in tropical forests	No

173   r_{birth} refers here to the r-vector length, i.e. to the intensity of reproductive seasonality

174   [1]Krain   is   a   constant   and   refers   to   the   mean   monthly   rainfall

METHODS

1- Sample and data selection

Our data set includes published reports on reproductive seasonality in natural populations of Africa-dwelling papionins from the genera *Cercocebus*, *Lophocebus*, *Macaca, Mandrillus*, *Papio*, *Rungwecebus* and *Theropithecus* (Figure 1). We selected papers that reported the number of births per month (except for yellow baboons, *Papio cynocephalus*, from Mikumi National Park, where births were provided in three month-periods). We obtained birth seasonality data from 21 wild populations representing 11 species: see Table S1 for references associated with each population, and Figure 1 for their locations. We did not find any monthly birth data for nine species of interest for which only data from captivity were available (Kingdon et al., 2012; Swedell, 2011): five species of *Cercocebus* (*agilis*, *chrysogaster*, *galeritus*, *lunulatus*, *torquatus*), as well as *Lophocebus aterrimus*, *Mandrillus leucophaeus*, *Papio papio* and *Rungwecebus kipunji* (Figure 1).

Figure 1: Distribution of the sampled species in Africa, and locations of the populations considered in this study.

The species sampled in this study are depicted with relatively larger icons and species names, as well as full coloured circles. Species for which we could not find birth seasonality data are represented with smaller icons and names, as well as empty circles. The coloured areas on the map, corresponding to each species as indicated by the coloured circles, show the geographical distribution of each species. Within each range, small circled numbers show the location of the populations included in this study. 1: Taï, 2: Udzungwa Mountains, 3: Kibale, 4: Akfadou, 5: Tigounatine, 6: Lékédi. 7: Moukalaba-Doudou. 8: Gashaka-Gumti. 9: Gilgil. 10: Queen Elizabeth. 11: Amboseli. 12: Mikumi. 13: Tana River. 14: Filoha. 15: Kasanka. 16: De Hoop. 17: Drakensberg. 18: Moremi. 19: Tokai. 20: Tsaobis. 21: Simien. The species distribution ranges and icons come from Julia Fischer et al., 2017; Kingdon et al., 2012.

2- Birth seasonality data

We were interested in quantifying two components of reproductive seasonality in each

population: (1) the mean population birth date, i.e. describing when most births mainly occur

during the year, and (2) the intensity of population birth seasonality, i.e. describing how

seasonal the births are. For our analysis, given the heterogeneity of the data (in some datasets,

precise birth dates were available, but in most datasets, we could only obtain a count of births

per month), we considered each birth to have occurred in the middle of the month (i.e., the

15th of each month, except for February where births were considered to occur on the 14th).

For the Mikumi population, we considered that births occurred in the middle of each 3 month-

period. We then used a circular statistic and represented each birth event on the annual circle

by a vector of length 1 and of angle θ representing its date ($15 \times 2 \times \pi/365.25$ for January,

$(14+31) \times 2 \times \pi/365.25$ for February, etc.). We computed the mean vector (r-vector) per

population, whose angle (converted in a date: μ_{birth}) indicates the mean day of the year in

which births occur (see Table S1) and is thus a measure of birth seasonality. We computed

μ_{birth} using the function 'circ.summary' from the 'CircStats' package (Agostinelli & Lund,

2018). For populations with a significant birth peak, μ_{birth} represents the date of the population

birth peak, i.e. when births are the most likely to occur in the annual cycle. The length (r_{birth})

of the r-vector measures the intensity of birth seasonality, i.e., the degree of uniformity of the

birth distribution across the annual cycle, varying from 0 to 1 (Di Bitetti & Janson, 2000;

Janson & Verdolin, 2005; Thompson & McCabe, 2013). When $r_{birth}=0$, births are evenly

spread across months (i.e., non-seasonal), while when $r_{birth}=1$, births all occur during the same

month of the year (extremely seasonal). After comparing several classical measures of

reproductive seasonality (Supplementary Materials, Appendix S1), we used only r_{birth} to

measure the intensity of reproductive seasonality, as this measure is more robust to

differences in sample size than other metrics, facilitating the comparison of seasonality

measures between populations (Janson & Verdolin, 2005; Thel, Chamaillé-Jammes, &

Bonenfant, 2022).

3- Environmental data

i.   Two indicators of environmental variation: rainfall and NDVI

In order to test our set of hypotheses, we considered environmental variation through

two components: rainfall and the Normalized Difference Vegetation Index (NDVI). NDVI

produces a quantitative index of vegetation productivity, where higher values indicate a

higher degree of vegetation cover (Didan, Barreto Munoz, Solano, & Huete, 2015). Climatic

seasonality in Africa (and in most tropical habitats) is mainly characterized by within-year

variation in rainfall (Alberts et al., 2005; Feng, Porporato, & Rodriguez-Iturbe, 2013; Van

Schaik, Terborgh, & Wright, 1993), which has been successfully used as an indicator of food

availability for several of our studied populations (Alberts et al., 2005; Hill, Lycett, &

Dunbar, 2000; Petersdorf et al., 2019; Tinsley Johnson, Snyder-Mackler, Lu, Bergman, &

Beehner, 2018). Yet, NDVI values have to be used with caution when comparing productivity

across environments (Pettorelli et al., 2005), which is why we opted to use rainfall to test

hypotheses H1.1-H1-8. For example, the mean annual NDVI value at De Hoop, one of our

driest habitats, was almost equal to that of Gashaka, one of our wettest habitats. We thus used

variation in rainfall, rather than in NDVI, to disentangle the various components of climatic

variation that may affect the intensity of reproductive seasonality (such as environmental

productivity, predictability and seasonality) when testing hypotheses H1.1-H1.8.

However, we opted to use NDVI as an index of food availability to calculate the

timing of the food peak when testing hypotheses H2.1-H2-3, as Africa-dwelling papionins

rely mainly on plants for their diet (Swedell, 2011). Specifically, the annual peak in NDVI –

and presumably in food availability - is likely to lag behind that of rainfall by a few weeks,

and this lag duration may vary depending on local climatic or environmental conditions

(Bercovitch & Harding, 1993; Dezeure et al., 2021; Jarvey, Low, Pappano, Bergman, &

Beehner, 2018). Consequently, we thought that using NDVI would be preferable than rainfall

to assess the timing of the annual food peak, and test hypotheses regarding its match to the

birth peak in each population. Nevertheless, the use of NDVI as an index of food productivity

in papinions, which are not herbivorous (except for Theropithecus gelada), could be arguable,

and results of this analysis would be discussed accordingly.

ii.   Data extraction

Daily rainfall was extracted from satellite data sensors with the Giovanni NASA website

(product TRMM 3B42) (Huffman, Bolvin, Nelkin, & Adler, 2016) using a 0.25×0.25 degree

resolution (corresponding to between 28×28km at the equator and 23×23km at 35° latitude).

The GPS coordinates used for this extraction are indicated per population in Table S2, and

were assessed either from indications about the home ranges of the habituated groups per

population when available in the literature, or alternatively from the geographical location

(Park, Reserve or nearby city) of the population (see also Figure 1). Monthly cumulative

rainfall (summed across daily values) was subsequently computed between January 1998 and

December 2019. We therefore gathered 22 years of rainfall data per population over the same

period of time.

We then extracted the mean NDVI per 16 day-period on a 500m × 500m resolution

within the same geographical areas used for rainfall extraction (see GPS coordinates in Table

S2) between March 2000 and March 2017 (data before and after these dates were not

available at the time of data extraction) using MODIS data (MODIS13A1 product) provided

by NASA (Didan et al., 2015). Daily NDVI was computed by linear interpolation and then

averaged to obtain a monthly value across 18 years.

iii.   Components of environmental variation

In order to test our hypotheses, we identified multiple components of rainfall variation within

and across years. First, we decomposed for a given site the observed rainfall value into three

components as follows: $Rainfall_{m,i} = K_{rain} + Rainfall\ S_m + Rainfall\ NS_{m,i}$, where m is the

month of the year (going from January to December) and i is the year (from 1998 to 2019).

K_{rain} is a constant, equalling the mean monthly rainfall across 22 years of records (Figure S1).

Rainfall S_m is the seasonal component of rainfall, i.e., the rainfall value, averaged across 22

years, for each month of the year, minus K_{rain} (Figure S1). For example, for a given site,

Rainfall S_1 (m=1) is the mean of all January rainfall values. The term Rainfall S thus captures

the seasonal component of rainfall variation in the annual cycle, i.e., its within-year variation.

Finally, Rainfall NS $_{m,i}$ is the non-seasonal component of rainfall, i.e., the difference between

the observed rainfall value in any month at a given site ($Rainfall_{m,i}$) and the predictable

component of rainfall variation for that particular site in that particular month (K_{rain} + Rainfall

S_m) (Figure S1). This captures the unpredictable, i.e., between-year, rainfall variation. Using

these measures, we assessed the following for each population (see Table S2 for the values

associated with each population):

-    Environmental productivity, or mean annual rainfall, equal to $12 \times K_{rain}$.

-    Magnitude of environmental seasonality. We computed the magnitude of within-year

298   rainfall variation, as the relative standard deviation (SD) of Rainfall S standardized for

299   environmental productivity, given by the formula: $\frac{100 \times SD(Rainfall\ S)}{12 \times K_{rain}}$. The higher the

300   value, the more seasonal is rainfall variation.

301   -   The number of rainy seasons per year. Using predictable rainfall variation (K_{rain} +

302   Rainfall S), we assessed graphically, for each population, the number of rainy seasons

303   per year.

304   -   The length of the rainy season. For environments with only one rainy season, we

305   further calculated the rainfall peak breadth (RPB), which is the minimum number of

306   consecutive months of the year during which 80% of the annual rainfall ($12*K_{rain}$)

307   occurs. This measure is meaningless for environments with more than one rainy

308   season, and we thus excluded the populations living in such environments from this

309   analysis.

310   -   Magnitude of environmental unpredictability. We computed the magnitude of

311   between-year rainfall variation, as the relative standard deviation of Rainfall NS,

312   standardized by environmental productivity, given by the formula: $\frac{100 \times SD(Rainfall\ NS)}{12 \times K\mathrm{rain}}$.

313   The higher the value, the more unpredictable rainfall variation is.

Using the literature, we categorized the habitat of each population into three types:

tropical forest, open savannah, and mosaic forest-grassland (Table S2).

In addition to the magnitude of environmental unpredictability, we were interested in

quantifying unpredictability in the timing of the annual rainfall peak, i.e., quantifying how

much the timing of rainfall varied between years. Details on the procedure can be found in

Appendix S2, but briefly, using circular statistics, we computed standard deviations of the

mean dates of the annual rainfall peak over the 20 years sampled. Values close to zero mean

that rainfall peak occurs the same month every year, and the higher the value, the more

variation in rainfall peak's timing between years.

Lastly, we computed the mean monthly NDVI across 18 years (i.e., K_{NDVI}) and the

seasonal component of NDVI variation for each month of the year (i.e., NDVI S), following

the same notation used to disentangle the components of rainfall variation. To characterise the

timing of the seasonal food peak, we used circular statistics to compute the mean annual

NDVI date, μ_{NDVI} (see Appendix S2 for methodology and Table S2 for values of μ_{NDVI}).

iv – Phylogenetic tree

We used the branch length of Version 3 of the 10kTreesPrimates consensus tree (Arnold,

Matthews, & Nunn, 2010). Two species of interest were absent from this tree: *Cercocebus*

sanjei and *Papio kindae*. For the former, we substituted *Cercocebus galeritus*, its closest

relative. For the latter, following recent genetic studies (Jordan et al., 2018; Rogers et al.,

2019), we added *Papio kindae* in the same branch as *Papio ursinus*, using the function

'bind.tip' from the package 'phytools' (Revell, 2012).

4- Statistical analysis

All statistical analyses were conducted in R version 3.5.0 (R Core Team, 2019).

i.   *Factors affecting the intensity of reproductive seasonality*

We considered eight potential environmental parameters associated with reproductive

seasonality, which are listed along each corresponding prediction in Table 1.

For each hypothesis, we plotted r_{birth} (the r-vector length measuring the intensity of

reproductive seasonality in a given population) versus the tested predictor. We then checked

the significance of the relationship between each predictor and r_{birth} while controlling for

phylogeny using Bayesian phylogenetic generalized linear mixed models, with a Beta

regression and a logit link function, with the package 'brms' (Bürkner, 2017). We included

the phylogenetic relationship between species as a covariance matrix, which was derived from

the phylogenetic tree.

Given the low number of populations in this study, and the collinearity between some

of our environmental parameters (multivariate models had variance inflated factors >3), we

were not able to run stable multivariate models. For example, the magnitude of environmental

seasonality was negatively associated with the duration of the rainy season (cor=-0.94, t=-

9.15, $p<10^{-4}$), and environmental productivity was negatively associated with both latitude

(cor=-0.55, t=-2.69, p=0.016) and the magnitude of environmental unpredictability (cor=-

0.80, t=-5.48, $p<10^{-4}$).

Therefore, our Beta regressions included as a response variable r_{birth} values, one fixed

effect (each environmental predictor in turn listed in Table 1, standardised if continuous) and

the phylogenetic matrix as a random effect. Given the high variation in sample size (i.e.,

number of births recorded) between populations, we used a weighed regression, where the

weight given to each data point equals to log(N) / minimum(log(N)) so that the population

with the lowest sample size counts for 1 observation, and the other populations count for more

observations depending on their sample size, following a logarithmic scale ; the logarithmic

scale was chosen to account for the diminishing return of increasing the sample size of

samples that are already large. Beyond a given sample size, further increases in sample size

do not affect much r_{birth} estimates, which are already stable and precise. For each model, we

set an informative prior and used 3 000 iterations, a burn-in of 1 000 and 3 chains. We

visually inspected for convergence and checked the absence of autocorrelations for the

posterior distributions of fixed and random effects. The predictors were considered

statistically significant when their associated 95% confidence intervals did not cross 0.

Finally, we extracted the phylogenetic signal in our dataset with the metric of Blomberg's

K, allowing us to compare it with other signals from other traits. To do so, we computed the

mean r_{birth} per species, and use the 'phylosig' function with 1000 simulations from the

'phytools' package (Revell, 2024).

ii.   Timings of conceptions, births and weaning in relation with NDVI seasonality

We tested H2 only for those populations for which a significant birth peak can be detected, as

it does not make sense to test which period of the reproductive cycle is matched with the

annual food peak if there is not a clear seasonal pattern of births in one population. We

therefore assessed whether each population had a significant birth peak using the Rayleigh

test for circular statistics, more precisely the 'r.test' function from 'CircStats' package

(Agostinelli & Lund, 2018). For each population, when the P-value associated with the

Rayleigh test was <0.05, meaning that the null hypothesis of a uniform birth distribution

could be rejected, the birth peak was considered significant. With this approach, some

populations with relatively low reproductive seasonality (low r_{birth}) but with a large number of

births are included, which should be taken into account when interpreting the results. Among

these populations with a significant birth peak (see Table S1), we investigated which

reproductive stage (H2.1, 'conception'; H2.2, 'lactation'; or H2.3, 'weaning') was

synchronized with the annual NDVI peak, i.e. μ_{NDVI}. Additional details are given in

Supplementary Materials, Appendix 3 & Table S3. We employed exact two-sample Fisher-

Pitman permutation tests, using the 'oneway_test' function from the 'coin' package (Hothorn,

Hornik, Van De Wiel, & Zeileis, 2006). This function tests if the observed monthly value of

NDVI during a target period, which depends on the hypothesis tested, is significantly higher

than monthly values of NDVI randomized across the entire year. For example, using mean

gestation length to infer the annual distribution of conception dates, we tested H2.1 by asking

if females tended to conceive during, soon before, or soon after the annual food peak, looking

at seasonal NDVI values respectively in (i) the six months surrounding μ_{conc}, (ii) the three

months before μ_{conc} and (iii) the three months after μ_{conc}.

RESULTS

1) How variable are patterns of reproductive seasonality?

The annual distribution of births for each population is shown in Figure 2, alongside seasonal

variation in rainfall and NDVI. The intensity of reproductive seasonality varies widely across

species: *Papio hamadryas* (Filoha population: r_{birth} =0.02) and most *Papio anubis* populations

(r_{birth} <0.22) show non-seasonal births while *Mandrillus sphinx* (Lékédi: r_{birth} =0.67 and

Moukalaba-Doudou: r_{birth} =0.80 resp.) and *Papio kindae* (Kasanka: r_{birth} =0.50) exhibit

pronounced birth seasonality (Figure 3). The phylogenetic signal associated with the intensity

of reproductive seasonality is substantial (Blomberg's K=1.83, pvalue<0.01), indicating that

more closely related species have more similar patterns of reproductive seasonality. Among

papionins, the genera *Mandrillus*, *Cercocebus* and *Macaca* show strong reproductive

seasonality, whereas the genera *Lophocebus*, *Papio* and *Theropithecus* show an overall lower

intensity of birth seasonality, which may be associated with greater flexibility within species

(Figure 3). Such flexibility is particularly pronounced in *Papio ursinus* populations, extending

from low (de Hoop: r_{birth} =0.10; Tokai: r_{birth} =0.22, and Tsaobis: r_{birth} =0.10) to moderate birth

seasonality (Drakensberg: r_{birth} =0.42; Moremi: r_{birth} =0.37) (Figure 3), while intra-specific

variation seems less marked for the other species represented by multiple populations in our

sample.

The timing of birth seasonality can also be surprisingly variable, even between species

that live in adjacent geographical ranges: for instance, *Papio kindae* from Kasanka give birth

mainly around July, whereas *Papio ursinus* from Moremi give birth primarily between July

and November (Figure 2).

Figure 2: Monthly distribution of births in relation to rainfall and NDVI seasonality

The proportion of births per month (left side of the y-axis) is represented with red bars. In addition, the darker red bar indicates the month of the mean birth date (μ_{birth}) for seasonal breeding populations. We indicate in darker red the value of the r-vector length (r_{birth}) for each population (top-right corner of each panel). We represent the mean monthly rainfall (equalled to K_{rain} + Rainfall S, in mm) in blue (right side of the y-axis), and the mean monthly NDVI (equal to K_{NDVI} + NDVI S and divided by 2 for graphical purposes) in green (left-side of the y-axis). The species and population names are indicated on top of each panel, along with the number of births observed (N).

434
435 **Figure** 3: Phylogenetic tree of the studied species including the variation in
436 the intensity of reproductive seasonality for each species.
437 The intensity of birth seasonality is here quantified using the r_{birth} value. For those species
438 represented by more than one population, the length of the black segment displays the
439 variation in the intensity of birth seasonality among populations.
440
2) What are the ecological parameters correlated with the intensity of reproductive

seasonality?

We detected a significant correlation between the magnitude of environmental

unpredictability and the intensity of reproductive seasonality, while controlling for species

relatedness, which supported our prediction H1.6 (Table 2): the higher the magnitude of

between-year variation in rainfall, the lower the intensity of reproductive seasonality (Figure

4F). We also found an effect of habitat productivity that contradicted our prediction H1.2

(Table 2): the lower the mean annual rainfall, the lower the intensity of reproductive

seasonality (Figure 4B). Lastly, we did not find any support for the other six hypotheses: there

was no effect of latitude, magnitude of rainfall seasonality, number of rainy seasons, breadth

of the rainfall peak, unpredictability in the timing of the annual peak of rainfall, or habitat on

the intensity of reproductive seasonality (Table 2, Figure 4).

Figure 4: Effect of multiple environmental factors on the intensity of reproductive seasonality.

We plotted the intensity of reproductive seasonality (r_{birth}) depending on latitude (Panel A), environmental productivity (indexed by mean annual rainfall in mm, Panel B), the magnitude of environmental seasonality (i.e. of the rainfall peak, Panel C), the number of rainy seasons (Panel D), the rainfall peak breadth (RPB, Panel E), the magnitude of environmental unpredictability (i.e. of variation in the non-seasonal component of rainfall, Panel F), the timing of environmental unpredictability (i.e. between-year variation in rainfall timings, Panel G), and habitat type (Panel H). For each panel, each dot represents a population (with the population name annotated), and the colour indicates the species (see legend at the bottom). The dashed black line represents the linear regression, and the shaded area displays 95% confidence intervals. On top of each panel, we indicated in italic

and between parentheses the significance of each predictor (NS for non-significant, S for
significant).

Table 2: Influence of several components of rainfall variation on the reproductive seasonality of Africa-dwelling papionin
populations

The table shows the posterior mean, the estimate error and the 95% (marginal) confidence intervals (CI) associated for each posterior distribution of the predictors of the Beta

regression brms models including species' relatedness as random effect, r_{birth} as response variable, weighted by the log-transformed number of observations, and each

predictor as the only fixed effect of a univariate model. Significant effects are indicated in bold. For categorical predictor, the tested category is indicated between parentheses.

Predictor		posterior mean	estimate error	CI	
				lower	upper
Latitude		0.11	0.34	-0.56	0.77
Habitat productivity		**0.38**	**0.18**	**0.01**	**0.73**
Magnitude of environmental seasonality		0.03	0.13	-0.23	0.28
Number or rainy seasons	(2)[†]	-0.22	0.40	-0.98	0.58
Rainfall peak breadth		-0.06	0.17	-0.40	0.27
Magnitude of environmental unpredictability		**-0.44**	**0.13**	**-0.70**	**-0.20**
Timing of environmental unpredictability		-0.21	0.14	-0.52	0.05
Habitat type	(Mosaic forest-savannah)[š]	-0.21	0.41	-1.05	0.59
	(Tropical forest)[š]	0.56	1.45	-2.3	3.45

473　[†] The reference category is 1 rainy season
474　[š] The reference category is open savannah habitat
475

3) Which stage of the reproductive cycle is timed with the annual food peak?

When considering all taxa in our sample as a whole, none of our three hypotheses were

clearly supported (H2.1, H2.2, H2.3). However, in five of the six populations with the

strongest reproductive seasonality (r_{birth} >0.5), females appeared to synchronize lactation with

the annual NDVI peak, and overall lactation was generally aligned with the NDVI peak in 8

of the 14 populations with a significant birth peak (P-value of the Rayleigh test <0.05) (Table

3, Table S4). In less seasonally breeding populations (r_{birth} <0.5), females were more variable

in the reproductive stages that were timed with the annual NDVI peak, ranging from

conception (6 of 14 populations, and mainly before than after conception: e.g. Amboseli or

Kibale) to weaning (3 of 8 populations: e.g., Moremi or Gilgil) to none (e.g., Simien) or all of

these stages (e.g., Udzungwa) (Table 3, Table S4). Moreover, the timing of the annual NDVI

peak compared to the mean conception, birth or weaning dates was highly variable between

populations (Table S4).

Table 3: Reproductive stage matched with the food peak in sampled

populations.

Cells are filled with NAs when the alignment of the food peak with this reproductive stage

was not tested (either because the data on weaning age in this population was missing, or

because the birth peak of the population is non-significant, i.e., with P-value of the

Rayleigh test >0.05). Cells are filled with X when there is no alignment, and shaded when

there is an alignment of the given reproductive stage with the food peak.

Species	Population	r_{birth}	Reproductive stages match with food peak		
			Conception (H2.1)	Lactation (H2.2)	Weaning (H2.3)
Cercocebus atys	Taï	0.8312	X	**Early**	X
Cercocebus sanjei	Udzungwu Mountains	0.5588	**Before**	**Mid**	**Before**
Lophocebus albigena	Kibale	0.2762	**Before**	**Mid**	X

Species	Population				
Macaca sylvanus	Akfadou	0.9350	X	Early	NA
	Tigounatine	0.9467	X	X	NA
Mandrillus sphinx	Lekedi	0.6766	Before	Mid	X
	Moukalaba-Doudou	0.7940	X	Whole	NA
Papio anubis	Gashaka-Gumti	0.2167	NA	NA	NA
	Gilgil	0.1873	X	X	After
	Queen Elizabeth	0.0971	NA	NA	NA
Papio cynocephalus	Amboseli	0.1344	Before	X	X
	Mikumi	0.1584	X	X	NA
	Tana River	0.1413	NA	NA	NA
Papio hamadryas	Filoha	0.0208	NA	NA	NA
Papio kindae	Kasanka	0.4971	After	X	NA
Papio ursinus	De Hoop	0.0969	NA	NA	NA
	Drakensberg	0.4143	X	Early	NA
	Moremi	0.3710	Before	Mid	After
	Tokai	0.2394	NA	NA	NA
	Tsaobis	0.1046	NA	NA	NA
Theropithecus gelada	Simien	0.1550	X	X	X

DISCUSSION

We revealed strong inter- and intra-specific variation in the intensity of reproductive seasonality as well as in the annual timing of births in Africa-dwelling papionins. Our study further emphasizes the importance of environmental unpredictability for the evolution of flexible reproductive seasonality. Lastly, we found that females from different populations of Africa-dwelling papionins match different reproductive stages with the annual food peak.

Papionins exhibit flexible reproductive seasonality

Of the sampled *Papio* populations, most showed very little seasonality and *Papio*

kindae departed from the overall baboon pattern in being relatively highly seasonal breeders.

Such diverse patterns of reproductive seasonality within a single genus have rarely been

reported outside primates (see for example *Cervus*: English et al., 2012; Loe et al., 2005;

Rutberg, 1987, *Damaliscus*: Rutberg, 1987, *Ovis*: Rutberg, 1987, *Ursus*: Spady, Lindburg, &

Durrant, 2007, *Mustela*: Heldstab et al., 2018, *Vulpes*: Heldstab et al., 2018), but apparently

occur in some other primate genera with large distribution ranges, such as *Alouatta* (Di Bitetti

& Janson, 2000; Janson & Verdolin, 2005), *Cercopithecus* (Heldstab et al., 2020; Janson &

Verdolin, 2005), *Cebus* (Janson & Verdolin, 2005) and *Macaca* (Heldstab et al., 2020; Janson

& Verdolin, 2005; Trébouet et al., 2021). Given the limited taxonomic scale of our study, it is

impossible to establish whether seasonal breeding was the ancestral state in papionins, but it

seems possible that the loss of seasonal reproduction is a derived state affecting the *Papio*

genus (Fig 3). The estimated phylogenetic signal is significant and shows that among the 11

sampled papionin species, the intensity of birth seasonality is more similar among two closely

related species. The value of this phylogenetic signal (Blomberg's K=1.83) is relatively high,

among the highest of many life history traits (such as age at maturity, adult mortality, clutch

size in birds, sexual dimorphism, etc.), and higher than behavioural traits (such as daily

movement distance, prey size, preferred body temperature, etc.), that are more labile

(Blomberg, Garland, & Ives, 2003). Despite this strong impact of phylogeny on the intensity

of reproductive seasonality, our study emphasizes the importance of the variations in birth

seasonality between two closely-related species, or even within a single species.

Importantly, the key adaptive trait that evolved in the *Papio* genus may not be simply

the loss of breeding seasonality *per se*, but the evolution of a flexible reproductive phenology.

A same papionin female can give birth at different timings for successive birth events,

depending on her own individual traits or physiological constraints, or alternatively depending

on the strategies of other females in the same social group, as shown in *Papio ursinus*

(Dezeure et al., 2021, 2023). This reproductive flexibility at the individual level necessarily

shapes population patterns of reproductive seasonality, leading to lower reproductive

seasonality. Reproductive flexibility at the population level could be defined as the ability for

different populations of a same species to exhibit diverse patterns of reproductive seasonality,

depending on the environmental conditions. Such flexibility is observed at the population

level in Papio ursinus, living in a large distributional range characterized by exceptional

ecological diversity, which includes cold and temperate climates, oceanic and mountainous

ecosystems, and tropical and arid savannahs. Indeed, populations of this species exhibit a

wide range of intensity of reproductive seasonality (r_{birth}=[0.10-0.41]), with significant

(Moremi) or non-significant (Tsaobis) birth peaks, and to some extent, various timings in their

birth peaks (Moremi: around November, versus Drakensberg: around September). However,

such population-level flexibility is often difficult to assess in many species, given the datasets

available (*Papio ursinus* is indeed the only species in our sample represented by more than

three populations with a reasonable number of births).

Flexibility in reproductive phenology may be facilitated by several mechanisms. First,

a slower life history may allow papionin species to spread the energetic costs of reproduction

over a prolonged period, such that pregnant or lactating females face only a small daily extra

energetic expenditure that can be afforded at any time, as suggested by a recent modelling

study (Burtschell et al., 2023). In addition, unlike many other mammals, cercopithecids do not

use strict photoperiodic cues to trigger their reproduction (Heldstab et al., 2020) but may

instead exhibit condition-induced reproduction, whereby conceptions (and/or cycle

resumption after lactation) are more likely to occur when females are in better condition

(Alberts et al., 2005; Beehner, Onderdonk, Alberts, & Altmann, 2006). Such reproductive

flexibility may have contributed to their historical ecological success via their ability to

colonize diverse environments, and may become a critical asset to facilitate their resilience to

climate change, associated with increasing environmental unpredictability (Feng et al., 2013).

Environmental unpredictability may drive flexible reproductive seasonality

We examined several climatic correlates of the intensity of reproductive seasonality across

our sample, and found that environmental unpredictability was a significant predictor, with

higher between-year rainfall variation being associated with lower reproductive seasonality.

So far, most studies investigating climatic effects on reproductive phenology have focused on

environmental seasonality, i.e., the magnitude of within-year environmental variation. In

primates, the effect of environmental unpredictability on reproductive seasonality, e.g.,

through climatic events such as el Niño or fruit mast years in South-East Asia, has been

suggested (Brockman & van Schaik, 2005a; van Schaik & van Noordwijk, 1985; Wiederholt

& Post, 2011) but had never been tested. In line with our results, a previous study across 70

ungulate populations showed that the birth peak is more spread out in environments with

more year-to-year environmental variation (English et al., 2012).

Unpredictable climates may considerably reduce the fitness benefits associated with

seasonal breeding, such as enhancing maternal condition and offspring survival. In Africa,

year-to-year climatic variation frequently takes the shape of an absence of rain during the

rainy season (Alberts et al., 2005), which could cause severe reproductive costs in seasonal

breeders who often synchronize lactation or weaning with the rainy season, subsequently

forcing females to wait until the next breeding season to initiate a new reproductive event. In

such conditions, other adaptive traits may be more advantageous than seasonal breeding to

face the energetic costs of reproduction, such as the capacity to store energy (Brockman &

van Schaik, 2005a; van Schaik & van Noordwijk, 1985), to expand the dietary repertoire via a

generalist diet or foraging innovations (Grueter, 2017), to increase daily foraging time

(Alberts et al., 2005; Grueter, 2017), to flexibly adjust lactation duration (Dezeure et al.,

2021; van Noordwijk, 2012), or to reproduce cooperatively (Cornwallis et al., 2017; Lukas &

Clutton-Brock, 2017). Papionin species show many such traits: they can store energy, they

have an eclectic omnivorous diet, and they are flexible foragers that typically rely on fallback

foods during the dry season (J. Altmann, Schoeller, Altmann, Muruthi, & Sapolsky, 1993; S.

A. Altmann, 2009; Swedell, 2011). As such, it is likely that rainfall unpredictability selected

these traits in papionin species (energy storage, omnivorous diet, slow life histories, etc.),

which in turn contributed to shape their flexible reproductive seasonality.

Two recent studies found no or little effect of climatic unpredictability on the intensity

of reproductive seasonality at the population level, one in *Papio ursinus* (Dezeure et al., 2023)

and one on *Papio cynocephalus* using a modelling approach (Burtschell et al., 2023),

questioning the robustness of the effect found in this study. This discrepancy may come from

the fact that unpredictable climates select for reproductive flexibility, rather than nonseasonal

breeding. In fact, the study by Burtschell et al., 2023 revealed that increasing climatic

unpredictability was associated with a lower variance, but not a lower mean, in the intensity

of reproductive seasonality. In addition, the effect of climatic unpredictability may be better

detected across space than time, i.e., by comparing different populations living in distinct

climates and environments, as is the case here, than by comparing the same population across

time, as was the case for Dezeure et al., 2023. Different study designs should be combined,

across time, taxonomic or spatial scales, to reveal the full complexity of selective pressures at

play.

The numerous pressures affecting the intensity of reproductive seasonality are often

hard to disentangle, meaning that the effects uncovered by correlational studies like ours may

sometimes reflect other co-varying pressures, and should thus be interpreted cautiously.

Specifically, environmental productivity, which is negatively correlated with climatic

unpredictability at our study sites (cor=-0.80, t=-5.48, $p<10^{-4}$), was also a significant predictor

of the intensity of reproductive seasonality, but in a direction opposed to our prediction, as

well as to previous results (Burtschell et al., 2023). In our sample, the least productive

climates are also the most unpredictable, and environmental productivity may thus represent a

confounding factor in the relationship between climatic unpredictability and seasonal

breeding. In species with a flexible phenology, females can start a new reproductive event

rapidly after a reproductive failure, without having to wait the next mating season. Such

failures are likely to be particularly frequent where environmental productivity is low,

contributing to spread reproductive events across the year cycle, and explaining how low

environmental productivity may contribute to decrease reproductive seasonality. In addition,

this study is based on datasets with heterogenous resolutions, including diverse numbers of

births and years of study (e.g. Amboseli: N=496, Nyears=33, versus Queen Elizabeth: N=35,

Nyears=2). Additional birth records in small datasets may change r_{birth}, and could thus alter

some of the results in our study.

Lastly, our decomposition of rainfall components further calls for a more rigorous

definition of the term seasonality, especially when it is used in a quantitative way. Indeed, a

'more seasonal' environment can either be an environment with higher within-year variation

(i.e. the amplitude of variation between the 'best' and the 'worst' month of the year), with

higher within-year over between-year variation (i.e. the amplitude of within-year variation

controlling for the intensity of unpredictable variations), with a shorter productive season (i.e.

the rainfall peak breadth), or with only a unimodal season (i.e. one rainy, or one warm season

per year). Similarly, climatic unpredictability can be broken into two components: (1) the

amount/magnitude of year-to-year variation (i.e., if the rainy season brings more or less

rainfall than usual), and (2) year-to-year variation in the timing of the rainy season (i.e., if the

rainy season occurs earlier or later than usual) (Clauss, Zerbe, Bingaman Lackey, Codron, &

Müller, 2020). These various components have rarely been disentangled in empirical studies

so far, and this study opens new methodological avenues to investigate various environmental

components that are likely to affect reproductive seasonality.

Females can match different reproductive stages with the food peak

The main pattern emerging from our investigations suggests that females from species with

high breeding seasonality match lactation with the peak in vegetation productivity (Table 3).

For species and populations with a more flexible reproductive phenology, females can match

different reproductive stages with the annual vegetation peak, with a possible preference

towards conception. This trend may reflect the condition-dependence of conception – a

proximate mechanism – rather than an adaptive, optimal strategy aimed at synchronizing the

vegetation peak with a particular reproductive stage. These results, obtained from (mostly)

tropical primates, echo the broader mammalian literature showing that most mammals from

temperate regions match lactation with the best season of the year (Bronson, 2009; Bronson &

Heideman, 1994), while patterns are more variable in tropical and long-lived mammals,

depending on body size, energy storage capacities and environmental predictability

(Brockman & van Schaik, 2005a; Janson & Verdolin, 2005; van Schaik & van Noordwijk,

1985). In addition, even though weaning is a vulnerable life-history stage in young primates,

which can be buffered when matched with the vegetation peak in a wild *Papio ursinus*

population (Dezeure et al., 2021), few populations seemed to adopt this strategy.

Several caveats apply to the test of H2. First, although NDVI is a relatively good

measure of plant productivity, highest values of NDVI do not necessarily coincide with the

annual food peak, especially when focusing on omnivorous/frugivorous species. Precise

phenological data from each population would be more accurate to quantify the annual food

peak. Second, our estimations of lactation peak and weaning might lack accuracy, due to

strong between-populations and between-individuals variation. Data quantifying maternal

energy expenditure during lactation (Rosetta, Lee, & Garcia, 2011), or isotopic measures of

trophic levels between mothers and infants (Carboni, Dezeure, Cowlishaw, Huchard, &

Marshall, 2022; Reitsema, 2012) would be necessary, for each population, to determine the

dynamics of lactation and weaning. Finally, additional unexplored factors can potentially

affect reproductive timing and further limit our ability to detect a clear pattern. For example,

for populations living at high altitudes like *Theropithecus gelada* from Simien and the *Papio*

ursinus from Drakensberg, seasonal variation in temperatures also constrain reproductive

phenology (Lycett et al., 1999; Tinsley Johnson et al., 2018).

The evolution of reproductive flexibility in Anthropoid primates may inform our

understanding of the reproductive phenology in early humans

Baboons and relatives represent an interesting model for understanding the evolution of

behavioural and reproductive plasticity of early humans (J. Fischer et al., 2019; King, 2022).

Although most great apes are nonseasonal breeders (Brockman & van Schaik, 2005b; Campos

et al., 2017), suggesting that their common ancestor also bred year-round, humans are distinct

from other apes by exhibiting much faster reproductive paces (which are similar to most

Africa-dwelling papionins: Swedell, 2011), and by living in a wider variety of environments,

rather than being restricted to tropical forests (Wells & Stock, 2007). The selective pressures

that have shaped reproductive phenology in the human lineage versus in other apes may

therefore differ, and the papionins, who have similarly left forested habitats to colonize

savannahs may provide valuable insights to understand the adaptation of early humans to such

diverse and unpredictable environments. Our results suggest that baboons have acquired a low

and flexible reproductive seasonality, as well as several other adaptive traits, when facing

more arid and unpredictable environments, such as a generalist omnivorous diet and the

frequent use of fallback foods, frequent foraging innovations, an increased ability to store fat

and to switch home ranges to more suitable areas. Similar reproductive, physiological and

behavioural adaptations to environmental unpredictability may have allowed early humans to

thrive and maintain fast reproductive paces during their colonization of a wide variety of

environments.

Conclusion

Our work revealed substantial variation in patterns of reproductive seasonality within and

across species of Africa-dwelling papionins, highlighting an exceptional flexibility in their

reproductive phenology. Among multiple dimensions of climatic variation, rainfall

unpredictability and productivity were the main predictors of the intensity of reproductive

seasonality, with arid and unpredictable climates being associated with less seasonal

reproduction. Among populations with a pronounced breeding seasonality, females often

match lactation with the annual vegetation peak, while phenology patterns are very diverse in

other populations. This study sheds new light on the selective pressures shaping reproductive

seasonality in long-lived tropical mammals, as well as on potential adaptations to

environmental unpredictability. It may further provide an original contribution to understand

why humans breed year-round, given their phylogenetic ties and convergences in life-history

and ecology with our taxonomic group.

ACKNOWLEDGMENTS

We thank Cassandra Raby for her kind help on extracting NDVI data, and Christophe

Dezeure for making the Figure 1 of this paper (and more generally for his paternal care). This

project was funded by a grant from the Agence Nationale de la Recherche (ANR ERS-17-

CE02-0008, 2018-2021) awarded to E. H.

DATA AVAILABILITY

The raw data and scripts used for this paper are available in the following Zenodo repository:

10.5281/zenodo.13312416.

REFERENCES

Agostinelli, C., & Lund, U. (2018). *Package "CircStats": Circular Statistics, from "Topics in Circular Statistics."*

Alberts, S. C., Hollister-smith, J. A., Mututua, R. S., Sayialel, S. N., Muruthi, P. M., Warutere, K. J., & Altmann, J. (2005). Seasonality and long-term change in a savanna environment. In D. K. Brockman & C. P. van Schaik (Eds.), *Seasonality in Primates: Studies of living and extinct human and non-human primates* (pp. 157–195). Cambridge: Cambridge Univ. Press.

Altmann, J. (1980). *Baboon Mothers and Infants*. The University of Chicago Press.

Altmann, J., & Alberts, S. C. (2005). Growth rates in a wild primate population: ecological influences and maternal effects. *Behavioral Ecology and Sociobiology*, *57*, 490–501. doi: 10.1007/s00265-004-0870-x

Altmann, J., Schoeller, D., Altmann, S. A., Muruthi, P., & Sapolsky, R. M. (1993). Body size and fatness of free‐living baboons reflect food availability and activity levels. *American Journal of Primatology*. doi: 10.1002/ajp.1350300207

Altmann, S. A. (2009). Fallback foods, eclectic omnivores, and the packaging problem. *American Journal of Physical Anthropology*, *140*(4), 615–629. doi: 10.1002/ajpa.21097

Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: A new online resource for primate phylogeny. *Evolutionary Anthropology*, *19*(3), 114–118. doi: 10.1002/evan.20251

Beehner, J. C., Onderdonk, D. A., Alberts, S. C., & Altmann, J. (2006). The ecology of conception and

pregnancy failure in wild baboons. *Behavioral Ecology*, *17*(5), 741–750. doi:

10.1093/beheco/arl006

Bercovitch, F. B., & Harding, R. S. O. (1993). Annual Birth Patterns of Savanna Baboons (Papio

cynocephalus anubis) over a Ten-Year Period at Gilgil, Kenya. *Folia Primatologica*, *61*, 115–

122.

Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for phylogenetic signal in comparative

data: Behavioral traits are more labile. *Evolution*, *57*(4), 717–745. doi: 10.1111/j.0014-

3820.2003.tb00285.x

Bobe, R., Martínez, F. I., & Carvalho, S. (2020). Primate adaptations and evolution in the Southern

African Rift Valley. *Evolutionary Anthropology*, *29*(3), 94–101. doi: 10.1002/evan.21826

Botero, C. A., Dor, R., McCain, C. M., & Safran, R. J. (2014). Environmental harshness is positively

correlated with intraspecific divergence in mammals and birds. *Molecular Ecology*, *23*(2), 259–

268. doi: 10.1111/mec.12572

Boyce, M. S. (1979). Seasonality and Patterns of Natural Selection for Life Histories. *The American*

Naturalist, *114*(4), 569–583. doi: 10.1086/283503

Brockman, D. K. (2005). What do studies of seasonality in primates tell us about human evolution? In

Seasonality in Primates: Studies of Living and Extinct Human and Non-Human Primates (pp.

543–570). Cambridge: Cambridge University Press. Retrieved from

https://www.researchgate.net/publication/230778889

Brockman, D. K., & van Schaik, C. (2005a). Seasonality and reproductive function. In D. K.

Brockman & C. P. van Schaik (Eds.), *Seasonality in Primates: Studies of Living and Extinct*

Human and Non-Human Primates (pp. 269–305). Cambridge University Press.

Brockman, D. K., & van Schaik, C. P. (2005b). *Seasonality in Primates: Studies of Living and Extinct*

Human and Non-Human Primates (D. K. Brockman & C. P. van Schaik, Eds.). Cambridge:

Cambridge University Press.

Brogi, R., Merli, E., Grignolio, S., Chirichella, R., Bottero, E., & Apollonio, M. (2022). It is time to

mate: population-level plasticity of wild boar reproductive timing and synchrony in a changing

environment. *Current Zoology*, *68*(4), 371–380. doi: 10.1093/cz/zoab077

Bronson, F. H. (2009). Climate change and seasonal reproduction in mammals. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *364*(1534), 3331–3340. doi:

10.1098/rstb.2009.0140

Bronson, F. H., & Heideman, P. D. (1994). Seasonal regulation of reproduction in mammals. In E.

Knobil & J. D. Neill (Eds.), *The physiology of reproduction* (Vol. 2, pp. 541–583). New York:

New York Raven Press. Retrieved from https://www.researchgate.net/publication/313495409

Bürkner, P. C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of*

Statistical Software, *80*. doi: 10.18637/jss.v080.i01

Burtschell, L., Dezeure, J., Huchard, E., & Godelle, B. (2023). Evolutionary determinants of

reproductive seasonality: A theoretical approach. *Peer Community Journal*, *3*(e56). doi:

10.24072/pci

Campos, F. A., Morris, W. F., Alberts, S. C., Altmann, J., Brockman, D. K., Cords, M., … Fedigan, L.

M. (2017). Does climate variability influence the demography of wild primates? Evidence from

long-term life-history data in seven species. *Global Change Biology*, *23*(3), 1–15. doi:

10.1111/gcb.13754

Carboni, S., Dezeure, J., Cowlishaw, G., Huchard, E., & Marshall, H. H. (2022). Stable isotopes reveal

the effects of maternal rank and infant age on weaning dynamics in wild chacma baboons.

Animal Behaviour, *193*, 21–32. doi: 10.1016/j.anbehav.2022.08.010ï

Cheney, D. L., Seyfarth, R. M., Fischer, J., Beehner, J., Bergman, T., Johnson, S. E., … Silk, J. B.

(2004). Factors affecting reproduction and mortality among baboons in the Okavango Delta,

Botswana. *International Journal of Primatology*, *25*(2), 401–428. doi: 0164-0291/04/0400-

0401/0

Clauss, M., Zerbe, P., Bingaman Lackey, L., Codron, D., & Müller, D. W. H. (2020). Basic

considerations on seasonal breeding in mammals including their testing by comparing natural

habitats and zoos. *Mammalian Biology*. doi: 10.1007/s42991-020-00078-y

Cornwallis, C. K., Botero, C. A., Rubenstein, D. R., Downing, P. A., West, S. A., & Griffin, A. S.

(2017). Cooperation facilitates the colonization of harsh environments. *Nature Ecology and*

Evolution, *1*(3). doi: 10.1038/s41559-016-0057

Dezeure, J., Baniel, A., Carter, A., Cowlishaw, G., Godelle, B., & Huchard, E. (2021). Birth timing

generates reproductive trade-offs in a non-seasonal breeding primate. *Proceedings of the Royal*

Society B, *288*, 1–9. doi: 10.1098/rspb.2021.0286

Dezeure, J., Burtschell, L., Baniel, A., Carter, A. J., Godelle, B., Cowlishaw, G., & Huchard, E.

(2023). Evolutionary Determinants of Nonseasonal Breeding in Wild Chacma Baboons.

American Naturalist, *201*(1), 106–124. doi: 10.1086/722082

Di Bitetti, M. S., & Janson, C. H. (2000). When will the stork arrive? Patterns of birth seasonality in

neotropical primates. *American Journal of Primatology*, *50*(2), 109–130. doi:

10.1002/(SICI)1098-2345(200002)50:2<109::AID-AJP2>3.0.CO;2-W

Didan, K., Barreto Munoz, A., Solano, R., & Huete, A. (2015). *MOD13A1 MODIS/Terra Vegetation*

Indices 16-Day L3 Global 500m SIN Grid V006 [Data set]. doi:

https://doi.org/10.5067/MODIS/MOD13A1.006

English, A. K., Chauvenet, L., Safi, K., & Pettorelli, N. (2012). Reassessing the Determinants of

Breeding Synchrony in Ungulates. *PLoS ONE*, *7*(7), 1–7. doi: 10.1371/journal.pone.0041444

Feng, X., Porporato, A., & Rodriguez-Iturbe, I. (2013). Changes in rainfall seasonality in the tropics.

Nature Climate Change, *3*(9), 811–815. doi: 10.1038/nclimate1907

Fischer, J., Higham, J. P., Alberts, S. C., Barrett, L., Beehner, J. C., Bergman, T. J., … Zinner, D.

(2019). The Natural History of Model Organisms: Insights into the evolution of social systems

and species from baboon studies. *ELife*, *1960*, 1–16.

Fischer, Julia, Kopp, G. H., Dal Pesco, F., Goffe, A., Hammerschmidt, K., Kalbitzer, U., … Zinner, D.

(2017). Charting the neglected West: The social system of Guinea baboons. *American Journal of*

Physical Anthropology, *162*, 15–31. doi: 10.1002/ajpa.23144

Grueter, C. C. (2017). Environmental Seasonality. In *The International Encyclopedia of Primatology*

(pp. 1–2). John Wiley & Sons, Inc. doi: 10.1002/9781119179313.wbprim0048

Heldstab, S. A. (2021a). Latitude, life history and sexual size dimorphism correlate with reproductive

seasonality in rodents. *Mammal Review*, *51*, 256–271.

Heldstab, S. A. (2021b, December 1). Habitat characteristics and life history explain reproductive

seasonality in lagomorphs. *Mammalian Biology*, Vol. 101, pp. 739–757. Springer Science and

Business Media Deutschland GmbH. doi: 10.1007/s42991-021-00127-0

Heldstab, S. A., Müller, D. W. H., Graber, S. M., Bingaman Lackey, L., Rensch, E., Hatt, J. M., …

Clauss, M. (2018). Geographical Origin, Delayed Implantation, and Induced Ovulation Explain

Reproductive Seasonality in the Carnivora. *Journal of Biological Rhythms*, *33*(4), 402–419. doi:

10.1177/0748730418773620

Heldstab, S. A., van Schaik, C. P., Müller, D. W. H., Rensch, E., Lackey, L. B., Zerbe, P., … Matsuda,

I. (2020). Reproductive seasonality in primates: patterns, concepts and unsolved questions.

Biological Reviews. doi: 10.1111/brv.12646

Hill, R. A., Lycett, J. E., & Dunbar, R. I. M. (2000). Ecological and social determinants of birth

intervals in baboons. *Behavioral Ecology*, *11*(5), 560–564. doi: 10.1093/beheco/11.5.560

Hothorn, T., Hornik, K., Van De Wiel, M. A., & Zeileis, A. (2006). A lego system for conditional

inference. *American Statistician*, *60*(3), 257–263. doi: 10.1198/000313006X118430

Huffman, G. J., Bolvin, D. T., Nelkin, E. J., & Adler, R. F. (2016). *TRMM (TMPA) Precipitation L3 1*

day 0.25 degree x 0.25 degree V7. Retrieved from 10.5067/TRMM/TMPA/DAY/7

Janson, C., & Verdolin, J. (2005). Seasonality of primate births in relation to climate. In D. K.

Brockman & C. P. van Schaik (Eds.), *Seasonality in Primates: Studies of Living and Extinct*

Human and Non-Human Primates (pp. 307–350). Cambridge: Cambridge University Press. doi:

10.1017/cbo9780511542343.012

Jarvey, J. C., Low, B. S., Pappano, D. J., Bergman, T. J., & Beehner, J. C. (2018). Graminivory and

Fallback Foods: Annual Diet Profile of Geladas (Theropithecus gelada) Living in the Simien

Mountains National Park, Ethiopia. *International Journal of Primatology*, *39*(1), 105–126. doi:

10.1007/s10764-018-0018-x

Jolly, C. J. (2001). A proper study for mankind: Analogies from the Papionin monkeys and their

implications for human evolution. *Yearbook of Physical Anthropology*, *44*, 177–204. doi:

10.1002/ajpa.10021

Jordan, V. E., Walker, J. A., Beckstrom, T. O., Steely, C. J., McDaniel, C. L., St Romain, C. P., …

Worley, K. C. (2018). A computational reconstruction of Papio phylogeny using Alu insertion

polymorphisms. *Mobile DNA*, *9*(1). doi: 10.1186/s13100-018-0118-3

King, G. E. (2022). Baboon perspectives on the ecology and behavior of early human ancestors.

Proceedings of the National Academy of Sciences of the United States of America, *119*(45). doi:

10.1073/pnas.2116182119

Kingdon, J., Happold, D., Butynski, T. M., Hoffmann, M., Happold, M., & Kalina, J. (2012).

Mammals of Africa.

Lee, P. C. (1996). The meanings of weaning: growth, lactation, and life history. *Evolutionary*

Anthropology, *5*(3), 87–98. doi: https://doi.org/10.1002/(SICI)1520-6505(1996)5:3<87::AID-

EVAN4>3.0.CO;2-T

Loe, L. E., Bonenfant, C., Mysterud, A., Gaillard, J. M., Langvatn, R., Klein, F., … Stenseth, N. C.

(2005). Climate predictability and breeding phenology in red deer: Timing and synchrony of

rutting and calving in Norway and France. *Journal of Animal Ecology*, *74*(4), 579–588. doi:

10.1111/j.1365-2656.2005.00987.x

Lukas, D., & Clutton-Brock, T. (2017). Climate and the distribution of cooperative breeding in

mammals. *Royal Society Open Science*, *4*(1). doi: 10.1098/rsos.160897

Lycett, J. E., Weingrill, T., & Henzi, S. P. (1999). Birth patterns in the Drakensberg Mountain

857   baboons (Papio cynocephalus ursinus). *South African Journal of Science*, *95*(8), 354–356.

858   McNutt, J. W., Groom, R., & Woodroffe, R. (2019). Ambient temperature provides an adaptive

859   explanation for seasonal reproduction in a tropical mammal. *Journal of Zoology*, *309*(3), 153–

860   160. doi: 10.1111/jzo.12712

861   Pereira, R. J. P., Dos Santos Zanetti, E., & Furlan Polegato, B. (2010). Female reproduction. In J. M.

862   Barbanti Duarte & S. Gonzalez (Eds.), *Neotropical Cervidology* (1st ed., Vol. 1).

863   FUNEP/IUCN. Retrieved from https://www.researchgate.net/publication/273143129

864   Petersdorf, M., Weyher, A. H., Kamilar, J. M., Dubuc, C., & Higham, J. P. (2019). Sexual selection in

865   the Kinda baboon. *Journal of Human Evolution*, *135*(102635). doi: 10.1016/j.jhevol.2019.06.006

866   Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J. M., Tucker, C. J., & Stenseth, N. C. (2005). Using

867   the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in

868   Ecology and Evolution*. doi: 10.1016/j.tree.2005.05.011

869   R Core Team. (2019). *R: A language and environment for statistical computing*. Retrieved from

870   https://www.r-project.org/

871   Reitsema, L. J. (2012). Introducing Fecal Stable Isotope Analysis in Primate Weaning Studies.

872   *American Journal of Primatology*, *74*(10), 926–939. doi: 10.1002/ajp.22045

873   Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things).

874   *Methods in Ecology and Evolution*, *3*, 217–223. doi: 10.1111/j.2041-210X.2011.00169.x

875   Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and

876   other things). *PeerJ*, *12*. doi: 10.7717/peerj.16505

877   Rhine, R. J., Norton, G. W., Wynn, G. M., & Wynn, R. D. (1989). Plant feeding of yellow baboons

878   (Papio cynocephalus) in Mikumi national park, Tanzania, and the relationship between seasonal

879   feeding and immature survival. *International Journal of Primatology*, *10*(4), 319–342. doi:

10.1007/BF02737420

Rogers, J., Muthuswamy, R., Harris, R. A., Mailund, T., Leppälä, K., Athanasiadis, G., … Worley, K.

C. (2019). The comparative genomics and complex population history of Papio baboons. *Science*

Advances, *5*, 1–14. Retrieved from http://advances.sciencemag.org/

Rosetta, L., Lee, P. C., & Garcia, C. (2011). Energetics during reproduction: A doubly labeled water

study of lactating baboons. *American Journal of Physical Anthropology*, *144*(4), 661–668. doi:

10.1002/ajpa.21475

Rutberg, A. T. (1987). Adaptive Hypotheses of Birth Synchrony in Ruminants: An Interspecific Test.

The American Naturalist, *130*(5), 692–710. doi: 10.1086/284739

Setchell, J. M., Lee, P. C., Wickings, E. J., & Dixson, A. F. (2002). Reproductive Parameters and

Maternal Investment in Mandrills (Mandrillus sphinx). *International Journal of Primatology*,

23(1), 51–68. doi: 10.1023/A:1013245707228

Spady, T. J., Lindburg, D. G., & Durrant, B. S. (2007). Evolution of reproductive seasonality in bears.

Mammal Review, *37*(1), 21–53. doi: 10.1111/j.1365-2907.2007.00096.x

Spinage, C. A. (1973). The role of photoperiodism in the seasonal breeding of tropical African

ungulates. *Mammal Review* , *3*(3), 71–85. doi: 10.1111/j.1365-2907.1973.tb00174.x

Struhsaker, T. T. (1997). Ecology of an African rain forest: logging in Kibale and the conflict between

conservation and exploitation. *Journal of Tropical Ecology*, *14*(4), 561–561. doi:

10.1017/S0266467498240399

Swedell, L. (2011). African papionins: diversity of social organization and ecological flexibility. In C.

J. Campbell, A. Fuentes, K. C. MacKinnon, S. Bearder, & R. Stumpf (Eds.), *Primates in*

perspective (Vol. 2, pp. 241–277). Oxford: Oxford University Press. Retrieved from

https://www.researchgate.net/publication/285789215

Thel, L., Chamaillé-Jammes, S., & Bonenfant, C. (2022). How to describe and measure phenology?

An investigation on the diversity of metrics using phenology of births in large herbivores. *Oikos*,

2022(4). doi: 10.1111/oik.08917

Thompson, M. E., & McCabe, G. M. (2013). Reproductive seasonality in wild Sanje mangabeys

(Cercocebus sanjei), Tanzania: Relationship between the capital breeding strategy and infant

survival. *Behaviour*, *150*(12), 1399–1429. doi: 10.1163/1568539X-00003102

Tinsley Johnson, E., Snyder-Mackler, N., Lu, A., Bergman, T. J., & Beehner, J. C. (2018). Social and

ecological drivers of reproductive seasonality in geladas. *Behavioral Ecology*, *29*(3), 574–588.

doi: 10.1093/beheco/ary008

Tonkin, J. D., Bogan, M. T., Bonada, N., Rios-Touma, B., & Lytle, D. A. (2017). Seasonality and

predictability shape temporal species diversity. *Ecology*, *98*(5), 1201–1216.

Trébouet, F., Malaivijitnond, S., & Reichard, U. H. (2021). Reproductive seasonality in wild northern

pig-tailed macaques (Macaca leonina). *Primates*. doi: 10.1007/s10329-021-00901-1

van Noordwijk, M. A. (2012). From maternal investment to lifetime maternal care. In *The evolution of*

primate societies (pp. 321–342). Chicago, London: The University of Chicago Press.

Van Schaik, C. P., Terborgh, J. W., & Wright, S. J. (1993). The phenology of tropical forests:

Adaptive significance and consequences for primary consumers. *Annual Review of Ecology and*

Systematics, *24*(1), 353–377. doi: 10.1146/annurev.es.24.110193.002033

van Schaik, C. P., & van Noordwijk, M. A. (1985). Interannual variability in fruit abundance and the

reproductive seasonality in Sumatran Long-tailed macaques (Macaca fascicularis). *J. Zool.,*

Lond. (A), *206*(4), 533–549. doi: 10.1111/j.1469-7998.1985.tb03557.x

Wells, J. C. K., & Stock, J. T. (2007). The biology of the colonizing ape. *Yearbook of Physical*

Anthropology, *50*(1), 191–222. doi: 10.1002/ajpa.20735

Wiederholt, R., & Post, E. (2011). Birth seasonality and offspring production in threatened neotropical

primates related to climate. *Global Change Biology*, *17*(10), 3035–3045. doi: 10.1111/j.1365-

2486.2011.02427.x

Wright, P. C. (1999). Lemur Traits and Madagascar Ecology: Coping With an Island Environment.

Yearbook of Physical Anthropology, *42*, 31–72. doi: 10.1002/(sici)1096-

8644(1999)110:29+<31::aid-ajpa3>3.0.co;2-0

Zerbe, P., Clauss, M., Codron, D., Bingaman Lackey, L., Rensch, E., Streich, J. W., … Müller, D. W.

H. (2012). Reproductive seasonality in captive wild ruminants: Implications for biogeographical

adaptation, photoperiodic control, and life history. *Biological Reviews*, *87*(4), 965–990. doi:

10.1111/j.1469-185X.2012.00238.x