# The climatic determinants of flexible reproductive seasonality in Africa-dwelling papionins

Jules Dezeure[1], Julie Dagorrette[2], Lugdiwine Burtschell[2,3], Shahrina Chowdhury[4,7,8], Dieter Lukas[5], Larissa Swedell[6,7,8,9,10], Elise Huchard[2]

[1]Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France

[2]Institute of Evolutionary Sciences of Montpellier (ISEM), Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France.

[3]AgroParisTech, Paris, France

[4]Department of Anthropology, Brooklyn College, CUNY, 2900 Bedford Ave, Brooklyn, NY 112

[5]Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

[6]Department of Anthropology, Queens College, City University of New York, Flushing, NY 11367-1597, USA

[7]New York Consortium in Evolutionary Primatology, New York, NY, USA

[8]Anthropology Program, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016, USA

[9]Biology and Psychology Programs, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016, USA

[10]Department of Archaeology, University of Cape Town, Rondebosch 7701, Cape Town, South Africa

## ABSTRACT

At a time when seasonal cycles are increasingly disrupted, the ecology and evolution of reproductive seasonality in tropical vertebrates remains poorly understood. In order to predict how changes in seasonality might affect these animals, it is important to understand which aspects of their diverse patterns of reproductive phenology are linked to either the equally diverse patterns of rainfall seasonality (within-year variations) or instead the marked climatic unpredictability (year-to-year variations) occurring across the intertropical belt. Here, we gather birth and climatic seasonality data from 19 populations of ten Africa-dwelling primate species from the papionin tribe, from strict seasonal to non-seasonal breeders, occupying a wide range of environments, including equatorial, tropical, temperate and arid climates. We investigate (1) the environmental measures that influence the intensity of reproductive seasonality, and (2) the reproductive stage that is synchronized with increased resource availability. Our results demonstrate wide variation in the intensity of birth seasonality between and within species. Across multiple measures of climatic variation, we found rainfall unpredictability to be the only clear predictor of the intensity of reproductive seasonality across populations, i.e., greater year-to-year variation in the amount of rainfall was associated with lower to no reproductive seasonality. Finally, we identified diverse patterns of reproductive phenology, with the most seasonal breeders generally aligning lactation with the peak in resource availability while other populations show more diverse patterns, where conception, lactation or weaning can all be synchronized with maximal food availability. This study sheds new light on the extent and ecological drivers of flexible reproductive phenology in long-lived tropical mammals, and may even contribute to our understanding of why humans give birth year-round.

## INTRODUCTION

Most animals face variation in their environment across the year (Boyce, 1979) in the form of

seasonal fluctuations in rainfall, temperature and resource availability that affect their energy

balance. Reproductive seasonality, the temporal clustering of reproductive events in the annual

cycle, is thought to be beneficial because it synchronizes the most energetically costly

reproductive stage with the seasonal food peak, thereby enhancing the condition and survival

probability of mothers and offspring (Bronson, 2009; Bronson & Heideman, 1994). Variation

in birth frequencies across the annual cycle is a continuous trait, ranging from a complete

absence of reproductive seasonality (i.e., random distribution of births throughout the year), as

in mountain gorillas (Campos et al., 2017), to cases in which all births occur within a few weeks

each year, as in many lemurs (Wright, 1999).

Comparative studies investigating determinants of variation in reproductive seasonality

across mammals have often been conducted at the level of the order (rodents:  Heldstab, 2021,

lagomorphs: Heldstab, 2021, ruminants: Rutberg, 1987; Zerbe et al., 2012, Carnivora: Heldstab

et al., 2018, primates: Di Bitetti & Janson, 2000; Heldstab et al., 2020; Janson & Verdolin,

2005) and thus focus on broad-scale macro-evolutionary patterns. These studies have typically

detected a relationship between geographic latitude and birth seasonality, suggesting that at

higher latitudes, birth seasonality is more pronounced, with a more intense birth peak (more

births concentrated over a shorter duration of time). However, important gaps remain in our

understanding of the determinants of reproductive seasonality. Few studies have attempted to

quantify the extent of variation in reproductive seasonality across multiple populations of the

same species (but see in African wild dogs (*Lycaon pictus*): McNutt, Groom, & Woodroffe,

2019, and in red-tailed monkeys (*Cercopithecus ascanius*): Struhsaker, 1997), or across closely

related species sharing relatively similar diets, body sizes and life histories (but see in several

ungulate species: Pereira, Dos Santos Zanetti, & Furlan Polegato, 2010; Spinage, 1973; Brogi

et al., 2022; macaque species: Trébouet, Malaivijitnond, & Reichard, 2021). Thus, studies that

control for major sources of variation in life history or broad dietary categories should be

particularly useful for identifying the climatic drivers of variation in reproductive phenology.

The well-known association between latitude and reproductive seasonality fails to

explain the diversity of reproductive seasonality patterns observed within restricted latitudinal

ranges, such as in the tropics (Heldstab et al., 2020; Janson & Verdolin, 2005). In addition,

latitude encapsulates multiple components of climatic variation, which need to be disentangled

in order to identify the main climatic factors at play (Burtschell, Dezeure, Huchard, & Godelle,

2023). First, latitude correlates positively with the degree of environmental seasonality,

measured as the magnitude of within-year variation (such as the difference between maximal

and minimal monthly rainfall in the annual cycle) (Botero, Dor, McCain, & Safran, 2014).

Further, latitude covaries negatively with environmental productivity, i.e., overall food

availability in a given environment. Variation in productivity may alter the benefits of seasonal

breeding, as populations living in more productive habitats may face less pressure to breed

seasonally (Burtschell et al., 2023).

Finally, environmental predictability, independently of latitude and seasonality (Tonkin,

Bogan, Bonada, Rios-Touma, & Lytle, 2017), could also influence breeding schedules. In

locations with intense year-to-year environmental variation, flexible reproduction may be more

advantageous than strictly seasonal reproduction (Brockman & van Schaik, 2005a; van Schaik

& van Noordwijk, 1985). Indeed, regular delays or decreases in the food peak may lead to

reproductive failures in strict seasonal breeders, thus reducing the fitness benefits of breeding

seasonally. However, few studies have investigated the effects of environmental

unpredictability on the intensity of reproductive seasonality, with mixed results so far. While

English *et al.* (2012) found that higher inter-annual variation in food availability decreased the

intensity of birth synchrony across 38 ungulate species, two studies of red deer, *Cervus elaphus*

L. (Loe *et al.*, 2005) and chacma baboons, *Papio ursinus* (Dezeure *et al.*, 2023) found no effect

of environmental unpredictability on reproductive seasonality. A recent modelling study

similarly detected limited effects of environmental unpredictability on evolutionary transitions

to nonseasonal breeding (Burtschell et al., 2023).

Aside from the selective pressures favouring a flexible reproductive phenology,

relatively little is known about how birth timing varies in relation to the annual resource peak

in long-lived species. In short-lived species, the full reproductive cycle, from conception to

offspring nutritional independence (weaning in mammals), can take place within a single

productive season (Bronson, 2009). However, this is not the case for long-lived species, in

which multiple stages of a female's reproductive cycle can be aligned with the annual food peak,

with varying fitness consequences (Dezeure et al., 2021). For example, females of some species

may have to reach a certain threshold of body condition for the onset of reproduction and

conception to take place (Brockman & van Schaik, 2005a), meaning that most conceptions are

expected to follow a peak of food availability (Brockman & van Schaik, 2005a). In some other

species, females may instead synchronize the costliest part of their reproductive cycle with the

most productive season so as to enhance maternal condition and survival (Bronson, 2009;

Bronson & Heideman, 1994), such that early- or mid-lactation occurs during the annual food

peak, as in many primates (J. Altmann, 1980; Brockman & van Schaik, 2005a; Janson &

Verdolin, 2005). Lastly, weaning is a critically vulnerable life stage, where juveniles must begin

to forage for themselves (J. Altmann, 1980; Lee, 1996). Accordingly, several species have been

shown to time their births so as to align weaning with the seasonal food peak (Brockman & van

Schaik, 2005a; Janson & Verdolin, 2005), as occurs in most lemurs (Wright, 1999). Overall,

the reasons underlying the observed variation in alignment of reproductive stages with the food

peak across species and populations remain largely unknown.

In this study, we attempt to address the above gap in our understanding by investigating

the evolutionary determinants of the intensity and timing of reproductive seasonality in Africa-

dwelling papionin monkeys. We focus on papionins for several reasons. First, they exhibit relatively similar body sizes (large-bodied), life history traits (slow) and diet (omnivorous) (Kingdon et al., 2012; Swedell, 2011), allowing us to investigate environmental effects on reproductive seasonality while controlling for these – potentially confounding – factors. Second, this taxonomic group displays a wide diversity of patterns of reproductive seasonality. Indeed, most baboon (*Papio* spp.) species are non-seasonal breeders (Bercovitch & Harding, 1993; Swedell, 2011) despite exhibiting variation in monthly birth frequencies (Cheney et al., 2004; Lycett, Weingrill, & Henzi, 1999), while mandrills (*Mandrillus sphinx*) (Setchell, Lee, Wickings, & Dixson, 2002), Kinda baboons (*Papio kindae*) (Petersdorf, Weyher, Kamilar, Dubuc, & Higham, 2019) and most mangabey species (i.e., *Cercocebus* and *Lophocebus* spp.) (Swedell, 2011) are seasonal breeders. Third, this species constellation exhibits great ecological flexibility, inhabiting arid areas, woodland savannahs, equatorial forests, and high altitude grasslands (J. Fischer et al., 2019; Kingdon et al., 2012; Swedell, 2011) (see also Figure 1). Fourth, within the African members of this tribe, baboons are one of the most well studied primates, with data available from multiple populations of each species within the genus *Papio* (J. Fischer et al., 2019).

Lastly, species from this taxonomic group, and in particular from the genus *Papio*, possess a variety of features shared with early hominins (Alberts et al., 2005; Brockman, 2005; Jolly, 2001): they are large, terrestrial and eclectic omnivorous primates (Alberts et al., 2005; Rhine, Norton, Wynn, & Wynn, 1989) that, unlike the great apes, have colonized African savannahs (Bobe, Martínez, & Carvalho, 2020) and give birth to a single offspring every one to three years (J. Altmann & Alberts, 2005; Swedell, 2011). Investigating the environmental determinants of their reproductive seasonality may thus shed new light on the evolution and maintenance of non-seasonal breeding in early hominins (King, 2022).

Here we ask three main questions regarding reproductive seasonality in the papionins

in our sample:

(i) What is the extent of inter- and intra-specific variation in patterns of reproductive

seasonality, specifically regarding the height and width of the birth peak, as well as its timing

in the annual cycle?

(ii) What are the main environmental factors responsible for variation in the intensity of

reproductive seasonality? We isolated eight components of environmental variation: latitude,

environmental productivity, magnitude of seasonal variation in rainfall, number of rainy

seasons, breadth of the rainy season, amount of between-year (unpredictable) variation in

rainfall, between-year variation in the timing of the rainfall season, and the type of habitat. We

tested the eight corresponding hypotheses (H1.1-1.8) and their associated predictions, which

are listed in Table 1.

(iii) In seasonally breeding populations, which stage of the reproductive cycle is synchronized

with the food peak? We tested whether females match the seasonal food peak with conceptions

(H2.1 - the 'conception hypothesis'), lactation (H2.2 - the 'lactation hypothesis'), or weaning

(H2.3 - the 'weaning hypothesis').

**METHODS**

*1- Sample and data selection*

Our data set includes published reports on reproductive seasonality in natural populations of

Africa-dwelling papionins from the genera *Cercocebus*, *Lophocebus*, *Macaca, Mandrillus*,

*Papio*, *Rungwecebus* and *Theropithecus* (Figure 1). We selected papers that reported the

number of births per month (except for yellow baboons, *Papio cynocephalus*, from Mikumi

National Park, where births were provided in three month-periods). We obtained birth

seasonality data from 21 wild populations representing 11 species: see Table S1 for references

associated with each population, and Figure 1 for their locations. We did not find any monthly

birth data for nine species of interest for which only data from captivity were available (Kingdon

et al., 2012; Swedell, 2011): five species of *Cercocebus* (*agilis*, *chrysogaster*, *galeritus*,

*lunulatus*, *torquatus*), as well as *Lophocebus aterrimus*, *Mandrillus leucophaeus*, *Papio papio*

and *Rungwecebus kipunji* (Figure 1).

*2- Birth seasonality data*

We were interested in quantifying two components of reproductive seasonality: the timing of

the birth peak and the intensity of birth seasonality. For our analysis of the monthly distribution

of births, we considered each birth to have occurred in the middle of the month (i.e., the 15th of

each month, except for February where births were considered to occur on the 14th). For the

Mikumi population, we considered that births occurred in the middle of each 3 month-period.

We then used a circular statistic and represented each birth on the annual circle by a vector of

length 1 and of angle θ representing its date ($15 \times 2 \times \pi / 365.25$ for January, $(14+31) \times 2 \times \pi / 365.25$

for February, etc.). We computed the mean vector (r-vector) per population, whose angle

(converted in a date: $\mu_{birth}$) indicates the mean day of the year in which births occur (see Table

S1) and is thus a measure of birth seasonality. We computed $\mu_{birth}$ using the function

'circ.summary' from the 'CircStats' package (Agostinelli & Lund, 2018). The length ($r_{birth}$) of

the r-vector measures the intensity of birth seasonality, i.e., the degree of uniformity of the birth

distribution across the annual cycle, varying from 0 to 1 (Di Bitetti & Janson, 2000; Janson &

Verdolin, 2005; Thompson & McCabe, 2013). When $r_{birth}=0$, births are evenly spread across

months (i.e., non-seasonal), while when $r_{birth}=1$, births all occur during the same month of the

year (extremely seasonal). After comparing several classical measures of reproductive

seasonality (Supplementary Materials, Appendix S1), we used only $r_{birth}$ to measure the

intensity of reproductive seasonality, as this measure is more robust to differences in sample

size than other metrics, facilitating the comparison of seasonality measures between

populations (Janson & Verdolin, 2005; Thel, Chamaillé-Jammes, & Bonenfant, 2022).

*3- Environmental data*

*i.*     *Two indicators of environmental variation: rainfall and NDVI*

In order to test our set of hypotheses, we considered environmental variation through

two components: rainfall and the Normalized Difference Vegetation Index (NDVI). NDVI

produces a quantitative index of vegetation productivity, where higher values indicate a higher

degree of vegetation cover (Didan, Barreto Munoz, Solano, & Huete, 2015). Climatic

seasonality in Africa (and in most tropical habitats) is mainly characterized by within-year

variation in rainfall (Alberts et al., 2005; Feng, Porporato, & Rodriguez-Iturbe, 2013; Van

Schaik, Terborgh, & Wright, 1993), which has been successfully used as an indicator of food

availability for several of our studied populations (Alberts et al., 2005; Hill, Lycett, & Dunbar,

2000; Petersdorf et al., 2019; Tinsley Johnson, Snyder-Mackler, Lu, Bergman, & Beehner,

2018). In addition, NDVI values have to be used with caution when comparing productivity

across environments (Pettorelli et al., 2005), which is why we opted to use rainfall to test

hypotheses H1.1-H1-8. For example, the mean annual NDVI value at De Hoop, one of our

driest habitats, was almost equal to that of Gashaka, one of our wettest habitats. We thus used

variation in rainfall, rather than in NDVI, to disentangle the various components of climatic

variation that may affect the intensity of reproductive seasonality (such as environmental

productivity, predictability and seasonality) when testing hypotheses H1.1-H1.8.

However, we opted to use NDVI as an index of food availability to calculate the timing

of the food peak when testing hypotheses H2.1-H2-3, as Africa-dwelling papionins rely mainly

on plants for their diet (Swedell, 2011). Specifically, the annual peak in NDVI – and presumably

in food availability - is likely to lag behind that of rainfall by a few weeks, and this lag duration

may vary depending on local climatic or environmental conditions (Bercovitch & Harding,

1993; Dezeure et al., 2021; Jarvey, Low, Pappano, Bergman, & Beehner, 2018). Consequently,

we thought that using NDVI would be preferable than rainfall to assess the timing of the annual

food peak, and test hypotheses regarding its match to the birth peak in each population.

*ii.      Data extraction*

Daily rainfall was extracted from satellite data sensors with the Giovanni NASA website

(product TRMM 3B42) (Huffman, Bolvin, Nelkin, & Adler, 2016) using a 0.25×0.25 degree

resolution (corresponding to between 28×28km at the equator and 23×23km at 35° latitude).

The GPS coordinates used for this extraction are indicated per population in Table S2, and were

assessed either from indications about the home ranges of the habituated groups per population

when available in the literature, or alternatively from the geographical location (Park, Reserve

or nearby city) of the population (see also Figure 1). Monthly cumulative rainfall (summed

across daily values) was subsequently computed between January 1998 and December 2019.

We therefore gathered 22 years of rainfall data per population over the same period of time.

We then extracted the mean NDVI per 16 day-period on a 500m × 500m resolution

within the same geographical areas used for rainfall extraction (see GPS coordinates in Table

S2) between March 2000 and March 2017 (data before and after these dates were not available

at the time of data extraction) using MODIS data (MODIS13A1 product) provided by NASA

(Didan et al., 2015). Daily NDVI was computed by linear interpolation and then averaged to

obtain a monthly value across 18 years.

*iii.     Components of environmental variation*

In order to test our hypotheses, we identified multiple components of rainfall variation within

and across years. First, we decomposed for a given site the observed rainfall value into three

components as follows: $\text{Rainfall}_{m,i} = K_{rain} + \text{Rainfall } S_m + \text{Rainfall } NS_{m,i}$, where m is the month

of the year (going from January to December) and i is the year (from 1998 to 2019). $K_{rain}$ is a

constant, equalling the mean monthly rainfall across 22 years of records (Figure S1). Rainfall

$S_m$ is the seasonal component of rainfall, i.e., the rainfall value, averaged across 22 years, for

each month of the year, minus $K_{rain}$ (Figure S1). For example, for a given site, Rainfall $S_1$ (m=1)

is the mean of all January rainfall values. The term Rainfall S thus captures the seasonal

component of rainfall variation in the annual cycle, i.e., its within-year variation. Finally,

Rainfall NS $_{m,i}$ is the non-seasonal component of rainfall, i.e., the difference between the

observed rainfall value in any month at a given site ($\text{Rainfall}_{m,i}$) and the predictable component

of rainfall variation for that particular site in that particular month ($K_{rain}$ + Rainfall $S_m$) (Figure

S1). This captures the unpredictable, i.e., between-year, rainfall variation. Using these

measures, we assessed the following for each population (see Table S2 for the values associated

with each population):

- Environmental productivity, or mean annual rainfall, equal to $12 \times K_{rain}$.

- Magnitude of environmental seasonality. We computed the magnitude of within-year

rainfall variation, as the relative standard deviation (SD) of Rainfall S standardized for

environmental productivity, given by the formula: $\frac{100 \times SD(Rainfall\ S)}{12 \times Krain}$. The higher the

value, the more seasonal is rainfall variation.

- The number of rainy seasons per year. Using predictable rainfall variation ($K_{rain}$ +

Rainfall S), we assessed graphically, for each population, the number of rainy seasons

per year.

- The length of the rainy season. For environments with only one rainy season, we further

calculated the rainfall peak breadth (RPB), which is the minimum number of

consecutive months of the year during which 80% of the annual rainfall ($12*K_{rain}$)

occurs. This measure is meaningless for environments with more than one rainy season,

and we thus excluded the populations living in such environments from this analysis.

-    Magnitude of environmental unpredictability. We computed the magnitude of between-

year rainfall variation, as the relative standard deviation of Rainfall NS, standardized by

environmental productivity, given by the formula: $\frac{100 \times SD(Rainfall\ NS)}{12 \times K\text{rain}}$. The higher the

value, the more unpredictable rainfall variation is.

Using the literature, we categorized the habitat of each population into three types:

tropical forest, open savannah, and mosaic forest-grassland (Table S2).

In addition to the magnitude of environmental unpredictability, we were interested in

quantifying unpredictability in the timing of the annual rainfall peak, i.e., quantifying how much

the timing of rainfall varied between years. Details on the procedure can be found in Appendix

S2, but briefly, using circular statistics, we computed standard deviations of the mean dates of

the annual rainfall peak over the 20 years sampled. Values close to zero mean that rainfall peak

occurs the same month every year, and the higher the value, the more variation in rainfall peak's

timing between years.

Lastly, we computed the mean monthly NDVI across 18 years (i.e., $K_{NDVI}$) and the

seasonal component of NDVI variation for each month of the year (i.e., NDVI S), following

the same notation used to disentangle the components of rainfall variation. To characterise the

timing of the seasonal food peak, we used circular statistics to compute the mean annual NDVI

date, $\mu_{NDVI}$ (see Appendix S2 for methodology and Table S2 for values of $\mu_{NDVI}$).

*iv – Phylogenetic tree*

We used the branch length of Version 3 of the 10kTreesPrimates consensus tree (Arnold,

Matthews, & Nunn, 2010). Two species of interest were absent from this tree: *Cercocebus*

*sanjei* and *Papio kindae*. For the former, we substituted *Cercocebus galeritus*, its closest

relative. For the latter, following recent genetic studies (Jordan et al., 2018; Rogers et al., 2019),

we added *Papio kindae* in the same branch as *Papio ursinus*, using the function 'bind.tip' from

the package 'phytools' (Revell, 2012).

## 4- *Statistical analysis*

302 All statistical analyses were conducted in R version 3.5.0 (R Core Team, 2019).

### i. *Factors affecting the intensity of reproductive seasonality*

We considered eight potential environmental determinants of reproductive seasonality,

which are listed along each corresponding prediction in Table 1.

For each hypothesis, we plotted $r_{birth}$ (the r-vector length measuring the intensity of

reproductive seasonality in a given population) versus the tested predictor. We then checked

the significance of the relationship between each predictor and $r_{birth}$ while controlling for

phylogeny using Bayesian phylogenetic generalized linear mixed models with the package

'MCMCglmm' (Hadfield, 2010). We included the phylogenetic relationship between species

as a covariance matrix, which was derived from the phylogenetic tree. We transformed $r_{birth}$

(which could vary from 0 to 1) by multiplying it by 10000 and taking its integer values, and

thereafter used a Poisson distribution.

Given the low number of populations in this study, and the collinearity between some

of our environmental parameters (multivariate models had variance inflated factors >3), we

were not able to run stable multivariate models. For example, the magnitude of environmental

seasonality was negatively associated with the duration of the rainy season (cor=-0.94, t=-9.15,

$p<10^{-4}$), and environmental productivity was negatively associated with both latitude (cor=-

0.55, t=-2.69, p=0.016) and the magnitude of environmental unpredictability (cor=-0.80, t=-

5.48, $p<10^{-4}$).

Therefore, our Poisson regressions included as a response variable the transformed $r_{birth}$

values, one fixed effect (each environmental predictor in turn listed in Table 1, standardised if

continuous) and the phylogenetic matrix as a random effect. For each model, we set an

informative prior and used 500 000 iterations, a burn-in of 100 000 and a thinning interval of

100. We visually inspected for convergence and checked the absence of autocorrelations for the

posterior distributions of fixed and random effects. A Bonferroni correction was applied due to

the number of tests, so that predictors were considered statistically significant when their

associated pMCMC values were less than 0.05/n, with n being the number of tests run for a

given research question.

*ii.*       *Timings of conceptions and births in relation with NDVI seasonality*

We assessed whether populations were seasonal breeders using the Rayleigh test for circular

statistic, more precisely the 'r.test' function from 'CircStats' package (Agostinelli & Lund,

2018). Populations were categorized as seasonal breeders when P-values associated with the

Rayleigh test were <0.05, meaning that the null hypothesis of a uniform birth distribution could

be rejected. Among these seasonally breeding populations (listed in Table S1), we investigated

which reproductive stage (H2.1, 'conception'; H2.2, 'lactation'; or H2.3, 'weaning') was

synchronized with the annual NDVI peak, i.e. $\mu_{NDVI}$. Additional details are given in

Supplementary Materials, Appendix 3 & Table S3. We employed exact two-sample Fisher-

Pitman permutation tests, using the 'oneway_test' function from the 'coin' package (Hothorn,

Hornik, Van De Wiel, & Zeileis, 2006). This function tests if the observed monthly value of

NDVI during a target period, which depends on the hypothesis tested, is significantly higher

than monthly values of NDVI randomized across the entire year. For example, using mean

gestation length to infer the annual distribution of conception dates, we tested H2.1 by asking

if females tended to conceive during, soon before, or soon after the annual food peak, looking

at seasonal NDVI values respectively in (i) the six months surrounding $\mu_{conc}$, (ii) the three

months before $\mu_{conc}$ and (iii) the three months after $\mu_{conc}$.

**RESULTS**

*1) How variable are patterns of reproductive seasonality?*

The annual distribution of births for each population is shown in Figure 2, alongside seasonal

variation in rainfall and NDVI. The intensity of reproductive seasonality varies widely across

species: hamadryas baboons (Filoha population: $r_{birth}$ =0.02) and most olive baboon populations

($r_{birth}$ <0.22) show non-seasonal births while mandrills (Lékédi: $r_{birth}$ =0.67 and Moukalaba-

Doudou: $r_{birth}$ =0.80 resp.) and Kinda baboons (Kasanka: $r_{birth}$ =0.50) exhibit pronounced birth

seasonality (Figure 3). The phylogenetic signal associated with the intensity of reproductive

seasonality is substantial (Lambda=0.61, 95% Highest Posterior Density Intervals = [0.002 ;

0.923]), indicating that more closely related species have more similar patterns of reproductive

seasonality. However, such a value is also compatible with some phylogenetic flexibility.

Among papionins, the genera *Mandrillus*, *Cercocebus* and *Macaca* show strong reproductive

seasonality, whereas the genera *Lophocebus*, *Papio* and *Theropithecus* show an overall lower

intensity of birth seasonality, which may be associated with greater flexibility within species

(Figure 3). Such flexibility is particularly pronounced in chacma baboons, extending from low

(de Hoop: $r_{birth}$ =0.10; Tokai: $r_{birth}$ =0.22, and Tsaobis: $r_{birth}$ =0.10) to moderate birth seasonality

(Drakensberg: $r_{birth}$ =0.42; Moremi: $r_{birth}$ =0.37) (Figure 3), while intra-specific variation is less

marked for the other species in our sample.

The timing of birth seasonality can also be surprisingly variable, even between species

that live in adjacent geographical ranges: for instance, Kinda baboons from Kasanka give birth

mainly around July, whereas chacma baboons from Moremi give birth primarily between July

and November (Figure 2).

*2) What are the ecological factors affecting the intensity of reproductive seasonality?*

We detected an effect of the magnitude of environmental unpredictability on the intensity of reproductive seasonality, while controlling for species relatedness, which supported our prediction H1.6 (Table 2): the higher the magnitude of between-year variation in rainfall, the lower the intensity of reproductive seasonality (Figure 4F). We also found an effect of habitat productivity that contradicted our prediction H1.2 (Table 2): the lower the mean annual rainfall, the lower the intensity of reproductive seasonality (Figure 4B). Note that these two effects lost significance when applying a Bonferroni correction for multiple tests, which was expected given our modest sample size. Lastly, we did not find any support for the other five hypotheses: there was no effect of latitude, magnitude of rainfall seasonality, number of rainy seasons, breadth of the rainfall peak, unpredictability in the timing of the annual peak of rainfall, or habitat on the intensity of reproductive seasonality (Table 2, Figure 4).

*3) Which stage of the reproductive cycle is timed with the annual food peak?*

When considering all taxa in our sample as a whole, none of our three hypotheses were clearly supported (H2.1, H2.2, H2.3). However, in five of the six populations with the strongest reproductive seasonality ($r_{birth}$ >0.5), females appeared to synchronize lactation with the annual NDVI peak, and overall lactation was generally aligned with the NDVI peak in 8 of the 14 populations with significant birth seasonality (P-value of the Rayleigh test <0.05) (Table 3, Table S4). In less seasonally breeding populations, females were more variable in the reproductive stages that were timed with the annual NDVI peak, ranging from conception (6 of 14 populations, and mainly before than after conception: e.g. Amboseli or Kibale) to weaning (3 of 8 populations: e.g., Moremi or Gilgil) to none (e.g., Simien) or all of these stages (e.g., Udzungwa) (Table 3, Table S4). Moreover, the timing of the annual NDVI peak compared to

396  the mean conception, birth or weaning dates was highly variable between populations (Table

397  S4).

## DISCUSSION

400  We revealed strong inter- and intra-specific variation in the intensity of reproductive seasonality

401  as well as in the annual timing of births in Africa-dwelling papionins. Our study further

402  emphasizes the importance of environmental unpredictability for the evolution of flexible

403  reproductive seasonality. Lastly, we found that females from different populations of Africa-

404  dwelling papionins match different reproductive stages with the annual food peak.

**Papionins exhibit flexible reproductive seasonality**

Of the sampled Papio populations, most showed very little seasonality, about a third showed

moderate seasonality, and Kinda baboons departed from the overall baboon pattern in being

highly seasonal  breeders. Such diverse patterns of reproductive seasonality within a single

genus have rarely been reported outside primates (see for example *Cervus*: English et al., 2012;

Loe et al., 2005; Rutberg, 1987, *Damaliscus*: Rutberg, 1987, *Ovis*: Rutberg, 1987, *Ursus*:

Spady, Lindburg, & Durrant, 2007, *Mustela*: Heldstab et al., 2018, *Vulpes*: Heldstab et al.,

2018), but apparently occur in some other primate genera with large distribution ranges, such

as *Alouatta* (Di Bitetti & Janson, 2000; Janson & Verdolin, 2005), *Cercopithecus* (Heldstab et

al., 2020; Janson & Verdolin, 2005), *Cebus* (Janson & Verdolin, 2005) and *Macaca* (Heldstab

et al., 2020; Janson & Verdolin, 2005; Trébouet et al., 2021). Given the limited taxonomic scale

of our study, it is impossible to establish whether seasonal breeding was the ancestral state in

papionins, but it seems possible that the loss of seasonal reproduction is a derived state affecting

the *Papio* genus (Fig 3).

Importantly, the key adaptive trait that evolved in the *Papio* genus may not be simply

the loss of breeding seasonality *per se*, but the evolution of a flexible reproductive phenology.

Such flexibility would imply that each population, or even each individual (Dezeure et al., 2021,

2023; Dezeure, Charpentier, & Huchard, 2022), may be able to adjust phenology to current

environmental conditions. In line with this, intraspecific variation is particularly pronounced in

chacma baboons, the only species in our sample represented by more than three populations

with a reasonable number of births. In addition, chacma baboons exhibit a large distributional

range characterized by exceptional ecological diversity, which includes cold and temperate

climates, oceanic and mountainous ecosystems, and tropical and arid savannahs.

Flexibility in reproductive phenology may be facilitated by several mechanisms. First,

a slower life history may allow papionin species to spread the energetic costs of reproduction

over a prolonged period, such that pregnant or lactating females face only a small daily extra

energetic expenditure that can be afforded at any time, as suggested by a recent modelling study

(Burtschell et al., 2023). In addition, unlike many other mammals, cercopithecids do not use

strict photoperiodic cues to trigger their reproduction  (Heldstab et al., 2020) but may instead

exhibit condition-induced reproduction, whereby conceptions (and/or cycle resumption after

lactation) are more likely to occur when females are in better condition (Alberts et al., 2005;

Beehner, Onderdonk, Alberts, & Altmann, 2006). Such reproductive flexibility may have

contributed to their historical ecological success via their ability to colonize diverse

environments, and may become a critical asset to facilitate their resilience to climate change,

associated with increasing environmental unpredictability (Feng et al., 2013).

**Environmental unpredictability may drive flexible reproductive seasonality**

We examined several possible climatic drivers of the intensity of reproductive seasonality

across our sample, and found that environmental unpredictability was a significant predictor,

with higher between-year rainfall variation being associated with lower reproductive seasonality. So far, most studies investigating climatic effects on reproductive phenology have focused on environmental seasonality, i.e., the magnitude of within-year environmental variation. In primates, the effect of environmental unpredictability on reproductive seasonality, e.g., through climatic events such as el Niño or fruit mast years in South-East Asia, has been suggested (Brockman & van Schaik, 2005a; van Schaik & van Noordwijk, 1985; Wiederholt & Post, 2011) but had never been tested. In line with our results, a previous study across 70 ungulate populations showed that the birth peak is more spread out in environments with more year-to-year environmental variation (English et al., 2012).

Unpredictable climates may considerably reduce the fitness benefits associated with seasonal breeding, such as enhancing maternal condition and offspring survival. In southern Africa, year-to-year climatic variation frequently takes the shape of an absence of rain during the rainy season (Alberts et al., 2005), which could cause severe reproductive costs in seasonal breeders who often synchronize lactation or weaning with the rainy season, subsequently forcing females to wait until the next breeding season to initiate a new reproductive event. In such conditions, other adaptive traits may be more advantageous than seasonal breeding to face the energetic costs of reproduction, such as a better ability to store energy (Brockman & van Schaik, 2005a; van Schaik & van Noordwijk, 1985), to expand the dietary repertoire via a generalist diet or foraging innovations (Grueter, 2017), to increase daily foraging time (Alberts et al., 2005; Grueter, 2017), to flexibly adjust lactation duration (Dezeure et al., 2021; van Noordwijk, 2012), or to reproduce cooperatively (Cornwallis et al., 2017; Lukas & Clutton-Brock, 2017). Papionin species show many such traits: they can store energy, they have an eclectic omnivorous diet, and they are flexible foragers that typically rely on fallback foods during the dry season (J. Altmann, Schoeller, Altmann, Muruthi, & Sapolsky, 1993; S. A. Altmann, 2009; Swedell, 2011).

Two recent studies found no or little effect of climatic unpredictability on the intensity

of reproductive seasonality at the population level, one in chacma baboons (Dezeure et al.,

2023) and one on yellow baboons using a modelling approach (Burtschell et al., 2023),

questioning the robustness of the effect found in this study. This discrepancy may come from

the fact that unpredictable climates select for reproductive flexibility, rather than nonseasonal

breeding. In fact, the study by Burtschell et al., 2023 revealed that increasing climatic

unpredictability was associated with a lower variance, but not a lower mean, in the intensity of

reproductive seasonality. In addition, the effect of climatic unpredictability may be better

detected across space than time, i.e., by comparing different populations living in distinct

climates and environments, as is the case here, than by comparing the same population across

time, as was the case for Dezeure et al., 2023. Different study designs should be combined,

across time, taxonomic or spatial scales, to reveal the full complexity of selective pressures at

play.

The numerous pressures affecting the intensity of reproductive seasonality are often

hard to disentangle, meaning that the effects uncovered by correlational studies like ours may

sometimes reflect other co-varying pressures, and should thus be interpreted cautiously.

Specifically, environmental productivity, which is negatively correlated with climatic

unpredictability at our study sites (cor=-0.80, t=-5.48, $p<10^{-4}$), was also a significant predictor

of the intensity of reproductive seasonality, but in a direction opposed to our prediction, as well

as to previous results (Burtschell et al., 2023). In our sample, the least productive climates are

also the most unpredictable, and environmental productivity may thus represent a confounding

factor in the relationship between climatic unpredictability and seasonal breeding. In species

with a flexible phenology, females can start a new reproductive event rapidly after a

reproductive failure, without having to wait the next mating season. Such failures are likely to

be particularly frequent where environmental productivity is low, contributing to spread

reproductive events across the year cycle, and explaining how environmental productivity may

contribute to decrease reproductive seasonality.

Lastly, our decomposition of rainfall components further calls for a more rigorous

definition of the term seasonality, especially when it is used in a quantitative way. Indeed, a

'more seasonal' environment can either be an environment with higher within-year variation

(i.e. the amplitude of variation between the 'best' and the 'worst' month of the year), with

higher within-year over between-year variation (i.e. the amplitude of within-year variation

controlling for the intensity of unpredictable variations), with a shorter productive season (i.e.

the rainfall peak breadth), or with only a unimodal season (i.e. one rainy, or one warm season

per year). Similarly, climatic unpredictability can be broken into two components: (1) the

amount/magnitude of year-to-year variation (i.e., if the rainy season brings more or less rainfall

than usual), and (2) year-to-year variation in the timing of the rainy season (i.e., if the rainy

season occurs earlier or later than usual) (Clauss, Zerbe, Bingaman Lackey, Codron, & Müller,

2020). These various components have rarely been disentangled in empirical studies so far, and

this study opens new methodological avenues to investigate various environmental components

that are likely to affect reproductive seasonality.

**Females can match different reproductive stages with the food peak**

The main pattern emerging from our investigations suggests that females from species with

high breeding seasonality match lactation with the peak in vegetation productivity (Table 3).

For species and populations with a more flexible reproductive phenology, females can match

different reproductive stages with the annual vegetation peak, with a possible preference

towards conception. This trend may reflect the condition-dependence of conception – a

proximate mechanism – rather than an adaptive, optimal strategy aimed at synchronizing the

vegetation peak with a particular reproductive stage. These results, obtained from (mostly)

tropical primates, echo the broader mammalian literature showing that most mammals from temperate regions match lactation with the best season of the year (Bronson, 2009; Bronson & Heideman, 1994), while patterns are more variable in tropical and long-lived mammals, depending on body size, energy storage capacities and environmental predictability (Brockman & van Schaik, 2005a; Janson & Verdolin, 2005; van Schaik & van Noordwijk, 1985). Finally, even though weaning is a vulnerable life-history stage in young primates, which can be buffered when matched with the vegetation peak in wild chacma baboons (Dezeure et al., 2021), few populations seemed to adopt this strategy. Our estimations of lactation peak and weaning might lack accuracy, due to strong between-populations and between-individuals variation. Data quantifying maternal energy expenditure during lactation (Rosetta, Lee, & Garcia, 2011), or isotopic measures of trophic levels between mother and infants (Reitsema, 2012; Reitsema, Partrick, & Muir, 2016) would be necessary, for each population, to determine the dynamics of lactation and weaning. Second, additional unexplored factors can potentially affect reproductive timing and further limit our ability to detect a clear pattern. For example, for populations living at high altitudes like the geladas from Simien and the chacma baboons from Drakensberg, seasonal variation in temperatures also constrain reproductive phenology (Lycett et al., 1999; Tinsley Johnson et al., 2018).

**The evolution of reproductive flexibility in Anthropoid primates may inform our understanding of the reproductive phenology in early humans**

Baboons and relatives represent an interesting model for understanding the evolution of behavioural and reproductive plasticity of early humans (J. Fischer et al., 2019; King, 2022). Although most great apes are nonseasonal breeders (Brockman & van Schaik, 2005b; Campos et al., 2017), suggesting that their common ancestor also bred year-round, humans are distinct from other apes by exhibiting much faster reproductive paces (which are similar to most Africa-

dwelling papionins: Swedell, 2011), and by living in a wider variety of environments, rather

than being restricted to tropical forests (Wells & Stock, 2007). The selective pressures that have

shaped reproductive phenology in the human lineage versus in other apes may therefore differ,

and the papionins, who have similarly left forested habitats to colonize savannahs may provide

valuable insights to understand the adaptation of early humans to such diverse and

unpredictable environments. Our results suggest that baboons have acquired a low and flexible

reproductive seasonality, as well as several other adaptive traits, when facing more arid and

unpredictable environments, such as a generalist omnivorous diet and the frequent use of

fallback foods, frequent foraging innovations, an increased ability to store fat and to switch

home ranges to more suitable areas. Similar reproductive, physiological and behavioural

adaptations to environmental unpredictability may have allowed early humans to thrive and

maintain fast reproductive paces during their colonization of a wide variety of environments.

**Conclusion**

Our work revealed substantial variation in patterns of reproductive seasonality within and

across species of Africa-dwelling papionins, highlighting an exceptional flexibility in their

reproductive phenology. Among multiple dimensions of climatic variation, rainfall

unpredictability and productivity were the main predictors of the intensity of reproductive

seasonality, with arid and unpredictable climates being associated with less seasonal

reproduction. Among populations with a pronounced breeding seasonality, females often match

lactation with the annual vegetation peak, while phenology patterns are very diverse in other

populations. This study sheds new light on the selective pressures shaping reproductive

seasonality in long-lived tropical mammals, as well as on potential adaptations to environmental

unpredictability. It may further provide an original contribution to understand why humans

breed year-round, given their phylogenetic ties and convergences in life-history and ecology

with our taxonomic group.

**ACKNOWLEDGMENTS**

We thank Cassandra Raby for her kind help on extracting NDVI data, and Christophe Dezeure

for making the Figure 1 of this paper (and more generally for his paternal care). This project

was funded by a grant from the Agence Nationale de la Recherche (ANR ERS-17-CE02-0008,

2018-2021) awarded to E. H.

**DATA AVAILABILITY**

The raw data and scripts used for this paper will be made available online upon submission, on

a Dryad repository.

**REFERENCES**

Agostinelli, C., & Lund, U. (2018). *Package "CircStats": Circular Statistics, from "Topics in Circular Statistics."*

Alberts, S. C., Hollister-smith, J. A., Mututua, R. S., Sayialel, S. N., Muruthi, P. M., Warutere, K. J., & Altmann, J. (2005). Seasonality and long-term change in a savanna environment. In D. K. Brockman & C. P. van Schaik (Eds.), *Seasonality in Primates: Studies of living and extinct human and non-human primates* (pp. 157–195). Cambridge: Cambridge Univ. Press.

Altmann, J. (1980). *Baboon Mothers and Infants*. The University of Chicago Press.

Altmann, J., & Alberts, S. C. (2005). Growth rates in a wild primate population: ecological influences and maternal effects. *Behavioral Ecology and Sociobiology*, *57*, 490–501. doi: 10.1007/s00265-004-0870-x

Altmann, J., Schoeller, D., Altmann, S. A., Muruthi, P., & Sapolsky, R. M. (1993). Body size and fatness of free-living baboons reflect food availability and activity levels. *American Journal of Primatology*. doi: 10.1002/ajp.1350300207

Altmann, S. A. (2009). Fallback foods, eclectic omnivores, and the packaging problem. *American Journal of Physical Anthropology*, *140*(4), 615–629. doi: 10.1002/ajpa.21097

Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: A new online resource for primate phylogeny. *Evolutionary Anthropology*, *19*(3), 114–118. doi: 10.1002/evan.20251

Beehner, J. C., Onderdonk, D. A., Alberts, S. C., & Altmann, J. (2006). The ecology of conception and pregnancy failure in wild baboons. *Behavioral Ecology*, *17*(5), 741–750. doi: 10.1093/beheco/arl006

Bercovitch, F. B., & Harding, R. S. O. (1993). Annual Birth Patterns of Savanna Baboons (Papio cynocephalus anubis) over a Ten-Year Period at Gilgil, Kenya. *Folia Primatologica*, *61*, 115–122.

Bobe, R., Martínez, F. I., & Carvalho, S. (2020). Primate adaptations and evolution in the Southern

African Rift Valley. *Evolutionary Anthropology*, *29*(3), 94–101. doi: 10.1002/evan.21826

Botero, C. A., Dor, R., McCain, C. M., & Safran, R. J. (2014). Environmental harshness is positively

correlated with intraspecific divergence in mammals and birds. *Molecular Ecology*, *23*(2), 259–

268. doi: 10.1111/mec.12572

Boyce, M. S. (1979). Seasonality and Patterns of Natural Selection for Life Histories. *The American*

*Naturalist*, *114*(4), 569–583. doi: 10.1086/283503

Brockman, D. K. (2005). What do studies of seasonality in primates tell us about human evolution? In

*Seasonality in Primates: Studies of Living and Extinct Human and Non-Human Primates* (pp.

543–570). Cambridge: Cambridge University Press. Retrieved from

https://www.researchgate.net/publication/230778889

Brockman, D. K., & van Schaik, C. (2005a). Seasonality and reproductive function. In D. K.

Brockman & C. P. van Schaik (Eds.), *Seasonality in Primates: Studies of Living and Extinct*

*Human and Non-Human Primates* (pp. 269–305). Cambridge University Press.

Brockman, D. K., & van Schaik, C. P. (2005b). *Seasonality in Primates: Studies of Living and Extinct*

*Human and Non-Human Primates* (D. K. Brockman & C. P. van Schaik, eds.). Cambridge:

Cambridge University Press.

Brogi, R., Merli, E., Grignolio, S., Chirichella, R., Bottero, E., & Apollonio, M. (2022). It is time to

mate: population-level plasticity of wild boar reproductive timing and synchrony in a changing

environment. *Current Zoology*, *68*(4), 371–380. doi: 10.1093/cz/zoab077

Bronson, F. H. (2009). Climate change and seasonal reproduction in mammals. *Philosophical*

*Transactions of the Royal Society B: Biological Sciences*, *364*(1534), 3331–3340. doi:

10.1098/rstb.2009.0140

Bronson, F. H., & Heideman, P. D. (1994). Seasonal regulation of reproduction in mammals. In E.

Knobil & J. D. Neill (Eds.), *The physiology of reproduction* (Vol. 2, pp. 541–583). New York:

New York Raven Press. Retrieved from https://www.researchgate.net/publication/313495409

Burtschell, L., Dezeure, J., Huchard, E., & Godelle, B. (2023). Evolutionary determinants of

reproductive seasonality: A theoretical approach. *Peer Community Journal*, *3*(e56). doi:

10.24072/pci

Campos, F. A., Morris, W. F., Alberts, S. C., Altmann, J., Brockman, D. K., Cords, M., … Fedigan, L.

M. (2017). Does climate variability influence the demography of wild primates? Evidence from

long-term life-history data in seven species. *Global Change Biology*, *23*(3), 1–15. doi:

10.1111/gcb.13754

Cheney, D. L., Seyfarth, R. M., Fischer, J., Beehner, J., Bergman, T., Johnson, S. E., … Silk, J. B.

(2004). Factors affecting reproduction and mortality among baboons in the Okavango Delta,

Botswana. *International Journal of Primatology*, *25*(2), 401–428. doi: 0164-0291/04/0400-

0401/0

Clauss, M., Zerbe, P., Bingaman Lackey, L., Codron, D., & Müller, D. W. H. (2020). Basic

considerations on seasonal breeding in mammals including their testing by comparing natural

habitats and zoos. *Mammalian Biology*. doi: 10.1007/s42991-020-00078-y

Cornwallis, C. K., Botero, C. A., Rubenstein, D. R., Downing, P. A., West, S. A., & Griffin, A. S.

(2017). Cooperation facilitates the colonization of harsh environments. *Nature Ecology and*

*Evolution*, *1*(3). doi: 10.1038/s41559-016-0057

Dezeure, J., Baniel, A., Carter, A., Cowlishaw, G., Godelle, B., & Huchard, E. (2021). Birth timing

generates reproductive trade-offs in a non-seasonal breeding primate. *Proceedings of the Royal*

*Society B*, *288*, 1–9. doi: 10.1098/rspb.2021.0286

Dezeure, J., Burtschell, L., Baniel, A., Carter, A. J., Godelle, B., Cowlishaw, G., & Huchard, E.

(2023). Evolutionary Determinants of Nonseasonal Breeding in Wild Chacma Baboons.

*American Naturalist*, *201*(1), 106–124. doi: 10.1086/722082

Dezeure, J., Charpentier, M. J. E., & Huchard, E. (2022). Fitness effects of seasonal birth timing in a

long-lived social primate living in the equatorial forest. *Animal Behaviour*, *185*, 113–126.

Di Bitetti, M. S., & Janson, C. H. (2000). When will the stork arrive? Patterns of birth seasonality in

neotropical primates. *American Journal of Primatology*, *50*(2), 109–130. doi:

10.1002/(SICI)1098-2345(200002)50:2<109::AID-AJP2>3.0.CO;2-W

Didan, K., Barreto Munoz, A., Solano, R., & Huete, A. (2015). *MOD13A1 MODIS/Terra Vegetation*

*Indices 16-Day L3 Global 500m SIN Grid V006 [Data set]*. doi:

https://doi.org/10.5067/MODIS/MOD13A1.006

English, A. K., Chauvenet, L., Safi, K., & Pettorelli, N. (2012). Reassessing the Determinants of

Breeding Synchrony in Ungulates. *PLoS ONE*, *7*(7), 1–7. doi: 10.1371/journal.pone.0041444

Feng, X., Porporato, A., & Rodriguez-Iturbe, I. (2013). Changes in rainfall seasonality in the tropics.

*Nature Climate Change*, *3*(9), 811–815. doi: 10.1038/nclimate1907

Fischer, J., Higham, J. P., Alberts, S. C., Barrett, L., Beehner, J. C., Bergman, T. J., … Zinner, D.

(2019). The Natural History of Model Organisms: Insights into the evolution of social systems

and species from baboon studies. *ELife*, *1960*, 1–16.

Fischer, Julia, Kopp, G. H., Dal Pesco, F., Goffe, A., Hammerschmidt, K., Kalbitzer, U., … Zinner, D.

(2017). Charting the neglected West: The social system of Guinea baboons. *American Journal of*

*Physical Anthropology*, *162*, 15–31. doi: 10.1002/ajpa.23144

Grueter, C. C. (2017). Environmental Seasonality. In *The International Encyclopedia of Primatology*

(pp. 1–2). John Wiley & Sons, Inc. doi: 10.1002/9781119179313.wbprim0048

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The

MCMCglmm R  Package. *Journal of Statistical Software*, *33*(2), 1–22. Retrieved from

https://www.jstatsoft.org/v33/i02/

Heldstab, S. A. (2021a). Latitude, life history and sexual size dimorphism correlate with reproductive

seasonality in rodents. *Mammal Review*, *51*, 256–271.

Heldstab, S. A. (2021b, December 1). Habitat characteristics and life history explain reproductive

seasonality in lagomorphs. *Mammalian Biology*, Vol. 101, pp. 739–757. Springer Science and

Business Media Deutschland GmbH. doi: 10.1007/s42991-021-00127-0

Heldstab, S. A., Müller, D. W. H., Graber, S. M., Bingaman Lackey, L., Rensch, E., Hatt, J. M., …

Clauss, M. (2018). Geographical Origin, Delayed Implantation, and Induced Ovulation Explain

Reproductive Seasonality in the Carnivora. *Journal of Biological Rhythms*, *33*(4), 402–419. doi:

10.1177/0748730418773620

Heldstab, S. A., van Schaik, C. P., Müller, D. W. H., Rensch, E., Lackey, L. B., Zerbe, P., … Matsuda,

I. (2020). Reproductive seasonality in primates: patterns, concepts and unsolved questions.

*Biological Reviews*. doi: 10.1111/brv.12646

Hill, R. A., Lycett, J. E., & Dunbar, R. I. M. (2000). Ecological and social determinants of birth

intervals in baboons. *Behavioral Ecology*, *11*(5), 560–564. doi: 10.1093/beheco/11.5.560

Hothorn, T., Hornik, K., Van De Wiel, M. A., & Zeileis, A. (2006). A lego system for conditional

inference. *American Statistician*, *60*(3), 257–263. doi: 10.1198/000313006X118430

Huffman, G. J., Bolvin, D. T., Nelkin, E. J., & Adler, R. F. (2016). *TRMM (TMPA) Precipitation L3 1*

*day 0.25 degree x 0.25 degree V7*. Retrieved from 10.5067/TRMM/TMPA/DAY/7

Janson, C., & Verdolin, J. (2005). Seasonality of primate births in relation to climate. In D. K.

Brockman & C. P. van Schaik (Eds.), *Seasonality in Primates: Studies of Living and Extinct*

*Human and Non-Human Primates* (pp. 307–350). Cambridge: Cambridge University Press. doi:

10.1017/cbo9780511542343.012

Jarvey, J. C., Low, B. S., Pappano, D. J., Bergman, T. J., & Beehner, J. C. (2018). Graminivory and

Fallback Foods: Annual Diet Profile of Geladas (Theropithecus gelada) Living in the Simien

Mountains National Park, Ethiopia. *International Journal of Primatology*, *39*(1), 105–126. doi:

10.1007/s10764-018-0018-x

Jolly, C. J. (2001). A proper study for mankind: Analogies from the Papionin monkeys and their

implications for human evolution. *Yearbook of Physical Anthropology*, *44*, 177–204. doi:

10.1002/ajpa.10021

Jordan, V. E., Walker, J. A., Beckstrom, T. O., Steely, C. J., McDaniel, C. L., St Romain, C. P., … Worley, K. C. (2018). A computational reconstruction of Papio phylogeny using Alu insertion polymorphisms. *Mobile DNA*, *9*(1). doi: 10.1186/s13100-018-0118-3

King, G. E. (2022). Baboon perspectives on the ecology and behavior of early human ancestors. *Proceedings of the National Academy of Sciences of the United States of America*, *119*(45). doi: 10.1073/pnas.2116182119

Kingdon, J., Happold, D., Butynski, T. M., Hoffmann, M., Happold, M., & Kalina, J. (2012). *Mammals of Africa*.

Lee, P. C. (1996). The meanings of weaning: growth, lactation, and life history. *Evolutionary Anthropology*, *5*(3), 87–98. doi: https://doi.org/10.1002/(SICI)1520-6505(1996)5:3<87::AID-EVAN4>3.0.CO;2-T

Loe, L. E., Bonenfant, C., Mysterud, A., Gaillard, J. M., Langvatn, R., Klein, F., … Stenseth, N. C. (2005). Climate predictability and breeding phenology in red deer: Timing and synchrony of rutting and calving in Norway and France. *Journal of Animal Ecology*, *74*(4), 579–588. doi: 10.1111/j.1365-2656.2005.00987.x

Lukas, D., & Clutton-Brock, T. (2017). Climate and the distribution of cooperative breeding in mammals. *Royal Society Open Science*, *4*(1). doi: 10.1098/rsos.160897

Lycett, J. E., Weingrill, T., & Henzi, S. P. (1999). Birth patterns in the Drakensberg Mountain baboons (Papio cynocephalus ursinus). *South African Journal of Science*, *95*(8), 354–356.

McNutt, J. W., Groom, R., & Woodroffe, R. (2019). Ambient temperature provides an adaptive explanation for seasonal reproduction in a tropical mammal. *Journal of Zoology*, *309*(3), 153–160. doi: 10.1111/jzo.12712

Pereira, R. J. P., Dos Santos Zanetti, E., & Furlan Polegato, B. (2010). Female reproduction. In J. M. Barbanti Duarte & S. Gonzalez (Eds.), *Neotropical Cervidology* (1st ed., Vol. 1). FUNEP/IUCN. Retrieved from https://www.researchgate.net/publication/273143129

Petersdorf, M., Weyher, A. H., Kamilar, J. M., Dubuc, C., & Higham, J. P. (2019). Sexual selection in

the Kinda baboon. *Journal of Human Evolution*, *135*(102635). doi: 10.1016/j.jhevol.2019.06.006

Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J. M., Tucker, C. J., & Stenseth, N. C. (2005). Using

the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in*

*Ecology and Evolution*. doi: 10.1016/j.tree.2005.05.011

R Core Team. (2019). *R: A language and environment for statistical computing*. Retrieved from

https://www.r-project.org/

Reitsema, L. J. (2012). Introducing Fecal Stable Isotope Analysis in Primate Weaning Studies.

*American Journal of Primatology*, *74*(10), 926–939. doi: 10.1002/ajp.22045

Reitsema, L. J., Partrick, K. A., & Muir, A. B. (2016). Inter-individual variation in weaning among

rhesus macaques (Macaca mulatta): Serum stable isotope indicators of suckling duration and

lactation. *American Journal of Primatology*, *78*(10), 1113–1134. doi: 10.1002/ajp.22456

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things).

*Methods in Ecology and Evolution*, *3*, 217–223. doi: 10.1111/j.2041-210X.2011.00169.x

Rhine, R. J., Norton, G. W., Wynn, G. M., & Wynn, R. D. (1989). Plant feeding of yellow baboons

(Papio cynocephalus) in Mikumi national park, Tanzania, and the relationship between seasonal

feeding and immature survival. *International Journal of Primatology*, *10*(4), 319–342. doi:

10.1007/BF02737420

Rogers, J., Muthuswamy, R., Harris, R. A., Mailund, T., Leppälä, K., Athanasiadis, G., … Worley, K.

C. (2019). The comparative genomics and complex population history of Papio baboons. *Science*

*Advances*, *5*, 1–14. Retrieved from http://advances.sciencemag.org/

Rosetta, L., Lee, P. C., & Garcia, C. (2011). Energetics during reproduction: A doubly labeled water

study of lactating baboons. *American Journal of Physical Anthropology*, *144*(4), 661–668. doi:

10.1002/ajpa.21475

Rutberg, A. T. (1987). Adaptive Hypotheses of Birth Synchrony in Ruminants: An Interspecific Test.

*The American Naturalist*, *130*(5), 692–710. doi: 10.1086/284739

Setchell, J. M., Lee, P. C., Wickings, E. J., & Dixson, A. F. (2002). Reproductive Parameters and

Maternal Investment in Mandrills (Mandrillus sphinx). *International Journal of Primatology*,

*23*(1), 51–68. doi: 10.1023/A:1013245707228

Spady, T. J., Lindburg, D. G., & Durrant, B. S. (2007). Evolution of reproductive seasonality in bears.

*Mammal Review*, *37*(1), 21–53. doi: 10.1111/j.1365-2907.2007.00096.x

Spinage, C. A. (1973). The role of photoperiodism in the seasonal breeding of tropical African

ungulates. *Mammal Review* , *3*(3), 71–85. doi: 10.1111/j.1365-2907.1973.tb00174.x

Struhsaker, T. T. (1997). Ecology of an African rain forest: logging in Kibale and the conflict between

conservation and exploitation. *Journal of Tropical Ecology*, *14*(4), 561–561. doi:

10.1017/S0266467498240399

Swedell, L. (2011). African papionins: diversity of social organization and ecological flexibility. In C.

J. Campbell, A. Fuentes, K. C. MacKinnon, S. Bearder, & R. Stumpf (Eds.), *Primates in*

*perspective* (Vol. 2, pp. 241–277). Oxford: Oxford University Press. Retrieved from

https://www.researchgate.net/publication/285789215

Thel, L., Chamaillé-Jammes, S., & Bonenfant, C. (2022). How to describe and measure phenology?

An investigation on the diversity of metrics using phenology of births in large herbivores. *Oikos*,

*2022*(4). doi: 10.1111/oik.08917

Thompson, M. E., & McCabe, G. M. (2013). Reproductive seasonality in wild Sanje mangabeys

(Cercocebus sanjei), Tanzania: Relationship between the capital breeding strategy and infant

survival. *Behaviour*, *150*(12), 1399–1429. doi: 10.1163/1568539X-00003102

Tinsley Johnson, E., Snyder-Mackler, N., Lu, A., Bergman, T. J., & Beehner, J. C. (2018). Social and

ecological drivers of reproductive seasonality in geladas. *Behavioral Ecology*, *29*(3), 574–588.

doi: 10.1093/beheco/ary008

Tonkin, J. D., Bogan, M. T., Bonada, N., Rios-Touma, B., & Lytle, D. A. (2017). Seasonality and

predictability shape temporal species diversity. *Ecology*, *98*(5), 1201–1216.

Trébouet, F., Malaivijitnond, S., & Reichard, U. H. (2021). Reproductive seasonality in wild northern

pig-tailed macaques (Macaca leonina). *Primates*. doi: 10.1007/s10329-021-00901-1

van Noordwijk, M. A. (2012). From maternal investment to lifetime maternal care. In *The evolution of*

*primate societies* (pp. 321–342). Chicago, London: The University of Chicago Press.

Van Schaik, C. P., Terborgh, J. W., & Wright, S. J. (1993). The phenology of tropical forests:

Adaptive significance and consequences for primary consumers. *Annual Review of Ecology and*

*Systematics*, *24*(1), 353–377. doi: 10.1146/annurev.es.24.110193.002033

van Schaik, C. P., & van Noordwijk, M. A. (1985). Interannual variability in fruit abundance and the

reproductive seasonality in Sumatran Long-tailed macaques (Macaca fascicularis). *J. Zool.,*

*Lond. (A)*, *206*(4), 533–549. doi: 10.1111/j.1469-7998.1985.tb03557.x

Wells, J. C. K., & Stock, J. T. (2007). The biology of the colonizing ape. *Yearbook of Physical*

*Anthropology*, *50*(1), 191–222. doi: 10.1002/ajpa.20735

Wiederholt, R., & Post, E. (2011). Birth seasonality and offspring production in threatened neotropical

primates related to climate. *Global Change Biology*, *17*(10), 3035–3045. doi: 10.1111/j.1365-

2486.2011.02427.x

Wright, P. C. (1999). Lemur Traits and Madagascar Ecology: Coping With an Island Environment.

*Yearbook of Physical Anthropology*, *42*, 31–72. doi: 10.1002/(sici)1096-

8644(1999)110:29+<31::aid-ajpa3>3.0.co;2-0

Zerbe, P., Clauss, M., Codron, D., Bingaman Lackey, L., Rensch, E., Streich, J. W., … Müller, D. W.

H. (2012). Reproductive seasonality in captive wild ruminants: Implications for biogeographical

adaptation, photoperiodic control, and life history. *Biological Reviews*, *87*(4), 965–990. doi:

10.1111/j.1469-185X.2012.00238.x

**FIGURES**

807
**Figure 1:** Distribution of the sampled species in Africa, and locations of the populations
considered in this study.

The species sampled in this study are depicted with relatively larger icons and species names, as well as full
coloured circles. Species for which we could not find birth seasonality data are represented with smaller icons and
names. The coloured areas on the map, corresponding to each species as indicated by the coloured circles, show
the geographical distribution of each species. Within each range, small circled numbers show the location of the
populations included in this study. 1: Taï, 2: Udzungwa Mountains, 3: Kibale, 4: Akfadou, 5: Tigounatine, 6:
Lékédi. 7: Moukalaba-Doudou. 8: Gashaka-Gumti. 9: Gilgil. 10: Queen Elizabeth. 11: Amboseli. 12: Mikumi. 13:
Tana River. 14: Filoha. 15: Kasanka. 16: De Hoop. 17: Drakensberg. 18: Moremi. 19: Tokai. 20: Tsaobis. 21:
Simien. The species distribution ranges and icons come from Julia Fischer et al., 2017; Kingdon et al., 2012.

**Figure 2**: Monthly distribution of births in relation to rainfall and NDVI seasonality

The proportion of births per month (left side of the y-axis) is represented with red bars. In addition, the darker red bar indicates the month of the mean birth date ($\mu_{birth}$) for seasonal breeding populations. We indicated in darker red the value of the r-vector length ($r_{birth}$) for each population (top-right corner of each panel). We represent the mean monthly rainfall (equalled to $K_{rain}$ + Rainfall S, in mm) in blue (right side of the y-axis), and the mean monthly NDVI (equalled to $K_{NDVI}$ + NDVI S divided by 2 for graphical purposes) in green (left-side of the y-axis). The species and population names are indicated on top of each panel, along with the number of births observed (N).

829
830 **Figure** 3: Phylogenetic tree of the studied species including the variation in the intensity of
831 reproductive seasonality for each species.

832 The intensity of birth seasonality is here quantified using the $r_{birth}$ value. For those species represented by more
833 than one population, the length of the black segment displays the variation in intensity of birth seasonality among
834 populations.

**Figure 4:** Effect of multiple environmental factors on the intensity of reproductive seasonality.

We plotted the intensity of reproductive seasonality ($r_{birth}$) depending on latitude (Panel A), environmental productivity (indexed by mean annual rainfall in mm, Panel B), the magnitude of environmental seasonality (i.e. of the rainfall peak, Panel C), the number of rainy seasons (Panel D), the rainfall peak breadth (RPB, Panel E), the

magnitude of environmental unpredictability (i.e. of variation in the non-seasonal component of rainfall (Panel F),
the timing of environmental unpredictability (i.e. between-year variation in rainfall timings, Panel G), and habitat
type (Panel H). For each panel, each dot represents a population (with the population name annotated), and the
colour indicates the species (see legend at the bottom). The dashed black line represents the linear regression, and
the shaded area displays 95% confidence intervals. On top of each panel, we indicated in italic and between
parentheses the significance of each predictor (NS for non-significant, * for P-values<0.05, ** for P-values<0.01).

# TABLES

**Table 1:** Hypotheses and predictions proposed on the effects of various environmental components on the intensity of reproductive seasonality ($r_{birth}$)

| Predictor | | Description | Hypothesis | Prediction | Results |
|---|---|---|---|---|---|
| Latitude | | Latitude, in degrees | H1.1 | When latitude increases, $r_{birth}$ increases | No |
| Environmental productivity | | Mean annual rainfall ($12 \times K_{rain}$) | H1.2 | When environmental productivity increases, $r_{birth}$ decreases | No (opposite effect) |
| Environmental seasonality | Magnitude of rainfall peak | Relative standard deviation of the seasonal component of rainfall (Rainfall S) | H1.3 | When the amount of within-year rainfall variation increases, $r_{birth}$ increases | No |
| | Modality of annual rainfall distribution | Number of rainy seasons (1 or 2) | H1.4 | When there is more than one rainy season, $r_{birth}$ decreases | No |
| | Breadth of the annual rainy season (broadness) | Rainfall peak breadth (RPB): minimum number of consecutive months with >80% of annual rainfall | H1.5 | When RPB increases, $r_{birth}$ decreases | No |
| Environmental unpredictability | Magnitude | Relative standard deviation of the non-seasonal component of rainfall (Rainfall NS) | H1.6 | When the amount of rainfall unpredictability (between-year variation) increases, $r_{birth}$ decreases | Yes |
| | Timing | Standard deviation of the yearly mean rainfall dates | H1.7 | When the level of rainfall unpredictability in terms of timing increases, $r_{birth}$ decreases | No |
| Habitat | | Type of habitat: tropical forest, open savannah, or mosaic forest-savannah | H1.8 | Populations living in open savannahs have lower $r_{birth}$ than the ones living in tropical forest | No |

$r_{birth}$ refers here to the r-vector length, i.e. to the intensity of reproductive seasonality

**Table 2**: Influence of several components of rainfall variation on the reproductive seasonality of African papionin populations

The table shows the posterior mean, the 95% HPD confidence intervals (CI) and the MCMC p-values (pMCMC) associated for each posterior distribution of the predictors of the Poisson MCMCglmm including species' relatedness as random effect, $r_{birth}$ as response variable (the r-vector length, transformed to be an integer value), and each predictor as the only fixed effect of a univariate model. Significant effects are indicated in bold. For categorical predictor, the tested category is indicated between parentheses.

| Predictor | | posterior mean | CI lower | CI upper | pMCMC |
|---|---|---|---|---|---|
| Latitude | | 0.11 | -0.49 | 0.67 | 0.686 |
| **Habitat productivity** | | **0.38** | **0.02** | **0.74** | **0.039** [+] |
| Magnitude of environmental seasonality | | 0.05 | -0.28 | 0.37 | 0.724 |
| Number or rainy seasons | (2)[†] | -0.19 | -0.96 | 0.60 | 0.617 |
| Rainfall peak breadth | | -0.06 | -0,52 | 0.45 | 0.741 |
| **Magnitude of environmental unpredictability** | | **-0.34** | **-0.62** | **-0.06** | **0.024** [+] |
| Timing of environmental unpredictability | | -0.18 | -0.46 | 0,09 | 0.184 |
| Habitat type | (Mosaic forest-savannah)[š] | 0.55 | -0.39 | 1.52 | 0.250 |
| | (Tropical forest) [š] | 1.11 | -0.63 | 2.47 | 0.152 |

[†] The reference category is 1 rainy season
[š] The reference category is open savannah habitat
[+] Note that when applying Bonferroni corrections for multiple tests, pMCMC>0.06 and is thus non-significant

**Table 3**: Reproductive stage matched with the food peak in sampled populations.

NAs are written when the alignment of the food peak with this reproductive stage was not tested (either because the data on weaning age in this population was missing, or because it is a non-seasonally breeding population, i.e. with P-value of the Rayleigh test >0.05). Xs are written when there is no alignment, while cells are shaded when there is an alignment of the given reproductive stage with the food peak.

| Species | Population | $r_{birth}$ | Reproductive stages match with food peak | | |
|---|---|---|---|---|---|
| | | | Conception | Lactation | Weaning |
| *Cercocebus atys* | Taï | 0.8312 | X | Early | X |
| *Cercocebus sanjei* | Udzungwu Mountains | 0.5588 | Before | Mid | Before |
| *Lophocebus albigena* | Kibale | 0.2762 | Before | Mid | X |
| *Macaca sylvanus* | Akfadou | 0.9350 | X | Early | NA |
| | Tigounatine | 0.9467 | X | X | NA |
| *Mandrillus sphinx* | Lekedi | 0.6766 | Before | Mid | X |
| | Moukalaba-Doudou | 0.7940 | X | Whole | NA |
| *Papio anubis* | Gashaka-Gumti | 0.2167 | NA | NA | NA |
| | Gilgil | 0.1873 | X | X | After |
| | Queen Elizabeth | 0.0971 | NA | NA | NA |
| *Papio cynocephalus* | Amboseli | 0.1344 | Before | X | X |
| | Mikumi | 0.1584 | X | X | NA |

| | Site | | | | |
|---|---|---|---|---|---|
| | Tana River | 0.1413 | NA | NA | NA |
| *Papio hamadryas* | Filoha | 0.0208 | NA | NA | NA |
| *Papio kindae* | Kasanka | 0.4971 | After | X | NA |
| *Papio ursinus* | De Hoop | 0.0969 | NA | NA | NA |
| | Drakensberg | 0.4143 | X | Early | NA |
| | Moremi | 0.3710 | Before | Mid | After |
| | Tokai | 0.2394 | NA | NA | NA |
| | Tsaobis | 0.1046 | NA | NA | NA |
| *Theropithecus gelada* | Simien | 0.1550 | X | X | X |