Cross-site harmonization of diffusion MRI data without matched training subjects

Alberto De Luca[1], Tine Swartenbroekx[2], Harro Seelaar[2], John van Swieten[2], Suheyla Cetin Karayumak[3], Yogesh Rathi[3], Ofer Pasternak[3], Lize Jiskoot[2], Alexander Leemans[1]

[1]Image Sciences Institute, University Medical Center Utrecht, Utrecht, the Netherlands
[2]Erasmus Medical Center, Rotterdam, the Netherlands
[3]Brigham and Women's Hospital, Harvard Medical School, Boston, USA

Abstract

Diffusion MRI (dMRI) data typically suffer of marked cross-site variability, which prevents naively performing pooled analyses. To attenuate cross-site variability, harmonization methods such as the rotational invariant spherical harmonics (RISH) have been introduced to harmonize the dMRI data at the signal level. A common requirement of the RISH method, is the availability of healthy individuals who are matched at the group level in terms of multiple demographics including age, sex etc. to learn harmonization features across sites while minimizing the impact of biological variabilities in this process. However, these subjects may not always be readily available when considering multiple independent cohorts with different population characteristics, particularly retrospectively.

To overcome these challenges, in this work we present a new approach to RISH harmonization that learns harmonization features while controlling for potential covariates using a voxel-based generalized linear model (RISH-GLM). By design, RISH-GLM allows to harmonize simultaneously data from any number of sites while also accounting for covariates of interest, thus not requiring matched training subjects. Additionally, RISH-GLM can harmonize data from multiple sites in a single step, whereas RISH is performed for each site independently. To demonstrate RISH-GLM, we considered data of training subjects from retrospective cohorts acquired with 3 different scanners. We performed 3 harmonization experiments of increasing complexity. First, we aimed to demonstrate that RISH-GLM is equivalent to conventional RISH when trained with data of matched training subjects. Secondly, we aimed to demonstrate that RISH-GLM can effectively learn harmonization with two groups of highly unmatched subjects. Thirdly, we evaluated the ability of RISH-GLM to simultaneously harmonized data from 3 different sites. Our results demonstrate that RISH-GLM can learn cross-site harmonization both from matched and unmatched groups of training subjects, and can effectively be used to harmonize data of multiple sites in one single step.

Introduction

Diffusion MRI (dMRI) has become a pivotal technique to investigate brain structure non-invasively[1]. The signal measured with dMRI originates from the motion of water molecules at the microscopic scale[2], which allows to infer microstructural properties of tissues when coupled with appropriate modelling[3]. For example, diffusion tensor imaging[4,5] (DTI), one of the most commonly applied dMRI techniques, provides metrics such as the mean diffusivity (MD) and the fractional anisotropy[4,6] (FA), which are related to the root mean square displacement and the anisotropy of the diffusion process, and reflect properties of the biological environment. Over time, these metrics have become established in the study of white matter microstructure in the study of neurological diseases[7], including Alzheimer's disease[8], small vessel disease[9,10] and frontotemporal dementia[11], among others.

A possible limitation to the use of dMRI in clinical research is that its measurements strongly depend on the employed MRI hardware and software[12,13]. As such, metrics derived from dMRI can typically be compared only within a single site, even when acquisition parameters are kept constant. In the latest years, several methods have been proposed to tackle such cross-site variability and harmonize dMRI. Broadly speaking, the goal of harmonization methods can be summarized as removing cross-site differences while not altering the sensitivity of dMRI to biological effects of interest. Two main families of post-processing dMRI harmonization methods have been proposed to date. The first family aims to remove batch effects during the analysis steps. Methods such as ComBat[14–16], for example, aim to remove cross-site differences on the final metrics derived from dMRI. As such, harmonization is applied independently to each considered dMRI metric, such as FA or MD. The second family of dMRI harmonization methods aims to remove cross-site effects on the acquired data before any quantification step. A popular example hereof is the rotational invariant spherical harmonics[17,18] (RISH), or recent deep learning alternatives[19,20]. These methods allow to harmonize dMRI data as part of the preprocessing pipeline. As such, data is harmonized only once, and can be then analyzed with any method of choice, as we have previously demonstrated in small vessel disease[21,22], for example. This second family of methods is the focus of this work, as it theoretically allows more freedom in subsequent analyses.

One of the challenges when applying dMRI harmonization methods such as RISH, is that they require healthy volunteer data. For example, the RISH method requires at least 16 training subjects matched at the group level for each site to be harmonized, or repeated scans of so-called travelling heads. This is particularly challenging for studies focusing on patient populations, where healthy controls may not always be recruited, or when retrospectively harmonizing multiple different cohorts with differences in average age or sex distribution. Furthermore, the concept of "matched" is ill-defined and very often interpreted as age and sex matching, but other factors such as education, presence of genetic mutations, and other factors that may affect the dMRI signal might need to be considered for specific studies.

In this study, we propose a generalization of the RISH framework that determines cross-site harmonization while also considering potential confounders. This new approach allows to control for potential biases when learning harmonization from unmatched groups of training subjects, and also allows to harmonize data from multiple sites in one single step. We demonstrate the potential of this framework through three experiments of growing complexity employing dMRI data of subjects from three different sites, while controlling for sex and age.

Methods

Theory: RISH and RISH-GLM

The conventional RISH harmonization method is based on the representation of diffusion MRI data acquired at a certain b-value (i.e., a diffusion "shell") with spherical harmonics. Spherical harmonics are a set of functions on the unit sphere characterized by an order and a degree. For each order i, rotational invariant representations (i.e., RISH coefficients) can be determined, and are indicated as L_i. Mirzaalian et al. and subsequent works[17,23] have shown that it is possible to harmonize dMRI data by learning the scaling factors between corresponding RISH coefficients determined in two groups of matched training subjects. The same framework can be also used to harmonize multi-shell data effectively by harmonizing each shell independently[18]. The harmonization essentially determines a voxel-wise scaling factor associated to each RISH feature to map a target site to a reference site. Mathematically, the scaling coefficient ϑ associated to the spherical harmonics of order i can be written as

$$\vartheta_i = {\sigma_{i,R}}/{\sigma_{i,T}}$$

where σ indicate the average RISH features of the reference (R) and target sites (T). Subsequently, dMRI datasets from the target site can be harmonized as follows: 1) dMRI data is converted to spherical harmonics; 2) their RISH features calculated and multiplied by the scaling factor ϑ_i; 3) harmonized dMRI data is reconstructed from the corrected spherical harmonics representation. The assumption behind this approach is that cross-site differences in σ purely originate from scanner-related properties. For this assumption to hold, the groups from which the σ values are computed must be matched across all factors that may affect the dMRI signal. Otherwise, systematic groups differences that are not accounted for, will be assimilated into the scaling factors, potentially biasing the factors and making them less generalizable to other subjects from these two sites.

To alleviate the matching requirement, we propose a new formulation based on a general linear model (GLM) that decomposes the effect of confounders such as age and sex on the RISH features:

$$\begin{bmatrix} \sigma_{i,R} \\ \sigma_{i,T} \end{bmatrix} = \begin{bmatrix} S_R & S_T & COV \end{bmatrix} \begin{bmatrix} \beta_{i,R} \\ \beta_{i,T} \\ \beta_{i,COV} \end{bmatrix}$$

where S are dummy variables assuming a value of 1 when a subject belongs to a certain site, and 0 otherwise. This approach, dubbed RISH-GLM, is based on two assumptions: 1) that the relationship between RISH features and covariates in the considered range is linear; 2) limited collinearity and interaction between different covariates. Of note, linearity in the relation between covariates and RISH-features, does not imply nor require linearity between covariates and diffusion metrics (e.g., fractional anisotropy). The RISH-GLM formulation allows to alleviate the need to match the average demographics at the group level across all sites. In practice, however, a reasonably even distribution of the effects of interest across sites is still needed to reliably estimate the associated coefficients β, from which the scaling coefficients can be determined as $\vartheta_i = {\beta_{i,1}}/{\beta_{i,2}}$. The formulation above can readily be extended to account for multiple sites by considering additional columns on the right side as follows:

$$\begin{bmatrix} \sigma_{i,R} \\ \sigma_{i,T1} \\ \vdots \end{bmatrix} = \begin{bmatrix} S_R & S_{T1} & \cdots & COV \end{bmatrix} \begin{bmatrix} \beta_{i,R} \\ \beta_{i,T1} \\ \vdots \end{bmatrix}$$

Data

We retrospectively collected data from FTD-RISC, which is a longitudinal study following up individuals at risk of frontotemporal dementia (FTD) curated at Erasmus Medical Center in Rotterdam (the Netherlands). In FTD-RISC, three major combinations of MRI hardware and software have been employed, which we refer to as Site 1, Site 2, Site 3. Data from all sites are used throughout this study to demonstrate the potential of GLM-RISH. Data of Site 1 was acquired with a 3T Philips Achieva scanner located at Leiden University Medical Center (the Netherlands) running on software release R5 with a 32ch head coil. Data of Site 2 was acquired with a GE Healthcare MR750 3T scanner located at Erasmus Medical Center (the Netherlands) with a 48ch head coil. Data of Site 3 was acquired with an older version of the scanner at Site 1, running on software release R4 and equipped with an 8ch head coil. For this study, only data of healthy control subjects and FTD-carriers who did not have FTD-related symptoms at the time of scanning were considered.

Experiments

We evaluated the performance of RISH-GLM as compared to conventional RISH with three experiments of growing complexity. We refer to RISH[matched/unmatched] and RISH-GLM[matched/unmatched] to indicate whether the methods were trained with matched or unmatched groups of participants from different sites, respectively. Table 1 shows an overview of the datasets used in the following three experiments.

Datasets	N (per site)	Age (years)	Sex (F)
Dataset 1	15 (site 1) vs 15 (site 2)	50.2 ± 12.2 vs 52.8 ± 11.1	47% vs 33%
Dataset 2	18 (site 1) vs 18 (site 2)	35.4 ± 6.1 vs 56.4 ± 8.5	44% vs 39%
Dataset 3	15 (site 1) vs 15 (site 2) vs 15 (site 3)	48.4 ± 10.2 vs 52.8 ± 10.8 vs 56.5 ± 8.9	40% vs 33% vs 53%
Dataset 4	15 (site 1) vs 15 (site 2) vs 15 (site 3)	52.3 ± 7.5 vs 55.5 ± 10.1 vs 50.5 ± 7.6	40% vs 40% vs 40%

Table 1: An overview of the datasets used to train RISH and RISH-GLM in the three experiments. N = sample size of each site. F = female

In experiment 1, we investigate whether RISH-GLM is equivalent to the conventional RISH method when trained on data of 15 individuals matched at the group level for both age and sex from Dataset 1. For each spherical harmonics' order, the scaling factors were computed with both RISH and RISH-GLM accounting for both age and sex, and visually compared. Subsequently, the training data from Site 2 was harmonized, and the diffusion tensor model was fit with REKINDLE[24] using MRIToolkit (https://github.com/delucaal/MRIToolkit). FA maps were computed and registered to a common space using the TBSS_PNL pipeline (https://github.com/pnlbwh/tbss). In short, this pipeline is based on the FSL TBSS[25] approach, but replaces the registration steps with ANTS[26] for additional spatial accuracy. Boxplots of FA computed on the white matter skeleton were compared before and after harmonization. Given FA is known to be strongly associated with age[27], we additionally visualized their relation by means of scatterplots, and evaluated their Pearson correlation coefficient before and after harmonization. As no ground truth exists for age and sex effects, we evaluated the agreement between the effects estimated by RISH-GLM on the whole datasets and age and sex effects within the individual sites by means of Pearson correlations. To estimate the effects within the individual sites, general linear models accounting for age, sex and intercept were estimated at the voxel-level for site 1 and site 2 independently.

Subsequently, in experiment 2, we evaluated the feasibility of removing scanner-related cross-site differences by learning harmonization with two age-unmatched groups of subjects from Dataset 2. RISH-GLM was trained by providing age and sex of each individual as covariates. As in the previous experiment, we compared the scaling factors corresponding to different RISH features, and visually inspected the spatial maps corresponding to age and sex effects. Harmonization was applied to the training set (Dataset 2). Boxplots of FA and scatterplots of FA as a function of age were derived as explained in experiment 1. In light of the large age difference between the two groups in Dataset 2, we anticipated to observe differences in FA between the two groups before harmonization. We expected such differences not to be completely removed after harmonization if the applied method truly removes only scanner-related differences and not biological effects. Furthermore, we expected a successful harmonization to recover a Pearson correlation coefficient between age and FA as similar as possible to the one determined in experiment 1. The same analyses were repeated on an independent dataset (Dataset 1) for validation purposes. We anticipated that RISH-GLM harmonization trained with age-unmatched subjects (Dataset 2) would generalize to age-matched subjects (Dataset 1), whereas RISH harmonization would not.

In experiment 3, we evaluated whether the RISH-GLM framework allows to effectively harmonize data from multiple sites from Dataset 3 in a single step. FA maps were calculated and aligned to a common space as previously explained. Next to boxplots of FA derived on the white matter skeleton, we additionally performed voxel-wise t-tests while correcting for age and sex to also evaluate the effectiveness of RISH-GLM at removing potential regional cross-site differences. Comparisons between the three sites were performed pair-wise (i.e., Site1 vs Site 2 and Site 3, Site 2 vs Site 3) using FSL *randomise* with threshold free cluster enhancement. Finally, to validate the one-step RISH-GLM harmonization on independent subjects, we applied the trained harmonization to age and sex-matched subjects of dataset 4, which shared only 2 out of 45 subjects with dataset 3. Here we evaluated the effectiveness of RISH-GLM at removing cross-site differences by means of boxplots of FA calculated on the white matter skeleton, and by evaluating the relation between FA and age as in the previous experiments.

Results

Experiment 1: RISH-GLM and RISH with matched training subjects (expect no differences)

Fig. 1 shows the scaling maps ϑ_i corresponding to spherical harmonics up to order 6 that were calculated with RISH and RISH-GLM using training subjects matched for age and sex at the group level from two different sites (hence the "[Matched]" suffix). The maps calculated by both methods are remarkably similar, as shown by the corresponding relative difference maps. The average relative differences between RISH[Matched] and RISH-GLM[Matched] are 0.38% for ϑ_0, 0.73% for ϑ_2, 1.32% for ϑ_4, and 1.45% for ϑ_6. While differences are globally less than 2%, local differences up to 20% can be appreciated in the bottom row of Fig. 1. Such differences are mostly located in areas of partial volume with cerebrospinal fluid or grey matter, but not in deep white not in white matter, and might originate from small imbalances in covariates or site-specific artefacts.

In dataset 1, age and sex effects estimated with RISH-GLM in white matter are generally in agreement with age effects calculated within site 1 and site 2 independently. The correlation coefficient between age effects estimated in the whole cohort and in site 1/2 are 0.80/0.74 for L0, 0.86/0.87 for L2, 0.62/0.83 for L4, 0.55/0.83 for L6, respectively. The correlation coefficient for sex effects estimated in the whole cohort and in site 1/2 are 0.75/0.70 for L0, 0.72/0.70 for L2, 0.53/0.89 for L4, 0.56/0.89 for L6, respectively.

Fig. 2 shows boxplots of average FA values of Site 1 and Site 2 obtained on the white matter skeleton before harmonization, and after harmonization with RISH and RISH-GLM. Before harmonization, a relative difference in average FA values between Site 1 and Site 2 equal to -9.7% (significant t-test, p≤0.05) can be observed, which is effectively removed by both RISH (0.5%, p=0.80) and RISH-GLM (1.7%, p=0.37). The second row of Fig. 2 shows the existence of a negative relation between age and FA within individual sites. When pooling unharmonized data together (first row), a biased correlation coefficient (rho) equal to -0.5 can be observed. After harmonization with both RISH and RISH-GLM, a linear negative relation between age and FA can be observed with equal correlation coefficients rho = -0.78. In both cases, the measurements from each site are well distributed around the regression line.

Fig. 1: Voxel-wise scaling maps calculated using group level-matched training subjects with RISH (first row), RISH-GLM (second row) and their difference (last row) for different orders of spherical harmonics (columns). Scaling maps calculated from both methods are globally similar, but local differences up to ±20% can be noticed particularly at the interface between grey matter and cerebrospinal fluid, and in the deep white matter for θ_4 and θ_6.

Fig. 2: The top row shows boxplots of average FA values per site, computed on the white matter skeleton, before harmonization (first column), after harmonization with RISH (middle column) and with RISH-GLM (last column). The second row shows scatterplots of the same average FA values as a function of age. Harmonization with both RISH and RISH-GLM was trained with group level-matched subjects.

Experiment 2: RISH-GLM and RISH with unmatched training subjects

Fig. 3 shows the scaling maps calculated for different orders of spherical harmonics with RISH and RISH-GLM trained with two groups of subjects that were not matched at the group level (hence the suffix "[Unmatched]"). In contrast to Fig. 1, where RISH and RISH-GLM produced similar scaling maps, large differences between the two methods can be observed when using unmatched training subjects, particularly for higher orders of spherical harmonics. Compared to the reference scaling maps calculated with RISH with matched subjects (Fig. 1), scaling maps calculated with unmatched subjects with RISH show large differences particularly around the ventricles, and at the interface between grey matter and cerebrospinal fluid for L0 and L2. For higher spherical harmonics orders (i.e., L4, L6), widespread differences in both the white and the grey matter can be observed. Compared to the reference (RISH[matched]), average relative differences of RISH[unmatched] and RISH-GLM[unmatched] are -10.2% and -3.3% for L0, -12.0% and -2.6% for L2, -10.5% and 2.9% for L4, -10.6% and 3.1% for L6, respectively, highlighting that scaling maps computed with RISH-GLM[unmatched] are on average closer to the reference(RISH[matched]) than RISH[unmatched].

Fig. 3: Voxel-wise scaling maps calculated using group age-unmatched training subjects with RISH (first row), RISH-GLM (second row) and their difference (last row) for different orders of spherical harmonics (columns). Compared to Fig. 1 (matched training groups), scaling maps calculated show different patterns with local differences up to ±50%.

Fig. 4: Age and sex effects estimated by RISH-GLM[Unmatched]. Age effects are prevalent at the interface between cerebrospinal fluid and white/grey matter for L0, and in the white matter for L2 and L4. The age effect map for L6 seems to be dominated by noise effects, given the low anatomical contrast, and the presence of clear stripes due to ghosting artefacts. Similar observations hold for sex effects.

Fig. 4 shows the spatial coefficients of age and sex effects associated to different spherical harmonics orders. Considering L0, it can be appreciated that age has a large effect around the ventricles and at the interface between grey matter and cerebrospinal fluid. At higher orders, age has a stronger effect on the brain white matter, particularly in the corpus callosum and corticospinal tracts. Similar observations also hold for sex effects. Importantly, both covariates have the largest effects in areas where RISH[matched] and RISH-GLM[unmatched] differ the most. In dataset 2, age and sex effects estimated with RISH-GLM in white matter are generally in agreement with age effects calculated within groups of Dataset 2 independently. The correlation coefficient between age effects estimated in the whole cohort and in site 1/2 are 0.58/0.80 for L0, 0.56/0.80 for L2, 0.25/0.90 for L4, 0.26/0.89 for L6, respectively. The correlation coefficient for sex effects estimated in the whole cohort and in site 1/2 are 0.66/0.68 for L0, 0.69/0.64 for L2, 0.50/0.81 for L4, 0.51/0.80 for L6, respectively.

In Fig. 5, we evaluated the average FA values calculated in the white matter skeleton of the unmatched groups. Before harmonization, differences in average FA values equal to -4.8% (p≤0.05) between the two sites can be observed. Importantly, such differences can originate from both scanner-specific differences as well as from true biological effects given the large age span. When looking at the relation between age and FA, no correlation is observed before harmonization. In this situation, applying RISH[Unmatched] while ignoring age and sex effects reduces the average FA difference between the two groups to 0.1% (p=0.95), but suppresses the expected difference between two groups with largely different age distributions. After harmonization with RISH-GLM[Unmatched], a relative difference in average FA values equal to 4.6% can be observed (p≤0.05), together with a linear negative relation between age and average FA that is in line with what was observed in Fig. 2 (correlation coefficient rho = -0.64 vs -0.78 in Fig. 2).

Subsequently, we have applied RISH[Unmatched] and RISH-GLM[Unmatched] to harmonize the data of the 2 matched groups of Dataset 1, for which no average difference in FA values is expected after harmonization. The boxplots shown in Fig. 6 showcase how RISH-GLM[Unmatched] can effectively remove the cross-site differences in FA between the two groups (average relative difference 0.5%, p=0.71), whereas RISH[Unmatched] does not (average relative difference -4.4%, p≤0.05). When looking at the relation between age and FA, the application of RISH-GLM[Unmatched] recovers the same age-FA relation observed also in Fig. 2 and Fig. 5.

Fig. 5: Boxplots of average FA values from two unmatched groups of healthy controls from Site1 and Site2, before harmonization, and after harmonization with RISH and RISH-GLM. Before harmonization, an average difference in FA between the two groups can be observed. While this could be plausible given the unmatched nature of the two groups, there is an unexpected lack of correlation between age and FA before harmonization. Harmonization with RISH[Unmatched] removes the difference between the two groups. Consequently, a mild negative correlation between age and FA is observed, but the datapoints from both sites are not well distributed around the regression line, indicating a biased fit. After harmonization with RISH-GLM[Unmatched], a difference between the two groups in average FA values is observed, as expected, as well as a clear negative relation between age and FA.

Fig. 6: Boxplots of average FA values from two matched groups from Site1 and Site2 before harmonization, and after applying RISH and RISH-GLM trained on unmatched data (Fig. 5). No differences in average FA values are observed after harmonization with RISH-GLM, as expected for two matched groups. The application of RISH-GLM also allows to reveal the same negative correlation between age and FA observed in the previous figures.

Experiment 3: One step harmonization

We investigated the ability of RISH-GLM to harmonize data from 3 sites in a single step. Example axial slices of the training scales derived with RISH and RISH-GLM between corresponding pairs of sites are shown in Supplementary Fig. S1. Scaling maps computed from both methods have similar appearance, it can be appreciated how the scaling computed with RISH-GLM are higher on average than those computed with RISH. The presence of ghosting artefacts can be appreciated in the scaling maps corresponding to L4 and L6, particularly with RISH-GLM. For this method, systematic artefacts can be learnt as part of cross-site differences, and can propagate to age and sex effect maps, as shown in Supp. Fig. 2. We subsequently evaluated differences in voxel-wise FA values between pairs of sites by means of permutations tests corrected by age and sex, obtaining the maps shown in Fig. 7. Before harmonization, 32.1%, 0.25% and 29.6% of the brain voxels were statistically different when comparing Site 1 to Site 2, Site 1 to Site 3, and Site 2 to Site 3, respectively. After harmonization, the amount of significantly different voxels was reduced to 0.1%, 0.1% and 0.4%, respectively.

To validate the method, we have applied RISH-GLM to harmonize the independent dataset 4. Results, which are reported in Fig. 8, demonstrate that RISH-GLM can effectively reduce the differences between all cohorts, and recover the same relation between age and FA previously observed in Fig. 2 and Fig. 5.

Fig. 7: Example axial (left) and coronal (right) slices showing differences in average FA values at the voxel-level between pairs of sites by applying a permutation test corrected for age and sex with a general linear model to three unmatched groups. Before harmonization, minimal to no differences are observed between Site1 and Site3. Conversely, extensive differences are observed between Site2 and the other sites. After harmonization with RISH-GLM in a single-step, most differences are effectively removed.

Training: multi-site with unmatched groups

Before harmonization   RISH-GLM[unmatched]

Fig. 8: A comparison of average FA values calculated on the white matter skeleton for three matched groups of subjects from Site1, Site2 and Site3 before and after harmonization in a single step with RISH-GLM. After harmonization, no statistically significant difference between the three sites can be observed, and the FA values from all sites are well distributed around the regression line in the correlation plot between age and FA.

Discussion

In this work, we have introduced a novel approach to perform RISH-based harmonization of the raw dMRI signal without need for training subjects matched at the group-level. Our approach does so by learning the voxel-wise effects of covariates on RISH scaling coefficients with a GLM (RISH-GLM). Our results demonstrate that RISH-GLM is effective in learning cross-site harmonization from unmatched training groups and can be used to harmonize data from multiple sites effectively in a single step.

The advent of dMRI harmonization methods such as RISH allows studies to overcome the barriers of single-site analyses, unharnessing the potential of retrospectively collected data to boost sample sizes. However, the applicability of such frameworks can be challenging when considering multiple cohorts selected with different inclusion criteria, because their requirement of matched training. Importantly, the results shown in Fig. 5 and Fig. 6 demonstrate that conventional RISH is unsuitable to learn cross-site harmonization with unmatched groups, as that would lead to a removal of both cross-site and biological

effects (e.g., aging, Fig. 5), or to the introduction of biases (Fig. 6). RISH-GLM can effectively tackle this limitation by lifting the need for matched training groups, as demonstrated in Fig. 6 and Fig. 8.

The ability to learn cross-site harmonization without matched cohorts could support several efforts in both clinical and neuroscientific research. Firstly, RISH-GLM could simplify the harmonization of cohorts with only partly overlapping covariates of interest. This could be for example age, in the case of studies pooling retrospective cohorts with distinct age ranges to cover the whole lifespan. Secondly, it could support further disentangling scanner effects from confounding biological effects than conventional RISH, particularly when learning harmonization from groups which are matched on average but feature relevant variance in covariates. In the study of dementia, for example, population cohorts, memory clinic cohorts, and population with different injury aetiologias could be potentially harmonized while accounting for key factors that could otherwise bias the harmonization process, such as differences in education, ethnicity, sex distributions, exposure to specific risk factors, lesion burden, etc.

While retrospective data is arguably the main application goal of post-processing dMRI harmonization methods such as RISH-GLM, they could prove advantageous also to support prospective studies. In general, accurate matching of acquisition hardware and sequence parameters can already reduce cross-site dMRI variability, as shown by previous initiatives in frontotemporal[28] and vascular dementia[29]. Yet, subtle differences between vendors, calibration, and characteristics of individual scanners might still dictate the existence of cross-site differences. Furthermore, even when prospective matching is achieved successfully at the start of a study, subsequent hardware changes and software updates might still introduce acquisition effects in long multiyear prospective studies. In the future, software versions and coil updates could be considered as covariates of interest in RISH-GLM to attempt limiting their effect on the acquired data.

It is important to acknowledge some limitations of this study. As explained in the methods section, RISH-GLM is based on assumptions that might not always hold true. An important assumption is the existence of a linear relation between covariates of interest and RISH features, at least within the range of covariates used for training. In the case of age, for example, little is known about its exact relation to RISH features. While validating this assumption throughout the lifespan will require dedicated studies, we observe that RISH-GLM is effective at recovering consistent relations between age and FA throughout the experiments of this work. Of note, the assumption of linearity between covariates and RISH features does not imply the need for linearity between covariates and derived metrics such as FA, given that RISH features are non-linear descriptors of the diffusion signal. This is particularly relevant as previous studies[27,30] of diffusion tensor imaging metrics across the lifespan have demonstrated the existence of a quadratic – locally linear – relation between age and fractional anisotropy, for example. Another assumption of RISH-GLM is that the effects of covariates is similar across training cohorts. As a consequence, training directly with non-homogeneous groups – such as patients with different pathologies – is discouraged. In this study, we used data from a retrospective cohort of individuals that were initially marked as at risk for frontotemporal dementia, which might have introduced a selection bias towards effects not explicitly accounted for. Furthermore, we only focused on modelling the effect of age and sex, and did not consider other relevant confounders such as education.

In conclusion, we have introduced a novel framework to learn cross-site harmonization while accounting for covariates of interests, and demonstrated its effectiveness to learn harmonization between two age unmatched groups with an average age difference of over two decades.

References

1. Le Bihan D, Johansen-Berg H. Diffusion MRI at 25: Exploring brain tissue structure and function. *Neuroimage*. 2012;61(2):324-341. doi:10.1016/j.neuroimage.2011.11.006

2. Stejskal EOO, Tanner JE. Spin Diffusion Measurements: Spin Echoes in the Presence of a Time-Dependent Field Gradient. *J Chem Phys*. 1965;42(288):288-292. doi:http://dx.doi.org/10.1063/1.1695690

3. Novikov DS, Kiselev VG, Jespersen SN. On modeling. *Magn Reson Med*. 2018;79(6):3172-3193. doi:10.1002/mrm.27101

4. Basser PJ, Pierpaoli C. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. *J Magn Reson B*. 1996;111:209-219. doi:10.1016/j.jmr.2011.09.022

5. Basser PJ, Mattiello J, LeBihan D. MR diffusion tensor spectroscopy and imaging. *Biophys J*. 1994;66(1):259-267. doi:10.1016/S0006-3495(94)80775-1

6. Pierpaoli C, Basser PJ. Toward a quantitative assessment of diffusion anisotropy. *Magn Reson Med*. 1996. http://onlinelibrary.wiley.com/doi/10.1002/mrm.1910360612/full. Accessed March 6, 2014.

7. Tae WS, Ham BJ, Pyun SB, Kang SH, Kim BJ. Current Clinical Applications of Diffusion-Tensor Imaging in Neurological Disorders. *Journal of Clinical Neurology*. 2018;14(2):129. doi:10.3988/jcn.2018.14.2.129

8. Chen Y, Wang Y, Song Z, Fan Y, Gao T, Tang X. Abnormal white matter changes in Alzheimer's disease based on diffusion tensor imaging: A systematic review. *Ageing Res Rev*. 2023;87:101911. doi:10.1016/j.arr.2023.101911

9. Duering M, Biessels GJ, Brodtmann A, et al. Neuroimaging standards for research into small vessel disease—advances since 2013. *Lancet Neurol*. 2023;22(7):602-618. doi:10.1016/S1474-4422(23)00131-X

10. de Luca A, Biessels GJ. Towards multicentre diffusion MRI studies in cerebral small vessel disease. *J Neurol Neurosurg Psychiatry*. 2021;0:jnnp-2021-326993. doi:10.1136/jnnp-2021-326993

11. Jiskoot LC, Bocchetta M, Nicholas JM, et al. Presymptomatic white matter integrity loss in familial frontotemporal dementia in the GENFI cohort: A cross-sectional diffusion tensor imaging study. *Ann Clin Transl Neurol*. 2018;5(9):1025-1036. doi:10.1002/acn3.601

12. Tax CM, Grussu F, Kaden E, et al. Cross-scanner and cross-protocol diffusion MRI data harmonisation: A benchmark database and evaluation of algorithms. *Neuroimage*. 2019;195(January):285-299. doi:10.1016/j.neuroimage.2019.01.077

13. Ning L, Bonet-Carne E, Grussu F, et al. Cross-scanner and cross-protocol multi-shell diffusion MRI data harmonization: Algorithms and results. *Neuroimage*. 2020;221(October 2019). doi:10.1016/j.neuroimage.2020.117128

14. Fortin JP, Cullen N, Sheline YI, et al. Harmonization of cortical thickness measurements across scanners and sites. *Neuroimage*. 2018;167(November 2017):104-120. doi:10.1016/j.neuroimage.2017.11.024

15. Fortin JP, Parker D, Tunç B, et al. Harmonization of multi-site diffusion tensor imaging data. *Neuroimage*. 2017;161:149-170. doi:10.1016/j.neuroimage.2017.08.047

16. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics*. 2007;8(1):118-127. doi:10.1093/biostatistics/kxj037

17. Cetin Karayumak S, Bouix S, Ning L, et al. Retrospective harmonization of multi-site diffusion MRI data acquired with different acquisition parameters. *Neuroimage*. 2019;184(September 2018):180-200. doi:10.1016/j.neuroimage.2018.08.073

18. De Luca A, Karayumak SC, Leemans A, et al. Cross-site harmonization of multi-shell diffusion MRI measures based on rotational invariant spherical harmonics (RISH). *Neuroimage*. 2022;259(February):119439. doi:10.1016/j.neuroimage.2022.119439

19. Nath V, Parvathaneni P, Hansen CB, et al. Inter-Scanner Harmonization of High Angular Resolution DW-MRI Using Null Space Deep Learning. *Math Vis*. 2019;(226249):193-201. doi:10.1007/978-3-030-05831-9_16

20. Moyer D, Ver Steeg G, Tax CMW, Thompson PM. Scanner invariant representations for diffusion MRI harmonization. *Magn Reson Med*. 2020;84(4):2174-2189. doi:10.1002/mrm.28243

21. de Brito Robalo BM, de Luca A, Chen C, et al. Improved Sensitivity and Precision in Multicentre Diffusion MRI Network Analysis Using Thresholding and Harmonization. *SSRN Electronic Journal*. 2022;36(October):103217. doi:10.2139/ssrn.4077955

22. de Brito Robalo BM, Biessels GJ, Chen C, et al. Diffusion MRI harmonization enables joint-analysis of multicentre data of patients with cerebral small vessel disease. *Neuroimage Clin*. 2021;32(November):102886. doi:10.1016/j.nicl.2021.102886

23. Mirzaalian H, Ning L, Savadjiev P, et al. Inter-site and inter-scanner diffusion MRI data harmonization. *Neuroimage*. 2016;135:311-323. doi:10.1016/j.neuroimage.2016.04.041

24. Tax CMW, Otte WM, Viergever MA, Dijkhuizen RM, Leemans A. REKINDLE: Robust extraction of kurtosis INDices with linear estimation. *Magn Reson Med*. 2015;73(2):794-808. doi:10.1002/mrm.25165

25. Smith SM, Jenkinson M, Johansen-Berg H, et al. Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *Neuroimage*. 2006;31(4):1487-1505. doi:10.1016/j.neuroimage.2006.02.024

26.   Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage*. 2011;54(3):2033-2044. doi:10.1016/j.neuroimage.2010.09.025

27.   Yeatman JD, Wandell BA, Mezer AA. Lifespan maturation and degeneration of human brain white matter. *Nat Commun*. 2014;5. doi:10.1038/ncomms5932

28.   Kalra S, Müller HP, Ishaque A, et al. A prospective harmonized multicenter DTI study of cerebral white matter degeneration in ALS. *Neurology*. 2020;95(8):E943-E952. doi:10.1212/WNL.0000000000010235

29.   Wilcock D, Jicha G, Blacker D, et al. MarkVCID cerebral small vessel consortium: I. Enrollment, clinical, fluid protocols. *Alzheimer's and Dementia*. 2021;17(4):704-715. doi:10.1002/alz.12215

30.   Lebel C, Walker L, Leemans A, Phillips L, Beaulieu C. Microstructural maturation of the human brain from childhood to adulthood. *Neuroimage*. 2008;40(3):1044-1055. doi:10.1016/j.neuroimage.2007.12.053

Supplementary Figures

Figure S1: Scaling maps calculated between pairs of sites with RISH, and in one single step with RISH-GLM on all three sites considered in Experiment 3.

Figure S2: Effect of covariates on RISH features of different orders as determined with RISH-GLM on data from all three sites in Experiment 3.