Cryo-EM Reveals the Mechanochemical Cycle of Reactive Full-length Human Dynein-1

Pengxin Chai[1,5], **Jun Yang**[1,5], **Indigo C. Geohring**[2], **Steven M. Markus**[1,*], **Yue Wang**[1,3,4,5,*], and **Kai Zhang**[1,*]

[1]Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, 06511, USA
[2]Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins, Colorado, USA
[3]Innovation Center for Brain Medical Sciences of the Ministry of Education, Huazhong University of Science and Technology, Wuhan, 430030, China
[4]Huazhong University of Science and Technology, Wuhan, 430030, China
[5]These authors contributed equally
[*]Correspondence: Email: steven.markus@colostate.edu (S.M.), yue.wang@yale.edu (Y.W.), jack.zhang@yale.edu (K.Z.)

Abstract

Dynein-mediated cargo transport plays pivotal roles in diverse cellular activities. Central to all these cellular processes is dynein's ATP-dependent mechanochemical cycle. Here, we perform a systematic cryo-electron microscopic investigation of the conformational landscape of full-length human dynein-1 actively undergoing nucleotide-induced conformational changes, both on and off microtubules. Our approach reveals over 30 high-resolution structures, categorized into nine distinct states of the AAA+ motor ring. This provides a dynamic and comprehensive view of dynein throughout its mechanochemical cycle. The novel intermediate states unveil a series of important mechanistic insights into dynein function, including a 'backdoor' phosphate release mechanism that coordinates linker straightening, a stepwise structural basis for the bidirectional communications between ATP binding/hydrolysis and microtubule-binding affinity, and the mechanism by which microtubule binding increases ATPase activity. Our findings also lead to a substantially revised model for the force-generating powerstroke, and clarify the means by which dynein achieves its unidirectional stepping on microtubules. These results substantially enhance our understanding of dynein function and provide a more complete model of its mechanochemical cycle.

Introduction

Dyneins are large cytoskeletal motor complexes that move toward the minus ends of microtubules (MTs) *(1)*. Cytoplasmic dynein-1 is the sole cytoplasmic isoform that drives retrograde intracellular cargo transport *(2)*, organelle positioning *(3)*, and cellular morphogenesis *(4)*, while dynein-2 and axonemal dyneins mediate intraflagellar transport *(5)* and ciliary motility *(6)*. Unlike the kinesin family which encompasses 45 different types of plus end-directed motors in humans *(7)*, a single cytoplasmic dynein-1 (dynein) is responsible for transporting diverse cellular cargos including membrane-bound organelles, RNAs, proteins, and viruses *(2)*. This is enabled by the processive retrograde transport machinery that includes dynein, dynactin, and a cargo adaptor (DDA complex) *(8–15)*, and is further regulated by cofactors like LIS1 *(16–21)* and NDEL1 *(22–24)*. Owing to its fundamental roles in a variety of cellular activities, mutations in dynein and its cofactors have been implicated in many human diseases, including neurodegenerative *(25)* and neurodevelopmental disorders *(26–28)*, viral infections *(29)*, and cancers *(30)*.

The dynein complex comprises two copies of the heavy chain (DHC) and several accessory chains. Each DHC consists of a tail domain, responsible for recruiting accessory chains, and a motor domain essential for motility *(2, 31–33)*. The motor domain includes an AAA+ ring that consists of six non-identical AAA+ domains, as well as a linker, coiled-coil stalk, and buttress appendages, a microtubule-binding domain (MTBD) located at the tip of the stalk, and a C-terminal domain (CTD). Importantly, the motor domain possesses ATPase activity and is responsible for converting the chemical energy stored in ATP into mechanical processes, collectively termed the mechanochemical cycle *(31)*.

The mechanochemical cycle of dynein is a fundamental process that underlies all dynein-mediated cellular activities. Our current understanding of this cycle primarily stems from biochemical *(34–37)*, biophysical *(38–41)*, and structural studies *(31, 42–47)* conducted on dynein motor domain fragments. Within the motor domain, AAA1 serves as the primary site for ATP hydrolysis *(35, 37)*. Throughout the catalysis cycle, the AAA+ ring adopts alternating closed and open conformations that are critical for force generation and microtubule-based motility *(31, 48)*. The linker domain can change from a bent to a straight conformation in a nucleotide-dependent manner, which has been deemed responsible for the force-generating "powerstroke" *(36, 40, 42, 49, 50)*. Both kinetic and structural evidence indicate that the release of phosphate (Pi) that is coupled with the opening of the AAA+ ring, and the AAA1 pocket specifically, triggers the straightening of the linker *(31, 36, 40, 48)*. However, the mechanism by which Pi is released from AAA1 remains elusive. Does it follow a "molecular backdoor" mechanism akin to myosin *(51)* and actin *(52)* which only requires minimal local conformational change, or does it require complete pocket opening? Furthermore, the correlation between Pi release, pocket opening, and linker straightening remains poorly understood *(31)*.

In contrast to the other two cytoskeletal motor proteins, kinesin and myosin, which bind to their respective tracks with their ATP hydrolysis domains *(51, 53)*, the MTBD of dynein

is 24 nm away from the AAA1 domain *(32)*. The nucleotide-bound states and pocket conformations of AAA1 have been shown to regulate the MTBD affinity for MTs *(42, 44, 45)*. Meanwhile, the MTs provide a track for the unidirectional stepping of dynein *(39, 41, 54)* and stimulate the AAA1 ATPase activity *(44, 55)*. The communication between AAA1 and the MTBD is mediated by the 15 nm-long stalk coiled-coil (stalk-CC) *(56–58)* which can adopt multiple registries corresponding to different MT-binding affinities *(38, 59–63)*. The registry change can be regulated by both AAA1 pocket conformations *(31)* and MT-binding *(57, 64, 65)*. However, due to the limited structural information available for dynein on and off MTs, the precise mechanism of communication between dynein and MTs remains unclear. Therefore, how dynein's mechanochemical cycle is accelerated by MTs *(66)*, and how dynein achieves unidirectional motility along MTs during the mechanochemical cycle *(54)* requires further investigation.

Previous studies have elucidated the structures of truncated dynein *(42, 44–47)* full-length "phi" dynein-1 *(10)*(similar to the Greek letter "Φ" *(67)*), phi dynein-2 *(68)*, outer-arm dynein (OAD)-MT *(69)*, and the DDA-MT *(15, 70)* complex, each captured under specific nucleotide conditions. Due to either limited resolution, or the loss of structural dynamics that is inherent to these conventional approaches, these structures provide valuable but limited insights into the atomic-level details of the mechanochemical cycle of full-length dynein. To overcome these limitations, here we employed strategies to identify the complete conformational landscape of full-length dynein progressing through its natural mechanochemical cycle. Firstly, we directly imaged full-length dynein in the presence of ATP (to visualize dynein "in reaction") as well as other nucleotide conditions, which allowed us to obtain structures of phi dynein and novel intermediate states from open/uninhibited dynein with much-improved resolution (2.2-3.6 Å). Furthermore, we imaged reconstituted MT-bound dynein complexes in different nucleotide conditions and obtained structures of the motor domains in four different states at high resolution (2.9-3.6 Å). This collection of motor domain structures (over 30) provides critical insights into several longstanding questions in the field. Our results reveal a "backdoor" channel through which Pi is released from AAA1, a mechanism to account for linker straightening, and how communication between AAA1 and the MTBD occurs, which we support with mutagenesis and functional studies. Comparison of dynein motor domains on and off MTs reveals the structural basis for enhanced ATPase activity upon MT-binding, the role of linker-AAA+ ring interactions in promoting unidirectional stepping, and a revised model for the dynein powerstroke. In summary, our comprehensive structural analysis of full-length dynein coupled with functional investigations significantly advances our understanding of the dynein mechanochemical cycle.

Results

Structural determination of full-length human dynein-1 off and on MTs

To explore the conformational landscape of dynein throughout its mechanochemical cycle ("in reaction"), we purified full-length human dynein-1 from insect cells (which had a roughly equal mixture of phi and open states) and prepared cryo-EM grids in the presence of 5 mM ATP (**Figure 1A, S1A**). We collected a large cryo-EM dataset and performed extensive in-silico 3D classification analysis followed by high-resolution refinement (**Figure S1B, Table S1-S5**). We obtained high-resolution cryo-EM maps of full-length phi dynein (**Figure 1C, S1B, Movie S1**), with the motor domain at 2.2 Å average resolution, and the majority of the tail region better than 4 Å. This significantly improved resolution compared to the previous one *(10)* allowed us to build a full atomic model for phi dynein (**Figure 1C**). The high-resolution structure of the motor domain provides unprecedented details, including holes within aromatic side chains, water molecules, and ions interacting with the nucleotide (**Figure 1F**). In addition to the phi dynein motor, we observed 10 distinct conformational states of the open motor domain, revealing snapshots of reactive intermediates during dynein's mechanochemical cycle (**Figure 1D**).

High-resolution structural determination of MT-bound dynein has been a challenge in the field. To address this, we recently developed a micrograph-based MT-signal subtraction method *(71, 72)*, leading to the successful reconstruction of OAD-MT and DDA-MT structures *(15, 69, 70)*. To capture complete MT-bound dynein-1 structures, we reconstituted full-length dynein-1 on microtubules in apo, ADP, and AMPPNP conditions, all of which increase dynein affinity for MTs *(73)*(**Figure 1B, S2, Table S6**). We obtained a structure of a two-headed full-length dynein bound to MTs at a resolution of 10 Å (**Figure 1E, S2**). Focused classification and refinement of the individual motor domains resulted in 4 distinct MT-bound motor domain structures at 2.9-3.6 Å (**Figure 1E, 1F, Movies S2**).

Overview of the conformational landscape of microtubule-unbound dynein

By focusing on the following four structural features, we obtained 11 distinct motor domain structures from phi and open dynein: (1) the pocket conformations of AAA1, (2) and of AAA3, (3) the linker position, and (4) the buttress-stalk conformations (**Figure S1, S3**). By primarily focusing on AAA1 and the overall AAA+ ring arrangement, these 11 structures were further categorized into 9 major states (state-0 to state-8) (**Figure 2A**). Among the 9 states, 5 of them are novel: states-0, -3, -4, -5, and -6. Below we describe in detail the salient features of each state, which are also summarized in **Table 1**.

Fig. 1. Cryo-EM analysis of full-length human dynein-1 off and on microtubules. **(A,B)** Schematic of cryo-EM pipeline depicting dynein in reaction with ATP without MTs (A) and dynein with MTs in different nucleotide conditions (B). **(C)** Cryo-EM reconstruction of phi dynein with the 3.8-7.0 Å tail region and 2.2 Å motor domain. **(D)** Representative cryo-EM reconstructions of the motor domain in different states from open dynein. The motor ring, linker, buttress, and stalk are colored grey, purple, orange, and yellow, respectively. **(E)** Cryo-EM map of full-length two-headed dynein bound to MTs in ADP condition at 10 Å resolution, with locally refined motor domains at 2.9-3.6 Å resolution. **(F)** High-resolution features are shown by representative density of amino acid residues and ligands.

State-0: The AAA+ ring in this state is not sealed and lacks clear density for AAA6 and the CTD likely due to these regions adopting highly flexible conformations **(Figure 2A, S3)**. AAA1 is in an apo state with a wide-open nucleotide-binding pocket, while the buttress and stalk are in a conformation that corresponds to the MTBD being in a low MT-binding affinity state. This distinct motor domain conformation potentially represents a novel mode of autoinhibition in which the motor is in a mechanochemical-incompetent state and may explain, at least in part, the weak processivity of open dynein in the absence of dynactin and a cargo adaptor *(10)*.

States-1, -2, -3, and 4: These motor domains all have a closed AAA1 pocket and are in low-MT binding affinity states, as determined by the buttress and stalk conformations

(31) **(Figure 2A)**. State-1 (phi-motor) has the highest resolution (2.2 Å), enabling accurate modeling and refinement of ADP, magnesium, and water molecules at the active site **(Figure 2B, 2C)**. States-2, -3, and -4, resolved at 2.8 Å, 3.0 Å, and 2.8 Å, respectively, all have clear ADP-Mg^{2+} density **(Figure S3)**. Unlike the autoinhibited state-1, states-2, -3, and -4 are actively engaged in the mechanochemical cycle and have different linker conformations. Specifically, state-2 possesses a bent linker that is stably docked between AAA2 and 3, similar to state-1 and previously reported pre-powerstroke motor structures (PDB: 4rh7 *(42)*, 5nug *(10)*, 6rla *(68)*). State-4 has a straight "hinge helix", which connects the N- and C-terminal regions of the linker, and an overall straight linker. However, the linker does not appear to be stably docked due to the closed AAA+ ring configuration in which the docking site at AAA5 is not exposed. The straight and undocked

linker and the apparent low MT binding affinity in state-4 suggest that this state is an intermediate during the dynein powerstroke. Finally, the hinge helix in state-3 adopts a bent conformation, while the rest of the linker region is flexible, indicating that state-3 represents an averaged map of many transient conformations between state-2 and state-4.

States-5, -6, and -7: All of these dyneins have ADP-bound to open AAA1 pockets, and account for about 60% of motor states during the active ATP cycle, indicating that ADP release from AAA1 pocket is rate-limiting. All three of these states exhibit varying degrees of pocket opening that are correlated with different buttress-stalk conformations (**Figure 2D**), consistent with the model whereby AAA1 pocket dynamics regulate MT affinity through the stalk-CC *(31, 42, 44, 45)*. State-7 has a post-powerstroke linker that is straight and stably docked in what we previously described as "post-2" *(69)* (in which the linker is upraised toward AAA3-AAA4), and a "kinked" buttress. The AAA1 pocket and AAA+ ring conformation are similar to dynein structures with high MT-binding affinities in previous (PDB: 7z8g *(15)*, 8fcy *(21)*, 4w8f *(47)*) and this work. These observations suggest that state-7 represents a conformation primed for MT-binding, or that the high MT-binding affinity state can be intrinsically adopted by the motor even in the absence of MTs. States-5 and -6 are two novel states that have intermediate open AAA1 pockets and overall straight but undocked linkers, which likely represent intermediates during the powerstroke (**Figure 2A**). The buttress in these two states exhibits an intermediate kink coupled with a flexible stalk-CC density, likely reflecting an intermediate MT-binding affinity. The stalk-CC is likely undergoing helix sliding *(38)* or local melting *(74)* in these two states, adopting a semi-α *(57)* or γ registry *(75)*. Comparison of these three states (-5, -6, and -7) to a previously reported ADP dynein-1 structure (PDB:3vkg *(44)*) reveals close alignment of the AAA1 pocket and linker position with state-7. However, the buttress-stalk conformation is most similar to the intermediate state-6, suggesting that 3vkg represents a further intermediate state between state-6 and -7.

State-8: State-8, which has a nucleotide-free AAA1 pocket, is transient in the presence of excess ATP, but can be obtained in the apo condition. The large and small subunits of the AAA1 are maximally separated in a wide-open state that permits ADP release (**Figure 2D**), similar to a previous yeast apo motor structure (PDB:4aki *(45)*) and OAD-MT structures (PDB: 7k58,7k5b *(69)*). The linker is docked in the canonical post-powerstroke "post-1" position, and the buttress-stalk is in a high MT-binding affinity conformation, indicating that state-8 is also primed for MT-binding.

Taken together, the range of conformational states we have observed for the full-length dynein-1 "in reaction" reflects not only the conformations observed for dynein structures solved in various nucleotide conditions but also novel intermediates never seen before (**Table 1**). Importantly, we observed that the linker, in addition to the previously described bent/docked conformation (pre-powerstroke) or straight/docked conformation (post-powerstroke), can adopt a straight but undocked conformation, likely representing crucial intermediate states during the powerstroke. These novel states demonstrate the importance of our approach and reveal novel insight into the mechanochemistry of the full-length dynein complex.

The effect of nucleotide conditions on the conformational distributions of full-length dynein

Given the range of conformations we observed for the dynein motor, even when bound to a given nucleotide (e.g., states-1 through 7 all have AAA1-bound ADP), we wondered whether incubating dynein with different nucleotide analogs would lock the motor in a single conformation, as has been noted in prior cryo-EM and crystallographic studies *(15, 21, 42, 44, 45, 47, 69, 76)*. To test this, we incubated full-length dynein with or without different analogs (i.e., AMPPNP, ADP, ATP-Vi, and apo) and performed cryo-EM classification (**Table S2-S5**). To our surprise, we found that multiple conformational states are present in each condition (**Figure 2E, S4**). It has been reported that dynein-2 adopts a closed AAA1 and a bent linker in the presence of ATP-vanadate (ATP-Vi; a mimic for ADP-Pi) *(42)*. Although we observed a significantly increased population of motors with closed AAA1 pocket states (states-1 through -4), we noted the linker could exist in both bent (states-1 and -2) and straight conformations (state-4), which is consistent with previous observations of dynein-c *(49, 77)*. As expected, in ADP and apo conditions, the proportion of molecules with open AAA1 pockets increased, but we also observed states with closed AAA1 pockets. Of note, in the apo condition, the wide-open apo AAA1 pocket conformation (state-8), which was very transient in the presence of ATP, was observed and co-existed with closed AAA1 states. With the exception of state-3, all states in ATP are also found in the AMPPNP condition.

We next determined the identity of the bound nucleotide in AAA1 for those dyneins with closed AAA1 pockets (states-1 to 4) under different conditions (**Figure 2F, S5**). In the presence of ATP, AAA1 is in the post-ATP hydrolysis state with a clear density for ADP-Mg^{2+}. With ATP-Vi, AAA1 exhibits density for ADP-Vi-Mg^{2+}, consistent with prior studies *(21, 42)*. This suggests that the phi-motor (state-1), despite being in an autoinhibited state, is capable of ATP hydrolysis and Pi release, the latter of which precedes Vi substitution. The non-hydrolyzable ATP analog AMPPNP has been widely used to lock cytoskeletal motors in an ATP-bound state *(78, 79)*. Surprisingly, in the presence of AMPPNP, the closed AAA1 pocket in both states-1 and -2 show clear density for ADP instead of AMPPNP, suggesting AMPPNP has a low affinity for AAA1. In both ADP and apo conditions, the closed AAA1 pocket has the same ADP-Mg^{2+} density as dynein in the ATP condition, indicating that the AAA1 pocket of full-length dynein can adopt a closed conformation even with low concentrations of ATP *(10)*. Taken

A

Full-length dynein-1 in ATP buffer

State-1
State-2
State-3
Hinge point
State-4
State-5
State-6
State-7
State-8
State-0

Auto-inhibited   Active Cycle

B

Phi-motor (state-1), AAA1, 2.2Å

AAA1S   AAA2L
AAA1L

ADP-Mg²⁺-Water

C

K1912
T2017
W3
T1913
W2
R2091   W1   D1958
D2320   W4
R2358   E1959

D

State-1 to 4	State-5	State-6	State-7	State-8
Closed ADP	**Intermediate Open1** ADP	**Intermediate Open2** ADP	**Open** ADP	**Wide Open** Apo

AAA1 pocket

AAA1-Large   AAA1-Small
AAA2-Large

Buttress-stalk

bulge   ß   →   semi-α   →   semi-α   →   kink   α   →   α

E

Conformational distribution of full-length dynein-1

Fraction

State-8 open AAA1
State-7
State-6
State-5
State-4 closed AAA1
State-3
State-2
State-1
State-0

ATP ATP-Vi AMPPNP ADP Apo

F

Buffer condition	ATP	ATP-Vi	AMPPNP	ADP	Apo
Closed AAA1's nucleotide density (state-1)					
Resolution	2.2Å	2.8Å	2.8Å	2.8Å	2.6Å
Assigned nucleotide	ADP-Mg²⁺	ADP-Vi-Mg²⁺	ADP	ADP-Mg²⁺	ADP-Mg²⁺

Fig. 2. The conformational landscape of MT-unbound dynein motor domains in reaction. (A) Nine major states of the dynein motor domain during the catalytic cycle are divided into auto-inhibited states (left) and states in the active cycle (right). Arrows indicate the proposed catalytic pathway supported by the literature as well as apparent AAA+ ring conformations. Percentages indicate the proportion of particles in each state. The apo AAA1 pocket state (state-8) is present at a very low proportion in the presence of ATP but can be obtained in apo conditions (shown is the structure obtained in apo condition). The linkers from states-5 and -6 are low passed filtered to 12Å and shown at low contours. Only one representative sub-class in each state is shown. The color scheme is consistent with Figure 1D. **(B)** Cryo-EM map of AAA1 pocket from 2.2 Å phi motors. AAA1L, AAA1S, and AAA2L are colored blue, light blue, and sky blue, respectively. On the right, carved transparent grey density is displayed for ADP (green), Mg^{2+} (light green), and water molecules (violet). **(C)** Side chain residues involved in hydrogen-bonding networks (dashed lines) at the AAA1 active site are displayed and colored by heteroatom. **(D)** Cartoon models showing the AAA1 dynamics and the coupled buttress-stalk conformational states. The isolated AAA1 pockets from different states are aligned on AAA1L. Vector maps depicting pairwise alpha carbon interatomic distances between the AAA1 domains of adjacent states depict the three-dimensional trajectories during the pocket opening process. The buttress-stalk regions are highlighted to show how the stalk-CC registries are coupled to motor ring states. **(E)** Plot depicting proportions of the various states of motor domain conformations from full-length dynein-1 in different nucleotide conditions.**(F)** EM densities for nucleotides bound to the closed AAA1 pockets under indicated nucleotide conditions.

Description	Buffer	State	AAA1 & motor ring	Linker	Buttress	Stalk-CC	MTBD affinity with MT	Analogous structures from published PDBs
Full length human dynein-1 unbound from MTs	5mM ATP	State-0	open	straight, docked	straight	bulging	low	N/A
	5mM ATP	State-1 Phi-motor	closed	bent, docked	straight	bulging	low	5nug, 6rla
	5mM ATP	State-2	closed	bent, docked	straight	bulging	low	4rh7
	5mM ATP	State-3	closed	bent, undocked	straight	bulging	low	N/A
	5mM ATP	State-4	closed	straight, undocked	straight	bulging	low	N/A
	5mM ATP	State-5	intermediate open	straight, undocked	intermediate	intermediate	intermediate	N/A
	5mM ATP	State-6	intermediate open	straight, undocked	intermediate	intermediate	intermediate	N/A
	5mM ATP	State-7	open	straight, docked at post-2	kink	canonical	high	3vkg*, 4w8f, 8fcy
	apo	State-8	wide open	straight, docked at post-1	kink	canonical	high	4aki
Full length human dynein-1 bound to MT	5mM ADP	State-7	open	straight, docked at post-1	kink	canonical	high	N/A
	5mM ADP	State-7	open	straight, docked at post-2	kink	canonical	high	N/A
	5mM AMPPNP	State-7	open	straight, docked at post-1	kink	canonical	high	7z8g
	apo	State-8	wide open	straight, docked at post-1	kink	canonical	high	7k58, 7k5b

Table1. Features of the motor domains in different states.

together, our findings demonstrate that incubation with a certain nucleotide analog does not necessarily lock full-length dynein in one conformation, but rather changes its conformational distribution. The structures found in these nucleotide conditions, combined with those in the ATP condition, provide a more comprehensive understanding of dynamic conformational changes during the dynein mechanochemical cycle.

Identification of a "molecular backdoor" for phosphate release from AAA1

States-1, -2, -3, and -4 in the presence of ATP all have closed AAA1 pockets and AAA+ ring conformations. However, the linker in each of these states ranges from pre-powerstroke to a variety of intermediate states during the powerstroke **(Figure 2A)**. We observed a similar range of linker states for those dyneins with a closed AAA1 pocket in the ATP-Vi and AMPPNP conditions **(Figure 2E, S4)**. Given the proposed role of AAA1 in governing linker position *(36, 40, 42, 49, 77)*, these observations suggest there may be structural differences in AAA1 that are not apparent from a simple large-scale assessment of the open or closed state of this pocket. We thus more closely investigated whether there exist any potential small-scale changes in AAA1 among the different states. This revealed that the sensor-I loop adopts multiple distinct conformations in the presence of ATP **(Figure 3A)**. This loop is essential for the ATPase activity of AAA+ proteins *(80)* and has been proposed to facilitate dynein Pi release through a backdoor mechanism *(31)*. Notably, this loop adopts the same conformation in dyneins with closed AAA1 pockets (states-1, -2, -4) incubated in ATP-Vi (or AMPPNP) **(Figure 3A)**. Interestingly, the sensor-I loop becomes completely disordered when the AAA1 pocket is open **(Figure 3B)**, such as in states-7 and -8.

Close inspection of the cryo-EM densities of the sensor-I loop reveals two residues within this loop, N2019 and Y2022, that exhibit significant conformational changes **(Figure 3C)**. In both ATP-Vi and AMPPNP conditions, these two residues consistently orient towards the AAA1 active site and make direct contact with the AAA1 walker-A motif and AAA2L. In the presence of ATP, these residues exhibit distinct conformations: in both states-1 and -2, N2019 points away from the active site ("outward"), while Y2022 adopts multiple conformations ("flexible density"). In state-3, which is absent in ATP-Vi and AMPPNP conditions, N2019 also points away from the pocket, and Y2022 has two alternative conformations. In state-4, the entire sensor-I loop exhibits stable density, resembling that observed in ATP-Vi and AMPPNP states. Thus, despite the closed AAA1 pocket for all these states, there indeed exist small-scale, residue-level changes that very likely coordinate dynein mechanochemistry.

To investigate the role of the dynamics of these sensor-I loop residues during the ATPase cycle, we compared the local environment among the different states **(Figure 3D, S5)**. This revealed that when both N2019 and Y2022 point outward, a

channel appears to form, creating a possible escape route for Pi release. This observation supports the notion of a "molecular backdoor"*(31)* in which the coordinated rearrangement of residues in the sensor-I loop can establish a closed or open backdoor during the Pi release process **(Figure 3E, Movies S3)**. To further test this model, we performed focused classification of the AAA1 active site for the phi-motor (state-1) in ATP and captured what may be Pi at two different points during its release **(Figure 3F**; see arrows**)**. Importantly, the residues at the sensor-I loop, including N2019 and Y2022, are highly conserved between dynein-1 in different species and among dyneins (e.g., axonemal dynein and dynein-2; **Figure 3G)**, suggesting they are likely important for some aspect of dynein mechanochemistry.

The dynamic nature of Y2022 suggests it plays a crucial role in mediating the release of Pi. To test the importance of this residue in dynein function, we utilized a well-characterized GST-dimerized yeast dynein motor construct *(41)* and mutated the corresponding residue (Y1902) to either phenylalanine or alanine. Compared to the wild-type, Y1902F exhibits similar basal and MT-stimulated ATPase levels while Y1902A was defective in both **(Figure 3H)**. The addition of MTs only marginally stimulated the ATPase activity of the Y1902A mutant (by 20%, compared to 3.4- and 3.1-fold for wild-type and Y1902F, respectively), suggesting MT-binding is uncoupled from ATPase activity. Unlike human dynein, this minimal GST-dynein fragment is processive without the addition of dynactin or adaptor proteins, permitting assessment of motility without complicating factors. Single-molecule assays revealed that, unlike Y1902F, the Y1902A mutant exhibits significant motility defects, as apparent from the reduced fraction of active motors and the much lower velocity for those motors that exhibit processive runs **(Figure 3I and J)**. Intriguingly, a large proportion of the Y1902A motors (84.4%) appear to exhibit bidirectional diffusive behavior along MTs (see red arrows in kymographs), which is further suggestive of uncoupled MT-binding and ATPase activity. These functional data demonstrate the importance of this tyrosine in the catalytic and motility behaviors of dynein.

Pi release triggers linker straightening during the active cycle

The backdoor Pi release mechanism prompted us to revisit the ATP turnover of the autoinhibited phi dynein, as well as open dynein. Our data indicate that phi dynein is indeed capable of hydrolyzing ATP, as apparent from the clear density of ADP-Mg^{2+}, as well as by the presence of Vi in the ATP-Vi conditions **(Figure 2B)**. Furthermore, the dynamic sensor-I loop conformation suggests that a Pi is in the process of escaping the pocket **(Figure 3D, 4A)**. Due to the autoinhibited state of the motor domains, ATP hydrolysis and subsequent Pi release do not lead to a change in linker conformations **(Figure 4A)**. Similar to phi dynein, states-2 and -3 of open, uninhibited dynein also have dynamic sensor-I loop conformations that are indicative of Pi release **(Figure 3C)**.

Fig. 3. The release of Pi from AAA1 through a molecular backdoor mechanism. (A) Comparison of molecular models of closed AAA1 pockets in different states and nucleotide conditions. The sensor-I loop displays conformational heterogeneity when dynein is in ATP condition but not in ATP-Vi and AMPPNP conditions. **(B)** Cryo-EM densities of sensor-I loop in closed and open AAA1 pockets, showing that this loop becomes disordered when the AAA1 pocket is in open states. **(C)** Structural comparison of the bound nucleotide and the sensor-I loop in closed AAA1 states. Transparent cryo-EM densities of the sensor-I loop are displayed in isosurface mode. The conformations of two residues, N2019 and Y2022, are highlighted. **(D)** A slice through the structures of the AAA1 pocket near the nucleotide-binding site, shown in surface representation. **(E)** Backdoor model for Pi release from dynein AAA1 controlled by N2019 and Y2022. **(F)** High-resolution 3D classification of the active site in the phi motor, showing the potential Pi group in various states of escape. **(G)** Conservation analysis of the sensor-I loop for dynein-1 in different species and dynein in different subfamilies (DNAH, dynein axonemal heavy chain; dynein-2, intraflagellar transport dynein). **(H)** Plot (mean ± SD, along with individual data points) depicting relative ATPase rates (in the absence or presence of 2 μM microtubules) for artificially dimerized yeast dynein motor domain fragments (from left to right, n = 6/5, 6/4, and 6/6 independent replicates from assays minus/plus MTs). Y1902 in yeast dynein corresponds to Y2022 in human dynein. **(I and J)** Representative kymographs (I) and plots (J; mean ± SD, along with individual data points) depicting indicated motility parameters determined from single-molecule motility assay (red arrows, diffusive motors; green arrows, processive motors; from left to right, n = 508/597, n = 148/880, and n = 679/837 processive/non-processive dynein molecules from three independent experiments). Microtubule polarity was determined with a wild-type dynein fragment (see Methods).

However, unlike phi dynein, the linker in state-3 shows a transitional conformation between bent and straight, suggesting a correlation between the sensor-I loop and linker position during Pi release. Moreover, after the Pi is released in state-4 via the backdoor mechanism, as indicated by the stable densities for N2019 and Y2022, the linker adopts a straight conformation. This is consistent with previous kinetic measurements that the linker adopts a straight conformation after the release of Pi *(40)*. Based on the Pi-free but closed state of the AAA1 pocket in state-4, we posit that subsequent to Pi release, and not prior to, the AAA1 pocket can adopt an open state.

Whether the linker is in a bent or straight state is determined by the hinge helix **(Figure 4B, C)** *(42)*. However, the precise mechanism that triggers the straightening of the hinge helix is unclear. Our structures reveal that the release of Pi from AAA1 is coordinated with angstrom-scale, but significant conformational changes of the sensor-I loop **(Figure 3)**. These changes could theoretically be propagated to the linker-AAA+ ring interface via AAA1L's central β-sheets, leading to large nanometer-scale conformational changes of the hinge helix, and subsequent linker straightening at an even larger scale **(Figure 4D)**. Close inspection of the linker-ring interfaces in states-3 and -4 reveal that the short linker helix 11 (H11) and the H11-H12 loop exhibit 1.7-5.1 Å displacements from state-3 to -4 **(Figure 4E)**. These movements remodel the interactions between the linker and AAA1L H2, AAA2L PS-I insert, and AAA2L H2 insert. One notable change is the apparent disruption of a salt bridge (R1943-E1620) between states-3 and -4 **(Figure 4F)**. This salt-bridge pair is conserved across dynein species and was also reported in the dynein-2 structure*(42)* **(Figure 4G)**, suggesting it likely plays a crucial role in maintaining a bent linker. In state-3, this salt bridge is further stabilized by a hydrogen bond between R1943 and Y1630 in linker H11. In state-4, as linker H11 shifts away from the AAA1L H2, the hydrogen bond and salt bridge are both broken, causing the R1943 side chain to flip. Additionally, the AAA2L PS-I insert and H2-insert no longer contact the linker in state-4. The loss of all these interactions relieves the bent hinge helix from its distorted and restrained state *(42, 49)*, thereby triggering the linker straightening **(Figure 4F, H)**.

The communication between AAA1 pocket and the MTBD

It has been shown that AAA1 pocket dynamics regulate

the affinity of the MTBD for MTs via registry changes in the stalk-CC *(31)*. However, the precise mechanism for this long-range (24 nm) communication is unclear. Our extensive collection of dynein's structural states **(Figure 2A)** reveals detailed pathways for the signal propagation from AAA1 to the MTBD **(Movies S4)**. From state-4 to -5, the AAA1 pocket opens slightly, resulting in a movement of AAA2L towards AAA2S, and a small movement of AAA1L, which triggers the outward rotational movement of AAA6 **(Figure 5A and B)**. This further induces the CTD to swing from AAA3 to AAA5, and a consequent rotation of AAA5 causes the buttress to twist and the stalk-CC to adopt a "semi-α" registry, which is indicative of the MTBD adopting a higher MT-binding affinity. When the AAA1 pocket further opens from state-5 to -6 **(Figure 2D)**, AAA2L completely loses its interactions with AAA1L and moves towards AAA3 **(Figure 5B)**. However, this propagated signal appears to stop at AAA3L **(Figure 5B**, note lack of movement beyond AAA3L in vector diagram), thus explaining the lack of change in the buttress-stalk conformation between state-5 and -6.

Upon transition to state-7, the AAA2L domain continues to move away from AAA1L, but in a different direction: toward AAA3L **(Figure 5B)**. This signal is propagated to the buttress-stalk via AAA4L and AAA5L, leading to the adoption of the α-registry by the stalk-CC. From state-7 to -8, the AAA1 pocket opens completely, with a sufficient separation between AAA1L and AAA1S that permits ADP release. This pocket widening slightly affects the conformations of AAA2-5L but does not affect the buttress-stalk conformation **(Figure 5B)**. Our results strongly support a model in which AAA1 conformational signals are propagated in two phases **(Figure 5C and D)**: first through AAA6, CTD, AAA5, and the buttress to stalk-CC (to adopt the semi-α registry), and then subsequently via AAA2, AAA3, and AAA4 to complete the adoption of the α registry by the stalk-CC.

Our results show how AAA1 transmits its conformational signals to the buttress-stalk region to change the stalk-CC registry to affect MTBD-MT affinity. However, previous studies have shown that locking the stalk-CC in different registries affects the ATPase activity, motility, and tension-sensing of dynein motors *(38, 75)*, supporting a model in which the MTBD can also transmit signals back to the AAA+ ring through the stalk-CC **(Figure 5E)**. To investigate the structural basis of the reverse communication, we performed

Fig. 4. A mechanism for linker straightening triggered by the Pi release during the active cycle. (A) Cartoon models showing the process for Pi release for phi and open dynein. Unlike autoinhibited phi dynein, the release of Pi in open dynein results in the linker straightening. **(B,C)** Cryo-EM maps of states-4 and -3 with accompanying molecular models of the linker. The hinge helix in the linker is indicated. **(D)** Structural comparison of the state-3 (grey) and state-4 (color by domains). The AAA+ ring between the two states shares the same conformation while the sensor-I loop and linker adopt different conformations.**(E)** Linker-ring interface remodeling from state-3 (grey) to state-4 (colors). From state-3 to -4, H11, and the H11-H12 loop shift away from the AAA1L-H2, AAA2L PS-I insert and AAA2L H2 insert. Distance change values for the indicated residues are included. **(F-H)** Mechanism of the hinge helix straightening from state-3 to -4. In state-3, the bent hinge helix is stabilized by the conserved salt bridge and hydrogen bond interactions between R1943 in AAA2L-H2 and the hinge helix (F, G). During the linker-ring remodeling, R1943 flips away from the hinge helix in state-4, breaking these interactions which leads to the straightening of the hinge helix (F). The cartoon models depicting this process are shown on the right (H).

cryo-EM analysis of a human dynein motor domain mutant in which the stalk-CC is locked in the α registry *(21)* (to mimic the MT-bound state; **Figure 5F, Table S7**). This revealed that in the presence of ATP, this mutant consistently has an open ADP-bound AAA1 pocket, and a kinked buttress conformation *(21)***(Figure 5G)**, resembling that of WT full-length dynein in state-7. Even in the presence of ATP-Vi, the AAA1 pocket is bound to ADP and remains in the same open conformation, consistent with this mutant not undergoing Vi-dependent photocleavage (a readout for Vi binding) *(21)*. These results suggest that locking the stalk-CC in the α-registry forces the AAA1 pocket to adopt an open conformation. Taken together, we provide a structural basis for a comprehensive understanding of the two-way communication between AAA1 and the MTBD.

Structures of MT-bound dynein motors reveal that MTs accelerate ADP release

To gain insight into the role of MTs during dynein's mechanochemical cycle, we obtained four distinct structures for MT-bound dynein motor domains **(Figure 6A)**, with AAA1 in either apo or ADP-bound states, all of which possess stably docked linkers and kinked buttress. Exclusion of nucleotide from the buffer resulted in AAA1 being in an apo state, both AAA3 and AAA4 being bound to ADP, and the linker being in the canonical "post-1" post-powerstroke state. In ADP-containing buffer conditions, we identified ADP in AAA1, AAA3, and AAA4, with the linker docking at one of two different positions (post-1 and post-2). In the presence of AMPPNP, both AAA3 and AAA4 are bound to AMPPNP, while AAA1 remains ADP-bound, and the linker is docked

Pengxin Chai, Jun Yang *et al.* |

Fig. 5. The mechanism for two-way communication between AAA1 and the MTBD. (A) Cartoon models illustrate the overall motor ring conformations from different states, each with varying degrees of the AAA1 pocket opening. The linker domain is hidden for better visibility of the motor ring states. **(B)** Vector maps depicting pairwise alpha carbon interatomic distances (for every 3rd residue) between the AAA domains of adjacent states. The length of the lines is proportional to the calculated interatomic distances. The major movements and directions are highlighted by dashed areas and arrows. **(C)** The cartoon model illustrates the signal propagation directions from AAA1 to the stalk-CC in two distinct phases. **(D)** Changes in dynein-MT binding affinity (Y-axis), as assessed by buttress-stalk conformations, as a consequence of AAA1 pocket conformations in the different states (X-axis). **(E)** Cartoon model illustrating the communication from the MTBD to AAA1 via the stalk-CC. **(F)** Cartoon model depicting the design of the α registry-locked dynein motor domain mutant. **(G)** Cryo-EM reconstructions of the α registry mutant in ATP or ATP-Vi conditions. In both conditions, the motor domains exhibit only open AAA1 pockets. As expected, the buttress-stalk conformation is consistent with a high MT-binding affinity.

at only post-1.

To assess the impact of MT binding on dynein motors, we compared these structures with the MT-unbound structures described above **(Figure S6)**. Strikingly, three of the MT-bound motor domain structures (i.e., those from apo and ADP conditions) matched very well with structures from the conformational landscape of MT-unbound dynein (Cα RMSD < 0.3 Å with state-7 or -8). In contrast, MT-bound dynein incubated in AMMPNP-containing buffer exhibits a 2.3 Å-5.1 Å Cα RMSD in comparison to the MT-unbound motors, with the closest match being to state-7. These findings indicate that the dynein motor domain can naturally sample the MT-bound state even in the absence of MTs. However, it is noteworthy that the MT-bound motors exhibit more stable cryo-EM densities in the buttress-stalk and linker regions compared to their MT-unbound counterparts. In addition, the state-5 and -6 structures with intermediate open pockets are no longer found in MT-bound motors, providing additional evidence that MT binding indeed stabilizes the stalk-CC registry, the AAA+ ring conformation, and the linker-AAA docking, all of which is likely triggered via the MTBD-MT contact.

How might MTs accelerate the ATPase activity based on these results? It has been proposed that ADP release is the rate-limiting step for dynein *(81)* and that MTs accelerate this step *(66)*. In agreement with this model, we find that approximately 45% of MT-unbound dynein motors exhibit ADP-bound intermediate AAA1 open pockets with their stalk-CC in the semi-α registry (state-5 and -6; **Figure 6B**). MT-binding shifts these two intermediates to state-7 by inducing the stalk-CC to adopt the α registry*(57)* **(Figure 6C, left)**, which leads to further opening of the motor ring and the AAA1 pocket for subsequent ADP release **(Figure 5)**. MT-binding can also induce the adoption of the "wide open" state of the AAA1 pocket, as observed in state-8, which allows ADP release. In support of this, nucleotide-free buffer conditions lead to the identification of both states-7 (i.e., ADP-bound AAA1) and -8 (i.e., apo AAA1) for MT-unbound dynein, but only lead to the presence of state-8 for MT-bound dynein **(Figure 6C, right)**. These results explain that two limiting steps, stalk-CC registry change *(64)*, and ADP release *(81)*, are strictly coupled and concurrently stimulated by MT-binding. In addition, our evidence for a Pi backdoor release mechanism in low-MT binding affinity states supports a previous model that MT-binding does not affect the Pi release step *(66)*.

Post-2 linker-ring docking mode provides a net bias for dynein minus-end stepping

The linker swing vector model has been proposed to explain the mechanism of dynein unidirectional motility *(50, 82, 83)*. However, which step during the mechanochemical cycle provides a net minus end-directed bias remains unknown. We analyzed the different linker-ring interaction interfaces **(Figure S7A)** and found that the two different

linker docking modes (post-1 and post-2) may directly impact dynein stepping behavior, similar to previous reports on OAD *(69)*. The observation of post-2 in dynein-1 suggests that this docking mode is conserved **(Figure S7 B-E)**.

Based on previous OAD-MT structures, it was proposed that dynein in post-2 mode would position its MTBD 8 nm forward of dynein in post-1 mode **(Figure 6D)**. To investigate the role of the two different docking modes in dynein stepping, we assessed how the linker position would correlate with dynein stepping behavior **(Figure 6E, Movie S5)**. When AAA1 is in the apo state, the dynein motor binds to MTs in post-1, which is the starting position (0 nm; **Figure 6Ei**). ATP binding and/or hydrolysis at AAA1 leads to the adoption of the pre-powerstroke state, and the detachment of dynein from MTs (due to the low affinity for MTs in this state; **Figure 6Eii**). The Pi backdoor release and linker straightening process likely induce the AAA+ ring to rotate relative to the tail rather than pull the tail region forward because the dynein is in the low MT-binding affinity state *(83)* **(Figure 6Eiii)**. After subsequent AAA1 pocket-opening, the motor can rebind to MTs with an ADP-bound open pocket. If the motor domain adopts the post-1 mode, the MTBD returns to its original position without any forward stepping **(Figure 6Eiv-a)**. By contrast, stepping with the motor domain in post-2 mode generates a net bias towards the minus-end of MTs, allowing dynein MTBD to bind 8 nm forward along the MT **(Figure 6Eiv-b)**. In support of this idea, the MT-unbound motors in state-7 (ADP-bound open AAA1) primarily adopt the post-2 mode in the presence of ATP **(Figure 2A)**. Finally, the transition from ADP-post-2 to ADP-post-1 – or to apo-post-1 – results in further translocation (total 8 nm) of the cargo-binding tail domain **(Figure 6Ev)**.

Discussion

Technical advances to explore the dynein mechanochemical cycle

In this work, we demonstrate that our improved cryo-EM pipeline **(Figure 1)** can identify the conformational landscape of full-length human dynein-1 in reaction with ATP. This approach contrasts with previous studies that employed truncations, mutations, and specific nucleotide conditions to lock dynein in one state *(42, 44, 45, 47)*. Our pipeline not only captures nine distinct motor states but also resolves them at high resolutions **(Figure 2)**. The 2.2 Å structure of the phi motor domain provides a solid structural basis to analyze the molecular interactions at an atomic level, particularly the residues involved at the reactive center **(Figure 2)**. Systematic analysis of the dynamic range of motor domain structures from full-length open dynein, both on and off MTs, permits an extremely comprehensive understanding of the mechanochemical cycle of dynein **(Figure 2-6)**. Our approach also resolves the MT-bound dynein motors at high resolutions, shedding light on the impacts of MT-binding on

Fig. 6. The mechanism for ATPase stimulation by MTs and dynein's unidirectional stepping along MTs. **(A)** High-resolution cryo-EM maps of MT-bound dynein motor domains in different nucleotide conditions. **(B)** The pie chart of the distribution of various states of dynein in active cycle with ATP. The Pi release and ADP release steps correlated with different states are shown. **(C)** The cartoon model depicts the mechanism of dynein mechanochemical cycle acceleration by MTs. MT-binding rapidly induces the stalk-CC registry from semi-α to α registry, therefore forcing the AAA1 pocket to adopt an open state (left). MT-binding, by stabilizing the stalk-CC registry, allows the further wide opening of the AAA1 pocket and leads to the final ADP release (right). **(D)** Dynein adopting post-2 linker docking mode intrinsically favors a forward step, based on the previous OAD-MT structures (PDB: 7k58, 7k5b *(69))*. **(E)** Proposed mechanism for dynein unidirectional stepping along MTs. **(i)** Dynein is bound to MTs in the apo state in post-1 mode (after completing a step). **(ii)** Upon ATP binding and/or hydrolysis, dynein unbinds MTs with a bent linker. **(iii)** Upon Pi release from AAA1, dynein with a straight but undocked linker (states-4 to 6) with low and intermediate MT-binding affinity. **(iv-b)** The preferred post-2 mode in ATP conditions (denoted by a star) creates a net bias towards the minus end of MTs and allows MTBD to take an 8 nm forward step. Note that adopting a post-1 mode upon MT-rebinding **(iv-a)** leads to a 0 nm step. The cartoon model from state-4 is shown in grey. **(v)** Switching from a post-2 to a post-1 mode and/or subsequent ADP release leads to an 8 nm forward movement of the entire dynein complex.

dynein (**Figure 6**). The numerous new intermediate states reported in this work, which are technically challenging to acquire using conventional approaches, substantially expand our knowledge of the dynein conformational landscape.

The release of Pi through a molecular backdoor mechanism

We provide structural and functional evidence that supports the existence of a "backdoor" for Pi release that involves small but significant rearrangements of the AAA1 sensor-I loop (**Figure 3**). In the presence of ATP, two residues in the sensor-I loop – N2019 and Y2022 – are found in various conformations in different motor states. We speculate that N2019 senses the Pi cleavage while Y2022 gates the release of Pi. In support of this idea, N2019 closely interacts with the vanadate group when AAA1 is bound to ADP-Vi (2.6 Å). The cleavage of the ATP forces N2019 to adopt outward conformation to accommodate the Pi. Meanwhile, Y2022, which exhibits multiple conformations, creates a gated escape route for Pi release. It likely performs this function via its large aromatic side chain, since the Y1902F was functional in ATPase and motility assays. When the Pi is fully released, both residues stably contact the walker-A motif, completing the Pi release step and allowing the subsequent pocket opening processes.

Furthermore, our data strongly support a mechanism whereby the conformational changes in the sensor-I loop transmit Pi-release signals to the process of linker straightening (**Figure 4**). However, this process is sterically restrained in the autoinhibited phi dynein in which Pi release does not lead to linker straightening. These findings also elucidate why the bent or straight linker are both found in the presence of ATP-Vi for human dynein-1 (this study), yeast dynein-1 (47), dynein-2 (42), and axonemal dynein-c (49, 77). As Pi is in the process of releasing through the backdoor, vanadate can attack the active site while the linker is in multiple intermediate states, therefore stabilizing dynein in a closed motor ring state with either a bent or straight linker.

A new perspective of dynein conformation distribution under various conditions

Our findings also show that a specific nucleotide environment influences the conformational distribution of the motor domain for full-length dynein. A recent structure of the DDA-LIS1 complex bound to MTs in the presence of AMPPNP revealed a pre-powerstroke state motor domain bound to LIS1 (70). In this motor-LIS1 complex with a bent linker, the AAA1 pocket binds ADP with a closed conformation, which is speculated to be induced by LIS1 (70). By contrast, our data demonstrates that this motor state (state-2), without LIS1 binding, can be directly observed under ATP, ADP, AMPPNP, and even apo conditions in the absence of LIS1 (**Figure S4**), indicating that this is an intrinsic property of the dynein motor domain. In the future, by comparing the conformational landscape of dynein with LIS1 or other regulators to the many structures reported in this study, the

impacts of regulators on dynein can be better characterized.

The powerstroke of dynein consists of multiple steps

The powerstroke of dynein involves the transition of the AAA+ ring from a closed (MT-unbound) to an open (MT-bound) conformation, coupled with the straightening of the linker from a bent to a straight conformation (31, 32, 84). This transition has been hypothesized to generate the force necessary for dynein to pull its cargo forward. The existence of state-4 in our dataset indicates that linker straightening occurs before pocket opening and even MT-binding (**Figure 3**). We also identify two additional states (state-5, and -6), with partially open ADP-bound AAA1 pockets that appear to represent intermediates before the MT-bound state (**Figure 4**). Note that all three states (4 through 6) feature straight (as apparent at low contour levels), but undocked linkers. Upon dynein binding to MTs, the straight linker is stably docked on the motor ring in one of the two modes (post-1 and post-2).

Based on the existence of these newly identified linker intermediates, we propose a new model for the force-generating powerstroke (**Movie S5**). We designate state-2, characterized by a closed AAA+ ring with an ADP-Pi bound AAA1 pocket (i.e., post-ATP hydrolysis and before Pi release), and a bent, stably docked linker, as the pre-powerstroke state (**Figure 6Eii**), consistent with a previous structure (42). We propose that the powerstroke proceeds through the following steps: 1) Pi is released from AAA1 via the backdoor (while the AAA1 pocket and AAA+ ring remain closed), and the linker transitions from a bent, stably docked state to a straight, unstable conformation. Note the MT-binding affinity remains low, and dynein is consequently unbound from MTs (**Figure 6Eiii**). 2) The AAA+ ring transitions from closed to intermediate open states while the linker remains straight and undocked. 3) Upon MT-binding (and adoption of a high MT-binding affinity state), the linker transitions to a conformation in which it is docked in a post-2 state (the preferred state for dynein in the moments before MT-binding), coincident with the AAA+ ring transitioning to an open state (**Figure 6Eiv-b**). 4) The linker transitions to a post-1 state in the ADP-bound or apo AAA1 state, resulting in the translocation of cargo by 8 nm (**Figure 6Ev**). In this model, the force necessary for dynein to pull its cargo is primarily generated during steps 3 and 4 as dynein is anchored on MTs.

Model for full-length human dynein-1's mechanochemical cycle

Our work significantly improves our understanding of dynein's mechanochemical cycle (**Figure 7**). We find that the motor domain from the autoinhibited phi dynein is competent for ATP hydrolysis and Pi release but is sterically hindered for ADP release. We have identified a potential additional level of autoinhibition for open dynein, characterized by an incomplete closure of the motor domain (state-0) that is likely catalytically inactive. The motors engaged

Fig. 7. Model for the mechanochemical cycle of full-length human dynein-1. The full-length dynein adopts two autoinhibited states: state-0 in the open conformation, and state-1 in phi conformation. Motor domains in autoinhibited states and the active states exist in an equilibrium. The mechanochemical cycle of the active motor states starts with the apo AAA1 state. ATP binding and/or hydrolysis induces pocket closure and triggers the MT release. The linker changes from straight, stably docked to bent, stably docked conformation. The backdoor release of Pi triggers the straightening of the linker. The presence of MTs actively engages in the pocket-opening process by inducing the stalk-CC registry to change from semi-α and α states. The linker conformation changes from the straight, unstably docked to straight, stably docked conformation upon MT-binding. The linker-docking mode transitions and ADP release completes the cycle and dynein takes a forward step.

in the mechanochemical cycle are predominantly in ADP-bound states, consistent with ADP-release being the limiting step, and with the fact that dynein only possesses basal ATPase levels off MTs *(66, 81)*. By comparing the conformational landscapes for dynein on and off MTs, we propose the following model to explain its mechanochemical cycle with MTs. Note that dynein can still progress through this cycle when unbound from MTs (**Figure 2**).

1) ATP binding to the apo AAA1 state (state-8) and/or subsequent hydrolysis induce pocket closure, triggering a registry change in the stalk-CC from α to β and a straight-to-bent shift in the linker *(31)*. 2) Following ATP hydrolysis, the sensor-I loop facilitates Pi release from AAA1 via a molecular backdoor mechanism. After the release of Pi, the linker is straightened but not stably docked on the closed mo-

tor ring. 3) The ADP-bound AAA1 pocket bound gradually transitions from closed to intermediate-open and open states. Throughout this process, the buttress and stalk respond to the pocket-opening signal across the motor ring, resulting in semi-α and α registry stalk-CC configurations. MT-binding significantly accelerates this step by actively triggering the adoption of the α registry of the stalk-CC that eventually facilitates ADP release. 4) Further widening of the AAA1 pocket ensures ADP release and dynein thus pulls its cargo one step forward (see above), completing the current cycle.

Our model, supported by previous structural studies and over 30 high-resolution structures in this work, clarifies many long-standing questions in the field *(31)*. Our approach to determining the conformational landscape of dynein "in reaction" – on and off MTs – has revealed valuable insight that

was impossible with prior approaches that relied on strategies that aimed for conformational homogeneity. Furthermore, in addition to providing numerous testable hypotheses (via mutagenesis combined with functional assays), our high-resolution intermediate states of the dynein motor provide a strong structural basis for computational simulation *(85)* and protein engineering *(86)*. Lastly, our work featuring high-resolution dynamic cryo-EM analysis provides a new paradigm that can be expanded to provide deeper mechanistic insight into the entire dynein transport machinery on MTs *(2)* (in complex with dynactin and a cargo adaptor), or regulation by its cofactors (e.g., LIS1 *(20)*, NDEL1 *(87)*).

Acknowledgements

We are very grateful to members of the Zhang and Markus laboratories for their valuable discussions. We thank R. Yang for the initial dynein-1 expression and preparations in the lab. This work was funded by the NIH/NIGMS (R35GM139483 to S.M.M., and R35GM142959 to K.Z.) and in part by a Collaboration Development Award Program (to K.Z.) from the Pittsburgh Center for HIV Protein Interactions (U54AI170791). We would like to thank K. Zhou, J. Lin, M, Llaguno, and S. Wu, for their help with cryo-EM data collection at Yale Cryo-EM facility. The Yale Cryo-EM Resource is funded in part by the NIH grant S10OD023603 awarded to F. Sigworth. We thank L. Wang, J. Kaminsky, and G. Hu at the Laboratory for BioMolecular Structure (LBMS) for help with cryo-EM data collection. The LBMS is supported by the DOE Office of Biological and Environmental Research (KP1607011).

Author contributions

K.Z. designed the study. J.Y. and Y.W. purified full-length human proteins. J.Y., Y.W. and P.C. prepared the cryo-EM samples and collected the data. P.C., J.Y., Y.W. and K.Z. processed the images, built PDB models and analyzed the structures. S.M. purified the human MT-B dynein motor domain. I.G. generated yeast strains, and purified proteins from yeast. I.G. and S.M. performed and analyzed ATPase and single molecule assays. P.C., J.Y. Y.W., K.Z., and S.M. generated figures and movies. P.C., J.Y. and K.Z. wrote the manuscript. P.C., J.Y., Y.W., K.Z. and S.M. edited and revised the manuscript.

Competing interests statement

The authors declare no competing interests.

Data availability

Cryo-EM maps and atomic coordinates have been deposited in the Electron Microscopy Data Bank under accession codes EMD-XXXX and in the Protein Data Bank under accession XXXX.

Methods

Full-length human dynein-1 purification

The plasmid encoding full-length human dynein-1 was graciously provided by the Andrew Carter Lab (Addgene plasmid 11903). This plasmid featured the dynein-1 heavy chain fused to an N-terminus for the His-ZZ-SNAPf tag, followed by the Tobacco Etch Virus (TEV) protease cleavage site. Recombinant baculoviruses were produced in sf9 insect cells via transfection of isolated Bacmid obtained from DH10MultiBac competent cells (Geneva Biotech) using Cellfectin® II (Gibco). Expression of dynein-1 in sf9 cells was accomplished by infecting them with P2 virus at a cell density of 2.5 million cells/ml. 28 ml of P2 virus was added into a 1.4 L culture of sf9 cells. Following 75 hours of incubation, the infected cells were harvested by centrifugation at 1000 rcf for 15 minutes at 4°C. The resulting cell pellet was flash-frozen in liquid nitrogen and stored at -80°C until further use. For dynein purification, frozen cell pellets from 1.4 L were thawed and resuspended in 100 ml of lysis buffer (50 mM HEPES pH 7.2, 100 mM NaCl, 1 mM DTT, 0.1 mM ATP, 10% glycerol) supplemented with 2 protease inhibitor tablets (Roche) and 2 mM PMSF. The cell suspension was lysed using a 40 mL Dounce tissue grinder (Wheaton) with 15-25 strokes on ice, followed by clarification via ultracentrifugation at 65k rpm with a Ti70 rotor for 1 hour at 4°C. The resulting supernatant was incubated with 3 mL of pre-equilibrated IgG Sepharose affinity beads (Cytiva) on a roller at 4°C for 3 hours. Subsequently, the beads were packed into a gravity column, washed sequentially with 200 mL of lysis buffer, followed by 200 mL of TEV buffer (50 mM Tris–HCl pH 7.4, 148 mM K-acetate, 2 mM Mg-acetate, 1 mM EGTA, 10% glycerol, 0.1 mM ATP, 1 mM DTT). The beads were then transferred into a 15 ml tube filled with TEV buffer and supplemented with 400 µg of house-made TEV protease, and incubated overnight on a roller at 4°C. The resulting flow-through was collected and concentrated using a pre-equilibrated 100K Amicon Ultra-15 concentrator (Millipore), then loaded onto a TSKgel G4000SWXL 7.8/300 column equilibrated with GF150 buffer (25 mM HEPES pH 7.2, 150 mM KCl, 1 mM MgCl2, 5 mM DTT, 0.1 mM ATP) at 4°C. Peak fractions were pooled and concentrated to 2.5-3 mg/ml for immediate use in cryo-EM grid preparation. The quality of dynein is monitored using negative staining EM (**Figure S1**).

α-*registry human dynein motor domain purification*

The human dynein motor domain locked in α-registry was expressed and purified from insect cells (ExpiSf9 cells; Life Technologies) as previously described *(21)*. Briefly, 4 ml of ExpiSf9 cells at 2.5 x 106 cells/ml, which were maintained in ExpiSf CD Medium (Life Technologies), were transfected with 5-9 µg of bacmid DNA using ExpiFectamine (Gibco) according to the manufacturer's instructions. 4-8 days following transfection, the cells were pelleted, and 1-2 ml of the resulting supernatant (P1) was used to infect 200 ml of ExpiSf9 cells (5 x 106 cells/ml). Approximately 65 hours later, the cells were harvested (2000 x g, 20 min), washed with lysis buffer (50 mM HEPES, pH 7.4, 100 mM NaCl, 10% glycerol, 1 mM DTT, 0.1 mM Mg-ATP, 1 mM PMSF), pelleted again (1810 x g, 20 min), and resuspended in an equal volume of same. The resulting cell suspension was drop frozen in liquid nitrogen and stored at -80ºC. For protein purification, 30 ml of additional lysis buffer supplemented with cOmplete protease inhibitor cocktail (Roche) was added to the frozen cell pellet, which was then rapidly thawed in a 37ºC water bath prior to incubation on ice. Cells were lysed in a dounce-type tissue grinder (Wheaton) using 50-60 strokes. Subsequent to clarification at 310,000 x g for 1 hour, the supernatant was applied to 2 ml of IgG sepharose fast flow resin (GE) pre-equilibrated in lysis buffer, and incubated at 4ºC for 3-5 hours. Beads were then washed 3 times with with 5-10 ml of lysis buffer, and 2 times with 5-10 ml of TEV buffer (50 mM Tris pH 7.4, 150 mM potassium acetate, 2 mM magnesium acetate, 1 mM EGTA, 10% glycerol, 1 mM DTT, 0.1 mM Mg-ATP). The beads were incubated with TEV protease overnight at 4ºC. The next morning, the recovered supernatant was collected, concentrated and used directly without freezing.

Purification of yeast dynein motors

Purification of yeast dynein (wild-type and mutants; ZZ-2xTEV-6His-GFP-3HA-GST-dyneinMOTOR-HALO, all under the control of the galactose-inducible promoter, GAL1p) was performed as previously described *(17, 88)*. Briefly, yeast strains were generated with plasmids encoding wild-type or mutant dyneins integrated into the yeast genome (after digestion with ApaI, as previously described *(17, 88)*. The strains were grown in YPA supplemented with 2% galactose, harvested, washed with cold water, and then resuspended in a small volume of water. The resuspended cell pellet was drop frozen into liquid nitrogen and then lysed in a coffee grinder (Hamilton Beach). After lysis, 0.25 volume of 4X lysis buffer (1X buffer: 30 mM HEPES, pH 7.2, 50 mM potassium acetate, 2 mM magnesium acetate, 0.2 mM EGTA) supplemented with 1 mM DTT, 0.1 mM Mg-ATP, 0.5 mM Pefabloc SC (concentrations for 1X buffer) was added, and the lysate was clarified at 310,000 x g for 1 hour. The supernatant was then bound to IgG sepharose 6 fast flow resin (Cytinva) for 2-4 hours at 4°C, which was subsequently washed three times in 5 ml lysis buffer, and twice in TEV buffer (50 mM Tris, pH 8.0, 150 mM potassium acetate, 2 mM magnesium acetate, 1 mM

EGTA, 0.005% Triton X-100, 10% glycerol, 1 mM DTT, 0.1 mM Mg-ATP, 0.5 mM Pefabloc SC). To fluorescently label the proteins, the bead-bound protein was incubated with either 10 µM JFX646-HaloTag or JFX549-HaloTag for 30-60 minutes at 4°C. The resin was then washed four more times in TEV digest buffer, then incubated in TEV buffer supplemented with TEV protease overnight at 4°C. Following TEV digest, the protein-containing supernatant was collected using a spin filtration device, aliquoted, flash frozen in liquid nitrogen, and stored at -80ºC.

Microtubules Preparation

Porcine tubulin, obtained either from Cytoskeleton or purified in-house, was resuspended in MT buffer (25 mM MES, 70 mM NaCl, 1 mM MgCl2, 1 mM EGTA, and 1 mM DTT, pH 6.5) to a final concentration of 10 mg/ml, flash-frozen, and stored at -80°C. Microtubules were polymerized using a previously described protocol *(15)*. In brief, tubulin was diluted in MT buffer supplemented with 3 mM GTP. The mixture was incubated on ice for 5 minutes and subsequently transferred to a 37°C incubator for 1 hour. Following polymerization, 5 µM paclitaxel (MilliporeSigma) was added to the mixture. The microtubules were then pelleted (20,000 rcf for 8 minutes at room temperature) and resuspended in MT buffer containing 5 µM paclitaxel.

ATPase measurements

Basal and microtubule-stimulated ATPase activities were measured using the EnzChek phosphate assay kit (Invitrogen). Assays were performed in motility buffer (30 mM HEPES, pH 7.2, 50 mM potassium acetate, 2 mM magnesium acetate, 1 mM EGTA, 10% glycerol) supplemented with 1 mM Mg ATP, with or without 2 µM taxol-stabilized microtubules and 5 nM 6His-GST-dyneinMOTOR (wild-type or mutants). Reactions were initiated with the addition of dynein, and the absorbance at 360 nm was monitored by plate reader (BMG LabTech CLARIOstar Plus) for at least 20 min. Background phosphate release levels (presumably from microtubules) for each reaction were measured for 2-5 min before addition of dynein motors. Relative ATPase levels were determined by fitting linear regressions to the time-resolved absorbance values, and determining the slope from each. Data shown are from 4-6 independent replicates, from two independent protein preparations.

Single-molecule motility assays

Single-molecule motility assays were performed as previously described *(17, 23)* with minor modifications. In brief, flow chambers constructed using slides and plasma cleaned and silanized coverslips attached with double-sided adhesive tape were coated with anti-tubulin antibodies (8 µg ml1, YL1/2; Accurate Chemical Scientific Corporation) and then blocked with 1% Pluronic F-127 (Fisher Scientific). Taxol-stabilized microtubules assembled from unlabelled porcine tubulin (Cytoskeleton) were introduced into the chamber. Af-

ter incubation for 2-5 min, the chamber was washed with motility buffer (see above) supplemented with 20 μM taxol and 1 mM DTT. Dynein, diluted in an oxygen-scavenging motility buffer (30 mM HEPES, pH 7.2, 50 mM potassium acetate, 2 mM magnesium acetate, 1 mM EGTA, 10% glycerol, 50 nM protocatechuate 3,4-dioxygenase, 2.5 mM protocatechuic acid, 1 mM Trolox, 1 mM cyclooctatetraene, 1 mM 4-nitrobenzyoyl alcohol) supplemented with 1 mM DTT, 20 μM taxol, and 1 mM Mg-ATP was then added. TIRFM images were immediately collected using an iLAS2 RING TIRF system on a Nikon Ti-2E inverted microscope equipped with a 1.49 NA 100X TIRF objective, a Ti2-SS-E motorized stage, piezo Z-control, and an iXon LIFE 897 cooled EM-CCD camera (Andor). 561 or 640 nm lasers housed in an LUN-F 3 (Nikon) were used along with a multi-pass quad filter cube set (C-TIRF for 405/488/561/638 nm; Chroma) and emission filters mounted in a filter wheel (600/50 nm and 700/75 nm; Chroma). To determine microtubule polarity (due to the diffusive/bidirectional behavior of the Y1902A mutant), wild-type GST-dyneinMOTOR-HaloJF549 (diluted in oxygen-scavenging motility buffer) was introduced into the imaging chamber immediately after acquiring a movie of the mutant GST-dyneinMOTOR-HaloJF646. To image non-fluorescent microtubules, we used interference reflection microscopy, as previously described *(89)*. Images of dynein motors were acquired at 2 second intervals for 8min. Velocity and run length values were determined from kymographs generated using the MultipleKymograph plugin for FIJI/ImageJ. Data shown are from 3 independent replicates, from 2 independent protein preparations.

Cryo-EM sample and grid preparation for dynein unbound from microtubules

For dynein samples in AMPPNP, ATP-Vi, and ADP conditions, the concentrated dynein obtained after gel filtration was diluted with GF150 buffer containing 0.1 mM ATP to a concentration of 2.0 mg/ml. Subsequently, ATP-vanadate, AMPPNP, or ADP was added to a final concentration of 5 mM, and the mixture was incubated with dynein for 1 hour on ice before vitrification. In the case of dynein in ATP condition, the protein was incubated with 5 mM ATP on ice and immediately vitrified. For dynein in apo buffer (GF150 buffer without ATP), the protein was diluted into the apo buffer and concentrated using an Amicon concentrator. This concentration process was repeated three times, resulting in an estimated 8000-fold dilution of the original 0.1 mM ATP. The final concentration of dynein in the apo buffer was adjusted to 2 mg/ml for vitrification. Prepared samples were applied to Quantifoil holy carbon grids (R2/1, 300 mesh gold or R2/2, 200 mesh) and incubated in the chamber of the Vitrobot Mark IV unit (FEI) for 5 seconds at 4°C and 100% humidity. Subsequently, the grids were blotted for 3-5 seconds and plunged into liquid ethane.

Cryo-EM sample and grid preparation for dynein bound to microtubules

TFor the reconstitution of dynein-MT complexes **(Figure S2)** in apo condition, dynein in apo buffer was mixed with microtubules (MTs) at room temperature in MT-binding buffer (30 mM HEPES pH 7.2, 60 mM KCl, 1 mM EGTA, 5 mM MgSO4, 1 mM DTT, 5 μM paclitaxel). The dynein-MT complex was then pelleted at 20,000r.c.f. for 8minutes. The optimal resuspension volume, which determined the final concentration used for cryo-EM vitrification, was estimated using negative staining electron microscopy. For dynein-MT complexes in ADP or AMPPNP condition, the final MT binding buffer was supplemented with 5 mM ADP or 5 mM AMPPNP. A volume of 3.5 μL of the dynein-MT complex was applied to Quantifoil holy carbon grids (R2/1, 300 mesh gold), followed by a 4-second wait time, blotting for 3-6 seconds, and subsequent freezing in liquid ethane using a Vitrobot Mark IV unit (FEI). The Vitrobot chamber was maintained at close to 95% humidity and 22°C during the process.

Cryo-EM data collection

All cryo-EM grids underwent initial screening at the Yale ScienceHill-Cryo-EM facility using a Glacios microscope (Thermo Fisher Scientific). Subsequent cryo-EM data were collected at three different sites: Yale ScienceHill-Cryo-EM facility, utilizing a Glacios microscope operated at 300 keV with a K3 detector; Yale WestCampus-Cryo-EM facility, where a Krios microscope operated at 300 keV equipped with a K3 detector and a Bioquantum Energy Filter was utilized; and the Laboratory for BioMolecular Structure at BNL with a Krios microscope operated at 300 keV with a K3 detector and a Bioquantum Energy Filter. Automatic data collection was facilitated by either SerialEM *(90)* or EPU software. In total, 33,302, 6,016, 7,488, 5,920, and 5,884 movies were acquired for dynein-ATP, dynein-ATPVi, dynein-AMPPNP, dynein-ADP, and dynein-Apo off microtubules, respectively. Additionally, 9,935, 11,828, and 5,690 movies were acquired for dynein-Apo-MT, dynein-ADP-MT, and dynein-AMPPNP-MT on microtubules, respectively. Detailed data collection parameters can be found in supplementary tables S1-S7 corresponding to different datasets.

Cryo-EM image processing for dynein unbound from microtubules

Preprocessing steps, including motion correction, CTF estimation, and particle picking, were conducted either in cryoSPARC Live *(91)* or via an in-house script utilizing MotionCor2 *(92)*, GCTF *(93)*, and Gautomatch. Cryo-EM scripts for real-time data transfer and on-the-fly preprocessing are available for download at https://github.com/JackZhang-Lab.

For the dynein-ATP off microtubules dataset **(Figure S1)**, the blob picker was utilized to select motor domains in phi or open conformations, with the cross-correlation score set to 0.01 to enable sampling of all possible motors during the picking step. A total of 8,534,500 particles were ex-

tracted using a box size of 400 pixels, subsequently binned to 200 pixels (1.664 Å pixel size) for faster computation. From this pool, a subset of 640,000 particles underwent subsequent rounds of 2D classification, ultimately yielding 164,658 high-quality particles selected for ab initio reconstruction (6 classes). During this phase, phi motor domains and several single motor domains were identified. Following the initial model acquisition, all original particles were employed for subsequent heterogeneous refinement to prevent the loss of motor particles during 2D classification. The original particles were divided into four subsets, with each subset subjected to heterogeneous refinement (12 classes). This initial heterogeneous refinement process was iterated three times, with updated references provided each time, successfully identifying phi-motors, single motor domains with bent or straight linkers, and several junk classes.

Each motor class from four subsets was merged for additional heterogeneous refinement or 3D classification without alignment. This step further explores the dynamics of the motor domain. For example, state-4, characterized by a straight linker, was initially obscured within the state-2 class due to their closed motor ring conformation. Similarly, state-5, state-6, and state-7 were initially grouped into a single class and later separated during subsequent heterogeneous refinement. The second-level heterogeneous refinement was conducted three times, with updated references provided each time. Further classification of individual classes ceased to reveal novel conformations due to limited particle numbers and achievable resolution. An increase in dataset size may unveil additional intermediate or transient conformations.

Following homogeneous refinement, most classes achieved resolutions at the Nyquist limit of 3.3 Å. Subsequently, particles were re-extracted at the full box size of 400 pixels (0.832 Å). For high-resolution refinement, two rounds of global and local CTF refinement followed by local refinement were performed. The resolution of single motor domains from open dynein reached 2.9-3.6 Å, while phi-motor domains achieved 2.6 Å using C2 symmetry. To further enhance the resolution of phi-motor domains, symmetry expansion of the phi-motor domain was executed to account for inter-motor domain dynamics. Particles were recentered on one of the motor domains and re-extracted using a box size of 640 pixels to incorporate delocalized high-resolution information. After localized refinement and another round of global and local CTF refinement, the phi-motor domain achieved 2.25 Å resolution.

For full-length phi dynein reconstruction, two sets of particles were considered. Firstly, junk particles from the initial refinement were merged and subjected to heterogeneous refinement using the dynein tail map (EMD-3703 (10)) and junk volumes as references, yielding classes with recognizable tail domains. Secondly, particles from phi dynein motor domains were recentered at the tail and re-extracted (box size 400 pixels binned to 200 pixels). These two sets of particles were merged for additional heterogeneous refinement

focusing on the tail region, resulting in classes with clear tail domain features. The selected class was re-extracted using a large box size of 1024 pixels (binned to 256 pixels) for full-length dynein refinement. Duplicated particles were removed during re-extraction. After another heterogeneous refinement, the class with both stable tail and motor domain was selected and refined to 6.8 Å resolution. To improve the resolution of the tail domain, particles were recentered on the tail and re-extracted using a box size of 384 pixels. A mask covering only the tail domain was created, and after local refinement, the entire tail domain reached 4.3 Å resolution, albeit with large variations in resolution among different tail regions. Masks covering the NDD region, middle, and neck regions were used to improve local resolution. The middle region of the tail was refined to 3.8 Å, revealing side chain density, while the NDD and neck regions reached 6.5 Å with resolved secondary structures. Localized refined maps were fitted to the consensus map and stitched together in ChimeraX (94) using the "vop max" command to create a composite map of full-length human phi dynein.

Datasets of dynein under ATPVi, AMPPNP, ADP, and apo conditions underwent processing using the same procedures.

Cryo-EM image processing of dynein bound to microtubules

For cryo-EM image processing of dynein-MT datasets, we employed a pipeline previously developed for MT tracing and MT-signal subtraction (**Figure S2**) (71, 72). Initially, MT particles were detected using template matching with a distance cut-off of 8 nm, with a relatively low cross-correlation score set to include MT particles with low contrast or located at cross-over locations. Following three rounds of 2D classification, false-picked MT particles, including carbon edges, and mis-centered MTs were rejected. The metadata of the selected particles was converted to Relion style using "csparc2star.py" in pyem (https://zenodo.org/records/3576630), and the coordinates were recentered and subjected to "multi-curve fitting". After MT-signal subtraction, original micrographs were replaced by MT-signal subtracted micrographs.

Subsequently, the blob-picker was used to detect motor particles in these new micrographs, which were then extracted using a box size covering the single motor domain. 2D classification was utilized to remove junk particles and residual MT particles, after which the selected particles underwent ab initio reconstruction. Similar to our above workflow, all original particles were used for initial heterogeneous refinement. 3D classification-based particle sorting greatly aided in identifying motor particles with low signal-to-noise ratios due to residual MT signal influence. After several rounds of sorting, classes with clear motor features were selected and subjected to local refinement, global, and local CTF refinement. Notably, the post-1 linker-docking mode was observed in three datasets, while the post-2 was exclu-

sively found in the dynein-MT-ADP dataset. The resolution of individual motor domains ranged from 2.9 Å to 3.6 Å (**Figure S2, table S6**).

For the reconstruction of full-length dynein bound to MT, particles were reextracted using a large box size covering two motor domains. The particles were directly reconstructed into a volume featuring the weak density of the second motor. Subsequently, this map was low-pass filtered to 60 Å for heterogeneous refinement, where some classes exhibited enhanced density for the second motor. We then manually fitted the motor map into the weak density to create a two-motor map. This map, together with previous reconstructions, was used for another round of heterogeneous refinement. The newly generated maps served as new references for iterative rounds of heterogeneous refinement. Ultimately, three main states were predominantly observed for each dynein-MT dataset: i) two stable and parallel heads, ii) one stable leading head, and iii) one stable trailing head. However, due to the limited number of particles and resolution, we were unable to detect two motors with different step sizes in our dynein-ADP-MT datasets (**Figure S2**).

Additionally, we attempted to reconstruct dynein-MTBD-MT by reverting the MT signal using the original micrographs. Unlike our previously determined out-arm dynein bound to protofilament structures featuring defined tubulin subunit density *(69)*, the protofilament of dynein-1 bound to MT exhibited smeared density. Furthermore, subsequent 3D classification without alignment failed to generate a relatively stable stalk-MTBD-protofilament class due to the limited number of particles. These limitations precluded us from performing localized refinement on the MTBD-MT region.

Model building and refinement

For model building and refinement, we utilized two previously reported human dynein motor structures (PDB: 5NUG *(10)*, 7Z8G *(15)*) as initial models for different states of dynein in ATP. Individual domains (linker, AAA large, AAA small) were rigidly docked into the cryo-EM map in UCSF ChimeraX. Subsequently, Namdinator *(95)*, a molecular dynamics flexible fitting tool, was employed to further refine the model into the cryo-EM map. Manual inspection and adjustments of the model were carried out in COOT v0.9.5 *(96, 97)*.

For water model building in the phi-motor, the maps underwent sharpening using two methods: i) different B-factor sharpening with local filtering based on local resolution in cryoSPARC, and ii) density modification with phenix.resolve-cryo-em *(98)* using only maps as input. The processed maps were up-sampled with a pixel size of 0.416 Å and cropped to a box size of 400 pixels. Initial solvent water molecules were placed using phenix.douse, resulting in approximately 2358 automatically placed water molecules. Subsequent manual inspection in COOT led to the addition or deletion of water molecules, resulting in a final placement of 1897 water molecules.

All models underwent iterative refinement using Phenix real-space refinement 1.21rc1-5190 and manual rebuilding in COOT. These refined models were utilized as initial models for dynein in other nucleotide conditions or dynein bound to MTs. The quality of the refined models was assessed using MolProbity *(99)* integrated into Phenix *(98)*, with statistics reported in **Table S1-S7**.

Plots, and molecular graphics

Plots were generated using GraphPad Prism. Molecular graphics were prepared using UCSF ChimeraX *(94)*.

References

1. S. M. King. The dynein microtubule motor. *Biochim Biophys Acta*, 1496(1): 60–75, 2000. ISSN 0006-3002 (Print) 0006-3002 (Linking). doi: 10.1016/s0167-4889(00)00009-4.

2. S. L. Reck-Peterson, W. B. Redwine, R. D. Vale, and A. P. Carter. The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol*, 19(6):382–398, 2018. ISSN 1471-0080 (Electronic) 1471-0072 (Linking). doi: 10.1038/s41580-018-0004-3.

3. T. Torisawa and A. Kimura. The generation of dynein networks by multi-layered regulation and their implication in cell division. *Front Cell Dev Biol*, 8:22, 2020. ISSN 2296-634X (Print) 2296-634X (Electronic) 2296-634X (Linking). doi: 10.3389/fcell.2020.00022.

4. D. Ricolo, J. Castro-Ribera, and S. J. Araujo. Cytoskeletal players in single-cell branching morphogenesis. *Dev Biol*, 477:22–34, 2021. ISSN 1095-564X (Electronic) 0012-1606 (Linking). doi: 10.1016/j.ydbio.2021.05.001.

5. N. Klena and G. Pigino. Structural biology of cilia and intraflagellar transport. *Annu Rev Cell Dev Biol*, 2022. ISSN 1530-8995 (Electronic) 1081-0706 (Linking). doi: 10.1146/annurev-cellbio-120219-034238.

6. S. M. King. Axonemal dynein arms. *Cold Spring Harb Perspect Biol*, 8 (11), 2016. ISSN 1943-0264 (Electronic) 1943-0264 (Linking). doi: 10.1101/cshperspect.a028100.

7. A. Bachmann and A. Straube. Kinesins in cell migration. *Biochem Soc Trans*, 43(1):79–83, 2015. ISSN 1470-8752 (Electronic) 0300-5127 (Linking). doi: 10.1042/BST20140280.

8. L. Urnavicius, C. K. Lau, M. M. Elshenawy, E. Morales-Rios, C. Motz, A. Yildiz, and A. P. Carter. Cryo-em shows how dynactin recruits two dyneins for faster movement. *Nature*, 554(7691):202–206, 2018. ISSN 1476-4687 (Electronic) 0028-0836 (Linking). doi: 10.1038/nature25462.

9. D. A. Grotjahn, S. Chowdhury, Y. Xu, R. J. McKenney, T. A. Schroer, and G. C. Lander. Cryo-electron tomography reveals that dynactin recruits a team of dyneins for processive motility. *Nat Struct Mol Biol*, 25(3):203–207, 2018. ISSN 1545-9985 (Electronic) 1545-9985 (Linking). doi: 10.1038/s41594-018-0027-7.

10. K. Zhang, H. E. Foster, A. Rondelet, S. E. Lacey, N. Bahi-Buisson, A. W. Bird, and A. P. Carter. Cryo-em reveals how human cytoplasmic dynein is auto-inhibited and activated. *Cell*, 169(7):1303–1314 e18, 2017. ISSN 1097-4172 (Electronic) 0092-8674 (Linking). doi: 10.1016/j.cell.2017.05.025.

11. L. Urnavicius, K. Zhang, A. G. Diamant, C. Motz, M. A. Schlager, M. Yu, N. A. Patel, C. V. Robinson, and A. P. Carter. The structure of the dynactin complex and its interaction with dynein. *Science*, 347(6229):1441–1446, 2015. ISSN 1095-9203 (Electronic) 0036-8075 (Linking). doi: 10.1126/science.aaa4080.

12. R. J. McKenney, W. Huynh, M. E. Tanenbaum, G. Bhabha, and R. D. Vale. Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science*, 345(6194):337–41, 2014. ISSN 1095-9203 (Electronic) 0036-8075 (Linking). doi: 10.1126/science.1254198.

13. S. Chowdhury, S. A. Ketcham, T. A. Schroer, and G. C. Lander. Structural organization of the dynein-dynactin complex bound to microtubules. *Nat Struct Mol Biol*, 22(4):345–7, 2015. ISSN 1545-9985 (Electronic) 1545-9985 (Linking). doi: 10.1038/nsmb.2996.

14. M. A. Schlager, H. T. Hoang, L. Urnavicius, S. L. Bullock, and A. P. Carter. In vitro reconstitution of a highly processive recombinant human dynein complex. *EMBO J*, 33(17):1855–68, 2014. ISSN 1460-2075 (Electronic) 0261-4189 (Linking). doi: 10.15252/embj.201488792.

15. S. Chaaban and A. P. Carter. Structure of dynein-dynactin on microtubules shows tandem adaptor binding. *Nature*, 2022. ISSN 1476-4687 (Electronic) 0028-0836 (Linking). doi: 10.1038/s41586-022-05186-y.

16. Z. M. Htet, J. P. Gillies, R. W. Baker, A. E. Leschziner, M. E. DeSantis, and S. L. Reck-Peterson. Lis1 promotes the formation of activated cytoplasmic dynein-1 complexes. *Nat Cell Biol*, 22(5):518–525, 2020. ISSN 1476-4679 (Electronic) 1465-7392 (Linking). doi: 10.1038/s41556-020-0506-z.

17. M. G. Marzo, J. M. Griswold, and S. M. Markus. Pac1/lis1 stabilizes an uninhibited conformation of dynein to coordinate its localization and activity. *Nat Cell Biol*, 22(5):559–569, 2020. ISSN 1476-4679 (Electronic) 1465-7392 (Linking). doi: 10.1038/s41556-020-0492-1.

18. M. M. Elshenawy, E. Kusakci, S. Volz, J. Baumbach, S. L. Bullock, and A. Yildiz. Lis1 activates dynein motility by modulating its pairing with dynactin. *Nat Cell Biol*, 22(5):570–578, 2020. ISSN 1476-4679 (Electronic) 1465-7392 (Linking). doi: 10.1038/s41556-020-0501-4.

19. R. J. McKenney. Lis1 cracks open dynein. *Nat Cell Biol*, 22(5):515–517, 2020. ISSN 1476-4679 (Electronic) 1465-7392 (Linking). doi: 10.1038/s41556-020-0500-5.

20. S. M. Markus, M. G. Marzo, and R. J. McKenney. New insights into the mechanism of dynein motor regulation by lissencephaly-1. *Elife*, 9, 2020. ISSN 2050-084X (Electronic) 2050-084X (Linking). doi: 10.7554/eLife.59737.

21. W. D. Ton, Y. Wang, P. Chai, C. Beauchamp-Perez, N. T. Flint, L. G. Lammers, H. Xiong, K. Zhang, and S. M. Markus. Microtubule-binding-induced allostery triggers lis1 dissociation from dynein prior to cargo transport. *Nat Struct Mol Biol*, 30(9):1365–1379, 2023. ISSN 1545-9985 (Electronic) 1545-9993 (Print) 1545-9985 (Linking). doi: 10.1038/s41594-023-01010-x.

22. Y. Zhao, S. Oten, and A. Yildiz. Nde1 promotes lis1-mediated activation of dynein. *Nat Commun*, 14(1):7221, 2023. ISSN 2041-1723 (Electronic) 2041-1723 (Linking). doi: 10.1038/s41467-023-42907-x.

23. K. Okada, B. R. Iyer, L. G. Lammers, P. A. Gutierrez, W. Li, S. M. Markus, and R. J. McKenney. Conserved roles for the dynein intermediate chain and ndel1 in assembly and activation of dynein. *Nat Commun*, 14(1):5833, 2023. ISSN 2041-1723 (Electronic) 2041-1723 (Linking). doi: 10.1038/s41467-023-41466-5.

24. J. P. Pandey, L. Shi, R. A. Brebion, and D. S. Smith. Lis1 and ndel1 regulate axonal trafficking of mitochondria in mature neurons. *Front Mol Neurosci*, 15: 841047, 2022. ISSN 1662-5099 (Print) 1662-5099 (Electronic) 1662-5099 (Linking). doi: 10.3389/fnmol.2022.841047.

25. Y. Chu, G. A. Morfini, L. B. Langhamer, Y. He, S. T. Brady, and J. H. Kordower. Alterations in axonal transport motor proteins in sporadic and experimental parkinson's disease. *Brain*, 135(Pt 7):2058–73, 2012. ISSN 1460-2156 (Electronic) 0006-8950 (Print) 0006-8950 (Linking). doi: 10.1093/brain/aws133.

26. H. T. Hoang, M. A. Schlager, A. P. Carter, and S. L. Bullock. Dync1h1 mutations associated with neurological diseases compromise processivity of dynein-dynactin-cargo adaptor complexes. *Proc Natl Acad Sci U S A*, 114(9): E1597–E1606, 2017. ISSN 1091-6490 (Electronic) 0027-8424 (Linking). doi: 10.1073/pnas.1620141114.

27. O. Reiner, R. Carrozzo, Y. Shen, M. Wehnert, F. Faustinella, W. B. Dobyns, C. T. Caskey, and D. H. Ledbetter. Isolation of a miller-dieker lissencephaly gene containing g protein beta-subunit-like repeats. *Nature*, 364(6439):717–21, 1993. ISSN 0028-0836 (Print) 0028-0836 (Linking). doi: 10.1038/364717a0.

28. M. G. Marzo, J. M. Griswold, K. M. Ruff, R. E. Buchmeier, C. P. Fees, and S. M. Markus. Molecular basis for dyneinopathies reveals insight into dynein regulation and dysfunction. *Elife*, 8, 2019. ISSN 2050-084X (Electronic) 2050-084X (Linking). doi: 10.7554/eLife.47246.

29. J. Scherer, J. Yi, and R. B. Vallee. Role of cytoplasmic dynein and kinesins in adenovirus transport. *FEBS Lett*, 594(12):1838–1847, 2020. ISSN 1873-3468 (Electronic) 0014-5793 (Linking). doi: 10.1002/1873-3468.13777.

30. I. Hassan Ibrahim, A. Balah, A. Gomaa Abd Elfattah Hassan, and H. Gamal Abd El-Aziz. Role of motor proteins in human cancers. *Saudi J Biol Sci*, 29(12):103436, 2022. ISSN 1319-562X (Print) 2213-7106 (Electronic) 2213-7106 (Linking). doi: 10.1016/j.sjbs.2022.103436.

31. H. Schmidt and A. P. Carter. Review: Structure and mechanism of the dynein motor atpase. *Biopolymers*, 105(8):557–67, 2016. ISSN 1097-0282 (Electronic) 0006-3525 (Linking). doi: 10.1002/bip.22856.

32. M. A. Cianfrocco, M. E. DeSantis, A. E. Leschziner, and S. L. Reck-Peterson. Mechanism and regulation of cytoplasmic dynein. *Annu Rev Cell Dev Biol*, 31:83–108, 2015. ISSN 1530-8995 (Electronic) 1081-0706 (Linking). doi: 10.1146/annurev-cellbio-100814-125438.

33. L. Rao and A. Gennerich. Structure and function of dynein's non-catalytic subunits. *Cells*, 13(4), 2024. ISSN 2073-4409 (Electronic) 2073-4409 (Linking). doi: 10.3390/cells13040330.

34. N. Numata, T. Shima, R. Ohkura, T. Kon, and K. Sutoh. C-sequence of the dictyostelium cytoplasmic dynein participates in processivity modulation. *FEBS Lett*, 585(8):1185–90, 2011. ISSN 1873-3468 (Electronic) 0014-5793 (Linking). doi: 10.1016/j.febslet.2011.03.036.

35. C. Cho, S. L. Reck-Peterson, and R. D. Vale. Regulatory atpase sites of cytoplasmic dynein affect processivity and force generation. *J Biol Chem*, 283(38):25839–45, 2008. ISSN 0021-9258 (Print) 0021-9258 (Linking). doi: M802951200[pii]10.1074/jbc.M802951200.

36. T. Kon, T. Mogami, R. Ohkura, M. Nishiura, and K. Sutoh. Atp hydrolysis cycle-dependent tail motions in cytoplasmic dynein. *Nat Struct Mol Biol*, 12(6):513–9, 2005. ISSN 1545-9993 (Print) 1545-9985 (Linking). doi: 10.1038/nsmb930.

37. T. Kon, M. Nishiura, R. Ohkura, Y. Y. Toyoshima, and K. Sutoh. Distinct functions of nucleotide-binding/hydrolysis sites in the four aaa modules of cytoplasmic dynein. *Biochemistry*, 43(35):11266–74, 2004. ISSN 0006-2960 (Print) 0006-2960 (Linking). doi: 10.1021/bi048985a.

38. T. Kon, K. Imamula, A. J. Roberts, R. Ohkura, P. J. Knight, I. R. Gibbons, S. A. Burgess, and K. Sutoh. Helix sliding in the stalk coiled coil of dynein couples atpase and microtubule binding. *Nat Struct Mol Biol*, 16(3):325–33, 2009. ISSN 1545-9985 (Electronic) 1545-9985 (Linking).

39. A. Gennerich, A. P. Carter, S. L. Reck-Peterson, and R. D. Vale. Force-induced bidirectional stepping of cytoplasmic dynein. *Cell*, 131(5):952–65, 2007. ISSN 0092-8674 (Print) 0092-8674 (Linking). doi: 10.1016/j.cell.2007.10.016.

40. T. Mogami, T. Kon, K. Ito, and K. Sutoh. Kinetic characterization of tail swing steps in the atpase cycle of dictyostelium cytoplasmic dynein. *J Biol Chem*, 282(30):21639–44, 2007. ISSN 0021-9258 (Print) 0021-9258 (Linking). doi: 10.1074/jbc.M701914200.

41. S. L. Reck-Peterson, A. Yildiz, A. P. Carter, A. Gennerich, N. Zhang, and R. D. Vale. Single-molecule analysis of dynein processivity and stepping behavior. *Cell*, 126(2):335–48, 2006. ISSN 0092-8674 (Print) 0092-8674 (Linking). doi: 10.1016/j.cell.2006.05.046.

42. H. Schmidt, R. Zalyte, L. Urnavicius, and A. P. Carter. Structure of human cytoplasmic dynein-2 primed for its power stroke. *Nature*, 518(7539):435–438, 2015. ISSN 1476-4687 (Electronic) 0028-0836 (Print) 0028-0836 (Linking). doi: 10.1038/nature14023.

43. A. P. Carter. Crystal clear insights into how the dynein motor moves. *J Cell Sci*, 126(Pt 3):705–13, 2013. ISSN 1477-9137 (Electronic) 0021-9533 (Linking). doi: 10.1242/jcs.120725.

44. T. Kon, T. Oyama, R. Shimo-Kon, K. Imamula, T. Shima, K. Sutoh, and G. Kurisu. The 2.8 a crystal structure of the dynein motor domain. *Nature*, 484(7394):345–50, 2012. ISSN 1476-4687 (Electronic) 0028-0836 (Linking). doi: 10.1038/nature10955.

45. H. Schmidt, E. S. Gleave, and A. P. Carter. Insights into dynein motor domain function from a 3.3-a crystal structure. *Nat Struct Mol Biol*, 19(5):492–7, S1, 2012. ISSN 1545-9985 (Electronic) 1545-9993 (Print) 1545-9985 (Linking). doi: 10.1038/nsmb.2272.

46. A. P. Carter, C. Cho, L. Jin, and R. D. Vale. Crystal structure of the dynein motor domain. *Science*, 331(6021):1159–65, 2011. ISSN 1095-9203 (Electronic) 0036-8075 (Linking). doi: 10.1126/science.1202393.

47. G. Bhabha, H. C. Cheng, N. Zhang, A. Moeller, M. Liao, J. A. Speir, Y. Cheng, and R. D. Vale. Allosteric communication in the dynein motor domain. *Cell*, 159(4):857–68, 2014. ISSN 1097-4172 (Electronic) 0092-8674 (Linking). doi: 10.1016/j.cell.2014.10.018.

48. J. T. Canty, R. Tan, E. Kusakci, J. Fernandes, and A. Yildiz. Structure and mechanics of dynein motors. *Annu Rev Biophys*, 50:549–574, 2021. ISSN 1936-1238 (Electronic) 1936-122X (Linking). doi: 10.1146/annurev-biophys-111020-101511.

49. A. J. Roberts, B. Malkova, M. L. Walker, H. Sakakibara, N. Numata, T. Kon, R. Ohkura, T. A. Edwards, P. J. Knight, K. Sutoh, K. Oiwa, and S. A. Burgess. Atp-driven remodeling of the linker domain in the dynein motor. *Structure*, 20 (10):1670–80, 2012. ISSN 1878-4186 (Electronic) 0969-2126 (Print) 0969-2126 (Linking). doi: 10.1016/j.str.2012.07.003.

50. S. A. Burgess, M. L. Walker, H. Sakakibara, P. J. Knight, and K. Oiwa. Dynein structure and power stroke. *Nature*, 421(6924):715–8, 2003. ISSN 0028-0836 (Print) 0028-0836 (Linking). doi: 10.1038/nature01377.

51. H. L. Sweeney and A. Houdusse. Structural and functional insights into the myosin motor mechanism. *Annu Rev Biophys*, 39:539–57, 2010. ISSN 1936-1238 (Electronic) 1936-122X (Linking). doi: 10.1146/annurev.biophys.050708.133751.

52. W. Oosterheert, F. E. C. Blanc, A. Roy, A. Belyy, M. B. Sanders, O. Hofnagel, G. Hummer, P. Bieling, and S. Raunser. Molecular mechanisms of inorganic-phosphate release from the core and barbed end of actin filaments. *Nat Struct Mol Biol*, 30(11):1774–1785, 2023. ISSN 1545-9985 (Electronic) 1545-9993 (Print) 1545-9985 (Linking). doi: 10.1038/s41594-023-01101-9.

53. J. D. Moore and S. A. Endow. Kinesin proteins: A phylum of motors for microtubule-based motility. *Bioessays*, 18(3):207–219, 1996. ISSN 0265-9247. doi: DOI10.1002/bies.950180308.

54. S. Can, S. Lacey, M. Gur, A. P. Carter, and A. Yildiz. Directionality of dynein is controlled by the angle and length of its stalk. *Nature*, 566(7744):407–410, 2019. ISSN 1476-4687 (Electronic) 0028-0836 (Linking). doi: 10.1038/s41586-019-0914-z.

55. M. A. DeWitt, C. A. Cypranowska, F. B. Cleary, V. Belyy, and A. Yildiz. The aaa3 domain of cytoplasmic dynein acts as a switch to facilitate microtubule release. *Nat Struct Mol Biol*, 22(1):73–80, 2015. ISSN 1545-9985 (Electronic) 1545-9985 (Linking). doi: 10.1038/nsmb.2930.

56. A. P. Carter and R. D. Vale. Communication between the aaa+ ring and microtubule-binding domain of dynein. *Biochem Cell Biol*, 88(1):15–21, 2010. ISSN 1208-6002 (Electronic) 0829-8211 (Linking). doi: 10.1139/o09-127.

57. N. Nishida, Y. Komori, O. Takarada, A. Watanabe, S. Tamura, S. Kubo, I. Shimada, and M. Kikkawa. Structural basis for two-way communication between dynein and microtubules. *Nat Commun*, 11(1):1038, 2020. ISSN 2041-1723 (Electronic) 2041-1723 (Linking). doi: 10.1038/s41467-020-14842-8.

58. S. Niekamp, N. Coudray, N. Zhang, R. D. Vale, and G. Bhabha. Coupling of atpase activity, microtubule binding, and mechanics in the dynein motor domain. *EMBO J*, 38(13):e101414, 2019. ISSN 1460-2075 (Electronic) 0261-4189 (Linking). doi: 10.15252/embj.2018101414.

59. Y. S. Kato, T. Yagi, S. A. Harris, S. Y. Ohki, K. Yura, Y. Shimizu, S. Honda, R. Kamiya, S. A. Burgess, and M. Tanokura. Structure of the microtubule-binding domain of flagellar dynein. *Structure*, 22(11):1628–38, 2014. ISSN 1878-4186 (Electronic) 0969-2126 (Linking). doi: 10.1016/j.str.2014.08.021.

60. Y. Nishikawa, T. Oyama, N. Kamiya, T. Kon, Y. Y. Toyoshima, H. Nakamura, and G. Kurisu. Structure of the entire stalk region of the dynein motor domain. *J Mol Biol*, 426(19):3232–3245, 2014. ISSN 1089-8638 (Electronic) 0022-2836 (Linking). doi: 10.1016/j.jmb.2014.06.023.

61. W. B. Redwine, R. Hernandez-Lopez, S. Zou, J. Huang, S. L. Reck-Peterson, and A. E. Leschziner. Structural basis for microtubule binding and release by dynein. *Science*, 337(6101):1532–1536, 2012. ISSN 1095-9203 (Electronic)

0036-8075 (Print) 0036-8075 (Linking). doi: 10.1126/science.1224151.

62. A. P. Carter, J. E. Garbarino, E. M. Wilson-Kubalek, W. E. Shipley, C. Cho, R. A. Milligan, R. D. Vale, and I. R. Gibbons. Structure and functional role of dynein's microtubule-binding domain. *Science*, 322(5908):1691–5, 2008. ISSN 1095-9203 (Electronic) 0036-8075 (Linking). doi: 10.1126/science.1164424.

63. S. E. Lacey, S. He, S. H. Scheres, and A. P. Carter. Cryo-em of dynein microtubule-binding domains shows how an axonemal dynein distorts the microtubule. *Elife*, 8, 2019. ISSN 2050-084X (Electronic) 2050-084X (Linking). doi: 10.7554/eLife.47145.

64. S. Uchimura, T. Fujii, H. Takazaki, R. Ayukawa, Y. Nishikawa, I. Minoura, Y. Hachikubo, G. Kurisu, K. Sutoh, T. Kon, K. Namba, and E. Muto. A flipped ion pair at the dynein-microtubule interface is critical for dynein motility and atpase activation. *J Cell Biol*, 208(2):211–22, 2015. ISSN 1540-8140 (Electronic) 0021-9525 (Print) 0021-9525 (Linking). doi: 10.1083/jcb.201407039.

65. L. Li, J. Alper, and E. Alexov. Cytoplasmic dynein binding, run length, and velocity are guided by long-range electrostatic interactions. *Sci Rep*, 6:31523, 2016. ISSN 2045-2322 (Electronic) 2045-2322 (Linking). doi: 10.1038/srep31523.

66. E. L. Holzbaur and K. A. Johnson. Microtubules accelerate adp release by dynein. *Biochemistry*, 28(17):7010–6, 1989. ISSN 0006-2960 (Print) 0006-2960 (Linking). doi: 10.1021/bi00443a034.

67. L. A. Amos. Brain dynein crossbridges microtubules into bundles. *Journal of Cell Science*, 93:19–28, 1989. ISSN 0021-9533.

68. K. Toropova, R. Zalyte, A. G. Mukhopadhyay, M. Mladenov, A. P. Carter, and A. J. Roberts. Structure of the dynein-2 complex and its assembly with intraflagellar transport trains. *Nat Struct Mol Biol*, 26(9):823–829, 2019. ISSN 1545-9985 (Electronic) 1545-9985 (Linking). doi: 10.1038/s41594-019-0286-y.

69. Q. Rao, L. Han, Y. Wang, P. Chai, Y. W. Kuo, R. Yang, F. Hu, Y. Yang, J. Howard, and K. Zhang. Structures of outer-arm dynein array on microtubule doublet reveal a motor coordination mechanism. *Nat Struct Mol Biol*, 28(10):799–810, 2021. ISSN 1545-9985 (Electronic) 1545-9985 (Linking). doi: 10.1038/s41594-021-00656-9.

70. K. Singh, C. K. Lau, G. Manigrasso, J. B. Gama, R. Gassmann, and A. P. Carter. Molecular mechanism of dynein-dynactin complex assembly by lis1. *Science*, 383(6690):eadk8544, 2024. ISSN 1095-9203 (Electronic) 0036-8075 (Linking). doi: 10.1126/science.adk8544.

71. P. Chai, Q. Rao, Y. Wang, and K. Zhang. High-resolution structural analysis of dyneins by cryo-electron microscopy. *Methods Mol Biol*, 2623:257–279, 2023. ISSN 1940-6029 (Electronic) 1064-3745 (Print) 1064-3745 (Linking). doi: 10.1007/978-1-0716-2958-1_16.

72. P. Chai, Q. Rao, and K. Zhang. Multi-curve fitting and tubulin-lattice signal removal for structure determination of large microtubule-based motors. *J Struct Biol*, 214(4):107897, 2022. ISSN 1095-8657 (Electronic) 1047-8477 (Linking). doi: 10.1016/j.jsb.2022.107897.

73. M. Miura, A. Matsubara, T. Kobayashi, M. Edamatsu, and Y. Y. Toyoshima. Nucleotide-dependent behavior of single molecules of cytoplasmic dynein on microtubules in vitro. *FEBS Lett*, 584(11):2351–5, 2010. ISSN 1873-3468 (Electronic) 0014-5793 (Linking). doi: 10.1016/j.febslet.2010.04.016.

74. Y. Nishikawa, M. Inatomi, H. Iwasaki, and G. Kurisu. Structural change in the dynein stalk region associated with two different affinities for the microtubule. *J Mol Biol*, 428(9 Pt B):1886–96, 2016. ISSN 1089-8638 (Electronic) 0022-2836 (Linking). doi: 10.1016/j.jmb.2015.11.008.

75. L. Rao, F. Berger, M. P. Nicholas, and A. Gennerich. Molecular mechanism of cytoplasmic dynein tension sensing. *Nat Commun*, 10(1):3332, 2019. ISSN 2041-1723 (Electronic) 2041-1723 (Linking). doi: 10.1038/s41467-019-11231-8.

76. J. P. Gillies, J. M. Reimer, E. P. Karasmanis, I. Lahiri, Z. M. Htet, A. E. Leschziner, and S. L. Reck-Peterson. Structural basis for cytoplasmic dynein-1 regulation by lis1. *Elife*, 11, 2022. ISSN 2050-084X (Electronic) 2050-084X (Linking). doi: 10.7554/eLife.71229.

77. A. J. Roberts, N. Numata, M. L. Walker, Y. S. Kato, B. Malkova, T. Kon, R. Ohkura, F. Arisaka, P. J. Knight, K. Sutoh, and S. A. Burgess. Aaa+ ring and linker swing mechanism in the dynein motor. *Cell*, 136(3):485–95, 2009. ISSN 1097-4172 (Electronic) 0092-8674 (Print) 0092-8674 (Linking). doi: 10.1016/j.cell.2008.11.049.

78. C. M. Yengo, E. M. De la Cruz, D. Safer, E. M. Ostap, and H. L. Sweeney. Kinetic characterization of the weak binding states of myosin v. *Biochemistry*, 41(26):8508–17, 2002. ISSN 0006-2960 (Print) 0006-2960 (Linking). doi: 10.1021/bi015969u.

79. S. Uemura, K. Kawaguchi, J. Yajima, M. Edamatsu, Y. Y. Toyoshima, and S. Ishiwata. Kinesin-microtubule binding depends on both nucleotide state and loading direction. *Proceedings of the National Academy of Sciences of the United States of America*, 99(9):5977–5981, 2002. ISSN 0027-8424. doi: 10.1073/pnas.092546199.

80. P. I. Hanson and S. W. Whiteheart. Aaa+ proteins: Have engine, will work. *Nature Reviews Molecular Cell Biology*, 6(7):519–529, 2005. ISSN 1471-0072. doi: 10.1038/nrm1684.

81. E. L. F. Holzbaur and K. A. Johnson. Adp release is rate limiting in steady-state turnover by the dynein adenosine-triphosphatase. *Biochemistry*, 28(13):5577–5585, 1989. ISSN 0006-2960. doi: DOI10.1021/bi00439a036.

82. S. Can, S. Lacey, M. Gur, A. Carter, and A. Yildiz. Dynein's directionality is controlled by the angle and length of its stalk. *Biophysical Journal*, 116(3):309a–309a, 2019. ISSN 0006-3495. doi: DOI10.1016/j.bpj.2018.11.1676.

83. J. Lin, K. Okada, M. Raytchev, M. C. Smith, and D. Nicastro. Structural mechanism of the dynein power stroke. *Nat Cell Biol*, 16(5):479–85, 2014.

ISSN 1476-4679 (Electronic) 1465-7392 (Print) 1465-7392 (Linking). doi: 10.1038/ncb2939.

84. A. J. Roberts, T. Kon, P. J. Knight, K. Sutoh, and S. A. Burgess. Functions and mechanics of dynein motor proteins. *Nat Rev Mol Cell Biol*, 14(11):713–26, 2013. ISSN 1471-0080 (Electronic) 1471-0072 (Print) 1471-0072 (Linking). doi: 10.1038/nrm3667.

85. M. Dutta and B. Jana. Computational modeling of dynein motor proteins at work. *Chemical Communications*, 57(3):272–283, 2021. ISSN 1359-7345. doi: 10.1039/d0cc05857b.

86. R. Ibusuki, T. Morishita, A. Furuta, S. Nakayama, M. Yoshio, H. Kojima, K. Oiwa, and K. Furuta. Programmable molecular transport achieved by engineering protein motors to move on dna nanotubes. *Science*, 375(6585):1159–1164, 2022. ISSN 1095-9203 (Electronic) 0036-8075 (Linking). doi: 10.1126/science.abj5170.

87. S. R. Garrott, J. P. Gillies, and M. E. DeSantis. Nde1 and ndel1: Outstanding mysteries in dynein-mediated transport. *Front Cell Dev Biol*, 10:871935, 2022. ISSN 2296-634X (Print) 2296-634X (Electronic) 2296-634X (Linking). doi: 10.3389/fcell.2022.871935.

88. K. H. Ecklund, T. Morisaki, L. G. Lammers, M. G. Marzo, T. J. Stasevich, and S. M. Markus. She1 affects dynein through direct interactions with the microtubule and the dynein microtubule-binding domain. *Nat Commun*, 8 (1):2151, 2017. ISSN 2041-1723 (Electronic) 2041-1723 (Linking). doi: 10.1038/s41467-017-02004-2.

89. M. Mahamdeh, S. Simmert, A. Luchniak, E. Schaffer, and J. Howard. Label-free high-speed wide-field imaging of single microtubules using interference reflection microscopy. *J Microsc*, 272(1):60–66, 2018. ISSN 1365-2818 (Electronic) 0022-2720 (Print) 0022-2720 (Linking). doi: 10.1111/jmi.12744.

90. D. N. Mastronarde. Automated electron microscope tomography using robust prediction of specimen movements. *Journal of Structural Biology*, 152(1):36–51, 2005. ISSN 1047-8477. doi: 10.1016/j.jsb.2005.07.007.

91. A. Punjani, J. L. Rubinstein, D. J. Fleet, and M. A. Brubaker. cryosparc: algorithms for rapid unsupervised cryo-em structure determination. *Nat Methods*, 14(3):290–296, 2017. ISSN 1548-7105 (Electronic) 1548-7091 (Linking). doi: 10.1038/nmeth.4169.

92. S. Q. Zheng, E. Palovcak, J. P. Armache, K. A. Verba, Y. Cheng, and D. A. Agard. Motioncor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods*, 14(4):331–332, 2017. ISSN 1548-7105 (Electronic) 1548-7091 (Linking). doi: 10.1038/nmeth.4193.

93. K. Zhang. Gctf: Real-time ctf determination and correction. *J Struct Biol*, 193(1):1–12, 2016. ISSN 1095-8657 (Electronic) 1047-8477 (Linking). doi: 10.1016/j.jsb.2015.11.003.

94. T. D. Goddard, C. C. Huang, E. C. Meng, E. F. Pettersen, G. S. Couch, J. H. Morris, and T. E. Ferrin. Ucsf chimerax: Meeting modern challenges in visualization and analysis. *Protein Sci*, 27(1):14–25, 2018. ISSN 1469-896X (Electronic) 0961-8368 (Linking). doi: 10.1002/pro.3235.

95. R. T. Kidmose, J. Juhl, P. Nissen, T. Boesen, J. L. Karlsen, and B. P. Pedersen. Namdinator - automatic molecular dynamics flexible fitting of structural models into cryo-em and crystallography experimental maps. *IUCrJ*, 6(Pt 4):526–531, 2019. ISSN 2052-2525 (Print) 2052-2525 (Linking). doi: 10.1107/S2052252519007619.

96. A. Casanal, B. Lohkamp, and P. Emsley. Current developments in coot for macromolecular model building of electron cryo-microscopy and crystallographic data. *Protein Sci*, 29(4):1069–1078, 2020. ISSN 1469-896X (Electronic) 0961-8368 (Linking). doi: 10.1002/pro.3791.

97. A. Brown, F. Long, R. A. Nicholls, J. Toots, P. Emsley, and G. Murshudov. Tools for macromolecular model building and refinement into electron cryo-microscopy reconstructions. *Acta Crystallographica Section D-Structural Biology*, 71:136–153, 2015. ISSN 2059-7983. doi: 10.1107/S1399004714021683.

98. P. V. Afonine, B. K. Poon, R. J. Read, O. V. Sobolev, T. C. Terwilliger, A. Urzhumtsev, and P. D. Adams. Real-space refinement in phenix for cryo-em and crystallography. *Acta Crystallogr D Struct Biol*, 74(Pt 6):531–544, 2018. ISSN 2059-7983 (Electronic) 2059-7983 (Linking). doi: 10.1107/S2059798318006551.

99. V. B. Chen, W. B. Arendall, J. J. Headd, D. A. Keedy, R. M. Immormino, G. J. Kapral, L. W. Murray, J. S. Richardson, and D. C. Richardson. Molprobity: all-atom structure validation for macromolecular crystallography. *Acta Crystallographica Section D-Structural Biology*, 66:12–21, 2010. ISSN 2059-7983. doi: 10.1107/S0907444909042073.

Figure S1. Cryo-EM data processing of full-length human dynein-1 in the presence of 5 mM ATP. (A) A representative negative staining micrograph of purified full-length human dynein-1 and corresponding 2D class averages. **(B)** A typical cryo-EM micrograph and the flowchart of cryo-EM image processing. **(C)** Image processing of full-length phi dynein and tail region. **(D, E)** Orientation distribution of phi dynein and a representative state from open dynein. **(F, G)** The Fourier shell correlation (FSC) curves of the motor domain in different states and the tail domain. Datasets for dynein in other nucleotide conditions (ATP-Vi, AMPPNP, ADP, apo) were processed similarly.

Figure S2. Cryo-EM data processing of full-length human dynein-1 bound to MTs. (A) Typical cryo-EM micrographs of dynein bound to MTs in apo, ADP, and AMPPNP conditions. **(B)** Microtubule signal subtraction and image processing flowchart. **(C)** Orientation distribution of dynein-MT-ADP reconstruction. **(D)** FSC curves of four MT-bound motors. **(E)** Plot for the full-

length dynein bound to MTs in different conformations (two stable heads, one stable trailing head, and one stable leading head).

Figure S3. Local resolution analysis and identification of bound nucleotides in AAA1, AAA3, and AAA4 in different states from the dynein-ATP dataset. The color scheme for the motor domain is consistent with Figure 5.

Figure S4. The conformational landscapes of motor domains in different nucleotide conditions. (A) Cartoon schematic depicting the experimental method for cryo-EM analysis of full-length dynein in different nucleotide conditions. **(B-F)** Conformation landscapes of motor domains in ATP, ATP-Vi, AMPPNP, ADP, apo conditions. Arrows indicate likely mechanochemical pathways. Percentages indicate the proportion of particles in each state. The missing states in each condition are shown in grey.

Figure S5. Cryo-EM densities of the nucleotide and sensor-I loop in various closed AAA1 states. (A) From this work. **(B)** From previously published structures (PDB:8FD6, 7MGM).

Figure S6. Structural comparisons of MT-bound and MT-unbound dynein motors. The nucleotide states in AAA1, AAA3, and AAA4 and the linker docking mode are listed.

Figure S7. Systematic analysis of linker-ring interactions among four MT-bound motor domains. (A) Cartoon and molecular models showing the interaction interfaces including linker-AAA2L, linker-AAA5L, and linker-AAA3-4-connector helix. **(B-E)** Comparison of linker docking modes between dynein-1 and outer-arm dynein.

Table S1. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM ATP.

Table S2. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM ATP-Vi.

Table S3. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM AMPPNP.

Table S4. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM ADP.

Table S5. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in apo condition.

Table S6. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 bound to MTs.

Table S7. Cryo-EM data collection, refinement, and validation statistics of engineered human dynein motor domain locked in α registry.

Supplementary Video 1: High-resolution cryo-EM map of phi dynein and representative local density maps.

Supplementary Video 2: High-resolution cryo-EM reconstruction of dynein bound to MTs.

Supplementary Video 3: The release of Pi from dynein AAA1 through the backdoor mechanism, triggering the linker straightening.

Supplementary Video 4: Communication between AAA1 pocket dynamics and MTBD.

Supplementary Video 5: The forward-stepping mechanism of dynein along MTs.

Supplementary Table 1. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM ATP.

Description	Full-length dynein-1 in the presence of 5mM ATP											
	Composite map of full length dynein-1	State1 Phi-motor	State2	State3	State4	State5a	State5b	State6	State7a	State7b	State7c	State0
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing												
Facility	Yale West Campus-Cryo-EM facility											
Microscope	Titan Krios											
Voltage (kV)	300											
Camera	K3											
Magnification	105k											
Pixel Size (Å)	0.416 (super resolution)											
Total Electron Exposure (e-/A^2)	40											
Defocus Range (μm)	1.5-2.7											
Symmetry Imposed	C1	C1*	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1
Num of mics	33302											
Initial Particles	4 670 501											
Final Particles	57,816	872,840	165,772	105,800	122,798	65,824	114,327	260,931	67,951	66,008	66,325	612,628
Refinement												
Initial models	5NUG, 7K8F ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio
Map pixel size	1.664	0.416	0.832	0.832	0.832	0.832	0.832	0.832	0.832	0.832	0.832	0.832
Map Resolution (Å) (FSC 0.143)	3.3	2.25	2.85	3.02	2.82	3.49	3.48	3.48	3.22	3.58	3.22	2.93
Map sharpening B-factor (Å^2)	0	60	76.2	56.4	72.8	63.9	63.8	56.5	62.9	59.9	63.3	91.7
Model Composition												
Non-hydrogen atoms	89407	25271	23712	23061	23048	21869	21803	21782	23123	21829	23121	21370
Protein residues	11063	2899	2,937	2,859	2,859	2,711	2,701	2,698	2,859	2,704	2,859	2,642
Ligands	MG:4 ADP:6 ATP:2	MG:2 ADP:3 ATP:1 Water:1897	MG:2 ADP:3 ATP:1 Water:3	MG:2 ADP:3 ATP:1	MG:2 ADP:3 ATP:1 Water:7	ADP:3 ATP:1	MG:1 ADP:2 ATP:2	ADP:2 ATP:2	MG:1 ADP:2 ATP:2	ADP:2 ATP:2	ADP:2 ATP:2	MG:1 ADP:2 ATP:1
Model vs. Data												

FSC Map to Model (Å) (FSC 0.5)	3	2.4	2.8	3.1	2.8	3.8	3.8	4.0	3.5	4.0	3.5	3.2
Correlation coefficient (mask)	0.7	0.86	0.85	0.90	0.87	0.82	0.82	0.76	0.86	0.85	0.85	0.82
B factors (A²)												
Protein	100.96	25.33	39.98	50.59	33.75	89.22	101.82	85.68	53.63	114.60	44.50	14.29
Ligand	1.53	6.87	12.02	21.32	10.08	64.61	68.47	60.08	40.73	87.63	30.44	3.05
Water	/	21.56	11.75	/	11.54	/	/	/	/	/	/	/
R.m.s deviation												
Bond length (Å)	0.006	0.004	0.002	0.002	0.002	0.004	0.003	0.005	0.003	0.004	0.003	0.005
Bond angles (º)	1.083	0.662	0.491	0.489	0.51	0.646	0.635	0.864	0.568	0.656	0.583	0.65
Validation												
Molprobity score	1.94	1.51	1.4	1.38	1.39	1.64	1.7	1.91	1.63	1.80	1.62	1.58
Clashscore	13.53	9.59	7.3	6.87	7.05	12.42	12.9	21.04	9.3	13.52	9.43	11.35
Rotamer outliers (%)	0.24	0.04	0.84	0.00	0.04	0.17	0.54	0.13	0.31	0.54	0.00	0.09
Ramachandran plot												
Outliers (%)	0.39	0	0	0	0	0	0.04	0	0.11	0.07	0.04	0.00
Allowed (%)	3.91	1.49	1.19	1.61	1.3	2.15	2.31	2.42	2.64	2.79	2.60	2.05
Favored (%)	95.7	98.51	98.81	98.39	98.70	97.85	97.66	97.58	97.26	97.14	97.36	97.95
Rama-Z (whole)	0.55	1.4	1.69	2.17	2.24	1.17	1.44	0.99	1.36	1.71	1.71	1.39

*after C2 symmetry expansion

Supplementary Table 2. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM ATP-Vi.

Description	Full-length dynein-1 in the presence of 5mM ATP-Vanadate			
	State1 Phi-motor	State2	State4	State0
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing				
Facility	Yale ScienceHill-Cryo-EM facility			
Microscope	Glacios			
Voltage (kV)	200			
Camera	K3			
Magnification	45k			
Pixel Size (Å)	0.432 (super resolution)			
Total Electron Exposure (e-/A^2)	40			
Defocus Range (μm)	1.5-2.7			
Symmetry Imposed	C1*	C1	C1	C1
Num of mics	6016			
Initial Particles	2 554 074			
Final Particles	266 214	69 963	248 329	130 308
Refinement				
Initial models				
Map pixel size	1.157	1.157	1.157	1.157
Map Resolution (Å) (FSC 0.143)	2.78	3.04	2.81	3.3
Map sharpening B-factor (Å^2)	72.6	59.5	73.5	74.6
Model Composition				
Non-hydrogen atoms	23714	23714	23,046	21,370
Protein residues	2937	2937	2,859	2,642
Ligands	AOV:1 MG:2 ATP:1 ADP:2	AOV:1 MG:2 ATP:1 ADP:2	AOV:1 MG:2 ATP:1 ADP:2	MG:1 ATP:1 ADP:2
Model vs. Data				
FSC Map to Model (Å) (FSC 0.5)	3	3.3	3	3.7
Correlation coefficient (mask)	0.84	0.86	0.84	0.76

B factors (Å^2)				
Protein	61.21	74.39	57.9	70.58
Ligand	45.53	55.35	46.15	39.34
Water	/	/	/	/
R.m.s deviation				
Bond length (Å)	0.002	0.002	0.003	0.003
Bond angles (º)	0.462	0.482	0.52	0.595
Validation				
Molprobity score	1.26	1.33	1.36	1.56
Clashscore	4.93	5.95	6.55	10.9
Rotamer outliers (%)	0	0	0	0.13
Ramachandran plot				
Outliers (%)	0	0	0	0
Allowed (%)	1.37	1.43	1.51	1.9
Favored (%)	98.63	98.57	98.49	98.1
Rama-Z (whole)	1.79	1.54	1.95	1.73

Supplementary Table 3. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM AMPPNP.

Description	Full-length dynein-1 in the presence of 5mM AMPPNP						
	State1 Phi-motor	State2	State4	State5	State6	State7	State0
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing							
Facility	Yale ScienceHill-Cryo-EM facility						
Microscope	Glacios						
Voltage (kV)	200						
Camera	K3						
Magnification	45k						
Pixel Size (Å)	0.434 (super resolution)						
Total Electron Exposure (e-/A^2)	40						
Defocus Range (μm)	1.5-2.7						
Symmetry Imposed	C1*	C1	C1	C1	C1	C1	C1
Num of mics	7488						
Initial Particles	4 670 501						
Final Particles	130832	40850	92310	58149	82309	47831	97743
Refinement							
Initial models	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio
Map pixel size	1.389	1.302	1.302	1.302	1.302	1.302	1.302
Map Resolution (Å) (FSC 0.143)	3.1	3.3	3.14	3.78	3.63	3.51	3.5
Map sharpening B-factor (Å^2)	74.5	63	78.1	79.4	79.3	77.3	84.2
Model Composition							
Non-hydrogen atoms	23708	23708	23040	21869	21869	23121	21370
Protein residues	2937	2937	2859	2711	2711	2859	2642
Ligands	MG: 1 ADP: 3 ATP: 1	MG: 1 ADP: 3 ATP: 1	MG: 1 ADP: 3 ATP: 1	ADP: 3 ATP: 1	ADP: 3 ATP: 1	ANP: 1 ADP: 2 ATP: 1	MG: 1 ADP: 2 ATP: 1
Model vs. Data							
FSC Map to Model (Å) (FSC 0.5)	3.3	3.5	3.4	4.2	4	3.9	3.9
Correlation coefficient (mask)	0.87	0.83	0.87	0.76	0.8	0.82	0.73
B factors (A^2)							

Protein	46.38	96.38	63.53	89.22	91.02	92.78	102.64
Ligand	26.32	68.76	38.31	65.61	67.75	60.2	62.88
Water	/	/	/	/	/	/	/
R.m.s deviation							
Bond length (Å)	0.003	0.005	0.002	0.005	0.003	0.003	0.003
Bond angles (º)	0.51	0.646	0.489	0.984	0.674	0.657	0.63
Validation							
Molprobity score	1.43	1.42	1.29	1.64	1.71	1.71	1.62
Clashscore	7.8	7.76	5.45	12.45	14.64	11.16	11.98
Rotamer outliers (%)	0	0.35	0.16	0.25	0.46	0.39	0.09
Ramachandran plot							
Outliers (%)	0	0	0	0	0	0.04	0
Allowed (%)	1.64	1.33	1.44	2.15	2.15	2.78	2.13
Favored (%)	98.36	98.67	98.56	97.85	97.85	97.19	97.87
Rama-Z (whole)	2.13	2.2	1.62	1.17	1.29	1.15	1.7

Supplementary Table 4. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in the presence of 5mM ADP.

Description	Full-length dynein-1 in the presence of 5mM ADP						
	State1 Phi-motor	State2	State5	State6	State7-post1	State7-post2	State0
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing							
Facility	The Laboratory for BioMolecular Structure in BNL						
Microscope	Krios						
Voltage (kV)	300						
Camera	K3						
Magnification	105k						
Pixel Size (Å)	0.4125(super resolution)						
Total Electron Exposure (e-/A^2)	50						
Defocus Range (µm)	1.5-2.7						
Symmetry Imposed	C1*	C1	C1	C1	C1	C1	C1
Number of mics	5920						
Initial Particles	2203744						
Final Particles	174680	38237	40399	93339	47280	41042	79738
Refinement							
Initial models	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio
Map pixel size	1.1	1.03	1.03	1.03	1.03	1.03	1.03
Map Resolution (Å) (FSC 0.143)	2.6	3.04	3.7	3.7	3.7	3.3	3.3
Map sharpening B-factor (Å^2)	63.3	56.6	70.1	54.7	64.1	59.1	71.7
Model Composition							
Non-hydrogen atoms	23709	23709	21869	21778	24615	24615	21370
Protein residues	2937	2937	2711	2698	3043	3043	2642
Ligands	MG:1 ATP:1 ADP:3	MG:1 ATP:1 ADP:3	ATP:1 ADP:3	ATP:1 ADP:3	MG:1 ATP:1 ADP:3	MG:2 ATP:1 ADP:3	MG:1 ATP:1 ADP:2
Model vs. Data							
FSC Map to Model (Å) (FSC 0.5)	2.9	3.4	4.1	4.3	4	3.7	3.7
Correlation coefficient (mask)	0.84	0.8	0.8	0.76	0.81	0.8	0.72
B factors (A^2)							

Protein	11.21	22.27	122.5	133.8	131.06	96.57	14.29
Ligand	1.53	11.48	92.99	94.71	94.12	60.87	3.05
Water	/	/	/	/	/	/	/
R.m.s deviation							
Bond length (Å)	0.005	0.005	0.003	0.004	0.003	0.003	0.004
Bond angles (º)	0.978	0.971	0.619	0.727	0.611	0.628	0.925
Validation							
Molprobity score	1.36	1.39	1.62	2.02	1.76	1.64	1.62
Clashscore	6.54	7.13	12.97	20.2	13.43	11.15	11.91
Rotamer outliers (%)	0	0.12	0.04	0.00	0	0	0.21
Ramachandran plot							
Outliers (%)	0.07	0	0	0	0.03	0	0
Allowed (%)	1.54	1.71	1.96	3.43	2.57	2.34	2.13
Favored (%)	98.4	98.29	98.04	96.57	97.40	97.66	97.87
Rama-Z (whole)	1.58	1.91	1.57	0.53	1.7	1.73	1.34

Supplementary Table 5. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 in apo condition.

Description	Full-length dynein-1 in apo buffer				
	State1 Phi-motor	State2	State7	State8	State0
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing					
Facility	Yale ScienceHill-Cryo-EM facility				
Microscope	Glacios				
Voltage (kV)	200				
Camera	K3				
Magnification	45k				
Pixel Size (Å)	0.434 (superresolution)				
Total Electron Exposure (e-/A^2)	43				
Defocus Range (μm)	1.5-2.7				
Symmetry Imposed	C1*	C1	C1	C1	C1
Number of mics	5884				
Initial Particles	3549872				
Final Particles	136324	65852	67545	50945	235803
Refinement					
Initial models	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio	5NUG ab initio
Map pixel size	1.157	1.302	1.302	1.302	1.302
Map Resolution (Å) (FSC 0.143)	2.86	2.97	3.6	3.8	3.2
Map sharpening B-factor (Å^2)	73.6	58.8	76.7	71.4	89.8
Model Composition					
Non-hydrogen atoms	23709	23709	23495	24474	21370
Protein residues	2937	2937	2909	3029	2642
Ligands	MG:2 ATP:1 ADP:3	MG:2 ATP:1 ADP:3	MG:2 ATP:1 ADP:3	MG:2 ATP:1 ADP:2	MG:1 ATP:1 ADP:2
Model vs. Data					
FSC Map to Model (Å) (FSC 0.5)	3.2	3.2	4.4	4.2	3.5
Correlation coefficient (mask)	0.84	0.83	0.65	0.83	0.77
B factors (A^2)					

Protein	11.21	22.27	37.72	141.21	14.29
Ligand	1.53	11.48	29.57	94.14	3.05
Water	/	/	/	/	/
R.m.s deviation					
Bond length (Å)	0.005	0.005	0.006	0.003	0.004
Bond angles (º)	0.977	0.971	1.013	0.745	0.925
Validation					
Molprobity score	1.37	1.4	1.66	1.8	1.61
Clashscore	6.64	7.21	11.55	16.73	11.82
Rotamer outliers (%)	0	0	0.43	0.41	0.17
Ramachandran plot					
Outliers (%)	0.07	0	0.1	0	0
Allowed (%)	1.54	1.71	2.28	2.35	2.09
Favored (%)	98.4	98.29	97.62	97.65	97.91
Rama-Z (whole)	1.58	1.91	1.57	1.39	1.34

Supplementary Table 6. Cryo-EM data collection, refinement, and validation statistics of full-length dynein-1 bound to MTs.

Description	Motor domain from full-length dynein-1 bound to microtubules			
	Dynein-MT-Apo, post-1	Dynein-MT-ADP, post-1	Dynein-MT-ADP, post-2	Dynein-MT-AMPPNP, post-1
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing				
Facility	Laboratory for BioMolecular Structure in BNL	Yale ScienceHill-Cryo-EM facility		Laboratory for BioMolecular Structure in BNL
Microscope	Titan Krios	Glacios	Glacios	Titan Krios
Voltage (kV)	300	200	200	300
Camera	K3	K2	K2	K3
Magnification	105k	45k	45k	105k
Pixel Size (Å)	0.4125 (super resolution)	0.434 (super resolution)	0.434 (super resolution)	0.4125 (super resolution)
Total Electron Exposure (e-/A^2)	40	40	40	40
Defocus Range (μm)	1.5-2.7	1.5-2.7	1.5-2.7	1.5-2.7
Symmetry Imposed	C1	C1	C1	C1
Number of mics	9 935	11 828	11 828	5 690
Initial Particles	1 387 854	3 573 833	3 573 833	1 291 794
Final Particles	136 496	139 709	127 909	241 362
Refinement				
Initial models	ab initio	ab initio	ab initio	7z8g ab initio
Map pixel size	0.825	0.868	0.868	0.825
Map Resolution (Å) (FSC 0.143)	3.18	3.6	3.5	2.88
Map sharpening B-factor (Å^2)	81.6	93.5	79	85.2
Model Composition				
Non-hydrogen atoms	24474	24615	24615	24601
Protein residues	3029	3043	3043	3038
Ligands	MG:1 ATP:1 ADP:2	MG:2 ATP:1 ADP:3	MG:2 ATP:1 ADP:3	MG:4 AMPPNP2 ATP:1
Model vs. Data				
FSC Map to Model (Å) (FSC 0.5)	3.4	3.9	3.8	3.2
Correlation coefficient (mask)	0.85	0.77	0.82	0.81

B factors (Å²)				
Protein	35.66	113.23	30.14	5.94
Ligand	27.47	93.48	7.26	2.41
Water	/	/	/	/
R.m.s deviation				
Bond length (Å)	0.005	0.003	0.004	0.003
Bond angles (º)	0.987	0.58	0.572	0.532
Validation				
Molprobity score	1.59	1.7	1.6	1.48
Clashscore	10.27	10.95	9.43	9.07
Rotamer outliers (%)	0.15	0	0.04	0.04
Ramachandran plot				
Outliers (%)	0.03	0	0.03	0.07
Allowed (%)	2.22	2.77	2.44	1.45
Favored (%)	97.75	97.23	97.53	98.48
Rama-Z (whole)	1.7	1.46	1.55	1.87

Supplementary Table 7. Cryo-EM data collection, refinement, and validation statistics of engineered human dynein motor domain locked in α registry.

Description	Engineered dynein motor domain locked in α registry			
	in ATP-Vi condition		in ATP condition	
	Class1 ADP-AAA1, ADP-AAA3	Class2 ADP-AAA1, ATP-AAA3	Class1 ADP-AAA1, ADP-AAA3	Class2 ADP-AAA1, ATP-AAA3
	EMD-XXXX	EMD-XXXX	EMD-XXXX	EMD-XXXX
	PDB-XXXX	PDB-XXXX	PDB-XXXX	PDB-XXXX
Data Collection and Processing				
Facility	Yale ScienceHill-Cryo-EM facility		Laboratory for BioMolecular Structure in BNL	
Microscope	Glacios		Titan Krios	
Voltage (kV)	200		300	
Camera	K2		K3	
Magnification	36k		105k	
Pixel Size (Å)	0.57 (super resolution)		0.4125 (super resolution)	
Total Electron Exposure (e-/A^2)	40		40	
Defocus Range (μm)	1.5-2.7		1.5-2.7	
Symmetry Imposed	C1		C1	
Number of mics	2466		9 498	
Initial Particles	638 774		2 954 198	
Final Particles	112 474	72 335	433 112	91 718
Refinement				
Initial models	ab initio	ab initio	ab initio	ab initio
Map pixel size	1.143	1.143	1.146	1.146
Map Resolution (Å) (FSC 0.143)	3.3	3.4	2.8	2.9
Map sharpening B-factor (Å^2)	61.4	53.6	79	64.4
Model Composition				
Non-hydrogen atoms	23158	23084	23158	23084
Protein residues	2866	2855	2866	2855
Ligands	ATP:1 ADP:3	ATP:2 ADP:2	ATP:1 ADP:3	ATP:2 ADP:2
Model vs. Data				
FSC Map to Model (Å) (FSC 0.5)	3.7	3.9	3.2	3.3
Correlation coefficient (mask)	0.8	0.8	0.75	0.79
B factors (A^2)				
Protein	82.16	119.44	19.63	60.63

Ligand	56.75	91.8	5.58	33.32
Water	/	/	/	/
R.m.s deviation				
Bond length (Å)	0.002	0.003	0.004	0.004
Bond angles (º)	0.584	0.607	0.672	0.709
Validation				
Molprobity score	1.69	1.72	1.8	1.71
Clashscore	8.89	11.09	11.03	11.5
Rotamer outliers (%)	0.04	0	0.04	0.04
Ramachandran plot				
Outliers (%)	0.04	0.14	0.04	0.14
Allowed (%)	3.29	2.75	3.57	2.57
Favored (%)	96.67	97.11	96.39	97.29
Rama-Z (whole)	0.82	1.06	0.75	0.81