# Genotypic and phenotypic consequences of domestication in dogs

**Authors:** Sweetalana[1], Jazlyn A Mooney[2,*] and Zachary A Szpiech[1,3,*]

[1]Department of Biology, Pennsylvania State University, University Park, PA, USA

[2]Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA, USA

[3]Institute for Computational and Data Sciences, Pennsylvania State University, USA

Email for all authors: sweetalana2000@gmail.com, jazlyn.mooney@gmail.com, szpiech@psu.edu

*Indicates co-correspondence

Mailing Addresses with zip codes for all authors:

[1]514A Wartik Laboratory, University Park, PA, USA 16802

[2]1050 Childs Way, Ray R. Irani Hall, Los Angeles, CA, USA, 90089

ORCID:
Sweetalana: 0009-0000-5450-5934
Zachary A Szpiech: 0000-0001-6372-8224
Jazlyn A Mooney: 0000-0002-2369-0855

Short running title: Domestication shapes dog genetics and traits

Keywords: runs of homozygosity, dogs, inbreeding, GWAS, complex traits

## Abstract

Runs of homozygosity (ROH) are genomic regions that arise when two copies of an identical ancestral haplotype are inherited from parents with a recent common ancestor. In this study, we performed a novel comprehensive analysis to infer genetic diversity among dogs and quantified the association between ROH and non-disease phenotypes. We found distinct patterns of genetic diversity across clades of breed dogs and elevated levels of long ROH, compared to non-domesticated dogs. These high levels of $F_{ROH}$ (inbreeding coefficient) are a consequence of recent inbreeding among domesticated dogs during breed establishment. We identified statistically significant associations between $F_{ROH}$ and height, weight, lifespan, muscled, white head, white chest, furnish, and length of fur. After correcting for population structure, we identified more than 45 genes across the three examined quantitative traits that exceeded the threshold for suggestive significance, indicating significant polygenic inheritance for the complex quantitative phenotypes in dogs.

## Introduction

Dogs are one of the most common domesticated animals and evolved ~15,000 years ago from wolves (Vilà et al. 1997; Horard-Herbin, Tresset, and Vigne 2014; A. H. Freedman et al. 2014). Over time, humans performed tightly controlled selective breeding that resulted in various phenotypic traits (Vaysse et al. 2011). This intense selection led to the emergence of hundreds of new breeds and dramatic loss of genetic variation (Mastrangelo et al. 2018; Cecchi et al. 2013; Yordy et al. 2019). Due to breed establishment with a small number of founders and strong selection for morphology characteristics, high levels of inbreeding are observed across most domestic dog breeds (Bannasch et al. 2021). These high levels of inbreeding result in long

stretches of identical-by-descent (IBD) haplotypes in individual genomes, also known as runs of

homozygosity (ROH). Additionally, the process by which breeds were established resulted in a

unique genomic architecture, making testing phenotypic associations with multiple breeds

plausible (Plassais et al. 2019; Boyko 2011; Sams and Boyko 2019).

In humans, ROH are associated with numerous complex disease and non-disease

phenotypes (Clark et al. 2019; Johnson, Evans, and Keller 2018; Moreno-Grau et al. 2021; Keller

et al. 2012; Bacolod et al. 2008), and in dogs ROH have been associated with disease phenotypes

(Mooney, Yohannes, and Lohmueller 2021; Sams and Boyko 2019; Boyko 2011). Previous

studies have shown the influence of genetic variants in different quantitative traits such as leg

length, wrinkled skin, coat color, hair length, and skeletal shape (Sutter et al. 2007; Boyko et al.

2010; Parker et al. 2009; Cadieu et al. 2009; Candille et al. 2007). However, associations with

non-disease complex phenotypes in dogs are relatively unexplored (Hayward et al. 2016). To

determine the relationship between ROH and non-disease phenotypes in dogs, we analyzed 556

canid whole-genome sequences and characterized the genomic distribution of ROH. After

accounting for breed structure, we test for associations between the total ROH coverage of the

genome (as measured by $F_{ROH}$) among 13 breed groups with 13 phenotypes. We observe a

significant association with height, weight, lifespan, muscled, White chest, White head and

furnish. Finally, we perform an ROH-mapping genome-wide association study and identify

multiple genetic variants associated with three quantitative traits, namely height, weight, and

lifespan. Understanding the distribution of ROH and its association with non-disease phenotypes

provides insights on complex phenotypic architectures. Further, our work could help breeders

and veterinarians to evaluate biological mechanisms underlying complex phenotype variation in

dogs.

## Methods

### **Data filtering and categorizations**

In this study, we utilized 722 canid whole genomes from (Plassais et al. 2019) (NCBI accession number: PRJNA448733) that were filtered using BCFtools (Danecek et al. 2021). We retained only biallelic SNPs and sites of genotype quality >= 20 along with missing genotype rate of less than 10%. We filtered to remove variants from sex chromosomes, which resulted in 4,053,761 biallelic SNPs for all 38 pairs of autosomal chromosomes. Next, we categorized all individuals based on their Breed/Common name (Plassais et al. 2019) using both the neighbor joining trees from (vonHoldt et al. 2010) and the American Kennel Club (AKC) classifications. Lastly, we removed all wild canids, unknown breed status individuals, mixed breed individuals, and individuals for which breed information could not be found. This resulted in 13 breed groups namely, Ancient Spitz (31 individuals and 11 breeds), Herding (76 individuals and 15 breeds), Mastiff-like (65 individuals and 15 breeds), Retrievers (50 individuals and 6 breeds), Scent Hound (23 individuals and 10 breeds), Small Terrier (102 individuals and 8 breeds), Terriers (16 individuals and 7 breeds), Spaniels (22 individuals and 8 breeds), Toy Dogs (18 individuals and 7 breeds), Working Dogs ( 44 individuals and 10 breeds), Sight Hound (21 individuals and 9 breeds), Village Dogs (75 individuals and 13 regions), and Chinese Indigenous dogs (15 individuals). Overall, our working dataset included 558 individuals and 106 different breeds (Table S1).

## Calling Runs of Homozygosity

To call runs of homozygosity (ROH) we used GARLIC v1.6.0a, a likelihood-based ROH inference method (Szpiech, Blant, and Pemberton 2017), which uses a logarithm of odds (LOD) score measure of autozygosity applied in a sliding window for the entire genome (Pemberton et al. 2012; Szpiech, Blant, and Pemberton 2017). GARLIC requires input data with genotype and sample information in the form of TPED and TFAM files, which were generated using PLINK 2.0 (C. C. Chang et al. 2015). Moreover, we generated a TGLS file to obtain per-genotype likelihoods. We employed genotype likelihood data in the form of GQ to account for phred-scaled probability being incorrect. Next, we allowed GARLIC to choose the best window size based on SNP density, which starts at the window size of 100 SNPs and increases by the step size of 10 SNPs. GARLIC provides in-built ROH length classification for which we chose 6 classes: <1 Mbp, 1 Mbp -2 Mbp, 2 Mbp - 3 Mbp, 3 Mbp - 4 Mbp, 4 Mbp - 5 Mbp and >5 Mbp. To account for sample size differences in allele frequency estimation, we set the number of resamples to 40. All other flags were set to default, and the following parameters were used: --auto-winsize --auto-overlap-frac --winsize 100 --centromere centromeres_file.txt --size-bounds 1000000 2000000 3000000 4000000 5000000 --tped input_file.tped --tfam input_file.tfam --tgls input_file.tgls –gl-type GQ --resample 40 --out – output_file. To eliminate segments that were noisy, very short, and common, we filtered to retain ROH segments longer than 0.5 Mbp.

We performed independent ROH calling on breeds with at least 10 individuals. Breeds with fewer than 10 individuals were combined in their respective breed groups (Table S2). Thereafter, the ROH calling of all breeds irrespective of the individual numbers were combined into their breed groups for further analysis. For each individual, we computed the associations

between total number of runs of homozygosity ($n_{ROH}$) and total length of runs of homozygosity

($s_{ROH}$). For each breed group, we computed the arithmetic mean and range of both total $n_{ROH}$ and

total $s_{ROH}$. For each individual and breed group, we computed the arithmetic mean of both nROH

and sROH for each ROH size classification separately. The longest total sROH ( 2141.22 Mbp)

and the highest total nROH (1039) were identified in two individuals - PER00747 and

PER00393, respectively - from the small terrier Yorkshire breed. PER00747 also had a  depth of

the sequencing coverage across the entire genome of approximately 2.03, and produced inflated

values of sROH. Thus, we dropped PER00747 and PER00393 given coverage was likely not

sufficient for yielding reliable results.

**Inbreeding coefficient**

For each sample and breed group, we computed a genomic inbreeding coefficient, $F_{ROH}$. $F_{ROH}$ is

defined as the fraction of autosomal genome in ROH regions (McQuillan et al. 2009):

124
$$F_{ROH} = \frac{s_{ROH}}{L_{Auto}}$$

Where $s_{ROH}$ is the sum of length of runs of homozygosity in an individual's genome and $L_{Auto}$ is

the length of autosomal genome in base pairs. Autosomal genome length of 2,203,764,842 base

pairs was used since variants were called with the CanFam3.1 (NCBI RefSeq assembly :

GCF_000002285.3) reference genome.

**ROH Sharing Matrix**

We generated an ROH sharing matrix, a square matrix that represents the sum of overlaps of

runs of homozygosity between two individuals, to account for shared ROH patterns among

breeds. For this, we first generated individual ROH bed files for each individual. BEDTools

(Quinlan and Hall 2010) was used to identify those overlapping regions between two sets of

ROH data. The following parameters were used: bedtools intersect -wao -a

Firstindividual_ROH_file.bed -b Secondindividual_ROH_file.bed.

**Phenotype data**

We updated the phenotypic data (Plassais et al. 2019) for 13 phenotypes (bulky, Drop ears,

furnish, hairless, height, large ears, length of fur, lifespan, Long legs, muscled, weight, White

chest, and White head). For height, weight and lifespan, we used the average values obtained

from the AKC ("American Kennel Club," n.d.) (Table S3). When the sex-specific information

was available, we incorporated that into the dataset. When the information was absent, we used

the same values for both males and females. We normalized all three continuous traits using min-

max scaling (Table S4) . The phenotypic dataset for height, weight, and lifespan was further

segmented into small and large categories based on the average weight (21.66kg) of all

individuals (Table S5, Table S6). Additionally, we averaged the values for the remaining 10

binary traits based on the provided phenotypic data within their respective breeds (Table S7)

Here, the binary trait observed in an individual was encoded as 1, whereas traits not observed in

an individual was encoded as 0. No phenotypic information was calculated for the Chinese

indigenous dogs and Village dogs.

**Association tests**

We computed the association between $F_{ROH}$ and 13 phenotypes. We used GMMAT, an R

package that performs association tests using GLMM, generalized linear mixed models (Chen et

al. 2016). To fit the GLMM, glmmkin was used and we examined continuous traits (height,

weight, and lifespan) as the quantitative phenotype traits, $F_{ROH}$ as the covariate and an ROH

sharing matrix as kins. We fit the model assuming a Gaussian distribution of the continuous

phenotypes and identity as the link function. For associations of height, weight, and lifespan with

$F_{ROH}$ we used the updated phenotypic values for all 466 individuals. We also used weight and

weight*$F_{ROH}$ as the covariates across all three segments of the data (All individuals, small

individuals, and large individuals) (Table S8). To fit the GLMM for the other 10 binary traits, we

used binomial distribution as the family and logit as the link function. Individuals with "NA"

binary phenotypic information were removed from analysis. Due to lack of phenotype

information, Chinese Indigenous dogs and Village dogs were excluded from the association tests.

**ROH-mapping GWAS**

To understand the biological mechanisms of how ROH-associated SNPs influence the

phenotypic traits, we performed Genome-Wide Association Studies (GWAS). We generated a

dataset that examined the occurrence of SNPs in ROH regions using BCFTools intersect and

BCFTools subtract (Danecek et al. 2021). Here, SNPs in the ROH regions were encoded as 1,

whereas SNPs outside ROH regions were encoded as 0. We removed sites that were not in the

ROH regions (coded as 0) for all individuals. Next, we fit separate GLMM models using

GMMAT for all three phenotypic traits (height, weight, and lifespan) where ROH associated

SNPs were used as the covariate, ROH sharing matrix as the kins, Gaussian distribution as the

family and identity as the link function (Chen et al. 2016). We also fit GLMM models for each

segment of the data (All individuals, small individuals, and large individuals) using weight and

weight*interaction as the covariates (Table S9). We used the option effectiveSize within the coda

R package (Plummer et al. 2006) to obtain the effective sample size adjusted for autocorrelation

between sites. To obtain the genome-wide significance and suggestive significance thresholds,

we used the Bonferroni correction with respect to effective sample size (Li et al. 2011; Yuan et

al. 2020; W. Yang et al. 2014):

$$Genome-Wide\ Significance\ Threshold = -log10(\frac{0.05}{Effective\ Sample\ Size})$$

$$Suggestive-Wide\ Significance\ Threshold = -log10(\frac{0.1}{Effective\ Sample\ Size})$$

To correct for population stratification, we calculated the genomic inflation factor, $\lambda$ (van den

Berg et al. 2019; Sofer et al. 2021). We corrected the log-scaled p-values by dividing them by the

genomic inflation factor (Table S9).  To visualize the GWAS summary statistics, we used an R

package, qqman to create Q-Q and Manhattan plots (Turner 2018).

## Results

### ROH distribution

We used GARLIC to assign each individual's ROH into our pre-specified ROH length classes.

This allowed us to explore the variance per-population within each class. Briefly, ROH were

categorized into 6 length classes: Class A (0.5 Mbp - 1 Mbp), Class B ( 1 Mbp - 2 Mbp), Class C

( 2 Mbp - 3 Mbp), Class D ( 3 Mbp - 4 Mbp), Class E (4 Mbp - 5 Mbp), and Class F ( > 5Mbp).

The mean $n_{ROH}$ and mean $s_{ROH}$ for 556 individuals was 37.06 and 94.17 Mbp, respectively. We

also reported the mean $n_{ROH}$ and mean $s_{ROH}$ for each ROH length classification: Class A (99.44

and 68.39 Mbp), Class B (50.29 and 70.36 Mbp), Class C (21.60 and 52.81 Mbp), Class D

(13.07 and 45.23 Mbp), Class E ( 9.21 and 41.15 Mbp), and Class F (28.77 and 287.05 Mbp)

(Table 1 and Table 2). Class A (0.5 Mbp - 1 Mbp) are the smallest ROH, and Class F are the

largest ROH ( > 5Mbp).

| Breed Group | Mean $n_{ROH}$ | Range $n_{ROH}$ | Mean $n_{ROH}$ for Class A (0.5-1 Mbp) | Mean $n_{ROH}$ for Class B (1 - 2 Mbp) | Mean $n_{ROH}$ for Class C (2 - 3 Mbp) | Mean $n_{ROH}$ for Class D (3 - 4 Mbp) | Mean $n_{ROH}$ for Class E (4 - 5 Mbp) | Mean $n_{ROH}$ for Class F (> 5 Mbp) |
|---|---|---|---|---|---|---|---|---|
| Ancient Spitz | 47.10 ± 58.20 | 3 - 338 | 133.42 ± 85.63 | 65.13 ± 40.00 | 24.71 ± 11.23 | 13.90 ± 4.31 | 9.87 ± 3.97 | 35.58 ± 18.29 |
| Retrievers | 38.83 ± 34.58 | 0 - 138 | 106.42 ± 13.75 | 53.96 ± 9.05 | 23.98 ± 6.39 | 13.1 ± 4.31 | 9.94 ± 3.32 | 26.3 ± 11.77 |
| Mastiff Like | 38.37 ± 26.65 | 0 - 117 | 78.6 ± 14.12 | 54 ± 17.53 | 27.46 ± 11.19 | 16.91 ± 7.55 | 12.18 ± 6.53 | 41.08 ± 19.83 |
| Herding | 37.88 ± 33.97 | 0 - 257 | 97.76 ± 32.29 | 48.47 ± 10.17 | 21 ± 7.22 | 13.91 ± 5.79 | 10.8 ± 5.51 | 35.32 ± 19.08 |
| Scent Hound | 40.25 ± 37.52 | 1 - 160 | 113.26 ± 21.76 | 53.65 ± 9.85 | 22.7 ± 6.35 | 13.3 ± 4.87 | 9.09 ± 4.92 | 27.48 ± 12.44 |
| Small Terrier | 40.93 ± 41.19 | 0 - 277 | 113.58 ± 39.05 | 60.27 ± 17.74 | 24.08 ± 9.18 | 14.31 ± 6.32 | 8.98 ± 4.61 | 24.38 ± 16 |
| Terriers | 30.56 ± 33.01 | 9 - 122 | 90.75 ± 15.75 | 56.25 ± 11.22 | 31.81 ± 7.75 | 18.88 ± 6.21 | 15.88 ± 4.66 | 53.75 ± 10.77 |
| Spaniels | 36.29 ± 35.23 | 5 - 144 | 101.18 ± 20.19 | 61 ± 9.23 | 28.73 ± 5.37 | 18.14 ± 5.35 | 14 ± 5.19 | 41.64 ± 14.62 |
| Sight Hound | 45.29 ± 48.55 | 0 - 192 | 138.67 ± 31.05 | 63.33 ± 20.63 | 21.67 ± 8.98 | 13.86 ± 7.57 | 8.19 ± 4.76 | 26 ± 15.86 |
| Toy Dogs | 41.19 ± 40.77 | 1 - 187 | 114.56 ± 31.06 | 56.17 ± 25.99 | 24.06 ± 13.48 | 14.78 ± 9.78 | 9.44 ± 6.46 | 28.11 ± 15.34 |
| Working Dogs | 38.29 ± 27.43 | 1 - 129 | 85.18 ± 16.07 | 50.02 ± 15.37 | 27.36 ± 12.12 | 17 ± 7.86 | 11.82 ± 5.51 | 38.34 ± 13.11 |
| Village Dogs | 20.5 ± 31.83 | 0 - 261 | 80.12 ± 33.69 | 23.79 ± 13.69 | 6.33 ± 5.77 | 3.51 ± 4.76 | 2.09 ± 2.64 | 7.15 ± 11.03 |
| Chinese Indigenous Dogs | 14.36 ± 22.91 | 0 - 103 | 61.53 ± 16.63 | 14.67 ± 5.79 | 4 ± 3.31 | 1.27 ± 1.29 | 1 ± 1.32 | 3.67 ± 5.37 |

**TABLE 2.** *sROH across all breed groups and ROH length classification. The table represents the range and mean ± standard deviation of the sum of runs of homozygosity for each breed group and ROH length classification.*

| Breed Group | Mean $s_{ROH}$ (Mbp) | Range $s_{ROH}$ (Mbp) | Mean $s_{ROH}$ for Class A (0.5-1 Mbp) | Mean $s_{ROH}$ for Class B (1-2 Mbp) | Mean $s_{ROH}$ for Class C (2-3 Mbp) | Mean $s_{ROH}$ for Class D (3-4 Mbp) | Mean $s_{ROH}$ for Class E (4-5 Mbp) | Mean $s_{ROH}$ for Class F (> 5 Mbp) |
|---|---|---|---|---|---|---|---|---|
| Ancient Spitz | 119.56 ± 162.09 | 13.73 - 1113.00 | 92.29 ± 59.41 | 90.67 ± 54.30 | 60.33 ± 27.20 | 48.30 ± | 44.14 ± 17.51 | 381.60 ± 255.7 |
| Retrievers | 89.28 ± 86.63 | 0 - 598.02 | 72.96 ± 9.44 | 75.50 ± 13.25 | 57.37 ± 16.00 | 45.22 ± 15.07 | 44.47 ± 14.90 | 240.17 ± 125.98 |
| Mastiff Like | 126.47 ± 176.91 | 0 - 1029.28 | 54.77 ± 9.71 | 76.97 ± 25.95 | 67.21 ± 27.55 | 58.57 ± 25.92 | 54.38 ± 29.38 | 446.89 ± 247.25 |
| Herding | 105.25 ± 139.34 | 0 - 861.74 | 67.24 ± 21.71 | 67.53 ± 14.78 | 51.27 ± 17.64 | 48.23 ± 20.12 | 48.26 ± 24.73 | 348.94 ± 206.94 |
| Scent Hound | 99.19 ± 113.14 | 4.59 - 946.54 | 77.78 ± 14.73 | 74.84 ± 15.22 | 55.26 ± 15.67 | 45.90 ± 16.82 | 40.62 ± 21.78 | 300.75 ± 159.58 |
| Small Terrier | 89.41 ± 96.13 | 0 - 690.33 | 78.20 ± 26.38 | 84.24 ± 24.77 | 59.07 ± 22.76 | 49.63 ± 22.10 | 40.18 ± 20.51 | 225.15 ± 170.89 |
| Terriers | 75.86 ± 116.57 | 28.77 - 951.29 | 63.93 ± 10.90 | 79.69 ± 16.00 | 77.98 ± 19.72 | 65.84 ± 21.84 | 71.06 ± 20.79 | 541.71 ± 161.42 |
| Spaniels | 94.03 ± 127.54 | 23.23 - 856.91 | 70.13 ± 12.83 | 85.21 ± 12.79 | 70.39 ± 13.56 | 62.52 ± 18.65 | 62.55 ± 23.16 | 427.24 ± 190.19 |
| Sight Hound | 94.65 ± 96.81 | 0 - 591.60 | 95.24 ± 21.50 | 86.87 ± 29.21 | 53.04 ± 22.07 | 47.98 ± 26.13 | 36.54 ± 21.32 | 248.26 ± 149.49 |
| Toy Dogs | 97.01 ± 107.57 | 4.12 - 602.43 | 78.92 ± 22.17 | 79.01 ± 36.61 | 58.31 ± 33.13 | 50.45 ± 33.72 | 42.33 ± 28.82 | 273.02 ± 162.01 |
| Working Dogs | 112.89 ± 130.13 | 4.74 - 658.42 | 58.67 ± 11.14 | 70.67 ± 22.08 | 66.54 ± 29.19 | 58.83 ± 27.51 | 52.76 ± 24.70 | 369.85 ± 139.08 |
| Village Dogs | 33.24 ± 59.75 | 0 - 554.84 | 54.07 ± 23.09 | 32.65 ± 19.59 | 15.30 ± 13.97 | 12.04 ± 16.34 | 9.35 ± 11.72 | 76.00 ± 127.67 |
| Chinese Indigenous Dogs | 19.82 ± 35.06 | 0 - 309.27 | 41.03 ± 11.34 | 20.00 ± 7.64 | 9.53 ± 7.68 | 4.27 ± 4.24 | 4.41 ± 5.87 | 39.67 ± 75.16 |

Domestic dog clades showed a distinct pattern of genetic diversity, with higher levels of long ROH (>5 Mbp) when compared with Chinese Indigenous Dogs (light green) and Village dogs (royal blue) (Figure 1 and Figure 2). The highest total $n_{ROH}$ (641) and the highest total $s_{ROH}$ (1643.98 Mbp) were both observed in individuals (Dingo01 and NGSD2, respectively) from the Ancient Spitz breed group (Table S10). The maximum length of a single ROH (1113.00 Mbp) was observed in a New Guinea Singing Dog (Table 2 and Table S10). Terriers had the smallest mean $n_{ROH}$ and $s_{ROH}$ when compared with the remaining 11 breed groups (Table 1 and Table 2).

**FIGURE 1.** *Relationship between number of runs of homozygosity (nROH) and ROH size class in Mbp for all breed groups. The x-axis represents the ROH size in Mbp. The y-axis represents the number of ROH. Each violin plot depicts the distribution of the total number of ROH within length ranges 0.5-1 Mbp, 1-2 Mbp, 2-3 Mbp, 3-4 Mbp, 4-5 Mbp and >5 Mbp for each breed group.*

**FIGURE 2.** *Relationship between sum of runs of homozygosity (sROH) and ROH size class in Mbp. The x-axis represents the ROH size in Mbp. The y-axis represents the sum of runs of homozygosity. Each violin plot depicts the distribution of the sum of ROH within length ranges 0.5-1 Mbp, 1-2 Mbp, 2-3 Mbp, 3-4 Mbp, 4-5 Mbp and >5 Mbp for each breed group.*

To further quantify the relationship between domesticated and non-domesticated populations, we examined $s_{ROH}$ and $n_{ROH}$ (Figure 3). Larger values $s_{ROH}$ have been shown in

populations that have experienced a bottleneck, while larger $n_{ROH}$ were found in small

populations (Ceballos et al. 2018). We observed larger values of mean $n_{ROH}$ and $s_{ROH}$ for the

domesticated breed dogs, relative to non-domesticated Chinese Indigenous Dogs and Village

Dogs (Figure 3). Further, the longest ROH were also observed in domestic dog clades. This

result corresponds with the higher degree of relatedness among all domestic breed group

individuals, and was expected given the known high levels of inbreeding during breed

establishment. Our results highlight Village and Chinese Indigenous dogs as more outbred

populations given lower ROH proportion when compared to breed dogs (Figure 3). We also

observed high levels of $n_{ROH}$ and $s_{ROH}$ within the Ancient Spitz clades. Our results suggest recent

bottlenecks in the Ancient Spitz clade's history.

*FIGURE 3.* *Relationship between sum of runs of homozygosity (sROH) and number of runs of homozygosity (nROH) for each breed group. The x-axis and the y-axis represents the sROH and nROH, respectively. The black line is y=x.*

Next, we examined the level of inbreeding within each clade by computing an inbreeding coefficient, $F_{ROH}$ for each individual and for the breed group (Table 3 and Table S11). To summarize the distribution of $F_{ROH}$ across all breed groups, we computed mean $F_{ROH}$ (Table 3). High levels of mean $F_{ROH}$ were observed in clades composed of breed dogs. $F_{ROH}$ values ranged from a mean of approximately 0.41 in Terriers to a mean of approximately 0.24 in Retrievers.

The lowest values were observed in Chinese Indigenous Dogs (mean ~0.05) and Village Dogs

(mean ~0.09). This result provides additional support for breed dogs sharing a more recent

common ancestor and more background relatedness relative to either the Village or Chinese

Indigenous dogs. Overall, our results from evaluating differences across $n_{ROH}$ and $s_{ROH}$ were

expected given what is known about the domestication of dogs, current populations of village

dogs, and the recent history of breed dogs and (Boyko 2011).

**TABLE 3.** $F_{ROH}$ *across all breed groups. The table represents the mean ± standard deviation of*
$F_{ROH}$ *for each breed group*

| Breed Group | Mean $F_{ROH}$ |
|---|---|
| Terriers | 0.41 ± 0.07 |
| Spaniels | 0.35 ± 0.09 |
| Mastiff-Like | 0.34 ± 0.15 |
| Ancient Spitz | 0.33 ± 0.17 |
| Working Dog | 0.31 ± 0.09 |
| Herding | 0.29 ± 0.11 |
| Scent Hound | 0.27 ± 0.08 |
| Toy Dogs | 0.26 ± 0.13 |
| Sight Hound | 0.26 ± 0.10 |
| Small Terrier | 0.24 ± 0.10 |
| Retriever | 0.24 ± 0.07 |
| Village Dogs | 0.09 ± 0.08 |
| Chinese Indigenous Dogs | 0.05 ± 0.04 |

### Association between $F_{ROH}$ and phenotypic traits

There exists a gap in the literature with respect to exploring the relationship between ROH and quantitative non-disease phenotypes. Thus, we next explored the relationship between $F_{ROH}$ and all 13 phenotypes (bulky, Drop ears, furnish, hairless, height, large ears, length of fur, lifespan, long legs, muscled, weight, White chest, and White head). Given that the effect of breed structure is particularly strong in dogs we and we were associating a trait with FROH we utilized with a kinship matrix quantified by pairwise ROH sharing between individuals, following the approach in Mooney et al. 2021. We identified three phenotypes for all individuals with significant associations: height ($\beta$ = 0.31 and p = *4.05 $\times$ 10$^{-3}$*), weight ($\beta$ = 0.19, p = *1.19 $\times$ 10$^{-2}$*), and lifespan ($\beta$ = -0.14 and p = *4.44 $\times$ 10$^{-2}$*) (Figure 4). Across breed dogs, for every 1% increase in $F_{ROH}$, height increases by 0.192 cm, weight increases by 0.14 kg and lifespan decreases by 0.014 years. Statistical analysis including normalized coefficient Beta ($\beta$), along with its associated p-values and confidence intervals for different variations of models for quantitative traits (height, weight, lifespan) are shown in Table S8.

**FIGURE 4.** *Relationship between effect size and quantitative phenotypic traits. The y-axis represents the phenotypes for which the associations were tested. The x-axis represents the normalized Beta coefficients (β) for their respective traits. A significant effect of FROH on a phenotype (nominal p < 0.05) is indicated with a red point. An effect size of greater than 0 indicates an increase of that trait with respect to $F_{ROH}$, and less than 0 represents the converse.*

Next, we determined whether there was a relationship between $F_{ROH}$ and the binary

phenotypes. We observed significant associations between $F_{ROH}$ and 5 of the binary phenotypes:

muscled (β = 7.96 and p =$9.68 \times 10^{-11}$) , White chest (β = 3.49 and p = $5.36 \times 10^{-5}$), White

head (β = 4.26 and p = $1.55 \times 10^{-4}$), length of fur (β = -1.88 and p = $1.27 \times 10^{-2}$), and furnish

($\beta$ = -2.50 and p = $8.24 \times 10^{-3}$) (Figure 5 and Table S12). Across all breed groups, as $F_{ROH}$

increases, White Chest, White Head and muscled phenotypes also increase, whereas length of fur

and furnish phenotypes decreases.

***FIGURE 5.*** *Relationship between effect size and categorical phenotypic traits. The y-axis*
*represents the phenotypes for which the associations were tested. The x-axis represents the Beta*
*coefficients ($\beta$) for their respective traits. A significant effect of $F_{ROH}$ on a phenotype (nominal p*
*< 0.05) is indicated with a red point. An effect size of greater than 0 indicates an increase of that*
*trait with respect to $F_{ROH}$, and less than 0 represents the converse.*

## ROH-mapping GWAS

To identify ROH-associated genomic regions that influence life span, weight, or height, a GLMM-based GWAS was performed (Figure 6, Figure 7, Figure 8, and Figure S1-S9). Out of the 157,832 SNPs for height, 27 SNPs were observed to be above SWS (Suggestive Wide Significance) threshold and 18 SNPs were observed to be above GWS (Genome Wide Significance) threshold when tested for all individuals. A single SNP, chr24:22647289, was detected to have the strongest associated genomic signal with height (p = $3.46 \times 10^{-6}$) corresponding to SNTA1 (Figure 6A). Three genes, CERS3 (7 SNPs) on chromosome 3; CBFA2T2 (16 SNPs) on chromosome 24; and SNTA1 (4 SNPs) on chromosome 24, were observed above the SWS threshold, and contained multiple SNPs within their transcription regions (Table S13). CERS3 was previously associated with Body Mass Index (BMI) in human studies (Costa-Urrutia et al. 2019). Moreover, CERS3, SNTA1, and CBFA2T2 are known to harbor common genetic variants that are found to be associated with height in humans (Kichaev et al. 2019; Yengo et al. 2022).

For small individuals, 34 SNPs were observed to be above SWS threshold and 17 SNPs were observed to be above GWS threshold when tested for association with height. A single SNP, chr9:45711401 was detected to have the strongest associated genomic signal (p =$1.43 \times 10^{-6}$) corresponding to the gene PITPNA (Figure 6B). On chromosome 3, CERS3 (3 SNPs), IGF1R (1 SNP), LRRC28 (1 SNP), and TTC23 (3 SNPs) contained common genetic variants observed above SWS threshold. On chromosome 9, PITPNA (4 SNPs), SLC43A2 (2 SNPs), VPS53 (2 SNPs), and RPH3AL (4 SNPs) contained common genetic variants observed above the SWS threshold. On chromosome 18, HGF (2 SNPs) was the only gene that contained

common genetic variants above the SWS threshold. On chromosome 22, COMMD6 (1 SNP)

was the only gene that contained a common genetic variant observed above the SWS threshold.

On chromosome 24, CBFA2T2 (7 SNPs) and SNTA1 (4 SNPs) contained common genetic

variants observed above the SWS threshold. Many of the genes (PITPNA, CERS3, LRRC28,

TTC23, HGF, CBFA2T2, SNTA1, SLC43A2, and VPS53) that we identified as reaching SWS

were previously identified as genes associated with height in humans (Yengo et al. 2022).

RPH3AL has been found to be associated with IGF1R in a network analysis with human patients

(Virakul et al. 2020). IGF1R has been previously associated with height in small dog breeds

(Hoopes et al. 2012). To our knowledge, no previous studies have identified an association

between height and COMMD6.

For large individuals, a single SNP, chr11:9943557 encoding for PRR16 was observed to

be above SWS threshold when tested for association with height (Figure 6C). PRR16 has also

been previously associated with height in humans (Yengo et al. 2022). An extended table

containing all the genes that were observed to be associated with height when using weight and

weight*interaction as the covariates are indicated in Table S13.

**FIGURE 6.** *Manhattan plots for GLMM-based GWAS for body height. A. All individuals B.*
*Small Individuals C. Large Individuals. The x-axis represents the genomic position. The y-axis*
*represents the log10 base transformed p-values. Single nucleotide polymorphisms are*
*represented by a single point. Red line indicates the suggestive wide significance (SWS)*
*threshold while the blue line indicates genome wide significance (GWS) threshold. Only genes*
*above the GWS threshold are indicated with the exception of IGF1R. All marked genes are not*
*drawn to scale.*

For weight, 66 SNPs were observed above SWS threshold with 45 SNPs above GWS

threshold. The majority of SNPs (30) that were above the SWS threshold and found within genes

were observed on chromosome 24 (Table S14). A single SNP, chr32:11413601, (p

$=3.24 \times 10^{-7}$) with the strongest association with weight was observed within a gene that

encodes for Protein Kinase D2, PKD2 (Figure 7A). Ablation of PKD2 was shown to be

associated with reduced absorption of fats in the intestines of mice (Trujillo-Viera et al. 2021).

PKD2 has also been linked to increase the capacity of an organism to absorb fat from a high fat

diet, in other words, this gene potentially promotes obesity (Trujillo-Viera et al. 2021).

Other GWS genes when tested for all individuals with weight were SNTA1, CBFA2T2,

TXNRD3, SPIDR, ZXDC, ZNF341, CERS3, and CHMP4B (Table S14). SNTA1 was found to

be one of the closest associated genes with Live Body Weight (LBW) at 4 months in Karachai

goats (Selionova et al. 2022). CBFA2T2 is a key regulator of adipogenesis which affects obesity

when the gene expressions are altered (Matulewicz et al. 2017; Ali et al. 2013; Luo et al. 2020).

TXNRD3, thioredoxin reductase 3 effects adipocyte differentiation through Wnt signaling

pathway, a pathway that suppresses the adipogenic transcription factors (Martínez-Montes et al.

2016; Kipp et al. 2012; S. Kang et al. 2007). CERS3 was previously associated with Body Mass

Index (BMI) in human studies (Costa-Urrutia et al. 2019). CHMP4B has been associated with

human offspring birth weight (Warrington et al. 2019). Moreover, SPIDR was associated with

increased total body fat and ZNF341 is associated with bone mineral content which is directly

linked to body weight (Shapses and Sukumar 2012; Blake et al. 2021). ZNF341 has also been

previously associated with weight in humans (Tachmazidou et al. 2017). However, to our

knowledge, no previous studies have associated obesity/body mass with ZXDC and CHMP4B.

For small individuals, 31 SNPs were observed above SWS threshold and 19 SNPS above

GWS threshold. A single SNP, chr24:22660667, (p = $5.88 \times 10^{-7}$) with the strongest

association with weight was observed within a gene that encodes for SNTA1 (Figure 7B).

ZNF341, PITPNA, CBFA2T2, and SLC43A2 were also observed above the GWS threshold.

Ablation of PITPNA is associated with decreased body weight in mice (Alb et al. 2003); and

ablation of SLC43A2 has been associated with 10% weight reduction in fetal mice (Alb et al.

2003; Guetg et al. 2015). In contrast, there were only two encoded genes in large individuals that

reached significance. PKD2 (chromosomes 32) and SPIDR (chromosome 35) were both

observed to be associated with weight above the SWS threshold (Figure 7C) .

**FIGURE 7.** *Manhattan plots for GLMM-based GWAS for body weight. A. All individuals B. Small Individuals C. Large Individuals. The x-axis represents the genomic position. The y-axis represents the log10 base transformed p-values. Single nucleotide polymorphisms are represented by a single point. Red line indicates the suggestive wide significance (SWS) threshold while the blue line indicates genome wide significance (GWS) threshold. All marked genes are not drawn to scale.*

For lifespan, 10 SNPs were observed above the SWS threshold with only 5 SNPs above

the GWS threshold when tested across all individuals ( Figure 8A, Table S15). The strongest

association SNP, chr20:789204, (p = 2.21 × $10^{-5}$) encoding for TXNRD3 was observed on

chromosome 20. TXNRD3 has been previously associated with lower survival rates for various

types of cancers, such as kidney renal cell carcinoma and pancreatic adenocarcinoma (Wu et al.

2021). However, to our knowledge, no previous studies have found that link lifespan with

TXNRD3. The next most significant SNP, chr33:18399003 (p = 2.68 × $10^{-5}$) was observed in

the transcription region encoding for D3 dopamine receptor, DRD3. DRD3 is associated with

schizophrenia, resulting in loss of life expectancy (Y. Kang et al. 2023; Hjorthøj et al. 2017). We

observed another significant signal near, TIGIT, which has been associated with solid tumors

from lung, colon, breast, uterine and renal cancer, resulting in reduced longevity (Johnston et al.

2014; Chauvin et al. 2015). We also observed a signal above the SWS threshold on chromosome

32 in SEPT11, this gene was previously identified as a tumor-promoting gene with a predicted

poor prognosis (Fu et al. 2023). Likewise, ZXDC has been previously associated with enhanced

cervical cancer metastasis which has a significant negative impact on patient survival (Mao et al.

2023).

For small individuals, RUNX2 (4 SNPs), on chromosome 12, was the single gene that

was  observed as a standout above the SWS threshold. (Figure 8B). RUNX2 has been previously

associated with multiple cancers such as breast cancer, prostate cancer, gastric cancer etc.,

resulting in poor prognosis (Zhao et al. 2021). For large individuals, TXNRD3, ZXDC,

CFAP100, and SLC41A3 were the observed genes above SWS threshold (Figure 8C). SLC41A3

has been previously associated with a probable indication for liver hepatocellular carcinoma

which is the second primary contributor of carcinoma-associated death (Zhao et al. 2021; Q.

Chang et al. 2021). To our knowledge, no previous studies have linked lifespan with CFAP100.

All associated genes that were observed with weight and weight*interaction covariates are listed

in Table S15.

Overall, our GWAS suggests that quantitative traits in dogs are polygenic with multiple

variants being associated with a trait. Our results also support previous research that has

demonstrated a much simpler genetic architecture of complex traits in breed dogs relative to

humans (Boyko et al. 2010). In the future, these genes could be used as candidates for further

functional exploration.

**FIGURE 8.** *Manhattan plots for GLMM-based GWAS for lifespan. A. All individuals B. Small Individuals C. Large Individuals. The x-axis represents the genomic position. The y-axis represents the log10 base transformed p-values. Single nucleotide polymorphisms are represented by a single point. Red line indicates the suggestive wide significance (SWS) threshold while the blue line indicates genome wide significance (GWS) threshold. All marked genes are not drawn to scale.*

## Discussion

In sum, our work provides an in-depth analysis of ROH patterns in domesticated and non-

domesticated dogs. All breed dogs, which have experienced domestication, showed distinct

patterns of genetic diversity in comparison to Village dogs and Chinese Indigenous dogs. We

observe a larger fraction of the genome in long ROH (>5 Mbp) in breed dogs (Table 2), which

corresponds to small population sizes <10 generations ago (Purfield et al. 2017; Thompson

2013). This increase in the amount of the genome within a long ROH also corresponds with

higher values of $F_{ROH}$. Breed dogs also share most of their common genetic variation and lack

unique genetic diversity (Boyko et al. 2010). Lower genetic diversity and larger $F_{ROH}$ values

provide evidence for elevated levels of inbreeding and background relatedness. These results are

in line with findings from (Bannasch et al. 2021), and are indicative of the strong bottlenecks

experienced during breed formation.

Conversely, the non-domesticated Village dogs and Chinese Indigenous dogs have the

lowest values for mean ROH distribution. The Village dogs were sampled from 13 different

regions across the world and have been used as out groups in previous studies (Plassais et al.

2019). Their use as an out group is consistent with our results, where we observe lower values

for mean $n_{ROH}$ and mean $s_{ROH}$ (Table 1 and Table 2) in Village dogs relative to breed dogs.

Interestingly, we also observe lower levels of mean $n_{ROH}$ and mean $s_{ROH}$ in Chinese Indigenous

dogs. Our results are in line with the origins of the Chinese Indigenous dogs. Previous work

highlighted relaxed trait selection during establishment of Chinese Indigenous dog population

(Q. Yang et al. 2019), and we believe that this relaxed selection has resulted in overall higher

levels of genetic diversity.

We also examined the relationship between ROH and non-disease traits. We observed

that the relationship between height and $F_{ROH}$ is positive within all breed dogs (Figure 4). This

positive effect size (β) opposes recently published work in human populations (Swinford et al.

2023; Joshi et al. 2015; McQuillan et al. 2012). This may be a result of combining the β from

multiple sizes of dogs (i.e. large and small dogs). When we partition by weight, our results

correspond to previous studies on height in breed dogs (Hoopes et al. 2012). As one might

expect, we also find a positive β when testing for the relationship between $F_{ROH}$ and weight.

Lastly, we observed a negative β with respect to the relationship between $F_{ROH}$ and lifespan,

indicating an association with inbreeding and survival to old age. This result is consistent with

the findings in dogs where an association with disease phenotypes was observed (Mooney,

Yohannes, and Lohmueller 2021). Further, we also partitioned dogs by weight and observed a

trend similar to (Urfer et al. 2019), where larger dogs have a shorter lifespan and smaller dogs

have a longer lifespan, though our results were not significant (Table S8).

We identified multiple SNPs within genes that were associated with height, weight, and

lifespan (Table S13, Table S14, and Table S15). Many of the genes that were observed in our

study were not present in previous work, highlighting the necessity to enhance the correction of

existing genetic structure and explore complex trait architecture in domesticated species. In

humans, there have been 25,551 associations with height, 2,233 associations with weight, and

664 associations with lifespan as of 2023 (Sollis et al. 2023). After correcting for inflation, we

recover a total of 50 associations with height, 84 associations with weight, and 21 associations

with lifespan. This corresponds to previous work which has suggested dogs having a more

simplified complex trait architecture (Boyko et al. 2010; Adam H. Freedman, Lohmueller, and

Wayne 2016).

There some limitations in the work that we would like to highlight. First, the phenotypic

values used in the analysis were calculated based on the average, restricting us to study

associations within breeds. Secondly, breed group categorization based on both neighbor joining

trees and AKC could lead to errors when grouping some of breeds. Additionally, we found that

there was noise in association tests for the last few chromosomes. We believe the noise could be

due to lower marker density on smaller chromosomes.

Despite these limitations, the significant relationship between $F_{ROH}$ and certain

phenotypic traits suggest that inbreeding and artificial selection played a role in shaping complex

non-disease traits within breed dogs. In sum, our work has further elucidated the genetic

structure of quantitative trait architecture in dogs. We also highlight, how domestication and

inbreeding, quantified in the form of ROH, affect the distribution of regions of the genome

associated with complex non-disease phenotypes.

## Author Contributions

ZAS and JAM designed and supervised the research project. S performed the research and

analyzed data. S along with the assistance of ZAS and JAM wrote the paper.

## Acknowledgements

This work was supported by the National Institute of General Medical Sciences of the National

Institutes of Health under Award Number R35GM146926 (ZAS and S). This work was also

supported by Eberly College of Science Startup Fund (ZAS and S). S would also like to thank

the entire Szpiech Lab for their invaluable support. JAM was supported by the startup funds from

Dornsife College of Letters, Arts and Sciences through the Department of Quantitative and

Computational Biology and the USC WiSE Gabilan Assistant Professorship.

## 483    Data Accessibility Statement

484    Whole Genome Sequencing is available on NCBI (Accession number: PRJNA448733). The

485    source phenotype data was obtained from (Plassais et al. 2019). All other data are contained

486    within the article and its supplementary information. Code can be found on GitHub at

487    (https://github.com/jaam92/QuantTraitInDogs.git).

# Bibliography

Alb, James G., Jr, Jorge D. Cortese, Scott E. Phillips, Roger L. Albin, Tim R. Nagy, Bruce A. Hamilton, and Vytas A. Bankaitis. 2003. "Mice Lacking Phosphatidylinositol Transfer Protein-Alpha Exhibit Spinocerebellar Degeneration, Intestinal and Hepatic Steatosis, and Hypoglycemia." *The Journal of Biological Chemistry* 278 (35): 33501–18.

Ali, Aus Tariq, Warren E. Hochfeld, Renier Myburgh, and Michael S. Pepper. 2013. "Adipocyte and Adipogenesis." *European Journal of Cell Biology* 92 (6-7): 229–36.

"American Kennel Club." n.d. American Kennel Club. https://www.akc.org/.

Bacolod, Manny D., Gunter S. Schemmann, Shuang Wang, Richard Shattock, Sarah F. Giardina, Zhaoshi Zeng, Jinru Shia, et al. 2008. "The Signatures of Autozygosity among Patients with Colorectal Cancer." *Cancer Research* 68 (8): 2610–21.

Bannasch, Danika, Thomas Famula, Jonas Donner, Heidi Anderson, Leena Honkanen, Kevin Batcher, Noa Safra, Sara Thomasy, and Robert Rebhun. 2021. "The Effect of Inbreeding, Body Size and Morphology on Health in Dog Breeds." *Canine Medicine and Genetics* 8 (1): 12.

Berg, Sanne van den, Jérémie Vandenplas, Fred A. van Eeuwijk, Marcos S. Lopes, and Roel F. Veerkamp. 2019. "Significance Testing and Genomic Inflation Factor Using High-density Genotypes or Whole-genome Sequence Data." *Journal of Animal Breeding and Genetics* = *Zeitschrift Fur Tierzuchtung Und Zuchtungsbiologie* 136 (6): 418.

Blake, Judith A., Richard Baldarelli, James A. Kadin, Joel E. Richardson, Cynthia L. Smith, and Carol J. Bult. 2021. "Mouse Genome Database (MGD): Knowledgebase for Mouse-Human Comparative Biology." *Nucleic Acids Research* 49 (D1): D981–87.

Boyko, Adam R. 2011. "The Domestic Dog: Man's Best Friend in the Genomic Era." *Genome Biology* 12 (2): 216.

Boyko, Adam R., Pascale Quignon, Lin Li, Jeffrey J. Schoenebeck, Jeremiah D. Degenhardt, Kirk E. Lohmueller, Keyan Zhao, et al. 2010. "A Simple Genetic Architecture Underlies Morphological Variation in Dogs." *PLoS Biology* 8 (8). https://doi.org/10.1371/journal.pbio.1000451.

Cadieu, E., M. W. Neff, P. Quignon, K. Walsh, K. Chase, H. G. Parker, B. M. Vonholdt, et al. 2009. "Coat Variation in the Domestic Dog Is Governed by Variants in Three Genes." 10.1126/science.1177808.

Candille, S. I., C. B. Kaelin, B. M. Cattanach, B. Yu, D. A. Thompson, M. A. Nix, J. A. Kerns, S. M. Schmutz, G. L. Millhauser, and G. S. Barsh. 2007. "A -Defensin Mutation Causes Black Coat Color in Domestic Dogs." 10.1126/science.1147880.

Ceballos, Francisco C., Peter K. Joshi, David W. Clark, Michèle Ramsay, and James F. Wilson. 2018. "Runs of Homozygosity: Windows into Population History and Trait Architecture." *Nature Reviews. Genetics* 19 (4): 220–34.

Cecchi, Francesca, Gisella Paci, Andrea Spaterna, and Roberta Ciampolini. 2013. "Genetic Variability in Bracco Italiano Dog Breed Assessed by Pedigree Data." *Italian Journal of Animal Science*, January. https://doi.org/10.4081/ijas.2013.e54.

Chalazan, Brandon, Denada Palm, Arvind Sridhar, Christina Lee, Maria Argos, Martha Daviglus, Jalees Rehman, Sreenivas Konda, and Dawood Darbar. 2021. "Common Genetic Variants Associated with Obesity in an African-American and Hispanic/Latino Population." *PloS One* 16 (5). https://doi.org/10.1371/journal.pone.0250697.

Chang, Christopher C., Carson C. Chow, Laurent Cam Tellier, Shashaank Vattikuti, Shaun M. Purcell, and James J. Lee. 2015. "Second-Generation PLINK: Rising to the Challenge of Larger and Richer Datasets." *GigaScience* 4 (1): s13742–015 – 0047–0048.

Chang, Qimeng, Yayun Xu, Jianfa Wang, Hui Jing, Longhua Rao, Weiguo Tang, Ziping Zhang,

and Xubo Wu. 2021. "SLC41A3 Exhibits as a Carcinoma Biomarker and Promoter in Liver
Hepatocellular Carcinoma." *Computational and Mathematical Methods in Medicine* 2021.
https://doi.org/10.1155/2021/8556888.
Chauvin, Joe-Marc, Ornella Pagliano, Julien Fourcade, Zhaojun Sun, Hong Wang, Cindy
Sander, John M. Kirkwood, et al. 2015. "TIGIT and PD-1 Impair Tumor Antigen–specific
CD8+ T Cells in Melanoma Patients." *The Journal of Clinical Investigation* 125 (5): 2046–
58.
Chen, Han, Chaolong Wang, Matthew P. Conomos, Adrienne M. Stilp, Zilin Li, Tamar Sofer,
Adam A. Szpiro, et al. 2016. "Control for Population Structure and Relatedness for Binary
Traits in Genetic Association Studies via Logistic Mixed Models." *American Journal of
Human Genetics* 98 (4): 653–66.
Clark, David W., Yukinori Okada, Kristjan H. S. Moore, Dan Mason, Nicola Pirastu, Ilaria
Gandin, Hannele Mattsson, et al. 2019. "Associations of Autozygosity with a Broad Range
of Human Phenotypes." *Nature Communications* 10 (1): 4957.
Costa-Urrutia, Paula, Valentina Colistro, Angélica S. Jiménez-Osorio, Helios Cárdenas-
Hernández, Jacqueline Solares-Tlapechco, Miryam Ramirez-Alcántara, Julio Iván
Granados, and Martha E. Rodríguez-Arellano. 2019. "Genome-Wide Association Study of
Body Mass Index and Body Fat in Mexican-Mestizo Children." *Genes (Basel)* 10 (11): 945.
Danecek, Petr, James K. Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O.
Pollard, Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools."
*GigaScience* 10 (2): giab008.
Freedman, Adam H., Kirk E. Lohmueller, and Robert K. Wayne. 2016. "Evolutionary History,
Selective Sweeps, and Deleterious Variation in the Dog." *Annual Review of Ecology,
Evolution, and Systematics* 47 (Volume 47, 2016): 73–96.
Freedman, A. H., I. Gronau, Schweizer RM, Vecchyo D. Ortega-Del, E. Han, P. M. Silva, M.
Galaverni, et al. 2014. "Genome Sequencing Highlights the Dynamic Early History of
Dogs." *PLoS Genetics* 10 (1). https://doi.org/10.1371/journal.pgen.1004016.
Fu, Lisheng, Xiaoyan Wang, Ying Yang, Meihua Chen, Adilijiang Kuerban, Haojie Liu, Yiwei
Dong, Qianqian Cai, Mingzhe Ma, and Xingzhong Wu. 2023. "Septin11 Promotes
Hepatocellular Carcinoma Cell Motility by Activating RhoA to Regulate Cytoskeleton and
Cell Adhesion." *Cell Death & Disease* 14 (4): 280.
Galis, Frietson, Inke Van der Sluijs, Tom J. M. Van Dooren, Johan A. J. Metz, and Marc
Nussbaumer. 2007. "Do Large Dogs Die Young?" *Journal of Experimental Zoology. Part B,
Molecular and Developmental Evolution* 308 (2): 119–26.
Guetg, Adriano, Luca Mariotta, Lukas Bock, Brigitte Herzog, Ralph Fingerhut, Simone M. R.
Camargo, and François Verrey. 2015. "Essential Amino Acid Transporter Lat4 (Slc43a2) Is
Required for Mouse Development." *The Journal of Physiology* 593 (5): 1273–89.
Hayward, Jessica J., Marta G. Castelhano, Kyle C. Oliveira, Elizabeth Corey, Cheryl Balkman,
Tara L. Baxter, Margret L. Casal, et al. 2016. "Complex Disease and Phenotype Mapping in
the Domestic Dog." *Nature Communications* 7 (1): 1–11.
Hjorthøj, Carsten, Anne Emilie Stürup, John J. McGrath, and Merete Nordentoft. 2017. "Years of
Potential Life Lost and Life Expectancy in Schizophrenia: A Systematic Review and Meta-
Analysis." *The Lancet Psychiatry* 4 (4): 295–301.
Hoopes, Barbara C., Maud Rimbault, David Liebers, Elaine A. Ostrander, and Nathan B. Sutter.
2012. "The Insulin-like Growth Factor 1 Receptor (IGF1R) Contributes to Reduced Size in
Dogs." *Mammalian Genome: Official Journal of the International Mammalian Genome
Society* 23 (11-12): 780–90.
Horard-Herbin, Marie-Pierre, Anne Tresset, and Jean-Denis Vigne. 2014. "Domestication and
Uses of the Dog in Western Europe from the Paleolithic to the Iron Age." *Animal Frontiers* 4
(3): 23–31.
Johnson, Emma C., Luke M. Evans, and Matthew C. Keller. 2018. "Relationships between

Estimated Autozygosity and Complex Traits in the UK Biobank." *PLoS Genetics* 14 (7):
e1007556.
Johnston, Robert J., Laetitia Comps-Agrar, Jason Hackney, Xin Yu, Mahrukh Huseni, Yagai
Yang, Summer Park, et al. 2014. "The Immunoreceptor TIGIT Regulates Antitumor and
Antiviral CD8+ T Cell Effector Function." *Cancer Cell* 26 (6): 923–37.
Joshi, P. K., T. Esko, H. Mattsson, N. Eklund, I. Gandin, T. Nutile, A. U. Jackson, et al. 2015.
"Directional Dominance on Stature and Cognition in Diverse Human Populations." *Nature*
523 (7561). https://doi.org/10.1038/nature14618.
Kang, Sona, Christina N. Bennett, Isabelle Gerin, Lauren A. Rapp, Kurt D. Hankenson, and
Ormond A. MacDougald. 2007. "Wnt Signaling Stimulates Osteoblastogenesis of
Mesenchymal Precursors by Suppressing CCAAT/Enhancer-Binding Protein α and
Peroxisome Proliferator-Activated Receptor γ." *The Journal of Biological Chemistry* 282
(19): 14515–24.
Kang, Yafei, Youming Zhang, Kexin Huang, and Zhenhong Wang. 2023. "The Genetic Influence
of the DRD3 rs6280 Polymorphism (Ser9Gly) on Functional Connectivity and Gray Matter
Volume of the Hippocampus in Patients with First-Episode, Drug-Naïve Schizophrenia."
*Behavioural Brain Research* 437 (February): 114124.
Keller, Matthew C., Matthew A. Simonson, Stephan Ripke, Ben M. Neale, Pablo V. Gejman,
Daniel P. Howrigan, Sang Hong Lee, et al. 2012. "Runs of Homozygosity Implicate
Autozygosity as a Schizophrenia Risk Factor." *PLoS Genetics* 8 (4): e1002656.
Kichaev, Gleb, Gaurav Bhatia, Po-Ru Loh, Steven Gazal, Kathryn Burch, Malika K. Freund,
Armin Schoech, Bogdan Pasaniuc, and Alkes L. Price. 2019. "Leveraging Polygenic
Functional Enrichment to Improve GWAS Power." *American Journal of Human Genetics*
104 (1): 65–75.
Kipp, Anna P., Mike F. Müller, Eva M. Göken, Stefanie Deubel, and Regina Brigelius-Flohé.
2012. "The Selenoproteins GPx2, TrxR2 and TrxR3 Are Regulated by Wnt Signalling in the
Intestinal Epithelium." *Biochimica et Biophysica Acta (BBA) - General Subjects* 1820 (10):
1588–96.
Li, Miao-Xin, Juilian M. Y. Yeung, Stacey S. Cherny, and Pak C. Sham. 2011. "Evaluating the
Effective Numbers of Independent Tests and Significant P-Value Thresholds in Commercial
Genotyping Arrays and Public Imputation Reference Datasets." *Human Genetics* 131 (5):
747–56.
Luo, Jun, Lei Dou, Zhengyan Yang, Zhi Zhou, and Hong Huang. 2020. "CBFA2T2 Promotes
Adipogenic Differentiation of Mesenchymal Stem Cells by Regulating CEBPA." *Biochemical*
*and Biophysical Research Communications* 529 (2): 133–39.
Mao, Yifang, Xingyu Jiang, Peng Guo, Ying Ouyang, Xiangfu Chen, Meng Xia, Lixin Wu, et al.
2023. "ZXDC Enhances Cervical Cancer Metastasis through IGF2BP3-Mediated Activation
of RhoA/ROCK Signaling." *iScience* 26 (8). https://doi.org/10.1016/j.isci.2023.107447.
Martínez-Montes, Angel M., Anixa Muiños-Bühl, Almudena Fernández, Josep M. Folch, Noelia
Ibáñez-Escriche, and Ana I. Fernández. 2016. "Deciphering the Regulation of Porcine
Genes Influencing Growth, Fatness and Yield-Related Traits through Genetical Genomics."
*Mammalian Genome: Official Journal of the International Mammalian Genome Society* 28
(3): 130–42.
Mastrangelo, Salvatore, Filippo Biscarini, Barbara Auzino, Marco Ragatzu, Andrea Spaterna,
and Roberta Ciampolini. 2018. "Genome-Wide Diversity and Runs of Homozygosity in the
'Braque Français, Type Pyrénées' Dog Breed." *BMC Research Notes* 11 (1): 1–6.
Matulewicz, Natalia, Magdalena Stefanowicz, Agnieszka Nikolajuk, and Monika Karczewska-
Kupczewska. 2017. "Markers of Adipogenesis, but Not Inflammation, in Adipose Tissue Are
Independently Related to Insulin Sensitivity." *The Journal of Clinical Endocrinology and*
*Metabolism* 102 (8): 3040–49.
McQuillan, Ruth, Harry Campbell, Veronique Vitart, and James Wilson. 2009. *Homozygosity,*

*Inbreeding and Health in European Populations*.

McQuillan, Ruth, Niina Eklund, Nicola Pirastu, Maris Kuningas, Brian P. McEvoy, Tõnu Esko,
Tanguy Corre, et al. 2012. "Evidence of Inbreeding Depression on Human Height." *PLoS*
*Genetics* 8 (7). https://doi.org/10.1371/journal.pgen.1002655.

Melhem, Nadine M., Cong Lu, Cara Dresbold, Frank A. Middleton, Lambertus Klei, Shawn
Wood, Stephen V. Faraone, et al. 2014. "Characterizing Runs of Homozygosity and Their
Impact on Risk for Psychosis in a Population Isolate." *American Journal of Medical*
*Genetics. Part B, Neuropsychiatric Genetics: The Official Publication of the International*
*Society of Psychiatric Genetics* 165B (6): 521.

Mooney, Jazlyn A., Abigail Yohannes, and Kirk E. Lohmueller. 2021. "The Impact of Identity by
Descent on Fitness and Disease in Dogs." *Proceedings of the National Academy of*
*Sciences of the United States of America* 118 (16).
https://doi.org/10.1073/pnas.2019116118.

Moreno-Grau, Sonia, Maria Victoria Fernández, Itziar de Rojas, Pablo Garcia-González, Isabel
Hernández, Fabiana Farias, John P. Budde, et al. 2021. "Long Runs of Homozygosity Are
Associated with Alzheimer's Disease." *Translational Psychiatry* 11 (1): 1–12.

Parker, H. G., B. M. VonHoldt, P. Quignon, E. H. Margulies, S. Shao, D. S. Mosher, T. C.
Spady, et al. 2009. "An Expressed fgf4 Retrogene Is Associated with Breed-Defining
Chondrodysplasia in Domestic Dogs." 10.1126/science.1173275.

Pemberton, Trevor J., Devin Absher, Marcus W. Feldman, Richard M. Myers, Noah A.
Rosenberg, and Jun Z. Li. 2012. "Genomic Patterns of Homozygosity in Worldwide Human
Populations." *American Journal of Human Genetics* 91 (2): 275–92.

Peritogiannis, Vaios, Angeliki Ninou, and Maria Samakouri. 2022. "Mortality in Schizophrenia-
Spectrum Disorders: Recent Advances in Understanding and Management."
*HealthcarePapers* 10 (12). https://doi.org/10.3390/healthcare10122366.

Plassais, Jocelyn, Jaemin Kim, Brian W. Davis, Danielle M. Karyadi, Andrew N. Hogan, Alex C.
Harris, Brennan Decker, Heidi G. Parker, and Elaine A. Ostrander. 2019. "Whole Genome
Sequencing of Canids Reveals Genomic Regions under Selection and Variants Influencing
Morphology." *Nature Communications* 10 (1): 1–14.

Plummer, Martyn, Nicky Best, Kate Cowles, and Karen Vines. 2006. "CODA: Convergence
Diagnosis and Output Analysis for MCMC." *R News* 6 (1): 7–11.

Purfield, Deirdre C., Sinead McParland, Eamon Wall, and Donagh P. Berry. 2017. "The
Distribution of Runs of Homozygosity and Selection Signatures in Six Commercial Meat
Sheep Breeds." *PloS One* 12 (5). https://doi.org/10.1371/journal.pone.0176780.

Quinlan, Aaron R. 2014. "BEDTools: The Swiss-Army Tool for Genome Feature Analysis."
*Current Protocols in Bioinformatics / Editoral Board, Andreas D. Baxevanis ... [et Al.]* 47
(September): 11.12.1–34.

Sams, Aaron J., and Adam R. Boyko. 2019. "Fine-Scale Resolution of Runs of Homozygosity
Reveal Patterns of Inbreeding and Substantial Overlap with Recessive Disease Genotypes
in Domestic Dogs."

Selionova, Marina, Magomet Aibazov, Tatiana Mamontova, Victor Malorodov, Alexander
Sermyagin, Natalia Zinovieva, and Ahmed A. Easa. 2022. "Genome-Wide Association
Study of Live Body Weight and Body Conformation Traits in Young Karachai Goats." *Small*
*Ruminant Research* 216 (November). https://doi.org/10.1016/j.smallrumres.2022.106836.

Shapses, Sue A., and Deeptha Sukumar. 2012. "Bone Metabolism in Obesity and Weight Loss."
*Annual Review of Nutrition* 32 (August): 287–309.

Sofer, Tamar, Xiuwen Zheng, Cecelia A. Laurie, Stephanie M. Gogarten, Jennifer A. Brody,
Matthew P. Conomos, Joshua C. Bis, et al. 2021. "Variant-Specific Inflation Factors for
Assessing Population Stratification at the Phenotypic Variance Level." *Nature*
*Communications* 12 (1): 1–14.

Sollis, Elliot, Abayomi Mosaku, Ala Abid, Annalisa Buniello, Maria Cerezo, Laurent Gil, Tudor

Groza, et al. 2023. "The NHGRI-EBI GWAS Catalog: Knowledgebase and Deposition Resource." *Nucleic Acids Research* 51 (D1): D977–85.

Sutter, Nathan B., Carlos D. Bustamante, Kevin Chase, Melissa M. Gray, Keyan Zhao, Lan Zhu, Badri Padhukasahasram, et al. 2007. "A Single IGF1 Allele Is a Major Determinant of Small Size in Dogs."

Swinford, Natalie, Brenna Gallagher, Jacob Sheehama, Meng Lin, Sean Prall, Brooke Scelza, and Brenna M. Henn. 2023. "Examination of Runs of Homozygosity in Relation to Height in an Endogamous Namibian Population." *American Journal of Biological Anthropology* 180 (1): 207.

Szpiech, Zachary A., Alexandra Blant, and Trevor J. Pemberton. 2017. "GARLIC: Genomic Autozygosity Regions Likelihood-Based Inference and Classification."

Tachmazidou, Ioanna, Dániel Süveges, Josine L. Min, Graham R. S. Ritchie, Julia Steinberg, Klaudia Walter, Valentina Iotchkova, et al. 2017. "Whole-Genome Sequencing Coupled to Imputation Discovers Genetic Signals for Anthropometric Traits." *American Journal of Human Genetics* 100 (6): 865–84.

Thompson, Elizabeth A. 2013. "Identity by Descent: Variation in Meiosis, Across Genomes, and in Populations." *Genetics* 194 (2): 301.

Trujillo-Viera, Jonathan, Rabih El-Merahbi, Vanessa Schmidt, Till Karwen, Angel Loza-Valdes, Akim Strohmeyer, Saskia Reuter, et al. 2021. "Protein Kinase D2 Drives Chylomicron-Mediated Lipid Transport in the Intestine and Promotes Obesity." *EMBO Molecular Medicine* 13 (5): e13548.

Turner, Stephen D. 2018. "Qqman: An R Package for Visualizing GWAS Results Using Q-Q and Manhattan Plots." *Journal of Open Source Software* 3 (25): 731.

Urfer, Silvan R., Mansen Wang, Mingyin Yang, Elizabeth M. Lund, and Sandra L. Lefebvre. 2019. "Risk Factors Associated with Lifespan in Pet Dogs Evaluated in Primary Care Veterinary Hospitals." *Journal of the American Animal Hospital Association* 55 (3): 130–37.

Vaysse, Amaury, Abhirami Ratnakumar, Thomas Derrien, Erik Axelsson, Gerli Rosengren Pielberg, Snaevar Sigurdsson, Tove Fall, et al. 2011. "Identification of Genomic Regions Associated with Phenotypic Variation between Dog Breeds Using Selection Mapping." *PLoS Genetics* 7 (10): e1002316.

Vilà, Carles, Peter Savolainen, Jesús E. Maldonado, Isabel R. Amorim, John E. Rice, Rodney L. Honeycutt, Keith A. Crandall, Joakim Lundeberg, and Robert K. Wayne. 1997. "Multiple and Ancient Origins of the Domestic Dog." *Science*, June. https://doi.org/10.1126/science.276.5319.1687.

Virakul, Sita, Poorichaya Somparn, Trairak Pisitkun, Peter J. van der Spek, Virgil A. S. H. Dalm, Dion Paridaens, P. Martin van Hagen, Nattiya Hirankarn, Tanapat Palaga, and Willem A. Dik. 2020. "Integrative Analysis of Proteomics and DNA Methylation in Orbital Fibroblasts From Graves' Ophthalmopathy." *Frontiers in Endocrinology* 11: 619989.

vonHoldt, Bridgett M., John P. Pollinger, Kirk E. Lohmueller, Eunjung Han, Heidi G. Parker, Pascale Quignon, Jeremiah D. Degenhardt, et al. 2010. "Genome-Wide SNP and Haplotype Analyses Reveal a Rich History Underlying Dog Domestication." *Nature* 464 (7290): 898–902.

Warrington, Nicole M., Robin N. Beaumont, Momoko Horikoshi, Felix R. Day, Øyvind Helgeland, Charles Laurin, Jonas Bacelis, et al. 2019. "Maternal and Fetal Genetic Effects on Birth Weight and Their Relevance to Cardio-Metabolic Risk Factors." *Nature Genetics* 51 (5): 804–14.

Wu, Wentao, Daning Li, Xiaojie Feng, Fanfan Zhao, Chengzhuo Li, Shuai Zheng, and Jun Lyu. 2021. "A Pan-Cancer Study of Selenoprotein Genes as Promising Targets for Cancer Therapy." *BMC Medical Genomics* 14 (1): 78.

Yang, Qianyong, Hao Chen, Junhua Ye, Chenlong Liu, Rongxing Wei, Congying Chen, and Lusheng Huang. 2019. "Genetic Diversity and Signatures of Selection in 15 Chinese

Indigenous Dog Breeds Revealed by Genome-Wide SNPs." *Frontiers in Genetics* 10 (November): 489469.

Yang, Wanneng, Zilong Guo, Chenglong Huang, Lingfeng Duan, Guoxing Chen, Ni Jiang, Wei Fang, et al. 2014. "Combining High-Throughput Phenotyping and Genome-Wide Association Studies to Reveal Natural Genetic Variation in Rice." *Nature Communications* 5 (1): 1–9.

Yengo, Loïc, Sailaja Vedantam, Eirini Marouli, Julia Sidorenko, Eric Bartell, Saori Sakaue, Marielisa Graff, et al. 2022. "A Saturated Map of Common Genetic Variants Associated with Human Height." *Nature* 610 (7933): 704–12.

Yordy, Jennifer, Cornelia Kraus, Jessica J. Hayward, Michelle E. White, Laura M. Shannon, Kate E. Creevy, Daniel E. L. Promislow, and Adam R. Boyko. 2019. "Body Size, Inbreeding, and Lifespan in Domestic Dogs."

Yuan, Jie, Xueqiang Wang, Yan Zhao, Najeeb Ullah Khan, Zhiqiang Zhao, Yanhong Zhang, Xiaorong Wen, Fusen Tang, Fengbin Wang, and Zichao Li. 2020. "Genetic Basis and Identification of Candidate Genes for Salt Tolerance in Rice by GWAS." *Scientific Reports* 10 (1): 9958.

Zhao, Weizhu, Haiying Yang, Jie Chai, and Ligang Xing. 2021. "RUNX2 as a Promising Therapeutic Target for Malignant Tumors." *Cancer Management and Research* 13 (March): 2539–48.