Silicone Wire Embolization-induced Acute Retinal Artery Ischemia and Reperfusion Model in Mouse: Gene Expression Provides Insight into Pathological Processes

Yuedan Wang[a#], Ying Li[a#], Jiaqing Feng[a#], Chuansen Wang[a], Yuwei Wan[a], Bingyang Lv[a], Yinming Li[b], Hao Xie[b], Ting Chen[a], Faxi Wang[b], Ziyue Li[a], Anhuai Yang[a*] & Xuan Xiao[a,b*]

a Department of Ophthalmology, Renmin Hospital of Wuhan University, Wuhan 430060, Hubei, China
b Department of Clinical Laboratory, Institute of Translational Medicine, Renmin Hospital of Wuhan University, Wuhan 430060, Hubei, China
# These authors contributed equally.
* **Correspondence to:** Xuan Xiao, Department of Ophthalmology, Renmin Hospital of Wuhan University, No. 238 Jie Fang Road, Wuhan 430060, Hubei, China. Tel: (86) 027-88041911, Email: xiaoxuan1111@whu.edu.cn
* **Correspondence to:** Anhuai Yang, Department of Ophthalmology, Renmin Hospital of Wuhan University, No. 238 Jie Fang Road, Wuhan 430060, Hubei, China. Email: yah0525@126.com

Abstract

Acute retinal ischemia and ischemia-reperfusion injury are the primary causes of retinal neural cell death and vision loss in retinal artery occlusion (RAO). The absence of an accurate mouse model for simulating the retinal ischemic process has hindered progress in developing neuroprotective agents for RAO. We developed a unilateral pterygopalatine ophthalmic artery occlusion (UPOAO) mouse model using silicone wire embolization combined with carotid artery ligation. The survival of retinal ganglion cells and visual function were evaluated to determine the duration of ischemia. Immunofluorescence staining, optical coherence tomography, and haematoxylin and eosin staining were utilized to assess changes in major neural cell classes and retinal structure degeneration at two reperfusion durations. Transcriptomics was employed to investigate alterations in the pathological process of UPOAO following ischemia and reperfusion, highlighting transcriptomic differences between UPOAO and other retinal ischemia-reperfusion models. The UPOAO model successfully replicated the acute interruption of retinal blood supply observed in RAO. 60-minutes of Ischemia led to significant loss of major retinal neural cells and visual function impairment. Notable thinning of the inner retinal layer, especially the ganglion cell layer, was evident post-UPOAO. Temporal transcriptome analysis revealed various pathophysiological processes related to immune cell migration, oxidative stress, and immune inflammation during the non-reperfusion and reperfusion periods. A pronounced increase in microglia within the retina and peripheral leukocytes accessing the retina was observed during reperfusion periods. Comparison of differentially expressed genes (DEGs) between the UPOAO and high intraocular pressure models revealed specific enrichments in

lipid and steroid metabolism-related genes in the UPOAO model. The UPOAO model
emerges as a novel tool for screening pathogenic genes and promoting further therapeutic
research in RAO.

Key words: Retinal artery occlusion; Mouse model; Ischemia and reperfusion injury;
Transcriptomics; Immune cell

1 Introduction
Retinal artery occlusion (RAO) is a severe ophthalmic disease characterized by a sudden
interruption of blood flow in the retinal artery, leading to retinal ischemia [1]. Over 60% of
RAO patients suffer from impaired vision, ranging from finger counting to complete vision
loss [2]. Additionally, RAO patients face increased risk of cardiovascular and cerebrovascular
events [3-5]. Retinal ischemia and hypoxia can lead to irreversible damage to retinal cells
within 90 minutes [6]. This damage is primarily due to the apoptosis of retinal ganglion cells
(RGCs), which is driven by inflammation and oxidative stress. Unfortunately, conservative
treatments for RAO, such as ocular massage and hyperbaric oxygen therapy, provide limited
therapeutic benefits [7]. Although thrombolysis is effective, its application is restricted by a
narrow therapeutic window (typically within 4.5 hours) [8]. Furthermore, the restoration of
blood flow to the ischemic area following thrombolysis can cause a sudden increase in
tissue oxidative levels, resulting in ischemia-reperfusion injury (IRI) in the retina [9]. IRI is a
common pathological condition that can induce inflammation, retinal tissue damage and
visual impairment [10]. Therefore, it is crucial to develop an animal model that accurately
simulates the pathological processes of RAO to thoroughly investigate the
pathophysiological changes and explore potential neuroprotective treatments following
retinal ischemia and reperfusion.

Various animal models have been employed to investigate the effects of retinal injury
resulting from ischemia and subsequent reperfusion in RAO [6, 11, 12]. Based on the
methods used for modelling, retinal ischemia-reperfusion models can be categorized into
two groups: 1) intravascular occlusion models, including photochemical-induced
thrombosis model and vascular intervention model; and 2) extravascular occlusion models,
including central retinal artery ligation (CRAL) model, unilateral common carotid artery
occlusion (UCCAO) model, and high intraocular pressure (HIOP) model. The photochemical-
induced thrombosis model and vascular intervention model have been reported to be
useful tools for assessing retinal injury after ischemia and reperfusion. However, both
models are limited by the need for trained interventional radiologists and the need for
advanced techniques [13, 14]. The CRAL model, which directly clips or ligates the central
retinal artery, also presents drawbacks, such as potential optic nerve damage and difficulty
in application to small animals such as mice, limiting its use in experiments with larger
sample sizes [6]. The UCCAO model, which induces retinal hypoperfusion by occluding the
unilateral common carotid artery (CCA), is more suitable for simulating chronic ischemic
retinal disease than acute retinal ischemia [15, 16]. The HIOP model is widely utilized in mice
for studying ischemia-reperfusion in acute primary angle-closure glaucoma (APACG) [17].
However, the mechanical compression of the retina caused by saline injection in this model
may also lead to retinal damage, potentially influencing IRI research. Therefore, it is critically

important to develop a simple and low-skill-required mouse model that can simulate acute
retinal ischemia and reperfusion injury in RAO patients.

To better simulate the retinal ischemic process and possible IRI following RAO, we
developed a novel vascular-associated mouse model called the unilateral pterygopalatine
ophthalmic artery occlusion (UPOAO) model. In this model, we employed silicone wire
embolization and carotid artery ligation to completely block the blood supply to the retina.
We characterized the major classes of retinal neural cells and evaluated visual function
following different durations of ischemia (30 minutes and 60 minutes) and reperfusion (3-
days and 7-days) after UPOAO. Additionally, we utilized transcriptomics to investigate the
transcriptional changes and elucidate the pathophysiological process of the UPOAO model
after ischemia and reperfusion. Finally, we highlighted the transcriptomic differences
between this model and other retinal ischemia-reperfusion models, including HIOP and
UCCAO, revealing the unique pathological processes that closely resemble retinal IRI in RAO.
The UPOAO model offers new insights into the mechanisms and pathways involved in
ischemia and reperfusion studies of RAO, providing a foundation for studying protective
strategies for ocular ischemic diseases.

2 Methods and materials
2.1 Animals
Eight-week-old male C57BL/6 mice were used in the experiments. Only male mice were
used to exclude the potential influence of oestrogenic hormones. The mice were provided
sufficient food and water and were maintained on a 12-hour dark-light cycle in a room with
regulated temperature conditions. All the experimental procedures were designed and
conducted according to the ethical guidelines outlined in the Association for Research in
Vision and Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic and
Vision Research. The study protocol and methods received approval from the Experimental
Animal Ethics Committee of Renmin Hospital of Wuhan University (approval number:
WDRM-20220305A).

2.2 Preparation and surgical procedure of the UPOAO model
Eight-week-old male C57BL/6 mice weighing 20-25 grams were used in this study. An
isoflurane-based anaesthesia system was used to induce and maintain general anaesthesia
in the mice during surgical procedures. The mice were anaesthetized with a 1.5-2%
concentration of isoflurane delivered in a mixture of nitrous oxide and oxygen via rubber
tubing. Body temperature was maintained at 37 ± 0.5°C throughout the surgery. Surgical
instruments were sterilized with 75% ethanol before use to ensure sterility.

We drew inspiration from the widely employed middle cerebral artery occlusion (MCAO)
model [18, 19], commonly used in cerebral ischemic injury research, which guided the
development of the UPOAO model. The mouse was gently positioned in a supine posture
on a heating blanket, and its neck was exposed. The preparation involved depilation of the
neck area, followed by skin disinfection, and finally, a midline incision along the neck was
made. After separating the neck gland using two tweezers, careful blunt dissection was

employed to separate the left CCA, internal carotid artery (ICA) and external carotid artery (ECA), avoiding the compression of nearby nerves and veins (**Fig. 1A**) (**Video 1**). Then, the CCA and ICA were ligated with a 6-0 suture. A knot was secured using a 6-0 suture at the distal end of the ECA, and a slipknot was created at the proximal end (**Fig. 1B**). To preserve the reperfusion process and avoid bleeding and mortality during surgery, a silicone wire embolus was inserted through the ECA instead of through the CCA. Ophthalmic scissors were used to make a small inverted "V"-shaped incision between the two suture knots on the ECA. A specialized silicone wire embolus, measuring 7 ± 0.1 mm in length and 0.21 ± 0.1 mm in diameter, was inserted through the incision, directing it into the ECA and further into the CCA (**Fig. 1C**). Next, the ligation on the ICA was removed, and the ECA was cut at the point where the artery had been previously incised with scissors. Subsequently, the silicone wire embolus was retracted to the bifurcation of the CCA, rotated counterclockwise, and inserted into the ICA and further into the pterygopalatine artery (PPA) (**Fig. 1D1**). The silicone tail of the silicone wire embolus was positioned near the bifurcation of the CCA (**Fig. 1D2**), effectively obstructing the ophthalmic artery (OA) (**Video 2**). Then, the slipknot was fastened, and the skin was sutured. The mouse was free to move around during arterial embolization.

Following a predetermined ischemia embolization time, the silicone wire embolus was carefully removed from the PPA without massive haemorrhage (**Fig. 1E**). The CCA suture was removed, restoring arterial reperfusion (**Fig. 1F**) (**Video 3**). Then, the skin wound was sutured, and the mice were routinely fed during reperfusion (**Fig. 1G**).

2.3 Staining of retinal vessels

To clearly visualize the retinal vessels, rhodamine-labelled canavalin A was used for mouse heart perfusion. Each mouse was anaesthetized with 1% sodium pentobarbital and placed in the supine position on a foam board resting on a tray. The sternum was opened to expose the heart. A needle attached to the perfusion tube was inserted into the left ventricle via the apex of the mouse heart and secured. The right atrial appendage was dissected with scissors to allow blood outflow. Saline solution was perfused through the left ventricle using a pump system to drain blood from the arteries. Subsequently, the prepared solution of rhodamine-labelled canavalin A was injected into the mice and circulated throughout the mouse vessels for 30 minutes. The mice were then sacrificed, and their eyes were removed and immersed in paraformaldehyde (PFA) for one hour in the dark. A flat-mounted retina was obtained, and retinal images were captured using a Leica SP8 confocal microscope equipped with a 10× objective.

2.4 Quantification of RGCs and microglia

The mice were euthanized by cervical dislocation, and their eyes were fixed in 4% PFA for 60 minutes at room temperature (RT). Subsequently, the cornea and lens were excised, and the intact retina was isolated for retinal flat mounting. The retina was immersed in PBS supplemented with 5% bovine serum albumin (BSA) and 0.5% Triton X-100 for overnight blocking and then incubated with Brn3a antibody (Santa Cruz Biotechnology, Dallas, Texas, USA) or Iba1 antibody (Wako, Japan). Following a two-day incubation with the primary

antibody, the retina was gently washed three times with PBS. Subsequently, the retina was incubated with Alexa Fluor 594 (AntGene, Wuhan, China) in a light-protected cassette for two days. After three rinses, the retina was uniformly sectioned into a four-leaf clover morphology and flattened using a coverslip. Finally, retinal flat mounts labelled with Brn3a were imaged using a Leica SP8 confocal microscope (Leica TCS SP8, Germany) equipped with a 20× objective, while those labelled with Iba1 were imaged utilizing a fluorescence microscope (BX63; Olympus, Tokyo, Japan). Each quadrant was consistently subdivided into central, middle and peripheral fields at regular intervals. The surviving RGCs and activated microglia were quantified and averaged using ImageJ software (National Institutes of Health, USA).

2.5 Electroretinogram (ERG)

Mice were dark-adapted overnight before ERG, and all subsequent procedures were carried out in the dark. Before ERG, the mice were anaesthetized with 1% sodium pentobarbital via intraperitoneal injection, and their pupils were dilated. A subcutaneous electrode was inserted into the posterior cervical skin, a tail electrode was affixed to the posterior end of the mouse tail, and the corneal electrode was gently placed on the central corneal surface. A RetiMINER-C visual electrophysiological system (3VMED Co., Ltd., Shanghai, China) was used for recording electrical responses. The a-waves, b-waves, and oscillatory potentials (OPs) after flash stimuli of 0.01, 0.03, 0.1, 0.3, 1.0, 3.0, and 10.0 cd.s/m^2 in scotopic adaptation were recorded. The amplitudes and implicit times of a-waves, b-waves, and OPs in response to various flash stimuli were analysed.

2.6 Optical Coherence Tomography Imaging (OCT) and Fluorescein Fundus Angiography (FFA)

The mice were placed on the platform of a Spectralis HRA + OCT device (Heidelberg Engineering, Heidelberg, Germany) for OCT imaging following anaesthesia. Pupil dilation was performed, and normal saline was applied regularly to maintain corneal moisture. The focal length of the device was adjusted until the mouse retina was clearly visible. The head of the mouse was gently repositioned to capture images of the peripheral fundus. The multiline mode was used to scan each layer of the retina, and four quadrants of view centred on the nipple in the upper left, lower left, upper right, and lower right were recorded. In this study, the total retinal thickness was manually segmented into three parts and encompassed the entire thickness from the nerve fibre layer to the photoreceptor layer. The ganglion cell complex (GCC) was defined as the combined thickness of the retinal nerve fibre layer (RNFL), ganglion cell layer (GCL), and inner plexiform layer (IPL). The inner nuclear layer (INL) and the outer plexiform layer (OPL) were combined for thickness analysis due to the difficulty in distinguishing between these layers. The remaining retinal layers, including the outer nuclear layer (ONL), inner segment/outer segment (IS/OS), and retinal pigment epithelium (RPE) layers, were measured together. The thickness of the retina at 1.5 papillary diameters (PD), 3.0 PD, and 4.5 PD from the centre of the optic disc was measured using the Heidelberg measuring tool.

For FFA, the mice were anaesthetized, and their pupils were dilated. Subsequently, fundus

221 angiography was performed immediately following an intraperitoneal injection of sodium
222 fluorescein. Images were acquired alternately for both eyes within 5 seconds of the injection.
223
2.7 Hematoxylin and eosin (HE)
Mouse eyes were fixed in FAS eyeball fixation solution (Service-bio, Wuhan, China) for 48
hours, followed by dehydration and subsequent embedding in paraffin. The paraffin blocks
were trimmed parallel to the optic nerve to obtain 3-4 µm thick sections where the optic
nerve was located. Six paraffin sections from each eyeball were stained with HE. Images
were captured using an Olympus fluorescence microscope. The retinal thickness within 500
µm of the optic disc was measured using Image-Pro Plus 6.0 software.
231
2.8 Staining of Retinal Sections
For immunofluorescence, mouse eyes were fixed in 4% PFA solution and subjected to
gradient dehydration using 10%, 20%, and 30% sucrose solutions. The following day, the eyes
were embedded in optimum cutting temperature (OCT) compound (SAKURA, USA) and
frozen at -80°C. Several 14-µm-thick frozen sections were obtained from each eyeball
through the use of a freezing microtome (Leica, Wetzlar, Germany). The section surface was
parallel to the optic nerve, and 3-6 frozen sections were mounted on a single slide. The
slides were stored at -20°C until use.
240
All sections were blocked with 5% BSA and 0.5% Triton X-100 in PBS for 2 hours. Primary
antibodies, as listed in **Table 1**, were diluted in PBS containing 5% BSA and 0.5% Triton X-100
and incubated at 4°C overnight. The following day, after three rinses, the sections were
incubated with secondary antibodies at RT for 2 hours in a cassette, followed by three
washes. The sections were then stained with DAPI (Service-bio, Wuhan, China) for 15
minutes. After the final three rinses, the frozen sections were sealed. A Leica SP8 instrument
equipped with a 40× objective was used to photograph frozen sections near the optic nerve
head. The fluorescence intensity of all slices of each eye was determined using ImageJ
software.
250
2.9 Surgical procedure for the UCCAO model and the HIOP model
Animals for the UCCAO model were prepared and managed according to the procedure
described for the UPOAO model. After blunt dissection, the left CCA was ligated with a 6-0
suture. After 60 minutes of ligation, the suture was removed, and the skin wound was
subsequently closed with sutures. Mice were routinely fed during the reperfusion period,
and the retinas of UCCAO mice were harvested 7-days post-surgery.
257
The HIOP model was generated based on previously established methods [20]. Briefly, mice
were anaesthetized by intraperitoneal injection of 1% sodium pentobarbital. Pupils were
dilated using tropicamide phenylephrine eye drops (0.5% tropicamide and 0.5%
deoxyadrenaline hydrochloride, Santen Pharmaceutical) applied 5 minutes in advance. An
insulin hypodermic needle attached to a silicone elastomer tube was inserted into the
anterior chamber of the unilateral experimental eye. The IOP was elevated for 60 minutes by
raising the height of the saline solution storage bag to 162 cm, after which the needle was

removed to immediately restore the IOP to baseline levels. Iris whitening and loss of retinal
red-light reflexes indicated retinal ischemia. After surgery, levofloxacin hydrochloride
ophthalmic gel was applied to the eyes of the mice. Retinas from HIOP mice were collected
7-days post-surgery.
269
2.10 Transcriptome sequence and analysis
Retinal samples from UPOAO, HIOP, and UCCAO mice were extracted and promptly frozen
in liquid nitrogen within 10 minutes of cervical dislocation. Total RNA extraction was
performed according to the protocol outlined in the TRIzol Reagent manual (Life
Technologies, CA, USA). RNA integrity and concentration were assessed using an Agilent
2,100 Bioanalyzer (Agilent Technologies, Inc., Santa Clara, CA, USA). The resulting RNA
samples were then pooled and subjected to sequencing on the Illumina HiSeq3000 platform
in a 150-bp paired-end read format. The raw RNA-sequencing (RNA-seq) reads were
preprocessed and quantified using the featureCounts function in the SubReads package
version 1.5.3, with default parameters.
280
281 RNA-seq analysis was conducted on samples from 5 mice in the non-perfusion group, 4
282 mice in the 3-days perfusion group, and 5 mice in the 7-days perfusion group of the
283 UPOAO model. Additionally, retinas from both the 7-days HIOP experimental eyes and
284 bilateral UCCAO eyes were collected (n=5) for analysis.
285
Data normalization and subsequent processing utilized the 'limma' package in R software
(version 4.2.3) [21]. Volcano plots and Venn diagrams were generated for visualization to
identify significantly differentially expressed genes (DEGs) and perform Gene Ontology (GO)
annotation analysis. DEGs with a |log2-fold change (FC)| > 1 and adjusted p value < 0.05
were significantly differentially expressed. Enrichment analysis focused on DEGs with an
adjusted p < 0.05 and an enriched gene count > 5. To visualize the expression patterns of
significant DEGs, a heatmap was generated using the 'pheatmap' package. Gene Set
Enrichment Analysis (GSEA) was conducted using the GSEA program (version 4.3.2) [22],
employing the default gene set m-subset (mh.all.v2023.1.mm.symbols.gmt) to explore
significant functional and pathway differences. Enriched pathways were classified based on
criteria such as adjusted p value (< 0.05), false discovery rate (FDR) q value (< 0.25), and
normalized enrichment score (|NES| > 1). Furthermore, protein−protein interaction (PPI)
analysis was performed using the Search Tool for the Retrieval of Interacting Genes (STRING)
database (https://string-db.org/) [23]. Genes with a greater degree of protein-level
interactions with others were further analysed using Cytoscape software (version 3.8.2) to
generate a downstream PPI map. PPI pairings with an interaction score > 0.7 were extracted
and visualized using Cytoscape 3.9.0 [24].
303
2.11 Real-time Quantitative polymerase chain reaction (RT-qPCR)
RNA was extracted from fresh mouse retinal samples and subjected to reverse transcription
using HiScript III RT SuperMix for qPCR (Vazyme, China) following the manufacturer＇s
instructions. RT−qPCR analysis was conducted on a CFX instrument (Bio-Rad) using AceQ
qPCR SYBR Green Master Mix (Vazyme, China). The PCR program comprised 40 cycles of 10

seconds at 95°C and 30 seconds at 60°C. Assays were performed in triplicate, and Ct values
were normalized to beta-actin levels. Relative quantification of target gene expression was
calculated using the 2-ΔΔCt method. The primer sequences are provided in **Table S1**.
312
2.12 Statistical Analysis
All the data were analysed using GraphPad Prism version 9.0 (GraphPad Software, San
Diego, CA, USA). Each experimental group included a minimum of three biological
replicates. T tests, including RGCs and microglia counting, OPs analysis, retinal thickness
measurements via HE staining, and fluorescence intensity quantification, were used to
compare the two groups. Two-way ANOVA (two-way analysis of variance) test was used for
more than two different experimental groups, including ERG waves and retinal thickness
measurements in OCT. All data are presented as the mean ± standard error of the mean
(SEM), and the statistical graphs are shown as scatter bar graphs and line charts. $P < 0.05$
was considered to indicate statistical significance.
323
3 Results
3.1 Silicone wire embolus insertion interrupts retinal blood flow
To reproduce the retinal ischemic process and potential IRI in RAO, we established a mouse
model of UPOAO by combining silicone wire embolization with carotid artery ligation.
Model validation was conducted through a series of experiments, with detailed descriptions
provided in the Materials and Methods section (**Fig. 1A-G**). To confirm blood flow
disruption, we performed cardiac perfusion with fluorescently labelled lectin and FFA.
Rhodamine-labelled canavalin A was used during in vivo perfusion with silicone wire
embolus, and both eyes were evaluated. The sham eye exhibited normal blood perfusion in
the retina, while the experimental eye showed no perfusion (**Fig. 1H, I**). FFA revealed
delayed and limited perfusion in the experimental lateral retina, primarily near the optic disc
(**Fig. 1J, K**). These findings indicate that the insertion of the silicone wire embolus effectively
impaired blood flow to the retina.
337
338 3.2 60-minute ischemia in UPOAO damage retinal structure and function
339 To investigate the optimal ischemic duration, the mice were subjected to ischemic periods
340 of 30 and 60 minutes, followed by reperfusion periods of 3-days and 7-days. Retinal
341 structure and visual function were assessed through flat-mounted retina analysis and flash
342 ERG.
343
In the 30-minute ischemia group, no significant RGCs death was observed after 3-days (**Fig.
2A, B**) or 7-days of reperfusion (**Fig. 2C, D**). However, after 60 minutes of ischemia followed
by either 3-days or 7-days of reperfusion, reductions in RGCs density were evident (**Fig. 2E-
H**). ERG results showed no statistical difference in amplitudes between bilateral eyes after 3-
and 7-days of 30-minutes reperfusion (**Fig. 3A-C**). However, the b-wave amplitude notably
decreased after 60 minutes of ischemia and 3-days of reperfusion (**Fig. 3D**). By 7-days of
reperfusion, the b-wave amplitude had halved compared to that of the sham eyes (**Fig. 3E**).
The appearance times of the a-waves and b-waves in the experimental and control eyes
remained consistent across all four groups (**Fig. S1**). Additionally, OPs have also been used

to sensitively monitor retinal ischemic effects by detecting changes before any alterations in
b-waves occur [25-27]. The amplitudes of OPs in the experimental eyes, particularly in the
60-minute ischemia and 7-days reperfusion group, significantly decreased to less than 50%
of those in the sham eyes (**Fig. 3F, G**). Based on RGCs survival and changes in ERG waves,
we determined that a 60-minute ischemic duration is optimal for the UPOAO model.
358
3.3 Evaluation of retinal thickness in UPOAO through OCT and HE
To assess retinal thickness non-invasively in the UPOAO model, we conducted OCT imaging
during 3-days and 7-days reperfusion periods. Following a 3-days reperfusion period, no
significant changes were observed in the GCC, INL + OPL, and ONL + IS/OS + RPE layers at
1.5, 3, and 4.5 PD from the optic disc (**Fig. 4B, Fig. S2A, B**). However, the total retinal
thickness decreased at 4.5 PD (**Fig. 4C**). After 7-days of reperfusion, the total retinal
thickness decreased at 1.5 PD, 3 PD, and 4.5 PD, primarily due to inner retinal thinning (**Fig.
4D-F, Fig. S2C, D**).
367
For a detailed analysis of inner retinal thickness, we extracted UPOAO mouse eyes and
stained them with HE. The thickness of the RNFL + GCL increased at 3-days of reperfusion
(**Fig. 4H**), followed by a decrease at 7-days of reperfusion (**Fig. 4L**). IPL thickness decreased
at both 3-days and 7-days of reperfusion (**Fig. 4I, M**), while INL thickness decreased only at
7-days of reperfusion (**Fig. 4N**). We hypothesized that the initial swelling of the GCL in
response to acute ischemia and hypoxia led to early thickening, followed by gradual
thinning due to tissue dysfunction. This hypothesis was supported by the observation of
RGCs loss and reduced GCC thickness evaluated by OCT at 7-days (**Fig. 4E**).
376
377   We observed by OCT that changes in retinal thickness became noticeable at 3-days post-
378   UPOAO and exhibited a significant decrease at 7-days. HE results further revealed that
379   alterations in inner retinal thickness constituted a substantial portion of the total retinal
380   thickness during the early reperfusion stage after UPOAO.
381
3.4 Survival of retinal neural cells in UPOAO
Given that mouse retinal function heavily relies on rod cells and that scotopic (low light)
vision is primarily governed by rod photoreceptors, the observed reductions in b-wave and
OPs-wave amplitudes indicate impaired synaptic transmission within the inner retina [28].
Our OCT and HE findings suggested structural injuries within the inner retina. Notably, the
thickness of the outer retinal layers in OCT remained unchanged. To further explore these
observations, we conducted immunofluorescence staining to investigate alterations in major
retinal cell types, primarily focusing on bipolar cells (BCs), photoreceptor cells, horizontal
cells (HCs), and cholinergic amacrine cells.
391
392   3.4.1 BCs loss and photoreceptor cells survival
393   PKCα serves as a marker delineating BCs, with their cell bodies primarily located in the
394   outermost part of the INL, axonal terminals extending into the innermost part of the IPL,
395   and dendrites confined to the OPL (**Fig. 5A, B**). In the experimental UPOAO eyes, BCs did
396   not show significant changes at the 3-days reperfusion mark but exhibited dramatic

alterations by the 7-days reperfusion period (**Fig. 5A2, B2**). In particular, the immunostaining density of BCs dendritic and axonal arbors notably decreased at the 7-days reperfusion mark. BCs account for approximately 40% of INL cells in mice, and their somata and axonal processes form a substantial part of the INL and IPL, consistent with the thinning of the INL observed in both OCT and HE at 7-days (**Fig. 4E, N**). Notably, the decrease in the number of PKCα$^+$ BCs at the 7-days reperfusion point coincided with the decrease in the b-waves amplitude in ERG (**Fig. 3E**), indicating functional interplay between BCs and other cells.

Recoverin marks the somata and outer segments of photoreceptor cells and is localized within the ONL (**Fig. S3A, B**). Interestingly, recoverin-positive photoreceptors remained relatively stable throughout the reperfusion periods, which aligns with the unchanged thickness of the outer retinal layers observed in OCT (**Fig. S2C, D**).

3.4.2 HCs and cholinergic amacrine cells loss

We evaluated HCs and cholinergic amacrine cells using calbindin and ChAT immunostaining, respectively. Calbindin immunostaining highlighted the somata of HCs within the INL, with terminal axon connections linearly positioned within the OPL (**Fig. 5E, F**). In the UPOAO experimental eyes, no significant change in the number of HCs was observed during the 3-days reperfusion period, while a notable reduction was observed after 7 days. By 7-days, a significant decrease in cell body numbers, a considerable reduction in axon density, and disrupted linear connections were observed (**Fig. 5F2**).

ChAT$^+$ cell bodies were located in the GCL and INL, with dendrites forming two narrow stratified bands within the IPL. The immunofluorescence density of ChAT$^+$ amacrine cells decreased notably after 3 days and even more prominently after 7 days (**Fig. 5I, J**). The fluorescence intensity of the two bands within the IPL markedly decreased and was nearly invisible after 7 days (**Fig. 5J2**).

3.5 Time course transcriptome analysis revealed features of different reperfusion periods in UPOAO

To explore the pathophysiological processes of UPOAO, we extracted retinas from both eyes for transcriptome sequencing. The samples included retinas from eyes subjected to 60 minutes of ischemia without reperfusion, 60 minutes of ischemia followed by 3-days of reperfusion, and 60 minutes of ischemia followed by 7-days of reperfusion, with the sham-treated eyes serving as controls.

In the non-perfusion group, 215 genes were upregulated and 204 genes were downregulated (**Fig. 6A**). GO enrichment analysis revealed that the DEGs were related to leukocyte migration, epidermis development, myeloid leukocyte migration and other cell migration pathways (**Fig. 6B**). The heatmap and box showed the upregulated and downregulated genes involved in immune cell migration-related pathways during the non-perfusion period in UPOAO (**Fig. 6C**). High-connectivity DEGs ('hub genes'), such as Dusp1 and Fos, were also enriched in these pathways (**Fig. S4A**). GSEA also showed similar results

(**Fig. S4B-D**). These results suggested that in the early stage of retinal ischemic injury, leukocytes from the microvasculature may infiltrate retinal tissue. More experimental validation will be performed to confirm this hypothesis.

In the 3-days reperfusion group, 372 genes were upregulated, and 170 genes were downregulated (**Fig. 6D**). GO enrichment analysis revealed that the DEGs at 3-days after reperfusion were related to energy metabolism, mitochondrial regulation, and oxidative stress pathways (**Fig. 6E**). The heatmap and box displayed the upregulated and downregulated genes involved in oxidative stress-related pathways at the 3-days reperfusion stage in UPOAO, including hub genes such as Mrpl18 and Mrps23 (**Fig. 6F, Fig. S5A**). GSEA revealed changes in pathways related to the binding membrane of organelles, the cell surface and the plasma membrane protein complex (**Fig. S5B**). Furthermore, 44 overlapping genes between the mitochondrial genes and DEGs at 3-days post-UPOAO were mainly related to mitochondrial transport and the mitochondrial tricarboxylic acid cycle (**Fig. S5C, D**). These results suggested that mitochondria and metabolism play significant roles in IRI.

In the 7-days reperfusion group, 429 genes were upregulated, and 310 genes were downregulated (**Fig 6G**). The DEGs at this stage were enriched in pathways related to immune effector processes, positive regulation of responses to external stimuli and inflammation (**Fig. 6H**). A heatmap was generated to show the upregulated and downregulated genes of immune inflammation-related pathways at 7-days after reperfusion in UPOAO, and most hub genes, such as Cd86, Cd48, Tlr4, and Tlr6 were also enriched in these pathways (**Fig. 6I, Fig. S6A**). GSEA also showed similar results (**Fig S6B**). Moreover, RT−qPCR confirmed the upregulation of immune inflammation-related gene expression (**Fig. S7**). Significantly, 112 genes that overlapped between immune-related genes and DEGs at 7-days post-UPOAO were primarily associated with immune receptor activity and phagocytic vesicles (**Fig. S6C, D**). These results underscore the predominant role of immune responses during this stage.

To explore the associations between changes observed at 3-days and 7-days after reperfusion, we analyzed the DEGs and identified 17 overlapping genes (**Fig. S8A**), which were mainly associated with the nitric oxide biosynthetic process (**Fig. S8B**). Our findings suggest that the immune inflammatory response observed after 7-days of reperfusion may represent the cumulative effect of the acute oxidative stress response during the initial 3-days reperfusion period.

3.6 Leukocyte Infiltration and Microglial Activation

Time course transcriptome analysis underscored the predominant role of immune inflammatory responses during the reperfusion stage. Previous reports have indicated that the inflammatory response in retinal ischemia-reperfusion injury (RIRI) is orchestrated by peripheral immune cells and resident immune cells within the retina [29, 30]. In our study, we demonstrated the presence of leukocyte infiltration and microglial activation in the UPOAO model. Immunofluorescence staining for CD45 was performed on retinas at

different time points post modelling to visualize the distribution and quantity of white blood

cells (**Fig. 7A-D**). Laser confocal Z-plane projections confirmed minimal leukocyte

infiltration in the retinas of sham eyes, which showed that CD45$^+$ cells in the vascular lumen

likely represented patrolling cells (**Fig. 7A**). A significant increase in CD45$^+$ cells was

observed in UPOAO model retinas after 1 day (**Fig. 7E**), with a progressive increase in

quantity over time (**Fig. 7F, G**). Notably, the majority of CD45$^+$ cells in UPOAO model retinas

at 1-day displayed a morphology similar to that of vascular leukocytes (**Fig. 7B**), suggesting

a large influx of peripheral white blood cells into the retinal tissue. In contrast, CD45$^+$ cells at

3-days and 7-days of reperfusion exhibited a combination of amoeboid and branched

morphologies (**Fig. 7C, D**), indicative of activated states.

Microglia, which are known to undergo activation and morphological changes in response

to various insults, were examined in the superficial retinas of sham and UPOAO eyes.

Immunofluorescence staining for Iba1 was conducted on flat-mounted retinas at 3-days

and 7-days post-reperfusion. In sham eyes, Iba1$^+$ microglia displayed small somas and

elongated dendrites that were evenly distributed within the retina (**Fig. 7H1, I1**). Conversely,

UPOAO retinas exhibited numerous activated Iba1$^+$ microglia with enlarged somas,

shortened dendrites, and an ameboid appearance (**Fig. 7H2, I2**). The number of Iba1$^+$

microglia in UPOAO retinas was approximately five times greater than that in sham eyes at

3-days (**Fig. 7J**), with a further increase observed at 7-days (**Fig. 7K**), although this increase

was less pronounced than that at the 3-days time point (**Fig. 7J**). These results indicated a

potential reduction in immune-mediated inflammation over time.

3.7 RNA-seq comparison between UPOAO and extravascular occlusion models

To further study the characteristics of UPOAO, we analyzed the transcriptomes of two

extravascular occlusion models: HIOP and UCCAO. The HIOP model induces ischemia

through anterior chamber perfusion with normal saline (**Fig. 8A**), while the UCCAO model

involves ligation of the unilateral common carotid artery (**Fig. 8H**). RNA-seq analysis of

retinas subjected to 60 minutes of ischemia followed by 7-days of reperfusion in the HIOP

model revealed enrichment of immune-related pathways, specifically highlighting increased

activity in leukocyte-mediated immunity and regulation of immune effector processes (**Fig.**

8B). This finding was consistent with the results of GO analysis, which revealed associations

of the overlapping DEGs between the HIOP and UPOAO models with toll-like receptor

binding, T-cell receptor binding, and immune-related functions (**Fig. 8C, D**). To further

explore these DEGs, we explored the remaining DEGs in the HIOP model in addition to the

overlapping genes. GO analysis revealed that the remaining DEGs in the HIOP model were

also enriched in immune responses, such as the T-cell receptor complex pathway, adaptive

immune response, and B-cell-mediated immunity function (**Fig. 8E**). Remarkably, upon

examining the remaining DEGs specific to the UPOAO model, GO analysis revealed DEGs

distinctly related to lipid and steroid metabolic processes (**Fig. 8F, G**). This observation

indicates that the UPOAO model, which is closely related to RAO, involves not only

conventional immune responses but also unique regulation of lipid and steroid metabolism.

Similarly, the UCCAO model exhibited associations with negative regulation of B-cell

proliferation, lymphocyte activation, and related processes (**Fig. 8I**). Additionally, the hub
genes identified through protein−protein interaction (PPI) analysis are presented (**Fig. 8J**).
However, the UCCAO model displayed minimal overlap with the UPOAO model, with only
15 overlapping DEGs identified in the Venn diagram analysis (**Fig. 8K**). Further analysis of
the UCCAO DEGs revealed two genes (Mettl14 and YTHDF2) related to m6A modifications
(**Fig. 8L**). Although arterial blood flow is occluded in both the UPOAO and UCCAO models,
our results revealed that the pathophysiological processes of the two models were
particularly different after 7-days of reperfusion.
537
4 Discussion
This study aimed to develop a more appropriate mouse model for investigating the
ischemia-reperfusion process in RAO and unravelling the underlying pathophysiological
mechanisms. We combined silicone wire embolization with carotid artery ligation to
effectively block the blood supply to the retinal artery, closely mimicking the characteristics
of the acute interruption of blood supply in RAO patients [31]. In the UPOAO model, a 60-
minute ischemia period was sufficient to simulate the injury of major retinal neural cells,
such as RGCs, BCs, HCs, and cholinergic amacrine cells, resulting in visual impairment.
Moreover, histologic examination demonstrated thinning of the inner layer of the retina,
especially the GCL. Time course transcriptome analysis revealed various pathophysiological
processes related to immune cell migration, oxidative stress, and immune inflammation
during non-reperfusion and reperfusion periods. The resident microglia within the retina
and peripheral leukocytes that access the retina were markedly increased during reperfusion
periods. Additionally, we compared the transcriptomic signatures of the UPOAO model with
those of two commonly used ischemia-reperfusion mouse models (HIOP and UCCAO),
highlighting the potential relevance of the UPOAO model to ocular vascular occlusive
diseases.
555
The UPOAO mouse model effectively simulates the characteristic features of RAO observed
in patients. RAO patients often exhibit distinctive patterns in dark-adapted ERGs, where b-
waves are reduced while a-waves remain stable [32-34]. In our study, we observed almost
the same phenomenon in ERG amplitude after 60 minutes of ischemia in the UPOAO model
(Table S2). This observation was further confirmed by immunofluorescence staining of
bipolar cells and photoreceptor cells. Importantly, a previous study linked the thinning of
the inner retina with alterations in ERG in patients with RAO [35]. Our results showed that
ischemic damage predominantly affected the inner retinal layers, while the outer layer was
almost unaffected (**Table S2**). This distinction is attributed to the reliance of the inner layers
on the retinal artery for blood supply, whereas the outer layers are supplied by the choroid.
Additionally, histological examination revealed increased retinal NFL and GCL thickness and
decreased IPL thickness after 3-days of reperfusion, indicating that ischemia-induced
oedema had not completely subsided.
569
This study comprehensively explored the transcriptomic signature after ischemia and
revealed that resident and peripheral immune cells may play major roles in pathological
processes. In the non-reperfusion group, the transcriptional changes were primarily

involved in the migration of immune cells such as leukocytes and neutrophils from blood
vessels into ischemic retinal tissue. This process is closely linked to injury during ischemia
and reperfusion, where leukocytes and neutrophils infiltrate neural tissue through the
vascular endothelium [36, 37]. Our results revealed an increase in CD45[+] leukocytes in the
retina and retinal microvasculature, suggesting that these leukocytes were recruited from
the bloodstream to the damaged retinal tissue following reperfusion. During leukocyte
access to the retina, chemokines, chemokine receptors, adhesion molecules and cytoskeletal
components play important roles in regulating endothelial permeability and facilitating
leukocyte adhesion to endothelial cells [38, 39]. Our findings of related gene expression are
consistent with this mechanism.
583
During the 3-days reperfusion period, we observed that an increased number of DEGs were
significantly involved in the critical pathological response to oxidative stress in the context of
IR tissues. The generation of reactive oxygen species (ROS) in mitochondria and subsequent
oxidative stress are widely recognized as major causes of retinal cell damage induced by IR
[40, 41]. This finding suggested that mitochondrial function, which is involved in oxidative
stress processes, may play an important role in the pathophysiology of IRI during the early
reperfusion stage. Oxidative stress can influence the permeability of the inner blood retina
barrier (iBRB), allowing leukocytes to access retinal tissues [42]. The infiltration and activation
of immune cells are recognized as the underlying causes of many retinal diseases, including
ischemic ophthalmia [42, 43]. In the UPOAO model, the increase in CD45[+] leukocytes and
Iba1[+] microglia in the retina after 3-days of reperfusion supports this viewpoint.
595
In the later stage of reperfusion (7-days), the DEGs were mainly enriched in immune
regulation and inflammation. With more upregulated DEGs than downregulated ones, we
hypothesize that the activation of the immune inflammatory response contributes to further
IRI in retinal tissue. Previous studies have shown that the retina can elicit immunological
responses during ischemia-reperfusion injury and that immune inflammation is an
important phenomenon in the progression of this injury [30]. The excessive ROS generated
by mitochondria in RGCs during the activation of inflammatory responses can damage cell
structure and visual function [44]. Microglia, considered the primary resident immune cells,
contribute to inflammatory responses and consequent neural damage [45]. Infiltrating
leukocytes can activate resident microglia, forming a feedback loop that exacerbates
inflammation [46]. Complex interactions between immune cells and retinal neurons after
RIRI have been reported [47]. Interestingly, in our UPOAO model, the death of major retinal
neural cells, such as RGCs, BCs, and HCs, was correlated with an increase in infiltrating
leukocytes and resident microglia. Therefore, we considered that the immune response and
neuroinflammation observed after 7-days of reperfusion may result from the cumulative
effects of acute oxidative stress and the responses of resident and peripheral immune cells
(**Fig. 9, Table S2**). In summary, we described the pathological processes at different time
points and highlighted the important role of resident and peripheral immune cell responses
in the UPOAO model. This finding offers valuable insights for subsequent screening of
pathogenic genes and potential immunotherapy approaches.
616

Compared with the HIOP and UCCAO models, the UPOAO model is a more appropriate
choice for studying retinal IRI in RAO. The HIOP model, widely used for investigating the
pathogenesis of APACG, exhibits retinal degeneration features similar to those of the
UPOAO model during both reperfusion periods [48]. In our UPOAO model, histologic and
immunohistochemical analysis demonstrated that 60 minutes of ischemia followed by 3-
days or 7-days of reperfusion can cause irreversible damage to retinal structure and visual
function. Specifically, we observed distinct alterations, such as thinning of the inner retina,
substantial RGCs apoptosis, and decreased b-wave amplitude. Multiple studies on the HIOP
model have reported consistent results [45, 49, 50]. However, some typical phenotypes in
the HIOP model, such as thinning of the outer nuclear layer and the entire neuroretina and
decreased a-wave amplitude, were not observed in the UPOAO model. Moreover, our
transcriptome sequencing revealed a specific subset of DEGs unique to the UPOAO model,
including Apoe, Abca1, Ldlr, Cyp39a1, and Bmp6. Notably, these DEGs were significantly
enriched in pathways associated with lipid and steroid synthesis. Steroid and lipid
metabolism homeostasis is disrupted under conditions of oxidative stress and inflammation
[51]. In the UPOAO model, the observed lipid and steroid biological processes may result
from the cumulative pathological responses triggered by IRI, as observed in the HIOP model.
In the UCCAO model, RNA-seq revealed that DEGs were enriched in two main pathways: 1)
epigenetic modification-related pathways, including nucleobase-containing compound
catabolic process, RNA catabolic process, mRNA catabolic process and N6-
methyladenosine (m6A) modification; and 2) cell death pathways, including regulation of
autophagosome assembly, negative regulation of neuron death, and negative regulation of
the neuronal apoptotic process. Epigenetic mechanisms may play a key role in the
pathophysiology of ocular disease [52]. m6A, one of the most common RNA modifications,
has been reported to regulate cell death processes, including apoptosis and autophagy, in
the pathological process of IRI [53]. These results suggest that epigenetic mechanisms may
significantly influence cell death during the 7-days reperfusion period in the UCCAO model.
Lee et al. evaluated the characteristics of UCCAO without reperfusion using visual,
histological and immunohistochemical approaches. Their results revealed delayed perfusion
of the ipsilateral retina, thinning of the inner retinal layer 10 weeks after surgery, and a
dramatic decrease in the amplitudes of b-waves on day 14 after UCCAO [15, 54]. This
finding suggested that UCCAO primarily represents a model of retinal hypoperfusion injury
and may not effectively reflect acute ischemic-induced structural and functional damage in
RAO (**Fig. 9**). These results led us to consider the UPOAO model an effective experimental
model for studying the pathological processes underlying acute ischemia and IRI.
652
Our research has certain limitations that should be acknowledged. First, our study focused
on the changes that occurred within 60 minutes of ischemia and within the first 7-days of
reperfusion in the UPOAO model. Further exploration is needed to understand the changes
induced by longer reperfusion periods. Additionally, we proposed possible pathological
mechanisms, including iBRB damage, oxidative stress and immune inflammation, mainly
based on the enrichment results of DEGs at three reperfusion time points. More molecular
experimental validation is required to substantiate these proposed mechanisms.
660

In summary, by thoroughly exploring the injury to major retinal neural cells, visual impairment and pathophysiological changes in the UPOAO model, we confirmed that this model can effectively simulate the acute ischemia-reperfusion processes in RAO. It serves as an ideal mouse model for investigating the underlying pathological mechanisms of ischemia and reperfusion. Furthermore, our UPOAO model holds great promise as a novel model for studies on pathogenic genes and potential therapeutic interventions for RAO.

Acknowledgements

The authors would like to express their gratitude to all participants involved in the experiments. The authors also wish to acknowledge the Eye Institute of Renmin Hospital of Wuhan University for providing us with the experimental facilities and environment. Special thanks are extended to Professor Shenqi Zhang for the technical support in the surgical procedure of UPOAO model.

Availability of data and materials

All data are available from the corresponding author upon reasonable request.

Authors' contributions

Xiao X and Yang AH conceived the project and supervised, guided the research. Li Y, Wang YD and Feng JQ wrote the draft, Feng JQ, Wang CS, Xie H and Li ZY revised the manuscript. Wang YD and Li Y constructed the animal models and obtained the data. Wang YD performed the immunofluorescence staining, HE staining. Li Y and Feng JQ performed the ERG, analyzed the RNA-seq data, and created the visualizations. Wan YW and Wang YD performed the FFA and OCT imaging and Wan YW analyzed the data. Lv BY and Wang YD took pictures and videos of the UPOAO model, and Feng JQ made schematic diagrams of modeling. Li YM, Xie H, Chen T, Wang FX provided support for the technical methods and contributed to the discussion, and reviewed the manuscript. All authors critically contributed to the manuscript, and have read and approved the final manuscript.

Funding

This work was supported by the National Nature Science Foundation of China (grant number: 82371079), National Key R&D Program of China (2023YFC2308404), Fundamental Research Funds for the Central Universities (2042023gf0013), Key research and development project of Hubei Province (grant number: 2022BCA009).

Ethics approval and consent to participate

All animal experiments were performed following the ethical guidelines outlined in the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research, with the approval of the Experimental Animal Ethics Committee of Renmin Hospital of Wuhan University (approval number: WDRM-20220305A).

References

1.   Scott, I.U., et al., *Retinal vascular occlusions.* Lancet, 2020. **396**(10266): p. 1927-1940.

2.   Hayreh, S.S. and M.B. Zimmerman, *Central retinal artery occlusion: visual outcome.* Am J
Ophthalmol, 2005. **140**(3): p. 376-91.

3.   Vestergaard, N., et al., *Risk of Stroke, Myocardial Infarction, and Death Among Patients*
With Retinal Artery Occlusion and the Effect of Antithrombotic Treatment. Transl Vis Sci
Technol, 2021. **10**(11): p. 2.

4.   Fallico, M., et al., *Risk of acute stroke in patients with retinal artery occlusion: a*
systematic review and meta-analysis. Eye (Lond), 2020. **34**(4): p. 683-689.

5.   Chen, T., et al., *Sex differences in major adverse cardiovascular and cerebrovascular*
event risk among central retinal artery occlusion patients. Sci Rep, 2023. **13**(1): p. 14930.

6.   Prasad, S.S., et al., *Retinal gene expression after central retinal artery ligation: effects of*
ischemia and reperfusion. Invest Ophthalmol Vis Sci, 2010. **51**(12): p. 6207-19.

7.   Mac Grory, B., et al., *Management of Central Retinal Artery Occlusion: A Scientific*
Statement From the American Heart Association. Stroke, 2021. **52**(6): p. e282-e294.

8.   Raber, F.P., et al., *Thrombolysis in central retinal artery occlusion: a retrospective*
observational study. J Neurol, 2023. **270**(2): p. 891-897.

9.   Hu, H., et al., *Efficacy of Intra-Arterial Thrombolysis for Acute Central Retinal Artery*
Occlusion: A Meta-Analysis. Eur Neurol, 2022. **85**(3): p. 186-194.

10.   Wan, P., et al., *LncRNA H19 initiates microglial pyroptosis and neuronal death in retinal*
ischemia/reperfusion injury. Cell Death Differ, 2020. **27**(1): p. 176-191.

11.   Shabanzadeh, A.P., et al., *Neurosurgical Modeling of Retinal Ischemia-Reperfusion Injury.*
J Stroke Cerebrovasc Dis, 2018. **27**(4): p. 845-856.

12.   Gao, Y., et al., *Novel Acute Retinal Artery Ischemia and Reperfusion Model in Nonhuman*
Primates. Stroke, 2020. **51**(8): p. 2568-2572.

13.   Morén, H., et al., *The porcine retinal vasculature accessed using an endovascular*
approach: a new experimental model for retinal ischemia. Invest Ophthalmol Vis Sci,
2009. **50**(11): p. 5504-10.

14.   Zhang, Y., et al., *Activation of the mitochondrial apoptotic pathway in a rat model of*
central retinal artery occlusion. Invest Ophthalmol Vis Sci, 2005. **46**(6): p. 2133-9.

15.   Lee, D., et al., *A mouse model of retinal hypoperfusion injury induced by unilateral*
common carotid artery occlusion. Exp Eye Res, 2020. **201**: p. 108275.

16.   Lee, D., et al., *Nicotinamide Mononucleotide Protects against Retinal Dysfunction in a*
Murine Model of Carotid Artery Occlusion. Int J Mol Sci, 2022. **23**(23).

17.   van Zyl, T., et al., *Cell atlas of aqueous humor outflow pathways in eyes of humans and*
four model species provides insight into glaucoma pathogenesis. Proc Natl Acad Sci U S
A, 2020. **117**(19): p. 10339-10349.

18.   Chiang, T., R.O. Messing, and W.H. Chou, *Mouse model of middle cerebral artery*
occlusion. J Vis Exp, 2011(48).

19.   Sasaki, M., O. Honmou, and J.D. Kocsis, *A rat middle cerebral artery occlusion model and*
intravenous cellular delivery. Methods Mol Biol, 2009. **549**: p. 187-95.

20.   Chen, Y.G., et al., *Increased nuclear factor-kappa B p65 immunoreactivity following*
retinal ischemia and reperfusion injury in mice. J Neurosci Res, 2003. **72**(1): p. 125-31.

21.   Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing*
and microarray studies. Nucleic Acids Res, 2015. **43**(7): p. e47.

22. Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.* Proc Natl Acad Sci U S A, 2005. **102**(43): p. 15545-50.

23. Szklarczyk, D., et al., *STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets.* Nucleic Acids Res, 2019. **47**(D1): p. D607-d613.

24. Kohl, M., S. Wiese, and B. Warscheid, *Cytoscape: software for visualization and analysis of biological networks.* Methods Mol Biol, 2011. **696**: p. 291-303.

25. Cao, W., et al., *Dextromethorphan attenuates the effects of ischemia on rabbit electroretinographic oscillatory potentials.* Documenta Ophthalmologica, 1993. **84**(3): p. 247-256.

26. Xu, J., et al., *Pregabalin Mediates Retinal Ganglion Cell Survival From Retinal Ischemia/Reperfusion Injury Via the Akt/GSK3β/β-Catenin Signaling Pathway.* Invest Ophthalmol Vis Sci, 2022. **63**(12): p. 7.

27. Takács, B., et al., *Electroretinographical Analysis of the Effect of BGP-15 in Eyedrops for Compensating Global Ischemia-Reperfusion in the Eyes of Sprague Dawley Rats.* Biomedicines, 2024. **12**(3).

28. Yang, Y., et al., *Tmem30a deficiency leads to retinal rod bipolar cell degeneration.* J Neurochem, 2019. **148**(3): p. 400-412.

29. Li, G., et al., *Diagnostic and Immunosuppressive Potential of Elevated Mir-424 Levels in Circulating Immune Cells of Ischemic Stroke Patients.* Aging Dis, 2018. **9**(2): p. 172-181.

30. Minhas, G., J. Sharma, and N. Khan, *Cellular Stress Response and Immune Signaling in Retinal Ischemia-Reperfusion Injury.* Front Immunol, 2016. **7**: p. 444.

31. Biousse, V., F. Nahab, and N.J. Newman, *Management of Acute Retinal Ischemia: Follow the Guidelines!* Ophthalmology, 2018. **125**(10): p. 1597-1607.

32. Block, F. and M. Schwarz, *The b-wave of the electroretinogram as an index of retinal ischemia.* Gen Pharmacol, 1998. **30**(3): p. 281-7.

33. Lima, L.H., et al., *Unilateral electronegative ERG in a presumed central retinal artery occlusion.* Clin Ophthalmol, 2010. **4**: p. 1311-4.

34. Suzuki, R., et al., *Degeneration and dysfunction of retinal neurons in acute ocular hypertensive rats: involvement of calpains.* J Ocul Pharmacol Ther, 2014. **30**(5): p. 419-28.

35. Shinoda, K., et al., *Changes in retinal thickness are correlated with alterations of electroretinogram in eyes with central retinal artery occlusion.* Graefes Arch Clin Exp Ophthalmol, 2008. **246**(7): p. 949-54.

36. Szabo, M.E., et al., *Ischemia and reperfusion-induced histologic changes in the rat retina. Demonstration of a free radical-mediated mechanism.* Invest Ophthalmol Vis Sci, 1991. **32**(5): p. 1471-8.

37. Szabo, M.E., et al., *Ischaemia- and reperfusion-induced Na+, K+, Ca2+ and Mg2+ shifts in rat retina: effects of two free radical scavengers, SOD and EGB 761.* Exp Eye Res, 1992. **55**(1): p. 39-45.

38. Johnson-Léger, C., M. Aurrand-Lions, and B.A. Imhof, *The parting of the endothelium: miracle, or simply a junctional affair?* J Cell Sci, 2000. **113 (Pt 6)**: p. 921-33.

39.   Springer, T.A., *Traffic signals for lymphocyte recirculation and leukocyte emigration: the*
792   *multistep paradigm.* Cell, 1994. **76**(2): p. 301-14.
793   40.   Fang, I.M., C.M. Yang, and C.H. Yang, *Chitosan oligosaccharides prevented retinal*
794   *ischemia and reperfusion injury via reduced oxidative stress and inflammation in rats.*
795   Exp Eye Res, 2015. **130**: p. 38-50.
796   41.   Oharazawa, H., et al., *Protection of the retina by rapid diffusion of hydrogen:*
797   *administration of hydrogen-loaded eye drops in retinal ischemia-reperfusion injury.*
798   Invest Ophthalmol Vis Sci, 2010. **51**(1): p. 487-92.
799   42.   Kaur, C., W.S. Foulds, and E.A. Ling, *Blood-retinal barrier in hypoxic ischaemic conditions:*
800   *basic concepts, clinical features and management.* Prog Retin Eye Res, 2008. **27**(6): p.
801   622-47.
802   43.   Eltzschig, H.K. and C.D. Collard, *Vascular ischaemia and reperfusion injury.* Br Med Bull,
803   2004. **70**: p. 71-86.
804   44.   Sanderson, T.H., et al., *Molecular mechanisms of ischemia-reperfusion injury in brain:*
805   *pivotal role of the mitochondrial membrane potential in reactive oxygen species*
806   *generation.* Mol Neurobiol, 2013. **47**(1): p. 9-23.
807   45.   Abcouwer, S.F., et al., *Inflammatory resolution and vascular barrier restoration after*
808   *retinal ischemia reperfusion injury.* J Neuroinflammation, 2021. **18**(1): p. 186.
809   46.   Au, N.P.B. and C.H.E. Ma, *Neuroinflammation, Microglia and Implications for Retinal*
810   *Ganglion Cell Survival and Axon Regeneration in Traumatic Optic Neuropathy.* Front
811   Immunol, 2022. **13**: p. 860070.
812   47.   Qin, Q., et al., *Inhibiting multiple forms of cell death optimizes ganglion cells survival*
813   *after retinal ischemia reperfusion injury.* Cell Death Dis, 2022. **13**(5): p. 507.
814   48.   Yao, F., et al., *Pathologically high intraocular pressure disturbs normal iron homeostasis*
815   *and leads to retinal ganglion cell ferroptosis in glaucoma.* Cell Death Differ, 2023. **30**(1):
816   p. 69-81.
817   49.   Zeng, Z., et al., *Pathologically high intraocular pressure induces mitochondrial*
818   *dysfunction through Drp1 and leads to retinal ganglion cell PANoptosis in glaucoma.*
819   Redox Biol, 2023. **62**: p. 102687.
820   50.   Yang, J., et al., *Overexpression of S100A4 protects retinal ganglion cells against retinal*
821   *ischemia-reperfusion injury in mice.* Exp Eye Res, 2020. **201**: p. 108281.
822   51.   Lam, T., et al., *Reversal of intramyocellular lipid accumulation by lipophagy and a p62-*
823   *mediated pathway.* Cell Death Discov, 2016. **2**: p. 16061.
824   52.   He, S., et al., *Review: Epigenetic mechanisms in ocular disease.* Mol Vis, 2013. **19**: p. 665-
825   74.
826   53.   Wang, J., Y. Li, and S. Zhang, *N(6)-methyladenosine modification: A vital role of*
827   *programmed cell death in myocardial ischemia/reperfusion injury.* Int J Cardiol, 2022.
828   **367**: p. 11-19.
829   54.   Lee, D., et al., *Retinal Degeneration in a Murine Model of Retinal Ischemia by Unilateral*
830   *Common Carotid Artery Occlusion.* Biomed Res Int, 2021. **2021**: p. 7727648.
831
832

833   Figures and Tables

834
Fig. 1 Modeling Procedure, Validation, and Cervical Artery Anatomy
(A1-G1) Schematic illustration of unilateral pterygopalatine ophthalmic artery occlusion
(UPOAO). (A2-G2) Practical operation of UPOAO. (A1, A2) Blunt separation and exposure of
the left cervical arteries. (B1, B2) Arterial suture ligation. (C1, C2) Insertion of the silicone wire
embolus. The artery was incised to create a hole, and the silicone wire embolus was inserted.
(D1, D2) Artery disconnection and movement of the silicone wire embolus. The artery was
cut along the incision, and the silicone wire embolus was retracted and reinserted. (E1, E2)
Removal of the silicone wire embolus and reperfusion. (F1, F2) Suture removal. The sutures
at both ends of the disconnected vessel were knotted, and the other two sutures were
removed. (G1, G2) Anatomic reduction and suturing of the skin. (H, I) Canavalin A label
vasculature of the UPOAO mouse retina. The silicone wire embolus was inserted into artery
before perfusing rhodamine-labeled canavalin A into the heart of UPOAO mouse. The sham
eye served as an unpracticed control eye, while the UPOAO lateral eye represented the
experimental eye. Retinal vessels in the sham eye (H) exhibited fluorescence filling, while
retinal vessels in the UPOAO lateral eye (I) remained unfilled. Scale bar = 1 mm. (J and K)
Fluorescein Fundus Angiography (FFA) performed before removing the silicone wire
embolus from the UPOAO mouse. The vessels in the sham lateral retina (J) were perfused,
while the lateral retinal perfusion in UPOAO (K) was delayed. (L) Schematic illustration of
cervical artery anatomy and ocular blood supply. Embolization of the PPA resulted in ocular
ischemia. The red arrow indicates the site of the silicone wire embolus occlusion. The

silicone wire embolus used a type 602156 wire, extended to 7mm with a diameter of
0.21mm. The blue arrows indicate the modelling locations of the HIOP model and the
UCCAO model, respectively. CCA: common carotid artery; ICA: internal carotid artery; ECA:
external carotid artery; PPA: pterygopalatine artery; MCA: middle cerebral artery; Infraorbital
art.: infraorbital artery; OA: ophthalmic artery; SPCA: short posterior ciliary artery; LPCA: long
posterior ciliary artery; CRA: central retinal artery.
861
862　**Table 1** Antibodies Used in Staining of Flat-Mounted Retina and Sections

Antibodies	Species	Dilution	Company
PKCα	rabbit	1:1000	Sigma
Recoverin	rabbit	1:1000	Millipore
Calbindin	rabbit	1:5000	SWANT
ChAT	goat	1:200	Millipore
Iba1	rabbit	1:500	Wako
CD45	rabbit	1:100	CST
Rhodamine-labeled canavalin A	/	1:125	MKBio
Goat anti-mouse IgG Alexa Fluor 488	goat	1:500	Jackson
Donkey anti-rabbit IgG Alexa Fluor 594	donkey	1:500	Jackson
Donkey anti-goat IgG Alexa Fluor 488	donkey	1:500	Jackson

Fig. 2 Staining and Quantification of Retinal Ganglion Cells (RGCs) at Different Ischemia and Reperfusion Times.

Flat-mounted retina RGCs were labeled with Brn3a staining. (A, B) Representative pictures of the peripheral retinal visual field (A), and quantification of surviving RGCs (B) in the 30-minute ischemia and 3-days reperfusion group. n = 3. (C, D) Representative pictures of the peripheral retinal visual field (C), and quantification of surviving RGCs (D) in the 30-minute ischemia and 7-days reperfusion group. n = 3. (E, F) Representative pictures of the peripheral retinal visual field (E), and quantification of surviving RGCs (F) in the 60-minute ischemia and 3-days reperfusion group. n = 5. (G, H) Representative pictures of the peripheral retinal visual field (G), and quantification of surviving RGCs (H) in the 60-minute ischemia and 7-days reperfusion group. n = 5. The results showed a significant loss of RGCs after 60 minutes of ischemia. Data were presented as means ± s.e.m, *: $p < 0.05$, **: $p < 0.01$,

***: p<0.001, ****: p<0.0001, t-test. Scale bar = 100 µm.

Fig. 3 Comparison of Electroretinographic (ERG) Dark-Adapted Responses at Different

Ischemia and Reperfusion Times.

Following the evaluation of surviving RGCs, visual function in sham and UPOAO

experimental eyes at various ischemia and reperfusion times was assessed using ERG. (A)

Representative waveforms in the four groups at the stimulus light intensities of 0.1, 1.0, and

10.0 cd.s/m2, respectively. (B) Quantification of a-wave and b-wave amplitudes in the 30-

minute ischemia and 3-days reperfusion group. n = 5. (C) Quantification of a-wave and b-

wave amplitudes in the 30-minute ischemia and 7-days reperfusion group. n = 5. (D)

Quantification of a-wave and b-wave amplitudes in the 60-minute ischemia and 3-days

reperfusion group. n = 7. (E) Quantification of a-wave and b-wave amplitudes in the 60-

minute ischemia and 7-days reperfusion group. n = 8. Dark-adapted responses showed

almost similar a-wave amplitudes but significantly decreased b-wave amplitudes in the 60-

minute ischemia groups. The amplitudes of b-waves declined at 3-days and even more

prominently at 7-days. (F, G) Representative OPs and quantification of amplitudes in the 60-

minute ischemic groups. n = 7 in the 3-days reperfusion group; n = 8 in the 7-days reperfusion group. The amplitudes of OPs decreased significantly at 7-days reperfusion. The decline in b-waves and OPs along with the loss of RGCs, supports the selection of a 60-minute ischemic duration as an appropriate choice. Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001, two-way ANOVA test for a-waves and b-waves; paired t-test for OPs.

Fig. 4 Changes in Retina Morphology in the UPOAO Model.
(A) Representative OCT images of the mouse retina at 3-days. The Green lines indicate the OCT scan area starting from the optic disc. Local magnification and layering are annotated. GCC: ganglion cell complex, including RNFL, GCL and IPL layers. (B and C) Quantification of GCC and total retinal thickness at 3-days. The thickness of the GCC and the entire retina in OCT was measured and compared at distances of 1.5 PD, 3.0 PD and 4.5 PD from the optic disc, respectively. n = 5. (D) Representative OCT images of the mouse retina at 7-days. (E and F) Quantification of GCC and total retinal thickness at 7-days. n = 5. (G) Representative HE images of the mouse retina at 3-days. (H, I and J) Quantification of NFL + GCL, IPL, and INL thickness at 3-days. Retinal thickness in HE was measured near the optic nerve head and compared. n = 3. (K) Representative HE images of the mouse retina at 7-days. (L, M and N) Quantification of NFL + GCL, IPL, and INL thickness at 7-days. n = 3. Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001, two-way ANOVA test in OCT and paired t-test in HE. PD: papillary diameters; RNFL: retinal nerve fiber layer; GCL: ganglion cell layer; IPL: inner plexiform layer; INL: inner nuclear layer; OPL: outer plexiform layer; ONL: outer nuclear layer; IS: inner segment; OS outer segment; RPE: retinal pigment epithelium. Scale bar = 50 μm.

919
Fig. 5 Changes in Bipolar Cells, Horizontal Cells, and Cholinergic Amacrine Cells in
UPOAO Mice.
(A) Representative images of mouse retina co-stained with DAPI and PKCα at 3-days (B)
Representative images of mouse retina co-stained with DAPI and PKCα at 7-days. (C)
Quantification of PKCα fluorescence density at 3-days. (D) Quantification of PKCα
fluorescence density at 7-days. n = 4. (E) Representative images of mouse retina co-stained
with DAPI and Calbindin at 3-days. (F) Representative images of mouse retina co-stained
with DAPI and Calbindin at 7-days. Horizontal cell somata are indicated by white asterisks.
(G) Quantification of Calbindin fluorescence density at 3-days. (H) Quantification of
Calbindin fluorescence density at 7-days. n = 3. (I) Representative images of mouse retina
co-stained with DAPI and ChAT at 3-days. (J) Representative images of mouse retina co-
stained with DAPI and ChAT at 7-days. Cholinergic amacrine cell somata are indicated by
white asterisks. (K) Quantification of ChAT fluorescence density at 3-days. (L) Quantification
of ChAT fluorescence density at 7-days. n = 3. Data were presented as means ± s.e.m, *:
p<0.05, **: p<0.01, ***: p<0.001,t- test. Scale bar = 50 µm.

Fig. 6 Transcriptomic Features at Different Times of Reperfusion.

RNA-seq evaluation was performed at 0-day, 3-days, and 7-days reperfusion periods in UPOAO revealing enrichment in pathways related to immune cells migration, oxidative stress, and immune inflammation. (A, D, G) Volcano plots display differential expression genes (DEGs) between UPOAO and sham eyes in the no-perfusion group (A), 3-days perfusion group (D), and 7-days perfusion group (G), respectively. Red dots: significant upregulated genes, green and blue dots: significant downregulated genes, grey dots: stable expressed genes, adjusted P < 0.05. log2FC = 1. (B, E, H) Gene ontology (GO) analysis of differential genes in the non-perfusion group (B), 3-days group (E), and 7-days group (H). The DEGs between UPOAO and sham eyes were enriched in pathways associated with immune cells migration (0d), oxidative stress (3d), and immune inflammation (7d), respectively. (C, F, I) Heatmap displaying the top 100 downregulated and top 100 upregulated DEGs between UPOAO and sham at non-perfusion, 3-days, and 7-days reperfusion, respectively. The box represents the genes related to immune cells migration

(C), oxidative stress (F), and immune inflammation (I). The ranking was determined by the magnitude of fold change. In each heatmap, the upper box represents the top 10 downregulated genes, while the lower box represents the top 10 upregulated genes.

Fig. 7 Peripheral Leukocyte Infiltration and Retinal Resident Microglial Activation. Rhodamine-labeled canavalin A was used for immediate cardiac perfusion to visualize blood vessels, followed by CD45 immunofluorescent staining to observe the relationship between blood vessels and CD45⁺ cells in sham (A), 1-day perfusion group (B), 3-days

perfusion group (C), and 7-days perfusion group (D). The presence of CD45⁺ cells within
blood vessels is indicated by white arrows. Scale bar = 50 µm. CD45⁺ cell counts were
performed in whole retinas of 1-day (E), 3-days (F), 7-days (G). The cellular morphology
and distribution of microglial cells in the superficial retina were assessed in 3-days (H) and
7-days (I). Activated microglial cells are indicated by white asterisks. Microglial cell counting
was conducted in the superficial retina of 3-days (J) and 7-days (K). Data points for CD45⁺
cells were derived from four flat-mounted retinas, and data points for microglial cells were
from five flat-mounted retinas. Data were presented as means ± s.e.m, *: $p<0.05$, **: $p<0.01$,
***: $p<0.001$, ****: $p<0.0001$, t- test. Scale bar = 50 µm.

Fig. 8 Transcriptomic Results Comparison between UPOAO, HIOP, and UCCAO Models.
RNA-seq comparison between UPOAO and extravascular occlusion models: HIOP and
UCCAO. (A) Schematic illustration of the HIOP model. (B) GO analysis of differential
expression genes(DEGs) in the HIOP model at 7-days perfusion. (C) Venn diagram
indicating the overlapping DEGs (146 genes) between HIOP and UPOAO models, as well as
the remaining DEGs specific to each group (593 genes in each group). (D) GO analysis of
the overlapping DEGs between HIOP and UPOAO models. (E) GO analysis of the remaining
DEGs in the HIOP model at 7-days perfusion, excluding the overlapping DEGs. (F) GO
analysis of the remaining DEGs, excluding the overlapping DEGs. (G) Heatmap showing the
inter-sample distribution of lipid and steroid-related DEGs from the analysis of the
remaining DEGs in the UPOAO model. (H) Schematic illustration of the UCCAO model. (I)
GO analysis of DEGs in the UCCAO model. (J) Hub genes identified through protein-protein
interaction (PPI) network analysis of the DEGs in the UCCAO model. (K) Venn diagram
indicating the overlapping DEGs between UCCAO and UPOAO models (15 genes). (L) Venn
diagram indicating the presence of two overlapping DEGs between UCCAO DEGs and m6A-
related genes.
986

988 **Fig. 9 The Characteristic Features of UPOAO Model during Ischemia-Reperfusion**
989 **Periods.**

Supplementary data

Fig.S1 Response Times of a-Waves and b-Waves in ERG at Different Light Intensities.

(A, B) Response times of a-waves and b-waves were measured under different light intensities at

3-days and 7-days post-UPOAO in the 30-minutes ischemia group. n = 5. (C, D) Response

times of a-waves and b-waves were measured under different light intensities at 3-days and 7-

days post-UPOAO in the 60-minute ischemia group. n = 7 at 3-days; n = 8 at 7-days. Data were

presented as means ± s.e.m, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$, two-way

ANOVA test.

1005

1006

Fig.S2 Quantification of INL+OPL Thickness and ONL+IS/OS+RPE Thickness in OCT during

3-days and 7-days Reperfusion of UPOAO Animals.

The INL+OPL thickness and the ONL+IS/OS+RPE thickness were measured and compared at

distances of 1.5 PD, 3.0 PD and 4.5 PD from the optic disc using OCT. (A and B) Quantification of

INL+OPL layer thickness (A) and ONL+IS/OS+RPE layer thickness (B) at 3-days. n = 5. No

significant differences were observed. (C and D) Quantification of INL+OPL layer thickness (C)

and ONL+IS/OS+RPE layer thickness (D) at 7-days. n = 5. No significant differences were

observed. Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****:

p<0.0001, two-way ANOVA test.

1016

1017

Fig.S3 Changes in Photoreceptor Cells in UPOAO.

(A) Representative images of mouse retina co-stained with DAPI and Recoverin at 3-days. (B) Representative images of mouse retina co-stained with DAPI and Recoverin at 7-days. n = 3. Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001, t-test. Scale bar = 50 μm.

1024

1025 **Fig.S4 Hub Genes in PPI Analysis and Gene Set Enrichment Analysis (GSEA) during the**

1026 **Non-Reperfusion Stage in UPOAO.**

1027 (A) Hub genes identified through PPI analysis of DEGs in the non-reperfusion group. (B-D)

1028 GSEA analysis of the pathway of the cell killing (B), lymphocyte mediated-immunity (C), and

1029 leukocyte mediated-immunity (D) in the non-reperfusion group.

Fig.S5 Hub Genes, GSEA Analysis at 3-days Reperfusion in UPOAO, and Relation with Mitochondrial Genes.

(A) Hub genes identified through PPI analysis of DEGs in the 3-days reperfusion group. (B) GSEA analysis of the pathways related to plasma membrane protein complex (top left), bounding membrane of organelle (top right), and cell surface (bottom). (C) Venn diagram showing the overlapping genes between mitochondrial genes and DEGs at 3-days reperfusion (44 genes). (D) GO analysis of the 44 overlapping DEGs.

1039

Fig.S6 Hub Genes, GSEA Analysis at 7-days Reperfusion in UPOAO, and Relation with Immune Genes.

(A) Hub genes identified through PPI analysis of DEGs in the 7-days reperfusion group. (B) GSEA analysis of the pathways related to the reactome innate immune system (top left), reactome immunoregulatory interactions between a lymphoid and a non-lymphoid cell (top right), reactome immune system (bottom left), and reactome adaptive immune system (bottom right). (C) Venn diagram showing the overlapping genes between immune genes and DEGs at 7-days reperfusion (112 genes). (D) GO analysis of the 112 overlapping DEGs.

1048

Fig.S7 Upregulation of Immune Inflammation-Related Gene Expression in the 7-days
Reperfusion Group.
The genes Abca1, Cd86, Cd48, Tlr4, Tlr6, and Tspo were significantly upregulated in the 7-
days reperfusion group, while other immune-inflammatory and chemotaxis-related genes
showed no significant differences. The data points were from the retina of four animals.
Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001,
Paired t-test.

1056

1057

1058 **Fig.S8 Co-Expressed Genes during 3-days and 7-days Reperfusion.**
1059 (A) Venn diagram showing the overlap of DEGs between the 3-days and 7-days reperfusion
1060 groups, along with a list of the 17 overlapping DEGs. (B) GO analysis of the 17 overlapping
1061 DEGs.

1062

1063

1064

1065

1066

1067

1068 **Table. S1 Primers Used in This Study.**

Gene Name		Sequence (5'->3')
Tlr3	Forward primer	CGATTCCCGACCAGCACTCT
	Reverse primer	CACATTTCCTTTCAGCCTGTGAT
Tlr4	Forward primer	TCCCTGCATAGAGGTAGTTCC
	Reverse primer	TCAAGGGGTTGAAGCTCAGA
Cd48	Forward primer	GGAACTGGATTTCAAGGTCATTCA
	Reverse primer	ATGGTCCAAGTGGGTCCTTAT
Ldlr	Forward primer	CCAATCGACTCACGGGTTCA
	Reverse primer	ACAGTGTCGACTTCTCTAGGC
Cd86	Forward primer	ATGGACCCCAGATGCACCA
	Reverse primer	TGTGCCCAAATAGTGCTCGT
Tspo	Forward primer	ACTGTATTCAGCCATGGGGTATG
	Reverse primer	CCAAGGGAACCATAGCGTCC
Abca1	Forward primer	GCAAGGCCGCACCATTATTT
	Reverse primer	TCTGAGAAACACTGTCCTCCTTTT
Apoe	Forward primer	CGGGTGCTCTGTTTTGGAGA
	Reverse primer	GGCTAGGCATCCTGTCAGC
S100a9	Forward primer	TGGCAACCTTTATGAAGAAAGAGA
	Reverse primer	GTGGGTTGTTCTCATGCAGC
β-actin	Forward primer	CACTGTCGAGTCGCGTCC
	Reverse primer	TCATCCATGGCGAACTGGTG

1069

Table. S2 Summary Table: Time Course of all the Morphological, Functional, Cellular, and Transcriptome Changes in the UPOAO Model.

	Changes in each group	30min-3d	30min-7d	60min-0d	60min-3d			60min-7d		
	RGCs counts	-	-	/	↓			↓		
ERG	Amplitudes of a-waves	-	-		-			- (0.001, 0.003, 0.01, 0.03, 0.1, 1.0 cd.s/m²) ↓ (0.3, 2.0, 10.0 cd.s/m²)		
	Response time of a-waves	-	-	/	-			-		
	Amplitudes of b-waves	-	-		↓			↓		
	Response time of b-waves	-	-		-			-		
	Amplitudes of OPs-waves		/		↓			↓		
					1.5 PD	3 PD	4.5 PD	1.5 PD	3 PD	4.5 PD
OCT	GCC thickness				-	-	-	↓	↓	↓
	INL+OPL thickness		/		-	-	-	-	-	-
	ONL+IS/OS+RPE thickness				-	-	-	-	-	-
	total retinal thickness				-	-	↓	↓	↓	↓
HE	RNFL+GCL thickness				↑			↓		
	IPL thickness				↓			↓		
	IINL thickness		/		-			↓		
	OPL thickness				-			-		
	ONL thickness				-			-		
	IS/OS+RPE thickness				-			-		
Retina neural cells	Bipolar cells (PKCα)				-			↓		
	Horizontal cells (Calbindin)				-			↓		
	Cholinergic amacrine cells (ChAT)		/		↓			↓		
	Photoreceptor cells (Recoverin)				-			-		
	Peripheral leukocyte infiltration (CD45)#		/		↑			↑		

Changes in each group		30min-3d	30min-7d	60min-0d	60min-3d	60min-7d
Microglia counts (Iba1)			/		↑	↑
RNA-seq	DEGs number		/	419 (up: 215; down: 204)	542 (up: 372; down: 170)	739 (up: 429; down: 310)
	Enrichment			Immune cells migration-related pathways	Oxidative stress-related pathways	Immune inflammation-related pathways

1072 -: no change;

1073 ↑ : increase;

1074 ↓ : decrease;

1075 /: not involved in the article;

1076 #: supplementary content: also increase after 60min-ischemia and 1d-reperfusion;

1077

Video 1. Blunt Dissection of Cervical Vessels. The neck was exposed, and the cervical

arteries were separated using blunt dissection techniques.

Video 2. Insertion of Silicone Wire Embolus and Induction of Ischemia. The silicone wire

embolus was inserted into the PPA, initiating the ischemic condition. PPA: the

pterygopalatine artery.

Video 3. Extraction of Silicone Wire Embolus and Reperfusion. The silicone wire embolus

was removed, allowing reperfusion to occur. The mouse's wound was sutured, and standard

feeding procedures were followed.

1086