Silicone Wire Embolization-induced Acute Retinal Artery Ischemia and Reperfusion

Model in Mouse: Gene Expression Provides Insight into Pathological Processes

Yuedan Wang[a#], Ying Li[a#], Jiaqing Feng[a#], Chuansen Wang[a], Yuwei Wan[a], Bingyang Lv[a],

Yinming Li[b], Hao Xie[b], Ting Chen[a], Faxi Wang[b], Ziyue Li[a], Anhuai Yang[a*] & Xuan Xiao[a,b*]

a Department of Ophthalmology, Renmin Hospital of Wuhan University, Wuhan 430060,

Hubei, China

b Department of Clinical Laboratory, Institute of Translational Medicine, Renmin Hospital

of Wuhan University, Wuhan 430060, Hubei, China

# These authors contributed equally.

* Correspondence to: Xuan Xiao, Department of Ophthalmology, Renmin Hospital of

Wuhan University, No. 238 Jie Fang Road, Wuhan 430060, Hubei, China. Tel: (86) 027-

88041911, Email: xiaoxuan1111@whu.edu.cn

* Correspondence to: Anhuai Yang, Department of Ophthalmology, Renmin Hospital of

Wuhan University, No. 238 Jie Fang Road, Wuhan 430060, Hubei, China. Email:

yah0525@126.com

Abstract

Acute retinal ischemia and ischemia-reperfusion injury are the primary causes of retinal

neural cell death and vision loss in retinal artery occlusion (RAO). The absence of an

accurate mouse model for simulating the retinal ischemic process has hindered progress

in developing neuroprotective agents for RAO. We developed a unilateral pterygopalatine

ophthalmic artery occlusion (UPOAO) mouse model using silicone wire embolization

combined with carotid artery ligation. The survival of retinal ganglion cells and visual

function were evaluated to determine the duration of ischemia. Immunofluorescence

staining, optical coherence tomography, and haematoxylin and eosin staining were

utilized to assess changes in major neural cell classes and retinal structure degeneration

at two reperfusion durations. Transcriptomics was employed to investigate alterations in

the pathological process of UPOAO following ischemia and reperfusion, highlighting

transcriptomic differences between UPOAO and other retinal ischemia-reperfusion

models. The UPOAO model successfully replicated the acute interruption of retinal blood

supply observed in RAO. 60-minutes of Ischemia led to significant loss of major retinal

neural cells and visual function impairment. Notable thinning of the inner retinal layer,

especially the ganglion cell layer, was evident post-UPOAO. Temporal transcriptome

analysis revealed various pathophysiological processes related to immune cell migration,

oxidative stress, and immune inflammation during the non-reperfusion and reperfusion

periods. A pronounced increase in microglia within the retina and peripheral leukocytes

accessing the retina was observed during reperfusion periods. Comparison of

differentially expressed genes (DEGs) between the UPOAO and high intraocular

pressure models revealed specific enrichments in lipid and steroid metabolism-related

genes in the UPOAO model. The UPOAO model emerges as a novel tool for screening

pathogenic genes and promoting further therapeutic research in RAO.

Key words: Retinal artery occlusion; Mouse model; Ischemia and reperfusion injury;

Transcriptomics; Immune cell

1 Introduction

Retinal artery occlusion (RAO) is a severe ophthalmic disease characterized by a sudden

interruption of blood flow in the retinal artery, leading to retinal ischemia [1]. Over 60% of

RAO patients suffer from impaired vision, ranging from finger counting to complete vision

loss [2]. Additionally, RAO patients face increased risk of cardiovascular and

cerebrovascular events [3-5]. Retinal ischemia and hypoxia can lead to irreversible

damage to retinal cells within 90 minutes [6]. This damage is primarily due to the

apoptosis of retinal ganglion cells (RGCs), which is driven by inflammation and oxidative

stress. Unfortunately, conservative treatments for RAO, such as ocular massage and

hyperbaric oxygen therapy, provide limited therapeutic benefits [7]. Although

thrombolysis is effective, its application is restricted by a narrow therapeutic window

(typically within 4.5 hours) [8]. Furthermore, the restoration of blood flow to the ischemic

area following thrombolysis can cause a sudden increase in tissue oxidative levels,

resulting in ischemia-reperfusion injury (IRI) in the retina [9]. IRI is a common

pathological condition that can induce inflammation, retinal tissue damage and visual

impairment [10]. Therefore, it is crucial to develop an animal model that accurately

simulates the pathological processes of RAO to thoroughly investigate the

pathophysiological changes and explore potential neuroprotective treatments following

retinal ischemia and reperfusion.

Various animal models have been employed to investigate the effects of retinal injury

resulting from ischemia and subsequent reperfusion in RAO [6, 11, 12]. Based on the

methods used for modelling, retinal ischemia-reperfusion models can be categorized into

two groups: 1) intravascular occlusion models, including photochemical-induced

thrombosis model and vascular intervention model; and 2) extravascular occlusion

models, including central retinal artery ligation (CRAL) model, unilateral common carotid

artery occlusion (UCCAO) model, and high intraocular pressure (HIOP) model. The

photochemical-induced thrombosis model and vascular intervention model have been

reported to be useful tools for assessing retinal injury after ischemia and reperfusion.

However, both models are limited by the need for trained interventional radiologists and

the need for advanced techniques [13, 14]. The CRAL model, which directly clips or

ligates the central retinal artery, also presents drawbacks, such as potential optic nerve

damage and difficulty in application to small animals such as mice, limiting its use in

experiments with larger sample sizes [6]. The UCCAO model, which induces retinal

hypoperfusion by occluding the unilateral common carotid artery (CCA), is more suitable

for simulating chronic ischemic retinal disease than acute retinal ischemia [15, 16]. The

HIOP model is widely utilized in mice for studying ischemia-reperfusion in acute primary

angle-closure glaucoma (APACG) [17]. However, the mechanical compression of the

retina caused by saline injection in this model may also lead to retinal damage,

potentially influencing IRI research. Therefore, it is critically important to develop a simple

and low-skill-required mouse model that can simulate acute retinal ischemia and

reperfusion injury in RAO patients.

To better simulate the retinal ischemic process and possible IRI following RAO, we

developed a novel vascular-associated mouse model called the unilateral pterygopalatine

ophthalmic artery occlusion (UPOAO) model. In this model, we employed silicone wire

embolization and carotid artery ligation to completely block the blood supply to the retina.

We characterized the major classes of retinal neural cells and evaluated visual function

following different durations of ischemia (30 minutes and 60 minutes) and reperfusion (3-

days and 7-days) after UPOAO. Additionally, we utilized transcriptomics to investigate

the transcriptional changes and elucidate the pathophysiological process of the UPOAO

model after ischemia and reperfusion. Finally, we highlighted the transcriptomic

differences between this model and other retinal ischemia-reperfusion models, including

HIOP and UCCAO, revealing the unique pathological processes that closely resemble

retinal IRI in RAO. The UPOAO model offers new insights into the mechanisms and

pathways involved in ischemia and reperfusion studies of RAO, providing a foundation for

studying protective strategies for ocular ischemic diseases.

2 Methods and materials

2.1 Animals

Eight-week-old male C57BL/6 mice were used in the experiments. Only male mice were

used to exclude the potential influence of oestrogenic hormones. The mice were provided

sufficient food and water and were maintained on a 12-hour dark-light cycle in a room

with regulated temperature conditions. All the experimental procedures were designed

and conducted according to the ethical guidelines outlined in the Association for

Research in Vision and Ophthalmology (ARVO) Statement for the Use of Animals in

Ophthalmic and Vision Research. The study protocol and methods received approval

from the Experimental Animal Ethics Committee of Renmin Hospital of Wuhan University

(approval number: WDRM-20220305A).

2.2 Preparation and surgical procedure of the UPOAO model

Eight-week-old male C57BL/6 mice weighing 20-25 grams were used in this study. An

isoflurane-based anaesthesia system was used to induce and maintain general

anaesthesia in the mice during surgical procedures. The mice were anaesthetized with a

1.5-2% concentration of isoflurane delivered in a mixture of nitrous oxide and oxygen via

rubber tubing. Body temperature was maintained at $37 \pm 0.5°C$ throughout the surgery.

Surgical instruments were sterilized with 75% ethanol before use to ensure sterility.

We drew inspiration from the widely employed middle cerebral artery occlusion (MCAO)

model [18, 19], commonly used in cerebral ischemic injury research, which guided the

development of the UPOAO model. The mouse was gently positioned in a supine posture

on a heating blanket, and its neck was exposed. The preparation involved depilation of

the neck area, followed by skin disinfection, and finally, a midline incision along the neck

was made. After separating the neck gland using two tweezers, careful blunt dissection

was employed to separate the left CCA, internal carotid artery (ICA) and external carotid

artery (ECA), avoiding the compression of nearby nerves and veins (**Fig. 1A**) (**Video 1**).

Then, the CCA and ICA were ligated with a 6-0 suture. A knot was secured using a 6-0

suture at the distal end of the ECA, and a slipknot was created at the proximal end (**Fig.**

1B). To preserve the reperfusion process and avoid bleeding and mortality during surgery,

a silicone wire embolus was inserted through the ECA instead of through the CCA.

Ophthalmic scissors were used to make a small inverted "V"-shaped incision between

the two suture knots on the ECA. A specialized silicone wire embolus, measuring 7 ± 0.1

mm in length and 0.21 ± 0.1 mm in diameter, was inserted through the incision, directing

it into the ECA and further into the CCA (**Fig. 1C**). Next, the ligation on the ICA was

removed, and the ECA was cut at the point where the artery had been previously incised

with scissors. Subsequently, the silicone wire embolus was retracted to the bifurcation of

the CCA, rotated counterclockwise, and inserted into the ICA and further into the

pterygopalatine artery (PPA) (**Fig. 1D1**). The silicone tail of the silicone wire embolus was

positioned near the bifurcation of the CCA (**Fig. 1D2**), effectively obstructing the

ophthalmic artery (OA) (**Video 2**). Then, the slipknot was fastened, and the skin was

sutured. The mouse was free to move around during arterial embolization.

Following a predetermined ischemia embolization time, the silicone wire embolus was

carefully removed from the PPA without massive haemorrhage (**Fig. 1E**). The CCA

suture was removed, restoring arterial reperfusion (**Fig. 1F**) (**Video 3**). Then, the skin

wound was sutured, and the mice were routinely fed during reperfusion (**Fig. 1G**).

2.3 Staining of retinal vessels

To clearly visualize the retinal vessels, rhodamine-labelled canavalin A was used for

mouse heart perfusion. Each mouse was anaesthetized with 1% sodium pentobarbital

and placed in the supine position on a foam board resting on a tray. The sternum was

opened to expose the heart. A needle attached to the perfusion tube was inserted into

the left ventricle via the apex of the mouse heart and secured. The right atrial appendage

was dissected with scissors to allow blood outflow. Saline solution was perfused through

the left ventricle using a pump system to drain blood from the arteries. Subsequently, the

prepared solution of rhodamine-labelled canavalin A was injected into the mice and

circulated throughout the mouse vessels for 30 minutes. The mice were then sacrificed,

and their eyes were removed and immersed in paraformaldehyde (PFA) for one hour in

the dark. A flat-mounted retina was obtained, and retinal images were captured using a

Leica SP8 confocal microscope equipped with a 10× objective.

2.4 Quantification of RGCs and microglia

The mice were euthanized by cervical dislocation, and their eyes were fixed in 4% PFA

for 60 minutes at room temperature (RT). Subsequently, the cornea and lens were

excised, and the intact retina was isolated for retinal flat mounting. The retina was

immersed in PBS supplemented with 5% bovine serum albumin (BSA) and 0.5% Triton

X-100 for overnight blocking and then incubated with Brn3a antibody (Synaptic Systems, Germany) or Iba1 antibody (Wako, Japan). Following a two-day incubation with the primary antibody, the retina was gently washed three times with PBS. Subsequently, the retina was incubated with Alexa Fluor 594 (AntGene, Wuhan, China) in a light-protected cassette for two days. After three rinses, the retina was uniformly sectioned into a four-leaf clover morphology and flattened using a coverslip. Retinal flat mounts labelled with Brn3a were photographed to count using a fluorescence microscope (BX51, Olympus, Japan) and representative pictures were imaged using a Leica SP8 confocal microscope (Leica TCS SP8, Germany). Retinas labelled with Iba1 were imaged utilizing a fluorescence microscope (BX63; Olympus, Tokyo, Japan). Each quadrant of the retina was systematically subdivided into central, middle, and peripheral fields (distance from the optic nerve head: central field: 0.1 mm to 0.5 mm, middle field: 0.9 mm to 1.3 mm, and peripheral field: 1.7 mm to 2.1 mm). The surviving RGCs and activated microglia of 12 fields in each retina were quantified and averaged using ImageJ software (National Institutes of Health, USA).

2.5 Electroretinogram (ERG)

Mice were dark-adapted overnight before ERG, and all subsequent procedures were carried out in the dark. Before ERG, the mice were anaesthetized with 1% sodium pentobarbital via intraperitoneal injection, and their pupils were dilated. A subcutaneous electrode was inserted into the posterior cervical skin, a tail electrode was affixed to the posterior end of the mouse tail, and the corneal electrode was gently placed on the

central corneal surface. A RetiMINER-C visual electrophysiological system (3VMED Co.,

Ltd., Shanghai, China) was used for recording electrical responses. The a-waves, b-

waves, and oscillatory potentials (OPs) after flash stimuli of 0.01, 0.03, 0.1, 0.3, 1.0, 3.0,

and 10.0 cd.s/m^2 in scotopic adaptation were recorded. The amplitudes and implicit times

of a-waves, b-waves, and OPs in response to various flash stimuli were analysed.

2.6 Optical Coherence Tomography Imaging (OCT) and Fluorescein Fundus

Angiography (FFA)

The mice were placed on the platform of a Spectralis HRA + OCT device (Heidelberg

Engineering, Heidelberg, Germany) for OCT imaging following anaesthesia. Pupil dilation

was performed, and normal saline was applied regularly to maintain corneal moisture.

The focal length of the device was adjusted until the mouse retina was clearly visible.

The head of the mouse was gently repositioned to capture images of the peripheral

fundus. The multiline mode was used to scan each layer of the retina, and four quadrants

of view centred on the nipple in the upper left, lower left, upper right, and lower right were

recorded. In this study, the total retinal thickness was manually segmented into three

parts and encompassed the entire thickness from the nerve fibre layer to the

photoreceptor layer. The ganglion cell complex (GCC) was defined as the combined

thickness of the retinal nerve fibre layer (RNFL), ganglion cell layer (GCL), and inner

plexiform layer (IPL). The inner nuclear layer (INL) and the outer plexiform layer (OPL)

were combined for thickness analysis due to the difficulty in distinguishing between these

layers. The remaining retinal layers, including the outer nuclear layer (ONL), inner

segment/outer segment (IS/OS), and retinal pigment epithelium (RPE) layers, were

measured together. The thickness of the retina at 1.5 papillary diameters (PD), 3.0 PD,

and 4.5 PD from the centre of the optic disc was measured using the Heidelberg

measuring tool.

For FFA, the mice were anaesthetized, and their pupils were dilated. Subsequently,

fundus angiography was performed immediately following an intraperitoneal injection of

sodium fluorescein. Images were acquired alternately for both eyes within 5 seconds of

the injection.

2.7 Hematoxylin and eosin (HE)

Mouse eyes were fixed in FAS eyeball fixation solution (Service-bio, Wuhan, China) for

48 hours, followed by dehydration and subsequent embedding in paraffin. The paraffin

blocks were trimmed parallel to the optic nerve to obtain 3-4 µm thick sections where the

optic nerve was located. Six paraffin sections from each eyeball were stained with HE.

Images were captured using an Olympus fluorescence microscope. The retinal thickness

within a range of 200-1100 µm from the optic disc was measured using Image-Pro Plus

6.0 software.

2.8 Staining of Retinal Sections

For immunofluorescence, mouse eyes were fixed in 4% PFA solution and subjected to

gradient dehydration using 10%, 20%, and 30% sucrose solutions. The following day, the

eyes were embedded in optimum cutting temperature (OCT) compound (SAKURA, USA)

and frozen at -80°C. Several 14-µm-thick frozen sections were obtained from each

eyeball through the use of a freezing microtome (Leica, Wetzlar, Germany). The section

surface was parallel to the optic nerve, and 3-6 frozen sections were mounted on a single

slide. The slides were stored at -20°C until use.

All sections were blocked with 5% BSA and 0.5% Triton X-100 in PBS for 2 hours.

Primary antibodies, as listed in **Table 1**, were diluted in PBS containing 5% BSA and 0.5%

Triton X-100 and incubated at 4°C overnight. The following day, after three rinses, the

sections were incubated with secondary antibodies at RT for 2 hours in a cassette,

followed by three washes. The sections were then stained with DAPI (Service-bio,

Wuhan, China) for 15 minutes. After the final three rinses, the frozen sections were

sealed. A Leica SP8 instrument equipped with a 40× objective was used to photograph

frozen sections near the optic nerve head. The fluorescence intensity of all slices of each

eye was determined using ImageJ software.

2.9 Surgical procedure for the UCCAO model and the HIOP model

Animals for the UCCAO model were prepared and managed according to the procedure

described for the UPOAO model. After blunt dissection, the left CCA was ligated with a 6-

0 suture. After 60 minutes of ligation, the suture was removed, and the skin wound was

subsequently closed with sutures. Mice were routinely fed during the reperfusion period,

and the retinas of UCCAO mice were harvested 7-days post-surgery.

The HIOP model was generated based on previously established methods [20]. Briefly,

mice were anaesthetized by intraperitoneal injection of 1% sodium pentobarbital. Pupils

were dilated using tropicamide phenylephrine eye drops (0.5% tropicamide and 0.5%

deoxyadrenaline hydrochloride, Santen Pharmaceutical) applied 5 minutes in advance.

An insulin hypodermic needle attached to a silicone elastomer tube was inserted into the

anterior chamber of the unilateral experimental eye. The IOP was elevated for 60

minutes by raising the height of the saline solution storage bag to 162 cm, after which the

needle was removed to immediately restore the IOP to baseline levels. Iris whitening and

loss of retinal red-light reflexes indicated retinal ischemia. After surgery, levofloxacin

hydrochloride ophthalmic gel was applied to the eyes of the mice. Retinas from HIOP

mice were collected 7-days post-surgery.

2.10 Transcriptome sequence and analysis

Retinal samples from UPOAO, HIOP, and UCCAO mice were extracted and promptly

frozen in liquid nitrogen within 10 minutes of cervical dislocation. Total RNA extraction

was performed according to the protocol outlined in the TRIzol Reagent manual (Life

Technologies, CA, USA). RNA integrity and concentration were assessed using an

Agilent 2,100 Bioanalyzer (Agilent Technologies, Inc., Santa Clara, CA, USA). The

resulting RNA samples were then pooled and subjected to sequencing on the Illumina

HiSeq3000 platform in a 150-bp paired-end read format. The raw RNA-sequencing

(RNA-seq) reads were preprocessed and quantified using the featureCounts function in

the SubReads package version 1.5.3, with default parameters.

RNA-seq analysis was conducted on samples from 5 mice in the non-perfusion group, 4

mice in the 3-days perfusion group, and 5 mice in the 7-days perfusion group of the

UPOAO model. Additionally, retinas from both the 7-days HIOP experimental eyes and

bilateral UCCAO eyes were collected (n=5) for analysis.

Data normalization and subsequent processing utilized the 'limma' package in R software

(version 4.2.3) [21]. Volcano plots and Venn diagrams were generated for visualization to

identify significantly differentially expressed genes (DEGs) and perform Gene Ontology

(GO) annotation analysis. DEGs with a |log2-fold change (FC)| > 1 and adjusted p value

< 0.05 were significantly differentially expressed. Enrichment analysis focused on DEGs

with an adjusted p < 0.05 and an enriched gene count > 5. To visualize the expression

patterns of significant DEGs, a heatmap was generated using the 'pheatmap' package.

Gene Set Enrichment Analysis (GSEA) was conducted using the GSEA program (version

4.3.2) [22], employing the default gene set m-subset (mh.all.v2023.1.mm.symbols.gmt) to

explore significant functional and pathway differences. Enriched pathways were classified

based on criteria such as adjusted p value (< 0.05), false discovery rate (FDR) q value (<

0.25), and normalized enrichment score (|NES| > 1). Furthermore, protein−protein

interaction (PPI) analysis was performed using the Search Tool for the Retrieval of

Interacting Genes (STRING) database (https://string-db.org/) [23]. Genes with a greater

degree of protein-level interactions with others were further analysed using Cytoscape

software (version 3.8.2) to generate a downstream PPI map. PPI pairings with an

interaction score > 0.7 were extracted and visualized using Cytoscape 3.9.0 [24].

2.11 Real-time Quantitative polymerase chain reaction (RT-qPCR)

RNA was extracted from fresh mouse retinal samples and subjected to reverse

transcription using HiScript III RT SuperMix for qPCR (Vazyme, China) following the

manufacturer's instructions. RT–qPCR analysis was conducted on a CFX instrument

(Bio-Rad) using AceQ qPCR SYBR Green Master Mix (Vazyme, China). The PCR

program comprised 40 cycles of 10 seconds at 95°C and 30 seconds at 60°C. Assays

were performed in triplicate, and Ct values were normalized to beta-actin levels. Relative

quantification of target gene expression was calculated using the 2-$\Delta\Delta$Ct method. The

primer sequences are provided in **Table S1**.

2.12 Statistical Analysis

All the data were analysed using GraphPad Prism version 9.0 (GraphPad Software, San

Diego, CA, USA). Each experimental group included a minimum of three biological

replicates. T tests, including RGCs and microglia counting, OPs analysis, retinal

thickness measurements via HE staining, and fluorescence intensity quantification, were

used to compare the two groups. Two-way ANOVA (two-way analysis of variance) test

was used for more than two different experimental groups, including ERG waves and

retinal thickness measurements in OCT. All data are presented as the mean ± standard

error of the mean (SEM), and the statistical graphs are shown as scatter bar graphs and

line charts. P < 0.05 was considered to indicate statistical significance.

3 Results

3.1 Silicone wire embolus insertion interrupts retinal blood flow

To reproduce the retinal ischemic process and potential IRI in RAO, we established a

mouse model of UPOAO by combining silicone wire embolization with carotid artery

ligation. Model validation was conducted through a series of experiments, with detailed

descriptions provided in the Materials and Methods section (**Fig. 1A-G**). To confirm blood

flow disruption, we performed cardiac perfusion with fluorescently labelled lectin and FFA.

Rhodamine-labelled canavalin A was used during in vivo perfusion with silicone wire

embolus, and both eyes were evaluated. The sham eye exhibited normal blood perfusion

in the retina, while the experimental eye showed no perfusion (**Fig. 1H, I**). FFA revealed

delayed and limited perfusion in the experimental lateral retina, primarily near the optic

disc (**Fig. 1J, K**). These findings indicate that the insertion of the silicone wire embolus

effectively impaired blood flow to the retina.

3.2 60-minute ischemia in UPOAO damage retinal structure and function

To investigate the optimal ischemic duration, the mice were subjected to ischemic

periods of 30 and 60 minutes, followed by reperfusion periods of 3-days and 7-days.

Retinal structure and visual function were assessed through flat-mounted retina analysis

and flash ERG.

In the 30-minute ischemia group, no significant RGCs death was observed after 3-days

(**Fig. 2A, B**) or 7-days of reperfusion (**Fig. 2C, D**). However, after 60 minutes of ischemia

followed by either 3-days or 7-days of reperfusion, reductions in RGCs density were

evident (**Fig. 2E-H**). ERG results showed no statistical difference in amplitudes between

bilateral eyes after 3- and 7-days of 30-minutes reperfusion (**Fig. 3A-C**). However, the b-

wave amplitude notably decreased after 60 minutes of ischemia and 3-days of

reperfusion (**Fig. 3D**). By 7-days of reperfusion, the b-wave amplitude had halved

compared to that of the sham eyes (**Fig. 3E**). The appearance times of the a-waves and

b-waves in the experimental and control eyes remained consistent across all four groups

(**Fig. S1**). Additionally, OPs have also been used to sensitively monitor retinal ischemic

effects by detecting changes before any alterations in b-waves occur [25-27]. The

amplitudes of OPs in the experimental eyes, particularly in the 60-minute ischemia and 7-

days reperfusion group, significantly decreased to less than 50% of those in the sham

eyes (**Fig. 3F, G**). Based on RGCs survival and changes in ERG waves, we determined

that a 60-minute ischemic duration is optimal for the UPOAO model.

3.3 Evaluation of retinal thickness in UPOAO through OCT and HE

To assess retinal thickness non-invasively in the UPOAO model, we conducted OCT

imaging during 3-days and 7-days reperfusion periods. Following a 3-days reperfusion

period, no significant changes were observed in the GCC, INL + OPL, and ONL + IS/OS

+ RPE layers at 1.5, 3, and 4.5 PD from the optic disc (**Fig. 4B, Fig. S2A, B**). However,

the total retinal thickness decreased at 4.5 PD (**Fig. 4C**). After 7-days of reperfusion, the

total retinal thickness decreased at 1.5 PD, 3 PD, and 4.5 PD, primarily due to inner

retinal thinning (**Fig. 4D-F, Fig. S2C, D**).

For a detailed analysis of inner retinal thickness, we extracted UPOAO mouse eyes and

stained them with HE. The thickness of the RNFL + GCL increased at 3-days of

reperfusion (**Fig. 4H**), followed by a decrease at 7-days of reperfusion (**Fig. 4L**). IPL

thickness decreased at both 3-days and 7-days of reperfusion (**Fig. 4I, M**), while INL

thickness decreased only at 7-days of reperfusion (**Fig. 4N**). We hypothesized that the

initial swelling of the GCL in response to acute ischemia and hypoxia led to early

thickening, followed by gradual thinning due to tissue dysfunction. This hypothesis was

supported by the observation of RGCs loss and reduced GCC thickness evaluated by

OCT at 7-days (**Fig. 4E**).

We observed by OCT that changes in retinal thickness became noticeable at 3-days

post-UPOAO and exhibited a significant decrease at 7-days. HE results further revealed

that alterations in inner retinal thickness constituted a substantial portion of the total

retinal thickness during the early reperfusion stage after UPOAO.

3.4 Survival of retinal neural cells in UPOAO

Given that mouse retinal function heavily relies on rod cells and that scotopic (low light)

vision is primarily governed by rod photoreceptors, the observed reductions in b-wave

and OPs-wave amplitudes indicate impaired synaptic transmission within the inner retina

[28]. Our OCT and HE findings suggested structural injuries within the inner retina.

Notably, the thickness of the outer retinal layers in OCT remained unchanged. To further

explore these observations, we conducted immunofluorescence staining to investigate

alterations in major retinal cell types, primarily focusing on bipolar cells (BCs),

photoreceptor cells, horizontal cells (HCs), and cholinergic amacrine cells.

3.4.1 BCs loss and photoreceptor cells survival

PKCα serves as a marker delineating BCs, with their cell bodies primarily located in the

outermost part of the INL, axonal terminals extending into the innermost part of the IPL,

and dendrites confined to the OPL (**Fig. 5A, B**). In the experimental UPOAO eyes, BCs

did not show significant changes at the 3-days reperfusion mark but exhibited dramatic

alterations by the 7-days reperfusion period (**Fig. 5A2, B2**). In particular, the

immunostaining density of BCs dendritic and axonal arbors notably decreased at the 7-

days reperfusion mark. BCs account for approximately 40% of INL cells in mice, and their

somata and axonal processes form a substantial part of the INL and IPL, consistent with

the thinning of the INL observed in both OCT and HE at 7-days (**Fig. 4E, N**). Notably, the

decrease in the number of PKCα$^+$ BCs at the 7-days reperfusion point coincided with the

decrease in the b-waves amplitude in ERG (**Fig. 3E**), indicating functional interplay

between BCs and other cells.

Recoverin marks the somata and outer segments of photoreceptor cells and is localized

within the ONL (**Fig. S3A, B**). Interestingly, recoverin-positive photoreceptors remained

relatively stable throughout the reperfusion periods, which aligns with the unchanged

thickness of the outer retinal layers observed in OCT (**Fig. S2C, D**).

3.4.2 HCs and cholinergic amacrine cells loss

We evaluated HCs and cholinergic amacrine cells using calbindin and ChAT

immunostaining, respectively. Calbindin immunostaining highlighted the somata of HCs

within the INL, with terminal axon connections linearly positioned within the OPL (**Fig. 5E,**

F). In the UPOAO experimental eyes, no significant change in the number of HCs was

observed during the 3-days reperfusion period, while a notable reduction was observed

after 7 days. By 7-days, a significant decrease in cell body numbers, a considerable

reduction in axon density, and disrupted linear connections were observed (**Fig. 5F2**).

ChAT$^+$ cell bodies were located in the GCL and INL, with dendrites forming two narrow

stratified bands within the IPL. The immunofluorescence density of ChAT$^+$ amacrine cells

decreased notably after 3 days and even more prominently after 7 days (**Fig. 5I, J**). The

fluorescence intensity of the two bands within the IPL markedly decreased and was

nearly invisible after 7 days (**Fig. 5J2**).

3.5 Time course transcriptome analysis revealed features of different reperfusion periods

in UPOAO

To explore the pathophysiological processes of UPOAO, we extracted retinas from both

eyes for transcriptome sequencing. The samples included retinas from eyes subjected to

60 minutes of ischemia without reperfusion, 60 minutes of ischemia followed by 3-days of

reperfusion, and 60 minutes of ischemia followed by 7-days of reperfusion, with the

sham-treated eyes serving as controls.

In the non-perfusion group, 215 genes were upregulated and 204 genes were

downregulated (**Fig. 6A**). GO enrichment analysis revealed that the DEGs were related to

leukocyte migration, epidermis development, myeloid leukocyte migration and other cell

migration pathways (**Fig. 6B**). The heatmap and box showed the upregulated and

downregulated genes involved in immune cell migration-related pathways during the non-

perfusion period in UPOAO (**Fig. 6C**). High-connectivity DEGs ('hub genes'), such as

Dusp1 and Fos, were also enriched in these pathways (**Fig. S4A**). GSEA also showed

similar results (**Fig. S4B**-D). These results suggested that in the early stage of retinal

ischemic injury, leukocytes from the microvasculature may infiltrate retinal tissue. More

experimental validation will be performed to confirm this hypothesis.

In the 3-days reperfusion group, 372 genes were upregulated, and 170 genes were

downregulated (**Fig. 6D**). GO enrichment analysis revealed that the DEGs at 3-days after

reperfusion were related to energy metabolism, mitochondrial regulation, and oxidative

stress pathways (**Fig. 6E**). The heatmap and box displayed the upregulated and

downregulated genes involved in oxidative stress-related pathways at the 3-days

reperfusion stage in UPOAO, including hub genes such as Mrpl18 and Mrps23 (**Fig. 6F**,

Fig. S5A). GSEA revealed changes in pathways related to the binding membrane of

organelles, the cell surface and the plasma membrane protein complex (**Fig. S5B**).

Furthermore, 44 overlapping genes between the mitochondrial genes and DEGs at 3-

days post-UPOAO were mainly related to mitochondrial transport and the mitochondrial

tricarboxylic acid cycle (**Fig. S5C, D**). These results suggested that mitochondria and

metabolism play significant roles in IRI.

In the 7-days reperfusion group, 429 genes were upregulated, and 310 genes were

downregulated (**Fig 6G**). The DEGs at this stage were enriched in pathways related to

immune effector processes, positive regulation of responses to external stimuli and

inflammation (**Fig. 6H**). A heatmap was generated to show the upregulated and

downregulated genes of immune inflammation-related pathways at 7-days after

reperfusion in UPOAO, and most hub genes, such as Cd86, Cd48, Tlr4, and Tlr6 were

also enriched in these pathways (**Fig. 6I, Fig. S6A**). GSEA also showed similar results

(**Fig S6B**). Moreover, RT–qPCR confirmed the upregulation of immune inflammation-

related gene expression (**Fig. S7**). Significantly, 112 genes that overlapped between

immune-related genes and DEGs at 7-days post-UPOAO were primarily associated with

immune receptor activity and phagocytic vesicles (**Fig. S6C, D**). These results

underscore the predominant role of immune responses during this stage.

To explore the associations between changes observed at 3-days and 7-days after

reperfusion, we analyzed the DEGs and identified 17 overlapping genes (**Fig. S8A**),

which were mainly associated with the nitric oxide biosynthetic process (**Fig. S8B**). Our

findings suggest that the immune inflammatory response observed after 7-days of

reperfusion may represent the cumulative effect of the acute oxidative stress response

during the initial 3-days reperfusion period.

3.6 Leukocyte Infiltration and Microglial Activation

Time course transcriptome analysis underscored the predominant role of immune

inflammatory responses during the reperfusion stage. Previous reports have indicated

that the inflammatory response in retinal ischemia-reperfusion injury (RIRI) is

orchestrated by peripheral immune cells and resident immune cells within the retina [29,

30]. In our study, we demonstrated the presence of leukocyte infiltration and microglial

activation in the UPOAO model. Immunofluorescence staining for CD45 was performed

on retinas at different time points post modelling to visualize the distribution and quantity

of white blood cells (**Fig. 7A-D**). Laser confocal Z-plane projections confirmed minimal

leukocyte infiltration in the retinas of sham eyes, which showed that CD45$^+$ cells in the

vascular lumen likely represented patrolling cells (**Fig. 7A**). A significant increase in

CD45$^+$ cells was observed in UPOAO model retinas after 1 day (**Fig. 7E**), with a

progressive increase in quantity over time (**Fig. 7F, G**). Notably, the majority of CD45$^+$

cells in UPOAO model retinas at 1-day displayed a morphology similar to that of vascular

leukocytes (**Fig. 7B**), suggesting a large influx of peripheral white blood cells into the

retinal tissue. In contrast, CD45$^+$ cells at 3-days and 7-days of reperfusion exhibited a

combination of amoeboid and branched morphologies (**Fig. 7C, D**), indicative of activated

states.

Microglia, which are known to undergo activation and morphological changes in response

to various insults, were examined in the superficial retinas of sham and UPOAO eyes.

Immunofluorescence staining for Iba1 was conducted on flat-mounted retinas at 3-days

and 7-days post-reperfusion. In sham eyes, Iba1$^+$ microglia displayed small somas and

elongated dendrites that were evenly distributed within the retina (**Fig. 7H1, I1**).

Conversely, UPOAO retinas exhibited numerous activated Iba1$^+$ microglia with enlarged

somas, shortened dendrites, and an ameboid appearance (**Fig. 7H2, I2**). The number of

Iba1$^+$ microglia in UPOAO retinas was approximately five times greater than that in sham

eyes at 3-days (**Fig. 7J**), with a further increase observed at 7-days (**Fig. 7K**), although

this increase was less pronounced than that at the 3-days time point (**Fig. 7J**). These

results indicated a potential reduction in immune-mediated inflammation over time.

3.7 RNA-seq comparison between UPOAO and extravascular occlusion models

To further study the characteristics of UPOAO, we analyzed the transcriptomes of two

extravascular occlusion models: HIOP and UCCAO. The HIOP model induces ischemia

through anterior chamber perfusion with normal saline (**Fig. 8A**), while the UCCAO model

involves ligation of the unilateral common carotid artery (**Fig. 8H**). RNA-seq analysis of

retinas subjected to 60 minutes of ischemia followed by 7-days of reperfusion in the

HIOP model revealed enrichment of immune-related pathways, specifically highlighting

increased activity in leukocyte-mediated immunity and regulation of immune effector

processes (**Fig. 8B**). This finding was consistent with the results of GO analysis, which

revealed associations of the overlapping DEGs between the HIOP and UPOAO models

with toll-like receptor binding, T-cell receptor binding, and immune-related functions (**Fig.**

8C, D). To further explore these DEGs, we explored the remaining DEGs in the HIOP

model in addition to the overlapping genes. GO analysis revealed that the remaining

DEGs in the HIOP model were also enriched in immune responses, such as the T-cell

receptor complex pathway, adaptive immune response, and B-cell-mediated immunity

function (**Fig. 8E**). Remarkably, upon examining the remaining DEGs specific to the

UPOAO model, GO analysis revealed DEGs distinctly related to lipid and steroid

metabolic processes (**Fig. 8F, G**). This observation indicates that the UPOAO model,

which is closely related to RAO, involves not only conventional immune responses but

also unique regulation of lipid and steroid metabolism.

Similarly, the UCCAO model exhibited associations with negative regulation of B-cell

proliferation, lymphocyte activation, and related processes (**Fig. 8I**). Additionally, the hub

genes identified through protein–protein interaction (PPI) analysis are presented (**Fig. 8J**).

However, the UCCAO model displayed minimal overlap with the UPOAO model, with

only 15 overlapping DEGs identified in the Venn diagram analysis (**Fig. 8K**). Further

analysis of the UCCAO DEGs revealed two genes (Mettl14 and YTHDF2) related to m6A

modifications (**Fig. 8L**). Although arterial blood flow is occluded in both the UPOAO and

UCCAO models, our results revealed that the pathophysiological processes of the two

models were particularly different after 7-days of reperfusion.

4 Discussion

This study aimed to develop a more appropriate mouse model for investigating the

ischemia-reperfusion process in RAO and unravelling the underlying pathophysiological

mechanisms. We combined silicone wire embolization with carotid artery ligation to

effectively block the blood supply to the retinal artery, closely mimicking the

characteristics of the acute interruption of blood supply in RAO patients [31]. In the

UPOAO model, a 60-minute ischemia period was sufficient to simulate the injury of major

retinal neural cells, such as RGCs, BCs, HCs, and cholinergic amacrine cells, resulting in

visual impairment. Moreover, histologic examination demonstrated thinning of the inner

layer of the retina, especially the GCL. Time course transcriptome analysis revealed

various pathophysiological processes related to immune cell migration, oxidative stress,

and immune inflammation during non-reperfusion and reperfusion periods. The resident

microglia within the retina and peripheral leukocytes that access the retina were markedly

increased during reperfusion periods. Additionally, we compared the transcriptomic

signatures of the UPOAO model with those of two commonly used ischemia-reperfusion

mouse models (HIOP and UCCAO), highlighting the potential relevance of the UPOAO

model to ocular vascular occlusive diseases.

The UPOAO mouse model effectively simulates the characteristic features of RAO

observed in patients. RAO patients often exhibit distinctive patterns in dark-adapted

ERGs, where b-waves are reduced while a-waves remain stable [32-34]. In our study, we

observed almost the same phenomenon in ERG amplitude after 60 minutes of ischemia

in the UPOAO model (Table S2). This observation was further confirmed by

immunofluorescence staining of bipolar cells and photoreceptor cells. Importantly, a

previous study linked the thinning of the inner retina with alterations in ERG in patients

with RAO [35]. Our results showed that ischemic damage predominantly affected the

inner retinal layers, while the outer layer was almost unaffected (**Table S2**). This

distinction is attributed to the reliance of the inner layers on the retinal artery for blood

supply, whereas the outer layers are supplied by the choroid. Additionally, histological

examination revealed increased retinal NFL and GCL thickness and decreased IPL

thickness after 3-days of reperfusion, indicating that ischemia-induced oedema had not

completely subsided.

This study comprehensively explored the transcriptomic signature after ischemia and

revealed that resident and peripheral immune cells may play major roles in pathological

processes. In the non-reperfusion group, the transcriptional changes were primarily

involved in the migration of immune cells such as leukocytes and neutrophils from blood

vessels into ischemic retinal tissue. This process is closely linked to injury during

ischemia and reperfusion, where leukocytes and neutrophils infiltrate neural tissue

through the vascular endothelium [36, 37]. Our results revealed an increase in CD45[+]

leukocytes in the retina and retinal microvasculature, suggesting that these leukocytes

were recruited from the bloodstream to the damaged retinal tissue following reperfusion.

During leukocyte access to the retina, chemokines, chemokine receptors, adhesion

molecules and cytoskeletal components play important roles in regulating endothelial

permeability and facilitating leukocyte adhesion to endothelial cells [38, 39]. Our findings

of related gene expression are consistent with this mechanism.

During the 3-days reperfusion period, we observed that an increased number of DEGs

were significantly involved in the critical pathological response to oxidative stress in the

context of IR tissues. The generation of reactive oxygen species (ROS) in mitochondria

and subsequent oxidative stress are widely recognized as major causes of retinal cell

damage induced by IR [40, 41]. This finding suggested that mitochondrial function, which

is involved in oxidative stress processes, may play an important role in the

pathophysiology of IRI during the early reperfusion stage. Oxidative stress can influence

the permeability of the inner blood retina barrier (iBRB), allowing leukocytes to access

retinal tissues [42]. The infiltration and activation of immune cells are recognized as the

underlying causes of many retinal diseases, including ischemic ophthalmia [42, 43]. In

the UPOAO model, the increase in CD45[+] leukocytes and Iba1[+] microglia in the retina

after 3-days of reperfusion supports this viewpoint.

In the later stage of reperfusion (7-days), the DEGs were mainly enriched in immune

regulation and inflammation. With more upregulated DEGs than downregulated ones, we

hypothesize that the activation of the immune inflammatory response contributes to

further IRI in retinal tissue. Previous studies have shown that the retina can elicit

immunological responses during ischemia-reperfusion injury and that immune

inflammation is an important phenomenon in the progression of this injury [30]. The

excessive ROS generated by mitochondria in RGCs during the activation of inflammatory

responses can damage cell structure and visual function [44]. Microglia, considered the

primary resident immune cells, contribute to inflammatory responses and consequent

neural damage [45]. Infiltrating leukocytes can activate resident microglia, forming a

feedback loop that exacerbates inflammation [46]. Complex interactions between

immune cells and retinal neurons after RIRI have been reported [47]. Interestingly, in our

UPOAO model, the death of major retinal neural cells, such as RGCs, BCs, and HCs,

was correlated with an increase in infiltrating leukocytes and resident microglia.

Therefore, we considered that the immune response and neuroinflammation observed

after 7-days of reperfusion may result from the cumulative effects of acute oxidative

stress and the responses of resident and peripheral immune cells (**Fig. 9, Table S2**). In

summary, we described the pathological processes at different time points and

highlighted the important role of resident and peripheral immune cell responses in the

UPOAO model. This finding offers valuable insights for subsequent screening of

pathogenic genes and potential immunotherapy approaches.

Compared with the HIOP and UCCAO models, the UPOAO model is a more appropriate

choice for studying retinal IRI in RAO. The HIOP model, widely used for investigating the

pathogenesis of APACG, exhibits retinal degeneration features similar to those of the

UPOAO model during both reperfusion periods [48]. In our UPOAO model, histologic and

immunohistochemical analysis demonstrated that 60 minutes of ischemia followed by 3-

days or 7-days of reperfusion can cause irreversible damage to retinal structure and

visual function. Specifically, we observed distinct alterations, such as thinning of the inner

retina, substantial RGCs apoptosis, and decreased b-wave amplitude. Multiple studies on

the HIOP model have reported consistent results [45, 49, 50]. However, some typical

phenotypes in the HIOP model, such as thinning of the outer nuclear layer and the entire

neuroretina and decreased a-wave amplitude, were not observed in the UPOAO model.

Moreover, our transcriptome sequencing revealed a specific subset of DEGs unique to

the UPOAO model, including Apoe, Abca1, Ldlr, Cyp39a1, and Bmp6. Notably, these

DEGs were significantly enriched in pathways associated with lipid and steroid synthesis.

Steroid and lipid metabolism homeostasis is disrupted under conditions of oxidative

stress and inflammation [51]. In the UPOAO model, the observed lipid and steroid

biological processes may result from the cumulative pathological responses triggered by

IRI, as observed in the HIOP model. In the UCCAO model, RNA-seq revealed that DEGs

were enriched in two main pathways: 1) epigenetic modification-related pathways,

including nucleobase-containing compound catabolic process, RNA catabolic process,

mRNA catabolic process and N6-methyladenosine (m6A) modification; and 2) cell death

pathways, including regulation of autophagosome assembly, negative regulation of

neuron death, and negative regulation of the neuronal apoptotic process. Epigenetic

mechanisms may play a key role in the pathophysiology of ocular disease [52]. m6A, one

of the most common RNA modifications, has been reported to regulate cell death

processes, including apoptosis and autophagy, in the pathological process of IRI [53].

These results suggest that epigenetic mechanisms may significantly influence cell death

during the 7-days reperfusion period in the UCCAO model. Lee et al. evaluated the

characteristics of UCCAO without reperfusion using visual, histological and

immunohistochemical approaches. Their results revealed delayed perfusion of the

ipsilateral retina, thinning of the inner retinal layer 10 weeks after surgery, and a dramatic

decrease in the amplitudes of b-waves on day 14 after UCCAO [15, 54]. This finding

suggested that UCCAO primarily represents a model of retinal hypoperfusion injury and

may not effectively reflect acute ischemic-induced structural and functional damage in

RAO (**Fig. 9**). These results led us to consider the UPOAO model an effective

experimental model for studying the pathological processes underlying acute ischemia

and IRI.

Our research has certain limitations that should be acknowledged. First, our study

focused on the changes that occurred within 60 minutes of ischemia and within the first

7-days of reperfusion in the UPOAO model. Further exploration is needed to understand

the changes induced by longer reperfusion periods. Additionally, we proposed possible

pathological mechanisms, including iBRB damage, oxidative stress and immune

inflammation, mainly based on the enrichment results of DEGs at three reperfusion time

points. More molecular experimental validation is required to substantiate these proposed

mechanisms.

In summary, by thoroughly exploring the injury to major retinal neural cells, visual

impairment and pathophysiological changes in the UPOAO model, we confirmed that this

model can effectively simulate the acute ischemia-reperfusion processes in RAO. It

serves as an ideal mouse model for investigating the underlying pathological

mechanisms of ischemia and reperfusion. Furthermore, our UPOAO model holds great

promise as a novel model for studies on pathogenic genes and potential therapeutic

interventions for RAO.

Acknowledgements

The authors would like to express their gratitude to all participants involved in the

experiments. The authors also wish to acknowledge the Eye Institute of Renmin Hospital

of Wuhan University for providing us with the experimental facilities and environment.

Special thanks are extended to Professor Shenqi Zhang for the technical support in the

surgical procedure of UPOAO model.

Availability of data and materials

All data are available from the corresponding author upon reasonable request.

Authors' contributions

Xiao X and Yang AH conceived the project and supervised, guided the research. Li Y,

Wang YD and Feng JQ wrote the draft, Feng JQ, Wang CS, Xie H and Li ZY revised the

manuscript. Wang YD and Li Y constructed the animal models and obtained the data.

Wang YD performed the immunofluorescence staining, HE staining. Li Y and Feng JQ

performed the ERG, analyzed the RNA-seq data, and created the visualizations. Wan

YW and Wang YD performed the FFA and OCT imaging and Wan YW analyzed the data.

Lv BY and Wang YD took pictures and videos of the UPOAO model, and Feng JQ made

schematic diagrams of modeling. Li YM, Xie H, Chen T, Wang FX provided support for

the technical methods and contributed to the discussion, and reviewed the manuscript.

All authors critically contributed to the manuscript, and have read and approved the final

manuscript.

Funding

This work was supported by the National Nature Science Foundation of China (grant

number: 82371079), National Key R&D Program of China (2023YFC2308404),

Fundamental Research Funds for the Central Universities (2042023gf0013), Key

research and development project of Hubei Province (grant number: 2022BCA009).

Ethics approval and consent to participate

All animal experiments were performed following the ethical guidelines outlined in the

ARVO Statement for the Use of Animals in Ophthalmic and Vision Research, with the

approval of the Experimental Animal Ethics Committee of Renmin Hospital of Wuhan

University (approval number: WDRM-20220305A).

References

1.   Scott, I.U., et al., *Retinal vascular occlusions.* Lancet, 2020. **396**(10266): p. 1927-

1940.

2.   Hayreh, S.S. and M.B. Zimmerman, *Central retinal artery occlusion: visual outcome.*

Am J Ophthalmol, 2005. **140**(3): p. 376-91.

3.   Vestergaard, N., et al., *Risk of Stroke, Myocardial Infarction, and Death Among*

Patients With Retinal Artery Occlusion and the Effect of Antithrombotic Treatment.

Transl Vis Sci Technol, 2021. **10**(11): p. 2.

4.   Fallico, M., et al., *Risk of acute stroke in patients with retinal artery occlusion: a*

systematic review and meta-analysis. Eye (Lond), 2020. **34**(4): p. 683-689.

5.   Chen, T., et al., *Sex differences in major adverse cardiovascular and cerebrovascular*

event risk among central retinal artery occlusion patients. Sci Rep, 2023. **13**(1): p.

14930.

6.   Prasad, S.S., et al., *Retinal gene expression after central retinal artery ligation:*

effects of ischemia and reperfusion. Invest Ophthalmol Vis Sci, 2010. **51**(12): p. 6207-

19.

7.   Mac Grory, B., et al., *Management of Central Retinal Artery Occlusion: A Scientific*

Statement From the American Heart Association. Stroke, 2021. **52**(6): p. e282-e294.

8.   Raber, F.P., et al., *Thrombolysis in central retinal artery occlusion: a retrospective*

observational study. J Neurol, 2023. **270**(2): p. 891-897.

9.   Hu, H., et al., *Efficacy of Intra-Arterial Thrombolysis for Acute Central Retinal Artery*

Occlusion: A Meta-Analysis. Eur Neurol, 2022. **85**(3): p. 186-194.

10.   Wan, P., et al., *LncRNA H19 initiates microglial pyroptosis and neuronal death in*

retinal ischemia/reperfusion injury. Cell Death Differ, 2020. **27**(1): p. 176-191.

11.   Shabanzadeh, A.P., et al., *Neurosurgical Modeling of Retinal Ischemia-Reperfusion*

Injury. J Stroke Cerebrovasc Dis, 2018. **27**(4): p. 845-856.

12.   Gao, Y., et al., *Novel Acute Retinal Artery Ischemia and Reperfusion Model in*

Nonhuman Primates. Stroke, 2020. **51**(8): p. 2568-2572.

13.   Morén, H., et al., *The porcine retinal vasculature accessed using an endovascular*

approach: a new experimental model for retinal ischemia. Invest Ophthalmol Vis Sci,

2009. **50**(11): p. 5504-10.

14.   Zhang, Y., et al., *Activation of the mitochondrial apoptotic pathway in a rat model of*

central retinal artery occlusion. Invest Ophthalmol Vis Sci, 2005. **46**(6): p. 2133-9.

15.   Lee, D., et al., *A mouse model of retinal hypoperfusion injury induced by unilateral*

common carotid artery occlusion. Exp Eye Res, 2020. **201**: p. 108275.

16.   Lee, D., et al., *Nicotinamide Mononucleotide Protects against Retinal Dysfunction in a*

Murine Model of Carotid Artery Occlusion. Int J Mol Sci, 2022. **23**(23).

17.   van Zyl, T., et al., *Cell atlas of aqueous humor outflow pathways in eyes of humans*

and four model species provides insight into glaucoma pathogenesis. Proc Natl Acad

Sci U S A, 2020. **117**(19): p. 10339-10349.

18.   Chiang, T., R.O. Messing, and W.H. Chou, *Mouse model of middle cerebral artery*

occlusion. J Vis Exp, 2011(48).

19.   Sasaki, M., O. Honmou, and J.D. Kocsis, *A rat middle cerebral artery occlusion model*

and intravenous cellular delivery. Methods Mol Biol, 2009. **549**: p. 187-95.

20.   Chen, Y.G., et al., *Increased nuclear factor-kappa B p65 immunoreactivity following*
retinal ischemia and reperfusion injury in mice. J Neurosci Res, 2003. **72**(1): p. 125-
31.

21.   Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-*
sequencing and microarray studies. Nucleic Acids Res, 2015. **43**(7): p. e47.

22.   Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach*
for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005.
102(43): p. 15545-50.

23.   Szklarczyk, D., et al., *STRING v11: protein-protein association networks with*
increased coverage, supporting functional discovery in genome-wide experimental
datasets. Nucleic Acids Res, 2019. **47**(D1): p. D607-d613.

24.   Kohl, M., S. Wiese, and B. Warscheid, *Cytoscape: software for visualization and*
analysis of biological networks. Methods Mol Biol, 2011. **696**: p. 291-303.

25.   Cao, W., et al., *Dextromethorphan attenuates the effects of ischemia on rabbit*
electroretinographic oscillatory potentials. Documenta Ophthalmologica, 1993. **84**(3):
p. 247-256.

26.   Xu, J., et al., *Pregabalin Mediates Retinal Ganglion Cell Survival From Retinal*
Ischemia/Reperfusion Injury Via the Akt/GSK3β/β-Catenin Signaling Pathway. Invest
Ophthalmol Vis Sci, 2022. **63**(12): p. 7.

27.   Takács, B., et al., *Electroretinographical Analysis of the Effect of BGP-15 in Eyedrops*
for Compensating Global Ischemia-Reperfusion in the Eyes of Sprague Dawley Rats.
Biomedicines, 2024. **12**(3).

28. Yang, Y., et al., *Tmem30a deficiency leads to retinal rod bipolar cell degeneration.* J

Neurochem, 2019. **148**(3): p. 400-412.

29. Li, G., et al., *Diagnostic and Immunosuppressive Potential of Elevated Mir-424 Levels*

in Circulating Immune Cells of Ischemic Stroke Patients. Aging Dis, 2018. **9**(2): p.

172-181.

30. Minhas, G., J. Sharma, and N. Khan, *Cellular Stress Response and Immune*

Signaling in Retinal Ischemia-Reperfusion Injury. Front Immunol, 2016. **7**: p. 444.

31. Biousse, V., F. Nahab, and N.J. Newman, *Management of Acute Retinal Ischemia:*

Follow the Guidelines! Ophthalmology, 2018. **125**(10): p. 1597-1607.

32. Block, F. and M. Schwarz, *The b-wave of the electroretinogram as an index of retinal*

ischemia. Gen Pharmacol, 1998. **30**(3): p. 281-7.

33. Lima, L.H., et al., *Unilateral electronegative ERG in a presumed central retinal artery*

occlusion. Clin Ophthalmol, 2010. **4**: p. 1311-4.

34. Suzuki, R., et al., *Degeneration and dysfunction of retinal neurons in acute ocular*

hypertensive rats: involvement of calpains. J Ocul Pharmacol Ther, 2014. **30**(5): p.

419-28.

35. Shinoda, K., et al., *Changes in retinal thickness are correlated with alterations of*

electroretinogram in eyes with central retinal artery occlusion. Graefes Arch Clin Exp

Ophthalmol, 2008. **246**(7): p. 949-54.

36. Szabo, M.E., et al., *Ischemia and reperfusion-induced histologic changes in the rat*

retina. Demonstration of a free radical-mediated mechanism. Invest Ophthalmol Vis

Sci, 1991. **32**(5): p. 1471-8.

37.   Szabo, M.E., et al., *Ischaemia- and reperfusion-induced Na+, K+, Ca2+ and Mg2+*

shifts in rat retina: effects of two free radical scavengers, SOD and EGB 761. Exp

Eye Res, 1992. **55**(1): p. 39-45.

38.   Johnson-Léger, C., M. Aurrand-Lions, and B.A. Imhof, *The parting of the endothelium:*

miracle, or simply a junctional affair? J Cell Sci, 2000. **113 (Pt 6)**: p. 921-33.

39.   Springer, T.A., *Traffic signals for lymphocyte recirculation and leukocyte emigration:*

the multistep paradigm. Cell, 1994. **76**(2): p. 301-14.

40.   Fang, I.M., C.M. Yang, and C.H. Yang, *Chitosan oligosaccharides prevented retinal*

ischemia and reperfusion injury via reduced oxidative stress and inflammation in rats.

Exp Eye Res, 2015. **130**: p. 38-50.

41.   Oharazawa, H., et al., *Protection of the retina by rapid diffusion of hydrogen:*

administration of hydrogen-loaded eye drops in retinal ischemia-reperfusion injury.

Invest Ophthalmol Vis Sci, 2010. **51**(1): p. 487-92.

42.   Kaur, C., W.S. Foulds, and E.A. Ling, *Blood-retinal barrier in hypoxic ischaemic*

conditions: basic concepts, clinical features and management. Prog Retin Eye Res,

2008. **27**(6): p. 622-47.

43.   Eltzschig, H.K. and C.D. Collard, *Vascular ischaemia and reperfusion injury.* Br Med

Bull, 2004. **70**: p. 71-86.

44.   Sanderson, T.H., et al., *Molecular mechanisms of ischemia-reperfusion injury in brain:*

pivotal role of the mitochondrial membrane potential in reactive oxygen species

generation. Mol Neurobiol, 2013. **47**(1): p. 9-23.

45. Abcouwer, S.F., et al., *Inflammatory resolution and vascular barrier restoration after*

retinal ischemia reperfusion injury. J Neuroinflammation, 2021. **18**(1): p. 186.

46. Au, N.P.B. and C.H.E. Ma, *Neuroinflammation, Microglia and Implications for Retinal*

Ganglion Cell Survival and Axon Regeneration in Traumatic Optic Neuropathy. Front

Immunol, 2022. **13**: p. 860070.

47. Qin, Q., et al., *Inhibiting multiple forms of cell death optimizes ganglion cells survival*

after retinal ischemia reperfusion injury. Cell Death Dis, 2022. **13**(5): p. 507.

48. Yao, F., et al., *Pathologically high intraocular pressure disturbs normal iron*

homeostasis and leads to retinal ganglion cell ferroptosis in glaucoma. Cell Death

Differ, 2023. **30**(1): p. 69-81.

49. Zeng, Z., et al., *Pathologically high intraocular pressure induces mitochondrial*

dysfunction through Drp1 and leads to retinal ganglion cell PANoptosis in glaucoma.

Redox Biol, 2023. **62**: p. 102687.

50. Yang, J., et al., *Overexpression of S100A4 protects retinal ganglion cells against*

retinal ischemia-reperfusion injury in mice. Exp Eye Res, 2020. **201**: p. 108281.

51. Lam, T., et al., *Reversal of intramyocellular lipid accumulation by lipophagy and a*

p62-mediated pathway. Cell Death Discov, 2016. **2**: p. 16061.

52. He, S., et al., *Review: Epigenetic mechanisms in ocular disease.* Mol Vis, 2013. **19**: p.

665-74.

53. Wang, J., Y. Li, and S. Zhang, *N(6)-methyladenosine modification: A vital role of*

programmed cell death in myocardial ischemia/reperfusion injury. Int J Cardiol, 2022.

367: p. 11-19.

54.   Lee, D., et al., *Retinal Degeneration in a Murine Model of Retinal Ischemia by*

Unilateral Common Carotid Artery Occlusion. Biomed Res Int, 2021. **2021**: p.

7727648.

Figures and Tables

Fig. 1 Modeling Procedure, Validation, and Cervical Artery Anatomy

(A1-G1) Schematic illustration of unilateral pterygopalatine ophthalmic artery occlusion

(UPOAO). (A2-G2) Practical operation of UPOAO. (A1, A2) Blunt separation and

exposure of the left cervical arteries. (B1, B2) Arterial suture ligation. (C1, C2) Insertion of

the silicone wire embolus. The artery was incised to create a hole, and the silicone wire

embolus was inserted. (D1, D2) Artery disconnection and movement of the silicone wire

embolus. The artery was cut along the incision, and the silicone wire embolus was

retracted and reinserted. (E1, E2) Removal of the silicone wire embolus and reperfusion.

(F1, F2) Suture removal. The sutures at both ends of the disconnected vessel were

knotted, and the other two sutures were removed. (G1, G2) Anatomic reduction and

suturing of the skin. (H, I) Canavalin A label vasculature of the UPOAO mouse retina.

The silicone wire embolus was inserted into artery before perfusing rhodamine-labeled

canavalin A into the heart of UPOAO mouse. The sham eye served as an unpracticed

control eye, while the UPOAO lateral eye represented the experimental eye. Retinal

vessels in the sham eye (H) exhibited fluorescence filling, while retinal vessels in the

UPOAO lateral eye (I) remained unfilled. Scale bar = 1 mm. (J and K) Fluorescein

Fundus Angiography (FFA) performed before removing the silicone wire embolus from

the UPOAO mouse. The vessels in the sham lateral retina (J) were perfused, while the

lateral retinal perfusion in UPOAO (K) was delayed. (L) Schematic illustration of cervical

artery anatomy and ocular blood supply. Embolization of the PPA resulted in ocular

ischemia. The red arrow indicates the site of the silicone wire embolus occlusion. The

silicone wire embolus used a type 602156 wire, extended to 7mm with a diameter of

0.21mm. The blue arrows indicate the modelling locations of the HIOP model and the

UCCAO model, respectively. CCA: common carotid artery; ICA: internal carotid artery;

ECA: external carotid artery; PPA: pterygopalatine artery; MCA: middle cerebral artery;

Infraorbital art.: infraorbital artery; OA: ophthalmic artery; SPCA: short posterior ciliary

artery; LPCA: long posterior ciliary artery; CRA: central retinal artery.

Table 1 Antibodies Used in Staining of Flat-Mounted Retina and Sections

Antibodies	Species	Dilution	Company
PKCα	rabbit	1:1000	Sigma
Recoverin	rabbit	1:1000	Millipore
Calbindin	rabbit	1:5000	SWANT
ChAT	goat	1:200	Millipore
Iba1	rabbit	1:500	Wako
CD45	rabbit	1:100	CST
Rhodamine-labeled canavalin A	/	1:125	MKBio
Goat anti-mouse IgG Alexa Fluor 488	goat	1:500	Jackson
Donkey anti-rabbit IgG Alexa Fluor 594	donkey	1:500	Jackson
Donkey anti-goat IgG Alexa Fluor 488	donkey	1:500	Jackson

Fig. 1 Staining and Quantification of Retinal Ganglion Cells (RGCs) at Different Ischemia and Reperfusion Times.

Flat-mounted retina RGCs were labeled with Brn3a staining. (A, B) Representative pictures of the peripheral field (A), and quantification of surviving RGCs in all fields (B) in the 30-minute ischemia and 3-days reperfusion group. n = 3. (C, D) Representative pictures of the peripheral field (C), and quantification of surviving RGCs in all fields (D) in

the 30-minute ischemia and 7-days reperfusion group. n = 3. (E, F) Representative

pictures of the peripheral field (E), and quantification of surviving RGCs in all fields (F) in

the 60-minute ischemia and 3-days reperfusion group. n = 5. (G, H) Representative

pictures of the peripheral field (G), and quantification of surviving RGCs in all fields (H) in

the 60-minute ischemia and 7-days reperfusion group. n = 5. The results showed a

significant loss of RGCs after 60 minutes of ischemia. Data were presented as means ±

s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001, t-test. Scale bar = 100 μm.

Fig. 2 Comparison of Electroretinographic (ERG) Dark-Adapted Responses at Different

Ischemia and Reperfusion Times.

Following the evaluation of surviving RGCs, visual function in sham and UPOAO

experimental eyes at various ischemia and reperfusion times was assessed using ERG.

(A) Representative waveforms in the four groups at the stimulus light intensities of 0.1,

1.0, and 10.0 cd.s/m2, respectively. (B) Quantification of a-wave and b-wave amplitudes

in the 30-minute ischemia and 3-days reperfusion group. n = 5. (C) Quantification of a-

wave and b-wave amplitudes in the 30-minute ischemia and 7-days reperfusion group. n

= 5. (D) Quantification of a-wave and b-wave amplitudes in the 60-minute ischemia and

3-days reperfusion group. n = 7. (E) Quantification of a-wave and b-wave amplitudes in

the 60-minute ischemia and 7-days reperfusion group. n = 8. Dark-adapted responses

showed almost similar a-wave amplitudes but significantly decreased b-wave amplitudes

in the 60-minute ischemia groups. The amplitudes of b-waves declined at 3-days and

even more prominently at 7-days. (F, G) Representative OPs and quantification of

amplitudes in the 60-minute ischemic groups. n = 7 in the 3-days reperfusion group; n = 8

in the 7-days reperfusion group. The amplitudes of OPs decreased significantly at 7-days

reperfusion. The decline in b-waves and OPs along with the loss of RGCs, supports the

selection of a 60-minute ischemic duration as an appropriate choice. Data were

presented as means ± s.e.m, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$, two-way

ANOVA test for a-waves and b-waves; paired t-test for OPs.

Fig. 3 Changes in Retina Morphology in the UPOAO Model.

(A) Representative OCT images of the mouse retina at 3-days. The Green lines indicate

the OCT scan area starting from the optic disc. Local magnification and layering are

annotated. GCC: ganglion cell complex, including RNFL, GCL and IPL layers. (B and C)

Quantification of GCC and total retinal thickness at 3-days. The thickness of the GCC

and the entire retina in OCT was measured and compared at distances of 1.5 PD, 3.0 PD

and 4.5 PD from the optic disc, respectively. n = 5. (D) Representative OCT images of

the mouse retina at 7-days. (E and F) Quantification of GCC and total retinal thickness at

7-days. n = 5. (G) Representative HE images of the mouse retina at 3-days. (H, I and J)

Quantification of NFL + GCL, IPL, and INL thickness at 3-days. Retinal thickness in HE

was measured near the optic nerve head and compared. n = 3. (K) Representative HE

images of the mouse retina at 7-days. (L, M and N) Quantification of NFL + GCL, IPL,

and INL thickness at 7-days. n = 3. Data were presented as means ± s.e.m, *: p<0.05, **:

Fig. 5 Changes in Bipolar Cells, Horizontal Cells, and Cholinergic Amacrine Cells in

UPOAO Mice.

(A) Representative images of mouse retina co-stained with DAPI and PKCα at 3-days (B)

Representative images of mouse retina co-stained with DAPI and PKCα at 7-days. (C)

Quantification of PKCα fluorescence density at 3-days. (D) Quantification of PKCα

fluorescence density at 7-days. n = 4. (E) Representative images of mouse retina co-

stained with DAPI and Calbindin at 3-days. (F) Representative images of mouse retina

co-stained with DAPI and Calbindin at 7-days. Horizontal cell somata are indicated by

white asterisks. (G) Quantification of Calbindin fluorescence density at 3-days. (H)

Quantification of Calbindin fluorescence density at 7-days. n = 3. (I) Representative

images of mouse retina co-stained with DAPI and ChAT at 3-days. (J) Representative

images of mouse retina co-stained with DAPI and ChAT at 7-days. Cholinergic amacrine

cell somata are indicated by white asterisks. (K) Quantification of ChAT fluorescence

density at 3-days. (L) Quantification of ChAT fluorescence density at 7-days. n = 3. Data

were presented as means ± s.e.m, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, t- test. Scale bar =

50 µm.

Fig. 6 Transcriptomic Features at Different Times of Reperfusion.

RNA-seq evaluation was performed at 0-day, 3-days, and 7-days reperfusion periods in

UPOAO revealing enrichment in pathways related to immune cells migration, oxidative

stress, and immune inflammation. (A, D, G) Volcano plots display differential expression

genes (DEGs) between UPOAO and sham eyes in the no-perfusion group (A), 3-days

perfusion group (D), and 7-days perfusion group (G), respectively. Red dots: significant

upregulated genes, green and blue dots: significant downregulated genes, grey dots:

stable expressed genes, adjusted P < 0.05. log2FC = 1. (B, E, H) Gene ontology (GO)

analysis of differential genes in the non-perfusion group (B), 3-days group (E), and 7-

days group (H). The DEGs between UPOAO and sham eyes were enriched in pathways

associated with immune cells migration (0d), oxidative stress (3d), and immune

inflammation (7d), respectively. (C, F, I) Heatmap displaying the top 100 downregulated

and top 100 upregulated DEGs between UPOAO and sham at non-perfusion, 3-days,

and 7-days reperfusion, respectively. The box represents the genes related to immune

cells migration (C), oxidative stress (F), and immune inflammation (I). The ranking was

determined by the magnitude of fold change. In each heatmap, the upper box represents

the top 10 downregulated genes, while the lower box represents the top 10 upregulated

genes.

Fig. 7 Peripheral Leukocyte Infiltration and Retinal Resident Microglial Activation.

Rhodamine-labeled canavalin A was used for immediate cardiac perfusion to visualize

blood vessels, followed by CD45 immunofluorescent staining to observe the relationship

between blood vessels and CD45+ cells in sham (A), 1-day perfusion group (B), 3-days

perfusion group (C), and 7-days perfusion group (D). The presence of CD45$^+$ cells within

blood vessels is indicated by white arrows. Scale bar = 50 μm. CD45$^+$ cell counts were

performed in whole retinas of 1-day (E), 3-days (F), 7-days (G). The cellular morphology

and distribution of microglial cells in the superficial retina were assessed in 3-days (H)

and 7-days (I). Activated microglial cells are indicated by white asterisks. Microglial cell

counting was conducted in the superficial retina of 3-days (J) and 7-days (K). Data points

for CD45$^+$ cells were derived from four flat-mounted retinas, and data points for microglial

cells were from five flat-mounted retinas. Data were presented as means ± s.e.m, *:

$p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$, t- test. Scale bar = 50 μm.

1000

Fig. 8 Transcriptomic Results Comparison between UPOAO, HIOP, and UCCAO Models.

RNA-seq comparison between UPOAO and extravascular occlusion models: HIOP and

UCCAO. (A) Schematic illustration of the HIOP model. (B) GO analysis of differential

expression genes (DEGs) in the HIOP model at 7-days perfusion. (C) Venn diagram

indicating the overlapping DEGs (146 genes) between HIOP and UPOAO models, as

well as the remaining DEGs specific to each group (593 genes in each group). (D) GO

analysis of the overlapping DEGs between HIOP and UPOAO models. (E) GO analysis

of the remaining DEGs in the HIOP model at 7-days perfusion, excluding the overlapping

DEGs. (F) GO analysis of the remaining DEGs, excluding the overlapping DEGs. (G)

Heatmap showing the inter-sample distribution of lipid and steroid-related DEGs from the

analysis of the remaining DEGs in the UPOAO model. (H) Schematic illustration of the

UCCAO model. (I) GO analysis of DEGs in the UCCAO model. (J) Hub genes identified

through protein-protein interaction (PPI) network analysis of the DEGs in the UCCAO

model. (K) Venn diagram indicating the overlapping DEGs between UCCAO and UPOAO

models (15 genes). (L) Venn diagram indicating the presence of two overlapping DEGs

between UCCAO DEGs and m6A-related genes.

1017

1018

1019   Fig. 9 The Characteristic Features of UPOAO Model during Ischemia-Reperfusion

1020   Periods.

1021

1022

1023

1024

1025

1026

1027 **Supplementary data**

1028

Fig.S1 Response Times of a-Waves and b-Waves in ERG at Different Light Intensities.

(A, B) Response times of a-waves and b-waves were measured under different light

intensities at 3-days and 7-days post-UPOAO in the 30-minutes ischemia group. n = 5. (C, D)

Response times of a-waves and b-waves were measured under different light intensities at 3-

days and 7-days post-UPOAO in the 60-minute ischemia group. n = 7 at 3-days; n = 8 at 7-

days. Data were presented as means ± s.e.m, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****:

$p<0.0001$, two-way ANOVA test.

1036

1037

Fig.S2 Quantification of INL+OPL Thickness and ONL+IS/OS+RPE Thickness in OCT during

3-days and 7-days Reperfusion of UPOAO Animals.

The INL+OPL thickness and the ONL+IS/OS+RPE thickness were measured and compared

at distances of 1.5 PD, 3.0 PD and 4.5 PD from the optic disc using OCT. (A and B)

Quantification of INL+OPL layer thickness (A) and ONL+IS/OS+RPE layer thickness (B) at 3-

days. n = 5. No significant differences were observed. (C and D) Quantification of INL+OPL

layer thickness (C) and ONL+IS/OS+RPE layer thickness (D) at 7-days. n = 5. No significant

differences were observed. Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***:

p<0.001, ****: p<0.0001, two-way ANOVA test.

1047

1048

1049   **Fig.S3 Changes in Photoreceptor Cells in UPOAO.**

1050   (A) Representative images of mouse retina co-stained with DAPI and Recoverin at 3-days. (B)

1051   Representative images of mouse retina co-stained with DAPI and Recoverin at 7-days. n = 3.

1052   Data were presented as means ± s.e.m, *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001, t-

1053   test. Scale bar = 50 μm.

1054

1055

1056 Fig.S4 Hub Genes in PPI Analysis and Gene Set Enrichment Analysis (GSEA) during the

1057 Non-Reperfusion Stage in UPOAO.

1058 (A) Hub genes identified through PPI analysis of DEGs in the non-reperfusion group. (B-

1059 D) GSEA analysis of the pathway of the cell killing (B), lymphocyte mediated-immunity

1060 (C), and leukocyte mediated-immunity (D) in the non-reperfusion group.

1061

Fig.S5 Hub Genes, GSEA Analysis at 3-days Reperfusion in UPOAO, and Relation with

Mitochondrial Genes.

(A) Hub genes identified through PPI analysis of DEGs in the 3-days reperfusion group.

(B) GSEA analysis of the pathways related to plasma membrane protein complex (top

left), bounding membrane of organelle (top right), and cell surface (bottom). (C) Venn

diagram showing the overlapping genes between mitochondrial genes and DEGs at 3-

days reperfusion (44 genes). (D) GO analysis of the 44 overlapping DEGs.

1069

1070

Fig.S6 Hub Genes, GSEA Analysis at 7-days Reperfusion in UPOAO, and Relation with

Immune Genes.

(A) Hub genes identified through PPI analysis of DEGs in the 7-days reperfusion group.

(B) GSEA analysis of the pathways related to the reactome innate immune system (top

left), reactome immunoregulatory interactions between a lymphoid and a non-lymphoid

cell (top right), reactome immune system (bottom left), and reactome adaptive immune

system (bottom right). (C) Venn diagram showing the overlapping genes between

immune genes and DEGs at 7-days reperfusion (112 genes). (D) GO analysis of the 112

overlapping DEGs.

1080

Fig.S7 Upregulation of Immune Inflammation-Related Gene Expression in the 7-days

Reperfusion Group.

The genes Abca1, Cd86, Cd48, Tlr4, Tlr6, and Tspo were significantly upregulated in the

7-days reperfusion group, while other immune-inflammatory and chemotaxis-related

genes showed no significant differences. The data points were from the retina of four

animals. Data were presented as means ± s.e.m, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****:

$p<0.0001$, Paired t-test.

1088

1089

1090 **Fig.S8 Co-Expressed Genes during 3-days and 7-days Reperfusion.**

1091　(A) Venn diagram showing the overlap of DEGs between the 3-days and 7-days

1092　reperfusion groups, along with a list of the 17 overlapping DEGs. (B) GO analysis of the

1093　17 overlapping DEGs.

1094

1095

1096

1097

1098

1099

1100　**Table. S1 Primers Used in This Study.**

Gene Name		Sequence (5'->3')
Tlr3	Forward primer	CGATTCCCGACCAGCACTCT
	Reverse primer	CACATTTCCTTTCAGCCTGTGAT
Tlr4	Forward primer	TCCCTGCATAGAGGTAGTTCC
	Reverse primer	TCAAGGGGTTGAAGCTCAGA
Cd48	Forward primer	GGAACTGGATTTCAAGGTCATTCA
	Reverse primer	ATGGTCCAAGTGGGTCCTTAT
Ldlr	Forward primer	CCAATCGACTCACGGGTTCA
	Reverse primer	ACAGTGTCGACTTCTCTAGGC
Cd86	Forward primer	ATGGACCCCAGATGCACCA
	Reverse primer	TGTGCCCAAATAGTGCTCGT
Tspo	Forward primer	ACTGTATTCAGCCATGGGGTATG
	Reverse primer	CCAAGGGAACCATAGCGTCC
Abca1	Forward primer	GCAAGGCCGCACCATTATTT
	Reverse primer	TCTGAGAAACACTGTCCTCCTTTT
Apoe	Forward primer	CGGGTGCTCTGTTTTGGAGA
	Reverse primer	GGCTAGGCATCCTGTCAGC

S100a9	Forward primer	TGGCAACCTTTATGAAGAAAGAGA
	Reverse primer	GTGGGTTGTTCTCATGCAGC
β-actin	Forward primer	CACTGTCGAGTCGCGTCC
	Reverse primer	TCATCCATGGCGAACTGGTG

1101

Table. S2 Summary Table: Time Course of all the Morphological, Functional, Cellular, and Transcriptome Changes in the UPOAO Model.

Changes in each group	30min-3d	30min-7d	60min-0d	60min-3d 1.5 PD	60min-3d 3 PD	60min-3d 4.5 PD	60min-7d 1.5 PD	60min-7d 3 PD	60min-7d 4.5 PD
RGCs counts	-	-	/		↓			↓	
ERG — Amplitudes of a-waves	-	-			-			- (0.001, 0.003, 0.01, 0.03, 0.1, 1.0 cd.s/m^2) ↓ (0.3, 3.0, 10.0 cd.s/m^2)	
ERG — Response time of a-waves	-	-	/		-			-	
ERG — Amplitudes of b-waves	-	-			↓			↓	
ERG — Response time of b-waves	-	-			-			-	
ERG — Amplitudes of OPs-waves		/			↓			↓	
OCT — GCC thickness				-	-	-	↓	↓	↓
OCT — INL+OPL thickness		/		-	-	-	-	-	-
OCT — ONL+IS/OS+RPE thickness				-	-	-	-	-	-
HE — total retinal thickness				-	-	↓	↓	↓	↓
HE — RNFL+GCL thickness					↑			↓	
HE — IPL thickness					↓			↓	
HE — INL thickness					-			↓	
HE — OPL thickness		/			-			-	
HE — ONL thickness					-			-	
HE — IS/OS+RPE thickness					-			-	

Changes in each group		30min-3d	30min-7d	60min-0d	60min-3d	60min-7d
Retina neural cells	Bipolar cells (PKCα)				-	↓
	Horizontal cells (Calbindin)				-	↓
	Cholinergic amacrine cells (ChAT)		/		↓	↓
	Photoreceptor cells (Recoverin)				-	-
Peripheral leukocyte infiltration (CD45) [#]			/		↑	↑
Microglia counts (Iba1)			/		↑	↑
RNA-seq	DEGs number		/	419 (up: 215; down: 204)	542 (up: 372; down: 170)	739 (up: 429; down: 310)
	Enrichment			Immune cells migration-related pathways	Oxidative stress-related pathways	Immune inflammation-related pathways

1104   -: no change;

1105   ↑: increase;

1106   ↓: decrease;

1107   /: not involved in the article;

1108   #: supplementary content: also increase after 60min-ischemia and 1d-reperfusion;

1109

Video 1. Blunt Dissection of Cervical Vessels. The neck was exposed, and the cervical

arteries were separated using blunt dissection techniques.

Video 2. Insertion of Silicone Wire Embolus and Induction of Ischemia. The silicone wire

embolus was inserted into the PPA, initiating the ischemic condition. PPA: the

pterygopalatine artery.

Video 3. Extraction of Silicone Wire Embolus and Reperfusion. The silicone wire embolus

was removed, allowing reperfusion to occur. The mouse's wound was sutured, and

standard feeding procedures were followed.

1118

A **B** **C** **D** **E** **F** **G**

A1 B1 C1 D1 E1 F1 G1

A2 B2 C2 D2 E2 F2 G2

H **I** **L**

Location of modeling
in the HIOP model

Infraorbital art.

SPCA
LPCA

MCA
OA
CRA

**Location of embolism
in the UPOAO model**

ECA
PPA

ICA
Palatine portion of PPA

J **K**

**Location of ligation
in the UCCAO model**

CCA

602156 silicone wire plug

Diameter of silicone: 0.21 ± 0.01mm

No. 0.4 line

Length of silicone: 7mm ± 0.1mm

UCCAO

①Closer to the disease process
②Simulate acute and complete blood flow disruption

UPOAO

30-60 Minutes Timer

HIOP

Air or saline was injected

Increased pressure

Without mechanical pressure from increased intraocular pressure

Pathological Process of Reperfusion Periods

1d_Leukocyte Migration

Chemotaxis to corresponding site

3d_Oxidative Stress

Blood   **Retina**

Microglial activation

Leukocytes migration

7d_Immune Inflammation

Leukocytes   Microglia

↕ Interaction

RGCs   ACs   BCs   HCs

Neuroinflammatory

Neuronal cells death

Vision Loss