Subcellular imaging of lipids and sugars using genetically encoded proximity sensors

William M. Moore[1], Roberto J. Brea[1§], Caroline Knittel[1], Ellen Wrightsman[1], Brandon Hui[1],
Jinchao Lou[2], Christelle F. Ancajas[2], Michael D. Best[2], Neal K. Devaraj[1], and Itay Budin[1*]

[1] Department of Chemistry and Biochemistry, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093

[2] Department of Chemistry, The University of Tennessee, Knoxville, Tennessee 37996

§ Current address: Centro Interdisciplinar de Química e Bioloxía (CICA), Universidade da Coruña, Rúa As Carballeiras, 15701 A Coruña, Spain

* To whom correspondence should be addressed: ibudin@ucsd.edu

Abstract

Live cell imaging of lipids and other metabolites is a long-standing challenge in cell biology. Bioorthogonal labeling tools allow for the conjugation of fluorophores to several phospholipid classes, but cannot discern their trafficking between adjacent organelles or asymmetry across individual membrane leaflets. Here we present fluorogen-activating coincidence sensing (FACES), a chemogenetic tool capable of quantitatively imaging subcellular lipid pools and reporting their transbilayer orientation in living cells. FACES combines bioorthogonal chemistry with genetically encoded fluorogen-activating proteins (FAPs) for reversible proximity sensing of conjugated molecules. We first validate this approach for quantifying discrete phosphatidylcholine pools in the ER and mitochondria that are trafficked by lipid transfer proteins. We then show that transmembrane domain-containing FAPs can be used to reveal the membrane asymmetry of multiple lipid classes that are generated in the trans-Golgi network. Lastly, we demonstrate that FACES is a generalizable tool for subcellular bioorthogonal imaging by measuring changes in mitochondrial *N*-acetylhexosamine levels. These results demonstrate the use of fluorogenic tags for spatially-defined molecular imaging.

Introduction

Lipid composition determines the biophysical properties of cell membranes and varies across every scale of biological organization: from tissues, to organelles, and even between individual leaflets of the same bilayer[1–3]. Investigating membrane chemical heterogeneity remains a challenge due to the paucity of tools for imaging lipids in live cells. Bioorthogonal 'click' chemistry has emerged as one tool for fluorescence labeling of specific lipid classes – including phosphatidylcholine (PC)[4], phosphatidylserine (PS)[5], phosphatidylinositol (PI)[6], phosphatidic acid[7], and glycosphingolipids[8] – that can be metabolically tagged with azide-bearing head group components (Fig. 1a). Azido (N_3) lipids are commonly conjugated to fluorophores using strain-promoted azide-alkyne cycloaddition (SPAAC) reactions, unveiling their cellular distribution once unreactive dyes are washed out. While powerful, this approach indiscriminately labels all lipids in the cell, and is thus limited by the resolution constraints of light microscopy: this approach cannot discriminate between lipid pools in organelles at close proximity or in individual leaflets of the same membrane, for example. To overcome some of these limitations, organelle-targeted fluorophores have been elegantly applied to image subcellular lipid pools[9]. Currently, these are limited to just a few chemical environments in the cell, such as rhodol-containing groups that localize to both endoplasmic reticulum (ER) and Golgi membranes, and thus lack additional specificity or resolution. They also depend on the self-segregation of dyes within the cell based on chemical cues, like membrane potential, which can be perturbed by lipid manipulation. We currently lack generalizable strategies to measure intracellular lipid heterogeneities in live cells and therefore cannot determine the molecular mechanisms by which they are generated.

We hypothesized that challenges for lipid imaging could be overcome using protein-based lipid sensors, whose localization is precisely controlled through signal sequences or fusion to other

protein components. Previously, natural lipid binding proteins derived from microbial toxins, such as perfringolysin O[10] (cholesterol) or equinatoxin II[11] (sphingomyelin, SM) have been expressed as fusions with fluorescent proteins (FP), to detect membranes enriched in the target lipids. In these modalities, lipids are qualitatively detected by binding of the probe to a membrane, but this limits their application to large compartments for which membrane binding is observable and does not allow for quantitative detection. We asked whether 'turn on' fluorophores, termed fluorogen-activating proteins (FAPs), could provide a more generalizable strategy. FAPs are expressed as fusion constructs with other proteins or targeting sequences, similar to FPs, but are not themselves fluorescent[12]. Instead, they interact with fluorogens, small cell-permeable molecules that act as fluorophores only upon specific binding to a FAP (Fig. 1b). Depending on the FAP used, fluorogen binding can be irreversible or reversible; the former provides brighter fluorescence for conventional imaging, while the latter generates stochastic blinking that can be harnessed for super-resolution imaging[13]. Recently, FAP domains have been applied towards detecting and controlling protein proximity in living cells through reversible dimerization[14], demonstrating that this technology can be harnessed in new applications beyond the capacity of FPs.

We sought to harness the unique properties of fluorogens through Fluorogen-Activating Coincidence Sensing (FACES), which uses the FAP protein for proximity-dependent reporting of fluorogen conjugated bioorthogonal metabolites, including lipids. To accomplish this, we synthesized fluorogenic dyes based on malachite green (MG) that are functionalized with a dibenzocyclooctyne (DBCO) group for SPAAC reactions with azido-labeled metabolites in living cells (Fig. 1c). In the FACES workflow (Fig. 1d), cells first express a FAP protein fused to an FP, like GFP, that is targeted to an organelle or site of interest. Cells are then incubated with azido-labeled metabolites for metabolic incorporation. Immediately before imaging, incorporated metabolites are rapidly labeled with fluorogen-DBCO, which remains non-fluorescent unless it is

proximal to the expressed FAP. Excess dye is removed from cells by washing with culture medium, leveraging the reversibility of the FAP-fluorogen complex to prevent detection of unreacted fluorogens. Subsequent ratiometric fluorescence imaging allows organelle-specific detection of fluorogen conjugates by coincidence sensing with the FAP receptor. The abundance of the target molecule is then quantified at the locale of interest by measuring the ratio of FAP-fluorogen fluorescence, which represents the bound state of the FAP and is at red shifted wavelengths for MG, to that of a green of blue reporter FP, which is used to normalize the rate of detection to the concentration of the FAP receptor.

Here we validate the FACES strategy through imaging of discrete N_3-lipid pools in the ER, mitochondria, and trans-Golgi network (TGN). Ratiometric imaging with FACES enabled changes in N_3-lipid content of membranes to be quantified, which we use to characterize the role of ER-mitochondrial PC transporters. We then show that transmembrane-anchored FAPs can probe membrane asymmetry generated during vesicle formation at the TGN. Lastly, we demonstrate that FACES is a broadly applicable tool compatible with non-lipid metabolites by measuring fluctuations in mitochondrial N-acetylhexosamine (HexNAz) levels.

Results

Development and validation of FACES

We synthesized derivatives of MG with a linker and DBCO group, compatible with copper-free SPAAC reactions, including compound **1a** (Fig. 1c; Supplementary Fig. 1). MG non-covalently binds to single-chain antibody scFvs FAP and the complex emits far-red fluorescence. We tested the compatibility and fluorogenic properties of **1a** with FACES using the FAP clone dH6.2, a lower affinity scFvs variant developed for reversible equilibrium binding to MG[15]. In all

experiments, we fused dH6.2 – from here on simply referred to as FAP – to GFP for ratiometric

imaging. A secretion signal and KDEL retention sequence were first used to target FAP-GFP to

the endoplasmic reticulum (ER) as the major site of PC synthesis[16] (Fig 1c), while GFP-KDEL,

an identical construct but lacking the FAP domain, was used as a control. HeLa cells transfected

with either FAP-GFP-KDEL or GFP-KDEL were grown for 24h in the presence or absence of 50

µM N_3-Cho. At concentrations up to 100 µM, N_3-Cho was incorporated into PC at a frequency up

to 0.1 mol %, but cells showed no detectable N_3-Cho incorporation into SM (Supplementary Fig.

2). Prior to staining, cells were washed and grown in normal growth media for 30 min to allow

for the metabolic consumption of any remaining N_3-Cho. Cells were then transferred into

FBS-free growth media containing 500 nM **1a** and imaged while in the staining solution (Fig.

2a). Images were collected using identical acquisition and processing settings.

Results from pre-washed cells demonstrated that **1a** was cell permeable and generated strong

far-red fluorescence in cells expressing FAP-GFP-KDEL. No far-red fluorescence was detected

in cells expressing GFP-KDEL, indicating that **1a** fluorescence was not activated directly by

conjugation to N_3-PC. Next we tested the reversibility of the FAP-fluorogen complex by washing

cells with growth medium and re-imaging (Fig. 2b). Three washes were sufficient to dissociate

free **1a** from FAP-GFP-KDEL cells in the absence of N_3-Cho, resulting in the loss of far-red

fluorescence. In contrast, far-red fluorescence was retained in FAP-GFP-KDEL cells that were

fed 50 µM N_3-Cho, which implicated the SPAAC reaction product **1a**-PC as the source of

FAP-dependent fluorescence. Similar experiments with a higher affinity FAP variant retained

far-red fluorescence after washing in the absence of N_3-Cho labeling (Supplementary Fig. 3),

indicating that the reversible binding to the MG product is essential for substrate-specific

detection.

We further quantified these results in repeated experiments using confocal laser scanning microscopy (CFLSM) to assess linear fluorescent detection. Cells transfected with FAP-GFP-KDEL were fed 10, 30, or 50 µM N_3-Cho for 24h, while FAP-GFP-KDEL (no N_3-Cho) and GFP-KDEL (+ 50 µM N_3-Cho) conditions were used as controls. All cells were stained with 500 nM **1a** and were washed as previously described; the **1a**-PC reaction product was confirmed by LC/MS/MS (Supplementary Fig. 4). Fluorescence intensity profiles revealed that far-red fluorescence from **1a** was tightly correlated with GFP fluorescence in FAP-GFP-KDEL cells fed 50 µM N_3-Cho (Fig. 2c). In contrast, cells from either control condition, lacking the FAP or N_3-Cho component, had low levels of far-red fluorescence that did not correlate with GFP localization (Fig. 2c). The average fluorescence intensity from 10 cells, using 10 individual ER segments per cell, was calculated as a ratiometric average relative to GFP. There was no intensity difference between either control condition, while signal from FAP-GFP-KDEL cells fed 50 µM N_3-Cho was 10-fold higher (Fig. 2d). Additionally, **1a**-PC fluorescence was linearly dependent on N_3-Cho concentration in the medium (Fig. 2e). We further tested whether the FACES signal was a consequence of free N_3-Cho, as opposed to membrane incorporated N_3-PC, by using compound D609 to pharmacologically perturb PC metabolism[17–19]. Cells treated with 50 µM D609 during the 24h N_3-Cho incorporation period decreased FACES fluorescence by 50% relative to untreated cells, despite being exposed to identical concentrations of N_3-Cho and **1a** (Fig. 2f). Together, these results indicate that FACES is capable of linear fluorescent quantification of N_3-PC, via its SPAAC reaction product **1a**-PC, in the ER of living cells.

FACES quantifies lipid transport between organelle pools

PC is the most abundant lipid in mammalian cells and is a component of all organelle membranes, which make disentangling its subcellular transport mechanisms challenging. To demonstrate the utility of the FACES method in illuminating organelle selective lipid pools, we

used the OTC mitochondrial targeting sequence (MTS) to direct FAP-GFP to the mitochondrial matrix. Cells expressing MTS-FAP-GFP and the ER marker Sec61b-mCherry showed that **1a**-PC in mitochondria did not overlap with the ER (Fig, 3a). Reciprocally, cells co-expressing FAP-GFP-KDEL and MTS-BFP show that **1a**-PC in the ER did not overlap with mitochondria (Fig. 3b).

When quantifying the performance of FACES in the mitochondrial matrix using CFLSM, we found that switching the C6 alkane linker in **1a** to a polyethylene glycol (PEG)-2 linker (compound **1b**) was beneficial for dissociating unreacted dye from this compartment. Similar to the ER-targeted protein, MTS-FAP-GFP detection of **1b**-PC generated fluorescence that linearly correlated with N_3-Cho concentration (Supplementary Fig. 5). Using super-resolution lattice-structured illumination microscopy (lattice SIM^2), we further tested whether the matrix soluble FAP could resolve structural features of the IMM. When imaged by lattice SIM^2, FACES signal from **1b**-PC occurred in a banding pattern distinct from the general distribution of the probe (GFP signal) in the mitochondrial matrix (Fig. 3c). Fluorescence intensity profiles drawn across individual mitochondria show undulating peaks, with alternating maxima between **1b**-PC and GFP. We interpreted this localization pattern as reflecting regions of high cristae density in the IMM that contrast with void areas of the matrix[20], as previously observed for mitochondrial PC[9].

Utilizing the ability to quantify PC in mitochondria and ER, we tested how defects in lipid transport differentially affect these pools. For this, we generated stable cell lines expressing either FAP-GFP-KDEL (ER) or MTS-FAP-GFP (mitochondria) for further experiments. PC is synthesized in the ER and is transported to mitochondria by lipid transport proteins (LTPs)[16,21]. We used siRNA to silence two LTPs: VPS13a, a bridge-type LTP that interacts with VAPB at ER-mitochondria contact sites and transports bulk phospholipids[22], and StarD7, a small

shuttle-type LTP that localizes to the mitochondrial intermembrane space and is thought to

transport PC between outer and inner membranes[23–25] (Fig. 2d). Silencing resulted in a >50%

reduction in both VPS13a and StarD7 protein content in each experiment (Supplementary Fig.

6). VPS13a and StarD7 siRNAs reduced **1b**-PC fluorescence in mitochondria, by 50% and 35%

respectively, relative to negative control (NC) siRNA treated cells (Fig. 3e). However, only

knockdown of VPS13a increased **1a**-PC fluorescence in the ER (Fig. 3f), consistent with its

proposed role as an ER to mitochondrial PC LTP and that of StarD7 as an intra-mitochondrial

LTP. The magnitude of **1b**-PC reduction in mitochondria was larger upon knockdown of VPS13a

than for that of StarD7, suggesting that the former is a dominant mechanism for bulk PC

transport to mitochondria.

<u>FACES measures transbilayer asymmetry arising in the TGN</u>

It has long been recognized that membrane compartmentalization inherently generates

directionality between exoplasmic and cytoplasmic leaflets, which is reflected both in membrane

protein topology and the transbilayer distribution of lipids[26]. Detailed lipidomic investigation of

membrane asymmetry has only been carried out in the PM of erythrocytes, whose lack of

intracellular compartments allow for calculation of lipid accessibility to externally added

phospholipases[3,27]. While asymmetry is an emerging concept in membrane biology, the

transbilayer distribution of phospholipids in intracellular membranes is poorly understood. There

exists few, if any, tools capable of measuring membrane asymmetry in living cells and we asked

if the spatial specificity of FACES could provide this capacity.

A potential site of asymmetry generation in the secretory pathway is at the TGN, where vesicles

bud off from the Golgi for trafficking of proteins and lipids to the highly asymmetric PM. We

explored asymmetric lipid detection in the TGN by reciprocal tagging of the transmembrane

protein TGN38, at N- and C-termini, to selectively orient FAPs toward either the exoplasmic or

cytoplasmic facing leaflets of the TGN (Fig. 4a). TGN38 cycles to and from the PM via secretory

vesicles and recycling endosomes[28], thereby labeling a split population of trans-Golgi cisternae

and trans-Golgi network vesiculated compartments[29]. Imaging of stable cells expressing

GFP-TGN38-FAP, oriented towards the cytoplasmic leaflet, and FAP-TGN38-GFP, oriented

towards the luminal leaflet, detected fluorogen fluorescence of **1a**-PC on both leaflets in

cisternae and vesicles (Fig. 4b). However, the abundance shifted between these compartments,

comparatively enriching in the cytoplasmic leaflet in TGN vesicles. Lateral segregation between

luminal **1a**-PC enriched and depleted regions could be observed at sites of vesicular budding

from the TGN (Supplementary Fig. 7a). These results are consistent with previous

measurements of PC in freeze-fractured yeast cells, also showing a cytoplasmic leaflet

enrichment arising in the Golgi[30].

We next asked if FACES could measure the asymmetry of non-choline lipids at the TGN.

Recently, metabolic incorporation of azide containing L-serine[5] (C-L-Ser-N_3) and *myo*-inositol[6]

(Ino-N_3) have been shown to label phosphatidylserine (PS) and phosphatidylinositol (PI),

respectively. Both these lipid classes are synthesized in the ER and have been proposed to be

trafficked to the cytoplasmic leaflet of the trans-Golgi and TGN through lipid transfer proteins[31,32].

When analyzed with FACES, we observed **1a**-PI in Golgi cisternae that was evenly distributed

between luminal and cytoplasmic leaflets (Fig. 4c, Supplementary Fig. 8a). In TGN vesicles, the

distribution of **1a**-PI shifted to the cytoplasmic leaflet, similar to that for PC. For **1a**-PS, using

C-L-Ser-N_3, there was an absence of signal on both leaflets of Golgi cisternae and a high,

exclusively cytoplasmic abundance on TGN vesicles (Fig. 4c, Supplementary Fig. 8b). This is

consistent with previous observations that the PS-binding C2 domain of lactadherin, when

expressed in the cytoplasm, robustly labels TGN vesicles, but not Golgi cisternae[33]. As a control

for these cytosolic leaflet-enriched phospholipids, we imaged the distribution of surface glycans

in cells fed with peracetylated *N*-azidoacetylmannosamine (Ac$_4$ManNAz)[34], which is metabolized

into sialic acid (Sia) within the cell. We measured an exclusively luminal distribution of **1a**-Sia

(Fig. 4c, Supplementary Fig. 8c) in both cisternae and vesicles, consistent with the synthesis of

protein and lipid surface glycans in the Golgi lumen[35]. Lastly, co-staining cells expressing

mCer3-TGN38-FAP (Blue FP variant) with equimolar amounts **1a** and BODIPY-DBCO, a

conventional fluorophore, demonstrated that FACES can spectrally separate organelle-specific

lipid pools that overlap with total cellular phospholipid distributions (Supplementary Fig. 9)

Previous analyses of TGN vesicles[36] and the PM[37] suggest that loss of luminal phospholipids

could be substituted with SM, which is predominantly synthesized in the TGN and packaged in

secretory vesicles. Consistent with this model, we observed that the SM probe SMeqt-GFP

co-localized with vesicles enriched with cytoplasmic leaflet **1a**-PC (Fig. 4d). These vesicles

shuttled rapidly between the Golgi and PM (Supplementary Movie 1). Similar results were

obtained for vesicles labeled with the secretory cargo mCherry-GPI (Supplementary Fig. 10,

Supplementary Movie 2), which also co-traffics with SM in secretory vesicles[11]. Overall, these

results support a model in which phospholipids redistribute to the cytoplasmic leaflet during

TGN vesicularization (Fig 4d), while the luminal leaflet becomes enriched in newly synthesized

sphingolipids.

<u>FACES reports on changes to mitochondrial glycans</u>

Based on our detection of surface glycans in the Golgi lumen, we asked whether FACES could

be used as a general tool compatible for imaging of azido-sugars in living cells.

N-azidoacetylgalactosamine-tetraacylated (Ac$_4$GalNAz) is widely used as a metabolic reporter

for glycoconjugates containing *N*-acetylhexosamines (HexNAc)[38]. Ac$_4$GalNAz fed to cells is

converted to UDP-*N*-azidoacetylgalactosamine (UDP-GalNAz) by the hexosamine salvage

pathway and is then further epimerized to UDP-*N*-azidoacetylglucosamine (UDP-GlcNAz). This

results in robust labeling of *N*- and *O*- linked glycoproteins on the cell surface[38,39], secretory

compartments where they are formed, and endosomal compartments where they are recycled

(Fig 5a). Meanwhile, intracellular protein glycosylation is modulated by nucleocytoplasmic and

mitochondrial *O*-linked β-*N*-acetylglucosamines (*O*-GlcNAc), which are added to

serine/threonine residues by *O*-GlcNAc transferase (OGT) and removed by *O*-GlcNAcase

(OGA)[38]. *O*-GlcNAc cycling has emerged as a regulator of metabolism, signaling, and immune

evasion of cancer cells[40–42] and mitochondrial motility in neurons[43]. However, intracellular

glycosylations are not readily observed by standard bioorthogonal imaging approaches due to

the much more extensive incorporation of $Ac_4GalNAz$ into surface glycans; very little signal from

conjugated conventional fluorophores co-localizes with mitochondria, for example (Fig. 5b).

Currently, detection of *O*-GlcNAcylation is performed in fixed cells or bulk extracts using

O-GlcNAc sensitive antibodies[44].

Using MTS-FAP-GFP expressing cells, we tested the efficacy of FACES in reporting

mitochondrial-specific changes in HexNAc. Cells were grown in the presence or absence of

either 100 μM $Ac_4GalNAz$ or 100 μM *N*-azidoacetylglucosamine-tetraacylated ($Ac_4GlcNAz$) for

24h. $Ac_4GlcNAz$ is not efficiently incorporated by the hexosamine salvage pathway[38] and thus

was used as a less metabolically active surrogate. SPAAC staining with **1b** generated

fluorescence signal intensities 10-fold greater in cells fed 100 μM $Ac_4GalNAz$ compared to unfed

control cells (Fig. 5c). In contrast, **1b** fluorescence from cells fed $Ac_4GlcNAz$, which is less

efficiently incorporated into glycosylated proteins, was significantly lower than that for

$Ac_4GalNAz$, indicating that the FACES signal from **1b**-HexNAz is from sugars that have been

metabolically incorporated and not due to free azido sugars.

Next we tested whether FACES can report changes to mitochondrial O-GlcNAc levels resulting

from pharmacological treatment or nutritional state. We found that the OGA inhibitor Thiemet-G

increased **1b**-HexNAz fluorescence, suggesting that the fluorogen signal was responsive to the

activity of O-GlcNAc modifying enzymes (Fig 5d). In cancer cells, glucose uptake is positively

correlated with O-GlcNAc levels, which in-turn reprogram cellular metabolism tailored towards

glucose availability[45]. We observed that switching cells from high (30 mM) to low glucose (5.5

mM) levels during $Ac_4GalNAz$ labeling reduced **1b**-HexNAz fluorescence by 30% (Fig. 5e). This

is similar to glucose-dependent changes in mitochondrial O-GlcNAc levels previously reported

in breast cancer cells[46]. Taken together, these data demonstrate that FACES can report

changes in mitochondrial HexNAz incorporation as a function of cell metabolism and

O-GlcNAcylation in living cells, and support the hypothesis that mitochondrial O-GlcNAcylation

is a post-translational modification that is responsive to the cell's dietary state.

Discussion

Studying cellular roles for non-proteinaceous biomolecules has long been hindered by the lack

of robust tools for their imaging in live cells. Here we introduce a new approach, FACES, that

combines the chemical specificity of metabolic incorporation with the spatial control of genetic

protein tagging for imaging of cellular metabolites. FACES harnesses reversible fluorogenic

systems, where fluorescence increases many orders of magnitude upon specific binding to a

FAP. It combines this turn on fluorescence with the specificity of bioorthogonal reactions for

molecular sensing. We highlight the use of this approach for subcellular imaging of

phospholipids, their transport between organelles, and orientations across membrane leaflets.

We propose that FACES is a generalizable tool that can be applied to other biomolecules.

Metabolic labeling has emerged as a particularly important tool for glycobiology and we

demonstrate that FACES is responsive to labeled glycoproteins in mitochondria, which have

been proposed to act as regulators of metabolic function[47].

Compared to other bioorthogonal imaging modalities, FACES has several key advantages for

mechanistic cell biology. For organelle-defined imaging, protein targeting sequences are more

specific and generalizable than corresponding chemically-encoded localization tags. We show

that this capacity can be applied to quantitative characterization of lipid distribution phenotypes

affected by trafficking networks across membrane contact sites. Beyond organelle-specific

imaging, FAPs can also be fused to specific proteins, allowing FACES-based detection of

molecules in defined nanoenvironments. Demonstrated is one such application, where FAP

domains are introduced on both sides of a TMD to read out bilayer asymmetry. Here, FACES

provides effective discrimination beyond what is possible with direct fluorescence imaging, as

the distance between two bilayer leaflets is less than 4 nm. Fluorogens show no fluorescence

when unbound to their corresponding FAP, so quantitative detection of conjugate products

occurs through the resulting fluorescence signal, not a change in its localization as for lipid

binding proteins. Because of this capacity, FACES is the only method, to our knowledge, that

allows for measurement of membrane asymmetry in live cells.

Applications of FACES could be further expanded through increases in the repertoire of

metabolic labeling, bioorthogonal chemistry, and the spectral plasticity of the fluorogen

employed. Conjugation of fluorogens to biomolecules is dependent on tagged metabolite

incorporation, which are continuously expanding across biological systems[48] and molecular

classes[49]. One concern with any metabolic labeling approach is that tagged molecules might not

fully mimic native compounds, which is especially relevant for lipids with minimal headgroups.

For several azido lipids, previous studies suggest that compounds localize across a range of

cells as expected[4,5], though it is challenging to preclude all differences between labeled and

unlabeled species. An additional limitation common to bioorthogonal approaches is the time

required for conjugation and washing out of unreacted fluorogens; these might make the

analysis of rapid redistributions of the underlying labeled species, like those relevant for lipid

signaling[50], challenging. Tuning fluorogenic reagents for faster conjugations[51] and washing could

decrease the time for processing steps to capture dynamic movements. Lastly, FACES could

also be broadened to utilize other fluorogen systems, including those based on the Photoactive

Yellow Protein[52] and bacterial lipocalin[53]. The MG-based system employed here was chosen

because of its very low background fluorescence in the unbound state. Because of the single

wavelength emission of the bound MG, measurements in two environments were taken across

multiple cells in separate experiments. Further development of multi-color FACES systems

would enable simultaneous detection of multiple subcellular pools.

Our experiments demonstrate that FACES is an especially powerful tool for investigating cellular

mechanisms by which lipids are transported between organelles and across individual

membranes, an intense area of investigation. A plethora of shuttle and bridge-like LTPs that

perform lipid exchange at contact sites[54] and flippases/floppases/scramblases that act on bilayer

asymmetry[55] have been identified and biochemically tested. However, these transporters are

characterized by overlapping affinities, apparent redundancies, and close interactions with lipid

metabolic enzymes; how they act in concert to shape the membrane heterogeneity that defines

eukaryotic cells is still unknown. Here, we show that FACES can report the asymmetry of

several labeled phospholipid classes in post-Golgi vesicles. The Golgi has long been proposed

as a site of asymmetry generation, given the co-occurrence of secretory cargo sorting to the PM

and sphingolipid synthesis[56]. Golgi membranes also host phospholipid flippases; genetic and

biochemical studies in yeast have suggested that TGN vesicle budding is tied to the generation

of lipid asymmetries by P4-ATPase that flip phospholipids[57]. Vesicles that are defined by

phospholipid asymmetry are enriched in GPI-anchored secretory cargoes that associate with

the exoplasmic leaflet of the PM, consistent with these models. Lipid asymmetry generated at the TGN could also be required to sort transmembrane proteins based on the asymmetric TMD sequences that are required for PM targeting[3,58]. As a tool that provides spatially-defined molecular quantitation, FACES could be broadly powerful for untangling how lipid trafficking, generation of membrane asymmetry, and transport of cargoes occurs throughout eukaryotic cells.

Acknowledgements

Gulcin Perkunnaz provided helpful discussions and reagents, Jennifer Santini and the UCSD Microscopy Core (NS047101, OD030505) provided assistance with microscopy, Chris Burd and Jennifer Lippincott-Schwartz provided plasmids. The work was supported by the National Institutes of Health (R35-GM142960 to I.B. and R35GM141939 to N.K.D.) and the National Science Foundation (CHE-2310263 to M.D.B.). R.J.B. acknowledges Agencia Estatal de Investigación and the Ministerio de Ciencia e Innovación (RYC2020-030065-I).

Methods

Synthesis of fluorogen reagents: Synthesis schematic of compounds **1a** and **1b** are shown in Supplementary Fig. 1b and synthetic protocols are detailed in the Supplementary Text. Products were validated by LC-MS and ^{1}H/^{13}C NMR.

Generation of constructs: FAPs were cloned from pcDNA3.1-KozATG-dH6.2-2XG4S-actin[15], pcDNA3.1-kappa-myc-dK-2xG4S-TMst[59], and pcDNA3.1-kappa-myc-dL5-2xG4S-mCer3-KDEL[60] (Addgene). These FAP sequences and the pcDNA3.1 backbone were used for generating the FAP-GFP-KDEL, GFP-KDEL, and MTS-FAP-GFP constructs used in transfection

experiments (Fig. 2). The Ig Kappa secretion signal was used in conjunction with a KDEL

sequence for ER-targeted constructs, while the targeting sequence from OTC was used for

MTS constructs. For stable expression, FAP-GFP-KDEL and MTS-FAP-GFP coding sequences

were subcloned into the sleeping beauty expression vector SBbi-PUR[61] (Fig. 3, Fig. 5).

Constructs utilizing TGN38 were derived from mCer3-TGN38, which was a gift from the

Lippincott-Schwartz lab. In transfection experiments, the dH6.2 sequence was inserted into

either the N-terminal or C-terminal side of the TGN38 transmembrane domain to generate

mCer3-FAP-TGN38 and mCer3-TGN38-FAP constructs, respectively (Fig. 4d, Supplementary

Figures 7 and 10). For stable cell lines, a reciprocal tagging approach was taken to generate

FAP-TGN38-GFP and GFP-TGN38-FAP constructs assembled into the SBtet-Pur expression

vector (Fig. 4b-c, Supplementary Fig. 8). The GPI-anchor marker protein was generated by

subcloning a mCherry-SBP-GPI sequence, received from the Lippincott-Schwartz lab, into

pcDNA3.1. The GFP-EqtSM plasmid was received as a gift from the Burd lab. Additional

Sec61mCherry and Mito-BFP plasmids were received from the Lippincott-Schwartz lab. All

constructs were validated by whole plasmid sequencing.

<u>Cell culture:</u> HeLa (ATCC CCL-2) cells were grown in Dulbecco's Modified Eagle Medium

(Gibco 11995-065) containing glucose (4.5 g/L), glutamine (4.5 g/L), sodium pyruvate (110

mg/L), phenol red, 10% FBS, and 1% PenStrep, at 37°C and 5% CO_2 atmosphere. For

microscopy, cells were seeded into 35 mm glass bottom dishes (MatTek) coated with fibronectin

(7.5 μg/mL). For transient expression, cells were transfected with Lipofectamine 3000 in

Opti-MEM media for 4 h, and changed into normal growth medium containing azido-containing

bioorthogonal metabolite for 24 h prior to SPAAC labeling and imaging. Stable cell lines

expressing FAPs were generated using the Sleeping Beauty transposon insertion system. After

transfection, cells were grown under selection with puromycin (10 μg/mL) for 2 weeks until all

showed fluorescence. Cells were routinely tested for mycoplasma by MycoStrip (InvivoGen).

 <u>Metabolic labeling:</u> 1-azidoethyl-choline (N_3-Cho, Jena BioScience) and resuspended in PBS to

stock concentration of 50 mM. $Ac_4ManNAz$, $Ac_4GalNAz$ and $Ac_4GlcNAz$ (Click Chemistry Tools)

were resuspended in DMSO to stock concentration of 100 mM. L-Serine analogue C-L-Ser-N_3

and *myo*-Inositol analogue Inos-N_3 were synthesized as previously reported, verified by $^1H/^{13}C$

NMR and high resolution mass spectrometry, and resuspended in PBS to a stock concentration

of 50 mM and 100 mM, respectively. In all cases, metabolic incorporation occurred for 24 h.

Afterwards, cells were washed once with HBSS buffer, and incubated in fresh growth media for

30 min to allow for metabolic consumption of any remaining bioorthogonal metabolites. For

experiments with peracetylated sugars, 1 h incubation was used.

<u>SPAAC conjugation:</u> Culture media was removed by aspiration and replaced with staining media

for SPAAC labeling. The staining media contained 500 nM **1a** or **1b** diluted in serum-free DMEM

clear (Gibco 31053-028), and incubated for 15-20 min at 37°C and 5% CO_2 atmosphere. Cells

were washed with 4 volumes of culture media containing 10% FBS, 3 times, at 15 min intervals

to remove unreacted dye. For experiments with the mitochondrial-localized FAP, cells were

incubated for an additional 45 min, followed by a fourth wash step prior to imaging.

<u>Lipid analysis:</u> Lipids were extracted with a modified Bligh and Dyer protocol. Briefly, cell pellets

from T75 culture flasks were resuspended in 200 μL PBS and extracted with 2 mL 1:2

$CHCl_3$:MeOH (v/v) and cell debris were removed by centrifugation. Extracts were transferred to

new vials and 1 mL chloroform was added, followed by 1 mL H_2O, and shaking. The lower

organic phase was collected after centrifugation and the aqueous phase was extracted twice

more with 650 μL and pooled. Lipid extracts were then dried under N_2 and resuspended in 100

μL 1:2 $CHCl_3$:MeOH for thin-layer chromatography (TLC) and 500 μL methanol for LC-MS/MS.

For TLC, entire lipid extracts were spotted onto 10x10 cm HPTLC silica gel 60 plates,

developed with 14:5:1 $CHCl_3:MeOH:NH_4$, and imaged with a Typhoon FLA 9500 using 633 nm

laser. For LC-MS/MS, 5 µL sample was injected, andHRMS spectra of the total lipid extracts

from HeLa cells were recorded on a Q Exactive Orbitrap mass spectrometer (Thermo Fisher

Scientific) connected to a Vanquish Flex Analytical UHPLC System (Thermo Fisher Scientific)

equipped with a C8 column (100 x 2.1 mm, particle size 1.9 µm). Following HPLC method was

used for all runs using 0.1% formic acid in water (Solvent A) and 0.1% formic acid in methanol

(Solvent B): 0-1 min 30% B, 1-4 min 30%-90% B, 4-15 min 90-99% B, 15-18 min 99%B, 18-20

min 30% B. Lipid standards (POPC, N_3-DSPC) were supplied by Avanti Polar Lipids.

Image collection: Confocal micrographs were acquired on a Zeiss LSM 880 microscope

equipped with plan-apochromat 64x/1.4 NA or 20x/0.8 NA objectives. For far-red fluorogen-FAP

signals, a 633 nm HeNe laser was used at 1-2% power. FP or other fluorophores were excited

with 488 nm Argon (GFP), 405 nm (BFP/mCer3) or 561 nm (mCherry) diode lasers. All images

used for intensity quantification were acquired in LSM mode using a QUASAR GaAsP detector

using identical settings. SIM imaging of mitochondrial **1b**-PC was performed on an Zeiss Elyra 7

Lattice-SIM[2] system equipped with a plan-apochromat 100x/1.46 NA objective using 642 nm

and 488 nm diode lasers (10% power each). Images were acquired in ZEN Black using Weak

Live default settings with a pixel dwell time of 843 ms.

Image quantification: All intensity were calculated using ImageJ on unprocessed images. For

proof of concept experiments using transfected cells (Fig. 2), line profiles drawn were across 10

individual ER segments per cell and the total fluorescence ratio between fluorogen and GFP

emissions were tabulated across them. For per-cell data (Fig. 2d, 2f), the 10 segments were

averaged and 10 cells were analyzed per condition. Raw per segment fluorescence intensity is

shown for all 100 segments in Fig. 2e. For mitochondrial and ER quantification in Fig. 3, fields of

cells stably expressing to corresponding FAPs were analyzed by segmenting individual cells

and computing the total fluorescence ratio between fluorogen and GFP emission across the

entire cell, given the highly specific distribution of the signal for the compartment of interest. For

TGN cisternae quantification (Fig. 4, 5), line segments were drawn across the Golgi apparatus

of individual cells so that they did not intersect any TGN vesicles. For TGN vesicle

quantification, 5 line segments were drawn across Golgi vesicles that did not intersect Golgi

cisternae. For each compartment, the total fluorescence ratio between fluorogen and GFP were

calculated across each and averaged together for a single cell value. Values were measured

this way for 25 cells per condition (luminal vs. cytosolic for PC, PS, PI, and SiA). Mitochondrial

quantification in Fig. 5 was carried out identically as in Fig. 3. Images highlighting fluorogen

distribution (Figs. 2a, 2b, 3a, 3b, 4b, 4d; Supplementary Figs. 3a, 3b, 7, 9, 10; Supplementary

Movies 1, 2) were acquired with an Airyscan detector and processed using default settings in

Zen Black; these were not used for quantification due to non-linear processing. All statistical

analyses were performed in GraphPad Prism 9.

Gene knockdown: Pre-validated Silencer Select siRNAs were supplied by Thermo Fisher:

STARD7 (s32370), VPS13a (s531256), negative control 1 (4390843). Cells were transfected

with Lipofectamine RNAiMAX (ThermoFisher) complexed with 25 pmol siRNA for 48h prior to

analysis. Total protein was extracted using RIPA Lysis and Extraction buffer (ThermoFisher) and

measured by Bradford assay (Bio-Rad). A total 20 µg protein per extract was boiled in 1X

reducing sample buffer and used for SDS-PAGE (BioRad, 4561083). Validation of gene

knockdowns was performed by Western blotting using primary antibodies against STARD7

(Proteintech, 15689) 1:2500, VPS13a (Atlas antibodies, HPA021662) 1:1000, and Actin

(Invitrogen, PA5-78715) 1:5000. HRP-conjugated goat anti-rabbit IgG secondary antibody

(Abcam, 6721) 1:10,000 and Pierce ECL western blotting substrate (Thermo Fisher, 32209)

were used for detection.

Figure 1. Conceptual development of FACES

a, N_3-Cho is metabolically incorporated into azido-PC lipids distributed throughout all cellular membranes. **b,** Fluorogens form reversible non-covalent fluorescent complexes with genetically encoded FAPs. **c,** Structure of the fluorogen malachite green linked to dibenzocyclooctyne for SPAAC labeling. **d,** Conceptual schematic detailing the FACES method. (1) Cells expressing a FAP-GFP fusion protein targeted to an organelle of interest are metabolically labeled with N_3-Cho, or other bioorthogonal metabolite. (2) After incorporation, metabolites are conjugated via a SPAAC reaction with fluorogen-DBCO. During this time free dye can interact with the FAP to generate far-red fluorescence. (3) After SPAAC labeling, excess free dye is reversibly disassociated from the FAP complex and removed from cells by washing. (4) Subsequent imaging of the fluorogen-FAP complex is used to detect the specific biomolecule pool. Co-measurement of GFP fluorescence is used to normalize the fluorogen signal to FAP sensor density.

Figure 2. Imaging N$_3$-PC in the ER using compound 1a

a, b, HeLa cells transfected with either FAP-GFP-KDEL (top two rows) or GFP-KDEL (bottom

row) were grown in the presence (top and bottom rows) or absence (middle row) of 50 μM

N$_3$-Cho and labeled with 500 nM **1a**. Cells were imaged while in the labeling solution **(a)** and

after washing with culture medium **(b)**. In washed cells, fluorgen fluorescence is only visible

when both N$_3$-PC and the FAP domain are present, while in unwashed cells unreacted

fluorogens also fluoresce. Scale = 20 μm. **c,** Merged confocal images of transfected HeLa cells

post washing. White line indicates the segment used for plotting the fluorescent intensity profile

of **1a** and GFP shown below. Scale = 10 μm. **d,** Quantified fluorescence intensity profiles from

confocal images expressed as ratiometric average; n = 10 cells. **e,** Linear regression of

quantified fluorescence from transfected HeLa cells fed 0, 10, 30, 50 μM N$_3$-Cho and labeled

with 500 nM **1a**; n = 100 individual ER segments derived from 10 cells. **f,** Quantified

fluorescence of cells incubated with the PC synthesis inhibitor D609 alongside N$_3$-Cho; n = 10

cells.

Figure 3. Using FACES to analyze PC trafficking between the mitochondrion and the ER

a, HeLa cells co-transfected with an ER-localized FAP (FAP-GFP-KDEL) and a mitochondrial marker (mito-BFP) show fluorogen fluorescence in the far-red channel that does not overlap with mitochondria. **b,** HeLa cells co-transfected with a mitochondrial matrix FAP (MTS-FAP-GFP) and an ER marker (Sec61-mCherry) show fluorgen fluorescence that does not overlap with the ER. For both **(a)** and **(b)**, 50 µM N_3-Cho and 500 nM **1a** were used for PC labeling. Scale = 5 µM. **c,** Super-resolution live cell imaging of **1b**-PC (magenta) in the mitochondrial matrix with Elyra 7 lattice SIM[2]. The graph below indicates fluorescence intensity profiles of **1b**-PC (magenta) and GFP (green) along the white dashed line (inset). Scale = 2 µm and 0.5 µm, respectively. **d-f,** Knockdown of two putative ER-mitochondrial lipid transporters alters the distribution of PC between the two organelles. Stable cell lines expressing

MTS-FAP-GFP **(e)** and FAP-GFP-KDEL **(f)** were used to quantify N_3-PC levels in the mitochondrion (Mito) and ER, respectively; n = 70 cells per condition; ***, $p < 0.01$. Images corresponding to each condition are shown in Supplementary Fig. 5a.

Figure 4. Imaging lipid asymmetry generated at the TGN with FACES

a, Illustrated schematic representing the constructs used for reciprocal tagging of TGN38 with a FAP exposed to the cytosolic or luminal leaflet. The constructs also include the cyan fluorescent protein mCer3 on the opposing leaflet to account for sensor abundance. **b,** Example composite images of individual Golgi from HeLa cells stably expressing the cytosolic (top) or luminal

(bottom) TGN38 FAP sensors. Cells were treated with a tracer for PC (100 µM N_3-Cho), PI (50 µM Inos-N_3), PS (250 µM C-L-Ser-N_3), or Sia (50 µM Ac_4ManNAz). Images were acquired using 441 nm and 633 nm laser lines. Magenta regions are enriched in the fluorogen conjugated metabolite, green regions correspond to low metabolite levels, while white regions show intermediate values, relative to the sensor levels. **1a**-PC detection in the TGN using 100 µM N_3-Cho and 500 nM **1a**. Scale bar = 10 µm. **c**, Quantification of the fluorescence intensity ratio of fluorogen to mCer3 for each metabolite and each orientation (cisternae, TGN-C; vesicles, TGN-V). Values were calculated post-subtraction of the background fluorescence using unfed null controls. Error bars indicate SD; *n* = 25 cells per metabolite per leaflet. Raw values and statistical analyses are provided in Supplementary Fig. 7. **d**, Co-localization of SMeqt-GFP and cytoplasmic **1a**-PC in individual TGN vesicles, as detected by mCer3-TGN38-FAP. Here, 488 and 633 nm laser lines were used for imaging to avoid imaging of mCer3. The SMeqt-GFP channel is colored cyan and the **1a**-PC channel is colored magenta. Scale = 10 µm. **e,** Conceptual model representing the proposed asymmetric distribution of PC and SM lipids that is generated alongside TGN vesicle formation.

Fig. 5. Intracellular imaging of *N*-acetylhexosamine sugars with FACES

a, Illustration depicting the metabolic fate of Ac$_4$GalNAz (yellow square) and Ac$_4$GlcNAz (blue square) incorporation into cellular glycoconjugates containing *N*-acetylhexosamine sugars. Ac$_4$GalNAz fed to cells is converted to UDP-GalNAz in the cytosol and further epimerized to UDP-GlcNAz, thereby robustly labeling both UDP-HexNAz pools (UDP-GalNAz + UDP-GlcNAz). Conversely, Ac$_4$GlcNAz is not efficiently recycled to UDP-GlcNAz. **b**, UDP-HexNAz is incorporated into diverse extracellular glycoproteins proteins on the cell surface. Structures for an *N*-linked glycoprotein and *O*-linked mucin are shown as examples. **c**, UDP-HexNAz is incorporated into intracellular *O*-GlcNAcylated proteins. Protein glycosylation is catalyzed by *O*-GlcNAc Transferase (OGT) and can be reversed by *O*-GlcNAcase (OGA). OGA

is pharmacologically inhibited by Thiamet-G (TMG). **d,** Bioorthogonal imaging of HeLa cells fed 100 µM GalNAz for 24h and labeled with 5 µM TAMRA-DBCO (cell surface) and 5 µM BODIPY-DBCO (intracellular), relative to the mitochondrial marker Mito-BFP. Scale = 20 µm. **e-i**, Stable MTS-FAP-GFP cells detect mitochondrial HexNAz. **e**, Quantified mitochondrial HexNAz incorporation in cells fed either 100 µM Ac_4GalNAz or 100 µM Ac_4GlcNAz for 24h, versus unfed controls. Error bars indicate SD; n = 100 cells. Cells were labeled with 500 nM **1b** and washed as previously described. Representative images of Ac_4GalNAz fed cells and control are shown on the right; scale = 20 µm. **f**, Response of mitochondrial HexNAz to OGA inhibition using Thiamet-G (TMG). Cells were treated with 50 µM Thiamet-G (or DMSO control) and fed 100 µM Ac_4GalNAz for 24h. Displayed values are normalized to the average background fluorescence of unfed cells, which were used to control for **1b** labeling. Error bars indicate SD; n = 59 cells. **g**, Response of mitochondrial HexNAz to glucose availability. Cells were fed 100 µM Ac_4GalNAz and grown in DMEM containing either low (5.5 mM) or high (30 mM) glucose for 24h. Displayed values are normalized to unfed cells used as control for **1b** labeling. Error bars indicate SD; n = 63 cells. Representative LSM images (right) show **1b**-HexNAz fluorescent intensities displayed using the FireLUT. Scale = 50 µm.

Supplementary Figures

a

DBCO-C$_6$-Malachite Green (**1a**)

DBCO-PEG$_2$-Malachite Green (**1b**)

b

c

DBCO-C$_6$-Malachite Green (**1a**)

DBCO-PEG$_2$-Malachite Green (**1b**)

Supplementary Fig. 1. Synthesis of DBCO-linked fluorogens

a, Chemical structures of all the Malachite Green (Chromatic forms) derivatives used in this study. **b**, Synthesis of the chromatic forms of DBCO-C$_6$-Malachite Green (**1a**) and DBCO-PEG$_2$-Malachite Green (**1b**). **c**, HPLC/ELSD spectrum corresponding to the chromatic forms of DBCO-C$_6$-Malachite Green (**1a**) and DBCO-PEG$_2$-Malachite Green (**1b**). Retention times were verified by mass spectrometry.

Supplementary Fig. 2. N₃-Cho is metabolized into N₃-PC but not N₃-SM

Cells were fed 100 µM N_3-Cho for 24 h and extracted lipids were analyzed by LC-MS/MS. **a**, Comparative extracted ion chromatograms (EIC) from full scan/all ion fragmentation mode. MS2 fragments: choline (black), N_3-Cho (red). EIC of POPC (green) and N_3-POPC (blue) were detected as representative PC species in the sample. **b**, Sphingoid base fragment 264.269 was detected in neutral loss scan mode NL = 59 (choline containing lipids) but not in NL = 114 (N_3-Cho containing lipids), indicating N_3-Cho was not incorporated into SM in a detectable amount under these labeling conditions.

Supplementary Fig. 3. 1a-FAP complex reversibility depends on the heavy chain scFvs FAP clone dH6.2

Cells that were not fed with any azido metabolites were transfected with either dH6.2-GFP-KDEL or light chain scFvs FAP variant dK-GFP-KDEL and stained with 500 nM **1a**. dH6.2 was shown to have reversible binding to MG[15,59], while dK features nanomolar affinity[15,59] and is thus irreversibly bound under experimental conditions. **a**, Cells imaged while in staining solution show co-localization of sensor (GFP) and fluorogen (**1a**) signal for both constructs. Scale = 10 μm. **b**, Cells imaged after washing 3x with DMEM show only fluorogen signal for the dK construct (top), as FAP-bound 1a is not washed out. Line profiles show the relative change in **1a** fluorescence intensity compared to GFP. Specific detection of SPAAC-conjugated fluorogens thus requires a lower affinity FAP like dH6.2. Scale = 25 μm.

Supplementary Fig. 4. 1a-PC conjugates are formed in cells

a, HR-MS/MS analysis of **1a**-PC band from thin layer chromatography detecting $[M+H]^{2+}=747.4713$ and MS2 fragment 449.2146. Cells stained with **1a** were not washed prior to lipid extraction to highlight conversion of **1a** to **1a**-PC. **b,** Comparative HR-MS/MS analysis of the **1a**-DSPC standard previously synthesized by reaction of **1a** with N_3-DSPC detecting $[M+H]^{2+}=752.4949$ and MS2 fragment 449.2146.

Supplementary Fig. 5. Linear quantification of 1b-PC in the mitochondrial matrix

a, HeLa cells stably expressing MTS-FAP-GFP were fed different concentrations of N_3-Cho for 24h and stained with 500 nM **1b**. Images show that fluorogen fluorescence (middle) is dependent on azido metabolic incorporation. Scale = 20 μm. **b**, Quantification of **1b**/GFP fluorescence in cells expressing MTS-FAP-GFP with increasing concentrations of N_3-Cho. **c**, Linear increase of **1b**/GFP fluorescence only up to 100 μM N_3-Cho.

Supplementary Fig. 6. LTP silencing affects 1a-PC abundance in the mitochondria and ER
a-b, Representative LSM images of siRNA transfected cells stably expressing either
MTS-FAP-GFP **(a)** and FAP-GFP-KDEL **(b)** Null control (NC) siRNA shows robust **1a**-PC/**1b**-PC
signal in both compartments. Both STARD7 and VPS13a show reduction in mitochondrial
1b-PC signal, but only VPS13a knockdown shows a concomitant increase in ER **1a**-PC signal.
Scale = 50 µm. **c**, Silencing efficiency of the two siRNAs compared to the NC as determined by
western blotting.

Supplementary Fig. 7. TGN vesicle formation is associated with a loss of luminal PC

Golgi (top) and highlighted region (bottom) are shown from an individual cell were transfected with mCer3-FAP-TGN38, fed 100 µM N_3-Cho for 24h, and labeled with 500 nM **1a**. 3D rendering of Airyscan z-stacked images show a lack of **1a**-PC labeling on the luminal leaflet of vesicles (arrow, outline) and tubes (arrow, filled) connected to TGN cisternae. Airyscan images were acquired with 405 and 633 lasers for excitation and localization of mCer3 and **1a**-PC, respectively. Scale = 10 µm (top) and 2 µm (bottom).

Supplementary Fig. 8. FACES quantification of phospholipid asymmetry across the TGN

Cells stably expressing Tet inducible FAP-TGN38-GFP (luminal leaflet) or GFP-TGN38-FAP (cytosolic leaflet) constructs were grown under 1 μg/mL doxycycline and fed different azido metabolic labeling reagents for 24h prior to staining with 500 nM **1a**. **a,** cells fed 100 μM N_3-Cho. **b**, cells feld 50 μM Inos-N_3. **c**, cells fed 250 μM C-L-Ser-N_3. **d**, cells fed 50 μM Ac$_4$ManNAz. LSM images were used for quantification. n = 25 cells per condition.

Supplementary Fig. 9. Multicolor imaging of global and targeted phospholipid pools

Cells were metabolically labeled with 100 µM N_3-Cho, 250 µM C-L-Ser-N_3, or 50 µM Inos-N_3 for 24h prior to SPAAC conjugation. **a**, HeLa cells stained with 2 µM BODIPY-DBCO, which reacts with intracellular azido lipids. Images reflect maximal projection of Airyscan z-stacked images. Scale bars, 20 µm. **b**, Merged Airyscan images of HeLa cells stably expressing pTet:mCer3-TGN38-FAP double stained with equimolar (1 µM each) **1a**:BODIPY-DBCO. Scale bars, 20 µm **c**, Higher magnification images show overlapping phospholipid distributions between total lipids labeled with BODIPY and those labeled by **1a** and detected in the cytosolic leaflet of the TGN. For experiments **b-c**, 488 and 633 nm laser lines were used for imaging to avoid excitation of mCer3. Scale = 10 µm.

Supplementary Fig. 10. GPI-anchored proteins co-localize with cytoplasmic 1a-PC in TGN vesicles

Cells were cotransfected with mCherry-GPI and mCer3-TGN38-FAP constructs, fed 100 μM N$_3$-Cho for 24h, and labeled with 500 nM **1a.** Airyscan images were acquired using only 561 and 633 lasers for excitation of m-Cherry-GPI and **1a**-PC, respectively. Maximal projection of z-stacked images show co-localization between mCherry-GPI cargo and cytosolic facing **1a**-PC in TGN cisternae and vesicles (inset). Scale = 25 μm.

Supplementary Movie 1: Mobility of TGN vesicles containing cytosolic PC and luminal SM

Airyscan time-lapse of Golgi from the cell shown in Fig. 4d expressing GFP-EqtSM (cyan) and mCer3-TGN38-FAP detecting **1a**-PC detection in the cytoplasmic leaflet (magenta). Images reflect maximal intensity values from z-stacked images spanning 1.5 µm. Scale = 10 µm.

https://drive.google.com/file/d/1uWdM-LxW6N6hfmS_1oAvt9gelVmVm2tq/view?usp=sharing

Supplementary Movie 2: Mobility of TGN vesicles containing cytosolic PC and protein cargoes

Airyscan time-lapse of Golgi from the cell shown in Supplementary Fig. 9 expressing mCherry-SBP-GPI (yellow) and mCer3-TGN38-FAP detecting **1a**-PC detection in the cytoplasmic leaflet (magenta). Scale = 20 µm.

https://drive.google.com/file/d/1S8RHj0p4P12DSThlM9wqXF7-yQTGs2qc/view?usp=sharing

Supplementary Text: Synthesis of the Malachite Green Probes

Commercially available 4-nitrobenzaldehyde, *N,N*-dimethylaniline, niobium(V) chloride (NbCl$_5$), 10% Pd/C, O-(7-azabenzotriazol-1-yl)- 1,1,3,3-tetramethyluronium hexafluoro-phosphate (HATU), *N,N*-diisopropylethylamine (DIEA), tetrachloro-*p*-benzoquinone, and glacial acetic acid (AcOH) were obtained from Sigma-Aldrich. DBCO-C$_6$-CO$_2$H, azido-PEG$_2$-CH$_2$CO$_2$H, and DBCO-PEG$_2$-CO$_2$H were obtained from BroadPharm. Deuterated chloroform (CDCl$_3$) was obtained from Cambridge Isotope Laboratories. All reagents obtained from commercial suppliers were used without further purification unless otherwise noted. Analytical thin-layer chromatography was performed on E. Merck silica gel 60 F$_{254}$ plates. Silica gel flash chromatography was performed using E. Merck silica gel (type 60SDS, 230-400 mesh). Solvent mixtures for chromatography are reported as v/v ratios. HPLC analysis was carried out on an Eclipse Plus C8 analytical column with *Phase A*/*Phase B* gradients [*Phase A*: H$_2$O with 0.1% formic acid; *Phase B*: MeOH with 0.1% formic acid]. HPLC purification was carried out on Zorbax SB-C18 semipreparative column with *Phase A*/*Phase B* gradients [*Phase A*: H$_2$O with 0.1% formic acid; *Phase B*: MeOH with 0.1% formic acid]. Proton nuclear magnetic resonance (^{1}H NMR) spectra were recorded on a Jeol ECZL 400 MHz spectrometer, and were referenced relative to residual proton resonances in CDCl$_3$ (at d 7.26 ppm) or CD$_3$OD (at d 4.87 or 3.31 ppm). ^{1}H NMR splitting patterns are assigned as singlet (s), doublet (d), triplet (t), quartet (q) or pentuplet (p). All first-order splitting patterns were designated on the basis of the appearance of the multiplet. Splitting patterns that could not be readily interpreted are designated as multiplet (m) or broad (br). Carbon nuclear magnetic resonance (^{13}C NMR) spectra were recorded on a Jeol ECZL 400 MHz spectrometer, and were referenced relative to residual proton resonances in CDCl$_3$ (at d 77.16 ppm) or CD$_3$OD (at d 49.15 ppm). Electrospray Ionization-Time of Flight (ESI-TOF) spectra were obtained on an Agilent 6230 TOF-MS.

<u>4,4'-((4-nitrophenyl)methylene)bis(*N,N*-dimethylaniline) [O$_2$N-Malachite Green *(Leuco form)*]</u>[62,63].

A mixture of 4-nitrobenzaldehyde (0.755 g, 5.0 mmol), *N,N*-dimethylaniline (1.90 mL, 15.0

mmol), and NbCl$_5$ (0.135 g, 0.5 mmol) was heated at 130 °C for 2 h under Ar. Several color

changes were observed, from yellow to orange, and finally purple. Afterwards, the reaction

mixture was cooled to room temperature and diluted in EtOAc (20 mL) and H$_2$O (20 mL). The

aqueous layer was then extracted with EtOAc (3 × 10 mL), and the combined organic layers

were dried over anhydrous MgSO$_4$ and concentrated *in vacuo*. The crude was absorbed over

SiO$_2$, and then purified by flash column chromatography (10-20% EtOAc in hexanes), affording

1.486 g of O$_2$N-Malachite Green *(Leuco form)* as an orange solid [79%, Rf = 0.42 (20% EtOAc

in hexanes)]. ^{1}H NMR (CDCl$_3$, 400.13 MHz, d): 8.12 (d, *J* = 8.7 Hz, 2H, 2 × CH), 7.30 (d, *J* = 8.7

Hz, 2H, 2 × CH), 6.96 (d, *J* = 8.7 Hz, 4H, 4 × CH), 6.69 (d, *J* = 8.2 Hz, 4H, 4 × CH), 5.47 (s, 1H,

1 × CH), 2.94 (s, 12H, 4 × CH$_3$). ^{13}C NMR (CDCl$_3$, 100.61 MHz, d): 153.3, 149.2, 146.3, 131.1,

130.2, 130.0, 123.6, 112.8, 55.0, 40.8. MS (ESI-TOF) [m/z (%)]: 376 ([MH]$^+$, 34), 188 ([MH]$^{2+}$,

100).

<u>4,4'-((4-aminophenyl)methylene)bis(*N,N*-dimethylaniline) [H$_2$N-Malachite Green *(Leuco form)*]</u>. A

suspension of 4,4'-((4-nitrophenyl)methylene)bis(*N,N*-dimethylaniline) [O$_2$N-Malachite Green

(*Leuco form*)), 150.0 mg, 0.40 mmol] and 10% Pd/C (42.6 mg) in absolute EtOH (8 mL) was

stirred for 2 h under H$_2$. Then, the mixture was filtered through a pad of celite, washing with

EtOH (3 × 2 mL). Afterwards, the filtrate was concentrated *in vacuo*, affording 137.2 mg of

H$_2$N-Malachite Green (*Leuco form*) as a white solid [99%, Rf = 0.42 (20% EtOAc in hexanes)].

The product was used directly without any further purification. ^{1}H NMR (CDCl$_3$, 400.13 MHz, d):

7.00 (d, *J* = 8.7 Hz, 4H, 4 × CH), 6.92 (d, *J* = 8.3 Hz, 2H, 2 × CH), 6.69 (d, *J* = 8.7 Hz, 4H, 4 ×

CH), 6.61 (d, *J* = 8.3 Hz, 2H, 2 × CH), 5.29 (s, 1H, 1 × CH), 3.86 (br s, 2H, 1 × NH$_2$), 2.92 (s,

12H, 4 × CH$_3$). ^{13}C NMR (CDCl$_3$, 100.61 MHz, d): 148.8, 144.2, 135.8, 133.8, 130.2, 130.0,

115.2, 112.8, 54.2, 41.0. MS (ESI-TOF) [m/z (%)]: 346 ([MH]$^+$, 11), 173 ([MH]$^{2+}$, 100).

<u>DBCO-C$_6$-Malachite Green (*Leuco form*).</u> A solution of DBCO-C$_6$-CO$_2$H (19.3 µL, 57.9 mmol) in 100 µL of CH$_2$Cl$_2$ was stirred at 0 ºC for 10 min, and then HATU (24.2 mg, 63.7 mmol) and DIEA (40.4 µL, 231.6 µmol) were successively added. After 10 min stirring at 0 ºC, a H$_2$N-Malachite Green *(Leuco form)* (20.0 mg, 57.9 mmol) solution in 100 µL of CH$_2$Cl$_2$ containing DIEA (20.2 µL, 115.8 µmol) was added. After 1 h stirring at rt, the solvent was removed under reduced pressure to give a blue oil. The corresponding residue was dissolved in MeOH (400 µL), filtered using a 0.2 µm syringe-driven filter, and the crude solution was purified by HPLC, affording 29.2 mg of DBCO-C$_6$-Malachite Green *(Leuco form)* as a blue oil [72%, t$_R$ = 10.8 min (Zorbax SB-C18 semipreparative column, 50% *Phase A* in *Phase B*, 1 min, then 50-5% *Phase A* in *Phase B*, 5 min, and then 5% *Phase A* in *Phase B*, 10 min)]. ^{1}H NMR (CDCl$_3$, 400.13 MHz, d): 7.78 (s, 1H, 1 × NH), 7.72-7.64 (m, 1H, 1 × CH), 7.50-7.28 (m, 7H, 7 × CH), 7.25-7.18 (m, 4H, 4 × CH), 7.08-6.94 (m, 6H, 6 × CH), 6.93-6.72 (m, 2H, 2 × CH), 5.38 (s, 1H, 1 × CH), 5.17 (d, *J* = 13.8 Hz, 1H, 0.5 × CH$_2$), 3.69 (d, *J* = 13.8 Hz, 1H, 0.5 × CH$_2$), 2.96 (s, 12H, 4 × CH$_3$), 2.29 (dt, *J$_1$* = 16.0 Hz, *J$_2$* = 6.0 Hz, 1H, 0.5 × CH$_2$), 2.10 (dt, *J$_1$* = 17.1 Hz, *J$_2$* = 6.8 Hz, 1H, 0.5 × CH$_2$), 2.02-1.81 (m, 2H, 1 × CH$_2$), 1.62-1.40 (m, 3H, 1.5 × CH$_2$), 1.39-1.21 (m, 1H, 0.5 × CH$_2$). ^{13}C NMR (CDCl$_3$, 100.61 MHz, d): 173.8, 171.2, 151.7, 148.0, 132.5, 130.3, 129.8, 129.1, 128.6, 128.5, 128.4, 128.0, 127.3, 125.6, 123.1, 122.6, 119.9, 115.1, 108.0, 55.6, 54.7, 44.0, 36.8, 34.6, 25.1, 24.0. MS (ESI-TOF) [m/z (%)]: 661 ([MHI]$^+$, 5), 331 ([MH]$^{2+}$, 100).

<u>DBCO-C$_6$-Malachite Green *(Chromatic form)* (**1a**).</u> To a blue solution of DBCO-C$_6$-Malachite Green *(Leuco form)* (4.0 mg, 6.1 µmol) and tetrachloro-*p*-benzoquinone (1.8 mg, 7.3 µmol) in CHCl$_3$ (226 µL), glacial acetic acid (4.5 µL, 0.45 µmol) was added. The solution was stirred at 35 ºC for 2 h. Afterwards, the volatiles were removed under reduced pressure, obtaining a dark green residue. The crude was absorbed over SiO$_2$, and then purified by flash column chromatography (0-7% MeOH in CH$_2$Cl$_2$), affording 3.8 mg of DBCO-C$_6$-Malachite Green

(*Chromatic form*) as a dark green solid [90%, Rf = 0.57 (10% MeOH in CH_2Cl_2)]. ^1H NMR (CD$_3$OD, 400.13 MHz, d): 7.91-7.76 (m, 4H, 4 × CH), 7.71-7.64 (m, 1H, 1 × NH), 7.57-7.47 (m, 2H, 2 × CH), 7.44 (d, *J* = 9.0 Hz, 4H, 4 × CH), 7.37 (d, *J* = 8.7 Hz, 2H, 2 × CH), 7.33-7.19 (m, 4H, 4 × CH), 7.06 (d, *J* = 9.4 Hz, 4H, 4 × CH), 4.93-4.91 (m, 2H, 1 × CH$_2$), 3.33 (s, 12H, 4 × CH$_3$), 2.52-2.45 (m, 2H, 1 × CH$_2$), 2.44-2.34 (m, 2H, 1 × CH$_2$), 1.83-1.65 (m, 4H, 2 × CH$_2$). ^{13}C NMR (CD$_3$OD, 100.61 MHz, d): 178.9, 174.7, 158.4, 145.6, 141.9, 137.7, 135.8, 135.6, 134.2, 133.5, 130.8, 130.4, 130.0, 129.7, 129.2, 128.9, 128.8, 128.3, 126.5, 125.6, 124.3, 123.1, 120.7, 120.2, 114.8, 114.5, 114.2, 40.9, 37.7, 34.4, 33.1, 30.8, 26.0, 25.5. MS (ESI-TOF) [m/z (%)]: 660 ([MH -Cl]$^+$, 51), 659 ([M -Cl]$^+$, 100), 330 ([MH -Cl]$^{2+}$, 53).

DBCO-PEG$_2$-Malachite Green *(Leuco form)*. A solution of DBCO-PEG$_2$-CO$_2$H (13.4 mg, 28.9 mmol) in 200 µL of CH$_2$Cl$_2$/DMF (1:1) was stirred at 0 ºC for 10 min, and then HATU (12.1 mg, 31.8 mmol) and DIEA (20.2 µL, 115.8 µmol) were successively added. After 10 min stirring at 0 ºC, a H$_2$N-Malachite Green *(Leuco form)* (10.0 mg, 28.9 mmol) solution in 200 µL of CH$_2$Cl$_2$/DMF (1:1) containing DIEA (10.1 µL, 57.9 µmol) was added. After 1 h stirring at rt, the solvent was removed under reduced pressure to give a blue oil. The corresponding residue was dissolved in MeOH (200 µL), filtered using a 0.2 µm syringe-driven filter, and the crude solution was purified by HPLC, affording 13.2 mg of DBCO-PEG$_2$-Malachite Green *(Leuco form)* as a blue oil [55%, t_R = 50.9 min (Zorbax SB-C18 semipreparative column, 95% *Phase A* in *Phase B*, 1 min, then 95-50% *Phase A* in *Phase B*, 30 min, and then 50-0% *Phase A* in *Phase B*, 22 min)]. ^1H NMR (CDCl$_3$, 400.13 MHz, d): 8.76 (s, 1H, 1 × NH), 7.60-7.54 (m, 1H, 1 × CH), 7.53-7.46 (m, 1H, 1 × CH), 7.46-7.34 (m, 6H, 6 × CH), 7.25-7.17 (m, 4H, 4 × CH), 7.09-6.96 (m, 6H, 6 × CH), 6.95-6.72 (m, 2H, 2 × CH), 6.28 (t, *J* = 5.6 Hz, 1H, 1 × NH), 5.36 (s, 1H, 1 × CH), 5.09 (d, *J* = 13.9 Hz, 1H, 0.5 × CH$_2$), 3.87-3.71 (m, 2H, 1 × CH$_2$), 3.69-3.56 (m, 5H, 2.5 × CH$_2$), 3.45 (t, *J* = 5.4 Hz, 2H, 1 × CH$_2$), 3.34-3.18 (m, 2H, 1 × CH$_2$), 2.95 (s, 12H, 4 × CH$_3$), 2.87-2.75 (m, 1H, 0.5 × CH$_2$), 2.53-2.41 (m, 2H, 1 × CH$_2$), 2.39-2.26 (m, 1H, 0.5 × CH$_2$), 2.17-2.02 (m, 1H,

0.5 × CH$_2$), 1.98-1.83 (m, 1H, 0.5 × CH$_2$). ^{13}C NMR (CDCl$_3$, 100.61 MHz, d): 172.7, 172.1,

169.9, 151.3, 148.1, 132.3, 130.3, 129.8, 129.5, 128.9, 128.4, 128.2, 127.9, 127.2, 125.6, 123.2,

122.6, 119.9, 114.8, 108.0, 70.7, 69.9, 69.8, 67.4, 55.8, 54.8, 39.1, 38.0, 31.5, 30.3. MS

(ESI-TOF) [m/z (%)]: 793 ([MH]$^+$, 4), 397 ([MH]$^{2+}$, 100).

<u>DBCO-PEG$_2$-Malachite Green *(Chromatic form)* (**1b**)</u>. To a blue solution of

DBCO-PEG$_2$-Malachite Green *(Leuco form)* (5.0 mg, 6.3 µmol) and tetrachloro-*p*-benzoquinone

(1.9 mg, 7.6 µmol) in CHCl$_3$ (235 µL), glacial acetic acid (4.7 µL, 0.47 µmol) was added. The

solution was stirred at 35 °C for 2 h. Afterwards, the volatiles were removed under reduced

pressure, obtaining a dark green residue. The crude was absorbed over SiO$_2$, and then purified

by flash column chromatography (0-10% MeOH in CH$_2$Cl$_2$), affording 4.8 mg of

N$_3$-PEG$_2$-Malachite Green *(Chromatic form)* as a dark green solid [92%, Rf = 0.44 (10% MeOH

in CH$_2$Cl$_2$)]. ^{1}H NMR (CDCl$_3$, 400.13 MHz, d): 11.33 (s, 1H, 1 × NH), 8.20 (d, *J* = 8.4 Hz, 2H, 2 ×

CH), 7.85-7.69 (m, 1H, 1 × NH), 7.63 (d, *J* = 7.3 Hz, 1H, 1 × CH), 7.58 (d, *J* = 7.6 Hz, 1H, 1 ×

CH), 7.42-7.27 (m, 9H, 9 × CH), 7.25-7.17 (m, 3H, 3 × CH), 6.86 (d, *J* = 8.5 Hz, 4H, 4 × CH),

5.13 (d, *J* = 13.9 Hz, 1H, 0.5 × CH$_2$), 3.91 (t, *J* = 5.9 Hz, 2H, 1 × CH$_2$), 3.76-3.57 (m, 5H, 2.5 ×

CH$_2$), 3.56-3.45 (m, 2H, 1 × CH$_2$), 3.31 (s, 12H, 4 × CH$_3$), 2.94 (t, *J* = 5.9 Hz, 2H, 1 × CH$_2$),

2.87-2.75 (m, 1H, 0.5 × CH$_2$), 2.68-2.53 (m, 1H, 0.5 × CH$_2$), 2.46-2.26 (m, 4H, 2 × CH$_2$). ^{13}C

NMR (CDCl$_3$, 100.61 MHz, d): 175.7, 174.4, 172.4, 156.5, 146.3, 141.1, 137.4, 134.2, 133.4,

133.3, 132.3, 129.3, 127.7, 127.6, 125.8, 124.9, 124.4, 122-1, 121.9, 120.0, 119.9, 117.3, 113.6,

133.5, 98.1, 98.0, 70.4, 70.3, 68.9, 67.3, 47.0, 41.2, 40.4, 38.0, 32.1, 29.9, 29.5, 22.9*, 14.3*.

MS (ESI-TOF) [m/z (%)]: 791 ([MH -Cl]$^+$, 3), 396 ([MH -Cl]$^{2+}$, 100). *Artifact.

¹H (left) and ¹³C NMR (right) spectra of synthesized compounds and key intermediates

References

1. van Meer, G., Voelker, D. R. & Feigenson, G. W. Membrane lipids: where they are and how they behave. *Nat. Rev. Mol. Cell Biol.* **9**, 112–124 (2008).

2. Levental, I. & Lyman, E. Regulation of membrane protein structure and function by their lipid nano-environment. *Nat. Rev. Mol. Cell Biol.* **24**, 107–122 (2023).

3. Lorent, J. H. *et al.* Plasma membranes are asymmetric in lipid unsaturation, packing and protein shape. *Nat. Chem. Biol.* **16**, 644–652 (2020).

4. Jao, C. Y., Roth, M., Welti, R. & Salic, A. Metabolic labeling and direct imaging of choline phospholipids in vivo. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 15332–15337 (2009).

5. Ancajas, C. F. *et al.* Cellular labeling of phosphatidylserine using clickable Serine probes. *ACS Chem. Biol.* **18**, 377–384 (2023).

6. Ricks, T. J. *et al.* Labeling of Phosphatidylinositol Lipid Products in Cells through Metabolic Engineering by Using a Clickable myo-Inositol Probe. *Chembiochem* **20**, 172–180 (2019).

7. Bumpus, T. W. & Baskin, J. M. A Chemoenzymatic Strategy for Imaging Cellular Phosphatidic Acid Synthesis. *Angew. Chem. Int. Ed Engl.* **55**, 13155–13158 (2016).

8. Bussink, A. P. *et al.* N-azidoacetylmannosamine-mediated chemical tagging of gangliosides. *J. Lipid Res.* **48**, 1417–1421 (2007).

9. Tamura, T. *et al.* Organelle membrane-specific chemical labeling and dynamic imaging in living cells. *Nat. Chem. Biol.* **16**, 1361–1367 (2020).

10. Maekawa, M., Yang, Y. & Fairn, G. D. Perfringolysin O Theta Toxin as a Tool to Monitor the Distribution and Inhomogeneity of Cholesterol in Cellular Membranes. *Toxins* **8**, (2016).

11. Deng, Y., Rivera-Molina, F. E., Toomre, D. K. & Burd, C. G. Sphingomyelin is sorted at the trans Golgi network into a distinct class of secretory vesicle. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6677–6682 (2016).

12. Gallo, E. Fluorogen-Activating Proteins: Next-Generation Fluorescence Probes for

Biological Research. *Bioconjug. Chem.* **31**, 16–27 (2020).

13. Kozma, E. & Kele, P. Fluorogenic probes for super-resolution microscopy. *Org. Biomol.*

Chem. **17**, 215–233 (2019).

14. Bottone, S. *et al.* A fluorogenic chemically induced dimerization technology for controlling,

imaging and sensing protein proximity. *Nat. Methods* **20**, 1553–1562 (2023).

15. Yan, Q. *et al.* Localization microscopy using noncovalent fluorogen activation by genetically

encoded fluorogen-activating proteins. *Chemphyschem* **15**, 687–695 (2014).

16. Horibata, Y. & Sugimoto, H. Differential contributions of choline phosphotransferases CPT1

and CEPT1 to the biosynthesis of choline phospholipids. *J. Lipid Res.* **62**, 100100 (2021).

17. Ng, M. N. P., Kitos, T. E. & Cornell, R. B. Contribution of lipid second messengers to the

regulation of phosphatidylcholine synthesis during cell cycle re-entry. *Biochim. Biophys.*

Acta **1686**, 85–99 (2004).

18. Adibhatla, R. M., Hatcher, J. F. & Gusain, A. Tricyclodecan-9-yl-xanthogenate (D609)

mechanism of actions: a mini-review of literature. *Neurochem. Res.* **37**, 671–679 (2012).

19. Larsen, E. C., Hatcher, J. F. & Adibhatla, R. M. Effect of tricyclodecan-9-yl potassium

xanthate (D609) on phospholipid metabolism and cell death during oxygen-glucose

deprivation in PC12 cells. *Neuroscience* **146**, 946–961 (2007).

20. Stephan, T., Roesch, A., Riedel, D. & Jakobs, S. Live-cell STED nanoscopy of

mitochondrial cristae. *Sci. Rep.* **9**, 12419 (2019).

21. John Peter, A. T., Petrungaro, C., Peter, M. & Kornmann, B. METALIC reveals

interorganelle lipid flux in live cells by enzymatic mass tagging. *Nat. Cell Biol.* **24**, 996–1004

(2022).

22. Kumar, N. *et al.* VPS13A and VPS13C are lipid transport proteins differentially localized at

ER contact sites. *J. Cell Biol.* **217**, 3625–3639 (2018).

23. Horibata, Y. *et al.* StarD7 Protein Deficiency Adversely Affects the Phosphatidylcholine

Composition, Respiratory Activity, and Cristae Structure of Mitochondria. *J. Biol. Chem.*

291, 24880–24891 (2016).

24. Horibata, Y. & Sugimoto, H. StarD7 mediates the intracellular trafficking of phosphatidylcholine to mitochondria. *J. Biol. Chem.* **285**, 7358–7365 (2010).

25. Horibata, Y. *et al.* Identification of the N-terminal transmembrane domain of StarD7 and its importance for mitochondrial outer membrane localization and phosphatidylcholine transfer. *Sci. Rep.* **7**, 8793 (2017).

26. Rothman, J. E. & Lenard, J. Membrane asymmetry. *Science* **195**, 743–753 (1977).

27. Verkleij, A. J. *et al.* The asymmetric distribution of phospholipids in the human red cell membrane. A combined study using phospholipases and freeze-etch electron microscopy. *Biochim. Biophys. Acta* **323**, 178–193 (1973).

28. Bos, K., Wraight, C. & Stanley, K. K. TGN38 is maintained in the trans-Golgi network by a tyrosine-containing motif in the cytoplasmic domain. *EMBO J.* **12**, 2219–2228 (1993).

29. Luzio, J. P. *et al.* Identification, sequencing and expression of an integral membrane protein of the trans-Golgi network (TGN38). *Biochem. J* **270**, 97–102 (1990).

30. Iyoshi, S. *et al.* Asymmetrical distribution of choline phospholipids revealed by click chemistry and freeze-fracture electron microscopy. *ACS Chem. Biol.* **9**, 2217–2222 (2014).

31. Cabukusta, B. *et al.* A lipid transfer protein knockout library reveals ORP9-ORP11 dimer mediating PS/PI(4)P exchange at the ER-trans Golgi contact site to promote sphingomyelin synthesis. *bioRxiv* 2023.06.02.543249 (2023) doi:10.1101/2023.06.02.543249.

32. Phillips, S. E., Ile, K. E., Boukhelifa, M., Huijbregts, R. P. H. & Bankaitis, V. A. Specific and nonspecific membrane-binding determinants cooperate in targeting phosphatidylinositol transfer protein beta-isoform to the mammalian trans-Golgi network. *Mol. Biol. Cell* **17**, 2498–2512 (2006).

33. Fairn, G. D. *et al.* High-resolution mapping reveals topologically distinct cellular pools of phosphatidylserine. *J. Cell Biol.* **194**, 257–275 (2011).

34. Saxon, E. & Bertozzi, C. R. Cell surface engineering by a modified Staudinger reaction.

Science **287**, 2007–2010 (2000).

35. Fleischer, B. Mechanism of glycosylation in the Golgi apparatus. *J. Histochem. Cytochem.*

31, 1033–1040 (1983).

36. Klemm, R. W. *et al.* Segregation of sphingolipids and sterols during formation of secretory

vesicles at the trans-Golgi network. *J. Cell Biol.* **185**, 601–612 (2009).

37. Doktorova, M. *et al.* Cell Membranes Sustain Phospholipid Imbalance Via Cholesterol

Asymmetry. *bioRxiv* 2023.07.30.551157 (2023) doi:10.1101/2023.07.30.551157.

38. Boyce, M. *et al.* Metabolic cross-talk allows labeling of O-linked

β-*N*-acetylglucosamine-modified proteins via the *N*-acetylgalactosamine salvage pathway.

Proceedings of the National Academy of Sciences **108**, 3141–3146 (2011).

39. Hang, H. C., Yu, C., Kato, D. L. & Bertozzi, C. R. A metabolic labeling approach toward

proteomic analysis of mucin-type O-linked glycosylation. *Proc. Natl. Acad. Sci. U. S. A.* **100**,

14846–14851 (2003).

40. Ma, Z. & Vosseller, K. Cancer Metabolism and Elevated O-GlcNAc in Oncogenic Signaling*.

J. Biol. Chem. **289**, 34457–34465 (2014).

41. Tan, E. P. *et al.* Sustained O-GlcNAcylation reprograms mitochondrial function to regulate

energy metabolism. *J. Biol. Chem.* **292**, 14940–14962 (2017).

42. Zhu, Q. *et al.* O-GlcNAcylation promotes tumor immune evasion by inhibiting PD-L1

lysosomal degradation. *Proceedings of the National Academy of Sciences* **120**,

e2216796120 (2023).

43. Pekkurnaz, G., Trinidad, J. C., Wang, X., Kong, D. & Schwarz, T. L. Glucose regulates

mitochondrial motility via Milton modification by O-GlcNAc transferase. *Cell* **158**, 54–68

(2014).

44. Dupas, T. *et al.* An overview of tools to decipher O-GlcNAcylation from historical

approaches to new insights. *Int. J. Biochem. Cell Biol.* **151**, 106289 (2022).

45. Onodera, Y., Nam, J.-M. & Bissell, M. J. Increased sugar uptake promotes oncogenesis via

EPAC/RAP1 and O-GlcNAc pathways. *J. Clin. Invest.* **124**, 367–384 (2014).

46. Jóźwiak, P. *et al.* Mitochondrial O-GlcNAc Transferase Interacts with and Modifies Many Proteins and Its Up-Regulation Affects Mitochondrial Function and Cellular Energy Homeostasis. *Cancers* **13**, (2021).

47. Ma, J. *et al.* O-GlcNAcomic Profiling Identifies Widespread O-Linked β-N-Acetylglucosamine Modification (O-GlcNAcylation) in Oxidative Phosphorylation System Regulating Cardiac Mitochondrial Function. *J. Biol. Chem.* **290**, 29141–29153 (2015).

48. Guzmán, L. E. *et al.* Bioorthogonal Metabolic Labeling of the Virulence Factor Phenolic Glycolipid in Mycobacteria. *ACS Chem. Biol.* (2024) doi:10.1021/acschembio.3c00724.

49. Fantoni, N. Z., El-Sagheer, A. H. & Brown, T. A Hitchhiker's Guide to Click-Chemistry with Nucleic Acids. *Chem. Rev.* **121**, 7122–7154 (2021).

50. Liang, D. *et al.* A real-time, click chemistry imaging approach reveals stimulus-specific subcellular locations of phospholipase D activity. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 15453–15462 (2019).

51. Nguyen, S. S. & Prescher, J. A. Developing bioorthogonal probes to span a spectrum of reactivities. *Nat Rev Chem* **4**, 476–489 (2020).

52. Gautier, A. Fluorescence-Activating and Absorption-Shifting Tags for Advanced Imaging and Biosensing. *Acc. Chem. Res.* **55**, 3125–3135 (2022).

53. Muslinkina, L. *et al.* Structure-Based Rational Design of Two Enhanced Bacterial Lipocalin Blc Tags for Protein-PAINT Super-resolution Microscopy. *ACS Chem. Biol.* **15**, 2456–2465 (2020).

54. Wong, L. H., Gatta, A. T. & Levine, T. P. Lipid transfer proteins: the lipid commute via shuttles, bridges and tubes. *Nat. Rev. Mol. Cell Biol.* **20**, 85–101 (2019).

55. Sakuragi, T. & Nagata, S. Regulation of phospholipid distribution in the lipid bilayer by flippases and scramblases. *Nat. Rev. Mol. Cell Biol.* **24**, 576–596 (2023).

56. Surma, M. A., Klose, C. & Simons, K. Lipid-dependent protein sorting at the trans-Golgi network. *Biochim. Biophys. Acta* **1821**, 1059–1067 (2012).

57. Sebastian, T. T., Baldridge, R. D., Xu, P. & Graham, T. R. Phospholipid flippases: building asymmetric membranes and transport vesicles. *Biochim. Biophys. Acta* **1821**, 1068–1077 (2012).

58. Sharpe, H. J., Stevens, T. J. & Munro, S. A comprehensive comparison of transmembrane domains reveals organelle-specific properties. *Cell* **142**, 158–169 (2010).

59. Carpenter, M. A. *et al.* Protein Proximity Observed Using Fluorogen Activating Protein and Dye Activated by Proximal Anchoring (FAP-DAPA) System. *ACS Chem. Biol.* **15**, 2433–2443 (2020).

60. Telmer, C. A. *et al.* Rapid, specific, no-wash, far-red fluorogen activation in subcellular compartments by targeted fluorogen activating proteins. *ACS Chem. Biol.* **10**, 1239–1246 (2015).

61. Kowarz, E., Löscher, D. & Marschalek, R. Optimized Sleeping Beauty transposons rapidly generate stable transgenic cell lines. *Biotechnol. J.* **10**, 647–653 (2015).

62. Montagut, A. M., Gálvez, E., Shafir, A., Sebastián, R. M. & Vallribera, A. Triarylmethane Dyes for Artificial Repellent Cotton Fibers. *Chemistry* **23**, 3810–3814 (2017).

63. Kim, J., Park, E. Y., Moon, S.-Y., Kim, J.-M. & Park, K. Patterned color images with a triphenylmethane-derived acrylate polymer. *Macromol. Res.* **16**, 81–83 (2008).