A novel induced pluripotent stem cell model of Schwann cell differentiation reveals *NF2-related gene regulatory networks of the extracellular matrix*

Olivia Lazaro[1,*], Sihong Li[1,2,*], William Carter[1], Oluwamayowa Awosika[1], Sylvia Robertson[1], Brooke E. Hickey[3], Steven P. Angus[3,4,5], Austin House[1], Wade D. Clapp[3,4,7], Abdul S. Qadir[1,#], Travis S. Johnson[1,3,4,6,#], Steven D. Rhodes[3,4,7,8#]

[1]Indiana Bioscience Research Institute

[2]Indiana University School of Public Health

[3]Department of Pediatrics, Herman B Wells Center for Pediatric Research, Indiana University School of Medicine, Indianapolis, IN, USA

[4]Melvin and Bren Simon Comprehensive Cancer Center, Indiana University School of Medicine, Indianapolis, IN USA

[5]Department of Pharmacology and Toxicology, Indiana University School of Medicine, Indianapolis, IN, USA

[6]Center for Computational Biology and Bioinformatics

[7]Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN, USA

[8]Division of Pediatric Hematology/Oncology/Stem Cell Transplant, Indiana University School of Medicine, Indianapolis, IN, USA

[#]**Corresponding authors:** Abdul S. Syed (asyed@indianabiosciences.org), Travis S. Johnson (johnstrs@iu.edu), Steven D. Rhodes (sdrhodes@iu.edu)

Abstract: Schwann cells are vital to development and maintenance of the peripheral nervous

system and their dysfunction has been implicated in a range of neurological and neoplastic

disorders, including *NF2*-related schwannomatosis. We developed a novel human induced

pluripotent stem cell (hiPSC) model to study Schwann cell differentiation in health and disease.

We performed transcriptomic, immunofluorescence, and morphological analysis of hiPSC

derived Schwann cell precursors (SPCs) and terminally differentiated Schwann cells (SCs)

representing distinct stages of development. To validate our findings, we performed integrated,

cross-species analyses across multiple external datasets at bulk and single cell resolution. Our

hiPSC model of Schwann cell development shared overlapping gene expression signatures with

human amniotic mesenchymal stem cell (hAMSCs) derived SCs and *in vivo* mouse models, but

also revealed unique features that may reflect species-specific aspects of Schwann cell biology.

Moreover, we identified gene co-expression modules that are dynamically regulated during

hiPSC to SC differentiation associated with ear and neural development, cell fate determination,

the *NF2* gene, and extracellular matrix (ECM) organization. By cross-referencing results

between multiple datasets, we identified new genes potentially associated with *NF2* expression.

Our hiPSC model further provides a tractable platform for studying Schwann cell development in

the context of human disease.

Keywords: Schwann cells, NF2, Schwannoma, hiPSC, RNA-seq, scRNA-seq, gene co-

expression analysis, systems biology, extracellular matrix

Abbreviations:

hiPSC	Human induced pluripotent stem cells
SC	Schwann cells
SPC	Schwann cell precursors
iSC	Immature SCs
mSC	Myelinating SCs
tSC	Transition SCs
pmSC	Pro-myelinating SCs
nmSC	Not-myelinating SCs
nm(R)SC	Non-myelinating (Remak) SCs

INTRODUCTION

Schwann cells are glial cells that originate in the neural crest and play a crucial role in myelinating peripheral nerves[1]. Myelination enhances the speed and efficiency of nerve conduction by forming a single spiraling myelin sheath around an axon [2]. Non-myelinating Schwann cells also provide unmyelinated axons with trophic support and mechanical cushioning [3]. Deregulated Schwann cell development can give rise to tumors of the peripheral and cranial nerves called schwannomas, which can have unpredictable clinical behavior, significant morbidities, and exceedingly limited therapeutic options available for their treatment [4].

Vestibular schwannomas are tumors that involve the eight cranial nerves, which are critical for hearing and balance. Affected individuals often suffer significant morbidities including tinnitus, sensorineural hearing loss, and impairment of balance and coordination [5]. Most vestibular schwannomas occur unilaterally; however, approximately 10% of cases involve both sides and are frequently associated with an autosomal dominant, heritable genetic condition called *NF2*-related schwannomatosis (*NF2*-SWN, formally known as neurofibromatosis type 2 [5, 6]. *NF2*-SWN is caused by pathogenic variants in the *NF2* tumor suppressor gene, which encodes Merlin, a cytoskeletal adaptor protein with diverse biochemical functions [7]. The development of bilateral vestibular schwannomas is a hallmark of *NF2*-SWN [8].

To better understand the molecular mechanisms underlying schwannoma genesis in *NF2*-SWN, the development of new model systems for normal and diseased Schwann cells is imperative. Human induced pluripotent stem cells (hiPSCs) have demonstrated promise in modeling a multitude of neurological diseases and cell types, including but not limited to hiPSC-derived astrocytes [9-12], microglia [13-15], and neurons [16, 17]. Direct conversion of human fibroblasts to hiPSC-deviled neurons has also been achieved [17]. These iPSC-derived cells have been employed to interrogate neuropsychiatric diseases [18-20], neurodegenerative

diseases [13, 20, 21], and gliomas [22-24]. However, beyond gliomas, considerably less

progress has been made in the field of iPSC modeling of peripheral nervous system derived

tumors including schwannoma [25].

To address this unmet need, we have developed a novel hiPSC Schwann cell model that can

be used to study Schwann cell differentiation in health and disease. Here, we performed

transcriptomic profiling of hiPSCs at multiple stages of Schwann cell differentiation to unveil

broad networks of differentially expressed genes (DEGs) associated with Schwann cell lineage

maturation and compare them to developmental stages outlined in other studies (**Table 1**). To

further characterize complex gene regulatory networks (GRNs) associated with Schwann cell

differentiation, we employed a systems biology approach, utilizing weighted gene co-expression

network analysis (WGCNA) to identify modules of transcriptionally associated genes [26] and to

study their temporal dynamics in unison. We further refine the putative roles of these modules

using functional enrichment analysis [27]. To validate our findings, we compared the DEGs and

gene modules derived from our hiPSC induced Schwann cells with two independent single cell

and bulk transcriptomic datasets of human Schwann cell differentiation [28] and mouse sciatic

nerve development [29].

METHODS

hiPSC generation

hiPSC lines were generated from neonatal human dermal fibroblasts (HDFn) from Invitrogen

(Catalog #: C-004-5C) using an Epi5™ Episomal iPSC Reprogramming Kit (ThermoFisher

Scientific) according to manufacturer's protocol. Transduced cells were plated on growth factor

reduced Matrigel (Corning, 356231) coated culture dishes in the presence of ROCK inhibitor and

maintained in Essential 8 (E8) medium. iPSC colonies were mechanically isolated and expanded

until at least passage 12 before differentiation. iPSCs were characterized by staining with

pluripotency markers and embryoid body (EB) formation. *Mycoplasma* testing was performed

prior to Schwann cell differentiation.

Schwann cell differentiation from hiPSCs

The human iPSC line IBRI.101.J was differentiated into Schwann cells using a six-stage

protocol over 31 days as described previously [30]. To start differentiation, the IBRi 101.J hiPSC

line was seeded on Matrigel-coated culture plate in E8 media. When the cells reached 50-60%

confluency, the culture medium was replaced with Schwann cell basal 1 (SCB1) media (NDM

containing 1x N2, 1x B27, 0.005% BSA, 2 mM GlutaMAX, 0.11 mM b-mercaptoethanol in

advanced DMEM/F12 and Neurobasal medium (1:1 mix)) with 3 µM CHIR and 20 µM

SB431542 for 6 days. On day 7, cells were fed with SCB1 media with 3 µM CHIR, 20 µM

SB431542 and 50 ng/mL neuregulin 1 (NRG1) through day 24. SCB1 media was replaced with

fresh medium daily, and the cells were routinely dissociated and expanded once they reached

80% confluence with Accutase treatment. On day 25, media was switched to Schwann cell

basal 2 (SCB2) media (DMEM, 1x NEAA, 1x BME and 2% FBS) with 200 ng/mL NRG1, 4

ng/mL Forskolin, 100 nM RA-all Trans and 10 ng/mL platelet derive growth factor subunit B

(PDGF-BB) for 3 days, then to SCB2 media with 200 ng/mL NRG1 and 10 ng/mL PDGF-BB for

2 days, and finally to SCB2 media with 200 ng/mL NRG1 for 2 more days. By day 31, terminal

differentiation to Schwann-like cells was completed. Samples were collected at days 0, 3, 6, 12,

18, 24 and at a uniform 70% confluency in all cases. Cell pellets were collected, and flash

frozen at -80°C from three wells for subsequent RNA extraction and RNA-seq, while one well

was retained for continued passaging to the next stage.

Immunoflourescence staining of hiPSC-derived Schwann cell precursors (SCPs) and SCs

We used immunofluorescence (IF) analysis to characterize hiPSC differentiation to SCPs and

SCs [31]. For immunostaining, cells were fixed with 4% Paraformaldehyde for 10 minutes. The

fixed cells were washed with PBS three times and then blocked and permeabilized with 0.3%

Triton X-100 (Sigma) in 1% Bovine Serum Albumin (Life Technologies) and 10% FBS for 1 hour

at room temperature. Cells were then incubated with primary antibodies 1:200 dilution of S100β

(Abcam #AB52642), 1:50 dilution of p75 (Abcam #AB3125), 1:100 dilution of HOXb7 (Santa

Cruz #sc-81292), and 1:500 dilution of GAP43 (Abcam #AB12274) at 4 °C overnight, then with

conjugated secondary antibodies at 1:500 dilution for 1 hour at room temperature. The

coverslips were mounted on slides with DAPI to stain the nuclei (VectorLabs). Images were

captured with an LSM 780 confocal microscope running Zen Black software.

Immunocytochemistry analysis confirmed the expression of S100β, p75, HOXb7, and GAP43.

RNA extraction, library preparation and sequencing

RNA was extracted using phenol-chloroform extraction with TRIzol LS Reagent (Life

Technologies Corporation, #10296028). 1mL of TRIzol LS Reagent was added to flash frozen

cell pellets followed by 0.2 mL of chloroform. Tubes were shaken by hand for 15 seconds and

incubated at room temperature for 2-3 minutes. Centrifugation was performed on samples at

4°C for 15 minutes at 13,200 rpm. The aqueous phase was transferred to an RNase-free tube

and an equal amount of 70% ethanol was added to each sample for a final concentration of

35%. Samples were vortexed and binding and washing of RNA was carried out according to the

instructions of Invitrogen's PureLink™ RNA Micro Scale Kit #12183016. Samples were eluted in

20 µL RNase-free water. The quality and quantity of RNA was determined using a NanoDrop™

Spectrophotometer (Thermo Scientific).

A total of 21 samples, 3 replicates for each of the 7 timepoints, were shipped overnight on dry

ice to Novogene for quality control, library preparation and RNA-sequencing as further

described below.

Initial processing of RNA-sequencing data

CASAVA (Consensus Assessment of Sequence And Variation) was used for base recognition

through the Illumina Genome Analyzer. Raw reads were removed if they contained adapter

sequences, greater than 10% undetermined bases, or more than 50% low-quality bases (quality

scores less than 5). Sequenced reads were then aligned to the GRCh38 Genome using HISAT2

v2.0.5 and quantified with feature counts v1.5.0-p3. FPKM values were calculated from the read

counts and CDS lengths for each gene across all timepoints and replicates.

Differential Gene Expression analysis

Differential gene expression analysis was performed with edgeR v3.22.5. The $\log_2$ fold change

(Log_2FC) was calculated between groups and the p-values (P) were corrected using the

Benjamini & Hochberg method (BH) to account for the false discovery rate (FDR). Each

timepoint (day 3, 6, 12, 18, 24, 31) was compared to baseline (day 0), resulting in 6 pairwise

comparisons. DEGs with $\text{Log}_2\text{FC} > 5.0$ and FDR < 0.001 were identified for each comparison.

Functional enrichment analysis was performed on DEGs from each comparison using

clusterProfiler v3.8.1 to test the statistical enrichment for genes within the Reactome, DO

(Disease Ontology), and DisGeNET databases. Terms with FDR < 0.05 were considered

enriched. Functional enrichment on other gene sets derived from this analysis was performed

using ToppGene [32].

Network Construction and Module Detection with WGCNA and lmQCM

Statistical analysis was performed using R (v4.1.1.). Genes with the lowest 10% mean FPKM

across all timepoints were removed. Ensembl IDs for the filtered genes were converted to

HGNC gene symbols using biomaRt (v2.52.0) and the Ensembl gene IDs from the Genome

assembly GRCh38.p14. For HGNC symbols with multiple corresponding Ensembl IDs, the

Ensembl ID with the highest average FPKM was used. WGCNA module generation was

conducted with an unsigned Pearson correlation coefficient (PCC) matrix. Soft thresholding

power was calculated to reduce noise for gene correlations and emphasize correlations

between genes per WGCNA documentation. A block size of 22000 was used for network

construction and module detection as described previously [26]. Eigengenes were subsequently

calculated for each module detected by WGCNA. Gene modules whose eigengene values

changed significantly (P<0.05) for at least one timepoint were identified by ANOVA. Enrichment

analysis was performed using clusterProfiler to identify significantly associated ontology terms

for each module. All ontologies contained in clusterProfiler were evaluated, including BP

(biological process), CC (cellular component), and MF (molecular function).

Evaluating Schwann cell differentiation from amniotic mesenchymal stem cells via single

cell RNA sequencing

Single cell RNA-seq (scRNA-seq) data of Schwann cells (SCs) differentiated from human

amniotic mesenchymal stem cells (hAMSCs) was downloaded from Gene Expression Omnibus

(GSE161066) [28]. Seurat [33] was used for initial data cleaning and quality control, removing

cells with >2,500 or <200 unique feature counts as well as cells with >5% mitochondrial RNAs.

The hAMSC and SCs samples were each normalized with SCTransform [34, 35] and then

integrated with Harmony [36, 37]. Uniform manifold approximation and projection (UMAP) [38]

was used as a final dimensionality reduction step. The integrated object was clustered, markers

were identified for each cluster ($|Log_2FC|>0.58$ and *FDR<0.05*), and the relative cluster

proportions were compared between hAMSCs and SCs. DGE analysis was also performed

between hAMSC and SCs to identify DEGs specific to either group (|Log$_2$FC|>0.58 and

FDR<0.05).

Evaluation of Schwann cell differentiation in mouse sciatic nerve

Bulk RNAseq data (FPKM values) from sciatic nerve harvested at various developmental stages

(E17.5, P1, P5, P14, P24, P60) and peripheral nerves distal to the DRG at E13.5 were

downloaded from the SNAT data portal/GEO (GSE137870) [29]. For gene names with more

than one ID, the ID was selected with the highest average expression across all cells in the

dataset. Because gene to gene comparisons are biases by sequence length, we converted the

FPKM values to TPM values for gene filtering. Genes with a TPM value greater than the median

TPM value of all genes were retained for downstream analysis. Next, differential gene

expression analysis was performed using edgeR (3.34.1). Six comparisons were performed,

one for each timepoint (E17.5, P1, P5, P14, P24, P60) compared to the E13.5 reference

timepoint. Genes were considered differentially expressed if they had an |Log$_2$FC|>5 and an

FDR<0.001. Genes were mapped to their appropriate human homolog using biomaRt (2.48.3)

and the Dec 2021 Ensembl archive for both human and mouse.

Evaluation of myelinating compared to non-myelinating mouse Schwann cells

Bulk RNAseq data (FPKM values) from sorted Schwann cell populations enriched for

myelinating versus non-myelinating Schwann cells were downloaded from the SNAT data

portal/GEO (GSE137947) [29]. The initial processing and homolog mapping steps were as

described above. DEGs were compared between myelinating and non-myelinating Schwann

cells. Genes were considered differentially expressed if they had an |Log$_2$FC|>5 and an

FDR<0.001.

Cellular subtyping of Schwann cells from mouse sciatic nerve

The integrated Seurat (4.3.0) object from the SNAT data portal (SS2_all) was downloaded [29]

for analysis and comparison with additional datasets. UMAP was calculated on the integrated

components using the first 10 dimensions. Marker genes were identified using the

FindAllMarkers function. The homolog mapping steps were as previously described. In addition

to identifying markers from the previously defined clusters, we also determined DEGs for

various differentiation trajectories. Specifically, we identified DEGs using the FindMarkers

function contrasting immature Schwann cells (iSCs) from myelinating Schwann cells (mSCs),

i.e., iSC vs. tSC and nm(R)SC, and DEGs contrasting iSCs from non-myelinating Schwann cells

(nmSCs), i.e., iSC vs. pmSC, mSC cluster 1, mSC cluster 2, and mSC cluster 3. Genes were

considered differentially expressed if they had an $|Log_2FC|>0.58$ and an *FDR<0.05*.

Identification of conserved gene expression programs in differentiating Schwann cells

across myelination status and species

To examine the similarity of transcriptional programs between mouse and hiPSCs during

Schwann cell differentiation, we compared DEGs from our SC-IU dataset with the SNAT RNA-

seq dataset for each of the six time point comparisons. The Jaccard index (JI) was calculated

for up-regulated DEGs for each pairwise comparison between the SC-IU and SNAT datasets

(i.e., E17.5 up-regulated DEGs (SNAT) and Day 3 up-regulated DEGs (SC-IU) JI, E17.5 up-

regulated DEGs (SNAT) and Day 6 up-regulated DEGs (SC-IU) JI, etc.). The same process was

conducted for the down-regulated DEGs, resulting in a total of 36 total JI values for the up-

regulated and downregulated DEGs, respectively. We then averaged and normalized the JI

values to obtain a square matrix with row and column sums of 1 to account for any skew toward

either end of the differentiation process. We then repeated this process with less stringent DEG

calling criteria ($|Log_2FC|>0.58$ and an *FDR<0.05*). The consistency of the JI matrix was

evaluated using a Student's t-test of the diagonal of the matrix vs. the off-diagonal and a

Spearman correlation of the JI with the distance of each cell from the diagonal. Next, we

compared the union of up-regulated and down-regulated DEGs from both the SC-IU and SNAT

datasets to differentiation and developmental gene modules from the WGCNA analysis using

the original criteria ($|Log_2FC|>5$ and an *FDR<0.001*). Finally, mSC markers and nmSC markers

from the SNAT scRNA-seq dataset were compared to the mSC DEGs identified from the SNAT

myelination bulk RNA-seq dataset. We use these comparisons to evaluate the consistency of

genes across modalities, species, myelination status, and analysis technique to identify high-

confidence targets for the study of Schwann cell differentiation.

RESULTS

Morphological and immunofluorescence-based validation of hiPSC induced

differentiation to SPCs and SCs

To confirm robust differentiation of hiPSCs into Schwann cells precursors (SCPs) and

subsequently SCs, we performed morphological and immunofluorescence (IF) staining of cells

collected at seven timepoints along the hiPSC to SC continuum (**Fig. 1A**, **Table 1**). We

observed morphological changes during the differentiation process where the cells exhibited

increasing levels of cohesion into aggregates and reduced levels of random arrangements of

individual cells (**Fig. 1A**). Schwann cells are derived from the neural crest and characterized by

the expression of distinct makers including S100β, p75, HOXb7, and GAP43 (**Fig. 1B**). A

gradual increase in the expression of S100β, p75, HOXb7, and GAP43 was observed

throughout the differentiation process reaching a maximum at the SC stage (**Fig. 1B, Fig S1**).

Collectively, these findings indicate that our methodology can efficiently generate SCPs and

SLPs from hiPSC with high fidelity.

Differential gene expression during SC differentiation

To characterize gene expression dynamics during Schwann cell differentiation, we performed

RNA sequencing of cells collected at seven discrete stages along the hiPSC to SC continuum.

As anticipated, replicates from each stage of differentiation clustered together in PCA space

(**Fig. 1C**). Notably, the hiPSC stage had clear separation from all other stages (**Fig. 1C**) and the

neural rosette (NR) through SCP stages of differentiation clustered more closely together

compared to other stages (**Fig. 1C**). There were considerable transcriptional differences

between all stages of differentiation, most notably hiPSCs versus later stages of differentiation

(**Fig. 1D, Fig. S1-S3**). We identified 18 genes that were significantly up-regulated across all 6

comparisons (**Table 2**). Of these, *ANXA1* and *CDH6* were also among the top 5 most significant

up-regulated DEGs (lowest FDR adjusted p-values) within any of the timepoint comparisons

(**Table 2**). Notably, *CDH6* is associated with cell-cell adhesion and *ANXA1* has been implicated

in Schwann cell proliferation and migration [47]. Of the 30 total top 5 up-regulated DEGs per

timepoint comparison, 18 were up-regulated across all timepoints These 18 genes were

enriched for ontology terms related to cell adhesion (GO:0007155, *P<0.001*) and anatomical

structure formation involved in morphogenesis (GO:0048646, *P<0.001*). In total, 50 significantly

down-regulated DEGs were identified at each timepoint comparison (**Table 2**). However, only

four were also found in the top 5 most significant down-regulated DEGs (lowest FDR adjusted

p-values) at any of the timepoint comparisons (**Table 2**). Of the top 5 down-regulated DEGs per

timepoint comparison, 18 genes were down-regulated across all timepoints and were enriched

for ontology terms related to the regulation of cell fate specification (GO:0042659, *P<0.001*).

Gene co-expression modules associated with differentiation, axon guidance and ECM

Building on the results of our differential gene expression analysis, we next performed weighted

gene co-expression network analysis (WGCNA) to identify co-regulated gene modules during

hiPSC differentiation to SCs. WGCNA identified 24 modules (**Fig. 2A**), and 11 of these had

significantly altered eigengene values during differentiation (i.e., *P<0.05*, **Table 3**). Of the 11

differentiation-associated modules, 10 had significantly enriched ontology terms associated with

them (**Table 3**). The most significant module identified by ANOVA was the brown module

(*P<0.001*, **Table 3**, **Fig. 2B**) which had an eigengene peak at day 6, i.e., a significant increase

from day 3 to day 6 (*P<0.001*, **Fig. 2B**) followed by a decrease from day 6 to day 12 (*P<0.001*,

Fig. 2B). Given the close relationship between Schwann cell development and neurons, we

were intrigued that the brown module was functionally enriched for axon guidance (P<0.001,

Fig. 2C), forebrain development (*P<0.001*, **Fig. 2C**), and sensory organ morphogenesis

(*P<0.001*, **Fig. 2C**).

The next two most significant modules were red (*P<0.001*) and turquoise (*P<0.001*) (**Table 3**).

The red module eigengene steadily increased from hiPSC to SCP stage (*P<0.001*, i.e., day 0 to

day 24, **Fig. S4**) then dramatically decreased from SCP to SC stage (*P<0.001*, i.e., day 24 to

day 31, **Fig. S4**) and was functionally enriched for biosynthetic pathways (**Fig. S4**). In contrast,

the turquoise module eigengene, which was functionally enriched for functionally enriched for

exocytosis (*P<0.001*, **Fig. S5**) and regulation of neuron projection development (*P<0.001*, **Fig.**

S5) decreased dramatically from the hiPSC to preneural rosette (pre-NR) stage (*P<0.001*, i.e.,

day 0 to day 3, **Fig. S5**) and remained suppressed through terminal SC differentiation (*P=0.732*,

i.e., day 3 to day 31, **Fig. S5**). Another pair of modules that showed distinct expression patterns

were the pink and blue modules (*P<0.001* and *P<0.001* respectively, **Table 3**). Similar to the

brown module that had a distinct eigengene peak at day 6, the pink module peaked at day 12,

i.e., a significant increase from day 6 to day 12 (*P<0.001*, **Fig. 2D**) and decreased thereafter

from day 12 to day 18 (*P<0.001*, **Fig. 2D**). Not surprisingly, this module had numerous ontology

terms related to cell fate commitment and ear development (**Fig. 2E**). The blue module

eigengene also had a peak early in hiPSC to SC differentiation at day 3, i.e., a significant

increase from day 0 to day 3 (*P<0.001*, **Fig. 2F**) and decrease from day 3 to day 6 (*P<0.001*,

Fig. 2F). However, this module eigengene gradually increased again from NR through

completion of SC maturation (*P<0.001*, **Fig. 2F**). Notably, the blue module contains the *NF2*

tumor suppressor gene and had significant functional enrichment for extracellular matrix

organization (*P<0.001*, **Fig. 2G**), extracellular structure organization (*P<0.001*, **Fig. 2G**), and

cell-substrate adhesion (*P<0.001*, **Fig. 2G**). Taken together with the remaining modules (**Fig.**

S6-S11), the results of our WGCNA reveal that hiPSC differentiation to SCs is associated with

dynamic changes in gene expression that are reflective of Schwann cell biological function and

show associations with neural morphology, ear development, and cellular organization.

Signatures of hAMSC to SC differentiation are unique at the single cell level

To evaluate the robustness and generalizability of our findings, we extended our analysis to

examine whether similar gene modules and pathways were retained in other complementary

models of human Schwann cell differentiation. Wei and colleagues performed single cell RNA-

sequencing of SCs derived from human amniotic mesenchymal stem cells (hAMSCs) to identify

heterogenous SC populations [28] (SC-SUMC: GSE161066, **Table 1**). The scRNA-seq data

contained 3,892 hAMSCs and 1,729 SCs. After quality control (QC) and dimensionality

reduction, we identified a total of 16 clusters (**Fig. 3A**). DGE analysis revealed clear differences

between hAMSC and SCs. Some of the most upregulated genes in SC compared to hAMSCs

included ECM related genes like: *COL1A1* and *COL3A1* (**Fig. 3B**, **Table 4**). To further

characterize each cluster, we performed cluster-specific marker analysis using Seurat to identify

distinct gene sets highly expressed in each cluster (**Fig. 3C**). We observed that some clusters

had a higher proportion of either hAMSCs or SCs, suggesting that they may represent different

stages of differentiation (**Fig. 3D**). For example, cluster 8 and 14 comprised of a higher

proportion of SC sample compared to the hAMSC sample (**Fig. 3D**) and both of these clusters

overexpressed ECM-related markers. Cluster 8 overexpressed *VIM*, and cluster 14

overexpressed *ITGA1*, *COL1A1*, and *ADAMTS6* (**Fig. 3C**, **Table S1**).

Evaluation of Schwann cell differentiation in mouse sciatic nerve

To further validate our hiPSC model of Schwann cell differentiation *in vitro*, we further compared our transcriptomic data with an *in vivo* dataset comprehensively cataloging Schwann cell development in the mouse sciatic nerve. Gerber and colleagues generated gene expression profiles from Schwann cells isolated from mouse sciatic nerve throughout embryogenesis and postnatal development, which are publicly available through the Sciatic Nerve Atlas, i.e., SNAT, [29]. By integrating bulk and single cell RNA-seq data from SNAT, we identified unique and conserved gene signatures of Schwann cell differentiation across species and modalities.

In our SC-IU dataset, a total of 50 genes were significantly up-regulated across all timepoint comparisons, i.e., were contained in each comparison (**Fig. 4A-F, Table S2, Table S3**). These genes represented myelination (GO:0042552, *P<0.001*) and collagen-containing extracellular matrix (GO:0062023, *P<0.001*). The 362 down-regulated DEGs significant at all comparisons were enriched for neuron development (GO:0048666, *P<0.001*) and synaptic signaling (GO:0099536, *P<0.001*). Furthermore, we identified 9 gene co-expression modules that were significantly altered over developmental stage from E13.5 to P60. (**Table S4**, **Fig. S12-S21**). Notably some of these modules were also enriched for terms related to neural development like the SNAT-brown module (GO:0010975, Regulation of neuron projection development, *P<0.001,* **Table S4**).

Differential gene expression and enrichment in myelinating versus non-myelinating murine Schwann cells

We next sought to investigate the gene expression profiles of myelinating (mSCs) and non-myelinating (nmSCs) in the Sciatic Nerve Atlas. mSCs had 478 up-regulated DEGs and 471 down-regulated DEGs in comparison to nmSCs (**Fig. 4G, Table S5**). These up-regulated DEGs were enriched for structural constituents of myelin sheath (GO:0019911, *P=0.039*), myelination

(GO:0042552, *P<0.001*), and sterol biosynthetic processes (GO:0016126, *P<0.001*). Down-

regulated DEGs were highly enriched for mitotic cell cycle (GO:0000278, *P<0.001*), cell division

(GO:0051301, *P<0.001),* and positive regulation of cell cycle processes (GO:0090068,

P<0.001).

The myelinating and non-myelinating Schwann cell subsets remained largely distinct even upon

recomputing the UMAP dimensionality reduction (**Fig. 5A**)[29]. Two distinct trajectories

emerged, both beginning from proliferating Schwann cells (prol. SC) and terminating at either

mSC cluster 3 or nm(R)SC (**Fig. 5A**)[29]. We observed 261 up-regulated and 116 down-

regulated DEGs in myelinating Schwann cells (mSC) as compared to immature Schwann cells

(iSCs) (**Table S6**). As expected, the up-regulated DEGs in mSCs were enriched for terms

related to regulation of nervous system development (GO:0051960, *P<0.001*) and myelination

(GO:0042552, *P<0.001*). However, the most significantly enriched term was sterol biosynthetic

process (GO:0016126, *P<0.001*). The robust upregulation of cell cycle inhibitors like *CDKN1A*

(**Fig. 5B**) in conjunction with the downregulation of cyclins such as *CDK6* (**Fig. 5B**) further

suggest a state of proliferative arrest within this cluster. Using as few as three transcriptomic

markers, we could distinguish the stages of Schwann cell differentiation and the bifurcation of

mSCs and nmSCs (**Fig. 5C & D, Table S7**). Again, we identified ECM-related markers in the

different differentiation stage iSC (*POSTN*), tSC (*ID3*), and mSC (*SPP1*) clusters (**Table S7**).

Differentiating Schwann cells have conserved pathways related to ECM

As anticipated, the transcriptional signatures of Schwann cell differentiation were not identically

conserved between hiPSCs and mice, likely due to both species and model specific differences.

Without normalization, there was no significant correlation between the developmental stages

within the SNAT dataset and the iPSC differentiation timepoints (**Fig. S22**). However, after

accounting for differences in the row and column DEG counts, a moderate association emerged

(PCC=0.44, *P=0.007*, **Fig 6A**). The one-to-one correspondences in DEGs between timepoints

(E17.5-Day 3, P1-Day 6, P5-Day 12, P14-Day 18, P24-Day 24, and P60-Day 31) were also

moderately associated (*P=0.084*, **Fig. 6B**).

We focused on the blue module and turquoise modules and compared them to the up- and

down-regulated DEGs from the SC-IU and SNAT datasets. The blue module contained *NF2*

and the eigengene trended upward throughout the differentiation process, whereas the

turquoise module trended downward. We found that the up-regulated DEGs from both SC-IU

and SNAT had high overlap with the blue module, but low overlap with one another (**Fig. 6C**). In

contrast the SC-IU up-regulated DEGs had lower overlap with the turquoise module, while

SNAT up-regulated DEGs had a similar number of DEGs in the turquoise module (**Fig. 6D**).

Both the SC-IU and SNAT down-regulated DEGs had lower overlap with the blue module (**Fig.**

6E) than with the turquoise module (**Fig. 6F**). Notably 21 genes were up-regulated in both

datasets and contained within the blue module (**Table S2**). These genes represented specific

molecular functions including extracellular matrix structural constituent (GO:0005201, *P=0.005*)

and S100 protein binding (GO:0044548, *P=0.008*) as well as biological processes including

caveola assembly (GO:0070836, *P=0.006*) and regulation of transforming growth factor beta

receptor signaling pathway (GO:0017015, *P=0.006*).

In addition to these unique conserved processes during murine Schwann cell development and

hiPSC to Schwann cell differentiation, mSCs had distinct molecular profiles in comparison to

nmSCs that were recapitulated in both RNA-seq and scRNA-seq data modalities. Notably, mSC

up-regulated DEGs from SNAT RNA-seq had high overlap with mSC markers from SNAT

scRNA-seq (**Fig. 6G**). Similarly, nmSC up-regulated DEGs from SNAT RNA-seq exhibited a

high degree of overlap with nmSC markers from SNAT scRNA-seq (**Fig. 6H**).

DISCUSSION

We have developed a state-of-the-art platform to study Schwann cell development using

hiPSCs. We demonstrate that hiPSCs successfully differentiated into SCs based on marker

gene expression and morphology. Furthermore, we identify similar transcriptomic profiles in our

hiPSC to SC differentiation model as observed in other external Schwann cell differentiation

studies [28, 29]. Notably, we observed correlation of DEGs and gene expression signatures

including those relating to ECM and TGF-β signaling between hiPSC derived SCs and those

arising in situ from the developing mouse sciatic nerve [29]. Intriguingly, TGF-β1 signaling has

been shown to coordinate the production of abundant ECM proteins by *Nf1* deficient Schwann

cells, contributing to plexiform neurofibroma pathogenesis in neurofibromatosis type 1 (NF1)

[39]. However, the role of ECM and TGF- β signaling in schwannoma development remains

largely understudied.

Key differences between our hiPSC model and mouse Schwann cell development also

emerged, which may be attributable to both species-specific differences as well as the model

system employed (*in vitro* versus *in vivo*). Transcriptional discordance between SCs derived

from hAMSCs versus hiPSCs may in part be attributable to unique cells of origin and the

experimental differences related the reprograming process. While hiPSCs have been utilized

successfully to overcome limitation of mouse models to enhance our understanding of

psychiatric disorders [40] and Alzheimer's disease [21], further investigation is needed to

validate the fidelity of this platform in recapitulate the pathogenesis of *NF2*-SWN and other

Schwann cell associated disorders.

Notwitstanding these disparities, our analysis did reveal distinctive gene signatures associated

with Schwann cell development that were conserved across species and model systems. We

identified gene co-expression modules related to differentiation, particularly the 'blue module'

that contained the *NF2* tumor suppressor gene. This module demonstrated functional

enrichment for multiple terms related to ECM and actin filaments, which is consistent with the

established relationship between Merlin and F-actin dependent cytoskeletal defects in *NF2-*

associated schwannoma [41]. The emergence of ECM-related genes was a consistent finding

across datasets, analytical techniques, and experimental methodologies employed in the study

of differentiating Schwann cells. Notably, two vesicle-associated genes within the 'blue module',

CAV1 and *CAV2*, were found to be upregulated in both hiPSC and murine derived SCs and

recently *NF2* was shown to regulate vesicle trafficking in schwannoma [42]. Additionally,

collagens *COL1A1* and *COL3A1*, were also overrepresented in our analysis and recently shown

to be associated with recurrence of vestibular schwannoma after radiation therapy [43]. Other

ECM-related genes identified, albeit with lesser substantiation in schwannoma included *ITGA1*,

POSTN, *SPP1*, *ID3*, and *ADAMTS6*. One of the most promising candidates was lysyl oxidase,

i.e., *LOX*, which recurred in multiple analyses. LOX is an essential protein in the ECM that

mediates conversion of collagen precursors into reactive forms capable of cross-linking and

stabilizing ECM proteins to modulate cell adhesion and motility [44].

Cell cycle inhibitors, such as *CDKN2B*, were also present in the 'blue module' and conserved in

cross species analysis. *CDKN2B* is a key component of the *INK4/ARF* tumor suppressor locus

and can induce proliferation when deleted or mutated in a multitude of cancer types [45, 46].

Furthermore, we identified hiPSC-specific genes of interest that were highly up-regulated during

SC differentiation and differentially expressed at all timepoints including *ANXA1,* which has

been implicated in promoting Schwann cell proliferation and migration in response to nerve

injury [47], yet its relationship to Merlin in Schwann cell biology remains unknown.

In summary, we developed a novel hiPSC Schwann cell differentiation model and performed

integrated analyses across five datasets from three independent studies, encompassing both

scRNA-seq and RNA-seq. Our findings indicate that our hiPSC model robustly recapitulates

Schwann cell ontologies and identifies critical genes implicated in Schwann cell differentiation

warranting further investigation. Additionally, our data revealed a pronounced association

between SC development and ECM across multiple datasets, that remains largely

uncharacterized with respect to Merlin biology and schwannoma development. Future studies

leveraging this hiPSC platform to conduct CRISPR-Cas9 editing of *NF2* and other associated

genes relevant to SC differentiation/function, generate hiPSC organoid co-cultures, and perform

bulk and single cell mulitomic analysis hold promise to reveal new insights into Schwann cell

biology, the Schwann cell-ECM relationship, and schwannomas pathogenesis in *NF2*-SWN.

ACKNOWLEDGEMENTS

We would like to acknowledge the help and support of the Indiana University School of Medicine

Center of Computational Biology and Bioinformatics, the Indiana University Information

Technology Services for their high-performance computing resources, and the Indiana

Biosciences Research Institute for their lab space and technical support.

461
462   **FUNDING**
463
Funding from the Indiana Biosciences Research Institute to A.S. and T.S.J., 1R21CA264339 to

T.S.J. S.D.R. is supported by the Francis S. Collins Scholars Program in Neurofibromatosis

Clinical and Translational Research funded by the Neurofibromatosis Therapeutic Acceleration

Program (NTAP, 2004757180), a K08 Mentored Clinical Scientist Research Career

Development Award funded by the NIH/National Institute of Neurological Disorders and Stroke

(NINDS) (K08-NS128266-02), and startup funding from the Department of Pediatrics at the

Indiana University (IU) School of Medicine and the IU Simon Comprehensive Cancer Center

(IUSCCC).

472
473   **DISCLOSURES**
474
475   O.L., W.C., A.H., A.Q., T.S.J., are partially supported by the Indiana Biosciences Research

476   Institute. A.H. currently holds a research position at Eli Lilly and Company.

477
478   **AUTHOR CONTRIBUTIONS**

480   S.D.R, A.S.Q, and T.J. designed the experiments and evaluated the data. W.C, A.H, A.S.Q, and

481   B.E.H. performed the experiments. O.L, S.L, and T.J analyzed all data and prepared the graph.

482   T.J, S.L, A.S.Q, and S.D.R wrote the initial draft of the manuscript. S.P.A and D.W.C. gave

483   advice during the project achievement. All co-authors provided feedback and revisions.

 TABLES

485 **Table 1** Table of datasets used for analysis in this study

Dataset name	Samples	Description	Publication	PMID	Accession
SC-IU	21 samples RNA-seq (3 replicates across 7 timepoints) RNA-seq	hiPSC to SC differentiation sampled at seven timepoints Organism: Human	This study	This study	This study
SC-SUMC	2 samples (1 hAMSC sample with 3,892 cells and 1 SC sample with 1,729 cells) scRNA-seq	hAMSC to SC differentiation sampled at hAMSC and SC stages Organism: Human	Wei, Z. frontiers in Physiology. (2021)	34093220	GSE161066
SNAT	28 samples (4 replicates across 7 timepoints) RNA-seq; 8 FACS samples (4 mSC and 4 nmSC) RNA-seq; 8 samples (2 replicates across 4 stages: P1, P5, P14, and P60) scRNA-seq	SCP to SC differentiation; Comparison of mSC and nmSCs; Cellular subtyping of Schwann cells during stages of development. Organism: Mouse	Gerber, D. eLife. (2021)	33890853	GSE137870

487 **Table 2** Top 5 up- and down-regulated DEGs for each of the 6 timepoints (k) (days 3, 6, 12, 18,

488 24, and 31) compared to day 0 hiPSCs as the reference.

Day k vs. Day 0	Gene	Log2FC	P	FDR
Day 3 Up	*COL8A1*	11.33	0	0
Day 3 Up	*POSTN*	10.15	0	0
Day 3 Up	*SERPINE1*	9.22	0	0
Day 3 Up	*IGFBP3*	7.61	0	0
Day 3 Up	*TGFBI*	7.15	0	0
Day 3 Down	*CD24*	-6.37	5.14E-260	8.69E-257
Day 3 Down	*EPCAM*	-7.93	1.19E-229	1.07E-226
Day 3 Down	*HLA-A*	-5.53	5.65E-213	4.16E-210
Day 3 Down	*PMEL*	-7.55	7.08E-213	5.08E-210
Day 3 Down	*ESRG*	-9.76	7.22E-212	5.06E-209
Day 6 Up	*HAPLN1*	6.97	8.78E-236	2.53E-231

Day 6 Up	CDH6	5.91	1.68E-227	2.43E-223
Day 6 Up	PRTG	4.91	4.72E-225	4.54E-221
Day 6 Up	WLS	5.36	1.57E-184	7.55E-181
Day 6 Up	GREB1L	6.90	2.12E-183	8.73E-180
Day 6 Down	TDGF1	-5.01	8.98E-198	6.47E-194
Day 6 Down	ADCY2	-6.77	1.89E-190	1.09E-186
Day 6 Down	LINC00678	-5.15	2.72E-151	7.83E-148
Day 6 Down	AC064802.1	-4.65	9.03E-148	2.37E-144
Day 6 Down	DUSP6	-4.65	7.95E-145	1.64E-141
Day 12 Up	ANXA1	10.95	4.31E-276	6.09E-272
Day 12 Up	CPE	5.72	2.35E-191	1.67E-187
Day 12 Up	PLK2	5.42	1.95E-190	1.11E-186
Day 12 Up	PRTG	4.52	1.72E-185	8.13E-182
Day 12 Up	CDH6	5.53	4.22E-181	1.71E-177
Day 12 Down	TDGF1	-8.16	0	0
Day 12 Down	AC064802.1	-8.52	7.05E-207	6.65E-203
Day 12 Down	NANOG	-7.30	7.96E-173	2.82E-169
Day 12 Down	USP44	-5.13	5.67E-164	1.46E-160
Day 12 Down	AC009446.1	-7.39	2.73E-162	6.42E-159
Day 18 Up	ANXA1	10.79	0	0
Day 18 Up	COL8A1	10.92	0	0
Day 18 Up	CAV1	6.12	0	0
Day 18 Up	SERPINE1	8.14	5.48E-295	3.85E-291
Day 18 Up	POSTN	8.01	2.79E-294	1.56E-290
Day 18 Down	TDGF1	-10.73	2.01E-273	7.06E-270
Day 18 Down	USP44	-5.52	1.62E-253	4.54E-250
Day 18 Down	ZSCAN10	-6.42	3.59E-216	4.03E-213
Day 18 Down	ESRG	-5.16	8.92E-199	7.82E-196
Day 18 Down	VSNL1	-7.28	3.70E-179	2.16E-176
Day 24 Up	ANXA1	11.19	0	0
Day 24 Up	POSTN	8.43	0	0
Day 24 Up	CAV1	6.16	0	0
Day 24 Up	PLK2	5.77	0	0
Day 24 Up	SPP1	4.22	1.63E-291	9.33E-288
Day 24 Down	USP44	-5.49	8.04E-274	2.29E-270
Day 24 Down	ESRG	-5.53	8.55E-249	1.22E-245
Day 24 Down	ZSCAN10	-7.03	1.28E-236	1.52E-233
Day 24 Down	PMEL	-4.70	6.92E-211	6.17E-208
Day 24 Down	B3GNT7	-6.79	7.86E-201	5.98E-198

Day 31 Up	CTGF	6.05	0	0
Day 31 Up	VIM	6.55	0	0
Day 31 Up	COL5A2	7.04	0	0
Day 31 Up	COL8A1	11.75	0	0
Day 31 Up	POSTN	9.95	0	0
Day 31 Down	CD24	-6.94	0	0
Day 31 Down	HLA-A	-6.02	8.07E-296	7.22E-293
Day 31 Down	SFRP2	-5.66	3.19E-289	2.61E-286
Day 31 Down	L1TD1	-5.78	4.22E-275	3.26E-272
Day 31 Down	EPCAM	-7.92	1.74E-267	1.18E-264

490   **Table 3** Significantly altered gene modules in relation to timepoint evaluated by ANOVA. The

491   number of genes in the module, ANOVA P, and most significant GO term are listed for each

492   module. Modules with no significant GO terms are blank.

Module	Genes	P	Top GO term	File
Brown	1449	7.66E-17	Axon guidance	File S2
Red	845	2.63E-14	Organic hydroxy compound biosynthetic process	File S3
Turquoise	4615	5.23E-14	Nuclear chromosome segregation	File S4
Cyan	100	1.38E-11		File S5
Pink	692	2.11E-11	Cell fate commitment	File S6
Blue	3797	2.16E-10	Extracellular matrix organization	File S7
Black	833	1.73E-8	Intraciliary transport particle B	File S8
Magenta	159	4.99E-5	Endoplasmic reticulum protein-containing complex	File S9
Yellow	1288	1.52E-2	Cytoplasmic translation	File S10
Midnight Blue	78	4.19E-2	Transcription coregulator activity	File S11
Green Yellow	146	4.26E-2	Methyltransferase activity	File S12

494   **Table 4** Top 10 up- and down-regulated DEGs in hAMSCs versus SCs from the SC-SUMC

495   dataset.

Gene	Proportion hAMSCs>0	Proportion SCs>0	Log_2FC	P	FDR
ISG15	0.04	0.44	0.66	1.00E-300	0.00E+00
IFI27	0.02	0.32	0.61	4.98E-256	5.71E-252
IFI6	0.01	0.27	0.45	1.37E-210	1.58E-206
COL3A1	0.07	0.40	0.55	2.62E-202	3.00E-198
COL1A1	0.06	0.34	0.49	6.85E-174	7.86E-170

LGALS1	0.49	0.81	0.68	2.36E-171	2.71E-167
EEF1A1	0.97	0.88	-0.58	1.66E-158	1.90E-154
BST2	0.02	0.22	0.34	1.86E-139	2.14E-135
TPT1	0.96	0.88	-0.55	1.79E-133	2.05E-129
TPM1	0.31	0.60	0.61	1.85E-118	2.12E-114
TMSB10	0.88	0.95	0.47	2.17E-99	2.49E-95
S100A6	0.47	0.70	0.55	3.15E-99	3.62E-95
RPL3	0.83	0.66	-0.45	8.45E-81	9.70E-77
CXCL8	0.71	0.48	-0.55	9.51E-79	1.09E-74
IGF2	0.20	0.01	-0.34	6.67E-76	7.65E-72
RPS3A	0.80	0.61	-0.42	1.52E-72	1.75E-68
SAT1	0.35	0.12	-0.37	7.64E-72	8.77E-68
MT2A	0.98	0.98	-0.58	1.37E-70	1.57E-66
RPS12	0.89	0.77	-0.40	1.54E-70	1.77E-66
RPS6	0.76	0.57	-0.40	6.27E-67	7.20E-63

 FIGURES

Fig. 1 Workflow and validation of directed differentiation of hiPSCs into human Schwann

cells. **A)** Schematic representation of the differentiation of hiPSCs into Schwann cell precursors

(SCPs) and Schwann cells (SCs) (Upper panel). Representative bright-field images showing the

process of differentiation into SCPs and SCs. Scale bar, 100 μm (middle panel). Steps involve

sequencing and data analysis (lower panel). **B)** Schwann cell differentiation was confirmed by

positive immunocytochemical staining for S100β (green), p75 (red) in upper panel and HOXb7

(green), GAP43 (red) in lower panel after 31 days of differentiation. DAPI (blue) was used to stain

the cell nuclei. Scale bars, 100 μm. **C)** Timepoints can be stratified into stages of differentiation

by PCA. **D)** Heatmap of differentially expressed genes across different timepoints where the rows

(genes) were clustered using hierarchical clustering.

Fig. 2 WGCNA reveals dynamic changes in module eigengene values during hiPSC to SC

differentiation related to tissue structure and neural development. **A)** Gene correlation matrix

with different modules annotated. **B)** Brown module eigengene values are plotted over the

course of the study timepoints. **C)** Significant ontology terms from functional enrichment

analysis for the brown module. **D)** Pink module eigengene changes over the study timepoints.

E) Significant ontology terms from functional enrichment analysis for the pink module. **F)** Blue

module (containing *NF2*) eigengene changes over the study timepoints. **G)** Significant ontology

terms from functional enrichment analysis for the blue module (containing *NF2*).

Fig. 3 SC-SUMC scRNA-seq analysis. **A)** UMAP clustering of hAMSCs and SCs from the

integrated SC-SUMC dataset (GSE161066) reveals 16 clusters. **B)** Volcano plot of differentially

expressed genes when compared between hAMSCs and SCs. **C)** Heatmap of cluster specific

522   markers. **D)** Proportion of hAMSCs and SCs comprising each cluster from the integrated SC-

523   SUMC dataset.

 Fig. 4 Differentially expression analysis of the SNAT RNA-seq dataset (GSE137868). Volcano

 plots depicting differentially expressed genes when comparing the E17.5 **(A)**, P1 **(B)**, P5 **(C)**,

 P14 **(D)**, P24 **(E)**, and P60 **(F)** stages to E13.5. Volcano plot depicting differentially expressed

 genes when comparing myelinating vs. non-myelinating Schwann cells **(G)**.

 Fig. 5 SNAT scRNA-seq analysis. **A)** Scatterplot of 8 clusters identified from the SNAT study

 representing different stages of Schwann cell development along both myelinating and non-

 myelinating trajectories. **B)** Volcano plot depicting differentially expressed genes comparing

 mSC vs iSC. **C)** Volcano plot depicting differentially expressed genes between nmSC versus

 iSC. **D)** Heatmap of cluster specific markers (top 3).

 Fig. 6 Comparison of DEGs identified between multiple datasets. **A)** Normalized JI matrix of

 averaged SC-IU DEGs (|log$_2$FC| > 0.58 and FDR < 0.05) throughout SC differentiation and

 SNAT DEGs (|log$_2$FC| > 0.58 and FDR < 0.05) through various embryonic and postnatal

 developmental states with comparison of the JI matrix diagonal to the non-diagonal **(B)**, **C)**

Venn diagram comparing genes comprising the blue module, SC-IU up-regulated DEGs during

SC differentiation, and SNAT up-regulated DEGs throughout development, **D)** Venn diagram

comparing genes comprising the turquoise module, SC-IU up-regulated DEGs during SC

differentiation, and SNAT up-regulated DEGs throughout development, **E)** Venn diagram

comparing genes comprising the blue module, SC-IU down-regulated DEGs during SC

differentiation, and SNAT down-regulated DEGs throughout development, **F)** Venn diagram

comparing genes comprising the turquoise module, SC-IU down-regulated DEGs during SC

differentiation, and SNAT down-regulated DEGs throughout development, **G)** Comparison of

genes up-regulated in mSC vs. iSC and nmSC vs. iSC from SNAT scRNA-seq and DEGs

upregulated in mSC vs. nmSC from SNAT RNA-seq, **H)** Comparison of genes down-regulated

in mSCs vs. and nmSC vs. iSC from SNAT scRNA-seq and DEGs down-regulated in mSC vs.

nmSC from SNAT RNA-seq.

REFERENCES

1. Jessen, K.R., R. Mirsky, and A.C. Lloyd, *Schwann Cells: Development and Role in Nerve Repair.* Cold Spring Harb Perspect Biol, 2015. **7**(7): p. a020487.

2. Salzer, J.L. and B. Zalc, *Myelination.* Curr Biol, 2016. **26**(20): p. R971-R975.

3. Goncalves, N.P., et al., *Schwann cell interactions with axons and microvessels in diabetic neuropathy.* Nat Rev Neurol, 2017. **13**(3): p. 135-147.

4. Carlson, M.L. and M.J. Link, *Vestibular Schwannomas.* N Engl J Med, 2021. **384**(14): p. 1335-1348.

5. Evans, D.G., *Neurofibromatosis type 2 (NF2): a clinical and molecular review.* Orphanet J Rare Dis, 2009. **4**: p. 16.

6. Agnihotri, S., et al., *The genomic landscape of schwannoma.* Nat Genet, 2016. **48**(11): p. 1339-1348.

7. Rouleau, G.A., et al., *Alteration in a new gene encoding a putative membrane-organizing protein causes neuro-fibromatosis type 2.* Nature, 1993. **363**(6429): p. 515-21.

8. Evans, D.G., et al., *A clinical study of type 2 neurofibromatosis.* Q J Med, 1992. **84**(304): p. 603-18.

9. Leventoux, N., et al., *Human Astrocytes Model Derived from Induced Pluripotent Stem Cells.* Cells, 2020. **9**(12).

10. Perriot, S., et al., *Differentiation of functional astrocytes from human-induced pluripotent stem cells in chemically defined media.* STAR Protoc, 2021. **2**(4): p. 100902.

11. Perriot, S., et al., *Human Induced Pluripotent Stem Cell-Derived Astrocytes Are Differentially Activated by Multiple Sclerosis-Associated Cytokines.* Stem Cell Reports, 2018. **11**(5): p. 1199-1210.

12. Voulgaris, D., P. Nikolakopoulou, and A. Herland, *Generation of Human iPSC-Derived Astrocytes with a mature star-shaped phenotype for CNS modeling.* Stem Cell Rev Rep, 2022. **18**(7): p. 2494-2512.

13. Abud, E.M., et al., *iPSC-Derived Human Microglia-like Cells to Study Neurological Diseases.* Neuron, 2017. **94**(2): p. 278-293 e9.

14. Hasselmann, J. and M. Blurton-Jones, *Human iPSC-derived microglia: A growing toolset to study the brain's innate immune cells.* Glia, 2020. **68**(4): p. 721-739.

15. McQuade, A., et al., *Development and validation of a simplified method to generate human microglia from pluripotent stem cells.* Mol Neurodegener, 2018. **13**(1): p. 67.

16. Autar, K., et al., *A functional hiPSC-cortical neuron differentiation and maturation model and its application to neurological disorders.* Stem Cell Reports, 2022. **17**(1): p. 96-109.

17. Vierbuchen, T., et al., *Direct conversion of fibroblasts to functional neurons by defined factors.* Nature, 2010. **463**(7284): p. 1035-41.

18. Brennand, K.J., et al., *Modelling schizophrenia using human induced pluripotent stem cells.* Nature, 2011. **473**(7346): p. 221-5.

19. Page, S.C., et al., *Electrophysiological measures from human iPSC-derived neurons are associated with schizophrenia clinical status and predict individual cognitive performance.* Proc Natl Acad Sci U S A, 2022. **119**(3).

20. Park, I.H., et al., *Disease-specific induced pluripotent stem cells.* Cell, 2008. **134**(5): p. 877-86.

21. Penney, J., W.T. Ralvenius, and L.H. Tsai, *Modeling Alzheimer's disease with iPSC-derived brain cells.* Mol Psychiatry, 2020. **25**(1): p. 148-167.

22. Krieger, T.G., et al., *Modeling glioblastoma invasion using human brain organoids and single-cell transcriptomics.* Neuro Oncol, 2020. **22**(8): p. 1138-1149.

23.   Sancho-Martinez, I., et al., *Establishment of human iPSC-based models for the study and targeting of glioma initiating cells.* Nat Commun, 2016. **7**: p. 10743.

24.   Schloo, C. and L.M. Kutscher, *Modeling brain and neural crest neoplasms with human pluripotent stem cells.* Neuro Oncol, 2023. **25**(7): p. 1225-1235.

25.   Zhang, S., et al., *Development of human induced pluripotent stem cell (iPSC) line from a 60year old female patient with multiple schwannoma.* Stem Cell Res, 2017. **19**: p. 31-33.

26.   Langfelder, P. and S. Horvath, *WGCNA: an R package for weighted correlation network analysis.* BMC Bioinformatics, 2008. **9**: p. 559.

27.   Garcia-Moreno, A., et al., *Functional Enrichment Analysis of Regulatory Elements.* Biomedicines, 2022. **10**(3).

28.   Wei, Z., et al., *A Subpopulation of Schwann Cell-Like Cells With Nerve Regeneration Signatures Is Identified Through Single-Cell RNA Sequencing.* Front Physiol, 2021. **12**: p. 637924.

29.   Gerber, D., et al., *Transcriptional profiling of mouse peripheral nerves to the single-cell level to build a sciatic nerve ATlas (SNAT).* Elife, 2021. **10**.

30.   Kim, H.S., et al., *Schwann Cell Precursors from Human Pluripotent Stem Cells as a Potential Therapeutic Target for Myelin Repair.* Stem Cell Reports, 2017. **8**(6): p. 1714-1726.

31.   Liu, Z., et al., *Specific marker expression and cell state of Schwann cells during culture in vitro.* PLoS One, 2015. **10**(4): p. e0123278.

32.   Chen, J., et al., *ToppGene Suite for gene list enrichment analysis and candidate gene prioritization.* Nucleic Acids Res, 2009. **37**(Web Server issue): p. W305-11.

33.   Hao, Y., et al., *Integrated analysis of multimodal single-cell data.* Cell, 2021. **184**(13): p. 3573-3587 e29.

34.   Choudhary, S. and R. Satija, *Comparison and evaluation of statistical error models for scRNA-seq.* Genome Biol, 2022. **23**(1): p. 27.

35.   Lause, J., P. Berens, and D. Kobak, *Analytic Pearson residuals for normalization of single-cell RNA-seq UMI data.* Genome Biol, 2021. **22**(1): p. 258.

36.   Korsunsky, I., et al., *Fast, sensitive and accurate integration of single-cell data with Harmony.* Nat Methods, 2019. **16**(12): p. 1289-1296.

37.   Tran, H.T.N., et al., *A benchmark of batch-effect correction methods for single-cell RNA sequencing data.* Genome Biol, 2020. **21**(1): p. 12.

38.   McInnes, L., J. Healy, and J. Melville, *Umap: Uniform manifold approximation and projection for dimension reduction.* arXiv preprint arXiv:1802.03426, 2018.

39.   Jiang, C., et al., *Basement membrane proteins in extracellular matrix characterize NF1 neurofibroma development and response to MEK inhibitor.* J Clin Invest, 2023. **133**(12).

40.   Deng, B., *Mouse models and induced pluripotent stem cells in researching psychiatric disorders.* Stem Cell Investig, 2017. **4**: p. 62.

41.   Bashour, A.M., et al., *The neurofibromatosis type 2 gene product, merlin, reverses the F-actin cytoskeletal defects in primary human Schwannoma cells.* Mol Cell Biol, 2002. **22**(4): p. 1150-7.

42.   Hennigan, R.F., et al., *The NF2 tumor suppressor regulates microtubule-based vesicle trafficking via a novel Rac, MLK and p38(SAPK) pathway.* Oncogene, 2013. **32**(9): p. 1135-43.

43.   Shi, Q., et al., *Collagen Family Genes Associated with Risk of Recurrence after Radiation Therapy for Vestibular Schwannoma and Pan-Cancer Analysis.* Dis Markers, 2021. **2021**: p. 7897994.

44.   Kagan, H.M. and W. Li, *Lysyl oxidase: properties, specificity, and biological roles inside and outside of the cell.* J Cell Biochem, 2003. **88**(4): p. 660-72.

45.   Tu, Q., et al., *CDKN2B deletion is essential for pancreatic cancer development instead of unmeaningful co-deletion due to juxtaposition to CDKN2A.* Oncogene, 2018. **37**(1): p. 128-138.

46.   Wrensch, M., et al., *Variants in the CDKN2B and RTEL1 regions are associated with high-grade glioma susceptibility.* Nat Genet, 2009. **41**(8): p. 905-8.

649   47.   Xia, W., et al., *ANXA1 directs Schwann cells proliferation and migration to accelerate nerve*
650   *regeneration through the FPR2/AMPK pathway.* FASEB J, 2020. **34**(10): p. 13993-14005.