Prefrontal dopamine circuits are required for active avoidance learning but not for fear

learning

Zachary Zeidler*[1], Marta Fernández Gómez *[2], Tanya A. Gupta[2], Meelan Shari, Scott A. Wilke[2#],

and Laura A. DeNardo[1#]

[1]Department of Physiology; [2]Semel Institute for Neuroscience and Human Behavior, Department

of Psychiatry, David Geffen School of Medicine, University of California, Los Angeles, California.

[#]Correspondence: Dr. Laura A. DeNardo, ldenardo@ucla.edu; Dr. Scott A. Wilke

swilke@mednet.ucla.edu.

* these authors contributed equally to this work

Abstract

The medial prefrontal cortex (mPFC) resolves approach-avoidance conflicts and mediates

associative processes required for learning to avoid threats. Dopamine (DA) projections from the

ventral tegmental area (VTA) to the mPFC carry information about aversive outcomes that may

inform prefrontal computations. However, the role of prefrontal DA in learning based on aversive

outcomes remains poorly understood. Here, we used platform mediated avoidance (PMA) to

study the role of mPFC DA in threat avoidance learning in mice. We show that activity within

dopaminergic VTA terminals in the mPFC is required for signaled avoidance learning, but not for

escape, conditioned fear, or to recall a previously learned avoidance strategy. Taken together,

these data indicate that mPFC DA is necessary to rapidly form associations between predictive

cues and actions that preempt aversive outcomes but not for cue-outcome learning.

Introduction

To thrive in a complex environment, animals must seek beneficial outcomes while avoiding

threats. Natural environments are highly dynamic, requiring individuals to continuously update

predictive relationships between stimuli, actions, and outcomes. With respect to threats, this

learning must occur rapidly and reliably to avoid injury or death. Excessive risk taking enhances

mortality rates. Conversely, excessive avoidance can be maladaptive by limiting opportunities to

engage in beneficial pursuits. Disorders featuring maladaptive avoidance are observed with both

excessive and insufficient avoidance[1–4]. Thus, it is critical to understand the neurobiological

mechanisms that mediate rapid threat avoidance learning.

The medial prefrontal cortex (mPFC) is essential for integrating learned information about

the environment to guide actions, in particular when resolving approach-avoidance conflicts and

when establishing action-outcome relationships[5–7]. Activity in mPFC is likewise critical for both

learning to avoid threats and for driving avoidance behavior, especially in situations that involve

conflicting motivational drives[8–16]. While much progress has been made in understanding the

mPFC mechanisms that drive avoidance behavior, less is known about the prefrontal

mechanisms underlying avoidance learning. In particular, the neural signals that drive mPFC

circuit changes underlying avoidance learning remain largely unknown.

Mesocortical DA is poorly studied compared to mesolimbic DA circuits due to challenges

associated with the lower density of cortical dopaminergic fibers as well as the lack of tools to

monitor DA with high temporal resolution in isolation from other catecholamines[17]. Most VTA DA

neurons targeting mPFC do not collateralize and represent a subpopulation with distinct

molecular, cellular and functional properties[18–24]. In contrast to the reward-related functions of

mesolimbic DA projections, VTA-mPFC projecting DA neurons preferentially respond to and

orient behavior toward aversive stimuli[24–29]. VTA-mPFC dopaminergic axons exhibit minimal

reward response but a substantial shock response[30]. Similarly, activating mPFC DA terminals

during a competitive appetitive-aversive stimulus assay biased behavior toward aversive

responding[31]. Phasic stimulation of VTA-mPFC DA terminals promoted deviation from previously

learned rules in a set-shifting task[32], highlighting a key role for mPFC DA in behavioral flexibility.

Altogether, there is a compelling role for mPFC DA in processing aversive stimuli and in

behavioral flexibility. But the role of mPFC DA in enabling the flexible behavior required to learn

appropriate behavioral responses to aversive stimuli remains unknown.

In this study, we utilized platform mediated avoidance[14,15] (PMA) to investigate the

hypothesis that activity within VTA-mPFC dopaminergic projections is critical for learning to

preemptively avoid aversive outcomes. We further hypothesized that these processes are distinct

from associative learning between a stimulus and an aversive outcome that lacks an action-

outcome contingency. Using a combination of terminal-specific optogenetic inhibition of VTA-

mPFC DA projections and distinct behavioral assays, we outline a critical role for mPFC DA

projections in threat avoidance learning that is independent of aversive associative learning.

These results suggest that DA modulates mPFC activity to rapidly and adaptively utilize predictive

cues to drive behaviors that preempt aversive outcomes.

Results

Platform mediated avoidance learning

Deciding to avoid potential threats often comes at the cost of rewarding behaviors and requires

mPFC activity[8,9,14]. To model threat avoidance under conditions of conflict, we trained mice in

platform mediated avoidance (PMA), in which a shock-conditioned tone prompts mice to stop

collecting rewards and navigate to a safety platform. We first trained thirsty mice that sweetened

condensed milk is available in the PMA chamber and then performed three days of PMA training

under these conditions. In this assay, the reward area was located on the opposite side of the

chamber from the safety platform (Figure 1A). On each day, nine 30-second tones each co-

terminated with a 2-second mild foot shock (Figure 1B) and we video recorded mouse behavior

from above, using supervised deep learning to track eight points on their body[33] (Figure 1C).

Successful trials were classified as trials in which mice accessed the safety platform before shock

onset and remained there for the duration of the shock. Mice steadily increased the percent of

successful trials across three days of PMA training (Figure 1D). They also steadily increased the

percent time spent on the safety platform during each tone (Figure 1E). Freezing behavior during

the tones was low and did not change significantly across days (Figure 1F). Thus, mice first learn

to associate a cue with a foot shock and subsequently learn to preemptively avoid footshocks in

the context of a competing opportunity to receive rewards.

Activity in dopaminergic terminals in mPFC is required for rapid avoidance learning

To determine if activity in VTA-mPFC dopaminergic axon terminals is required for avoidance

learning, we employed an optogenetic approach to inhibit axon terminals during tone+shock

presentations across three days of PMA training. First, we injected an AAV encoding the Cre-

conditional inhibitory opsin Jaws[34] or a control fluorophore (GFP) into the VTA of TH-Cre mice. In

these mice, the bacterial recombinase Cre is expressed under the control of the promoter for

tyrosine hydroxylase – an enzyme necessary for dopamine synthesis – so Jaws expression is

restricted to dopaminergic VTA neurons. Then we placed bilateral optic fibers over mPFC (Figure

2A). In these mice, we observed a subset of brightly labeled GFP+ neurons in VTA and GFP+

axons in mPFC (Figure 2B). We then trained mice in PMA as described above. During each tone

(including during the shock period), we inhibited VTA-mPFC dopaminergic axon terminals using

constant 635 nm laser light delivered through bilateral head mounted optic fibers (Figure 2C).

We analyzed mouse behavior across the three days of PMA to determine whether

disrupting activity in VTA-mPFC DA terminals affected conditioned freezing behavior, a measure

of the tone-shock association, and platform mediated avoidance behavior, a measure of the

action-outcome association. First, we analyzed the fraction of successful trials when mice

preemptively avoided foot shocks (Figure 2D). Inhibiting activity in VTA-mPFC DA terminals

caused a significant slowing of avoidance learning. Mice expressing Jaws had significantly fewer

successful trials than GFP control mice, especially on the first day and the beginning of the second

day of PMA (Figure 2E).

Interestingly, Jaws mice had a significantly higher proportion of escape trials (Figure 2F)

when they leaped to the platform once the shock began. The biggest effects were apparent early

in learning, at the end of day 1 and the beginning of day 2. VTA-mPFC DA terminal inhibition had

no effect on freezing behavior in the context of PMA (Figure 2G). In the same mice, we showed

that inhibiting VTA-mPFC DA terminals did not induce conditioned place preference or aversion

(Figure S1). Together, these data reveal that VTA-mPFC activity is most important in the early

stages of learning. Instead of preventing learning, our findings suggest that inhibiting mPFC DA

activity slows the formation of the predictive relationship between the avoidance action and the

aversive outcome without affecting the formation of the association between the tone and the

shock.

Even though Jaws mice could eventually learn PMA, we wondered if they developed

qualitative differences in approach and avoidance behaviors as a result of inhibiting VTA-mPFC

DA activity during learning. To examine this, we quantified the patterns of behavior during tone

presentations, including interactions with the platform and with the reward port. Jaws animals

exhibited fractured avoidance behavior, spending significantly shorter bouts of time on the

platform overall and in particular during the tone period (Figure 2H,I). Both groups of mice

exhibited a behavioral strategy in which they infrequently triggered the reward during the tone

period and but did so more frequently during the inter-tone intervals, when threat levels were

lower. We observed no differences in reward port interactions during the tone period (Figure 2K),

but Jaws mice had significantly more interactions with the reward port than controls during the

inter-tone interval (ITI) (Figure 2J). Together, these data indicate that inhibition of VTA-mPFC DA

terminals does not alter cue-shock associative learning, nor does it prevent learning about the

location of the safety platform. Instead, the data indicate that VTA-mPFC inhibition impairs the

ability to use that information to guide preemptive avoidance behavior.

VTA-mPFC activity is required for rapid avoidance learning even without motivational conflict

To determine if the effects we observed were specific to 1) avoidance under conflict and 2) threat avoidance learning vs. memory retrieval, we performed additional optogenetics experiments in separate groups of mice. First, we tested whether VTA-mPFC activity was necessary during a PMA assay without a reward conflict. As before, we injected an AAV expressing Cre-dependent Jaws or GFP into the VTA of TH-Cre mice and implanted bilateral optic fibers above mPFC (Figure 3A). Because we observed the largest effects of inhibition during the first day of PMA, we performed a single day of PMA training during which 12 tone-shock trials were paired with delivery constant 635 nm light to mPFC and no reward was available. As before, we observed that Jaws mice had a significant decrease in the fraction of successful trials, a significant increase in escape trials, and no change in the percent time freezing during the tone (Figure 3C), suggesting that mPFC DA activity is still required for learning to avoid threats even without a motivational conflict .

To determine to what extent VTA-mPFC dopaminergic projection activity is necessary for avoidance behavior versus for learning associations between conditioned stimuli and adaptive behaviors, we inhibited VTA-mPFC DA terminals during PMA learning and then examined avoidance behavior during a PMA memory retrieval session the following day. During retrieval, mice were presented with 10 tones without foot shocks or laser inhibition. Jaws mice spent significantly less time on the safety platform during tones compared to controls (Figure 3D), indicating the VTA-mPFC DA terminal inhibition impaired learning as opposed to preventing the expression of avoidance behavior. We observed no group differences in conditioned freezing during the tones, indicating further that VTA-mPC DA terminal activity is not needed to learn the tone-shock association.

VTA-mPFC activity is not required to maintain a learned avoidance strategy

We next investigated whether inhibiting VTA-mPFC axon terminals would affect PMA retrieval

after learning had occurred (Figure 3E). We trained mice in PMA (without conflict) without laser

inhibition (Figure 3F). The next day, mice had a PMA retrieval session. In this case, half the tones

were paired with laser inhibition and laser on and off trials were randomly interleaved. This

manipulation had no effect on the percent time on the platform or percent time spent freezing

during the tones (Figure 3G), supporting the idea that VTA-mPFC activity is only required during

the early stages of avoidance learning to enable the mPFC to associate action-outcome

contingencies.

Activity in dopaminergic terminals in mPFC is not required for cued fear conditioning

We next sought to determine which aspects of VTA-mPFC DA terminal activity are specific to

PMA, which requires animals to first learn that a tone predicts a shock (cue-outcome contingency)

and then to learn that during the tone, they can enter the platform to avoid the foot shock (action-

outcome contingencies). To do so, we examined the role of mPFC DA terminals in cued fear

conditioning, which features similar tone-shock associative learning without any action-outcome

contingency. To test the role of VTA-mPFC DA terminals in cued fear conditioning, we injected

an AAV expressing cre-dependent Jaws or a fluorophore control (GFP) into VTA of TH-Cre mice

and implanted optic fibers bilaterally above mPFC (Figure 4A). On the training day, we paired

tones with continuous delivery of far-red light through the optic fibers in mPFC. After three

baseline tones, 12 subsequent tones were paired with a 2-second footshock (identical to PMA).

On the test day, mice were presented with six tones without shocks or laser illumination (Figure

4B). On the training day, mice in both groups significantly increased their freezing across the

session, but we observed no difference between the GFP and Jaws groups (Figure 4C). On the

test day, there was no difference in the time spent freezing during tone presentation (Figure 4D).

Together, these results indicate that VTA-mPFC activity is not required to learn the association

between cues and shocks.

Discussion

Here we discovered that VTA-mPFC DA activity is critical for rapidly learning to avoid aversive

outcomes in PMA. On the other hand, activity in the same projection is not required for developing

a conditioned freezing response during auditory fear conditioning, indicating that activity in this

pathway is not required for establishing cue-outcome relationships. These findings extend our

understanding of the role of VTA-mPFC DA projections beyond aversive processing and

behavioral flexibility to include a causal function in learning the predictive relationships between

cues, actions and aversive outcomes.

191   The mPFC is heavily implicated in mediating approach-avoidance conflicts[8,9,9,11,15] and in

the recall and expression of cued fear[35–40], but the mechanisms that drive the circuit plasticity

required for learning to avoid aversive outcomes are poorly understood. DA signaling plays a

critical role in aversive learning but has mostly been studied without regional specificity and/or in

the context of fear conditioning[41]. While fear conditioning paradigms involve associative learning

between a stimulus and an aversive outcome, they lack the possibility of taking action to avoid

the shock, i.e. action-outcome contingencies, which require mPFC activity[42–44]. Studies that used

midbrain lesions, global pharmacological inhibition of DA signaling and microdialysis identified a

key role for DA in threat avoidance behaviors[45–47]. While these studies built a critical foundation

of knowledge, the approaches lack the spatiotemporal precision to reveal the causal role of mPFC

DA in avoidance behavior. Previous research using microdialysis lacked the temporal resolution

to align mPFC DA with cues and behavior, while other research using voltammetry could not

separate DA from fluctuations in other catecholamines (e.g., norepinephrine)[17]. Therefore, the

nuanced relationship between VTA-mPFC DA activity, threatening cues, and threat-induced

behaviors, and how these relationships may change across learning was not understood. Using

temporally precise manipulations, we showed that inhibiting VTA-mPFC DA terminals delayed the

acquisition of avoidance behavior without affecting the learning or recall of the cue-shock

relationship.

While VTA-mPFC DA activity supports behavioral flexibility in some scenarios[27,32,48–52], we

do not find evidence that inhibiting VTA-mPFC DA activity slows avoidance learning by impairing

behavioral flexibility. In support of this idea, Jaws mice did not exhibit elevations in competing

behaviors - including freezing and collecting rewards - during conditioned tones. Instead, Jaws

mice exhibited fractured avoidance behavior - once they entered the safety platform, they left

more quickly than controls. These data suggest that reducing mPFC DA does not impair

deviations from existing behavioral schemes. Instead, mPFC DA likely biases prefrontal

computations to rapidly adopt behavioral strategies that prioritize actions that preempt aversive

outcomes, including entering and then staying on the platform until the end of the shock-

conditioned tone.

Interestingly, Jaws mice collected more rewards than GFP mice during the ITI even though

no laser manipulation was present. Over the course of learning, control mice had slightly longer

platform bouts during the ITI, suggesting they prioritized safety seeking over reward seeking even

when no overt threats were present. One interpretation of these findings is that Jaws mice have

less generalized fear than controls. If true, this would suggest that while mPFC DA activity is

critical for learning the predictive relationships between cues, actions and outcomes, it also

promotes generalization of conditioned fear outside the cue periods.

Our study causally manipulated the activity of VTA-mPFC terminals during presentations

of tones, foot shocks, and foot shock omissions on successful trials. Since we performed these

experiments in TH-Cre mice, the effects we observed were likely related to DA activity. However,

recent studies showed that a subset of TH+ VTA cells also co-express the vesicular glutamate

transporter VGLUT2[53] and mesocortical DA neurons may co-release glutamate[54]. A more

complete understanding of how these projections drive plasticity will require temporally precise

manipulations during isolated events (e.g. only during shocks), examination of how activity arising

from VTA-mPFC terminals influences mPFC microcircuit activity, and investigation of to what

extent observed effects arise from DA vs. other neurotransmitters.

Protracted disturbances in the balance of reward seeking vs. avoidance is a hallmark of

numerous psychiatric disorders[3,55,56]. Environmental factors (e.g., stress or trauma) can promote

maladaptive learning that precipitates pathological avoidance[57–60]. Primary treatments include

cognitive behavioral therapy and drugs whose targets cluster in neuromodulatory systems,

including in the dopamine system. But once instantiated, excessive avoidance can interfere with

therapy and exacerbate features of psychiatric illness[61]. Establishing a mechanistically detailed

and temporally specific understanding of the role of mPFC DA in specific types of learning

promises to enable more precise interventions that combine pharmacologic, brain stimulation

and/or psychotherapeutic modalities during precise windows of vulnerability or capacity for

change. Therefore, it is critical to understand the neurobiology of avoidance learning if we are to

effectively intervene on psychiatric disorders.

Materials and Methods

All experiments were conducted in accordance with procedures established by the administrative

panels on laboratory animal care at the University of California, Los Angeles.

Animals Male and female C57Bl6/J mice (JAX stock #000664) or tyrosine hydroxylase Cre (TH-

Cre; line Fl12; www.gensat.org) mice were group housed (two to five per cage) and kept on a

12/12 h light/dark cycle (lights on 7AM to 7PM). Mice were sexed by examination of external

genitalia   at   P21.   Animals   received   *ad   libitum*   food   and   water.

Surgery Mice were induced in 5% isoflurane in oxygen until loss of righting reflex and transferred

to a stereotaxic apparatus where they were maintained under 2% isoflurane in oxygen. Mice were

warmed with a circulating water heating pad throughout surgery and eye gel was applied to the

eyes. The head was shaved and prepped with three scrubs of alternating betadine and then 70%

ethanol. Following a small skin incision, a dental drill was used to drill through the skull above the

injection targets. A syringe pump (Kopf, 693A) with a Hamilton syringe was used for injections.

Injections were delivered at a rate of 100 nl/min and the syringe was left in the brain for 10 min

following injection. For optogenetics experiments, animals were injected with 500nL of AAV-CAG-

FLEX- JAWS-KGC-GFP-ER2 or AAV5-CAG-FLEX-EGFP-WPRE at a titer of 5×10^{12} vg/mL

bilaterally into the VTA (AP: -2.9 mm, ML: +/- 0.45 mm, DV: -4.5 mm from bregma). Bilateral optic

fibers (200 μm core, Newdoon) were implanted over the prelimbic (PL) area of the mPFC (AP:

+1.8, ML: +/- 0.35, DV: -1.8 mm from bregma). For pain management mice received 5 mg/kg

carprofen diluted in 0.9% saline subcutaneously. Mice received one injection during surgery and

daily injections for 2 d following surgery. Optogenetic experiments waited at least four weeks after

surgery before experimentation.

Behavior

Platform mediated avoidance

Three days prior to starting PMA, citric acid (2% w/v) was added to animals' drinking water. This

self-initiates a reduction in water consumption and increases performance on liquid reward tasks

similar to forced water restriction[62]. Animals were then placed into a 20x22cm arena with metal

rod flooring (Lafeyette Instruments) and a plexiglass platform covering one-quarter of the floor

space. Opposite the platform was a reward port and reward pump (Campden Instruments). The

reward port featured an infrared beam that triggered 25 μL of 10% sweetened condensed milk

(Nestle) reward on a variable interval schedule (mean interval: 30 seconds). Once a day for two

days, animals were placed in the arena and allowed to freely explore and consume reward for 45

minutes. The following day, PMA training began. For three days, animals were placed into the

arena with a 180 second baseline period followed by delivery of a 30 second, 75 dB, 4kHz tone

that co-terminated with a 2 second, 0.15 mA shock delivered to the floor (Lafeyette Instruments).

Eight subsequent tone-shock pairings were delivered at randomly chosen intervals between 80

and 160 seconds. Training on PMA continued for two more days for a total of three days. Data

presented in tone tone blocks are the average performance over three tones.

Cued fear conditioning

Experimentally naïve animals were placed in the same arena as PMA without the platform or

reward port. On training day, animals received three, 10 second 4 kHz tones to assess baseline

freezing. The subsequent twelve tones were paired with a footshock identical to PMA. The

following day (recall), animals were exposed to six, 30 second 4 kHz tones unpaired with shock.

Optogenetics

JAWS activation was achieved via continuous red light (635 nm, Shanghai Laser and Optics

Century) delivered through the entirety of the tone-shock periods. For PMA, laser illumination was

concurrent with all the tones. For fear conditioning, this occurred on the first day (training).

Real-time place preference

After PMA, a subset of animals were placed in a real-time place preference (RTPP) assay.

Animals were placed in a 68 x 23 cm chamber for 20 minutes. During the first ten minutes, the

animals freely explored the arena to establish a baseline side preference. During the subsequent

ten minutes, a closed loop video monitoring system (Bioviewer) tracked the location of the animal

and triggered a 635 nm laser when the animal entered the "laser" side of the arena. The ratio of

time spent in the "non-laser" side of the arena to the "laser" side of the arena was calculated for

both baseline (laser off) and test (laser on) periods.

Histology Animals were deeply anesthetized with isoflurane and transcardially perfused with

phosphate buffered saline (PBS), followed by 4% paraformaldehyde (PFA). Brains were removed

and post-fixed overnight in 4% PFA before being moved to PBS. Brains were then sectioned via

vibratome at 60 microns. To detect VTA axons expressing JAWS-GFP in the mPFC, anti-GFP

immunohistochemistry was performed. Sections were incubated with chicken anti-GFP polyclonal

antibody (1:2000, Aves Labs) overnight at 4℃. Sections were next rinsed and incubated with

donkey anti-chicken AlexaFluor 488 conjugate (1:500; Jackson ImmunoResearch) at room

temperature for 2 hr. Sections were washed before being mounted with DAPI. Images of

fluorescent expression and implant targeting were taken using a Leica DM6 scanning microscope

(Leica Microsystems) using a 10x objective.

Data Analysis

Behavioral Analysis

High resolution, overhead videos of PMA and fear conditioning were collected at 50 Hz (Teledyne

FLIR) and animal positioning extracted via supervised deep learning networks (cite DLC). Animal

position information was transformed into freezing and ROI features using BehaviorDEPOT

software[63].

Statistical analysis

Modeling was performed with custom Python code. All other statistical testing was performed in

Graphpad Prism.

Data and code availability

Custom MATLAB code available upon request.

Acknowledgements

We thank Dr. Peter Balsam for helpful discussions of this work. This work was supported by a
Klingenstein-Simons Fellowship in Neuroscience, a Whitehall Foundation Research Grant and
1R01MH127214-01A1 (LAD), a Whitehall Foundation Research Grant and 1R01MH131858-
01A1 (SAW), T32NS115753 and F31MH133387 (ZEZ) and T32NS115753 (TG).

Author Contributions

LAD, SAW, ZEZ, MFG and TG conceptualized the experiments. ZEZ, MFG and MS performed

experiments. TG analyzed behavioral data. LAD, SAW and ZEZ wrote the manuscript. All authors

edited the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

1. Chandler, R. A., Wakeley, J., Goodwin, G. M. & Rogers, R. D. Altered Risk-Aversion and
Risk-Seeking Behavior in Bipolar Disorder. *Biological Psychiatry* **66**, 840–846 (2009).

2. Cáceda, R., Nemeroff, C. B. & Harvey, P. D. Toward an Understanding of Decision Making
in Severe Mental Illness. *JNP* **26**, 196–213 (2014).

3. Penninx, B. W. J. H., Pine, D. S., Holmes, E. A. & Reif, A. Benzodiazepines for the long-term
treatment of anxiety disorders? – Authors' reply. *The Lancet* **398**, 120 (2021).

4. Letkiewicz, A. M., Kottler, H. C., Shankman, S. A. & Cochran, A. L. Quantifying aberrant
approach-avoidance conflict in psychopathology: A review of computational approaches.
Neuroscience & Biobehavioral Reviews **147**, 105103 (2023).

5. Pastor, V. & Medina, J. H. Medial prefrontal cortical control of reward- and aversion-based
behavioral output: Bottom-up modulation. *European Journal of Neuroscience* **53**, 3039–3062
(2021).

6.   Mair, R. G., Francoeur, M. J., Krell, E. M. & Gibson, B. M. Where Actions Meet Outcomes:

Medial Prefrontal Cortex, Central Thalamus, and the Basal Ganglia. *Frontiers in Behavioral*

Neuroscience **16**, (2022).

7.   Matsumoto, K. & Tanaka, K. The role of the medial prefrontal cortex in achieving goals.

Current Opinion in Neurobiology **14**, 178–185 (2004).

8.   Diehl, M. M., Bravo-Rivera, C. & Quirk, G. J. The study of active avoidance: A platform for

discussion. *Neuroscience and Biobehavioral Reviews* **107**, 229–237 (2019).

9.   Diehl, M. M. *et al.* Active avoidance requires inhibitory signaling in the rodent prelimbic

prefrontal cortex. *eLife* (2018) doi:10.7554/eLife.34657.

10.   Diehl, M. M. *et al.* Divergent projections of the prelimbic cortex bidirectionally regulate active

avoidance. *eLife* **9**, 1–13 (2020).

11.   Moscarello, J. M. & LeDoux, J. E. Active avoidance learning requires prefrontal suppression

of   amygdala-mediated   defensive   reactions.   *Journal   of   Neuroscience*   (2013)

doi:10.1523/JNEUROSCI.2596-12.2013.

12.   Jercog, D. *et al.* Dynamical prefrontal population coding during defensive behaviours. *Nature*

595, 690–694 (2021).

13.   Kajs, B. L., Loewke, A. C., Dorsch, J. M., Vinson, L. T. & Gunaydin, L. A. Divergent encoding

of active avoidance behavior in corticostriatal and corticolimbic projections. *Scientific Reports*

12, 1–11 (2022).

14.   Bravo-Rivera, C., Roman-Ortiz, C., Brignoni-Perez, E., Sotres-Bayon, F. & Quirk, G. J. Neural

Structures Mediating Expression and Extinction of Platform-Mediated Avoidance. *Journal of*

Neuroscience (2014) doi:10.1523/JNEUROSCI.0191-14.2014.

15.   Bravo-Rivera, C., Roman-Ortiz, C., Montesinos-Cartagena, M. & Quirk, G. J. Persistent active

avoidance   correlates   with   activity   in   prelimbic   cortex   and   ventral   striatum.   *Frontiers   in*

Behavioral Neuroscience (2015) doi:10.3389/fnbeh.2015.00184.

16.   Capuzzo, G. & Floresco, S. B. Prelimbic and Infralimbic Prefrontal Regulation of Active and

387 Inhibitory Avoidance and Reward-Seeking. *Journal of Neuroscience* **40**, 4773–4787 (2020).

388 17. Megan E. Fox & R. Mark Wightman. Contrasting Regulation of Catecholamine

389 Neurotransmission in the Behaving Brain: Pharmacological Insights from an Electrochemical

390 Perspective. *Pharmacol Rev* **69**, 12 (2017).

391 18. Ikemoto, S. Brain reward circuitry beyond the mesolimbic dopamine system: A

392 neurobiological theory. *Neuroscience & Biobehavioral Reviews* **35**, 129–150 (2010).

393 19. Lammel, S. *et al.* Unique Properties of Mesoprefrontal Neurons within a Dual

394 Mesocorticolimbic Dopamine System. *Neuron* **57**, 760–773 (2008).

395 20. Chiodo, L. A., Bannon, M. J., Grace, A. A., Roth, R. H. & Bunney, B. S. Evidence for the

396 absence of impulse-regulating somatodendritic and synthesis-modulating nerve terminal

397 autoreceptors on subpopulations of mesocortical dopamine neurons. *Neuroscience* **12**, 1–16

398 (1984).

399 21. M J Bannon & R H Roth. Pharmacology of mesocortical dopamine neurons. *Pharmacol Rev*

400 **35**, 53 (1983).

401 22. Björklund, A. & Dunnett, S. B. Dopamine neuron systems in the brain: an update. *Trends in*

402 *Neurosciences* **30**, 194–202 (2007).

403 23. Lammel, S., Ion, D. I., Roeper, J. & Malenka, R. C. Projection-Specific Modulation of

404 Dopamine Neuron Synapses by Aversive and Rewarding Stimuli. *Neuron* **70**, 855–862

405 (2011).

406 24. Lammel, S. *et al.* Input-specific control of reward and aversion in the ventral tegmental area.

407 *Nature* **491**, 212–217 (2012).

408 25. Kim, C. K. *et al.* Simultaneous fast measurement of circuit dynamics at multiple sites across

409 the mammalian brain. *Nature methods* **13**, 325–328 (2016).

410 26. Weele, C. M. V., Siciliano, C. A. & Tye, K. M. Dopamine tunes prefrontal outputs to orchestrate

411 aversive processing. *Brain Research* **1713**, 16–31 (2019).

412 27. Vander Weele, C. M. *et al.* Dopamine enhances signal-to-noise ratio in cortical-brainstem

encoding of aversive stimuli. *Nature* **563**, 397–401 (2018).

28. Abercrombie, E. D., Keefe, K. A., DiFrischia, D. S. & Zigmond, M. J. Differential Effect of

Stress on In Vivo Dopamine Release in Striatum, Nucleus Accumbens, and Medial Frontal

Cortex. *Journal of Neurochemistry* **52**, 1655–1658 (1989).

29. Lerner, T. N., Holloway, A. L. & Seiler, J. L. Dopamine, Updated: Reward Prediction Error and

Beyond. *Current Opinion in Neurobiology* **67**, 123–130 (2021).

30. Kim, C. K. *et al.* Simultaneous fast measurement of circuit dynamics at multiple sites across

the mammalian brain. *Nat Methods* **13**, 325–328 (2016).

31. Vander Weele, C. M., Siciliano, C. A. & Tye, K. M. Dopamine tunes prefrontal outputs to

orchestrate aversive processing. *Brain Res* **1713**, 16–31 (2019).

32. Ellwood, I. T. *et al.* Tonic or Phasic Stimulation of Dopaminergic Projections to Prefrontal

Cortex Causes Mice to Maintain or Deviate from Previously Learned Behavioral Strategies.

Journal of Neuroscience **37**, 8315–8329 (2017).

33. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with

deep learning. *Nature Neuroscience* **21**, (2018).

34. Chuong, A. S. *et al.* Noninvasive optical inhibition with a red-shifted microbial rhodopsin.

Nature Neuroscience **17**, 1123–1129 (2014).

35. Do-Monte, F. H., Quiñones-Laracuente, K. & Quirk, G. J. A temporal shift in the circuits

mediating retrieval of fear memory. *Nature* **519**, 460–463 (2015).

36. Vidal-Gonzalez, I., Vidal-Gonzalez, B., Rauch, S. L. & Quirk, G. J. Microstimulation reveals

opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear.

Learn. Mem. **13**, 728–733 (2006).

37. Burgos-Robles, A., Vidal-Gonzalez, I. & Quirk, G. J. Sustained Conditioned Responses in

Prelimbic Prefrontal Neurons Are Correlated with Fear Expression and Extinction Failure. *The*

Journal of Neuroscience **29**, 8474–8482 (2009).

38. Sierra-Mercado, D., Padilla-Coreano, N. & Quirk, G. J. Dissociable Roles of Prelimbic and

Infralimbic Cortices, Ventral Hippocampus, and Basolateral Amygdala in the Expression and

Extinction of Conditioned Fear. *Neuropsychopharmacology* **36**, 529–538 (2011).

39. Corcoran, K. A. & Quirk, G. J. Activity in Prelimbic Cortex Is Necessary for the Expression of

Learned, But Not Innate, Fears. *Journal of Neuroscience* **27**, 840–844 (2007).

40. Lee, Y. K. & Choi, J.-S. Inactivation of the Medial Prefrontal Cortex Interferes with the

Expression But Not the Acquisition of Diff erential Fear Conditioning in Rats. *Exp Neurobiol*

21, 23–29 (2012).

41. Zafiri, D. & Duvarci, S. Dopaminergic circuits underlying associative aversive learning.

Frontiers in Behavioral Neuroscience **16**, (2022).

42. Simon, N. W. & Moghaddam, B. Neural processing of reward in adolescent rodents.

Developmental Cognitive Neuroscience **11**, 145–154 (2015).

43. Alexander, W. H. & Brown, J. W. Medial prefrontal cortex as an action-outcome predictor.

Nature Neuroscience **14**, 1338–1344 (2011).

44. Matsumoto, M., Matsumoto, K., Abe, H. & Tanaka, K. Medial prefrontal cell activity signaling

prediction errors of action values. *Nature Neuroscience* **10**, 647–656 (2007).

45. Fibiger, H. C., Zis, A. P. & Phillips, A. G. Haloperidol-induced disruption of conditioned

avoidance responding: Attenuation by prior training or by anticholinergic drugs. *European*

Journal of Pharmacology **30**, 309–314 (1975).

46. Arnt, J. Pharmacological Specificity of Conditioned Avoidance Response Inhibition in Rats:

Inhibition by Neuroleptics and Correlation to Dopamine Receptor Blockade. *Acta*

Pharmacologica et Toxicologica **51**, 321–329 (1982).

47. Wadenberg, M.-L., Ericson, E., Magnusson, O. & Ahlenius, S. Suppression of conditioned

avoidance behavior by the local application of (−)sulpiride into the ventral, but not the dorsal,

striatum of the rat. *Biological Psychiatry* **28**, 297–307 (1990).

48. Floresco, S. B. Prefrontal dopamine and behavioral flexibility: shifting from an "inverted-U"

toward a family of functions. *Front. Neurosci.* **7**, (2013).

49. Luo, R. *et al.* A dopaminergic switch for fear to safety transitions. *Nature Communications* **9**,

1–11 (2018).

50. Peters, G. J., David, C. N., Marcus, M. D. & Smith, D. M. The medial prefrontal cortex is critical

for memory retrieval and resolving interference. *Learn. Mem.* **20**, 201–209 (2013).

51. Nakayama, H., Ibañez-Tallon, I. & Heintz, N. Cell-Type-Specific Contributions of Medial

Prefrontal Neurons to Flexible Behaviors. *J. Neurosci.* **38**, 4490–4504 (2018).

52. Del Arco, A., Park, J., Wood, J., Kim, Y. & Moghaddam, B. Adaptive Encoding of Outcome

Prediction by Prefrontal Cortex Ensembles Supports Behavioral Flexibility. *J. Neurosci.* **37**,

8363–8373 (2017).

53. Vrettou, M. *et al.* Vesicular glutamate transporter 2 expression in the ventral tegmental area

of outbred male rats following exposure to nicotine and alcohol. *Drug and Alcohol*

Dependence Reports **8**, 100180 (2023).

54. Antonieta Lavin *et al.* Mesocortical Dopamine Neurons Operate in Distinct Temporal Domains

Using Multimodal Signaling. *J. Neurosci.* **25**, 5013 (2005).

55. Loijen, A., Vrijsen, J. N., Egger, J. I. M., Becker, E. S. & Rinck, M. Biased approach-avoidance

tendencies in psychopathology: A systematic review of their assessment and modification.

Clinical Psychology Review **77**, 101825 (2020).

56. Robin L., A. & Martin, P. P. Neural systems underlying approach and avoidance in anxiety

disorders. *Dialogues in Clinical Neuroscience* **12**, 517–531 (2010).

57. Dunsmoor, J. E. & Paz, R. Fear Generalization and Anxiety: Behavioral and Neural

Mechanisms. *Biological Psychiatry* **78**, 336–343 (2015).

58. Assary, E., Vincent, J. P., Keers, R. & Pluess, M. Gene-environment interaction and

psychiatric disorders: Review and future directions. *Seminars in Cell & Developmental*

Biology **77**, 133–143 (2018).

59. Giovanniello, J., Bravo-Rivera, C., Rosenkranz, A. & Matthew Lattal, K. Stress, associative

learning, and decision-making. *Neurobiology of Learning and Memory* **204**, 107812 (2023).

60. Teicher, M. H., Samson, J. A., Anderson, C. M. & Ohashi, K. The effects of childhood

maltreatment on brain structure, function and connectivity. *Nature Reviews Neuroscience* **17**,

652–666 (2016).

61. Labrenz, F., Woud, M. L., Elsenbruch, S. & Icenhour, A. The Good, the Bad, and the Ugly—

Chances, Challenges, and Clinical Implications of Avoidance Research in Psychosomatic

Medicine. *Frontiers in Psychiatry* **13**, (2022).

62. Anne E. Urai *et al.* Citric Acid Water as an Alternative to Water Restriction for High-Yield

Mouse Behavior. *eNeuro* **8**, ENEURO.0230-20.2020 (2021).

63. Gabriel, C. J. *et al.* Behavior DEPOT is a simple, flexible tool for automated behavioral

detection based on marker less pose tracking. *eLife* **11**, 1–33 (2022).

Figure 1. Mice learn to perform a platform mediated avoidance assay.

A. Experimental timeline.

B. Behavioral training protocol.

C. Schematic of PMA chamber (left) and frames from videos (right).

D-F. Behavior across 3 days of PMA. On each day, data is analyzed in bins of 3 tones.

D. Successful trials ($F_{Trial}(1.27,6.34)=17.99$, p=0.004; $F_{Day}(1.96,9.79)=7.64$, p=0.01; n=6 mice).

E. Time on platform during tones ($F_{Trial}(1.37,6.82)=21.03$, p=0.002; $F_{Day}(1.74,8.71)=8.24$, p=0.01; n=6 mice).

F. Freezing during tones ($F_{Trial}(1.72,8.61)=1.31$, p=0.31; $F_{Day}(1.38,6.9)=1.83$, p=0.22; n=6 mice).

Figure 2. Optogenetic inhibition of DAergic axon terminals in mPFC impairs avoidance learning.

A. Schematic of viral and fiberoptic targeting locations.

B. Coronal sections from a representative brain showing Jaws-eGFP expression in VTA and bilateral fiber placement in PL.

C. Optogenetic inhibition during PMA. 635nm laser light was presented coincidently with a 30 second 4kHz tone that co-terminated with a 2-second footshock.

D. Schematics showing a successful trial when mice preemptively avoided the shock vs. an escape trial when mice leaped to the platform after shock begins.

E. Fraction of successful trials across days in GFP vs. Jaws mice (F_{time}(4.26, 49) = 17, P<0.0001; F_{opsin}(1, 13) = 5.66, p=0.03; $F_{interaction}$(8, 92) = 2.197, p=0.06).

F. Fraction of escape trials across days in GFP vs. Jaws mice (F_{time} (4,43, 49.86) = 5.43, p=0.0007; F_{opsin} (1, 13) = 4.78, p=0.04; $F_{interaction}$ (8, 90) = 1.04; p=0.4).

G. Fraction time freezing during tone across days in GFP vs. Jaws mice (F_{time} (4.32, 46.98) = 8.41, P<0.0001; F_{opsin} (1, 12) = 1.4, p=0.2; $F_{interaction}$ (8, 87) = 1.01; p=0.4).

H. Platform bout duration across days in GFP vs. Jaws mice (F_{time}(1.88, 19.83) = 0.5, p=0.5; F_{opsin}(1, 12) = 6.8, p=0.02; $F_{interaction}$(2, 21) = 0.7, p=0.4).

I. Platform bout duration during tone periods across days in GFP vs Jaws mice (F_{time}(4.26, 49) = 17, p=0.005; F_{opsin}(1, 13) = 5.66, p=0.04; $F_{interaction}$(8, 92) = 2.197, p=0.06; GFP vs. Jaws Day 1 p = 0.9, Day 2 = 0.02, Day 3 = 0.4.

J. Number of reward beam breaks across days in GFP and Jaws mice (F_{time}(1.9, 19.9) = 4.8, p=0.01; F_{opsin}(1, 12) = 8, p=0.01; $F_{interaction}$(2, 21) = 0.02, p=0.97; GFP vs Jaws Day 1 p=0.01; Day 2 p=0.1; Day 3 p=0.3).

K. Number of reward beam breaks during the tone period across days in GFP and Jaws mice (F_{time}(1.9, 32.1) = 2.5, p=0.09; F_{opsin}(1, 33) = 5.66, p=0.9; $F_{interaction}$(2, 33) = 0.2, p=0.7). E-K statistical testing performed with a mixed effects model, n=7 GFP, n=7-8 Jaws.

Values are mean +/- SEM. Posthoc testing performed with Tukey's test.

Figure 3. Optogenetic inhibition of VTA-mPFC DA terminals during PMA without motivational conflict.

A. Schematic of viral and fiber optic targeting locations.

B. Protocol for optogenetic inhibition during PMA training. On training day, 635nm laser light was presented coincidently with a 30 second 4kHz tone that co-terminated with a 2-second footshock. On retrieval day, mice were presented with 10 tones without shocks or laser.

C. Fraction of successful trials across days in control vs. Jaws mice (F_{time}(4.323, 49.71) = 16.66, P<0.0001; F_{opsin}(1, 13) = 5.092, p=0.04; $F_{interaction}$(8, 92) = 2.197, p=0.03; Mixed effects model, N=7 Jaws, N=8 control). Fraction of escape trials across days in control vs. Jaws mice (F_{time} (3.975, 44.71) = 2.33, p=0.07; F_{opsin} (1, 12) = 1.68, p=0.22; $F_{interaction}$ (8, 90) = 3.09; p=0.004. Mixed effects model, N=7 Jaws, N=8 control mice). Fraction time freezing during tone across days in control vs. Jaws mice (F_{time} (4.39, 46.6) = 8.39, P<0.0001; F_{opsin}(1, 11) = 0.60 p=0.46; $F_{interaction}$ (8, 85) = 0.88; p=0.54. Mixed effects model, N=7 Jaws, N=7 control mice).

D. Fraction time on platform and fraction of time freezing during tones on retrieval day (platform: p=0.002, freezing: p=0.6, two-tailed t-test, n=10 control, N=8 Jaws mice).

E. Protocol for optogenetic inhibition during PMA retrieval. Mice are trained in PMA without laser inhibition. The next day, mice are presented with 10 randomly interleaved tones, half of which are paired with constant 635 nm light.

F. Successful trials during PMA training.

G. Percent time on platform and percent time freezing during the tone on interleaved light on and light off trials.

*p<0.05, **p<0.01. Plots show mean +/- SEM.

Figure 4. mPFC DA reflects but is not required for associative learning in cued fear conditioning.

A. AAV injection and fiber placement for optogenetic inhibition of VTA-mPFC axon terminals.

B. Experimental protocol.

C. Freezing during fear conditioning did not differ between GFP and Jaws groups. Two-way repeated measures ANOVA (F_{time} (4.7, 33.3)=18.2, P<0.001; F_{opsin} (1, 7)=0.6, p=0.4; $F_{interaction}$ (14, 98)=0.5, p=0.9).

D. Freezing during fear memory retrieval did not differ between GFP and Jaws groups. Two-way RM ANOVA (F_{time} (2.8, 20.2)=5.5, p=0.006; F_{opsin} (1, 7)=1.2, p=0.2; $F_{interaction}$ (5, 35)=0.8, p=0.5).

* P<0.05, ** P<0.01.