AgeML: Age modelling with Machine Learning

Jorge Garcia Condado, Iñigo Tellaetxe Elorriaga, Jesus M. Cortes, Asier Erramuzpe

Abstract— **A successful approach to age modeling involves the supervised prediction of age using machine learning from subject features. Used for exploring the relationship between healthy and pathological ageing in brain and multiple body systems, as well as the interactions between them, we lack a standard for prediction of age from any generic system. In this work we developed AgeML, an OpenSource software for age-prediction following well-established and tested methodologies from any type of tabular clinical data. The objective is to set standards for reproducibility and standardization of reporting in supervised age modelling tasks. AgeML allows for modelling age and calculating age deltas, the difference between predicted and chronological age, measuring correlations between age deltas and factors, visualising differences in age deltas of different clinical populations and classifying clinical populations based on age deltas. Using the software AgeML, we're demoing its capabilities on a hybrid dataset, reproduce published work, and unveil novel relationships between body organs and polygenetic risk scores. AgeML made easy for standardization and reproducibility.**

Index Terms— **Brain Age, Ageing, Machine Learning, Open Source, Standardization, Reproducibility, Polygenetic Risk Scores**

I. INTRODUCTION

Age modelling can be implemented by supervised machine learning regression in which clinical, behavioural, cognitive or brain/body morphological features of a subject are used to predict their chronological age [1]. Age models can be built using body organ specific features to create organ specific models, or organ ages [1]. Models have been built for many organs such as the brain [2]–[4], heart [5], eye [6], but also

The project that gave rise to these results received the support of a fellowship from "la Caixa" Foundation (ID 100010434) to JGC. The fellowship code is LCF/BQ/DI21/11860030. JMC is funded by Ikerbasque: The Basque Foundation for Science, the Department of Economic Development and Infrastructure of the Basque Country (Elkartek Program, grant KK-2021-00009), and the Health Department of the Basque Country (grants 2022111031 and 2023111002). AE is supported by Ikerbasque, the Basque Foundation for Science, and the Spanish Ministry of Science and Innovation, grant RYC2021-032390-I. *(JCG and ITE contributed equally to this work) (Corresponding author: AE)*

JGC and ITE are with the Computational Neuroimaging Lab, Biobizkaia Health Research Institute, Barakaldo, Spain, 48903 and the Biomedical Research Doctorate Program, University of the Basque Country, Leioa, Spain, 48940.

JMC is with the Computational Neuroimaging Lab, Biobizkaia Health Research Institute, Barakaldo, Spain, 48903 and the Department of Cell Biology and Histology, University of the Basque Country, Leioa, Spain, 48940 and IKERBASQUE, the Basque Foundation for Science, Bilbao, Spain, 48009.

AE is with the Computational Neuroimaging Lab, Biobizkaia Health Research Institute, Barakaldo, Spain, 48903 and IKERBASQUE, the Basque Foundation for Science, Bilbao, Spain, 48009. (email: asier.erramuzpe@gmail.com)

from cognitive performance [7]. These models can integrate information from a single source of data or from multiple sources, thereby facilitating the development of multi-modal age models. The emergence of such age models is attributed to the accessibility of large-scale healthcare databases in the recent decade, exemplified by the UKBiobank [8].

Researchers are focused on elucidating the age gap or delta, the discrepancy between the predicted age and the chronological age of a subject. Age models are trained on two different types of populations: healthy participants devoid of chronic disorders, or on a sample representative of the general population. These deltas, representing the residuals in the regression task, have been shown to correlate with lifestyle factors. Participants engaging in unhealthy lifestyle practices, including smoking, excessive alcohol consumption, or sedentary behavior, exhibit elevated deltas [1]. Age deltas have also been recently associated to specific genetic mutations [9]. The variation in age deltas across organs at different time points, and their multi-variate integration, can help us understand large-scale physiological multi-organ networks [1], of special relevance for the field of network physiology. Participants diagnosed with diseases exhibit higher deltas in specific organs; for example, variations in brain age deltas can distinguish individuals with a neurodegenerative disease such as Alzheimer's disease from healthy controls [10]. However, from the practical perspective of implementing models and calculating deltas, there still exist many problems and questions in this emerging multidisciplinary field, particularly concerning the implementation, standardization, and reproducibility across institutions.

AgeML is an OpenSource Python package that makes age modelling with machine learning easy. AgeML has three main goals: standardization, reproducibility and easy access. Standardization is the first element necessary to enable the comparison of studies. Reproducibility is crucial for fostering trust within the community and ensuring the consistency of results. It is vital to provide an easy start for individuals interested in age modelling, particularly for clinicians with potentially less theoretical backgrounds in machine learning, is essential. We believe that AgeML could serve as a foundational platform for addressing the research questions within the field, led by developers and researchers themselves.

II. RELATED WORK

Previous works on machine learning based age modelling have focused on understanding whether these deviations from a healthy ageing trajectory come from a sudden change, an increase in the delta at a specific time point, or an accumulation of small changes [11]. A debate exists regarding

Fig. 1. **Age modelling with Machine Learning (AgeML), general overview, and major capabilities.** *Green panels:* the `model_age` module requires only a features file to operate, which should contain the chronological age of the subject and the set of features $\mathbf{x}_i$ for each individual. A system file consisting of sets of features (e.g.: a set that encompasses the musculoskeletal system and/or another set referring to renal features) can be provided to train an age prediction model for each specific system. For training a model that distinguishes participants based on a categorical value, a covariates file can be provided, enabling the performance of covariate correction. If a clinical file is provided indicating which participants are healthy controls, the age prediction model is exclusively trained using their features. This pipeline displays the age distribution of the healthy controls, the Pearson correlation (ρ) and significance of each feature and chronological age (separated by system and covariate as required), and the predicted age as well as the bias corrected age. Finally, an ages file in CSV format is generated, incorporating the participant's chronological ages, their predicted uncorrected, corrected ages and the delta, the difference between corrected and chronological age. The age prediction pipeline includes the following components: a scaler selection step, an optional feature extension step, a model selection step, and the availability of optional hyperparameter optimization. *Red panel:* the `factor_correlations` module analyzes the Pearson correlation ρ strength and significance between the specified numerical factors (e.g.: weekly exercise hours, years of education) and the calculated age delta. Additionally, covariate correction can be applied, and this analysis can be performed for each clinical group as needed. *Blue panel:*, the `clinical_groups` module creates a box plot illustrating the age deltas for each clinical group, reporting the FDR and Bonferroni corrected T-test p-values and T-statistics of all groups. Although not depicted in the figure, the age distributions of all clinical groups are plotted, and a T-test is performed to assess group differences, reporting their p-values. *Purple panel:* The `clinical_classify` module employs Logistic Regression with *n*-fold cross-validation and a defined sigmoid threshold to differentiate between two clinical categories identified in the clinical file. This classification is exclusively based on the age deltas within the ages file. If the ages file comprises multiple systems, individual models are trained for each system, and a comprehensive model encompassing all system-specific age deltas is also trained.

whether research should prioritize minimizing the error in age estimation tasks or, alternatively, focus on developing models that are generalizable for subsequent classification tasks [12]–[14], ie., differentiating between individuals at higher risk of developing a pathological condition. There is also large variability between reported average deltas of disease groups across different studies [15]. Caution is essential in selecting training data, as it predominantly comes from white populations, and biases have been observed when applying these models to non-white populations [16]. Biases have also been found when applying a model across different age ranges than those it was trained in [17]. To tackle all these open questions, standardization in age modeling procedures is needed.

A previous effort aimed at developing a tool specific for brain age modeling is the Brain Age Standardized Evaluation (BASE) initiative [18]. BASE focuses on age modelling from brain magnetic resonance images, encompassing various scenarios such as scanning at multi-site, generalize to unseen site, test-retest, and longitudinal data. It also employs an evaluation protocol involving repeated training of brain age prediction models, assessing performance metrics for accuracy and robustness, to ultimately ensure reproducibility. AgeML expands upon these functionalities, enabling multimodal inputs of different organ features, which aligns with a more contemporary approach of system medicine viewing the body as a network of interacting organs. This makes AgeML more accessible to a broader research community. Moreover, AgeML is open-source developed with transparency to ensure that standards are led by the community. Additionally, AgeML is specifically focused on analysis procedures for understanding clinical populations, thereby offering added value to the medical research community.

III. METHODS

AgeML is developed as a Python package readily available to install through PyPI [19]. An overview of all AgeML functionalities can be seen in Fig. 1. There are three main strategies to work with AgeML. The first one is a simple command line interface that can be launched

after installation by using the command `ageml`. This interface interactively asks for the required individual inputs at each different stage. The second strategy is by using the four available commands, each for a different pipeline: `model_age` for age modelling, `factor_correlation` to study the relationships between deltas and clinical outcomes, `clinical_groups` to study the deltas of different clinical groups and `clinical_classify` to use deltas as predictors in a classification task. Lastly, `ageml` can be used as an importable Python library, allowing for the implementation of custom functionality. The Open Source code can be found in GitHub: https://github.com/compneurobilbao/ageml.

A. Age Modelling

Age modeling can be effectively applied through supervised machine learning regression, where subject features are employed to predict known chronological age of the subject (Φ), corresponding to the subject's birth date. Let Ω denote the predicted age, and $\mathbf{X}$ represent the model features. The aim us to find $f()$, our age modeling pipeline, such that:

$$\Omega = f(\mathbf{X}). \tag{1}$$

The pipeline consists of three steps: a scaler, a feature extension step, and a model. The available scalers are all imported from the Scikit-learn Python library [20] and include the max absolute scaler, min-max scaler, normalizer, power transformer, quantile transformer, robust scaler, and standard scaler. For feature extension, the polynomial feature extension from Scikit-learn [20] is used. The various models implemented consist of Linear Regression, Ridge, LASSO, Support Vector Regression (SVR) and Random Forests (RF); and XGBoost [21].

The pipeline is trained by minimizing the mean squared error between the predicted and chronological age. The pipeline undergoes training via cross-validation, with a default of 5-folds. The reported metrics are derived from these cross-validation folds. Users have the option to configure the hyperparameters of the model manually, or they can choose for hyperparameter tuning. In the case of hyperparameter tuning, a grid search is conducted alongside cross-validation, and users can specify the number of points to search for.

A well-known bias in age modelling is the tendency for younger participants to be assigned higher ages, while older healthy controls are assigned lower ages than their actual ages. This phenomenon is attributed to a regression to the mean problem [17], [22], [23]. This bias can be conceptualized as our best estimate for a subject about whom we have no prior information being the mean age. The predicted age bias can be can be corrected [22], by taking into account the actual participant's age through fitting the following linear model :

$$\Omega = \alpha \times \Phi + \beta. \tag{2}$$

The coefficients α and β correspond to the slope and intercept of the linear model. The training set is used to fit these coefficients α and β. Selecting the appropriate method for age bias correction is of paramount importance to prevent the introduction of new biases, such as inflating correlation effects. An alternative method to that used in this study is to use age as a covariate when studying group differences [22]. The fitted α and β are then used to correct the predictions in a test set using:

$$\hat{\Omega} = \Omega + [\Phi - \alpha \times \Phi + \beta]. \tag{3}$$

Research in this context typically focused on the age gap or delta (δ), which is defined as the difference between the subject's chronological age and the predicted age:

$$\delta = \hat{\Omega} - \Phi. \tag{4}$$

AgeML provides the capability to simultaneously train distinct models for each system defined, thereby obtaining age deltas for each system. This is achieved by specifying which features belong to each system. Additionally, separate models can be trained based on a covariate, such as sex, with AgeML training distinct models for males and females separately.

Users also have the option to specify which participants are considered as healthy controls for model training purposes. The model is then trained exclusively on these designated participants and subsequently applied to calculate deltas for the remaining participants. The predicted age (Ω), the corrected age ($\hat{\Omega}$) and the age delta (δ) are saved and can be used as inputs for further steps or for independent further analysis by the researcher, referred as the *ages* file.

To ensure standardization and reproducibility, after each model is trained, both training and test metrics are stored. These metrics emcompass Mean Absolute Error (MAE), Root Mean Square Error (RMSE), R^2, and Pearson's correlation (ρ). Recognizing that MAE and RMSE are strongly influenced by the age range of the training data, an additional metric referred to as the *dummy regressor* (DR) is also calculated, by obtaining MAE and RMSE using the training mean predicted age as the predictor for each individual subject, thus somehow normalizing this effect.

B. Factor Correlation

Previous research has addressed the relationship between the deltas and other external factors such as lifestyle choices or genetics. To visualize and quantify this relations, a Pearson correlation is calculated between each factor and each delta as well as the corresponding p-values. To account for multiple comparisons, p-values can be adjusted using either Bonferroni or False Discovery Rate (FDR) corrections. Additionally, users have the option to provide covariates and remove covariate effects from the factors prior to conducting the correlation analysis. The covariate correction is calculated as the coefficients in the maximum likelihood solution of a general linear model, ie.,

$$\mathbf{\Gamma} = (\mathbf{Z}^T \mathbf{Z})^{-1} \mathbf{Z}^T \mathbf{\Theta}. \tag{5}$$

where $\mathbf{\Theta}$ represents the subject factors and $\mathbf{Z}$ the covariates. Notice that $\mathbf{\Gamma}$ is calculated using only the healthy controls. In cases where there are distinct clinical populations, the same $\mathbf{\Gamma}$ coefficients calculated for the healthy controls are used

also for covariate correction in the clinical populations. The participant's covariate-corrected factors are then given by:

$$\hat{\Theta} = \Theta - Z\Gamma. \qquad (6)$$

The correlation analysis between factors and deltas is subsequently carried out using the corrected factors rather than the non-corrected ones.

C. Clinical Groups

AgeML offers the capability to examine the participant's age deltas labeled into various clinical populations. It assesses the distributions of age deltas for each provided clinical population, displaying the mean and standard deviation for each group. Additionally, AgeML determines whether the delta distributions exhibit significant differences between groups, with the option to apply Bonferroni or FDR correction for multiple comparisons. Similar to the factor correlation analysis, covariate correction can be applied to the deltas previous to calculating distributions for each clinical group.

AgeML also integrates a quality control step to check whether the age distributions of participants within each clinical group exhibit significant differences as training on a specific distribution and subsequently making predictions in a different one with a different age can potentially introduce biases [17]. This is particularly evident when considering young and old populations, as many phenotypes follow quadratic trends, as observed in brain volumes [24], and cognitive performance. Hence, training on a young population and applying the model to an older one can be problematic and susceptible to bias.

D. Clinical Classification

In scenarios involving multiple organ-age systems, we represent $\Delta \equiv \delta_i$ as the collection of deltas for each subject, and use them as the input for a classifier designed to categorize participants into various clinical groups. Employing a cross-validation strategy, a logistic regressor is trained to differentiate between two specific labels. The sigmoid function is then given by $\sigma(\Delta) = 1/\left(1 + e^{-\beta_0 - \sum_i \beta_i \delta_i}\right)$, that after fitting the logistic regressor, the coefficient β_i accounts for the contribution of δ_i in the classification. The vector $\beta \equiv \beta_i$, previously normalized between 0 and 1 by dividing each component by the maximum β_i, is stored and printed to assess which organ delta participates with a higher relevance in the classification task.

E. Data

The UKBiobank database [8] was used for this project (pre-approved ID 79105). Overall, UKBiobank consists of a total of 502,504 individuals, of which 229,122 are males. These participants, aged between 37 and 73 years during recruitment (2006–2010), underwent thorough assessments, including questionnaires, physical examinations, blood and urine sample analyses, and genome-wide genotyping. The extensive data collection occurred across 22 assessment centers located throughout the United Kingdom.

Body phenotypes: We created age models using 78 different body phenotypes, which align with those in [1]. The different body phenotypes were grouped into seven distinct organ systems: Musculoskeletal, Cardiovascular, Pulmonary, Immune, Renal, Hepatic, and Metabolic. These groups were used to create distinct age models for each organ system. The 78 distinct phenotypes were also considered as features to create an unified age model, referred to as Body.

Clinical categories: We established them using ICD-10, ICD-9 codes and self-reported pathologies, translated into clinical categories by using the Clinical Classifications Software Refined [25]. Data was used based only on first occurrence labels. The distinction between chronic and non-chronic participants was established using the Chronic Condition Indicator Refined [26], resulting in a healthy population of N=6,388. Eight clinical categories were finally selected, in relation with the organ systems previously defined: blood diseases (BLD), circulatory system diseases (CIR), endocrine, nutritional and metabolic diseases (END), mental, behavioral and neurodevelopmental disorders (MBD), Neoplasms (NEO), diseases of the musculoskeletal system (MUS), diseases of the nervous system (NVS), respiratory system diseases (RSP). Participants with multiple comorbidities were excluded to mitigate potential mixed effects.

Polygenetic Risk Scores (PRS): To investigate the relationship between organ deltas and genetics, if any, a factor correlation analysis was carried out with PRS derived from the genetic data of the UKBiobank. Two types of PRS were used: Standard PRS and Enhanced PRS [27]. Standard PRS are sourced from external UKBiobank GWAS data, while Enhanced PRS are derived from a combination of external and internal UK Biobank data. Notably, Enhanced PRS encompass a larger number of categories (53) compared to Standard PRS (39). Standard PRS were available for 6,323 out of the 6,388 participants, whereas Enhanced PRS were only accessible for 1,234 of these individuals. Prior to conducting the factor correlation analysis, the PRS were adjusted for the covariates recruitment site, sex, and ethnicity.

TABLE I
AGE DISTRIBUTION SUMMARY OF THE CLINICAL CATEGORIES

CLINICAL CATEGORY	$\mu_{age} \pm \sigma_{age}$ (YEARS)	AGE RANGE (YEARS)	N (PARTICIPANTS)
CN	51.39 ± 7.31	[40-70]	6,388
CIR	56.00 ± 7.90	[40-70]	3,907
END	55.43 ± 7.78	[40-70]	1,616
MBD	50.86 ± 7.04	[40-70]	1,184
MUS	54.11 ± 7.67	[39-70]	5,603
NVS	53.10 ± 7.63	[39-70]	983
RSP	51.41 ± 7.57	[40-70]	4,376
NEO	56.09 ± 7.75	[40-70]	2,215
BLD	51.76 ± 7.33	[40-69]	338

CN: Healthty Controls, CIR: Circulatory system disease, MBD: neurodevelopmental disorders, MUS: Musculoskeletal system disorders, RSP: respiratory system diseases, NEO: neoplasms, BLD: blood diseases

Additional information regarding the total number of participants per category and their age distributions can be found in Table I.

TABLE II

MEAN ABSOLUTE ERROR (MAE) AND PEARSON'S CORRELATION (ρ) OF SVR MODELS FOR PREDICTING THE CHRONOLOGICAL AGES OF HEALTHY CONTROLS USING DIFFERENT PHYSIOLOGICAL SYSTEMS, COMPARING THE WORK OF TIAN ET AL. 2023 [1] WITH OURS.

METRIC	GENDER	MODEL	BODY	CARDIOVASCULAR	PULMONARY	MUSCULOSKELETAL
MAE*	FEMALE	OURS	3.86 ± 0.21	5.380 ± 0.340	5.34 ± 0.27	4.98 ± 0.17
		TIAN ET AL. 2023 [1]	3.708 ± 0.0015	5.620 ± 0.0016	5.675 ± 0.0008	5.267 ± 0.0010
	MALE	OURS	4.61 ± 0.21	5.80 ± 0.25	5.55 ± 0.23	5.61 ± 0.22
		TIAN ET AL. 2023 [1]	4.463 ± 0.0016	6.323 ± 0.0011	6.130 ± 0.0008	6.052 ± 0.0012
ρ^*	FEMALE	OURS	0.753 ± 0.034	0.424 ± 0.065	0.443 ± 0.064	0.519 ± 0.045
		TIAN ET AL. 2023 [1]	0.787 ± 0.0002	0.436 ± 0.0004	0.421 ± 0.0003	0.513 ± 0.0002
	MALE	OURS	0.638 ± 0.033	0.296 ± 0.072	0.382 ± 0.051	0.372 ± 0.061
		TIAN ET AL. 2023 [1]	0.715 ± 0.0002	0.317 ± 0.0004	0.375 ± 0.0003	0.403 ± 0.0003

METRIC	GENDER	MODEL	IMMUNE	RENAL	HEPATIC	METABOLIC
MAE*	FEMALE	OURS	5.70 ± 0.22	5.16 ± 0.24	5.13 ± 0.29	4.97 ± 0.24
		TIAN ET AL. 2023 [1]	5.965 ± 0.0019	5.220 ± 0.0012	5.466 ± 0.0010	5.302 ± 0.0022
	MALE	OURS	5.78 ± 0.22	5.41 ± 0.31	5.75 ± 0.25	5.85 ± 0.24
		TIAN ET AL. 2023 [1]	6.330 ± 0.0022	5.830 ± 0.0011	6.294 ± 0.0012	6.514 ± 0.0019
ρ^*	FEMALE	OURS	0.284 ± 0.087	0.475 ± 0.069	0.476 ± 0.084	0.524 ± 0.064
		TIAN ET AL. 2023 [1]	0.303 ± 0.0005	0.532 ± 0.0002	0.451 ± 0.0004	0.498 ± 0.0005
	MALE	OURS	0.297 ± 0.067	0.437 ± 0.058	0.308 ± 0.065	0.247 ± 0.065
		TIAN ET AL. 2023 [1]	0.301 ± 0.0007	0.453 ± 0.0003	0.322 ± 0.0004	0.230 ± 0.0009

* the reported $\pm$ are 95% confidence intervals for our work and Tian et al. 2023 [1] report standard deviations.

IV. RESULTS

A. Reproducibility

To demonstrate that AgeML can replicate previous research findings, we chose to replicate the study by Tian et al. [1], in which brain age models are presented in combination with multiple organ system age models. The different age models are trained on healthy participants with no known chronic disease. The feature set consists of the same 78 body phenotypes described in Methods, and participants with any missing phenotypes are excluded from the analysis. Training is carried out using a Support Vector Regressor (SVR) with a 20-fold cross-validation scheme (with a linear kernel, a regularization parameter of 1.0 and a gap tolerance of 0.001). Comparisons of the results between AgeML (Ours) and the one in Tian et al. 2023 [1] are given in Table II.

AgeML results within 95% confidence intervals recovered all the previous results reported in Tian et al. 2023 [1], despite some minor discrepancies possibly attributed to the differences in sample sizes (Ours N=6,388; Tian's N=28,589) due to differences in inclusion criteria, primarily attributed to the more stringent definition of chronic conditions in our study.

B. Genetic Analysis

PRSs measure genetic risks for various diseases, and some diseases are closely related to specific organs, possibly influencing organ age. For example, it is expected that genetic predisposition to factors affecting the heart, such as cardiovascular diseases, coronary artery diseases, ischemic stroke, or hypertension, would lead to higher cardiovascular ages. Similarly, protective genetic factors that promote stronger bones with higher mineral density can lead to a reduction in musculoskeletal age. Here, and for the group of healthy

participants, the correlation between age deltas for each organ and both Standard PRS and Enhanced PRS were compared, cf. Fig 2 shows statistically significant correlations between PRS and organ deltas (detailed correlation values and corresponding p-values are given in code repository). Very remarkably, for the results corresponding to enhanced PRS, we found that many metabolic genetic propensities negatively impact metabolic age, and in contrast, a protective effect of glomerular filtration. The differences in statistical significance between Standard PRS and Enhanced PRS may be due to the lower number of participants available for analysis in the latter.

C. Clinical data

TABLE III

DISEASE CATEGORIES THAT HAVE A STATISTICALLY SIGNIFICANT DIFFERENCE IN THEIR AGE DELTA COMPARED TO HEALTHY CONTROLS, FOR DIFFERENT SYSTEMS (BONFERRONI CORRECTED, $p < 0.05$).

SYSTEM DELTA	CLINICAL CATEGORY DIFFERENT TO CN				
CARDIOVASCULAR	CIR	MBD	MUS	RSP	NEO
PULMONARY	CIR	MBD	RSP	NEO	
METABOLIC	MUS	NEO			
IMMUNE	MBD				

CN: Healthty Controls, CIR: Circulatory system disease, MBD: neurodevelopmental disorders, MUS: Musculoskeletal system disorders, RSP: respiratory system diseases, NEO: neoplasms, BLD: blood diseases

Deltas for each organ systems were calculated for all participants. The average deltas for each clinical group and organ were calculated, and then compared across groups (Table III). The significance of these differences indicates that the participants belonging to clinical groups related to specific organs exhibit significant variations as compared to healthy

Fig. 2. **Factor correlation analysis between Polygenetic Risk Scores (PRS) and system-specific age deltas**. Each link represents a statistical significant Bonferroni-corrected Pearson's correlation ($p < 0.05$) and only those with $|\rho| > 0.05$ are plotted. The width of the link indicates the strength of the correlation. Red links denote positive correlations, which are associated with higher age deltas or accelerated aging. Blue links, on the other hand, represent negative correlations, suggesting protective or age-slowing effects. The left purple panel shows results for Standard PRSs (6,323 participants), while the right gray panel does it for Enhanced PRSs (1,234 participants). Note that several PRS labels are shared across panels, reassuring the significance of the correlations. All the Pearson's correlation values and their associated p-values can be found in the code repository.

controls. For example, individuals in the clinical category of circulatory diseases show a higher delta for the cardiovascular system compared to healthy controls. Detailed information about the differences in average deltas and their corresponding p-values between clinical groups and organs can be found in the code repository.

Logistic regressor classifiers were trained to differentiate between healthy controls and each clinical category based on all organ deltas as input. Fig 3 displays the logistic regressor weights, which have been normalized for each case. Similar to the differences observed in average deltas for organ systems in specific clinical categories, the most discriminant features between healthy controls and clinical categories related to specific organs are the deltas for those respective organs. Thus for instance, when classifying between healthy controls and cardiovascular disease, the most discriminant feature is the cardiovascular delta. To avoid biases in predictions, the majority class was under-sampled to match the size of the minority class.

V. DISCUSSION

The first step towards standardization is showing the reproducibility of previous findings. In this context, we have replicated the study presented by Tian et al. [1]. AgeML facilitates standardization by providing a tool designed for use with explicitly defined pipelines. Moreover, the reporting of

Fig. 3. **Normalized Logistic Regressor weights of system-specific deltas for classification between healthy controls versus a specific disease category**. The highest values for each column indicate the most relevant system-specific delta used by the Logistic Regressor for classifying healthy controls against the column disease category. Normalization is done per column (classifier), dividing the absolute values by the absolute highest one. Thus, the cells with the value 1.0 correspond to the absolute highest weight per column. In the situation of class imbalance, the majority class was undersampled to match the minority class occurrences.

standarized metrics by AgeML constitutes a crucial issue for usability. AgeML aims to establish the minimum guidelines for clear communication of training outcomes.

Previous research like BASE [18] focused on brain age modeling has shown the need for standardization and reproducibility. While BASE is tailored to brain age modeling, it highlights the significance of integrating standards into the field. AgeML not only replicates many of the benchmarks established in BASE with respect to metric reporting but

also extends it further by enhancing ease of access, embracing open-source development, and demonstrating applicability across various types of clinical or organ data.

We have shown that genetic information, specifically Polygenic Risk Scores (PRS), is associated with distinct forms of organ aging, aligning with prior GWAS studies [9]. This is of critical importance for quantifying the extent to which an individual's organ age-variations can be attributed to genetics. Moreover, it paves the road for exploring therapeutic targets aimed at decelerating aging [9]. A logical continuation of this work is to develop organ-specific PRS based on genomic data to assess genetic predisposition to either significantly rejuvenated or minimally aged organs as a measure of aging resilience. Efforts in this direction are already in progress, notably in the context of brain age PRS [28].

As expected the age deltas of each organ systems vary among clinical populations. This variation is particularly evident in the deltas associated with the cardiovascular system and its corresponding clinical category of cardiovascular disease. Similar to previous work [1], our results confirm that while large age deltas are not exclusive to any single disease, distinct patterns of organ deltas are associated with each specific disease.

The practical integration of age modeling in clinical environments is currently in a conceptual phase. An initial proposition involves using age deltas as a flag, where during routine analyses, these deltas are computed, and an alert is triggered if any of the calculated deltas are high (previously determined a normal or standardized distribution of deltas). Although organ deltas lack specificity to particular diseases (eg., patients with Parkinson's, Alzheimer's, or depression might exhibit an elevated brain delta), they can alert clinicians to potential deviations from healthy aging normative patterns. By computing deltas for various organs and achieving a multimodal perspective, it is also feasible to mitigate the issue of non-specificity, thereby enhancing the feasibility of age deltas in clinical settings.

AgeML is a first approach to setting a standard in the community and a minimal software tool for basic age modelling. The AgeML project will continue by integrating more features. Currently AgeML supports only tabular data; however, future plans aim to incorporate additional data modalities, including imaging data (e.g., MRI or CT scans) routinely used in medical settings, as well as time series data, such as electrocardiograms (ECG). The next step involves integrating foundational models that have been trained on extensive datasets, such as the UKBiobank. Following that, the strategy includes providing transfer learning capabilities to retrain models with local datasets. In medical contexts, the demand for interpretability is paramount and should be addressed. The factor correlation pipeline could be upgraded to introduce causal analysis. Ultimately, it is anticipated that community-driven efforts will emerge, aiming to incorporate a broader range of desired features and improvements.

VI. CONCLUSION

We have shown that AgeML is capable to replicate prior studies in the field, thereby paving the way for standardization of age modeling strategies. We have proven the AgeML efficacy by investigating the interplay between age-related organ variations and genetics, and in particular, by assessing the correlation between Polygenic Risk Scores and specific organ deltas. Consistent with previous work, statistical significant differences in organ deltas occur between healthy controls and clinical populations, especially when focusing on organ-systems related to specific diseases.

The ultimate objective of our study extends beyond merely developing a tool; it aims to trigger a community of developers who contribute new features and help lead towards age-modeling standardization. The major goal of AgeML is to standardize procedures, reduce the barriers to entry for age modeling, and guarantee the reproducibility of research findings. The AgeML project is Open-Source, designed to foster a welcoming environment and a community to work together to improve and validate existing methodologies. We are actively inviting new developers interested in contributing to the growth and expansion of the package.

REFERENCES

[1] Y. E. Tian, V. Cropley, A. B. Maier, N. T. Lautenschlager, M. Breakspear, and A. Zalesky, "Heterogeneous aging across multiple organ systems and prediction of chronic disease and mortality," *Nature Medicine*, Apr. 2023. [Online]. Available: https://www.nature.com/articles/s41591-023-02296-6

[2] K. Franke, G. Ziegler, S. Klöppel, and C. Gaser, "Estimating the age of healthy subjects from T1-weighted MRI scans using kernel methods: Exploring the influence of various parameters," *NeuroImage*, vol. 50, no. 3, pp. 883–892, Apr. 2010. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S1053811910000108

[3] P. Bonifazi, A. Erramuzpe, I. Diez, I. Gabilondo, M. P. Boisgontier, L. Pauwels, S. Stramaglia, S. P. Swinnen, and J. M. Cortes, "Structure–function multi-scale connectomics reveals a major role of the fronto-striato-thalamic circuit in brain aging," *Human Brain Mapping*, vol. 39, no. 12, pp. 4663–4677, 2018. [Online]. Available: https://onlinelibrary.wiley.com/doi/abs/10.1002/hbm.24312

[4] A. Erramuzpe, R. Schurr, J. D. Yeatman, I. H. Gotlib, M. D. Sacchet, K. E. Travis, H. M. Feldman, and A. A. Mezer, "A Comparison of Quantitative R1 and Cortical Thickness in Identifying Age, Lifespan Dynamics, and Disease States of the Human Cortex," *Cerebral Cortex (New York, N.Y.: 1991)*, vol. 31, no. 2, pp. 1211–1226, Jan. 2021.

[5] A. O. Ladejobi, J. R. Medina-Inojosa, M. Shelly Cohen, Z. I. Attia, C. G. Scott, N. K. LeBrasseur, B. J. Gersh, P. A. Noseworthy, P. A. Friedman, S. Kapa, and F. Lopez-Jimenez, "The 12-lead electrocardiogram as a biomarker of biological age," *European Heart Journal - Digital Health*, vol. 2, no. 3, pp. 379–389, Sep. 2021. [Online]. Available: https://academic.oup.com/ehjdh/article/2/3/379/6248088

[6] J. Ma, X. Xu, M. Li, Y. Zhang, L. Zhang, P. Ma, J. Hou, Y. Lei, J. Liu, X. Huangfu, Y. Yang, X. Yi, G. Cheng, J. Bai, X. Zhong, X. Xu, and Y. Wang, "Predictive models of aging of the human eye based on ocular anterior segment morphology," *Journal of Biomedical Informatics*, vol. 120, p. 103855, Aug. 2021. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S1532046421001842

[7] J. Garcia Condado, J. M. Cortes, and for the Alzheimer's Disease Neuroimaging Initiative, "NeuropsychBrainAge: A biomarker for conversion from mild cognitive impairment to Alzheimer's disease," *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, vol. 15, no. 4, p. e12493, Oct. 2023. [Online]. Available: https://alz-journals.onlinelibrary.wiley.com/doi/10.1002/dad2.12493

[8] C. Sudlow, J. Gallacher, N. Allen, V. Beral, P. Burton, J. Danesh, P. Downey, P. Elliott, J. Green, M. Landray, B. Liu, P. Matthews, G. Ong, J. Pell, A. Silman, A. Young, T. Sprosen, T. Peakman, and R. Collins, "UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age," *PLOS Medicine*, vol. 12, no. 3, p. e1001779, Mar. 2015. [Online]. Available: https://dx.plos.org/10.1371/journal.pmed.1001779

[9] J. Wen, Y. E. Tian, I. Skampardoni, Z. Yang, E. Mamourian, F. Anagnostakis, B. Zhao, A. W. Toga, A. Zaleskey, and C. Davatzikos, "The Genetic Architecture of Biological Age in Nine Human Organ Systems," Genetic and Genomic Medicine, preprint, Jun. 2023. [Online]. Available: http://medrxiv.org/lookup/doi/10.1101/2023.06.08.23291168

[10] C. Gaser, K. Franke, S. Klöppel, N. Koutsouleris, H. Sauer, and Alzheimer's Disease Neuroimaging Initiative, "BrainAGE in Mild Cognitive Impaired Patients: Predicting the Conversion to Alzheimer's Disease," PLoS ONE, vol. 8, no. 6, p. e67346, Jun. 2013. [Online]. Available: https://dx.plos.org/10.1371/journal.pone.0067346

[11] D. Vidal-Pineiro, Y. Wang, S. K. Krogsrud, I. K. Amlien, W. F. Baaré, D. Bartres-Faz, L. Bertram, A. M. Brandmaier, C. A. Drevon, S. Düzel, K. Ebmeier, R. N. Henson, C. Junqué, R. A. Kievit, S. Kuhn, E. Leonardsen, U. Lindenberger, K. S. Madsen, F. Magnussen, A. M. Mowinckel, L. Nyberg, J. M. Roe, B. Segura, S. M. Smith, O. Sorensen, S. Suri, R. Westerhausen, A. Zalesky, E. Zsoldos, K. B. Walhovd, and A. Fjell, "Individual variations in 'brain age' relate to early-life factors more than to longitudinal brain change," eLife, vol. 10, p. e69995, Nov. 2021. [Online]. Available: https://elifesciences.org/articles/69995

[12] V. M. Bashyam, G. Erus, J. Doshi, M. Habes, I. M. Nasrallah, M. Truelove-Hill, D. Srinivasan, L. Mamourian, R. Pomponio, Y. Fan, L. J. Launer, C. L. Masters, P. Maruff, C. Zhuo, H. Völzke, S. C. Johnson, J. Fripp, N. Koutsouleris, T. D. Satterthwaite, D. Wolf, R. E. Gur, R. C. Gur, J. Morris, M. S. Albert, H. J. Grabe, S. Resnick, R. N. Bryan, D. A. Wolk, H. Shou, and C. Davatzikos, "MRI signatures of brain age and disease over the lifespan based on a deep brain network and 14 468 individuals worldwide," Brain, vol. 143, no. 7, pp. 2312–2324, Jul. 2020. [Online]. Available: https://academic.oup.com/brain/article/143/7/2312/5863667

[13] T. Hahn, L. Fisch, J. Ernsting, N. R. Winter, R. Leenings, K. Sarink, D. Emden, T. Kircher, K. Berger, and U. Dannlowski, "From 'loose fitting' to high-performance, uncertainty-aware brain-age modelling," Brain, vol. 144, no. 3, pp. e31–e31, Apr. 2021. [Online]. Available: https://academic.oup.com/brain/article/144/3/e31/6214475

[14] S. More, G. Antonopoulos, F. Hoffstaedter, J. Caspers, S. B. Eickhoff, and K. R. Patil, "Brain-age prediction: A systematic comparison of machine learning workflows," NeuroImage, vol. 270, p. 119947, Apr. 2023. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S1053811923000940

[15] L. Baecker, R. Garcia-Dias, S. Vieira, C. Scarpazza, and A. Mechelli, "Machine learning for brain age prediction: Introduction to methods and clinical applications," eBioMedicine, vol. 72, p. 103600, Oct. 2021. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S2352396421003935

[16] A. Puc, V. Struc, and K. Grm, "Analysis of Race and Gender Bias in Deep Age Estimation Models," in 2020 28th European Signal Processing Conference (EUSIPCO). Amsterdam, Netherlands: IEEE, Jan. 2021, pp. 830–834. [Online]. Available: https://ieeexplore.ieee.org/document/9287219/

[17] A. G. de Lange, M. Anatürk, J. Rokicki, L. K. M. Han, K. Franke, D. Alnæs, K. P. Ebmeier, B. Draganski, T. Kaufmann, L. T. Westlye, T. Hahn, and J. H. Cole, "Mind the gap: Performance metric evaluation in brain-age prediction," Human Brain Mapping, vol. 43, no. 10, pp. 3113–3129, Jul. 2022. [Online]. Available: https://onlinelibrary.wiley.com/doi/10.1002/hbm.25837

[18] L. Dular and Z. Spiclin, "BASE: Brain Age Standardized Evaluation," NeuroImage, vol. 285, p. 120469, Jan. 2024. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S1053811923006183

[19] PyPI. (2021) Python package index - pypi. [Online]. Available: https://pypi.org/

[20] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and Édouard Duchesnay, "Scikit-learn: Machine learning in python," Journal of Machine Learning Research, vol. 12, no. 85, pp. 2825–2830, 2011. [Online]. Available: http://jmlr.org/papers/v12/pedregosa11a.html

[21] T. Chen and C. Guestrin, "XGBoost: A Scalable Tree Boosting System," in Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. San Francisco California USA: ACM, Aug. 2016, pp. 785–794. [Online]. Available: https://dl.acm.org/doi/10.1145/2939672.2939785

[22] A.-M. G. de Lange and J. H. Cole, "Commentary: Correction procedures in brain-age prediction," NeuroImage: Clinical, vol. 26, p. 102229, 2020. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S2213158220300668

[23] E. R. Butler, A. Chen, R. Ramadan, T. T. Le, K. Ruparel, T. M. Moore, T. D. Satterthwaite, F. Zhang, H. Shou, R. C. Gur, T. E. Nichols, and R. T. Shinohara, "Pitfalls in brain age analyses," Human Brain Mapping, vol. 42, no. 13, pp. 4092–4101, Sep. 2021. [Online]. Available: https://onlinelibrary.wiley.com/doi/10.1002/hbm.25533

[24] D. Dima, A. Modabbernia, E. Papachristou, G. E. Doucet, I. Agartz, M. Aghajani, T. N. Akudjedu, A. Albajes-Eizagirre, D. Alnæs, K. I. Alpert, M. Andersson, N. C. Andreasen, O. A. Andreassen, P. Asherson, T. Banaschewski, N. Bargallo, S. Baumeister, R. Baur-Streubel, A. Bertolino, A. Bonvino, D. I. Boomsma, S. Borgwardt, J. Bourque, D. Brandeis, A. Breier, H. Brodaty, R. M. Brouwer, J. K. Buitelaar, G. F. Busatto, R. L. Buckner, V. Calhoun, E. J. Canales-Rodríguez, D. M. Cannon, X. Caseras, F. X. Castellanos, S. Cervenka, T. M. Chaim-Avancini, C. R. K. Ching, V. Chubar, V. P. Clark, P. Conrod, A. Conzelmann, B. Crespo-Facorro, F. Crivello, E. A. Crone, U. Dannlowski, A. M. Dale, C. Davey, E. J. C. Geus, L. Haan, G. I. Zubicaray, A. Braber, E. W. Dickie, A. Di Giorgio, N. T. Doan, E. S. Dørum, S. Ehrlich, S. Erk, T. Espeseth, H. Fatouros-Bergman, S. E. Fisher, J. Fouche, B. Franke, T. Frodl, P. Fuentes-Claramonte, D. C. Glahn, I. H. Gotlib, H. Grabe, O. Grimm, N. A. Groenewold, D. Grotegerd, O. Gruber, P. Gruner, R. E. Gur, R. C. Gur, T. Hahn, B. J. Harrison, C. A. Hartman, S. N. Hatton, A. Heinz, D. J. Heslenfeld, D. P. Hibar, I. B. Hickie, B. Ho, P. J. Hoekstra, S. Hohmann, A. J. Holmes, M. Hoogman, N. Hosten, F. M. Howells, H. E. Hulshoff Pol, C. Huyser, N. Jahanshad, A. James, T. L. Jernigan, J. Jiang, E. G. Jönsson, J. A. Joska, R. Kahn, A. Kalnin, R. Kanai, M. Klein, T. P. Klyushnik, L. Koenders, S. Koops, B. Krämer, J. Kuntsi, J. Lagopoulos, L. Lázaro, I. Lebedeva, W. H. Lee, K. Lesch, C. Lochner, M. W. J. Machielsen, S. Maingault, N. G. Martin, I. Martínez-Zalacaín, D. Mataix-Cols, B. Mazoyer, C. McDonald, B. C. McDonald, A. M. McIntosh, K. L. McMahon, G. McPhilemy, S. Meinert, J. M. Menchón, S. E. Medland, A. Meyer-Lindenberg, P. Naaijen, P. Najt, T. Nakao, J. E. Nordvik, L. Nyberg, J. Oosterlaan, V. O. Foz, Y. Paloyelis, G. Pergola, E. Pomarol-Clotet, M. J. Portella, S. G. Potkin, J. Radua, A. Reif, D. A. Rinker, J. L. Roffman, P. G. P. Rosa, M. D. Sacchet, P. S. Sachdev, R. Salvador, P. Sánchez-Juan, S. Sarró, T. D. Satterthwaite, A. J. Saykin, M. H. Serpa, L. Schmaal, K. Schnell, G. Schumann, K. Sim, J. W. Smoller, I. Sommer, C. Soriano-Mas, D. J. Stein, L. T. Strike, S. C. Swagerman, C. K. Tamnes, H. S. Temmingh, S. I. Thomopoulos, A. S. Tomyshev, D. Tordesillas-Gutiérrez, J. N. Trollor, J. A. Turner, A. Uhlmann, O. A. Heuvel, D. Meer, N. J. A. Wee, N. E. M. Haren, N. Ent, T. G. M. Erp, I. M. Veer, D. J. Veltman, A. Voineskos, H. Völzke, H. Walter, E. Walton, L. Wang, Y. Wang, T. H. Wassink, B. Weber, W. Wen, J. D. West, L. T. Westlye, H. Whalley, L. M. Wierenga, S. C. R. Williams, K. Wittfeld, D. H. Wolf, A. Worker, M. J. Wright, K. Yang, Y. Yoncheva, M. V. Zanetti, G. C. Ziegler, P. M. Thompson, S. Frangou, and Karolinska Schizophrenia Project (KaSP), "Subcortical volumes across the lifespan: Data from 18,605 healthy individuals aged 3–90 years," Human Brain Mapping, vol. 43, no. 1, pp. 452–469, Jan. 2022. [Online]. Available: https://onlinelibrary.wiley.com/doi/10.1002/hbm.25320

[25] HCUP. (2020) Hcup clinical classifications software refined (ccsr) for icd-10-cm diagnoses, v2021.2. [Online]. Available: www.hcup-us.ahrq.gov/toolssoftware/ccsr/dxccsr.jsp

[26] HCUP. (2021) Hcup chronic condition indicator for icd-10-cm (beta version), v2021.1. [Online]. Available: www.hcup-us.ahrq.gov/toolssoftware/chronic/chronic_icd10/chronic_icd10.jsp

[27] D. J. Thompson, D. Wells, S. Selzam, I. Peneva, R. Moore, K. Sharp, W. A. Tarran, E. J. Beard, F. Riveros-Mckay, C. Giner-Delgado, D. Palmer, P. Seth, J. Harrison, M. Futema, Genomics England Research Consortium, G. McVean, V. Plagnol, P. Donnelly, and M. E. Weale, "UK Biobank release and systematic evaluation of optimised polygenic risk scores for 53 diseases and quantitative traits," Genetic and Genomic Medicine, preprint, Jun. 2022. [Online]. Available: http://medrxiv.org/lookup/doi/10.1101/2022.06.16.22276246

[28] J. Wen, B. Zhao, Z. Yang, G. Erus, I. Skampardoni, E. Mamourian, Y. Cui, G. Hwang, J. Bao, A. Boquet-Pujadas, Z. Zhou, Y. Veturi, M. D. Ritchie, H. Shou, P. M. Thompson, L. Shen, A. W. Toga, and C. Davatzikos, "The Genetic Architecture of Multimodal Human Brain Age," Genetics, preprint, Apr. 2023. [Online]. Available: http://biorxiv.org/lookup/doi/10.1101/2023.04.13.536818