1

CetZ1-dependent assembly and positioning of the motility machinery in
haloarchaea

Hannah J. Brown[1], Md Imtiazul Islam[2], Juanfang Ruan[2,3], Matthew A. B. Baker[2], Solenne
Ithurbide[1,4], Iain G. Duggin[1#]

[1]Australian Institute for Microbiology and Infection (AIMI), University of Technology
Sydney, Ultimo NSW 2007, Australia

[2]School of Biotechnology and Biomolecular Sciences (BABS), University of New South
Wales, Kensington NSW 2052, Australia

[3]Electron Microscope Unit, Mark Wainwright Analytical Centre, UNSW Sydney,
Kensington NSW 2052, Australia

[4]Present address: Département de Microbiologie, Infectiologie et Immunologie,
Université de Montréal, Montréal, QC, Canada.

[#]Correspondence: Iain.Duggin@uts.edu.au

Abstract

The tubulin-like CetZ proteins are archaeal cytoskeletal proteins that contribute to cell
shape and swimming motility in the halophilic archaeon *Haloferax volcanii*. Currently it
is unknown whether CetZs contribute to motility solely through their control of cell shape
or in other ways too. Here, we used cryo-electron and fluorescence microscopy to
observe *H. volcanii* cell surface filaments and the localisation of the motility machinery,
respectively, in strains with *cetZ1* or *cetZ2* deletion, overexpression, and polymerisation-
defective mutant backgrounds. Our results show that CetZ1 has an important role at the
poles of mature motile rod cells for the assembly of key motility proteins, including ArlD1,
a constituent of the archaellum base, and the chemotaxis sensory array adapter CheW1
and signal transducer CheY. Importantly, overproduction of CetZ1 and CetZ1-
mTurquoise2 inhibited motility and reduced the frequency of localisation of the motility
machinery markers but did not affect rod-shape in swimming cells. Our data suggest that
CetZ1 acts as a polar cytoskeletal structure that orchestrates the assembly and
positioning of both major components of the motility machinery. The multifunctionality
and dynamic redeployment of CetZ1 during motile cell development is reminiscent of
eukaryotic cytoskeletal proteins and the roles of tubulin at the base of the eukaryotic
flagellum.

Significance statement

This study demonstrates that archaeal CetZ tubulin-like cytoskeletal proteins control
swimming motility through promoting the assembly of motility structures at cell poles, in
addition to their established roles in rod morphology development. To achieve this, the
conserved CetZ1 protein assembles as a cap or patch-like structures at the cell poles of
mature motile rod cells of the model archaeal organism, *H. volcanii*. CetZ1 migration to
the cell poles for assembly of motility structures has a primitive resemblance to the roles
of tubulin in basal bodies essential to the development of eukaryotic flagella. Our
findings represent the discovery of both multifunctionality and dynamic redeployment in
an archaeal cytoskeletal protein, combined behaviours thought to be a characteristic of
the eukaryotic cytoskeleton.

Introduction

Cytoskeletal proteins are utilised as platforms or scaffolds for large protein complexes across the tree of life. Tubulin superfamily proteins are particularly widespread across all three domains, including tubulin in Eukarya, FtsZ in Bacteria and Archaea, and the Archaea-specific CetZ family[1-5]. In eukaryotes tubulin forms microtubules, which act as platforms in several different biological contexts, including for the anchoring and partitioning of chromosomes during cell division[6], intracellular transport of cargo[7, 8] and assembly of protein complexes[9] and surface appendages[10]. In bacteria, FtsZ forms a ring at mid-cell which is an essential platform required for the assembly of the divisome complex, consisting of more than thirty proteins[11]. In comparison to tubulin and FtsZ, CetZs are understudied in terms of their basic biological functions and their capacity to act as platforms or scaffolds for assembly of other protein complexes.

In the motile and pleomorphic model halophilic archaeon, *Haloferax volcanii*, CetZs have been shown to control cell shape changes and have also been implicated in swimming motility[12, 13]. *H. volcanii* has six CetZ paralogues, with CetZ1 and CetZ2 being the most highly conserved CetZs, and the only CetZs which have been functionally investigated. CetZ1 has a necessary role in rod-development, a morphological change which occurs in several conditions, including trace element depletion[14], in early log phase growth, and when cells are motile[12, 14]. Deletion of *cetZ1* and mutation of its GTPase active site, which is critical for CetZ1 polymerisation, both result in inhibition of rod-development[12, 13]. Importantly, deletion and mutation of *cetZ1* also causes a severe motility defect[12]. While deletion of *cetZ2* has not yet shown to result in any shape or motility phenotype[12], mutation of its GTPase active site prevents rod formation and causes a motility defect in a dominant inhibitory manner[12]. In addition to these observed motility phenotypes in *H. volcanii*, a survey of CetZs, motility, and rod shape across Haloarchaea found that the presence of both CetZ1 and CetZ2 in Haloarchaea species is highly correlated with motility and rod-shape[15]. With our current knowledge, the role of CetZs in motility is seemingly connected to their dominant function in cell shape control. It is not yet clear whether CetZs only contribute to motility through their control of rod-development, or whether they also participate in other mechanisms to promote swimming motility.

Previous studies have shown that CetZ1 localises strongly at poles of rod-shaped cells, either as small bright foci or as larger caps, and appeared especially bright and stable at the poles of motile rods sampled from soft agar[12, 13, 16]. As CetZs are hypothesised to be multifunctional[15], one possibility is that they may contribute to the assembly or positioning of motility protein complexes, such as the archaellum or chemosensory arrays, at cell poles. Like in bacteria, archaeal chemotaxis arrays are responsible for sensing extracellular signals and activating a signal cascade that results in a change in rotational direction of the archaellum[17-19], the whip-like structure equivalent to the

bacterial flagellum (reviewed in [20-24]). On a molecular level, chemotaxis proteins in
Haloarchaea are like the bacterial system, with methyl-accepting chemotaxis proteins
(MCPs), CheW, and CheA organised as a chemosensory array at the cell envelope. When
extracellular signals bind to the MCPs they induce a signal cascade that results in
autophosphorylation of CheA, which in turn phosphorylates CheY[18]. Phosphorylated
CheY shuttles between chemosensory arrays and the archaellum base (composed of
ArlD and ArlCE) where it binds and phosphorylates CheF[17, 19]. Together, ArlD and
ArlCE are responsible for signal transduction to the archaellum motor proteins (ArlH, ArlI,
ArlJ) and highly depend on these motor proteins for localisation at the cell poles[18]. ArlD
and ArlCE are thought to directly bind the archaellum motor[25, 26] but are also
presumed to bind a hypothesized polar cap or cytoplasmic cone structure at the cell
poles[18].

Polar caps are large cytoplasmic structures which are situated adjacent to the inner
membrane in motile euryarchaea. They have been identified in *Pyrococcus furiosus*[26],
Thermococcus kodakarensis[25], and *Halobacterium salinarum*[27]. In all cases, the
polar caps have been found to act as polar organising centres associated with archaella
bundles. In *P. furiosus* and *T. kodakarensis*, polar caps were also associated with
chemosensory arrays. A polar cap has not yet been identified in *H. volcanii*, and the
proteins which compose polar caps in euryarchaea are not yet known.

We hypothesized that CetZs, typical of scaffold forming cytoskeletal proteins, may be
able to act as a platform or polar cap-like structure in *H. volcanii*, and contribute to
motility via mechanisms independent from their control of cell shape. Hence, in this
study we investigated the potential for CetZs to contribute to motility via assembly and
positioning of archaellum and chemotaxis structures. The results are consistent with a
key role for CetZ1 in the assembly and positioning of motility protein complexes at the
poles of motile cells, and our study thus provides the first indication that CetZs have
structural and organisational roles for the assembly of protein complexes at the cell
surface.

Methods

Growth and genetic modification of H. volcanii

H. volcanii was routinely grown in Hv-Cab or Hv-YPCab medium[14] supplemented with 50 μg/mL uracil if necessary to fulfil auxotrophic requirements of strains, or with L-Tryptophan to induce expression of genes of interest from pTA962-based[28] expression plasmids. Cultures were grown at 42 °C with shaking (200 rpm). *H. volcanii* was transformed with demethylated plasmids extracted from *E. coli* strain C2925 using the previously described method[29].

H. volcanii H26 (ID621) was used as the wildtype background and as the parent strain of the Δ*cetZ1* (ID622) and Δ*cetZ2* (ID357) strains, which were generated as in-frame deletions of *cetZ1* (HVO_2204) or *cetZ2* (HVO_0745) using the previously described plasmids pTA131_2204_IF[16] and pTA131_CetZ2[12], respectively and 'pop-in pop-out' method[30]. For all experiments, *H. volcanii* strains contained the pTA962 shuttle vector (control strains) or vectors for expression of relevant genes. Strains are listed in Table S1.

Plasmid construction

All reagents for plasmid construction were obtained from New England Biolabs and used according to the manufacturer's instructions, and oligonucleotides were obtained from Integrated DNA Technologies. Plasmids and oligonucleotides used for cloning detailed in Tables S2 and S3, respectively. To generate a plasmid for the expression of CetZ2-EG-mTurquoise2, the *cetZ2* ORF was amplified from pTA962-CetZ2 using a reverse primer to remove the stop codon and incorporate a BamHI restriction site to the 3' end. The amplified ORF was then ligated between NdeI and BamHI in pHVID21[16].

For plasmids for dual expression of untagged CetZs and GFP tagged motility proteins, ORFs for ArlD1-GFP, GFP-CheW1, and CheY-GFP were amplified from pSVA3919, pSVA5031, and pSVA5611, respectively, incorporating a BglII restriction site at the 5' end, and a NotI site at the 3' end. After purification and restriction digestion, these ORFs were ligated between BamHI and NotI sites of pTA962-CetZ1, pTA962-CetZ2, pTA962-CetZ1.E218A, and pTA962-CetZ2.E212A, utilising the compatible ends generated by BglII and BamHI. The resulting vectors contained the p.*tna* promotor, immediately followed by the CetZ ORF and then the ArlD1, CheW1, or CheY-GFP fusion ORF which was maintained in-frame. The sequences of all constructed plasmids were confirmed by Sanger sequencing (Australian Genome Research Facility).

Soft-agar motility assays

To prepare cultures for soft-agar motility assays, single colonies were inoculated into Hv-Cab medium with 1 mM L-Tryptophan and grown to an approximate OD_{600} of 0.6 overnight.

Following this, cultures were passaged into fresh Hv-Cab medium with 1 mM L-
Tryptophan every 24 hours for three days. All cultures were normalised to the same
starting OD_{600} before spotting 2 µL of culture onto the soft agar.

Hv-Cab soft-agar containing 1 mM L-Tryptophan was prepared the day before inoculation
of cultures onto the agar. For experiments quantifying halo size, 0.25% agar was used,
and motility plates were incubated for 3 days at 42 °C. This concentration of agar resulted
in better uniformity of halo shape and reproducibility of halo size, which was quantified
by taking the mean of two perpendicular measurements of the halo diameter. Motility
assays prepared for subsequent microscopy of halo edge cells used 0.3% agar instead,
and were incubated for 4-5 days at 42 °C. This method sometimes resulted in irregularly
shaped motility halos but produced a more defined leading edge for easier extraction of
cells from the agar for analysis by microscopy.

Light microscopy

Cells were extracted from the edge of motility halos by stabbing the agar with a
micropipette (5-10 µL) near the bottom of the agar plate. After pipetting up and down
several times to liquify the soft-agar and release the cells, the sample was transferred to
a glass slide with a pad containing 1.5% agarose and 18% Buffered Salt Water (BSW)[30]
before covering with a #1.5 glass coverslip. All phase-contrast and fluorescence
microscopy were conducted on the V3 DeltaVision Elite (GE Healthcare) with filter sets
for GFP/FITC (excitation=464-492 nm, emission=500-523 nm) and mTurquoise2/CFP
(excitation=400-453 nm, emission=463-487 nm), an 100X Olympus UPLSAPO/NA 1.4
objective, and pco.edge 5.5 sCMOS camera. Exposure settings were always maintained
between all strains expressing the same fusion protein.

Light microscopy image analysis

To analyse cell shape characteristics and analysis of fluorescence and localisation of
fluorescent fusion proteins within cells, MicrobeJ v5.13I[31] was used, a plugin for FIJI
v2.14.0/1.54f[32]. Unless otherwise mentioned, all settings in MicrobeJ were left as
default. Phase-contrast images were used to detect cell outlines using the default
thresholding algorithm and a cell area restriction of 1.0-max μm^2. Fluorescent
localisation was analysed using the *foci* detection method, restricting foci area to 0.01-
0.75 μm^2. Due to large variance in focus intensity and background fluorescence between
genetic backgrounds and plasmids, the tolerance and Z-score parameters had to be
adjusted for each strain as needed for accurate foci detection, but foci detection
parameters were maintained between biological replicates. Heatmaps of foci
localisation were generated using the *XYCellDensity* function, and all other image
analysis outputs were extracted from the MicrobeJ results tables.

A FIJI macro and python scripts were written and used to conduct local cell curvature
analysis and correlation with fluorescence intensity, based on a previously used
approach[12]. In short, local curvature was analysed using phase-contrast images and
the Contour Curvature v1.1.x (www.gluender.de) plugin in FIJI with the number of contour
samples set to 100, and number of coefficient pairs set to 10. The local curvature at each
contour sample was recorded, and these values were used to calculate the relative
standard deviation of curvature for each cell. The mean fluorescence intensity along the
cell outline generated by Contour Curvature was measured using a 3-pixel line width and
was correlated with local curvature by calculating the Pearson Correlation Coefficient
between fluorescence intensity and local curvature for each individual cell. A summary
of the image acquisition and analysis pipeline for light microscopy is provided in Figure
S3.

Cryo-Electron Microscopy and analysis

To maximize the number of cells per sample for cryo-electron microscopy, cultures were
grown from a theoretical starting OD_{600} of 0.0005 in Hv-YPCab[14] medium and harvested
at OD_{600}~0.25 (after ~24 hours). A 4.5 µL droplet of the culture was applied to glow
discharged copper grids (Quantifoil R2/2, Quantifoil Micro Tools) and blotted for 2.5 s in
a 95% humidity chamber at 20 °C. The grids were then plunged into liquid ethane using a
Lecia EM GP device. The grids were imaged using a Talos Arctica cryo-TEM (Thermo Fisher
Scientific), operating at 200 kV with the specimen held at liquid nitrogen temperature.
Images were captured at 17,500X magnification using a Falcon 3EC direct electron
detector in linear mode. The number of surface filaments per cell (Fig. 1d, e) was
manually counted using Cryo-TEM images. This was conducted independently and
blindly by three individuals.

Results

CetZ1 and CetZ2 influence motility independently of cell shape control

CetZ1 and CetZ2 have previously been implicated in the control of cell motility[12], however it is not yet clear whether they solely contribute to motility through their control of cell shape, or whether they can influence motility independently of cell shape. To begin to investigate this, we first quantitatively measured the impact of *cetZ1* and *cetZ2* deletion and overexpression on motility in soft-agar, where cells swim out from a central inoculation site forming a halo (Fig. 1a). We also assessed the shape of motile cells withdrawn from the leading edge of the halo by image analysis of cell circularity, which is inversely related to cell elongation (Fig. 1b).

Deletion of *cetZ1* caused a severe motility defect, approximately 12% of wildtype motility (Fig. 1a), and prevented rod-formation in cells extracted from the leading edge of motility halos (Fig 1b), as previously reported with a different genetic background [12]. Importantly, overexpression of *cetZ1* in the wildtype or Δ*cetZ1* background reduced motility to approximately 75% of the wildtype control (carrying the vector only), despite cells at the leading edge still forming rods to the same degree as wildtype (Fig. 1a, b).

Deletion of *cetZ2* was previously reported to have no effect on motility or rod shape[12], which was confirmed here in motile cells (Fig. 1a). However, we also discovered that overexpression of c*etZ2*, whether in wildtype or Δ*cetZ2*, resulted in a mild increase in motility, approximately 115% of wildtype (Fig. 1a). This also did not result in a clear change in cell shape (Fig. 1b).

We also investigated the interdependency of the overexpression phenotypes. Overexpression of *cetZ1* in the Δ*cetZ2* background caused reduced motility like it did in the wildtype background, indicating that the effects of *cetZ1* are independent of *cetZ2* (Fig. S1a). Overexpression of *cetZ2* in the Δ*cetZ1* background failed to rescue the Δ*cetZ1* motility defect and did not clearly stimulate motility in this strain. Lastly, overexpression of the GTPase (T7 loop) mutants (*cetZ1*.E218A and *cetZ2*.E212A) inhibited motility and rod shape in the H26 background, as previously reported in H98[12], and had consistent effects in the *cetZ1* and *cetZ2* deletion strains (Fig. S1), suggesting that GTPase activity is necessary for the overexpression phenotypes of *cetZ1* and *cetZ2*.

Importantly, the above experiments indicated that CetZs can promote motility through mechanisms independent of cell shape. This led us to hypothesise that CetZs are involved in assembly or function of other cell envelope or motility structures such as the archaellum or chemosensory arrays.

CetZ1 and CetZ2 influence the frequency of cell surface filaments

To investigate whether CetZs influence the overall appearance of surface filaments such
as pili and archaella, we used cryo-EM (Fig. 1c) to compare the wildtype, *cetZ1* deletion
and *cetZ2* overexpression strains, which showed the most substantial and opposing
effects on motility. Deletion of *cetZ1* resulted in a significant reduction in the number of
surface filaments per cell, whereas overproduction of CetZ2 resulted in slightly more (Fig.
1d), which are thus consistent with the observed motility phenotypes and potential
changes in archaellum assembly.

By comparing surface filament frequency in rods and plate-shaped cells, we found that
in the rod-forming wildtype and *cetZ2* overexpression strains, rod-shaped cells had
significantly more surface filaments compared to plate-shaped cells (Fig. 1e), suggesting
as expected that cell shape correlates with the assembly of surface filaments.
Interestingly, the number of surface filaments in the Δ*cetZ1* strain was still lower than the
number of surface filaments in plate-shaped wildtype cells (Fig. 1d, e). Plates can be
motile as shown by timelapse imaging albeit with reduced speeds[12]. These
observations suggest that the motility defect of Δ*cetZ1* is not only linked to its rod shape
defect, but also a reduction in surface filaments such as the archaellum.

Functional CetZ-mTq2 fusions localise to cell poles in motile cells

CetZ1 was initially reported to localise strongly to cell poles in rod-shaped cells[12],
however since the CetZ1-GFP fusion was not fully functional as a sole copy in rod-
development, it remained unclear whether the polar localisation reflected a native
function of CetZ1. Subsequent studies of CetZ1 localisation using an almost fully
functional CetZ1-mTurquoise2 (mTq2) fusion have not investigated CetZ1 localisation in
motile cells[13, 16]. Therefore, we next used the existing CetZ1-mTq2 fusion and
generated a CetZ2-mTq2 fusion to observe the localisations of CetZ1 and CetZ2 in motile
cells extracted from the leading edge of soft-agar motility halos.

CetZ1-mTq2 and CetZ2-mTq2 were first tested for their capacity to support motility in their
respective *cetZ* knockout backgrounds (Fig. 2). CetZ1- and CetZ2-mTq2 fusions produced
motility halos that were on average 71% and 107% of the size of those produced by
expression of untagged *cetZ1* and *cetZ2*, respectively (Fig. 2a). These differences in
motility were statistically significant between Δ*cetZ1*+CetZ1 and Δ*cetZ1*+CetZ1-mTq2,
but not between Δ*cetZ2*+CetZ2 and Δ*cetZ2*+CetZ2-mTq2. Notably, neither CetZ1-mTq2
nor CetZ2-mTq2 had a substantial effect on the shape of motile rods when expressed in
the wildtype or knockout backgrounds, further supporting the notion that CetZ roles in
motility involve but extend beyond supporting mature rod shape of swimming cells (Fig.
2b).

In motile cells extracted from the halo edge, both proteins displayed bright patches at
cell poles (Fig. 2c-e). CetZ1-mTq2 showed additional patchy localisation around the inner
membrane and at mid-cell as previously described (Fig. 2c)[12, 13, 16]. These similarities

and differences in CetZ1-mTq2 and CetZ2-mTq2 localisations were clearly reflected in
heat map representations of foci localisation along normalized longitudinal and lateral
cell axes (Fig. 2d). No differences in localisation were observed when either CetZ-mTq2
fusion was expressed in the wildtype or their knockout genetic background. The polar
localisations of CetZ1 and CetZ2 are consistent with the hypothesis that they may
participate in the assembly or positioning of motility structures, like the archaellum, at
the cell poles.

CetZ1 influences the positioning and assembly of ArlD1

Given the above results, we proceeded to use fluorescence microscopy to determine
whether CetZ1 and CetZ2 affect the assembly or positioning of key motility proteins.
These included ArlD1 as a marker for the archaellum base, CheW1 as a marker for
chemosensory arrays, and CheY, the effector protein which shuttles between
chemosensory arrays and the archaellum base. These three proteins were considered
good candidates to understand how the CetZs influence assembly and positioning of
motility proteins, as their localisations in motile cells have been well characterised[17,
18] and expression of ArlD1-GFP, GFP-CheW1, and CheY-GFP did not cause substantial
motility or cell shape defects in wildtype (Fig. S2) or *cetZ* mutant backgrounds as
described below. For ArlD1, CheW1, and CheY, the number of foci per cell was used to
assess their degree of assembly, and for rod-forming strains, the percentage of foci
located at cell poles was used as an indicator of correct positioning. As some mutations
of CetZs (*cetZ1* deletion, CetZ1.E218A, and CetZ2.E212A) result in significant inhibition
to rod-formation, we also measured the degree to which curvature of the cell outline
correlated with fluorescent signal of the motility proteins in individual cells (Fig. S3), and
across the population of cells.

In wildtype motile cells, ArlD1-GFP localised as bright foci at the cell poles (Fig. 3a, S4c),
with most cells having two foci (Fig. 3e), one at each pole, as has been previously
observed[17]. The vast majority (>97%) of detected foci were located at cell poles (Fig.
S4b), and across the population, membrane curvature strongly correlated with
fluorescence (Fig. 3f). In the Δ*cetZ1* background, ArlD1-GFP localised as discrete foci
adjacent to the cell edges in some cells (Fig. 3b, S4c), however most cells either had one,
or no foci (Fig. 3e). In addition, there was no correlation between membrane curvature
and ArlD1 localisation (Fig. 3f). When *cetZ1* was overexpressed, the number of ArlD1-GFP
foci (Fig. 3e) and their association with membrane curvature (Fig. 3f) was reduced,
consistent with the reduced swimming motility of this strain (Fig. 1a). Production of
CetZ1.E218A at relatively high levels disrupts cell shape, resulting in a heterogeneous
population of irregularly shaped and enlarged plate-shaped cells (Fig. 3b, S1)[12]. In the
larger cells, ArlD1-GFP localised as many foci distributed throughout the cell and not at
the cell edges (Fig. S4c), while the smaller plate-shaped cells showed low or no localised
fluorescent signal. Image analysis showed that across the population, the number of

ArlD1 foci per cell was increased in comparison to wildtype cells (Fig. 3e), and there was
a negative correlation between the localisation of ArlD1 and membrane curvature (Fig.
3f). Together these data demonstrate that CetZ1 is important for correct positioning and
assembly of ArlD1, but that it is not the only factor important for assembly, as ArlD1 still
formed discrete foci in the absence of CetZ1.

Deletion of *cetZ2* (Fig. 3b) had no effect on the assembly (Fig. 3e) or localisation (Fig. 3f,
S5b, c) of ArlD1 at cell poles, consistent with the lack of motility and cell shape
phenotype exhibited by this strain (Fig. 1a, b)[12]. Interestingly, overproduction of the
CetZ2.E212A GTPase mutant resulted in a more moderate effect than the CetZ1.E218A
GTPase mutant. While rod shape was inhibited, as with production of CetZ2.E212A alone
(Fig. S1b, c), resulting cell shapes were not highly irregular. In addition, ArlD1 localisation
appeared more like that described for the Δ*cetZ1* background. Foci number per cell (Fig.
3e) was reduced compared to the wildtype background, and correlation with membrane
curvature was substantially reduced (Fig. 3f). When CetZ2 was overproduced (Fig. 3b),
ArlD1 positioning at cell poles was not substantially affected (Fig. 3f, S5b, c). However,
CetZ2 overproduction counterintuitively resulted in less ArlD1 foci per cell (Fig. 3e)
despite our observation that this strain had increased motility (Fig. 1a) and has overall
more surface filaments (Fig. 1c). One possible explanation is that the increased number
of surface filaments is not observable by light microscopy due to clustering or
localisation of filaments near one another. Overall, these observations point to a
synthetic role of CetZ2 in motility by promoting of archaellum assembly when it is
overproduced; *cetZ2* deletion showed no impact on ArlD1 positioning or assembly.

CheW1 assembly is highly dependent on CetZ1

Strikingly, we found that CetZ1 was had a major impact on the assembly (localisation) of
the chemotaxis array marker protein GFP-CheW1; deletion of *cetZ1* resulted in diffuse
localisation of GFP-CheW1, with only a small proportion of cells displaying 1-2 faint foci
(Fig. 4b, 4e) in comparison to two or more polar-localised foci in the wildtype background
(Fig. 4a, 4e, S6c). Fluorescence intensity of GFP-CheW1 also was not correlated with
membrane curvature as it is in the wildtype background, likely due to the strong reduction
in localisation of this protein in the absence of CetZ1 (Fig. 4f). Overproduction of CetZ1
reduced the number of foci per cell (Fig. 4e), and their localisation at areas of membrane
curvature (Fig. 4f, S6b, c), consistent with the resulting reduced motility (Fig. 1a). The
CetZ1.E218A GTPase mutant had a similar inhibitory effect on CheW1 localisation as it
did on ArlD1, substantially disrupting cell shape, CheW1 localisation (Fig. 4b, S6c), and
association with membrane curvature (Fig. 4f). Together these observations suggest
CetZ1 is a key driver of CheW1 assembly.

Deletion of *cetZ2* had no effect on foci number (Fig. 4e) or positioning of CheW1 (Fig. 4f).
Differently from the effect of CetZ2.E212A production on ArlD1 assembly, CheW1

localisation was diffuse (Fig. 4b), with the vast majority of cells having no discrete CheW1
foci (Fig. 4e), and GFP-CheW1 fluorescence showing no correlation with membrane
curvature (Fig. 4f, S6c). Additionally, overproduction of CetZ2 caused no difference in
CheW1 foci number (Fig. 4e), conversely to what was observed with ArlD1 (Fig. 3e). CetZ2
overproduction did however cause CheW1 to localise less strongly at the cell poles which
was observable by microscopy (Fig. 4b, S6b, c) and through the reduced correlation of
GFP-CheW1 fluorescence with membrane curvature (Fig. 4f).

CetZ1 is important for CheY assembly and positioning

Following these observations of the impacts of CetZs on the assembly of archaellum and
chemotaxis structures, we next wanted to determine whether CetZs also influence the
effector protein, CheY, which shuttles between chemosensory arrays and the archaellum
base as part of the signal transduction pathway which results in switching of rotation
direction of the archaellum. Generally, we observed that the effect of *cetZ1* deletion,
overexpression, and GTPase mutation on CheY localisation was like that of CheW1. All
three mutations disrupted CheY localisation (Fig. 5b, S7c) by reducing the number of
CheY foci per cell (Fig. 5e), as well as the correlation of CheY positioning with regions of
high membrane curvature (Fig. 5f). Like with ArlD1 and CheW1, production of
CetZ1.E218A also produced a sub-population of large and highly irregularly shaped cells
with several CheY foci (Fig. 5b). This might indicate that CetZ1 is important for CheY
shuttling directly or through its impact on CheW1 assembly and positioning (Fig. 4).

Deletion of *cetZ2* also had no effect on assembly or positioning of CheY, and production
of CetZ2.E212A had similar effects as those on CheW1 assembly and positioning (Fig. 5b,
e, f, S7b, c). CheY-GFP signal was diffuse in most cells, resulting in reduced foci number
and correlation between CheY localisation and membrane curvature. Overproduction of
CetZ2 resulted in less CheY foci per cell, and like CheW1, correlation between CheY
localisation and membrane curvature was reduced. The percentage of detected CheY
foci present at cell poles when CetZ2 was overproduced was also significantly reduced
compared to the wildtype background (Fig. S7b).

Discussion

Several genes in *Haloferax volcanii* have been shown to affect motility, including genes encoding archaella components and its assembly[17, 18, 33], chemotaxis[17, 19], glycosylation[34, 35], lipid modification[36, 37], adhesion[38], as well as transmembrane proteases[34, 39], KaiC-like ATPases[40], transcriptional regulators[41], a multitude of genes with unknown functions[42-45], and cytoskeletal proteins[12, 46, 47]. CetZ tubulin-like proteins in archaea have previously been implicated in the control of cell shape and motility in *H. volcanii*[12], however the molecular mechanisms used by CetZs to control these cellular processes have not yet been described. Rod shape, which is controlled by CetZ1[12, 13], is highly correlated with motility in Haloarchaea[12, 15, 48]. There are a few likely contributing factors to this correlation. Rod-shape itself may provide directionality and a more streamlined cell shape for swimming[49]. Additionally, rod-shaped cells have a polarity which can be a cue for the correct assembly and polar placement of polar motility machinery. For example, the MinD4 positioning protein oscillates between and localises at the cell poles to control the positioning of ArlD1 and CheW1[50]. In plate-shaped cells, which do not have such polarity, MinD4 circles instead of oscillates, and while ArlD1 is still assembled and localised, the archaellum filament and chemosensory arrays are not[17, 50], demonstrating the importance of rod shape and polarity in the assembly mechanism of these polar motility complexes.

Here, we identified that although some manipulations of CetZs (Δ*cetZ1,* +CetZ1.E218A, +CetZ2.E212A) exhibit both rod-shape and motility defects (also demonstrated in [12] and [13]), overproduction of either CetZ1 or CetZ2 result in motility phenotypes without affecting cell shape (Fig. 1a, b, S1). This shows that in addition to their role in controlling cell shape, CetZs are likely to also contribute to motility through separate mechanisms. Following these observations, we focused on investigating the role of CetZ1 and CetZ2 in assembly and positioning of motility proteins ArlD1, CheW1, and CheY, testing the hypothesis that they may contribute to motility by acting as a scaffold or platform for the assembly of archaella and chemosensory array protein complexes.

Although overproduction of CetZ2 resulted in an increase in motility and the number of surface filaments, it did not appear to promote the assembly or positioning of ArlD1, CheW1, or CheY at the resolutions achievable by fluorescence microscopy. Instead, overproduction of CetZ2 resulted in slightly less efficient localisation of these proteins at cell poles, which may be a result of tightly clustered proteins within proximity to one another. From these data alone, the mechanism behind hypermotility resultant from CetZ2 overproduction is still unclear.

As deletion of *cetZ2* had no impact on cell shape, motility (as previously observed in [12]), or assembly and positioning of ArlD1, CheW1, or CheY, we believe the motility phenotypes resulting from overproduction of CetZ2 and CetZ2.E212A are likely to be synthetic consequences of overproduction, and CetZ2 most likely does not have a

significant role in motility, at least under the conditions investigated where CetZ2 protein levels are expected to be low[51]. Therefore, we believe it is likely that the level of CetZ2 during mid-log phase or in motile rods is too low to result in any cell shape or motility phenotype when deleted. Here, and in previous work[12], the CetZ2 GTPase mutant has been demonstrated to function within cell shape and motility pathways, so perhaps CetZ2 has an important role in the control of cell shape or motility in other conditions instead.

The different effects of deletion, overproduction, or GTPase mutations of CetZ1 and CetZ2, in addition to their strong conservation throughout haloarchaea and their distinct phylogenetic groupings, demonstrates that CetZ1 and CetZ2 are likely to have distinct functions. Here we observed that CetZ1 overproduction in the absence of CetZ2 still reduced motility equivalently to the wildtype background. However, CetZ2 overproduction did not complement the motility defect caused by deletion of *cetZ1*, indicating CetZ1 and CetZ2 are not redundant proteins. As previously reported[12], CetZ2.E212A dominantly blocks rod shape even though deletion of *cetZ2* has no effect on the shape of motile cells, suggesting it inhibits CetZ1-mediated maturation of motile rods[12]. Further work must be carried out to understand the endogenous function of CetZ2.

CetZ1 demonstrated obvious effects on both the positioning and assembly of ArlD1, CheW1, and CheY, but had a particularly strong influence on the latter two chemotaxis proteins. Deletion or overexpression of *cetZ1*, as well as production of the CetZ1.E218A dominant-inhibitory GTPase mutant all reduced assembly and polar positioning of ArlD1, CheW1, and CheY to varying degrees, consistent with their observed motility defects. The reduction in swimming motility and localisation of motility markers upon overproduction of CetZ1 suggests that the excess CetZ interferes with the proper assembly of the motility proteins[52].

We also confirmed, using functional CetZ1-mTq2 fusions, that CetZ strongly localises to the poles of motile rods, which differs compared to the longitudinal filaments and dynamic foci seen in rods and during rod formation in liquid culture[12, 13]. Notably, a CetZ1-mTq2 fusion fully supported rod shape in swimming cells (ΔcetZ1 background), but motility was reduced compared to the untagged CetZ1. This further supports the notion that these two functions of CetZ1 occur independently. Together, these observations demonstrate that while CetZ1 is required for mature rod-shape of motile cells, it also has a mechanistic role in the positioning and assembly of archaella and chemotaxis arrays. Our findings are thus consistent with the general hypothesis that CetZ1 is an upstream player in the assembly of polar protein complexes in addition to aiding motility through its control of rod-shape.

Polar cap structures have been identified in several archaeal species, including *Thermococcus kodakarensis*[25], *Pyrococcus furiosus*[26], and a relative of *H. volcanii*,

Halobacterium salinarum[53]. Although hypothesized[18], a polar cap has not been
identified in *H. volcanii*, and it is not known what components constitute the cap
structures in other archaea. Together with the observation that CetZ1 strongly localises
at the cell poles of motile cells, our data suggests a new role of CetZ1 as a functional
component of a polar cap in *H. volcanii*. CetZ1 forms protofilaments which assemble
laterally into sheets and are predicted to interact with the inner membrane[12, 13]. We
predict that such sheet-like structures might assemble and help form the cell poles and
promote assembly of motility structures there.

Both CetZ1 and CetZ2 have even been identified as potential interacting partners with
several chemotaxis proteins (including CheW1 and CheY) by affinity purification mass
spectrometry in *H. salinarum*[54], which is known to have a polar cap[53], further
indicating the potential for CetZs to interact with protein complexes at cell poles in *H.*
volcanii. These previous findings are consistent with the strong effect of CetZ1 on
chemotaxis protein positioning and assembly we observed in *H. volcanii*. High resolution
structural and compositional analysis of cap structures in *H. volcanii* will be valuable
towards determining the exact structural role of CetZ1 at cell poles. With CetZ1 and
MinD4[50] both implicated in the positioning of archaella and chemosensory arrays at
cell poles, future research could also investigate how these two proteins coordinate their
activities, and where they lie within the hierarchy of proteins contributing to motility.

Our findings are the first to show that a CetZ family protein participates in multiple
distinct roles. Taken together with the diversity and multiplicity of tubulin superfamily
proteins in archaea[1, 15], it is likely that similar characteristics in early archaea led to
the vast functional complexity, diversity, and specialisation of tubulins in eukaryotes. Our
findings are also reminiscent of the fundamental roles of tubulin at the base of the
eukaryotic flagellum, which originated before the last eukaryotic common ancestor[55-
58]. Since flagellar basal body microtubules are derived from mitotic centrioles during
eukaryotic cell differentiation, it is interesting to consider parallels to CetZ1 longitudinal
filaments during rod formation[12, 13], followed by its polar localisation to promote
stable assembly of the motility machinery in mature rod cells.

Acknowledgements

The authors would like to acknowledge S. Albers for providing strains and plasmids, as well as L. Cole and A. Bottomley from the UTS Microbial Imaging Facility. We acknowledge the use of the Cryo-Electron Microscopy Facility through the Victor Chang Cardiac Research Institute Innovation Centre, funded by the NSW government and the Electron Microscope Unit within the Mark Wainwright Analytical Centre (MWAC) at UNSW Sydney. The work was supported by the Australian Research Council (DP160101076 and FT160100010 to IGD).

Author contributions

Conceptualisation – HJB & IGD; Data curation – HJB, MII & JR; Formal analysis – HJB, MII & JR; Investigation – HJB, MII, JR; Methodology – all authors; Supervision – IGD, SI & MABB; Original draft – HJB; Revision – HJB & IGD; Reviewing the manuscript – all authors. Funding Acquisition – IGD.

References

1. Aylett CH, Duggin IG: **The tubulin superfamily in archaea**. In: *Prokaryotic Cytoskeletons.* Springer; 2017: 393-417.
2. Nogales E, Downing KH, Amos LA, Löwe J: **Tubulin and FtsZ form a distinct family of GTPases**. *Nature Structural and Molecular Biology* 1998, **5**(6):451.
3. Vaughan S, Wickstead B, Gull K, Addinall SG: **Molecular evolution of FtsZ protein sequences encoded within the genomes of archaea, bacteria, and eukaryota**. *Journal of molecular evolution* 2004, **58**(1):19-29.
4. Erickson HP: **Evolution of the cytoskeleton**. *Bioessays* 2007, **29**(7):668-677.
5. Erickson HP: **FtsZ, a prokaryotic homolog of tubulin?** *Cell* 1995, **80**(3):367-370.
6. Barton NR, Goldstein L: **Going mobile: microtubule motors and chromosome segregation**. *Proceedings of the National Academy of Sciences* 1996, **93**(5):1735-1742.
7. Appert-Rolland C, Ebbinghaus M, Santen L: **Intracellular transport driven by cytoskeletal motors: General mechanisms and defects**. *Physics Reports* 2015, **593**:1-59.
8. Hirokawa N, Noda Y, Tanaka Y, Niwa S: **Kinesin superfamily motor proteins and intracellular transport**. *Nature reviews Molecular cell biology* 2009, **10**(10):682-696.
9. An S, Deng Y, Tomsho JW, Kyoung M, Benkovic SJ: **Microtubule-assisted mechanism for functional metabolic macromolecular complex formation**. *Proceedings of the National Academy of Sciences* 2010, **107**(29):12872-12876.
10. Hawkins T, Mirigian M, Yasar MS, Ross JL: **Mechanics of microtubules**. *Journal of biomechanics* 2010, **43**(1):23-30.
11. Lutkenhaus J, Pichoff S, Du S: **Bacterial cytokinesis: from Z ring to divisome**. *Cytoskeleton* 2012, **69**(10):778-790.
12. Duggin IG, Aylett CH, Walsh JC, Michie KA, Wang Q, Turnbull L, Dawson EM, Harry EJ, Whitchurch CB, Amos LA: **CetZ tubulin-like proteins control archaeal cell shape**. *Nature* 2015, **519**(7543):362.
13. de Silva RT, Shinde V, Brown HJ, Liao Y, Duggin IG: **Dynamic self-association of archaeal tubulin-like protein CetZ1 drives Haloferax volcanii morphogenesis**. *bioRxiv* 2024:2024.2004. 2008.588506.
14. de Silva RT, Abdul-Halim MF, Pittrich DA, Brown HJ, Pohlschroder M, Duggin IG: **Improved growth and morphological plasticity of Haloferax volcanii**. *Microbiology* 2021.
15. Brown HJ, Duggin IG: **Diversity and Potential Multifunctionality of Archaeal CetZ Tubulin-like Cytoskeletal Proteins**. *Biomolecules* 2023, **13**(1):134.
16. Ithurbide S, de Silva RT, Brown HJ, Shinde V, Duggin IG: **A vector system for single and tandem expression of cloned genes and multi-colour fluorescent tagging in Haloferax volcanii**. *bioRxiv* 2024:2024.2003. 2013.584740.
17. Li Z, Kinosita Y, Rodriguez-Franco M, Nußbaum P, Braun F, Delpech F, Quax TE, Albers S-V: **Positioning of the motility machinery in halophilic archaea**. *MBio* 2019, **10**(3):e00377-00319.
18. Li Z, Rodriguez‐Franco M, Albers SV, Quax TE: **The switch complex ArlCDE connects the chemotaxis system and the archaellum**. *Molecular microbiology* 2020, **114**(3):468-479.
19. Quax TE, Altegoer F, Rossi F, Li Z, Rodriguez-Franco M, Kraus F, Bange G, Albers S-V: **Structure and function of the archaeal response regulator CheY**. *Proceedings of the National Academy of Sciences* 2018, **115**(6):E1259-E1268.
20. Albers S-V, Jarrell KF: **The archaellum: how Archaea swim**. *Frontiers in Microbiology* 2015, **6**:23.
21. Jarrell KF, Albers S-V: **The archaellum: an old motility structure with a new name**. *Trends in microbiology* 2012, **20**(7):307-312.
22. Jarrell KF, Albers S-V, Machado J: **A comprehensive history of motility and Archaellation in Archaea**. *FEMS Microbes* 2021, **2**.
23. Albers S-V, Jarrell KF: **The archaellum: an update on the unique archaeal motility structure**. *Trends in microbiology* 2018, **26**(4):351-362.
24. Beeby M, Ferreira JL, Tripp P, Albers S-V, Mitchell DR: **Propulsive nanomachines: the convergent evolution of archaella, flagella and cilia**. *FEMS microbiology reviews* 2020, **44**(3):253-304.
25. Briegel A, Oikonomou CM, Chang YW, Kjær A, Huang AN, Kim KW, Ghosal D, Nguyen HH, Kenny D, Ogorzalek Loo RR: **Morphology of the archaellar motor and associated cytoplasmic cone in Thermococcus kodakaraensis**. *EMBO reports* 2017, **18**(9):1660-1670.

26. Daum B, Vonck J, Bellack A, Chaudhury P, Reichelt R, Albers S-V, Rachel R, Kühlbrandt W:
Structure and in situ organisation of the Pyrococcus furiosus archaellum machinery.
Elife 2017, **6**:e27470.

27. Kupper J, Marwan W, Typke D, Grünberg H, Uwer U, Gluch M, Oesterhelt D: **The flagellar**
bundle of Halobacterium salinarium is inserted into a distinct polar cap structure.
Journal of bacteriology 1994, **176**(16):5184-5187.

28. Allers T, Barak S, Liddell S, Wardell K, Mevarech M: **Improved strains and plasmid vectors**
for conditional overexpression of His-tagged proteins in Haloferax volcanii. *Applied and*
environmental microbiology 2010, **76**(6):1759-1769.

29. Cline SW, Lam WL, Charlebois RL, Schalkwyk LC, Doolittle WF: **Transformation methods**
for halophilic archaebacteria. *Canadian Journal of Microbiology* 1989, **35**(1):148-152.

30. Allers T, Ngo H-P, Mevarech M, Lloyd RG: **Development of additional selectable markers**
for the halophilic archaeon Haloferax volcanii based on the leuB and trpA genes.
Applied and environmental microbiology 2004, **70**(2):943-953.

31. Ducret A, Quardokus EM, Brun YV: **MicrobeJ, a tool for high throughput bacterial cell**
detection and quantitative analysis. *Nature microbiology* 2016, **1**(7):1-7.

32. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B: **Fiji: an open-source platform for biological-image**
analysis. *Nature methods* 2012, **9**(7):676.

33. Tripepi M, Esquivel RN, Wirth R, Pohlschröder M: **Haloferax volcanii cells lacking the**
flagellin FlgA2 are hypermotile. *Microbiology* 2013, **159**(Pt 11):2249.

34. Tripepi M, Imam S, Pohlschröder M: **Haloferax volcanii flagella are required for motility**
but are not involved in PibD-dependent surface adhesion. *Journal of bacteriology* 2010,
192(12):3093-3102.

35. Tripepi M, You J, Temel S, Önder Ö, Brisson D, Pohlschröder M: **N-glycosylation of**
Haloferax volcanii flagellins requires known Agl proteins and is essential for
biosynthesis of stable flagella. *Journal of bacteriology* 2012, **194**(18):4876-4887.

36. Abdul Halim MF, Pfeiffer F, Zou J, Frisch A, Haft D, Wu S, Tolić N, Brewer H, Payne SH, Paš
a‐Tolić L: **H aloferax volcanii archaeosortase is required for motility, mating, and C‐**
terminal processing of the S‐layer glycoprotein. *Molecular microbiology* 2013, **88**(6):1164-
1175.

37. Abdul Halim MF, Schulze S, DiLucido A, Pfeiffer F, Bisson Filho AW, Pohlschroder M: **Lipid**
Anchoring of Archaeosortase Substrates and Midcell Growth in Haloarchaea. *mBio*
2020, **11**(2):e00349-00320.

38. Esquivel RN, Pohlschroder M: **A conserved type IV pilin signal peptide H‐domain is**
critical for the post‐translational regulation of flagella‐dependent motility. *Molecular*
microbiology 2014, **93**(3):494-504.

39. Cerletti M, Martínez MJ, Giménez MI, Sastre DE, Paggi RA, De Castro RE: **The LonB**
protease controls membrane lipids composition and is essential for viability in the
extremophilic haloarchaeon H aloferax volcanii. *Environmental microbiology* 2014,
16(6):1779-1792.

40. Chatterjee P, Garcia M, Cote J, Yun K, Legerme G, Habib R, Tripepi M, Young C, Kulp DW,
Dyall-Smith M: **Involvement of ArlI, ArlJ, and CirA in Archaeal Type-IV Pilin-Mediated**
Motility Regulation. *bioRxiv* 2024:2024.2003. 2004.583388.

41. Hackley RK, Hwang S, Herb JT, Bhanap P, Lam K, Vreugdenhil A, Darnell CL, Pastor MM,
Martin JH, Maupin‐Furlow JA: **TbsP and TrmB jointly regulate gapII to influence cell**
development phenotypes in the archaeon Haloferax volcanii. *Molecular Microbiology*
2024.

42. Parente J, Casabuono A, Ferrari MC, Paggi RA, De Castro RE, Couto AS, Giménez MI: **A**
rhomboid protease gene deletion affects a novel oligosaccharide N-linked to the S-
layer glycoprotein of Haloferax volcanii. *Journal of Biological Chemistry* 2014,
289(16):11304-11317.

43. Collins M, Afolayan S, Igiraneza AB, Schiller H, Krespan E, Beiting DP, Dyall-Smith M, Pfeiffer
F, Pohlschroder M: **Mutations affecting HVO_1357 or HVO_2248 cause hypermotility in**
Haloferax volcanii, suggesting roles in motility regulation. *Genes* 2021, **12**(1):58.

44. Nagel C, Machulla A, Zahn S, Soppa J: **Several one-domain zinc finger μ-proteins of**
Haloferax volcanii are important for stress adaptation, biofilm formation, and
swarming. *Genes* 2019, **10**(5):361.

45.   Legerme G, Yang E, Esquivel RN, Kiljunen S, Savilahti H, Pohlschroder M: **Screening of a**
633   **Haloferax volcanii transposon library reveals novel motility and adhesion mutants**. *Life*
634   2016, **6**(4):41.

635   46.   Schiller H, Hong Y, Kouassi J, Rados T, Kwak J, DiLucido A, Safer D, Marchfelder A, Pfeiffer
636   F, Bisson A: **Identification of structural and regulatory cell-shape determinants in**
637   **Haloferax volcanii**. *Nature communications* 2024, **15**(1):1414.

638   47.   Schiller H, Kouassi J, Hong Y, Rados T, Kwak J, DiLucido A, Safer D, Marchfelder A, Pfeiffer
639   F, Bisson-Filho A: **Identification and characterization of structural and regulatory cell-**
640   **shape determinants in Haloferax volcanii**. *bioRxiv* 2023:2023.2003. 2005.531186.

641   48.   Schwarzer S, Rodriguez-Franco M, Oksanen HM, Quax TE: **Growth phase dependent cell**
642   **shape of Haloarcula**. *Microorganisms* 2021, **9**(2):231.

643   49.   Young KD: **The selective value of bacterial shape**. *Microbiology and molecular biology*
644   *reviews* 2006, **70**(3):660-703.

645   50.   Nußbaum P, Ithurbide S, Walsh JC, Patro M, Delpech F, Rodriguez-Franco M, Curmi PM,
646   Duggin IG, Quax TE, Albers S-V: **An oscillating MinD protein determines the cellular**
647   **positioning of the motility machinery in archaea**. *Current Biology* 2020, **30**(24):4956-4972.
648   e4954.

649   51.   Cerletti M, Giménez MI, Tröetschel C, D'Alessandro C, Poetsch A, De Castro RE, Paggi RA:
650   **Proteomic Study of the Exponential–Stationary Growth Phase Transition in the**
651   **Haloarchaea Natrialba magadii and Haloferax volcanii**. *Proteomics* 2018, **18**(14):1800116.

652   52.   Prelich G: **Gene overexpression: uses, mechanisms, and interpretation**. *Genetics* 2012,
653   **190**(3):841-854.

654   53.   Mormile MR, Biesen MA, Gutierrez MC, Ventosa A, Pavlovich JB, Onstott TC, Fredrickson JK:
655   **Isolation of Halobacterium salinarum retrieved directly from halite brine inclusions**.
656   *Environmental Microbiology* 2003, **5**(11):1094-1102.

657   54.   Schlesner M, Miller A, Besir H, Aivaliotis M, Streif J, Scheffer B, Siedler F, Oesterhelt D: **The**
658   **protein interaction network of a taxis signal transduction system in a halophilic**
659   **archaeon**. *BMC microbiology* 2012, **12**:1-20.

660   55.   Carvalho-Santos Z, Azimzadeh J, Pereira-Leal JB, Bettencourt-Dias M: **Evolution: Tracing**
661   **the origins of centrioles, cilia, and flagella**. *J Cell Biol* 2011, **194**(2):165-175.

662   56.   Mitchell DR: **The evolution of eukaryotic cilia and flagella as motile and sensory**
663   **organelles**. *Adv Exp Med Biol* 2007, **607**:130-140.

664   57.   Moran J, McKean PG, Ginger ML: **Eukaryotic Flagella: Variations in Form, Function, and**
665   **Composition during Evolution**. *BioScience* 2014, **64**(12):1103-1114.

666   58.   Loreng TD, Smith EF: **The Central Apparatus of Cilia and Eukaryotic Flagella**. *Cold Spring*
667   *Harb Perspect Biol* 2017, **9**(2).

Figure 1. CetZs can contribute to motility without influencing shape. H26 wildtype, Δ*cetZ1* and Δ*cetZ2* strains with pTA962 vector only, and wildtype with *cetZ1* or *cetZ2* expressed from pTA962 as indicated (+CetZ1/2) were inoculated onto Hv-Cab soft-agar with 1 mM L-Tryptophan. **A)** Their motility halo diameters were quantified after four days and normalized to the mean of WT *H. volcanii* H26. Individual points represent biological replicates. **B)** . Circularity of individual cells extracted from the edge of motility halos was measured and is represented as a frequency distribution for wildtype and knockout *cetZ* strains with pTA962. CetZ1 and CetZ2 (+CetZ1/2) were expressed in their respective knockout backgrounds. Data for WT +pTA962 is displayed on both graphs for reference. The number of individual cells analysed for each strain is noted in Fig. S1. **C)** Representative CryoEM image of H26 wildtype cells (left) and zoom (right) showing surface filaments (red tick marks). **D)** The number of surface filaments per cell were manually counted using images obtained in **C** for H26 wildtype, Δ*cetZ1*, and +CetZ2 strains. The number of cells analysed per strain is as follows: WT +pTA962, n=46; Δ*cetZ1* +pTA962, n=58; WT +CetZ2, n=66. **E)** For rod forming strains, WT and +CetZ2, the number of surface filaments per cell was determined for rod and plate-shaped cells. Data in **D** and **E** was obtained from the same samples. **A, D,** and **E** show mean and standard deviation. **A** and **D** use one-way ANOVA as a statistical test, and **E** uses an un-paired T-test. ****p<0.00005, **p<0.005, *p<0.05.

Figure 2. Localisation of CetZ1 and CetZ2 in motile cells. CetZ1-mTq2 and CetZ2-mTq2 fusion proteins were expressed in H26 wildtype cells and their respective knockout backgrounds. H26 wildtype, Δ*cetZ1*, and Δ*cetZ2* with pTA962, and Δ*cetZ1* +CetZ1, and Δ*cetZ2* + CetZ2 were used as controls. **A)** Strains were inoculated onto Hv-Cab soft-agar with 1 mM L-Tryptophan and their motility halo diameters were quantified. Individual points represent biological replicates. **(B)** Cells expressing CetZ1 or CetZ2-mTq2 fusions were extracted from the edge of the motility halo and their circularity was measured using phase-contrast microscopy images (not shown). **C)** Fluorescence microscopy was also conducted to observe CetZ1 and CetZ2 localisation in these cells. **D)** Heatmap representations of CetZ1 and CetZ2 localisation in motile cells. **E)** Proportion of detected CetZ foci located at cell poles. In **A** and **E**, mean and standard deviation are shown. One-way ANOVA was used as a statistical test, and only significant and/or relevant comparisons are shown on graphs. ****$p<0.00005$, ***$p<0.0005$, ns=not significant. The number of cells analysed for each strain in **B-E** was as follows: WT +pTA962, n=2046; WT +CetZ1-mTq2, n=1193; Δ*cetZ1* +CetZ1-mTq2, n=1672; WT +CetZ2-mTq2, n=1338; Δ*cetZ2* +CetZ2-mTq2, n=861.

Figure 3. ArlD1 localisation in CetZ mutant strains. ArlD1-GFP was expressed from pSVA3919 in H26
wildtype, Δ*cetZ1*, and Δ*cetZ2*, or was co-expressed with CetZ1, CetZ1.E218A, CetZ2, or CetZ2.E212A from
pHJB31, pHJB34, pHJB37, or pHJB40, respectively, in H26 wildtype. These strains were inoculated onto Hv-
Cab soft-agar with 1 mM L-Tryptophan and extracted from the halo-edge for **A-B)** phase-contrast and
fluorescence microscopy. Four biological replicates were carried out for each strain. Scale bar: 5 µm. **C)**
Motility halo diameters for each strain were quantified. Individual points represent biological replicates. **D)**
Phase-contrast images were used to measure circularity. Individual points represent the mean of all cell
circularities in a biological replicate. In **C** and **D** mean and standard deviation are shown. One-way ANOVA
was used as a statistical test. ****$p<0.00005$, ***$p<0.0005$, **$p<0.005$, *$p<0.05$. Only significant
comparisons are shown. **E)** Frequency distributions of the number of foci per cell in each strain, and **F)** the
Pearson Correlation Coefficient between local membrane curvature and fluorescence intensity. In **E** and **F**,
data for WT +ArlD1-GFP is displayed on all graphs for reference, and the key refers to both panels. For **D-F**
the number of cells pooled from four biological replicates for each strain is as follows: WT +ArlD1-GFP,
n=2240; Δ*cetZ1* +ArlD1-GFP, n=1097; WT +CetZ1 +ArlD1-GFP, n=1448; WT +CetZ1.E218A +ArlD1-GFP,
n=2007; Δ*cetZ2* +ArlD1-GFP, n=1249; WT +CetZ2 +ArlD1-GFP, n=1655; WT +CetZ2.E212A +ArlD1-GFP,
n=1466.

Figure 4. CheW1 localisation in CetZ mutant strains. GFP-CheW1 was expressed from pSVA5031 in H26 wildtype, Δ*cetZ1*, and Δ*cetZ2*, or was co-expressed with CetZ1, CetZ1.E218A, CetZ2, or CetZ2.E212A from pHJB32, pHJB35, pHJB38, or pHJB41, respectively, in H26 wildtype. These strains were inoculated onto Hv-Cab soft-agar with 1 mM L-Tryptophan and extracted from the halo-edge. **A-F** is as described in Figure 3. For **D-F** the number of cells pooled from four biological replicates for each strain is as follows: WT +GFP-CheW1, n=1792; Δ*cetZ1* +GFP-CheW1, n=2458; WT +CetZ1 +GFP-CheW1, n=2195; WT +CetZ1.E218A +GFP-CheW1, n=3834; Δ*cetZ2* +GFP-CheW1, n=3222; WT +CetZ2 +GFP-CheW1, n=2121; WT +CetZ2.E212A +GFP-CheW1, n=4055.

Figure 5. CheY localisation in CetZ mutant strains. CheY-GFP was expressed from pSVA5611 in H26
wildtype, Δ*cetZ1*, and Δ*cetZ2*, or was co-expressed with CetZ1, CetZ1.E218A, CetZ2, or CetZ2.E212A from
pHJB33, pHJB36, pHJB39, or pHJB42, respectively, in H26 wildtype. These strains were inoculated onto Hv-
Cab soft-agar with 1 mM L-Tryptophan and extracted from the halo-edge. **A-F** is as described in Figure 3.
For **D-F** the number of cells pooled from four biological replicates for each strain is as follows: WT +CheY-
GFP, n=2892; Δ*cetZ1* +CheY-GFP, n=1825; WT +CetZ1 +CheY-GFP, n=1490; WT +CetZ1.E218A +CheY-GFP,
n=1704; Δ*cetZ2* +CheY-GFP, n=1341; WT +CetZ2 +CheY-GFP, n=1833; WT +CetZ2.E212A +CheY-GFP,
n=2422.

739