What does your gut tell you? Influences of a zombie-making and generalist fungal entomopathogen on carpenter ant micro- and mycobiota

Sophia Vermeulen[a], Anna M Forsman[a,b], Charissa de Bekker*[a,c]

[a] Department of Biology, University of Central Florida, Orlando FL 32816, USA
[b] Department of Biology, Colby University, Waterville ME 04901, USA
[c] Microbiology, Department of Biology, Utrecht University, 3584 CH Utrecht, the Netherlands
*Corresponding author email: a.m.debekker@uu.nl

Highlights

- *Ophiocordyceps* and *Beauveria* infections impact the gut microbiota of carpenter ants differently
- The fungal taxa present in the ant gut are more diverse and dynamic across infection conditions compared to the bacterial taxa
- There is significant dysbiosis in the bacterial and fungal microbiota of ants that display *Ophiocordyceps*-manipulated summiting behavior
- Dysbiosis in *Ophiocordyceps*-infected ants might be contributing to observed host behavioral changes while a comparable dysbiosis is not observed in *Beauveria* infections

Abstract

Microbiome composition impacts many host aspects including health, nutrition, reproduction, and behavior. This warrants the recent uptick in insect microbiota research across species and ecosystems. Commensurate with this, the bacterial microbiome of the ant *Camponotus floridanus* has been well characterized across body regions and maturation levels. However, potential effects of entomopathogens on the gut microbiome, and the fungal communities therein, are yet to be assessed. Investigation of the microbiome during infection could provide insight into entomopathogenic infection and manipulation strategies and inform effective biopesticide strategies. Additionally, the mycobiome remains often overlooked despite playing a vital role in gut ecology with potential implications for health and infection outcomes. To improve our limited understanding of fungal infections in insects, and ants in particular, we characterized the effects of two entomopathogens with different infection strategies on the gut micro- and mycobiota of *C. floridanus* over time; *Ophiocordyceps camponoti-floridani* and *Beauveria bassiana.* Specialist, 'zombie-making' *O. camponoti-floridani* fungi hijack the behavior of *C. floridanus* ants over three weeks, causing them to find an elevated position, and fix themselves in place with their mandibles. This summiting behavior is adaptive to *Ophiocordyceps* as the ant transports the fungus to conditions that favor fruiting body development, spore production, dispersal, and transmission. In contrast, the generalist entomopathogen *B. bassiana* infects and kills the ant within a few days, without the induction of obvious fungus-adaptive behaviors. By comparing healthy ants with

Beauveria- and *Ophiocordyceps*-infected ants we aimed to 1) describe the dynamics of the micro- and mycobiome of *C. floridanus* during infection, and 2) determine if the effects on gut microbiota are distinctive between fungi that have different infection strategies. While *Beauveria* did not measurably affect the ant host micro-and mycobiome, *Ophiocordyceps* did, especially for the mycobiome. Moreover, ants that were sampled during *Ophiocordyceps*-adaptive summiting behavior had a significantly different micro- and mycobiome composition compared to healthy controls and those sampled before and after manipulation took place. This suggests that the host microbiome might have a role to play in the manipulation strategy of *Ophiocordyceps*.

Keywords
Ophiocordyceps, *Beauveria*, infection, microbiome, behavior manipulation

1. Introduction

Parasitism has evolved multiple times across the fungal Kingdom, giving rise to numerous species that established antagonistic interactions with plants, animals, and other fungi. The order Hypocreales contains a multitude of parasitic species including many insect-infecting fungi (i.e., fungal entomopathogens) (1). The genera *Beauveria* (family: Cordycipitaceae) and *Ophiocordyceps* (family: Ophiocordycipitaceae) represent well-studied examples with different pathogenic strategies. *Beauveria bassiana* is a generalist that can infect insects across orders. As such, strains of this species are isolated and studied for their potential use in the biocontrol of insect pests (2, 3). To initiate infection, *Beauveria* spores attach and penetrate through the insect host's cuticle. Once inside, the fungus proliferates as blastospores (i.e., single-celled, yeast-like cells), releases toxins, destroys host tissues, and kills the insect within a few days (3). This destructive, toxin-based infection strategy, paired with its wide host range classify *Beauveria* as a necrotroph-like fungal entomopathogen.

In contrast to *Beauveria*, *Ophiocordyceps* species are highly specialized entomopathogens that generally infect only one insect species (4-6). The 'zombie- making' fungi of the species complex *Ophiocordyceps unilateralis* sensu lato infect ants from the Camponotini tribe. These fungi slowly hijack ant behavior over the course of three to four weeks before eventually killing their host (7, 8). While the fungus is growing as single blastospores, infected individuals wander away from the nest and climb to elevated positions that provide favorable conditions for fungal growth (9, 10). The manipulated ant then proceeds to bite onto the vegetation, firmly attaching itself in the final moments before death. Following this summiting behavior, the fungus converts to a multicellular, mycelial growth, which eventually gives rise to a fruiting body that sprouts from the cadaver (11, 12). This fruiting body carries the spores that are released to infect new ants (4-6). These manipulated behaviors are a crucial aspect of the *Ophiocordyceps* life cycle that aid in circumventing social immunity behaviors of healthy nest mates (13) and place the ant in a microclimate that promotes fungal growth and spore dispersal (10, 14). Without this behavior the fruiting body cannot form (11, 15), and transmission fails. Therefore, *Ophiocordyceps* species within the *unilateralis* complex must treat their host in a much more delicate manner than *B. bassiana* to ensure that the ant does not succumb to the infection prematurely. As such, *Ophiocordyceps* is thought to begin its infection much like a biotrophic fungal plant pathogen, gaining resources from living cells, and only inducing subtle changes without causing severe tissue damage. However, once manipulated biting occurs, the fungus necessarily becomes more necrotrophic as it rapidly needs to consume insect tissue to gain enough energy for fruiting body development. As such, in contrast to *B. bassiana*, ant-infecting *Ophiocordyceps* fungi use parasitic strategies comparable to those of a hemi-biotroph (16).

Unraveling how *Ophiocordyceps* and other insect-infecting fungi interact with their insect hosts could prove valuable for the discovery of novel, more sustainable pesticides, and pesticide targets. Moreover, learning how pathogenic behavioral phenotypes are

induced leads to a better mechanistic understanding of insect behavior in general. As such, recent laboratory studies on *Ophiocordyceps*-ant interactions have used a combination of quantitative behavioral assays and various omics technologies to investigate how *Ophiocordyceps* affects ant circadian activity, communication, olfaction, immunity, and biogenic amine levels such as dopamine (8, 13, 17-19). The fungus appears to achieve this by secreting small bioactive molecules of which some are predicted to bind to G-protein coupled receptors that are involved in light and odor sensing or binding of biogenic amines (20). While these mechanistic studies are vital to understand how *Ophiocordyceps* manipulates its host through chemical means, they only focus on fungal interactions with the nervous and muscular tissue in the ant head. This approach leaves the interactions between the fungus and other host tissues, such as the gut and its microbiota, completely in the dark. Meanwhile, microbiome studies produce mounting evidence for the importance of (gut) microbiota in general host functioning, including nutrition, immunity, and behavior (21-25). As such, interactions between infectious agents, such as fungal entomopathogens and other parasites, and the gut microbiota of their host could be a vital part of disease progression and resulting phenotypes (22, 26). This could especially be true for fungal specialists like *Ophiocordyceps* that rely on altered behavioral phenotypes to increase their fitness outcomes. Most studies that examined the microbiota–brain axis so far have focused on mouse models to demonstrate links between gut dysbiosis and neurodegenerative diseases, emotional state and social disfunctions (27-31). Nevertheless, *Drosophila* models have also been able to link gut dysbiosis with changes in short-chain fatty acids, which are considered an Alzheimer's marker (32, 33). In addition, studies with honeybees produced results that corroborate findings obtained in autism-spectrum disorder research with mice and humans (24). Furthermore, the gut microbiome of *Drosophila* has been linked to octopamine-dependent aggression behaviors (34), while ant gut bacteria have been suggested to impact chemical communication and, subsequently, olfactory-dependent social behaviors (35).

While there is previous work that connects insect pathologies and behavior with microbiota composition, the majority of microbiome research, including studies done on ants, focuses on comparisons between species, ecosystems, diet composition, and social status (25, 36-42). Most of these studies have only focused on the bacterial microbiota, leaving the fungal microbiome (i.e., mycobiome) underexplored. Studies that did investigate insect mycobiomes suggest that fungal communities can also be impacted by disease and show different patterns in community composition when compared to the bacterial microbiome (43). Both the bacterial and fungal microbiome of grain beetles infected with tapeworms showed changes when compared to uninfected individuals (44). Moreover, the gut mycobiome of honeybees was shown to be much more variable than their bacterial microbiome (36) and together with the bacterial gut community to be responsible for shaping social status (39). In fact, mycobiome explorations in mosquitoes demonstrated that interkingdom interactions between fungal and bacterial microbiome communities shape these communities, with consequences for host growth and development (45). These studies suggest that research into insect mycobiomes would provide a more complete picture of the microbiome while potentially revealing a different side of microbiome-host interactions. Moreover, studies that investigate the effects of infectious diseases on gut microbiota are currently few and far between, while these could reveal a greater involvement in disease outcomes than previously foreseen. With regards to the topic of fungal pathogens of insects, the intersection between insect pathology and the microbiome is even considered to be an avenue of research that might improve the use of these fungi in the biocontrol of insect pests (26).

Currently, the interaction between the behavior-manipulating fungus *Ophiocordyceps camponoti-floridani* and its carpenter ant host *Camponotus floridanus* is one of the more in-depth studied examples of summit disease (8, 10, 13, 19, 20, 46). Additionally, the bacterial microbiome of *C. floridanus* has been well characterized across different body regions (25)

and maturation levels (47). Nevertheless, the fungal communities and effects from disease on *C. floridanus microbiota* have yet to be assessed. Since microbiome composition has been linked to behavior in other insects, and infection can seemingly change that composition, perhaps part of the behavioral changes observed in *Ophiocordyceps*-manipulated ants can be attributed to their microbiota. As such, we set out to investigate the bacterial and fungal gut microbiomes of *Ophiocordyceps*-infected *C. floridanus* using metabarcoding. We compared our findings to data obtained from *Beauveria*-infected ants to answer if 1) fungal infection alters the bacterial and fungal community composition of the gut microbiome in *C. floridanus* when compared to healthy nestmates, 2) the gut microbiome community changes as disease progresses, and 3) the microbiome community is affected differently depending on the infecting entomopathogen (*Ophiocordyceps* vs *Beauveria*). We hypothesized that the microbiome of infected ants would diverge from that of healthy nestmates, and that *Ophiocordyceps* and *Beauveria* infections would affect microbiome compositions differently since they are fungi with contrasting parasitic lifestyles. If indeed true, this could suggest that changes in the host gut microbiota caused by *Ophiocordyceps* infection could underly some of the behavioral changes observed.

2. Methods

2.1 Ant collection and housing

To conduct our metabarcoding analysis of carpenter ant guts upon fungal infection, we collected two *C. floridanus* colonies from the University of Central Florida's arboretum (Colony 1 and 2) and two additional colonies from the Chuluota Wilderness area (Colony 3 and 4). We received permits for these collections from the UCF Arboretum and Seminole County Natural Lands, respectively. All collected colonies were queenless and averaged between one to two thousand individuals consisting of majors, minors, and brood.

Upon collection, we placed each colony into a 9.4 L plastic container (Rubbermaid) with ad libitum water and food (i.e., frozen crickets and 15% sugar water). Several darkened 50 mL tubes (Greiner) served as nest chambers and talcum-lined walls prevented the ants from climbing out. To reset *C. floridanus* biological clocks to lab conditions, we placed the colonies in an incubator (I36VL, Percival) under constant light, 25 °C temperature, and 70% relative humidity for 48 hours. Subsequently, we entrained the colonies to 24h rhythms of light and temperature (i.e., environmental entrainment cues known as Zeitgebers). We used a 12 hr :12 hr light-dark cycle at 28°C and 20°C, respectively, and 70% relative humidity. The cycle began with lights on at Zeitgeber time (ZT) 0 and followed a four-hour increasing ramping step to full light and 28°C at ZT 4. These settings were followed by a four hour hold at peak light and temperature until ZT 8, and a four-hour decreasing ramp step to complete darkness and 20°C at ZT 12.

Seven days prior to infections, we separated two groups of 110 minor caste workers from each colony into new 9.4 L plastic containers with a plaster bottom. *Tillandsia* is the main biting substrate for *Ophiocordyceps*-infected *C. floridanus* in Central Florida (Will et al., 2023). To provide this substrate for manipulated individuals to latch onto, we draped *Tillandsia* on two upright 12 cm wooden skewers embedded in the plaster on one end of the container. On the other end of the container, we placed a solid dark plastic bottom of a micropipette tip box (TipOne) upside down to serve as a nest area. We fed all fragment colonies ad libitum on 15% sucrose solution and water throughout the experiment.

2.2 Fungal culturing and ant infections

To infect carpenter ants with a behavior-manipulating hemi-biotroph-like fungus, we used *O. camponoti-floridani* strain OJ 2.1, isolated in January 2021 from a naturally infected *C. floridanus* ant cadaver found in Ocala National Forest. Seven days prior to infections, we started an 8 mL liquid culture by inoculating Grace's Insect Medium (Invitrogen) supplemented with 2.5% Fetal Bovine Serum (FBS) (Invitrogen) with blastospores obtained

from a frozen glycerol stock. We incubated the culture at 28 °C in the dark at 50 rpm. For necrotroph-like fungal infections we used *B. bassiana* strain ARSEF 2860. To obtain fresh conidiospores, we inoculated potato dextrose agar (PDA) by rolling cryobeads containing frozen spores across the surface. For blastospore growth, we harvested the fresh spores from a fully colonized PDA plate and inoculated a 20 mL liquid culture of Sabouraud dextrose broth (Sigma) six days prior to infections. We incubated liquid cultures at 25 °C in the dark at 120 rpm.

To conduct *Ophiocordyceps* and *Beauveria* infections, we harvested fungal blastospores immediately prior to injections by filtering the cultures through sterile gauze. Following centrifugation and supernatant removal, we resuspended the cells in fresh Grace's media supplemented with 2.5% FBS. *Ophiocordyceps* was resuspended to a cell concentration of 4.6 x 10^7 blastospores/mL, while the *Beauveria* infection solution was brought to 3.5 x 10^7 blastospores/mL. We performed infections with 10 µL borosilicate capillary tubes (Drummond) pulled to a needle point with a PC-100 Narishige instrument.

To establish *Ophiocordyceps* infections, we injected two times 110 ants, obtained from two different colonies (colony 1 and 2), on the ventral side of the thorax under the first pair of legs with 0.5 uL of blastospore solution. The healthy control groups also consisted of two times 110 ants, taken from the same two colonies, which were sham-injected with 0.5 mL Grace's media supplemented with 2.5% FBS without fungal cells (Will et al., 2020). Following injection, we left ants undisturbed for three days after which we removed deceased ants, considering them a result of injection trauma. As a result, 79% of *Ophiocordyceps*-injected and 85% of sham-injected ants survived, which were kept in the infection experiment for further observation, data collection and sampling.

While injection of fungal blastopores is necessary for *Ophiocordyceps* infection, *Beauveria* infections are best established by pricking ants with a capillary needle that was briefly dipped into a fungal blastospore solution (13). As such, we conducted *Beauveria* infections in two times 110 ants, again obtained from two different colonies (colony 3 and 4), by pricking them on the ventral side of the thorax under the first pair of legs with a blastospore-coated capillary needle. In this case, the control groups of two times 110 ants, taken from the same colonies, were pricked with a needle dipped in a sham solution of Grace's-FBS 2.5% without fungal cells. We left ants undisturbed for one day after which we removed deceased ants and counted all surviving ants as part of the experiment, which comprised 97% of *Beauveria*-pricked and 98% of sham-pricked ants.

2.3 Daily observations, survival data collection and ant sampling

After leaving *Ophiocordyceps*-infected ants undisturbed for three days post injection (DPI), we checked survival and removed dead individuals daily between ZT 5 and 7 (i.e., while lights were on). At 19 DPI, 25% of the infected population had died, which we indicated as lethal time (LT) 25 (Figure 1A). At this time, none of the infected individuals had shown manipulated summiting and biting behaviors yet. However, most individuals that make it across this time point eventually progress towards the ultimate summiting behavior (8). As such, to investigate potential microbiome changes prior to manipulated biting, we included LT25 as our first sampling time point (Figure 1A).

For our second *Ophiocordyceps* infection time point, we wanted to include live individuals that displayed the characteristic manipulated summiting and biting behavior (i.e., "Live Manipulation" or LM). Previous studies on *O. camponoti-floridani* infections found that manipulated biting occurs around ZT 21 under laboratory conditions (Will et al., 2020). So, as we neared the three-week mark when these manipulations typically begin, we also made daily observations around ZT 21-23 to catch ants in the act of summiting and biting. At 21 DPI, the first manipulations occurred (Figure 1A). We observed ants overnight, checking every two hours and found manipulations started between ZT 18 and 22, but could be as late as ZT 1 in one case. We did not collect ants until ZT 22 to avoid disturbing other ants

Figure 1: Sampling regime and survival curves of C. floridanus ants infected with O. camponoti-floridani or B. bassiana. *Visualization of the parallel sampling timelines that were used to collect A) Ophiocordyceps-infected and B) Beauveria-infected ants for this study. The corresponding survival curves of both these infections are plotted underneath these timelines, demonstrating that both infections led to a significant reduction in survival probability for both infections as compared to healthy control ants obtained from the same colony (p < 0.0001). For both infections two independently collected ant colonies were used: colony 1 and 2 (C1 and C2) were used for Ophiocordyceps infections and colony 3 and 4 (C3 and C4) were used for infections with Beauveria. The two colony replicates for each infection showed a comparable survival probability. Samples were collected at the time when 25% of the ants had died (i.e., lethal time LT25), ants showed manipulated or sickness behavior (i.e., live manipulated LM and live sick LS), ants had died after showing these behaviors (i.e., dead manipulated DM and dead sick DS), and 24 hours after the ant had succumbed from the respective infections (i.e., 24 hours post death 24PD).*

that might be climbing and biting. At this time, live manipulated ants were still very responsive (i.e., moving legs and antennae rapidly when gently poked with forceps).

After manipulation, *Ophiocordyceps* is thought to rapidly switch its biotroph-like growth towards a more necrotrophic one as it begins to switch from single celled blastospores to a multicellular hyphal growth, killing the ant host in the process (7). To capture potential microbiome changes during this process, for our third time point, we included ants that had just succumbed to the *Ophiocordyceps* infection after displaying manipulated summiting behavior (i.e., "Dead Manipulation" or DM) (Figure 1A). Ants for this time point were collected by conducting hourly checks following manipulation to monitor time of death, which occurs within a few hours upon manipulated biting. We considered ants dead when they no longer moved in response to being poked with forceps.

For our fourth timepoint, we included ants 24h after manipulation and death when tissue damage due to cell death and fungal carbon and nutrient consumption is more

abundant (i.e., "24Hr Post Death" or 24PD (Figure 1A). To collect these samples, we moved dead manipulated individuals to an isolated container with an identical setup to the original one, kept within the same incubator for 24h. Isolation protected the dead ants from being removed/dismembered by their live nestmates as part of their social immune response.

To collect *Beauveria*-infected ants we aimed to collect comparable timepoints to the *Ophiocordyceps* experiment. However, *Beauveria's* much more rapid disease progression required a different monitoring and sampling approach. After leaving ants undisturbed for one day, we conducted survival and behavioral observations every 4 hours. As we neared the LT25 timepoint, observations were done every two hours to ensure precise timing. Because ant mortality at this point was a result from infection followed by rapid death, there was necessary overlap in the LT25 sample collection and later timepoints (i.e., 'sick' and "dead"). The *Beauveria* infection experiment reached LT25 at 4 DPI (Figure 1B). To distinguish this first sampling group (LT25) from the second collection timepoint (the 'sick' phenotype), only ants that were out of the nest and able to walk at LT25 were selected.

Since *Beauveria* does not manipulate, we collected *Beauveria*-infected ants at a distinct "sick" phenotype, starting at 3 DPI (i.e., "live sickness behavior" or LS) (Figure 1B). Ants that displayed the "sick" phenotype could no longer walk more than a few steps without falling but still actively responded to a poke with forceps. They were often found on their sides while moving their legs but unable to stand. We considered this a comparable timepoint to the *Ophiocordyceps* ants collected at live manipulation because once *Beauveria*-infected ants displayed this behavior, they similarly died within a few hours (Figure 1B).

To collect *Beauveria*-infected "dead" ants we isolated "sick" ants in identical container setups to prevent disturbance from nest mates and monitored them hourly until death, after which they were immediately sampled (i.e., "dead sickness behavior", or DS). To collect *Beauveria*-infected ants 24h post death (24PD), we left expired ants that had displayed the "sick" phenotype in isolation for 24h under the same incubator conditions as the remainder of the experiment (Figure 1B).

We sampled ten *Ophiocordyceps*-infected ants, ten *Beauveria*-infected ants, and ten sham-injected and -pricked control ants at each timepoint (i.e., LT25, LM/LS, DM/DS, and 24PD). Three additional *Ophiocordyceps*-infected ants were collected at live manipulation due to availability. Eventually, we reduced the control group to comprise twelve healthy sham samples randomly selected from across the four *Ophiocordyceps*-infection timepoints and twelve from the *Beauveria*-infection timepoints. To maintain microbiome integrity, we flash froze ants in liquid nitrogen immediately following collection.

2.4 DNA extraction, library preparation, and sequencing

To conduct bacterial 16S and fungal ITS metabarcoding on healthy and infected ant guts, we processed all samples in a randomized order. First, we separated gasters from frozen ants under sterile conditions and placed them in 2 mL Eppendorf tubes with two metal ball bearings (5/32" type 2B, grade 300, Wheels Manufacturing). We disrupted the cells in frozen gasters by placing the tubes in a Mini G Tissue Homogenizer (SPEX) for 60 seconds at max rpm. We extracted DNA from these samples using the ZymoBIOMICS DNA microprep kit (Zymo Research, D4301) following manufacturer's recommendations. Using the extracted DNA, we performed library preparations according to the 16S Metagenomic Sequencing Library Preparation protocol from Illumina. We also included negative DNA extraction controls to account for potential contaminants that we introduced into our samples during this process.

For bacterial metabarcoding, we targeted the V4 region of the 16S SSU rRNA using primers 515F (48) and 806R (49). The internal transcribed spacer (ITS) region is predominantly used for fungal metabarcoding (50, 51). As such, we targeted the ITS2 region using ITS primers ITS3_KYO2 (51) and ITS4 (52). We conducted amplicon PCR (PCR1) in 25 μL reactions using 2.5 μL of DNA sample, 5 μL each of 1 μM forward and reverse primer, and 12.5 μL of KAPA HiFi HotStart ReadyMix (Roche). We ran the 16S amplicon reactions

in duplicate using the following PCR conditions: initial denaturation at 95°C for 3 minutes, followed by 25 cycles of 95°C for 30 seconds, 50°C for 30 seconds and 72°C for 30 seconds, and a final extension at 72°C for 5 minutes. We performed the ITS reactions in singleton using the same PCR conditions, with the exception that the annealing temperature used was 47°C. Following PCR, reaction we cleaned PCR1 products by removing primer dimers with 1.5x Sera-Mag SpeedBead solution (Fisher Scientific) following the protocol described in (53). The duplicate 16S PCR1 reactions were combined during this process. We also included negative PCR1 controls to account for potential contaminants we introduced into our samples.

Prior to the index PCR (PCR2), we checked the fragment size of randomly selected 16S and ITS libraries using a Tapestation 4200 (Agilent) and D1000 HS reagents (Agilent). For both 16S and ITS, we detected expected fragment sizes between 300-400 bp. We uniquely dual-indexed all ITS libraries using 50 μL PCR reactions, consisting of 8 μL PCR1 product, 5 μL each of Nextera XT Index primers i5 and i7 (Illumina), 7 μL molecular grade water (Invitrogen), and 25 μL KAPA HiFi HotStart ReadyMix (Roche). The 16S libraries were of higher concentrations and, therefore, indexed in 25 μL PCR reactions consisting of 8.5 μL PCR1 product, 2 μL each of Nextera XT Index primers i5 and i7, and 12.5 μL KAPA HiFi HotStart ReadyMix. We used the following PCR conditions for both the ITS and 16S libraries: initial denaturation at 95°C for 3 minutes, followed by 8 cycles of 95°C for 30 seconds, 55°C for 30 seconds and 72°C for 30 seconds, and a final extension at 72°C for 5 minutes. We cleaned PCR2 products with magnetic beads at a 1.2x concentration to remove smaller base pair fragments following the protocol as described for PCR1.

Following PCR2, we quantified all libraries using a Qubit fluorometer and an Invitrogen Qubit 1X dsDNA HS assay kit (Fisher Scientific) to combine equimolar concentrations of each library for sequencing. After an additional 1.2x magnetic bead cleanup, we created a high and a low concentration pool, quantified the two pools using an NEBNext Library Quant Kit for Illumina (New England Biolabs), and determined average fragment size and adaptor dimer removal on a Tapestation 4200 (Agilent) with D1000 HS reagents (Agilent). Subsequently, we created a final pool by combining the 16S and ITS library pools in equimolar amounts for sequencing using an Illumina MiSeq instrument with a paired-end approach (2x300bp) and V3 reagents (Illumina).

2.5 Sequence data processing

Prior to importing our paired-end read data into QIIME2 (54) for further processing and quality control, we demultiplexed the samples on the MiSeq. Using the QIIME2 *cutadapt trim-paired* function, we removed reads lacking primer regions and trimmed forward and reverse primer regions from reads that did have them. We then proceeded to denoise, dereplicate, and merge the trimmed forward and reverse reads using the *dada2 denoise-paired* function and created amplicon sequence variants (ASVs).

For ITS sequences, we performed denoising with and without truncation to determine the most optimal procedure for our data. The fungal ITS region can be rather variable in length. As such, it is generally not recommended to trim reads at the 3' end to prevent bias in amplicon size (55). However, not trimming reverse reads left many taxa unidentified after the taxonomy assignment. Similarly, we attempted to process ITS forward reads following the same procedures for single end data without truncation. This still left many taxa unidentified. The ITS reads with truncation proved to have the most identified reads compared to the other methods. Therefore, despite truncation may introduce some bias in amplicon size, we used our truncated data for downstream analysis.

We taxonomically classified 16S sequences using the Green Genes 13.5 database at 99% identity with BLAST+ (QIIME2, *classify-consensus-blast*; (56) and created a phylogenetic reference tree with a Green Genes 13.8 SATé-enabled phylogenetic placement database (QIIME2, *fragment-insertion sepp*; (57). We classified ITS sequences using a naïve bayes classifier (QIIME2, *fit-classifier-naive-bayes*) trained on the fungal 9.0 UNITE database (58). After assigning taxonomy, we filtered out 16S reads not identified at the phyla

level, assigned as chloroplast or mitochondria, and those that did not meet the set minimum of three reads per feature. Rarefaction curves (package: vegan – *rarecurve*) indicated that a sequencing depth of 4000 was sufficient to capture most species. Of all our collected samples, 94 passed this cutoff (extraction and PCR controls not included) (Supplementary File 1, Figure 1 and Table 1). We filtered ITS reads similarly, excluding reads that were not identified to a phyla level and requiring a minimum of three reads per feature. We also filtered out reads identified as *Ophiocordyceps spp.* and *Beauveria spp.* as these were the infecting entomopathogens. Rarefaction curves indicated sufficient species coverage at a sequencing depth of 500. Out of all our samples, 81 samples passed this cutoff (extraction and PCR controls not included) (Supplementary File 1, Figure 1 and Table 1).

2.6 Statistical analyses

All statistical analyses were conducted in R version 4.2.1. We analyzed survival probability of ants across the infection period with the R package survival version 3.5-8 (59) and constructed the survival curves using survminer version 0.4.9 (60).

To determine alpha and beta diversity of overall biological (infection) groups and separate infection timepoints, we paired classified reads and taxonomy data with sample metadata and imported this into R as phyloseq objects. We calculated within-sample alpha diversity using observed richness and Shannon diversity (package: phyloseq – *estimate_richness*) (61). To test for differences between diversities, we used a Kruskal-Wallis rank sum test, followed by pairwise comparisons using the Dunn test (package: dunn.test - *dunn.test*, p-value adjustment method for multiple comparisons = Bonferroni) (62). We visualized alpha diversity differences by generating box plots using ggplot2 (63).

To detect differences between groups and timepoints we also used beta diversity metrics and performed pairwise permutational multivariate analyses of variance (PERMANOVA) (package: pairwiseAdonis – *pairwise.adonis,* method = bray curtis, permutations = 999, p-value adjustment method for multiple comparisons = Bonferroni) (64). We visualized beta diversity using non-metric multidimensional scaling (NMDS) ordination of Bray-Curtis distances (package: phyloseq – *plot_ordination*) (61). We produced taxa bar plots using ggplot2 (63).

Using these methods, we compared sham-injected control ants that were obtained from the same colonies as the *Ophiocordyceps* infection and the *Beauveria* infection and found that their microbiota had no significant differences in alpha or beta diversity. Therefore, we combined them into a single healthy control group for comparisons to the infected ant groups. Moreover, this indicated that the ants used in these infection experiments had similar microbiota despite being from different colonies and field collections, allowing for direct comparisons between the two fungal infection treatments.

To determine whether the type of fungal infection (i.e., necrotroph-like versus hemi-biotroph-like) affected the bacterial and fungal community of the gut microbiome differently, we grouped reads from timepoints LT25 through 24HR into two categories based on the infecting entomopathogen (treatment groups): *Ophiocordyceps-* or *Beauveria*-infected ants. For timepoint comparisons, we separated reads from the two fungal groups back into their respective timepoints: LT25, live or sick, dead, 24HR, and sham. In doing so, we discovered one outlier ITS *Beauveria* 24HR sample that had high NMDS1 and NMDS2 scores (Supplementary File 1, Figure 2). After the filtering steps, and removal of *Ophiocordyceps* and *Beauveria* reads, this sample was only left with unidentified reads and reads identified as *Malassezia*, a likely contaminant. Therefore, we removed this sample from both alpha and beta diversity analyses.

3. Results

3.1 Overall data quantity and quality

Sequencing of ant samples resulted in 6,670,956 paired-end sequences across all samples for 16S libraries and 5,968,530 paired-end sequences across all samples for ITS libraries.

After filtering, taxonomic assignment, and rarefaction, 1280 ASVs across 94 samples for 16S libraries and 833 ASVs across 81 samples for ITS libraries remained Supplementary File 1, Table 1).

To account for and monitor levels of contamination introduced during DNA extraction and library preparation, we included negative controls for DNA extraction (extraction controls) and library preparation (PCR controls). Both technical controls had a read coverage similar to biological samples and a detected comparable number of taxa. Consequently, we could not reliably use these controls to remove potential contaminants from our samples. To investigate the source of contaminant introduction, we grouped samples by the date they were extracted and the first PCR for library preparation was performed and analyzed alpha diversity metrics (Observed richness, Shannon). While certain extraction dates were significantly different, there was no gradual increase in richness or Shannon diversity over time (Supplementary File 1, Figure 3). This indicates that the reagents themselves were not contaminated during the process and that the taxa observed in the negative controls were likely introduced due to some level of cross-contamination during extraction. Grouping samples by PCR date did not show any significant differences (Supplementary File 1, Figure 4), indicating that no measurable additional cross-contamination was introduced during library preparation. Taken together, our biological samples have some level of cross-contamination that might obscure present differences between groups. However, since we processed all samples in a random order (both extractions and PCRs), significant differences between treatment groups and infection time points that we do detect cannot be attributed to batch effects from the days they were processed.

3.2 Differences in alpha diversity between infection treatments

When we compared the alpha diversity of gut microbiota in healthy ants to those infected with *Ophiocordyceps* and *Beauveria*, irrespective of specific infection time points, we did not find an effect of fungal infection on bacterial (16S) alpha diversity (Figure 2A). However, the mycobiome (ITS) in *Ophiocordyceps*-infected ants had a significantly higher Shannon diversity index (p-value = 0.01) compared to healthy controls (Figure 2B, Supplementary File 2, Table 2). This indicates that *Ophiocordyceps* infection might affect the relative abundance of fungal taxa present within the healthy ant host gut. Assuming no new taxa were introduced, this increased Shannon index suggests that the evenness between present fungal taxa increased upon *Ophiocordyceps* infection.

When we compared the effects of *Ophiocordyceps* and *Beauveria* infections, we found that the gut microbiota of both infection groups significantly differed in their alpha diversity for both the bacterial microbiome and mycobiome (Figure 2A and B). The bacterial microbiome in the guts of *Ophiocordyceps*-infected ants had a significantly lower observed richness (p=0.003) and Shannon diversity index (p = 0.0003) when compared to those infected with *Beauveria* (Figure 2A, Supplementary File 2, Table 2). However, we observed the reverse for the mycobiome. Ants infected with *Ophiocordyceps* had a significantly higher observed richness (p = 0.006) and Shannon diversity index (p = 0.01) in their mycobiome compared to those infected with *Beauveria* (Figure 2B, Supplementary File 2, Table 2). This suggests that these fungal pathogens, which have different infection strategies, affect the bacterial microbiome and the mycobiome differently.

Figure 2: Differences in bacterial and fungal gut microbiota between healthy C. floridanus ants and those infected with O. camponoti-floridani or B. bassiana.
A and B) Box plot representation of the bacterial (16S) (A) and fungal (ITS) (B) alpha diversity in the guts of healthy sham-treated control ants (yellow), ants infected with Ophiocordyceps (purple), and ants infected with Beauveria (blue). We performed a Kruskal-Wallis test to determine if groups were significantly different (p<0,05) from each other. When found significant, we performed pairwise comparisons between groups using the Dunn test. Adjusted p-values of significantly different observed richness and Shannon diversity between infection groups are indicated as follows: * p<0.05, ** p< 0.01, *** p<0.001. C and D) Beta diversity non-metric multidimensional scaling (NMDS) plots for the different infection treatment groups. The points represent microbiomes of individual samples, and their color again indicates which treatment group the sample belongs to. C) NMDS plot using a Bray Curtis distance matrix of the bacterial gut microbiome for different infection groups (stress = 0.161). D) NMDS plot using a Bray Curtis distance matrix of the fungal gut microbiome for different infection groups (stress = 0.22). E and F) Taxa bar plots showing the relative abundance of bacterial (E) and fungal (F) orders present in the guts of healthy sham-treated ants, ants infected with Beauveria, and ants infected with Ophiocordyceps.

3.3 Beta diversity between infection treatments

We also used beta diversity metrics to determine if there is a difference in taxa abundance in

the guts of healthy and fungal infected ants. In doing so, we obtained results that corroborated our alpha diversity results. We again found that bacterial community compositions were not significantly different between infected ants and healthy controls (Figure 2C). Nevertheless, the fungal community composition was again significantly different between the *Ophiocordyceps* infected ants and healthy controls (PERMANOVA, p = 0.003) (Figure 2D, Supplementary File 2, Table 3). Moreover, similar to the alpha diversity comparisons, the beta diversity metrics also indicated that *Ophiocordyceps* and *Beauveria* treatments were significantly different from each other in both bacterial and fungal community composition (PERMANOVA, p = 0.033 and p = 0.003, respectively) (Figure 2C and D, Supplementary File 2, Table 3).

In line with findings from previous microbiome work done on *C. floridanus* (25, 65), Enterobacteriaceae constituted the most abundant order of bacteria in the microbiome of ants from all treatment groups and were predominately composed of *Blochmannia,* a known endosymbiont of the C*amponotini* tribe. Taxa bar plots for the bacterial microbiome (Figure 2E) show a decrease in taxa of this order within the *Beauveria* infection group. This suggests that this might be the main cause of the significant difference between the bacterial gut microbiota of *Ophiocordyceps-* and *Beauveria-*infected ants. Taxa bar plots for the mycobiome (Figure 2F) showed that *Ophiocordyceps*-infected ants contained a greater percentage of unknown fungal taxa in their gut compared to the *Beauveria*-infected and healthy sham-treated ants, which likely accounts for the found difference in community composition. *Beauveria-* and sham-treated ants also contained a greater percentage of fungi within the *Ophiostomatales* order, mainly comprised of the species *Sporothrix insectorum,* which is regularly reported to be a pathogen of insects (66, 67)*.*

3.4 Differences in alpha diversity in *Ophiocordyceps*-infected ants over time
We additionally compared the alpha diversity of gut microbiota in healthy ants and different progressive *Ophiocordyceps* infection time points. These time points comprised 1) infected, live ants, while 25% had succumbed to the infection, before the ultimate manipulated summiting and biting phase (lethal time 25%; LT25), 2) live ants that displayed manipulated summiting and biting behavior (live manipulation; LM), 3) ants that died a few hours after manipulation took place (dead after manipulation; DM), and 4) ants that had been dead for 24 hours after manipulation (24hr post death after manipulation; 24PD) (Figure 3A and B). These comparisons indicated that the alpha diversity composition of the ant gut microbiome is significantly different during live behavioral manipulation (LM) compared to both earlier and later time points. Only the dead manipulated time point (DM), which was taken only a few hours later, was similar. Looking at the observed richness in the bacterial microbiome (Figure 3A), live ants that are showing the manipulated summiting and biting phenotype have significantly fewer species, or lower taxa evenness, in their gut compared to healthy controls (p = 0.005), infected ants prior to manipulation (p = 0.022), and those 24 hours after death (p = 0.03) (Supplementary File 2, Table 4). The mycobiome again shows the opposite pattern (Figure 3B). Live manipulated ants have an increased observed richness in the number of species present in their gut when compared to healthy shams (p = 0.014), infected ants before manipulation (p = 0.006), and those sampled 24 hours after dying in the manipulated summiting and biting position (p = 0.048) (Supplementary File 2, Table 4). Differences in the mycobiome alpha diversity are again more apparent than bacterial diversity differences since we observed the same significant differences in the Shannon diversity index across timepoints in *Ophiocordyceps*-infected ants (Figure 3B); Live manipulation samples had a higher Shannon diversity index compared to healthy sham samples (p < 0.0001), LT25 (p = 0.012), and 24PD samples (p = 0.034) (Supplementary File 2, Table 4). This suggests that the number of taxa and/or the evenness of the taxa in the

guts of ants that show *Ophiocordyceps*-adaptive summing and biting behaviors is significantly different compared to healthy ants, and even those in other stages of the infection.

Figure 3: Differences in bacterial and fungal gut microbiota of C. floridanus during infection progression and manipulation by O. camponoti-floridani.
A and B) Box plot representation of the bacterial (16S) (A) and fungal (ITS) (B) alpha diversity in the guts of healthy sham-treated control ants (yellow) and ants infected with Ophiocordyceps, which were sampled before (LT25), during (LM) and after manipulated summing and biting behavior (DM and 24PD) (different shades of purple). We performed a Kruskal-Wallis test to determine if groups were significantly different (p<0.05) from each other. When found significant, we performed pairwise comparisons between groups using the Dunn test. Adjusted p-values of significantly different observed richness and Shannon diversity between infection groups are indicated as follows: * p<0.05, ** p< 0.01, *** p<0.001. C and D) Beta diversity non-metric multidimensional scaling (NMDS) plots for the different Ophiocordyceps infection time points. The points represent microbiomes of individual samples, and their color again indicates which infection time point the sample belongs to. C) NMDS plot using a Bray Curtis distance matrix of the bacterial (16S) gut microbiome for different infection time points (stress = 0.135). D) NMDS plot using a Bray Curtis distance

matrix of the fungal (ITS) gut microbiome for different infection time points (stress = 0.195). E and F) Taxa bar plots at order level showing the relative abundance of bacterial (E) and fungal (F) microbiota present in the guts of healthy sham-treated ants, and ants infected with Ophiocordyceps prior (LT25), during (LM) and after manipulation (DM and 24PD).

3.5 Beta diversity between *Ophiocordyceps*-infected ants over time

To further investigate the potential difference in gut taxa abundance between behaviorally manipulated ants and other stages of infection, we also used beta diversity metrics. When looking at the bacterial microbiome of *Ophiocordyceps*-infected ants (Figure 3A), we only found a significant difference between the guts of live manipulated ants and those that were sampled 24 hours after manipulation and death (PERMANOVA, p = 0.04) (Supplementary File 2, Table 5). In contrast, the mycobiome of *Ophiocordyceps*-infected ants (Figure 3B) again showed significant differences in community composition at several timepoint comparisons. Guts from ants that were sampled during manipulated summing behavior had significantly different fungal taxa abundances from healthy controls (p = 0.01), infected ants prior to manipulation (LT25 p = 0.02) and ants that were sampled 24 hours after they succumbed to the infection (24PD p = 0.01) (Supplementary File 2, Table 5). This strengthens the suggestion from the alpha diversity analysis that the fungal microbiome is significantly different in infected ants when *Ophiocordyceps* induces manipulated summing and biting behavior. Taxa bar plots (Figure 3F) indicate a sudden increase in the relative abundance of unidentified taxa at the behavioral manipulation (LM) timepoint and a large decrease in taxa of the order *Ophiostomatales,* the most prevalent fungal taxa. In the samples from the two time points taken after behavioral manipulation (DM and 24PD), there is again an increase in the relative abundance of *Ophiostomatales* and a decrease in unidentified taxa, indicating a gradual shift in community composition.

3.6 Alpha and beta diversity in *Beauveria*-infected ants over time

We also looked into the gut micro- and mycobiota of ants infected with *Beauveria* at infection time points that we consider to be parallels to those we investigated for *Ophiocordyceps* infection. As such, we sampled *Beauveria*-infected ants at LT25, when showing non-manipulative "sick behavior" (LS), having died after showing sick behavior (DS) and 24 hours post death (24PD). While the gut taxa composition of *Ophiocordyceps*-infected ants that showed the manipulation phenotype (LM) was different from healthy and other infection stages, the gut composition of *Beauveria*-infected ants that showed the "sick" phenotype (LS) was not. Comparing bacterial and fungal alpha diversity across infection time points, we only detected a significant difference in the observed richness between the mycobiome of the healthy controls and the *Beauveria*-infected ants that did not show the sick phenotype yet (LT25, p = 0.034) (Supplementary File 2, Table 6). Yet, the Shannon diversity index did not corroborate this finding (Figure 4B). Additionally, our beta diversity analyses gave another, singular and distinct result. Using this metric, we found that only the bacterial microbiome of ants 24 hours after they had succumbed to their *Beauveria*-infection (24PD) was significantly different from the healthy controls (PERMANOVA, p=0.01) and the early infection time point in which a sick phenotype was not detected yet (LT25) (PERMANOVA, p = 0.04) (Figure 4C, Supplementary File 2, Table 7). This suggests that *Beauveria* infection does not affect the host bacterial or fungal microbiome consistently enough to detect significant differences between infection stages. The taxa bar plots of different *Beauveria* infection stages indicate that the mycobiome of *Beauveria*-infected ants (Figure 4F) is dominated by *Ophiostomatales* while the relative abundance of unidentified fungal taxa as seen in *Ophiocordyceps*-infected ants is a lot smaller. This suggests that the increased relative abundance of these ant gut taxa, especially during behavioral manipulation, might be

a specific hallmark of the infection strategies of *Ophiocordyceps*.

Figure 4: Bacterial and fungal gut microbiota of C. floridanus during infection by B. bassiana.

A and B) Box plot representation of the bacterial (16S) (A) and fungal (ITS) (B) alpha diversity in the guts of healthy sham-treated control ants (yellow) and ants infected with Beauveria, which were sampled at different time points during infection (different shades of teal). We performed a Kruskal-Wallis test to determine if groups were significantly different (p<0.05) from each other. When found significant, we performed pairwise comparisons between groups using the Dunn test. The adjusted p-value of the sole significantly different observed richness between 24 hour after death (24PD) and healthy sham control is indicated as * p<0.05. C and D) Beta diversity non-metric multidimensional scaling (NMDS) plots for the different Beauveria infection time points. The points represent microbiomes of individual samples, and their color again indicates which infection time point the sample belongs to. C) NMDS plot using a Bray Curtis distance matrix of the bacterial (16S) gut microbiome for different infection time points (stress = 0.159). D) NMDS plot using a Bray Curtis distance matrix of the fungal (ITS) gut microbiome for different infection time points (stress = 0.236). E and F) Taxa bar plots at order level showing the relative abundance of

bacterial (E) and fungal (F) microbiota present in the guts of healthy sham-treated ants, and ants infected with Beauveria prior to displaying a "sick" phenotype (LT25), during (LS) and after displaying the "sick" phenotype (DS and 24PD).

4. Discussion

Fungal pathogens can have vastly different infection strategies, ranging from necrotrophic to biotrophic. In addition to affecting the host and its tissues differently, these pathogens might impact its microbiota in different ways as well. To test this hypothesis for insect-infecting fungi specifically, we investigated the effect of infection with *O. camponoti-floridani,* a specialist behavior-manipulating hemi-biotroph-like pathogen of *C. floridanus*, and *B. bassiana*, a generalist non-manipulating necrotroph-like pathogen, on the gut microbiota of carpenter ant *C. floridanus*. When comparing the bacterial microbiome (16S) of healthy ants to *Ophiocordyceps*- and *Beauveria*-infected counterparts we did not find any significant differences. The bacterial community composition of infected ants apparently remains relatively similar to that of healthy controls. This would indicate that fungal infection is not impacting the ant gut microbiome as hypothesized. However, we found that the mycobiome (ITS) of the *Ophiocordyceps* infected ants had a significantly higher Shannon diversity index compared to the mycobiome of healthy controls with a corroborating significantly different community composition. Thus, based on the mycobiome, one would conclude that behavior manipulating *Ophiocordyceps* infection does alter the host gut microbiota. The mycobiome appears to be more dynamic in its composition compared to the bacterial microbiome, but also more diverse in gut colonizing taxa. *Blochmannia* was by far the most dominant colonizer of the carpenter ant gut both in this and other studies (25, 65), which might obscure changes other than those in the dominating species. Instead, there seems to be more evenness between the fungal taxa, which might make subtle differences easier to detect. Indeed, studies on other insect species also indicate that the mycobiome can be more variable than the bacterial microbiome (36, 68). Moreover, we detected an abundant presence of *S. insectorum*, which is commonly known as an entomopathogen (66, 67), but not necessarily as an insect gut symbiont. Similarly, known pathogens of the genus *Ophiocordyceps* itself have been detected in the microbiomes of plants and beetles (69, 70). As such, exploring the mycobiome can also reveal alternative lifestyles of presumed primary pathogens. Taken together, this all suggests that it is worthwhile to include the mycobiome in insect microbiota studies, which are currently still largely overlooked.

We also found that both the bacterial and fungal diversity and composition were significantly different when comparing *Ophiocordyceps* and *Beauveria* infections to each other. The bacterial microbiome of *Ophiocordyceps*-infected ants had a significantly lower species diversity compared to that of the *Beauveria*-infected ants. In contrast, the mycobiome of *Ophiocordyceps*-infected ants had a significantly higher species diversity. This indicates that these entomopathogenic fungi with contrasting infection strategies impact the overall gut microbiota differently, or that one of them causes a dysbiosis while the other one leaves the gut composition unchanged. Indeed, when we compared healthy and progressive infection time points from early infection until after host death, we did not detect consistent significant changes in the gut of *Beauveria*-infected ants when analyzing alpha and beta diversity. This suggests that *Beauveria* infection leaves the gut composition relatively unchanged, with perhaps some gradual changes leading up to the fungus' saprophytic growth after death as the significant differences in beta diversity between the 24 hours post death (24PD) time point and the healthy and early infection time point (LT25) suggest.

In contrast, in *Ophiocordyceps*-infected ants, the micro- and mycobiome both displayed a notable shift in species richness at live manipulation (LM). During this time, the ant host shows a characteristic summiting behavior adaptive to the development and spread of the fungus (9, 10). The alpha diversity for the LM timepoint in both microbiomes significantly differed from the healthy control, LT25 and 24PD timepoints, which may be an

indication of a gradual change across the infection. However, these other time points were not significantly different from one another, suggesting that something uniquely different might be happening with the gut biota during manipulated summiting behavior. At the LM timepoint, the bacterial microbiome again showed a sudden drop in species richness while there was a significant increase seen in the mycobiome. The only time point during which the gut biota was not significantly different compared to the LM timepoint was death after manipulation (DM). For *Ophiocordyceps*-infected ants, death occurred within a few hours of live manipulation, as compared to the 24+ hours to multiple days that separated the other timepoints. These few hours may not have been enough time for the micro- and mycobiota to significantly change. However, DM not being indicated as significantly different from 24PD, while LM was, suggests that there is already a gradual change during that short amount of time. While the beta diversity analysis strengthened these findings for the mycobiome, they did not do so entirely for the bacterial microbiome. This suggests again that the fungal composition of the gut appears to be more dynamic than the bacterial composition and that *Ophiocordyceps* infection is somehow significantly affecting that composition, especially during the manipulated summiting stage.

These different disease outcomes for the host gut micro- and mycobiota could be due to a direct effect of the pathogen on the host and its tissues. Alternatively, it could be a result of changed behavior upon infection. Many social insects, including *Camponotus* ants, engage in trophallaxis – the social exchange of fluids between individuals. Such social fluids contain (partially) digested and regurgitated food materials as well as that individual's endocrine molecules that play a role in maintaining the social network and decision-making within an ant colony. Trophallaxis is also hypothesized to play a role in inoculating and maintaining gut microbiota among colony members (71). Thus, a disconnect from the social network of the colony could have potential impacts on an individual's microbiome. Indeed, *Ophiocordyceps*-infected ants display diminished communication with nestmates (13), which consequently might reduce trophallaxis. As such, they might be disconnected from the social network in the colony with the consequence that their regular gut microbiota is not maintained. While the period between infection, manipulation and eventual death spans several weeks in *Ophiocordyceps, Beauveria* infection runs a more rapid course of 5 days to a week. Diminished communication was not readily observed in *C. floridanus* infections with *Beauveria* (13). However, even if this were to occur, the disruption might be too short to have a measurable influence on the micro- and mycobiota. Therefore, a prolonged detachment from the social-fluid exchange due to asocial behavior might result in the microbiome dysbiosis seen in *Ophiocordyceps*-infected ants, which could explain the differences between the microbiomes of *Ophiocordyceps*- versus *Beauveria*-infected ants. However, if this were the case, one might expect a more gradual change over time with significance compared to healthy controls being found for all later stages of infection (LM, DM and 24PD) and not just the single live manipulation time point.

Of course, we cannot completely exclude that our non-significant results are caused by the nature of our samples and sample size. The survival curve for the *Ophiocordyceps* infections (Figure 1A) followed expected trends with most infected ants dying once manipulations began (21 DPI). The survival curve for *Beauveria* infections (Figure 1B) also matched expected trends with a decline in survival starting between 3-5 DPI. However, the survival rate levelled out after 6 DPI with around 50% of the population still alive. This indicates that not all ants progressed through the infection the same way and some were able to clear it. The sick (LS), dead (DS), and 24 post death (24PD) *Beauveria* timepoint samples are unaffected by this, as they were sampled based on a distinguishable sick behavior phenotype. However, the LT25 timepoint samples may be a mixture of infected and uninfected individuals as there was no clear way to distinguish infection status for those ants. Moreover, sample size was limited for each timepoint and might not have been robust

enough to reveal the full effects of fungal infection over time. Yet, the presence of dysbiosis in the guts of *Ophiocordyceps*-infected ants that displayed summiting behavior was clearly significantly different compared to most other infection time points, including healthy ants. In fact, looking at the NMDS plots (Figure 3C and D), the LM samples nicely cluster together while the other biological groups seem to show a more dispersed level of variation. As such, we consider the finding that *Ophiocordyceps* infection affects the gut micro- and mycobiome of ants during manipulated summiting to be evident.

What the biological relevance could be of a distinctly altered gut microbiome composition during manipulated summiting behavior currently remains a topic of speculation. One possibility could be that the ant gut microbiome affects host behavior and could be (partially) involved in the expression of stereotypically altered behaviors during *Ophiocordyceps* infection. There is mounting evidence from human research that gut dysbiosis correlates with an increased chance of Alzheimer's, depression, and other behavioral pathologies (72). Some studies suggest that this occurs via microbial metabolites secreted in the gut interacting with neurological receptors and G protein coupled receptors (GPCRs) (73, 74). Transcriptomics analyses of *Ophiocordyceps*-infected ants that displayed manipulated summiting behavior showed enrichment for GPCRs among the genes that were downregulated during this infection stage as compared to healthy individuals (8). These GPCRs were annotated to have functions involved in the perception of light and smell, and the binding of biogenic amines such as dopamine. Most of the pathogen-related behavioral changes modulated by these GPCRs are likely due to the predicted binding of *Ophiocordyceps* secreted proteins that are upregulated during manipulated summiting (20). However, other GPCRs might be interacting with (altered levels of) microbial peptides secreted by the microbiome. Additionally, there is increasing evidence that gut microbiomes aid in insect host immune defense and microbiome homeostasis is thought to be an important factor in mediating the immune system (26, 75). For instance, recent research into honeybee core microbiomes showed that its dysbiosis was linked to higher susceptibility to pathogen invasion and consequently higher mortality rates (75). Considering that a hemibiotroph-like pathogen such as *Ophiocordyceps* needs to ensure prolonged colonization and successful manipulation of the ant without causing significant physical damage, this fungus could perhaps be affecting the microbiome such that the host immune response is effectively lowered. This hypothesis would be in line with metabolomics data that suggest a lowered immune response in the ant paired with the presence of neuroprotectants (19, 76). However, one would expect this strategy to be needed during the earlier time points in infection as well, where we did not detect any significant micro- nor mycobiome changes. Taken together, the potential pathogen-adaptive function of gut dysbiosis during *Ophiocordyceps* infection, if any, remains to be discovered. Nevertheless, both the hypotheses we propose here could provide an explanation as to why we do detect this dysbiosis in infections with the hemibiotroph-like, behavior-manipulating specialist *Ophiocordyceps* and not in those with the necrotroph-like, non-manipulating generalist *Beauveria*.

5. Conclusions

To our knowledge, all previous research done on the microbiota of ants has focused on their bacterial microbiome, making our study the first to investigate the fungal taxa present within the ant gut. Our data show that, while the bacterial taxa are often dominated by one species, the fungal taxa are highly diverse with more evenness between them. Moreover, our data also suggest that the fungal taxa composition is more dynamic, as microbiome changes were more prominently detected in the fungal aspects of the gut. This supports the notion that this side of the microbiome likely plays a more prominent role in insect functioning than previously credited and should be studied more intensely. Moreover, we provided the first evidence that the gut microbiome of *C. floridanus*, and with that perhaps also other insects, appears to be impacted differently by infecting entomopathogens with different lifestyles. More specifically, our data suggest that behavior-manipulating specialists like *O. camponoti-*

floridani might affect the host microbiome in a timed manner that aligns with pathogen-adaptive host behavioral phenotypes. As such, considering the role that microbiota might play in host behavior, part of the behavioral manipulation might be the result of dysbiosis of the gut microbiota. Research in which specific modulation of gut taxa is linked to host behavior and infection status will be needed to begin to unravel these potential intersections. This will improve our understanding of the gut-brain axis and provide knowledge that could improve the effective use of fungal pathogens in the biocontrol of insect pests.

CRediT authorship contribution statement

Sophia Vermeulen: Data curation, Formal Analysis, Investigation, Methodology, Writing – original draft. **Anna Forsman:** Formal Analysis, Methodology, Supervision, Writing – review and editing. **Charissa de Bekker:** Conceptualization, Formal Analysis, Funding acquisition, Methodology, Supervision, Writing – review and editing.

Declaration of competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank Ian Will, Biplabendu Das, and Erandeni Ponce de Leon for their assistance with ant collection, infection, and monitoring. We would also like to thank the Fall 2022 UCF Genomics Course undergraduate students for being involved with the DNA extractions. Funding to conduct this study was obtained from the National Science Foundation, CAREER IOS-1941546, awarded to Charissa de Bekker.

Appendix. Supplementary materials

Supplementary File 1: This file presents figures and tables that support our technical data and sample validation analyses.
Supplementary File 2: This file contains tables that present the output of the alpha and beta diversity statistical analyses performed in this study.

Data availability

All data generated in this study can be found…

References

1. Araújo JPM, Hughes DP. Chapter One - Diversity of entomopathogenic fungi: Which groups conquered the insect body? In: Lovett B, St. Leger RJ, editors. Advances in Genetics. 94: Academic Press; 2016. p. 1-39.
2. Maina U, Galadima I, Gambo F, Zakaria D. A review on the use of entomopathogenic fungi in the management of insect pests of field crops. J Entomol Zool Stud. 2018;6(1):27-32.
3. Mascarin GM, Jaronski ST. The production and uses of *Beauveria bassiana* as a microbial insecticide. World J Microbiol Biotechnol. 2016;32(11):177.
4. Araújo JPM, Evans HC, Geiser DM, Mackay WP, Hughes DP. Unravelling the diversity behind the *Ophiocordyceps unilateralis* (Ophiocordycipitaceae) complex: Three new species of zombie-ant fungi from the Brazilian Amazon. Phytotaxa. 2015;220(3):224-38.
5. Araújo PM, Evans HC, Kepler R, Hughes DP. Zombie-ant fungi across continents: 15 new species and new combinations within *Ophiocordyceps*. I. Myrmecophilous hirsutelloid species. Studies in Mycology. 2018(90):119-60.
6. Araújo JPM, Evans HC, Fernandes IO, Ishler MJ, Hughes DP. Zombie-ant fungi across continents: II. Myrmecophilous hymenostilboid species and a novel zombie lineage. Mycologia. 2020;112(6):1138-70.

7. de Bekker C, Ohm RA, Loreto RG, Sebastian A, Albert I, Merrow M, et al. Gene expression during zombie ant biting behavior reflects the complexity underlying fungal parasitic behavioral manipulation. BMC Genomics. 2015;16(1):620.

8. Will I, Das B, Trinh T, Brachmann A, Ohm RA, de Bekker C. Genetic underpinnings of host manipulation by *Ophiocordyceps* as revealed by comparative transcriptomics. G3 Genes|Genomes|Genetics. 2020;10(7):2275-96.

9. Andersen SB, Ferrari M, Evans HC, Elliot SL, Boomsma JJ, Hughes DP. Disease dynamics in a specialized parasite of ant societies. PLoS One. 2012;7(5):e36352.

10. Will I, Linehan S, Jenkins DG, de Bekker C. Natural history and ecological effects on the establishment and fate of Florida carpenter ant cadavers infected by the parasitic manipulator *Ophiocordyceps camponoti-floridani*. Functional Ecology. 2022;37(4):886-99.

11. Andersen SB, Gerritsma S, Yusah KM, Mayntz D, Hywel-Jones NL, Billen J, et al. The life of a dead ant: The expression of an adaptive extended phenotype. American Naturalist. 2009;174(3):424-33.

12. Fredericksen MA, Zhang Y, Hazen ML, Loreto RG, Mangold CA, Chen DZ, et al. Three-dimensional visualization and a deep-learning model reveal complex fungal parasite networks in behaviorally manipulated ants. Proceedings of the National Academy of Sciences. 2017;114(47):12590-5.

13. Trinh T, Ouellette R, de Bekker C. Getting lost: the fungal hijacking of ant foraging behaviour in space and time. Animal Behaviour. 2021;181:165-84.

14. Andriolli FS, Ishikawa NK, Vargas-Isla R, Cabral TS, de Bekker C, Baccaro FB. Do zombie ant fungi turn their hosts into light seekers? Behavioral Ecology. 2019;30(3):609-16.

15. Loreto RG, Elliot SL, Freitas MLR, Pereira TM, Hughes DP. Long-Term Disease Dynamics for a Specialized Parasite of Ant Societies: A Field Study. PLOS ONE. Aug 18, 2014;9(8).

16. de Bekker C, Beckerson William C, Elya C. Mechanisms behind the madness: How do zombie-making fungal entomopathogens affect host behavior to increase transmission? mBio. 2021;12(5):e01872-21.

17. Das B, Brachmann A, De Bekker C. Both behavior-manipulating and non-manipulating entomopathogenic fungi affect rhythmic gene expression in carpenter ant foragers upon infection. 2023.

18. de Bekker C, Das B. Hijacking time: How *Ophiocordyceps* fungi could be using ant host clocks to manipulate behavior. Parasite Immunol. 2022;44(3):e12909.

19. Will I, Attardo GM, de Bekker C. Multiomic interpretation of fungus-infected ant metabolomes during manipulated summit disease. Scientific Reports. 2023;13(1):14363.

20. Will I, Beckerson WC, de Bekker C. Using machine learning to predict protein–protein interactions between a zombie ant fungus and its carpenter ant host. Scientific Reports. 2023;13(1):13821.

21. Brinker P, Fontaine MC, Beukeboom LW, Falcao Salles J. Host, Symbionts, and the Microbiome: The Missing Tripartite Interaction. Trends in Microbiology. 2019;27(6):480-8.

22. Koch H, Schmid-Hempel P. Gut microbiota instead of host genotype drive the specificity in the interaction of a natural host-parasite system. Ecology Letters. 2012;15(10).

23. Feldhaar H. Bacterial symbionts as mediators of ecologically important traits of insect hosts. Ecological Entomology. 2011;36(5).

24. Liberti J, Engel P. The gut microbiota - brain axis of insects. Curr Opin Insect Sci. 2020;39:6-13.

25. Ramalho MO, Moreau CS, Bueno OC, Ramalho MO, Moreau CS, Bueno OC. The potential role of environment in structuring the microbiota of *Camponotus* across parts of the body. Advances in Entomology. 2019;7(3).

26. Boucias DG, Zhou Y, Huang S, Keyhani NO. Microbiota in insect fungal pathology. Applied Microbiology and Biotechnology. 2018;102(14):5873-88.

27. Buffington SA, Di Prisco GV, Auchtung TA, Ajami NJ, Petrosino JF, Costa-Mattioli M.

    Microbial reconstitution reverses maternal diet-induced social and synaptic deficits in offspring. Cell. 2016;165(7):1762-75.

28. Sgritta M, Dooling SW, Buffington SA, Momin EN, Francis MB, Britton RA, et al. Mechanisms Underlying Microbial-Mediated Changes in Social Behavior in Mouse Models of Autism Spectrum Disorder. Neuron. 2019;101(2):246-59.e6.

29. Sampson TR, Debelius JW, Thron T, Janssen S, Shastri GG, Ilhan ZE, et al. Gut Microbiota Regulate Motor Deficits and Neuroinflammation in a Model of Parkinson's Disease. Cell. 2016;167(6):1469-80.e12.

30. Vuong HE, Yano JM, Fung TC, Hsiao EY, Vuong HE, Yano JM, et al. The microbiome and host behavior. Annual Review of Neuroscience. 2017;40(Volume 40, 2017).

31. Shen L, Ji H-F. Associations between gut microbiota and Alzheimer's disease: Current evidences and future therapeutic and diagnostic perspectives. Journal of Alzheimer's Disease. 2019;68(1).

32. Dalile B, Van Oudenhove L, Vervliet B, Verbeke K. The role of short-chain fatty acids in microbiota-gut-brain communication. Nat Rev Gastroenterol Hepatol. 2019;16(8):461-78.

33. Kong Y, Jiang B, Luo X. Gut microbiota influences Alzheimer's disease pathogenesis by regulating acetate in *Drosophila* model. Future Microbiol. 2018;13:1117-28.

34. Jia Y, Jin S, Hu K, Geng L, Han C, Kang R, et al. Gut microbiome modulates *Drosophila* aggression through octopamine signaling. Nature Communications. 2021;12(1):2698.

35. Teseo S, van Zweden JS, Pontieri L, Kooij PW, Sørensen SJ, Wenseleers T, et al. The scent of symbiosis: gut bacteria may affect social interactions in leaf-cutting ants. Animal Behaviour. 2019;150:239-54.

36. Decker LE, San Juan PA, Warren ML, Duckworth CE, Gao C, Fukami T. Higher variability in fungi compared to bacteria in the foraging honey bee gut. Microbial Ecology. 2023;85(1):330-4.

37. Mendoza-Guido B, Rodríguez-Hernández N, Ivens ABF, von Beeren C, Murillo-Cruz C, Zuniga-Chaves I, et al. Low diversity and host specificity in the gut microbiome community of Eciton army ants (Hymenoptera: Formicidae: Dorylinae) in a Costa Rican rainforest. Myrmecological News. 2023;33:19-34.

38. Hu Y, Sanders JG, Łukasik P, D'Amelio CL, Millar JS, Vann DR, et al. Herbivorous turtle ants obtain essential nutrients from a conserved nitrogen-recycling gut microbiome. Nature Communications. 2018;9(1):964.

39. Yun J-H, Jung M-J, Kim PS, Bae J-W. Social status shapes the bacterial and fungal gut communities of the honey bee. Scientific Reports. 2018;8(1):2019.

40. Martins C, Moreau CS. Influence of host phylogeny, geographical location and seed harvesting diet on the bacterial community of globally distributed *Pheidole* ants. PeerJ. 2020;8:e8492.

41. Ramalho MO, Moreau CS. Untangling the complex interactions between turtle ants and their microbial partners. Animal Microbiome. 2023;5(1):1.

42. Russell JA, Moreau CS, Goldman-Huertas B, Fujiwara M, Lohman DJ, Pierce NE, et al. Bacterial gut symbionts are tightly linked with the evolution of herbivory in ants. Proceedings of the National Academy of Sciences. 2009;106(50).

43. Wu H, Rao Z-C, Cao L, De Clercq P, Han R-C. Frontiers | Infection of Ophiocordyceps sinensis Fungus Causes Dramatic Changes in the Microbiota of Its Thitarodes Host. Frontiers in Microbiology. 2020/12/03;11.

44. Fredensborg BL, Fossdal Í Kálvalíð I, Johannesen TB, Stensvold CR, Nielsen HV, Kapel CMO. Parasites modulate the gut-microbiome in insects: A proof-of-concept study. PLoS One. 2020;15(1):e0227561.

45. Hegde S, Khanipov K, Hornett EA, Nilyanimit P, Pimenova M, Saldaña MA, et al. Interkingdom interactions shape the fungal microbiome of mosquitoes. Animal Microbiome. 2024;6(1):11.

46. Das B, de Bekker C. Time-course RNASeq of *Camponotus floridanus* forager and nurse ant brains indicate links between plasticity in the biological clock and behavioral division of labor. BMC Genomics. 2022;23(1).

47. Ramalho MO, Bueno OC, Moreau CS. Species-specific signatures of the microbiome from Camponotus and Colobopsis ants across developmental stages. PLOS ONE. Nov 22, 2017;12(11).
48. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. Environmental Microbiology. 2016;18(5).
49. Apprill A, McNally SP, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial Ecology. 2015;75(2).
50. Tedersoo L, Bahram M, Zinger L, Nilsson RH, Kennedy PG, Yang T, et al. Best practices in metabarcoding of fungi: From experimental design to results. Mol Ecol. 2022;31(10):2769-95.
51. Toju H, Tanabe AS, Yamamoto S, Sato H. High-Coverage ITS Primers for the DNA-Based Identification of Ascomycetes and Basidiomycetes in Environmental Samples. PLOS ONE. 2012;7(7).
52. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. PCR Protocols. San Diego: Academic Press; 1990. p. 315-22.
53. Forsman AM, Hoenig BD, Gaspar SA, Fischer JD, Siegrist J, Fraser K. Evaluating the impacts of metabarcoding primer selection on DNA characterization of diet in an aerial insectivore, the Purple Martin. Ornithology. 2021;139(1).
54. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology. 2019;37(8):852-7.
55. Rolling T, Zhai B, Frame J, Hohl TM, Taur Y. Customization of a DADA2-based pipeline for fungal internal transcribed spacer 1 (ITS1) amplicon data sets. JCI Insight. 2022;7(1).
56. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and applications. BMC Bioinformatics. 2009;10(1):421.
57. Janssen S, McDonald D, Gonzalez A, Navas-Molina JA, Jiang L, Xu ZZ, et al. Phylogenetic placement of exact amplicon sequences improves associations with clinical information. mSystems. 2018;3(3).
58. Abarenkov K, Nilsson RH, Larsson K-H, Taylor Andy FS, May Tom W, Frøslev TG, et al. The UNITE database for molecular identification and taxonomic communication of fungi and other eukaryotes: sequences, taxa and classifications reconsidered. Nucleic Acids Research. 2023;52(D1):D791-D7.
59. Therneau TM. A Package for Survival Analysis in R. R package version 3.5-8. Modeling Survival Data: Extending the Cox Model. 2024.
60. Kassambara A, Kosinski M, Biecek P, Fabian S. Drawing Survival Curves using 'ggplot2. Package 'survminer' version 0.4.9. 2021.
61. McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. Plos one. 2013;8(4):e61217.
62. Dinno A. Dunn's Test of Multiple Comparisons Using Rank Sums. Package 'dunn.test' Version1.3.6. 2024.
63. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016.
64. Martinez Arbizu P. Pairwise Multilevel Comparison using Adonis. Package: pairwiseAdonis version 0.4.1. 2017.
65. Ramalho MO, Bueno OC, Moreau CS. Species-specific signatures of the microbiome from *Camponotus* and *Colobopsis* ants across developmental stages. PLOS ONE. 2017;12(11).
66. Li Z, Alves SB, Roberts DW, Fan M, Delalibera Jr I, Tang J, et al. Biological control of insects in Brazil and China: history, current programs and reasons for their successes using entomopathogenic fungi. Biocontrol Science and Technology. 2010;20(2):117-36.
67. Luz C, Netto MC, Rocha LF. In vitro susceptibility to fungicides by invertebrate-pathogenic and saprobic fungi. Mycopathologia. 2007;164(1):39-47.

68. Višňovská D, Pyszko P, Šigut M, Kostovčík M, Kolařík M, Kotásková N, et al. Caterpillar gut and host plant phylloplane mycobiomes differ: a new perspective on fungal involvement in insect guts. FEMS Microbiol Ecol. 2020;96(9).

69. Matsuura Y, Moriyama M, Łukasik P, Vanderpool D, Tanahashi M, Meng XY, et al. Recurrent symbiont recruitment from fungal parasites in cicadas. Proc Natl Acad Sci U S A. 2018;115(26):E5970-e9.

70. Zhong X, Peng Q-y, Li S-S, Chen H, Sun H-X, Zhang G-R, et al. Detection of *Ophiocordyceps sinensis* in the roots of plants in alpine meadows by nested-touchdown polymerase chain reaction. Fungal Biology. 2014;118(4):359-63.

71. Meurville MP, LeBoeuf AC. Trophallaxis: the functions and evolution of social fluid exchange in ant colonies (Hymenoptera: Formicidae). Myrmecological News. 2021;31:1-30.

72. Krautkramer KA, Fan J, Bäckhed F. Gut microbial metabolites as multi-kingdom intermediates. Nature Reviews Microbiology. 2021;19(2):77-94.

73. Aleti G, Troyer EA, Hong S. G protein-coupled receptors: A target for microbial metabolites and a mechanistic link to microbiome-immune-brain interactions. Brain, Behavior, & Immunity - Health. 2023;32:100671.

74. Cohen LJ, Esterhazy D, Kim S-H, Lemetre C, Aguilar RR, Gordon EA, et al. Commensal bacteria make GPCR ligands that mimic human signalling molecules. Nature. 2017;549(7670):48-53.

75. Raymann K, Moran NA. The role of the gut microbiome in health and disease of adult honey bee workers. Curr Opin Insect Sci. 2018;26:97-104.

76. Loreto RG, Hughes DP. The metabolic alteration and apparent preservation of the zombie ant brain. Journal of Insect Physiology. 2019;118:103918.