Endoplasmic Reticulum contact sites facilitate the coordinated division of *Salmonella*-

containing vacuole (SCV)

Umesh Chopra[1], Priyanka Bhansali[1], Subba Rao Gangi Setty[1], Dipshikha Chakravortty[1,2]

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, Karnataka

560012, India

[2]School of Biology, Indian Institute of Science Education and Research Thiruvananthapuram,

Maruthamala PO, Vithura, Thiruvananthapuram - 695551, Kerala, India

Corresponding author: Prof. Dipshikha Chakravortty

Email: dipa@iisc.ac.in

Abstract

Salmonella Typhimurium (STM) resides in a membrane-bound compartment called *Salmonella*

Containing Vacuole (SCV) in several infected cell types. Within host cells, the division of bacteria

and SCV are synchronous to maintain the single bacterium per vacuole. However, the mechanism

regulating the synchronous fission and the machinery is not well understood. The fission of several

intracellular organelles is regulated by the dynamic nature of the tubular endoplasmic reticulum

(ER). In this study, we have evaluated the role of ER in controlling SCV fission. Interestingly,

Salmonella-infected cells show the activation of unfolded protein response (UPR) with expanded

ER tubules compared to the uninfected cells. Further, changing the ratio/expression of ER

morphology regulators, such as reticulon-4a (Rtn4a) and CLIMP63, affected bacterial proliferation

significantly, suggesting a potential role for tubular ER in facilitating the SCV division. Live-cell

imaging analysis shows the marking of tubular ER precisely at the center of the majority of SCV division (78%) sites. We have investigated the role of SteA, a known *Salmonella* effector, in modulating the membrane dynamics and coordinating the SCV division. We observed that SteA resides on the SCV membranes and helps in making membrane contact sites between SCV and ER. Accordingly, the colocalization of ER with SCV enclosing steA mutant *Salmonella* was significantly reduced compared to SCV-formed by wild-type *Salmonella*. Depletion of *steA* in *Salmonella* resulted in profound defects in SCV division, resulting in multiple bacteria residing in a single vacuole with defects in proliferation compared to the wild-type strain in epithelial cells. However, during infection in mice, the STM*ΔsteA* mutant did not show any defect in colonization but affected the survival rate of mice. Overall, this study suggests a coordinated role of bacterial effectors in promoting the ER contact sites with SCVs and thus regulating the successful division of pathogenic vacuole.

Introduction

Salmonella enterica subsp. *enterica* serovar Typhimurium (STM) is an intracellular pathogen known to infect a wide range of hosts and result in diseases across species, including gastroenteritis to systemic infections[1,2]. *S.* Typhimurium enters the body through the fecal-oral route following the bacteria utilize an arsenal of strategies to overcome several stresses like the acidic pH of the stomach, gut flora, bile salts, etc., and finally reach the small intestinal epithelia[3]. *Salmonella* secretes several virulence proteins inside the cell which lead to actin rearrangement and membrane ruffling, that helps in the invasion of bacteria inside the host epithelial cell[4–7]. *Salmonella* effector sopD is known to facilitate the recruitment of dynamin2 (Dyn2) at the invasion site, which helps in the endocytosis of bacteria inside host cells[8].

Upon invasion into the host cell, *Salmonella* resides in *Salmonella* containing Vacuole (SCV), a membrane-enclosed compartment. Mature SCV shares several characteristics markers of lysosomes/late endosomes such as LAMP1 and Rab7 but is negative for acid hydrolases and cathepsins[9]. Inside the epithelial cells, several protruding structures from SCV known as *Salmonella*-induced filaments (SIFs)[10,11] are observed which as LAMP1 positive. The membrane composition of SCV is not very well understood but it contains molecules and proteins from several host organelles and are enriched with several bacterial effector proteins[12,13]. Our group as well as several others have previously shown that *Salmonella* resides as a single bacterium per vacuole state inside the host, suggesting that SCV membrane divides synchronously with the division of *Salmonella* but the mechanism behind SCV division is not known[14–17]. For successful SCV division, expansion of SCV membrane, pinching of SCV membrane precisely with bacterial

division, and lastly, scission of host-derived SCV membrane must be followed in an orchestrated

manner.

Several literatures have highlighted the importance of contact sites between several organelles

such as ER with mitochondria, plasma membrane and endosomes, and mitochondria with

lysosomes[18–21]. Contact sites between ER-mitochondria and ER endosomes are very widely

studied in the aspect of organelle division[18,22]. In the case of mitochondria that comprises of an

inner and an outer membrane, and both membranes divide simultaneously during its fission. The

inner mitochondrial membrane uses its machinery for fission while outer membrane constriction

is initiated by the ER followed by the recruitment of Drp1, which mediates the fission of

mitochondria[18,23,24]. Bacterial pathogens like *Chlamydia* and *Brucella* make contact sites with

cellular organelles to facilitate their proliferation inside the host cell[25–27]. Interestingly, the SCV

membrane also comprises of several ER proteins, and recently membrane contact sites between

SCV and ER have also been observed [28]. To understand the mechanism behind the SCV division,

we reasoned that SCV is partially like mitochondria as far as the origin of the membrane is

concerned. SCV and mitochondria (being an endosymbiont) encompass an outer membrane

derived from the eukaryotic host while the inner membrane of mitochondria is prokaryotic in

nature like the cell wall of *Salmonella*. To understand the same, we hypothesized that a process

similar to mitochondria could lead to SCV fission. Our results suggest that *Salmonella* infection

leads to activation of UPR, ER expansion, and an increase in tubulation. We have observed ER

tubules mark at the fission site of SCV division which possibly initiate the constriction. *Salmonella*

translocated effector steA localizes onto the SCV membrane and possibly helps in making contact

with ER, suggesting that steA can act as a factor to recruit the ER onto SCV and thus facilitate the

SCV division. STM*ΔsteA* resides as multiple bacteria in one big vacuole with defects in

proliferation. In the mice model, STM*ΔsteA* doesn't show any defect in colonization in the

secondary site of infection, however, there is a delay in the onset of mice death upon infection as

compared to STM WT. Overall, our study unravels for the first time, the coordinated role of ER

and bacterial effector steA in bacterial proliferation inside the cell.

Materials and Methods

Bacterial strains and culture conditions

Salmonella enterica subspecies *enterica* serovar Typhimurium (STM WT) wild-type strain ATCC 14028s was used in all experiments. The bacterial strain was cultured in Luria broth (LB-Himedia) with constant shaking (170 rpm) at 37°C orbital-shaker. Antibiotics like kanamycin (50 µg/ml), and ampicillin (50µg/ml) were used wherever required. A one-step gene inactivation method was used for the generation of knockout strains[29]. Wild type as well mutant strains were transformed with pFPV-m-cherry or pFPV-GFP plasmid for immunofluorescence assays.

Table:1 List of bacterial strains used in this study

Strain	Characteristic (Antibiotic resistance)	Source
STM WT 14028s	-	Kind gift by Prof. M. Hensel
STM*ΔssaV*	Kan^R	Laboratory stock
STM*ΔsteA*	Kan^R	This study
STM*ΔsteA:steA*	Kan^R +Amp^R	This study

Generation of Complement strain and steA-eGFP plasmid

Low copy number plasmid pQE-60 was used to prepare the steA-HA complement strain. Briefly, PCR amplification (using Thermo Q5 DNA polymerase) of the *steA* gene was performed using Wild Type STM colony with respective PCR primers. Forward cloning primer was chosen from a position of 250 base pairs upstream of the gene start site, to amplify the native promoter of the gene during PCR and to maintain native expression level of protein and sequence of HA tag was added in reverse primer. The amplified PCR product was purified using a Gel extraction kit (Qiagen). The PCR product and Plasmid were subjected to restriction digestion by using Xho1 (NEB) and Hind III (NEB) at 37°C for 2 hours. Double digested PCR product and plasmid were further subjected to Ligation using T4 DNA ligase (NEB) at 16°C for 12-14 hours. The ligated product was further electroporated in STMΔ*steA* strain to prepare the complemented strain i.e. STMΔ*steA*:pQE60-*steA* (hence referred to as STMΔ*steA:steA*). The expression level of complement was confirmed with western blotting and confirmed with DNA sequencing.

For preparation of steA-eGFP plasmid for transfection study, eGFP-N3 empty vector was used. The sequence of steA was amplified via PCR from STM WT using primer with restriction site for Xho1 and BamH1. The entire process for cloning was performed as discussed above.

Eukaryotic Cell Culture, Transfection and Plasmids

HeLa cells and RAW 264.7 murine macrophages were cultured in DMEM (Lonza) containing 10% FBS (Gibco) at a humified incubator maintaining a temperature of 37°C and 5% CO_2. Cells were

seeded in 6 wells, 12 wells, or 24 well plates at a confluency of 60-70% prior to infection. Cells

were maintained in the presence of 1% penstrep. Cells were given a PBS wash and supplemented

with fresh media without penstrep at least 6 hours prior to infection. Plasmid overexpression or

knockdown via shRNA was done using Lipofectamine 3000 using manufacturer protocol. Briefly,

300ng of plasmid was mixed in opti-MEM media for 5 min while 1ul of L3000 was mixed (for

one well of 24 well plate) in opti-MEM for 5 min separately and further mixed and incubated for

20 min at room temperature. Post incubation, this mixture was added to cells for 6 hours, for

transfection to take place. Cells were washed with PBS and supplemented with complete media

post 6 hours of treatment, and post 48h of transfection, infection was given. Overexpression

constructs such as: pHAGE2 mCherry-Rtn4a was a kind gift from Prof. Tom Rapoport (Addgene

plasmid 86683)[30] mCh Climp-63 and mCh sec61β was a kind gift from Prof. Gia Voeltz (Addgene

plasmid 136293, Addgene 49155)[31,32], RFP KDEL construct was a kind gift from Prof. Nagaraj

Balasubramanian[33], GFP-LAMP1 was provided by Prof. Mahak Sharma[34]. The shRNA plasmids

were obtained from the human genome-wide TRC shRNA library (purchased from Sigma-Aldrich,

USA, Catalog no. SH0411), available at shRNA resource centre, MCB, IISc. pLKO.1-puro non-

mammalian shRNA (SHC002, shControl) was used as a control in all the shRNA mediated

knockdown experiments. Plasmid sequences and details are added in supplementary table 3.

Protocol for infection (Gentamicin Protection Assay)

The cells were seeded at the required confluency and infected with *Salmonella* strains with actively

growing log phase culture (for epithelial cells) and stationary phase culture (for RAW 264.7

macrophages) at Multiplicity of infection (MOI) 10 for Intra Cellular Survival Assay (ICSA) and

confocal imaging. For the live cell imaging experiment, an MOI of 25 was used to increase the

event of finding infected cells for imaging. For confocal microscopy studies, cells were seeded on

sterile glass coverslips at least 24 hours before infection. Upon infecting the cells at the required

MOI, the plate was centrifuged at 600 rpm for 5 min to facilitate the adhesion, and then the plate

was incubated for 25 min at 37°C and 5% CO_2. Post incubation, bacteria-containing media was

removed, and cells were given 2 PBS washes to remove any extracellular bacteria. Fresh media

containing 100µg/ml of gentamicin was added into wells and incubated at 37°C for one hour.

Following this, cells were again given 2 PBS washes, and fresh media containing 25µg/ml of

gentamicin was added and this concentration was maintained till the end point of the experiment.

Cells were harvested at required time points such as 2h and 16h for ICSA, 2h, 6h, and 16h for

confocal microscopy-based experiments.

Confocal Microscopy

After an appropriate hour post-infection, cells were washed with PBS twice and fixed with 3.5%

PFA at room temperature for 15 min. After fixation, coverslips were washed with PBS, and cells

were stained with primary antibody prepared in 2% BSA in PBS carrying 0.01% of saponin (for

membrane permeabilization) for 1 hour at room temperature or 4°C overnight in a cold room as

required. Post incubation, coverslips were washed again twice with PBS and incubated with

specific secondary fluorophore-labeled antibodies and incubated for one hour at room temperature.

Coverslips were then mounted on clean glass coverslips using mounting media. Once the mounting

media was dried, coverslips were stabilized by sealing the periphery with transparent nail paint on

the sides. Images were acquired on a confocal scanning laser microscope (Zeiss 880 multiphoton

microscope or Leica sp8 falcon system with or without zoom as required). Zen black software was

used for all colocalization-based quantification and to calculate mean fluorescence intensity (MFI).

Calculations of CTCF, percent of ER tubules, SCV diameter, and Line scans were prepared using

ImageJ. Live cell microscopy was performed on Leica SP8 Falcon microscope and live cell video

was processed and prepared using LAS X software. Antibodies used in this study is mentioned in

supplementary table 2.

Bacterial enumeration after infection for Intracellular survival assay

After the appropriate hour of post-infection, cells were washed with PBS at RT and lysed using

0.1% Triton-X 100. Cell lysate was plated on SS agar to obtain colony forming unit (CFU) and it

was used for the calculation of Percent Invasion and Fold proliferation using this formula.

Percent Invasion: (CFU obtained at 2 hours post infection/ CFU obtained from pre-inoculum) *100

Fold proliferation: CFU obtained at 16-hour post-infection/CFU obtained at 2 hours PI

RNA isolation and Real-Time PCR (qRT-PCR)

RNA isolation was performed using TRIzol (Takara) following the manufacturer's protocol.

Quantification of RNA was done using nano-drop (Thermo-Fisher Scientific) and RNA samples

were run on 1% agarose gel to check the quality of RNA. 2 µg of RNA was subjected to DNase I

(Thermo Fischer Scientific) treatment at 37°C for 1 hr followed by addition of 0.5M EDTA (final

concentration 5mM) and heat inactivation at 65°C for 10 mins to inactivate the enzyme. Further

cDNA was prepared by using a cDNA synthesis kit (Takara) as per the manufacturer's instructions.

The expression profile of all genes of interest was evaluated using RT primers by using SYBR

green mix (Takara) in a Biorad Real-Time PCR machine. The expression value of the target gene

was normalized to housekeeping internal control β-actin and further compared with uninfected

cells. Primers used in this study are mentioned in supplementary Table 1.

***In vivo* animal experiment**

6-8 weeks old C57BL/6 female mice were infected by orally gavaging 10^7 CFU of STM WT,

STMΔ*steA* and STMΔ*steA:steA*. 5 days post-infection, mice were euthanized and organs such as liver, spleen, MLN, and blood were harvested to study the colonization in organs. To enumerate CFU,

organs were homogenized and further plated on *Salmonella-Shigella* agar. Obtained CFU values were normalized with the gram weight of the organ and then converted into a log scale. Blood was collected through heart puncture and plated on SS agar. For the mice survival assay, 10^8 CFU was administered to each mouse via oral gavage. Mice were monitored every day till the time none of the mice in the cohort survived. The animal experiments were carried out in accordance with the approved guidelines of the institutional animal ethics committee at the Indian Institute of Science, Bangalore, India (Registration No: 48/GO/ReRcBiBt-S/Re-L/99/CPCSEA). All procedures involving the use of animals were performed according to the Institutional Animal Ethics Committee (IAEC)-approved protocol, ethics number: CAF/Ethics/853/2021.

Statistical analysis:

Statistical analysis was performed using GraphPad prism software. For analysis, the student's t-test (unpaired parametric test) was performed. The results are indicative of mean± SD or mean± SEM. For animal experiments, the Mann-Whitney test was performed. Group size, experimental number, no of cells in each set, and P value of each experiment are described in figure legends.

Results

Salmonella infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum

Several bacterial and viral infections are known to activate UPR for their survival[35,36]. UPR helps to alleviate the ER stress by initiating several transcriptional and translational regulations which lead to the maintenance of ER homeostasis[37,38]. Inside the cell, UPR signaling is maintained by transmembrane proteins located in the ER membrane and their catalytic domain towards the cytosol. Inositol requiring enzyme-1 (IRE-1), Protein kinase c like ER kinase (PERK), and Activating transcription factor-6 (ATF-6) comprises the main arm of UPR. Recent reports have shown the contact of SCV with ER, also the presence of ER proteins in the membrane fraction of SCV suggests the crosstalk between the two[28,12]. Here, we investigated how _Salmonella_ infection modulates the ER dynamics. To understand the activation of UPR by any of these pathways, we used confocal microscopy to look at the downstream transcription factors such as XBP1, ATF4, and ATF6 respectively inside the nucleus upon infection. SPI2 mutants of _Salmonella_, STM_ΔssaV_ (which is unable to secret SPI-2 encoded effector protein in host cytosol) was kept as a control to understand if activation of UPR is mediated by bacterial effectors or dependent on bacterial invasion. Our results suggest that from as early as post 2h of _Salmonella_ infection, both the transcription factors, ATF4 and ATF6 in the cytosol translocated to the nucleus of infected cells as compared to uninfected cells, and STM_ΔssaV_ infected cells, suggesting that only wild type

Salmonella can activate UPR (fig 1A-1D). *Salmonella* induced ATF4 and ATF6 signaling was

retained till   16h post-infection (fig S1A-S1D). However, we didn't observe XBP1 recruitment

inside the nucleus (fig 1E,1F) upon infection at any time point post-infection (fig S1E, S1F)

suggesting that *Salmonella* infection upregulates UPR by PERK and ATF6 pathways. To alleviate

the ER stress, activation of UPR leads to the expansion of the ER, protein folding capacity, and

increased phospholipid production[40]. Increased bacterial proliferation requires a continuous source

of membrane for the increasing number of SCV and increased phospholipid production upon UPR

could facilitate the SCV proliferation inside the host cell. Therefore, we sought to determine if

Salmonella infection led to ER expansion during the pathogenesis phase. Our transcriptional

analysis suggests that upon infection, there is upregulation at the transcript level of ER shaping

proteins such as Climp63 (responsible for maintaining ER sheets) and Reticulon-4a (Rtn4a,

essential for maintaining ER tubules) (fig 1G, fig 1H). The upregulation was significantly observed

at the late time point of infection, i.e., 16h post-infection, suggesting that active proliferation of

Salmonella and not just invasion (as evident at 2h and 6h post-infection) leads to ER expansion.

Since an increase in the transcript levels of ER shaping proteins were observed at the later time of

infection, we tried to quantify the percentage of ER sheets and tubules using ER marker, RFP-

KDEL upon infection as a measure to understand ER expansion upon infection. Upon analysis of

microscopy images, we observed that *Salmonella*-infected cells showed an increased percentage

of ER tubules compared to uninfected cells (fig1I, fig1J). The changes in percent ER tubules were

only evident in infected cells as compared to un-infected and bystander cells (neighboring cells

which are not infected), suggesting that stabilized SCV and bacterial effectors are responsible for

this phenotype (fig S1G and S1H). Together, this data indicates that *Salmonella* infection leads to

changes in ER morphology.

Expansion of ER tubules facilitates SCV proliferation and its division

Dynamic ER tubules play an essential role in forming membrane contact sites with several organelles [41,42]. Fig1 indicates that actively proliferating bacteria increased tubular ER and upregulated the expression of Rtn4a. We also tried to understand if there is an advantage of the increased tubules to bacterial proliferation. To identify the role of ER tubules in mediating bacterial proliferation, we transiently overexpressed reticulon 4a (Rtn4a) in HeLa cells, which helps to increase ER tubules[43], and then quantified the proliferation of bacteria by intracellular survival assay (ICSA). As compared to empty vector and general ER marker sec61β-mCh control, increased ER tubulation by Rtn4a led to the higher proliferation of *Salmonella* with no significant change in percent invasion, suggesting an increase in ER tubules helps in bacterial proliferation (fig2A, figS2A). To validate if this change in proliferation was dependent solely on the function of Rtn4a, knockdown of Rtn4a via shRNA was performed (figS2B). Knockdown of Rtn4a led to reduced proliferation of bacteria as compared to scrambled control hence strengthening the crucial role of ER tubules in bacterial proliferation (fig2B). Climp63 overexpression led to an increase in ER sheets as opposed to tubules. We quantified the number of SCV per cell upon transiently overexpressing the Climp63 in HeLa cells. Our data indicated that the number of SCV present per cell decreases upon overexpressing ER sheets using Climp63 compared to controls (fig2C, fig2D). Division of *Salmonella* is linked with the fission of SCV membrane [44]. Since the intricate balance of ER sheets and tubules affects bacterial proliferation, we hypothesized that ER tubules might facilitate SCV fission. In the case of mitochondria, membrane-less organelle, and endosomes, ER marks the division site, and later fission is completed by Drp1 (in mitochondria) and coronin1c and TMCC 1 complex (in endosome)[18,45–47]. To investigate this, we performed live cell imaging in cells transfected by RFP-KDEL (to mark the ER) followed by infection with *Salmonella*.

Quantification of data suggests that ER tubules mark the division site precisely at the center of

dividing SCV in 78% of events (fig2E-G). STM WT infected RAW macrophage also showed

contact of ER on the dividing bacteria (S2C).This was also evident in fixed cell images where we

could observe Lamp1 enclosed bacterial division in SCV marked by ER tubules exactly in the

center in 56.7±6.61% of fission events, however, in 14±9.07% of events, ER was seen in proximity

to the center and hence named adjacent while 29.2±8.4% of dividing SCV did not possess ER

tubules on them (fig2H, fig2I). In the majority of STM WT-infected cells, SCV was observed near

the periphery of the cell, suggesting mature SCV migrates towards the cell periphery to make

contacts with ER tubules to facilitate its division (S2D).This data suggests the importance of ER

tubules in marking the fission site of SCV. Taken together this data suggests that ER tubules help

in the proliferation and division of SCV inside the cell and help in maintaining SCV number within

the cell.

Salmonella **translocated effector steA is crucial for maintaining SCV contact site with ER**

resulting in a single bacterium per vacuole

SCV division leads to a single bacterium per vacuole throughout the course of pathogenesis in cell

lines or in mouse liver model[14]. Domingues et al suggested the role of steA in maintaining the

membrane dynamics of SCV[48]. In accordance with the same, we also observed that steA mutant

strain of *Salmonella* (STM*ΔsteA*) resides as multiple bacteria in one single vacuole (fig 3A). Our

data suggests that compared to STM WT, 30±3% of mutant bacteria reside in one big vacuole,

suggesting that loss of function of steA leads to improper SCV fission (fig 3B). Live cell imaging

shows several points of contacts between SCV and ER, suggesting that SCV actively makes

membrane contact sites with ER for its survival and proliferation (fig2E, fig3C). SteA resides onto

the PI4P enriched SCV membrane (fig3C), and hence we asked are membrane contacts between

ER and SCV maintained by steA? Microscopy images show that steA-enriched SCV makes

contact with ER (fig3C). Quantification of the data suggests that ER colocalizes more with the

SCV of STM WT than the SCV formed by STM*ΔsteA*, suggesting that steA helps in making

contact sites of SCV with ER (fig3D, fig3E). This was also observed upon transfecting HeLa cells

with steA-eGFP construct. SteA-eGFP showed higher colocalization with ER marker RFP-KDEL

as compared to eGFP empty vector control, strengthening the fact that steA possibly associate with

ER (figS3A,S3B). This data suggests that steA might help in making contact sites with ER and

further recruiting ER onto the SCV. These contact sites can be used as a stealthy approach by

Salmonella to recruit membrane from ER, and along with other *Salmonella* effector proteins,

facilitating SCV fission.

STM*ΔsteA* shows defect in proliferation inside epithelial cells and pathogenicity in mice in

vivo model

Previous reports from our lab suggested that *Salmonella* utilizes the strategy of SCV division to

increase the number of SCV per cell, so it becomes difficult for the cellular system to eradicate the

growing SCV[14]. Since steA mutants form a big vacuole inside the cell, we asked how the mutant

strain proliferates in the cell. As compared to STM WT and steA complemented strain, STM*ΔsteA*

proliferated significantly less in the HeLa cell line while there was no difference in invasion rate

of bacteria inside cell (fig4A, figS4A). Less proliferation of mutant strain suggests that those bulky

vacuoles might be targeted by cellular machinery for degradation. Quantification of vacuole size

suggests that the diameter of SCV formed by STM*ΔsteA* is around $6.9\pm3.13\mu m$ whereas SCV of

STM WT is 0.4±0.11µm (fig4B, fig4C). Using LC3B as a marker to detect the extent of autophagy,

we observed that LC3B colocalized more with SCV of steA mutants as compared to SCV of STM

WT at 16h post-infection (fig4D,fig4E). Actively proliferating *Salmonella* also reduces the

number of lysosomes inside the cell to escape its fusion with acidic lysosomes[14.] In STM*ΔsteA*

infected cells, we observed a higher intensity of Lamp1 similar to uninfected cells, suggesting that

STM*ΔsteA* is unable to change the lysosomal dynamics as opposed to STM WT. Higher flux of

lysosomal proteins along with the recruitment of LC3B led to less proliferation of the STM*ΔsteA*

inside the cell (figS4B, figS4C). Next, we were interested in understanding the pathogenicity of

STM*ΔsteA* in an in vivo model. We orally gavaged C57BL/6 mice with 10^7 CFU of STM WT,

STM*ΔsteA*, and STM*ΔsteA:steA,* and 5 days post-infection, mice were euthanized, and organs such

as liver, spleen, and mesenteric lymph nodes were harvested since they act as a secondary site of

infection (fig4F-4I). We observed no significant change in the colonization of bacteria in these

organs, suggesting STM*ΔsteA* doesn't lose its capability to cross the gut barrier and colonizing and

dissemination in distal organs and blood (figS4D). However, upon performing a survival assay of

mice, we noticed the delayed onset of mice death in the STM*ΔsteA* infected cohort (fig4J), where

STM WT infected mice started succumbing to infection at 5^{th} day post infection, STM*ΔsteA*

infected cohort started succumbing to infection at 7^{th} day post infection. Taken together, this data

suggests that if the fission of SCV is altered, *Salmonella* resides as multiple bacteria in one big

vacuole and proliferates less. Those big/bulky vacuoles are further targeted by autophagic

machinery and cleared by active lysosomes in the cell culture system. However, in an in-vivo mice

model, colonization of bacterium is not hampered, but it affects the survival rate of mice,

suggesting that SteA acts as an essential effector for pathogenicity in mice.

Discussion:

Bacterial pathogens invade several cell types and reside in endocytic or phagocytic vacuoles[49]. Cell uses their extraordinary machinery to clear the pathogens, but by using several effector proteins, bacterial pathogens modulate cellular machinery and prolong their survival. A few strategies that several bacterial pathogens utilize are: modification of the membrane of the pathogen-containing vacuole, blocking the activation of xenophagy, and escaping its fusion with acidic lysosome[50–56]. Bacterial pathogens such as *Legionella, Brucella, Chlamydia*, and *Salmonella* are known to reside in vacuoles, but the morphology and membrane source of the vacuole are quite different. *Legionella* and *Chlamydia* reside in a spacious vacuole that comprises multiple bacteria in a single vacuole[57], contrary *Salmonella* resides in a tight-fitting vacuole as a single bacterium per vacuole[14]. The biogenesis of the SCV membrane is not very well known but proteome analysis of SCV suggests that the SCV membrane is highly modified and contain proteins from the lysosome, Golgi, and ER[12]. Several effectors of *Salmonella* help in recruiting membrane components, inhibiting the transport of M6PR, and making contact sites with organelles for its proliferation [58–60]. Our results suggest that *Salmonella* infection induces ER stress inside the cell and activates the UPR. A few ways by which UPR helps in alleviating stress are 1) by increasing the protein folding capacity of cells and 2) by expanding the ER. Our data suggests that wild-type STM infection leads to UPR activation while STM*ΔssaV* is unable to activate the UPR arm. This suggests that bacterial effectors are crucial for initiating ER stress, as it was evident as early as 2h post-infection. Our studies support the previous findings, which suggest *Salmonella*-induced ER stress is crucial for bacterial proliferation[61–63]. Further, UPR activation leads to an increase in phospholipid production[40], which could be useful for the bacteria as growing SCV requires a continuous source of membrane. *Salmonella*-infected cells show an increase in

expression of *Reticulon-4a* and *Climp63* with an increase in tubular ER, suggesting this is an active

strategy employed by bacteria for its survival. Exogenously overexpression of Rtn4a provides an

advantage in the proliferation of *Salmonella* however, reduction in ER tubules by knockdown of

Rtn4a or overexpression of ER sheets using Climp63 adversely affects the bacterial proliferation

and SCV number inside the cell, indicating ER tubules plays a crucial role in bacterial

proliferation. The intricate balance of ER sheets and tubules has been recently shown to affect the

dynamics of membrane-less organelle[42]. Our study also highlights the importance of ER in the

successful proliferation of *Salmonella*. Our live cell data showed the recruitment of ER tubules at

the center of dividing SCV. This was evident in HeLa as well as RAW macrophage cell lines,

suggesting that ER tubule recruitment marks the division site of SCV. ER is very well known for

marking the division site of organelles such as mitochondria, membrane-less organelle, and

endosomes[18,42,64]. Our study suggests that *Salmonella* hijacks the strategy of ER recruitment for

its survival and SCV division. We have identified the role of bacterial effector SteA in maintaining

the contact of SCV with ER in the formation of possible membrane contact sites. Earlier studies

have shown the presence of steA on SCV and its colocalization with Golgi [65–67]. In our study, we

have partially observed a similar phenotype and have observed a new role of steA in interaction

with ER. Inside the infected cell, the expression of steA is highly regulated as it can be mostly seen

around the bacteria onto SCV. To investigate the role of steA in making contact sites, we looked

for the colocalization of ER with SCV of STM WT and STM*ΔsteA* and observed the difference in

colocalization of ER with SCV, highlighting the importance of *Salmonella* effector steA in

mediating contact site between SCV and ER. Bacterial pathogens such as *Chlamydia* use its

effector protein for forming contacts of its inclusion bodies with ER [25] and also it targets host

cellular protein to facilitate this interaction [68]. *Salmonella* effectors sseJ and sseL have previously

been reported to recruit OSBP1 onto SCV which helps in making contact with VapA and VapB

(conserved ER proteins known for tethering of ER with cellular organelles) and maintain SCV

vacuolar integrity [69]. Our data suggests the potential role of steA in forming contact sites of SCV

with ER as a strategy to facilitate its proliferation.

The loss of function of steA led to a bulky vacuole carrying multiple bacteria inside and with

defects in proliferation. Evidence from McQuate et al., also suggests the defect in the proliferation

of STMΔsteA in epithelial as well as macrophages [66]. To understand the difference in proliferation

of STMΔsteA compared to STM WT, we have observed SCV formed by STMΔsteA colocalizes

more with autophagy marker LC3B, suggesting big bulky vacuoles are cleared by the cellular

system and hence lead to less proliferation inside the cell. STM WT also actively reduces lysosome

number inside the cell as an active strategy to block its fusion with the acidic lysosome [14]. But in

the case of STMΔsteA infected cell, we observed a higher mean intensity of LAMP1 which was

similar to intensity observed in uninfected cells suggesting that somehow STMΔsteA could not

modulate lysosomal dynamics and as a result, active lysosome could hinder the bacterial

proliferation in cell line model. Similarly, Liu et al.,2021 have highlighted the importance of steA

in bacterial proliferation inside cells by impairing the fusion of SCV with lysosomes[70]. Taken

together this data suggests the importance of steA in bacterial proliferation inside infected cell.

However, to our surprise, we did not observe any defect in the in vivo colonization of mouse organs

such as the liver, spleen, MLN, and blood. Survival analysis of mice suggests late onset of mice

death in STMΔsteA infected cohort upon infection as compared to STM WT. We have also

observed less weight reduction in STMΔsteA infected mice as compared to STM WT and

STMΔsteA:steA, and this difference in weight reduction could benefit mice survival (figS4E).

However, Geddes et al, 2005 have reported the importance of steA for efficient colonization in

mouse spleen using the Balb/c mice model infecting via the intraperitoneal route [65]. The difference

in colonization from our study can be dependent on the route of infection wherein, our study uses

a conventional way of oral gavaging for *Salmonella* infection which is a natural route of infection

in contrast to intraperitoneal injection. Taken together, our study suggests the importance of steA

in bacterial pathogenicity and proliferation.

Our study provides new insights into how bacteria use a strategy of remodeling the ER dynamics

for its benefit. Domingues et al.,2014 and Chen et al.,2021 have highlighted the importance of

steA for the active division of SCV [48,71]. Our data supports their findings and suggests the possible

role of steA in recruiting the ER and facilitating SCV fission. Interestingly, where wild-type STM

can easily divide along with SCV, while loss of function of steA hampers 30% SCV division with

marked less colocalization with ER, suggesting there can be several other effector proteins that

might facilitate the contacts between ER and SCV. As reported by Chen et al.,2021 about the

coordinated role of steA, sopD2, and sifA in successful SCV division, our data explains the newly

identified role of steA in the process. Since these effectors have coordinated roles, it will be

interesting to understand how individual bacterial effectors are crucial in the complete fission of

SCV. Several host proteins have also been linked with SCV division as the absence of those host

proteins such as PLEKHM1 and ESCRT-3 leads to SCV carrying multiple bacteria [59,72]. Within the

cell, SCV acts as an autonomous organelle that has acquired a vacuolar membrane from the host

system which also synchronously divides along with bacterial division. The machinery regulating

the SCV division could be difficult to identify because these fission events are very transient in

nature. As per literature evidence, both host system and bacterial effectors could be equally

involved in this process to execute successful fission. The complete knowledge of fission

machinery can be used as a targeted approach to inhibit bacterial proliferation during infection.

Declaration of interest:

The authors declare no competing interests.

Acknowledgement:

This work was funded by the DAE SRC fellowship (DAE00195),DST (FIST), and UGC is sincerely acknowledged. DC acknowledges the ASTRA Chair professorship grant from IISc and TATA innovation fellowship grant. UC sincerely acknowledges IISc GATE fellowship for financial support.

References:

1. Haraga, A., Ohlson, M.B., and Miller, S.I. (2008). Salmonellae interplay with host cells. *6*, 53–66. 10.1038/nrmicro1788.

2. Brenner, F.W., Villar, R.G., Angulo, F.J., Tauxe, R., and Swaminathan, B. (2000). GUEST COMMENTARY Salmonella Nomenclature.

3. Pradhan, D., and Devi Negi, V. (2019). Stress-induced adaptations in Salmonella: A ground for shaping its pathogenesis. Preprint at Elsevier GmbH, 10.1016/j.micres.2019.126311 10.1016/j.micres.2019.126311.

4. Criss, A.K., and Casanova, J.E. (2003). Coordinate regulation of Salmonella enterica serovar typhimurium invasion of epithelial cells by the Arp2/3 complex and Rho GTPases. Infect Immun *71*, 2885–2891. 10.1128/IAI.71.5.2885-2891.2003.

5. Zhou, D., Chen, L.M., Hernandez, L., Shears, S.B., and Galán, J.E. (2001). A Salmonella inositol polyphosphatase acts in conjunction with other bacterial effectors to promote host cell actin cytoskeleton rearrangements and bacterial internalization. Mol Microbiol *39*, 248–260. 10.1046/j.1365-2958.2001.02230.x.

6. Bakshi, C.S., Singh, V.P., Wood, M.W., Jones, P.W., Wallis, T.S., and Galyov, E.E. (2000). Identification of SopE2, a Salmonella Secreted Protein Which Is Highly Homologous to SopE and Involved in Bacterial Invasion of Epithelial Cells.

7. Hardt, W.-D., Chen, L.-M., Schuebel, K.E., Bustelo, X.R., and Galá, J.E. (1998). S. typhimurium Encodes an Activator of Rho GTPases that Induces Membrane Ruffling and Nuclear Responses in Host Cells.

8. Boddy, K.C., Zhu, H., D'Costa, V.M., Xu, C., Beyrakhova, K., Cygler, M., Grinstein, S., Coyaud, E., Laurent, E.M.N., St-Germain, J., et al. (2021). Salmonella effector SopD promotes plasma membrane scission by inhibiting Rab10. Nat Commun *12*. 10.1038/s41467-021-24983-z.

9. Gorvel, J.-P., and Méresse, S. (2001). Maturation steps of the Salmonella-containing vacuole.

10. Drecktrah, D., Levine-Wilkinson, S., Dam, T., Winfree, S., Knodler, L.A., Schroer, T.A., and Steele-Mortimer, O. (2008). Dynamic behavior of salmonella-induced

membrane tubules in epithelial cells. Traffic *9*, 2117–2129. 10.1111/j.1600-
0854.2008.00830.x.

11.   Garcia-del Portillo, Fran., Zwick, M.B., Yin Leung, K., and Brett Finlay, B. (1993).
Salmonella induces the formation of filamentous structures containing lysosomal
membrane glycoproteins in epithelial cells.

12.   Santos, J.C., Duchateau, M., Fredlund, J., Weiner, A., Mallet, A., Schmitt, C.,
Matondo, M., Hourdel, V., Chamot-Rooke, J., and Enninga, J. (2015). The COPII
complex and lysosomal VAMP7 determine intracellular Salmonella localization and
growth. Cell Microbiol *17*, 1699–1720. 10.1111/cmi.12475.

13.   Reuter, T., Vorwerk, S., Liss, V., Chao, T.C., Hensel, M., and Hansmeier, N. (2020).
Proteomic Analysis of Salmonella-modified Membranes Reveals Adaptations to
Macrophage Hosts. Molecular and Cellular Proteomics *19*, 900–912.
10.1074/mcp.RA119.001841.

14.   Eswarappa, S.M., Negi, V.D., Chakraborty, S., Sagar, B.K.C., and Chakravortty, D.
(2010). Division of the Salmonella-containing vacuole and depletion of acidic
lysosomes in Salmonella-infected host cells are novel strategies of Salmonella
enterica to avoid lysosomes. Infect Immun *78*, 68–79. 10.1128/IAI.00668-09.

15.   Buchmeiert, N.A., and Heffront, F. (1991). Inhibition of Macrophage Phagosome-
Lysosome Fusion by Salmonella typhimurium.

16.   Salmonella maintains the integrity of its intracellular vacuole through the action of
SifA.pdf.

17.   Holden, D.W. (2002). Trafficking of the Salmonella vacuole in macrophages.
Preprint, 10.1034/j.1600-0854.2002.030301.x 10.1034/j.1600-0854.2002.030301.x.

18.   Friedman J.R., Lackner L.L., West M. DiBenedetto J.R., Nunnari J., and Voeltz
G.K. (2011). ER tubules mark site of mitochondrial division. Science
10.1126/science.1207385

19.   Fernández-Busnadiego, R., Saheki, Y., and De Camilli, P. (2015). Three-
dimensional architecture of extended synaptotagmin-mediated endoplasmic
reticulum-plasma membrane contact sites. Proc Natl Acad Sci U S A *112*, E2004–
E2013. 10.1073/pnas.1503191112.

20.   Wong, Y.C., Ysselstein, D., and Krainc, D. (2018). Mitochondria-lysosome contacts
regulate mitochondrial fission via RAB7 GTP hydrolysis. Nature *554*, 382–386.
10.1038/nature25486.

21.   Wang, H.-J., Guay, G., Pogan, L., Sauvé, R., and Nabi, I.R. (2000). JCB Report
504   Calcium Regulates the Association between Mitochondria and a Smooth
505   Subdomain of the Endoplasmic Reticulum.

506   22.   Rowland, A.A., Chitwood, P.J., Phillips, M.J., and Voeltz, G.K. (2014). ER contact
507   sites define the position and timing of endosome fission. Cell *159*, 1027–1041.
508   10.1016/j.cell.2014.10.023.

509   23.   Ji, W.K., Chakrabarti, R., Fan, X., Schoenfeld, L., Strack, S., and Higgs, H.N.
510   (2017). Receptor-mediated Drp1 oligomerization on endoplasmic reticulum.
511   Receptor-mediated Drp1 oligomerization on endoplasmic reticulum *216*, 190538.
512   10.1101/190538.

513   24.   Smirnova, E., Griparic, L., Shurland, D., and Bliek, A.M. Van Der (2001).
514   Dynamin-related Protein Drp1 Is Required for Mitochondrial Division in
515   Mammalian Cells. *12*, 2245–2256.

516   25.   Stanhope, R., Flora, E., Bayne, C., and Derré, I. (2017). IncV, a FFAT motif-
517   containing Chlamydia protein, tethers the endoplasmic reticulum to the pathogen-
518   containing vacuole. Proc Natl Acad Sci U S A *114*, 12039–12044.
519   10.1073/pnas.1709060114.

520   26.   Boncompain, G., Müller, C., Meas-Yedid, V., Schmitt-Kopplin, P., Lazarow, P.B.,
521   and Subtil, A. (2014). The intracellular bacteria Chlamydia hijack peroxisomes and
522   utilize their enzymatic capacity to produce bacteria-specific phospholipids. PLoS
523   One *9*. 10.1371/journal.pone.0086196.

524   27.   Sedzicki, J., Tschon, T., Low, S.H., Willemart, K., Goldie, K.N., Letesson, J.J.,
525   Stahlberg, H., and Dehio, C. (2018). 3D correlative electron microscopy reveals
526   continuity of Brucellacontaining vacuoles with the endoplasmic reticulum. J Cell
527   Sci *131*. 10.1242/jcs.210799.

528   28.   Li, M., Tripathi-Giesgen, I., Schulman, B.A., Baumeister, W., and Wilfling, F.
529   (2023). In situ snapshots along a mammalian selective autophagy pathway. Proc
530   Natl Acad Sci U S A *120*. 10.1073/pnas.2221712120.

531   29.   Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal
532   genes in Escherichia coli K-12 using PCR products.

533   30.   Wang, S., Tukachinsky, H., Romano, F.B., and Rapoport, T.A. (2016). Cooperation
534   of the ER-shaping proteins atlastin, lunapark, and reticulons to generate a tubular
535   membrane network. 10.7554/eLife.18605.001.

31. Shibata, Y., Voss, C., Rist, J.M., Hu, J., Rapoport, T.A., Prinz, W.A., and Voeltz, G.K. (2008). The reticulon and Dp1/Yop1p proteins form immobile oligomers in the tubular endoplasmic reticulum. Journal of Biological Chemistry *283*, 18892–18904. 10.1074/jbc.M800986200.

32. Zurek, N., Sparks, L., and Voeltz, G. (2011). Reticulon short hairpin transmembrane domains are used to shape ER tubules. Traffic *12*, 28–41. 10.1111/j.1600-0854.2010.01134.x.

33. Singh, V., Erady, C., and Balasubramanian, N. (2018). Cell-matrix adhesion controls Golgi organization and function through Arf1 activation in anchorage-dependent cells. J Cell Sci *131*. 10.1242/jcs.215855.

34. Marwaha, R., Arya, S.B., Jagga, D., Kaur, H., Tuli, A., and Sharma, M. (2017). The Rab7 effector PLE KHM1 binds Arl8b to promote cargo traffic to lysosomes. Journal of Cell Biology *216*, 1051–1070. 10.1083/jcb.201607085.

35. Alshareef, M.H., Hartland, E.L., and McCaffrey, K. (2021). Effectors targeting the unfolded protein response during intracellular bacterial infection. Preprint at MDPI AG, 10.3390/microorganisms9040705 10.3390/microorganisms9040705.

36. Cirone, M. (2021). Er stress, upr activation and the inflammatory response to viral infection. Viruses *13*. 10.3390/v13050798.

37. Yamamoto, K., Sato, T., Matsui, T., Sato, M., Okada, T., Yoshida, H., Harada, A., and Mori, K. (2007). Transcriptional Induction of Mammalian ER Quality Control Proteins Is Mediated by Single or Combined Action of ATF6α and XBP1. Dev Cell *13*, 365–376. 10.1016/j.devcel.2007.07.018.

38. Yoshida, H., Okada, T., Haze, K., Yanagi, H., Yura, T., Negishi, M., and Mori, K. (2000). ATF6 Activated by Proteolysis Binds in the Presence of NF-Y (CBF) Directly to the cis-Acting Element Responsible for the Mammalian Unfolded Protein Response.

39. Li, M., Tripathi-Giesgen, I., Schulman, B.A., Baumeister, W., and Wilfling, F. (2023). In situ snapshots along a mammalian selective autophagy pathway. Proc Natl Acad Sci U S A *120*. 10.1073/pnas.2221712120.

40. Sriburi, R., Jackowski, S., Mori, K., and Brewer, J.W. (2004). XBP1: A link between the unfolded protein response, lipid biosynthesis, and biogenesis of the endoplasmic reticulum. Journal of Cell Biology *167*, 35–41. 10.1083/jcb.200406136.

41. Wu, H., Carvalho, P., and Voeltz, G.K. (2018). Here, there, and everywhere: The
importance of ER membrane contact sites. Science (1979) *361*.
10.1126/science.aan5835.

42. Lee, J.E., Cathey, P.I., Wu, H., Parker, R., and Voeltz, G.K. (2020). Endoplasmic
reticulum contact sites regulate the dynamics of membraneless organelles. Science
(1979) *367*. 10.1126/science.aay7108.

43. Voeltz, G.K., Prinz, W.A., Shibata, Y., Rist, J.M., and Rapoport, T.A. (2006). A
Class of Membrane Proteins Shaping the Tubular Endoplasmic Reticulum. Cell
124, 573–586. 10.1016/J.CELL.2005.11.047.

44. Buchmeiert, N.A., and Heffront, F. (1991). Inhibition of Macrophage Phagosome-
Lysosome Fusion by Salmonella typhimurium.

45. Hoyer, M.J., Chitwood, P.J., Ebmeier, C.C., Striepen, J.F., Qi, R.Z., Old, W.M., and
Voeltz, G.K. (2018). A Novel Class of ER Membrane Proteins Regulates ER-
Associated Endosome Fission. Cell *175*, 254-265.e14. 10.1016/j.cell.2018.08.030.

46. Smirnova, E., Griparic, L., Shurland, D.-L., and Van Der Bliek, A.M. (2001).
Dynamin-related Protein Drp1 Is Required for Mitochondrial Division in
Mammalian Cells.

47. Ji, W.K., Hatch, A.L., Merrill, R.A., Strack, S., and Higgs, H.N. (2015). Actin
filaments target the oligomeric maturation of the dynamin GTPase Drp1 to
mitochondrial fission sites. Elife *4*, 1–25. 10.7554/eLife.11553.

48. Domingues, L., Holden, D.W., and Mota, L.J. (2014). The Salmonella effector SteA
contributes to the control of membrane dynamics of Salmonella-containing
vacuoles. Infect Immun *82*, 2923–2934. 10.1128/IAI.01385-13.

49. Case, E.D.R., Smith, J.A., Ficht, T.A., Samuel, J.E., and De Figueiredo, P. (2016).
Space: A Final Frontier for Vacuolar Pathogens. Preprint at Blackwell Munksgaard,
10.1111/tra.12382 10.1111/tra.12382.

50. Birmingham, C.L., Smith, A.C., Bakowski, M.A., Yoshimori, T., and Brumell, J.H.
(2006). Autophagy controls Salmonella infection in response to damage to the
Salmonella-containing vacuole. Journal of Biological Chemistry *281*, 11374–
11383. 10.1074/jbc.M509157200.

51. Tilney L.G., Harb O.S., Connelly P.S., Robinson C.G., and Roy C.R.(2001).How
the parasitic bacterium Legionella pneumophila modifies its phagosome and
transforms it into rough ER implications for conversion of plasma membrane to the
ER membrane. Journal of cell science 114, 4637-4650 (2001)

52. Kimmey, J.M., and Stallings, C.L. (2016). Bacterial Pathogens versus Autophagy: Implications for Therapeutic Interventions. Preprint at Elsevier Ltd, 10.1016/j.molmed.2016.10.008 10.1016/j.molmed.2016.10.008.

53. Pujol, C., Klein, K.A., Romanov, G.A., Palmer, L.E., Cirota, C., Zhao, Z., and Bliska, J.B. (2009). Yersinia pestis can reside in autophagosomes and avoid xenophagy in murine macrophages by preventing vacuole acidification. Infect Immun *77*, 2251–2261. 10.1128/IAI.00068-09.

54. Rzomp, K.A., Scholtes, L.D., Briggs, B.J., Whittaker, G.R., and Scidmore, M.A. (2003). Rab GTPases are recruited to chlamydial inclusions in both a species-dependent and species-independent manner. Infect Immun *71*, 5855–5870. 10.1128/IAI.71.10.5855-5870.2003.

55. Mesquita, F.S., Thomas, M., Sachse, M., Santos, A.J.M., Figueira, R., and Holden, D.W. (2012). The Salmonella deubiquitinase SseI inhibits selective autophagy of cytosolic aggregates. PLoS Pathog *8*. 10.1371/journal.ppat.1002743.

56. Zhou, L., Li, Y., Gao, S., Yuan, H., Zuo, L., Wu, C., Huang, R., and Wu, S. (2021). Salmonella spvC Gene Inhibits Autophagy of Host Cells and Suppresses NLRP3 as Well as NLRC4. Front Immunol *12*. 10.3389/fimmu.2021.639019.

57. Bastidas, R.J., Elwell, C.A., Engel, J.N., and Valdivia, R.H. (2013). Chlamydial intracellular survival strategies. Cold Spring Harb Perspect Med *3*. 10.1101/cshperspect.a010256.

58. McGourty, K., Thurston, T.L., Matthews, S.A., Pinaud, L., Mota, L.J., and Holden, D.W. (2012). Salmonella inhibits retrograde trafficking of mannose-6-phosphate receptors and lysosome function. Science (1979) *338*, 963–967. 10.1126/science.1227037.

59. McEwan, D.G., Richter, B., Claudi, B., Wigge, C., Wild, P., Farhan, H., McGourty, K., Coxon, F.P., Franz-Wachtel, M., Perdu, B., et al. (2015). PLEKHM1 regulates salmonella-containing vacuole biogenesis and infection. Cell Host Microbe *17*, 58–71. 10.1016/j.chom.2014.11.011.

60. Sindhwani, A., Arya, S.B., Kaur, H., Jagga, D., Tuli, A., and Sharma, M. (2017). Salmonella exploits the host endolysosomal tethering factor HOPS complex to promote its intravacuolar replication 10.1371/journal.ppat.1006700.

61. Bernal-Bayard, J., Cardenal-Muñoz, E., and Ramos-Morales, F. (2010). The salmonella type III secretion effector, Salmonella Leucine-rich Repeat Protein (SlrP), targets the human chaperone ERdj3. Journal of Biological Chemistry *285*, 16360–16368. 10.1074/jbc.M110.100669.

62. Antoniou, A.N., Lenart, I., Kriston-Vizi, J., Iwawaki, T., Turmaine, M., McHugh, K., Ali, S., Blake, N., Bowness, P., Bajaj-Elliott, M., et al. (2019). Salmonella exploits HLA-B27 and host unfolded protein responses to promote intracellular replication. Ann Rheum Dis *78*, 74–82. 10.1136/annrheumdis-2018-213532.

63. Kriston-Vizi, J., Lenart, I., Iwawaki, T., Gould, K., Nesbeth, D., Powis, S.J., and Antoniou, A.N. (2022). Salmonella Exhibit Altered Cellular Localization in the Presence of HLA-B27 and Codistribute with Endo-Reticular Membrane. J Immunol Res *2022*. 10.1155/2022/9493019.

64. Lee, J.E., Cathey, P.I., Wu, H., Parker, R., and Voeltz, G.K. (2020). Endoplasmic reticulum contact sites regulate the dynamics of membraneless organelles. Science (1979) *367*. 10.1126/science.aay7108.

65. Geddes, K., Worley, M., Niemann, G., and Heffron, F. (2005). Identification of new secreted effectors in Salmonella enterica serovar typhimurium. Infect Immun *73*, 6260–6271. 10.1128/IAI.73.10.6260-6271.2005.

66. McQuate, S.E., Young, A.M., Silva-Herzog, E., Bunker, E., Hernandez, M., de Chaumont, F., Liu, X., Detweiler, C.S., and Palmer, A.E. (2017). Long-term live-cell imaging reveals new roles for Salmonella effector proteins SseG and SteA. Cell Microbiol *19*. 10.1111/cmi.12641.

67. Domingues, L., Ismail, A., Charro, N., Rodríguez-Escudero, I., Holden, D.W., Molina, M., Cid, V.J., and Mota, L.J. (2016). The Salmonella effector SteA binds phosphatidylinositol 4-phosphate for subcellular targeting within host cells. Cell Microbiol *18*, 949–969. 10.1111/cmi.12558.

68. Derré, I., Swiss, R., and Agaisse, H. (2011). The lipid transfer protein CERT interacts with the Chlamydia inclusion protein IncD and participates to ER-Chlamydia inclusion membrane contact sites. PLoS Pathog *7*. 10.1371/journal.ppat.1002092.

69. Kolodziejek, A.M., Altura, M.A., Fan, J., Petersen, E.M., Cook, M., Brzovic, P.S., and Miller, S.I. (2019). Salmonella Translocated Effectors Recruit OSBP1 to the Phagosome to Promote Vacuolar Membrane Integrity. Cell Rep *27*, 2147-2156.e5. 10.1016/j.celrep.2019.04.021.

70. Liu, K., Kong, L., Graham, D.B., Carey, K.L., and Xavier, R.J. (2021). SAC1 regulates autophagosomal phosphatidylinositol-4-phosphate for xenophagy-directed bacterial clearance. Cell Rep *36*. 10.1016/j.celrep.2021.109434.

71. Chen, D., Burford, W.B., Pham, G., Zhang, L., Alto, L.T., Ertelt, J.M., Winter, M.G., Winter, S.E., Way, S.S., and Alto, N.M. (2021). Systematic reconstruction of

673 an effector-gene network reveals determinants of Salmonella cellular and tissue
674 tropism. Cell Host Microbe *29*, 1531-1544.e9. 10.1016/j.chom.2021.08.012.

675 72. Göser, V., Kehl, A., Röder, J., and Hensel, M. (2020). Role of the ESCRT-III
676 complex in controlling integrity of the Salmonella-containing vacuole. Cell
677 Microbiol *22*. 10.1111/cmi.13176.

Graphical Abstract

Salmonella infection leads to ER stress and ER expansion

(A) SCV makes contact with ER using effector SteA

(B) ER tubules facilitate the SCV division

SCV

SteA recruitment on SCV

Induction of ER stress

Fig:1 *Salmonella* infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum

1G

1H

1I

1J

Fig:1-*Salmonella* infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum (A) Representative confocal microscopic images of uninfected and infected HeLa cells with STM WT mCh or STM*ΔssaV* mCh at MOI 10, 16h post-infection and immuno-stained against ATF4, LAMP1 and DAPI. (B) Represents the quantification of Cell Total Corrected Fluorescence (CTCF) intensity of ATF4 inside the nucleus, N=4, n=20-30 cell per experiment, data is representative of mean±SD (C) Representative confocal microscopic images of uninfected and infected HeLa cells with STM WT mCh or STM*ΔssaV* mCh at MOI 10, 16h post infection and immuno-stained against ATF6, LAMP1 and DAPI. (D) Represents the quantification of Cell Total Corrected Fluorescence (CTCF) intensity of ATF4 inside the nucleus, N=3, n=20-30 cell per experiment, data is representative of mean±SD (E) Representative confocal microscopic images of uninfected and infected HeLa cells with STM WT mCh or STM*ΔssaV* mCh at MOI 10, 16 hour post infection and immuno-stained against XBP1, LAMP1, and DAPI (F) Represents the quantification of Cell Total Corrected Fluorescence (CTCF) intensity of XBP1 inside the nucleus, N=3, n=20-30 cell per experiment, data representative of mean±SD (G) quantitative real-time PCR of *Rtn4a* (H) and *Climp63*, N=3, n=3, Data is representative of mean±SEM (I) confocal microscopic images of Un-infected and infected HeLa cell showing ER expansion (J) Represents the quantification of tubular ER, N=4,n>50 cells per experiment, data is representative of mean±SD. Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig 2: Expansion of ER tubules facilitates SCV proliferation and its division

2H

2I

Fig2 Expansion of ER tubules facilitates SCV proliferation and its division (A) Represents the ICSA upon overexpression of Rtn4a, N=3, n=3 mean± SEM(B) ICSA upon Rtn4a knockdown in HeLa cells, N=3, n=3 mean± SEM (C) Representative confocal microscopic images of SCV number per cell upon overexpression of mCh-EV, sec61βs-mCh, and mCh-Climp6 (D) quantification of the number of SCV per cell, representative of N=2, n=20-30 cells per experiment (E) snapshot of live cell imaging of *Salmonella* infected HeLa cells, White arrowhead represent the ER overlap at dividing bacteria. White line along the length of bacteria represent the portion used for line scan. (F) showing the line scan of images. Blue arrows represent the images for which the line scan is analyzed. (G) quantification of SCV fission events marked by ER n=11 fission events from 7 cells (H) Confocal microscopic images representing SCV fission events marked by ER, at center or adjacent (I) quantification of fission events in fixed images marked by ER, n=49 fission events from 31 cells. White arrowhead represent the ER overlap at dividing SCV. Student's t-test was used to analyze the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig 3: *Salmonella* translocated effector steA is crucial for maintaining SCV contacts with ER resulting in a single bacterium per vacuole state

3A

LAMP1 | STM WT | Merge
5 µm | 5 µm | 5 µm
STM*ΔsteA*
5 µm | 5 µm | 5 µm

3B

% of cells showing multiple bacteria per vacuole

***

STM WT | STM*ΔsteA*

3C

DAPI | LAMP1 GFP | steA-HA | mCh-Sec61β | Merge
10µm | | | | 1µm

3D

DAPI | RFP KDEL | STM WT GFP | LAMP1 | RFP KDEL+ LAMP1 | Merge
5 µm | 5 µm | 5 µm | 5 µm | 5 µm | 5 µm
| RFP KDEL | STM*ΔsteA* GFP | LAMP1 | |
5 µm | 5 µm | 5 µm | 5 µm | 5 µm | 5 µm

3E

Colocalization coefficient of RFP KDEL with SCV

****

STM WT | STM*ΔsteA*

Fig 3: *Salmonella* translocated effector steA is crucial for maintaining SCV contacts with ER resulting in a single bacterium per vacuole state (A) Confocal microscopic images showing SCV formed by STM WT and STM*ΔsteA* (B) Quantification of the number of cells showing multiple bacteria in one vacuole, representative N=3, n>50 cells per experiment, mean±SEM (C) Representative confocal image of infected HeLa cell showing contact between steA-HA and ER, arrowheads represent site of SCV fission and interaction between ER and steA-HA (D) Confocal images represents the colocalization of ER with SCV (E) Quantification of colocalization coefficient of ER marker RFP KDEL with SCV marker LAMP1, Data is representative N=2, n>50 cells per experiment, 16h post infection, mean±SD. Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig 4: STM*ΔsteA* shows defect in proliferation inside epithelial cells and pathogenicity in mice in vivo model

Fig 4: STM*ΔsteA* shows defect in proliferation inside epithelial cells and pathogenicity in mice in vivo model(A) Intracellular survival assay of STM WT, STM*ΔsteA*, and STM*ΔsteA:steA* in HeLa cells, data is representative of N=3,n=3, mean±SEM. (B) Line scan representation to quantify SCV diameter where red line indicates the area used for quantification (C) Quantification of SCV diameter, N=3,n>50 cells per experiment (D) Representative microscopic images showing colocalization of LC3B with SCV (E) Quantification of colocalization coefficient of LC3B with SCV, data is representative of N=3, n>50 cells per experiment, mean±SD (F) outline of experimental procedure of mice infection (G) CFU enumeration in Liver (H) Spleen (I) Mesentric lymph node, data is representative of one experiment, N=2, n=5 animal per cohort (J) survival growth curve of cohort of mice infected with STM WT, STM*ΔsteA*, and STM*ΔsteA:steA,* n=8 mice per cohort. Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05. Mann-Whitney test was performed for animal experiment.

Supplementary Fig:1 *Salmonella* infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum

Fig:1 *Salmonella* infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum Representative confocal microscopic images of uninfected, STM WT mCh and STMΔssaV mCh infected cells, immunostained with (S1A) ATF4, LAMP1, DAPI, 6h post infection (S1B) 2h post infection (S1C) ATF6, LAMP1 and DAPI, 6h post infection (S1D) 2h post infection (S1E) XBP1, LAMP1 and DAPI, 6h post infection (S1F) 2h post infection (S1G) Representing uninfected, infected and bystander cell, showing an expansion of ER tubules (S1G) Quantification of ER tubules, N=2, n>20 cells per experiment. . Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Supplementary Fig 2: Expansion of ER tubules facilitates SCV proliferation and its division

Fig2 Expansion of ER tubules facilitates SCV proliferation and its division (S2A) Percent invasion of STM WT in HeLa cells upon overexpressing mCh-EV, Sec61β-mCh, and Rtn4a-mCh (S2B) Quantitative PCR of *Rtn4a* upon knock down, N=3, n=3, Data is representative of mean±SD (S3C) Live cell imaging snapshots of RAW 264.7 murine macrophage cell line infected with STM WT showing ER overlap at fission site, N=1.(S2D) confocal images of infected HeLa cells showing SCV localization at cell periphery, 16h post infection. Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Supplementry Fig:3 *Salmonella* translocated effector steA is crucial for maintaining SCV contacts with ER resulting in a single bacterium per vacuole state

S3A

S3B

Fig 3: *Salmonella* translocated effector steA is crucial for maintaining SCV contacts with ER resulting in a single bacterium per vacuole state (S3A) Representative microscopic images showing colocalization of eGFP or steA-eGFP with RFP KDEL (ER marker) (S3B) quantification of colocalization coefficient or eGFP with RFP KDEL. Data is representive of N=2, n>50 cells per experiment, mean±SD. Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Supplementary Fig 4: STM*ΔsteA* shows defect in proliferation inside epithelial cells and pathogenicity in mice in vivo model

Fig 4: STM*ΔsteA* shows defect in proliferation inside epithelial cells and pathogenicity in mice in vivo model (S4A) percent invasion of STM WT, STMΔsteA, and STMΔsteA:steA in HeLa cells (S4B) Representative images of uninfected and infected HeLa cells with STM WT and STM*ΔsteA* showing intensity of LAMP (S4C) Quantification of mean fluorescence intensity of LAMP1, data is representative of N=3, n>50 cells per experiment, mean±SD (S4D) CFU analysis of STM WT and STMΔsteA in blood sample of C57BL/6 mice, 5[th] day post oral gavage , N=2, n=5 mice per cohort.(S4E) Represents weight reduction in mice upon infection with STM WT, STM*ΔsteA and* STM*ΔsteA:steA* till 8 days post infection, 8 mice per cohort. Student's t test was used to analyse the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Supplementary File: Table-1, List of Primers used in this study:

S.No	Primer Name	Sequence (5'-3')
1	steA-KO-FP	CAAATAGTTATGGTAGCGAGCTTTTATGTCGGCCGCCCAT CATATGAATATCCTCCTTAG
2	steA-KO-RP	TCAGTTTCTACCTATGCCAGAGCTTTATCAGGAAATAAGC GTGTAGGCTGGAGCTGCTTC
3	steA-eGFP cloning-FP	TTTTTCTCGAGATGCCATATACATCAGTTTCTACC
4	steA-eGFP cloning-RP	TTTTTGGATCCATAATTGTCCAAATAGTTATGG
5	HU-Rtn4a-RT-FP	TCGGGCTCAGTGGATGAGA
6	HU-Rtn4a-RT-RP	GCAGGACAGATGGGAAATCCT
7	HU-Climp63-RT-FP	GCGTCGAGCAGAAGGTGC
8	HU-Climp63-RT-RP	CATGGATCCCATCCGAGAGG
9	Xho1_fwd_Native_promoter_steA_cloning	TTTTTCTCGAGCGGCAGTGATTGCGTTGC
10	HindIII steA-HA_cloning_RP	TTTTTTAAGCTTTTATGCATAATCCGGAACATCATACGGAT AATAATTGTCCAAATAGTTATGGTAGCGAGC
11	HU-β-actin-FP	GCCGCCAGCTCACCAT
12	HU-β-actin-RP	TCGTCGCCCACATAGGAATC

Supplementary File: Table:2 List of Antibodies

Antibody	Catalog number
ATF-4	CST-11815s
ATF-6	ATF-6α Antibody (H-280): sc-22799
XBP-1	XBP-1 Antibody (M-186): sc-716
LAMP1	DSHB 1D4B
Anti-Salmonella	Invitrogen- PA1-20811
LC3B	Sigma-L7543

Supplementary File: Table: 3 List of plasmids used for overexpression

Plasmid	Source
1) pHAGE2 mCherry-Rtn4a (Plasmid-86683)	pHAGE2 mCherry-Rtn4a was a gift from Tom Rapoport (Addgene plasmid # 86683)
2) mCh-Climp63 (Plasmid #136293)	mCh-Climp63 was a gift from Gia Voeltz (Addgene plasmid # 136293)
3) RFP-KDEL	RFP KDEL construct was a kind gift from Prof. Nagaraj Balasubramanian
4) LAMP1-GFP	GFP-LAMP1 construct was provided by Prof. Mahak Sharma
5) Sec61β-mCh	mCh sec61β was a kind gift from Prof. Gia Voeltz (Addgene 49155)

List of shRNA plasmids used for knockdown

S.No.	Target Gene	TRC ID	Sequence
1	RTN4 reticulon 4 [*Homo sapiens* (human)]	TRCN0000179649	CCGGGCAGTGTTGATGTGGGTATTTCTCGAGAAATACCCACATCAACACTGCTTTTTTG
2	RTN4 reticulon 4 [*Homo sapiens* (human)]	TRCN0000147624	CCGGGCATATCTGGAATCTGAAGTTCTCGAGAACTTCAGATTCCAGATATGCTTTTTTG