Endoplasmic Reticulum contact sites facilitate the coordinated division of *Salmonella*-

containing vacuole

Umesh Chopra[1], Priyanka Bhansali[1], Subba Rao Gangi Setty[1], Dipshikha Chakravortty[1,2*]

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, Karnataka

560012, India

[2]School of Biology, Indian Institute of Science Education and Research Thiruvananthapuram,

Maruthamala PO, Vithura, Thiruvananthapuram - 695551, Kerala, India

[*]Corresponding Author- dipa@iisc.ac.in

Abstract

Salmonella Typhimurium (STM) resides in a membrane-bound compartment called *Salmonella*

containing vacuole (SCV) in several infected cell types. Within host cells, the division of bacteria

and SCV are synchronous to maintain the single bacterium per vacuole. However, the mechanism

regulating the synchronous fission and the machinery is not well understood. The fission of several

intracellular organelles is regulated by the dynamic nature of the tubular endoplasmic reticulum

(ER). In this study, we have evaluated the role of ER in controlling SCV fission. Interestingly,

Salmonella-infected cells show the activation of unfolded protein response (UPR) with expanded

ER tubules compared to the uninfected cells. Further, changing the expression of ER morphology

regulators, such as reticulon-4a (Rtn4a) and CLIMP63, affected bacterial proliferation

significantly, suggesting a potential role for tubular ER in facilitating the SCV division. Live-cell

imaging analysis shows the marking of tubular ER precisely at the center of the majority of SCV

division (78%) sites. We have investigated the role of SteA (a known *Salmonella* effector in

modulating the membrane dynamics) in coordinating the SCV division. We observed that SteA

resides on the SCV membranes and helps in making membrane contact sites between SCV and ER. Accordingly, the colocalization of ER with SCV enclosing SteA mutant *Salmonella* was significantly reduced compared to SCV-formed by wild-type *Salmonella*. Depletion of *steA* in *Salmonella* resulted in profound defects in SCV division, resulting in multiple bacteria residing in a single vacuole with defects in proliferation compared to the wild-type strain in epithelial cells. Also, during *in vivo* infection, the STM*ΔsteA* mutant shows a defect in colonization in the spleen and liver and affects the initial survival rate of mice. Overall, this study suggests a coordinated role of bacterial effector SteA in promoting the ER contact sites with SCVs and thus regulating the successful division of SCV.

Introduction

Salmonella enterica subsp. enterica serovar Typhimurium (STM) is an intracellular pathogen known to infect a wide range of hosts and result in diseases across species, including gastroenteritis to systemic infections [1], [2]. S. Typhimurium enters the body through the fecal-oral route following the bacteria utilize an arsenal of strategies to overcome several stresses like the acidic pH of the stomach, gut flora, bile salts, etc., and finally reach the small intestinal epithelia [3]. Salmonella secretes several virulence proteins inside the cell, which leads to actin rearrangement and membrane ruffling, which helps in the invasion of bacteria inside the host epithelial cell [4], [5], [6], [7]. Salmonella effector SopD is known to facilitate the recruitment of dynamin2 (Dyn2) at the invasion site, which helps in the endocytosis of bacteria inside host cells [8].

Upon invasion into the host cell, Salmonella resides in Salmonella containing vacuole (SCV), a membrane-enclosed compartment. Mature SCV shares several characteristic markers with lysosomes/late endosomes such as LAMP1 and Rab7 but is negative for acid hydrolases and cathepsins [9]. Inside the epithelial cells, several protruding structures from SCV known as Salmonella-induced filaments (SIFs) are observed, which are LAMP1 positive [10], [11]. The membrane composition of SCV is not very well understood, but it contains molecules and proteins from several host organelles and is enriched with several bacterial effector proteins [12], [13]. Our group, as well as several others, have previously shown that Salmonella resides as a single bacterium per vacuole state inside the host, suggesting that SCV membrane divides synchronously with the division of Salmonella but the mechanism behind SCV division is not known [14], [15], [16], [17]. For successful SCV division, expansion of SCV membrane, pinching of SCV membrane

precisely with bacterial division, and lastly, scission of host-derived SCV membrane must be

followed in an orchestrated manner.

Several literatures have highlighted the importance of contact sites between cellular organelles,

such as contact sites of ER with mitochondria, plasma membrane and endosomes, and contact sites

between mitochondria with lysosomes [18], [19], [20], [21]. Contact sites between ER-

mitochondria and ER endosomes are very widely studied in the aspect of organelle division [18],

[22]. In the case of mitochondria, which comprises an inner and an outer membrane, both

membranes divide simultaneously during its fission. The inner mitochondrial membrane uses its

machinery for fission, while outer membrane constriction is initiated by the ER, followed by the

recruitment of Drp1, which mediates the fission of mitochondria [18], [23], [24]. Bacterial

pathogens like *Chlamydia* and *Brucella* make contact sites with cellular organelles to facilitate

their proliferation inside the host cell [25], [26], [27]. Interestingly, the SCV membrane also

comprises several ER proteins, and recently, membrane contact sites between SCV and ER have

also been observed [28]. To understand the mechanism behind the SCV division, we reasoned that

SCV is partially like mitochondria. SCV and mitochondria (being an endosymbiont) encompass

an outer membrane derived from the eukaryotic host, while the inner membrane of mitochondria

is prokaryotic in nature, like the cell wall of *Salmonella*. To understand the same, we hypothesized

that a process similar to mitochondria could lead to SCV division. Our results suggest that

Salmonella infection leads to activation of UPR, ER expansion, and an increase in ER tubulation.

We have observed ER tubules mark at the fission site of SCV division which possibly initiate the

constriction. *Salmonella* translocated effector SteA localizes onto the SCV membrane and possibly

helps in making contact with ER, suggesting that SteA can act as a factor to recruit the ER onto

SCV and thus facilitate the SCV division. STM*ΔsteA* resides as multiple bacteria in one big

vacuole with defects in proliferation. During *in vivo* infection, STM*ΔsteA* shows defects in

colonization in the spleen and liver if injected through the intraperitoneal route. Also, there is a

delay in the onset of mice death upon infection with STM*ΔsteA* as compared to STM WT, but

eventually, mice succumbed to death on the 8th day post-infection. Overall, our study unravels,

for the first time, the coordinated role of ER and bacterial effector SteA in successful SCV division

inside the cell.

Materials and Methods

Bacterial strains and culture conditions

Salmonella enterica subspecies *enterica* serovar Typhimurium (STM WT) wild-type strain ATCC

14028s was used in all experiments. The bacterial strain was cultured in Luria broth (LB-Himedia)

with constant shaking (170 rpm) at 37°C orbital-shaker. Antibiotics like kanamycin (50 µg/ml) and

ampicillin (50µg/ml) were used wherever required. A one-step gene inactivation method was used

for the generation of knockout strains [29]. Wild type as well mutant strains were transformed with

pFPV-m-cherry or pFPV-GFP plasmid for immunofluorescence assays.

Table:1 List of bacterial strains used in this study

Strain	Characteristic (Antibiotic resistance)	Source
STM WT 14028s	-	Kind gift by Prof. M. Hensel
STMΔ*ssaV*	Kan^R	Laboratory stock
STMΔ*steA*	Kan^R	This study
STMΔ*steA:steA*	Kan^R +Amp^R	This study

Generation of Complement strain and SteA-eGFP plasmid

Low copy number plasmid pQE-60 was used to prepare the SteA-HA complement strain. Briefly,

PCR amplification (using Thermo Q5 DNA polymerase) of the *steA* gene was performed using

Wild Type STM colony with respective PCR primers. Forward cloning primer was chosen from a

position of 250 base pairs upstream of the gene start site to amplify the native promoter of the gene

during PCR and to maintain the native expression level of protein, and the sequence of HA tag was

added in reverse primer. The amplified PCR product was purified using a Gel extraction kit

(Qiagen). The PCR product and Plasmid were subjected to restriction digestion by using Xho1

(NEB) and Hind III (NEB) at 37°C for 2 hours. The double-digested PCR product and plasmid

were further subjected to Ligation using T4 DNA ligase (NEB) at 16°C for 12-14 hours. The

ligated product was further electroporated in STM*ΔsteA* strain to prepare the complemented strain,

i.e., STM*ΔsteA*:pQE60-*steA* (hence referred to as STM*ΔsteA:steA*). The expression level of

complement was confirmed with western blotting and confirmed with DNA sequencing.

For preparation of SteA-eGFP plasmid for transfection study, eGFP-N3 empty vector was used.

The sequence of *steA* was amplified via PCR from STM WT using primer with restriction site for

Xho1 and BamH1. The entire process for cloning was performed as discussed above.

Eukaryotic Cell Culture, Transfection and Plasmids

HeLa cells and RAW 264.7 murine macrophages were cultured in DMEM (Lonza) containing 10%

FBS (Gibco) at a humified incubator, maintaining a temperature of 37°C and 5% CO_2. Cells were

seeded in 6 wells, 12 wells, or 24 well plates at a confluency of 60-70% prior to infection. Cells

were maintained in the presence of 1% penicillin and streptomycin (penstrep). Cells were given a

PBS wash and supplemented with fresh media without penstrep at least 6 hours prior to infection.

Plasmid overexpression or knockdown via shRNA was done using Lipofectamine 3000 using

manufacturer protocol. Briefly, 300ng of plasmid was mixed in opti-MEM media for 5 min while

1ul of L3000 was added (for one well of 24 well plates) in opti-MEM for 5 min separately and

further mixed and incubated for 20 min at room temperature. Post incubation, this mixture was

added to cells for 6 hours for transfection to take place. Cells were washed with PBS and

supplemented with complete media post 6 hours of treatment, and post 48h of transfection,

infection was given. Overexpression constructs such as pHAGE2 mCherry-Rtn4a was a kind gift

from Prof. Tom Rapoport (Addgene plasmid 86683) [30] mCh Climp-63 and mCh sec61β was a

kind gift from Prof. Gia Voeltz (Addgene plasmid 136293, Addgene 49155) [31], [32], RFP KDEL

construct was a kind gift from Prof. Nagaraj Balasubramanian [33], GFP-LAMP1 was provided

by Prof. Mahak Sharma [34]. The shRNA plasmids were obtained from the human genome-wide

TRC shRNA library (purchased from Sigma-Aldrich, USA, Catalog no. SH0411), available at

shRNA resource centre, MCB, IISc. pLKO.1-puro non-mammalian shRNA (SHC002, shControl)

was used as a control in all the shRNA mediated knockdown experiments. Plasmid sequences and

details are added in supplementary table 2.

Protocol for infection (Gentamicin Protection Assay)

The cells were seeded at the required confluency and infected with *Salmonella* strains with actively

growing log phase culture (for epithelial cells) and stationary phase culture (for RAW 264.7

macrophages) at Multiplicity of infection (MOI) 10 for Intra Cellular Survival Assay (ICSA) and

confocal imaging. For the live cell imaging experiment, an MOI of 25 was used to increase the event of finding infected cells for imaging. For confocal microscopy studies, cells were seeded on sterile glass coverslips at least 24 hours before infection. Upon infecting the cells at the required MOI, the plate was centrifuged at 600 rpm for 5 min to facilitate the adhesion, and then the plate was incubated for 25 min at 37°C and 5% CO_2. Post incubation, bacteria-containing media was removed, and cells were given 2 PBS washes to remove any extracellular bacteria. Fresh media containing 100µg/ml of gentamicin was added into wells and incubated at 37°C for one hour. Following this, cells were again given 2 PBS washes, and fresh media containing 25µg/ml of gentamicin was added, and this concentration was maintained till the end point of the experiment. Cells were harvested at required time points, such as 2h and 16h for ICSA, 2h, 6h, and 16h for confocal microscopy-based experiments.

Confocal Microscopy

After an appropriate hour post-infection, cells were washed with PBS twice and fixed with 3.5% PFA at room temperature for 15 min. After fixation, coverslips were washed with PBS, and cells were stained with primary antibody prepared in 2% BSA in PBS carrying 0.01% of saponin (for membrane permeabilization) for 1 hour at room temperature or 4°C overnight in a cold room as required. Post incubation, coverslips were washed again twice with PBS and incubated with specific secondary fluorophore-labeled antibodies and incubated for one hour at room temperature. Coverslips were then mounted on clean glass coverslips using mounting media. Once the mounting media was dried, coverslips were stabilized by sealing the periphery with transparent nail paint on the sides. Images were acquired on a confocal scanning laser microscope (Zeiss 880 multiphoton microscope or Leica sp8 falcon system with or without zoom as required). Zen black software was used for all colocalization-based quantification and to calculate mean fluorescence intensity (MFI).

Calculations of CTCF, percent of ER tubules, SCV diameter, and Line scans were prepared using

ImageJ. The ER tubule percentage is calculated by calculating the sheet percentage from the total

cell area. Sheet area / total area *100 = Sheet area percentage and further 100 - Sheet area

percentage = tubule area percentage [35]. Live cell microscopy was performed on the Leica SP8

Falcon microscope, and live cell video was processed and prepared using LAS X software.

Antibodies used in this study are mentioned in supplementary table 3.

165 Bacterial enumeration after infection for Intracellular survival assay

After the appropriate hour of post-infection, cells were washed with PBS at RT and lysed using

0.1% Triton-X 100. Cell lysate was plated on SS agar to obtain colony forming unit (CFU) and it

was used for the calculation of Percent Invasion and Fold proliferation using this formula.

Percent Invasion: (CFU obtained at 2 hours post-infection/ CFU obtained from pre-inoculum)

*100

Fold proliferation: CFU obtained at 16-hour post-infection/CFU obtained at 2 hours PI

172 RNA isolation and Real-Time PCR (qRT-PCR)

RNA isolation was performed using TRIzol (Takara) following the manufacturer's protocol.

Quantification of RNA was done using nano-drop (Thermo-Fisher Scientific) and RNA samples

were run on 1% agarose gel to check the quality of RNA. 2 µg of RNA was subjected to DNase I

(Thermo Fischer Scientific) treatment at 37°C for 1 hour followed by addition of 0.5M EDTA

(final concentration 5mM) and heat inactivation at 65°C for 10 mins to inactivate the enzyme.

Further, cDNA was prepared by using a cDNA synthesis kit (Takara) as per the manufacturer's

instructions. The expression profile of all genes of interest was evaluated using RT primers by

using SYBR green mix (Takara) in a Biorad Real-Time PCR machine. The expression value of

the target gene was normalized to housekeeping internal control β-actin and further compared with

uninfected cells. Primers used in this study are mentioned in Supplementary Table 4.

***In vivo* animal experiment**

6-8 weeks old C57BL/6 female mice were infected by orally gavaging 10^7 CFU of STM WT,

STM*ΔsteA* and STM*ΔsteA:steA*. 5 days post-infection, mice were euthanized, and organs such as

liver, spleen, MLN, and blood were harvested to study the colonization in organs. For

intraperitoneal injection, 10^3 CFU of bacteria was administered per animal, and 3 days post-

infection, organs were homogenized and further plated on *Salmonella-Shigella* agar. Obtained

CFU values were normalized with the gram weight of the organ and then converted into a log

scale. Blood was collected through heart puncture and plated on SS agar. For the mice survival

assay, 10^8 CFU was administered to each mouse via oral gavage. Mice were monitored every

day till the time none of the mice in the cohort survived. The animal experiments were carried

out in accordance with the approved guidelines of the institutional animal ethics committee at the

Indian Institute of Science, Bangalore, India (Registration No: 48/1999/CPCSEA). All

procedures involving the use of animals were performed according to the Institutional Animal

Ethics Committee (IAEC)-approved protocol.

Statistical analysis:

Statistical analysis was performed using GraphPad Prism software. For analysis, the student's t-

test (unpaired parametric test) was performed. The results are indicative of mean± SD or mean±

SEM. To calculate the statistical significance of bacterial colonization *in vivo*, the Mann-

Whitney test was performed. Group size, experimental number, no of cells in each set, and P

value of each experiment are described in figure legends.

Results

Salmonella infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum

Several bacterial and viral infections are known to activate UPR for their survival [36], [37]. UPR helps to alleviate the ER stress by initiating several transcriptional and translational regulations which lead to the maintenance of ER homeostasis [38], [39]. Inside the cell, UPR signaling is maintained by transmembrane proteins located in the ER membrane and their catalytic domain towards the cytosol. Inositol requiring enzyme-1 (IRE-1), Protein kinase c like ER kinase (PERK), and Activating transcription factor-6 (ATF-6) comprises the main arm of UPR. Recent reports have shown the contact of SCV with ER, also the presence of ER proteins in the membrane fraction of SCV suggests the crosstalk between them [12], [28]. Here, we investigated how *Salmonella* infection modulates the ER dynamics. To understand the activation of UPR by any of these pathways, we used confocal microscopy and quantified the translocation of downstream transcription factors, such as XBP1, ATF4, and ATF6, inside the nucleus upon infection. SPI2 mutants of *Salmonella*, STM*ΔssaV* (which is unable to secret SPI-2 encoded effector protein in host cytosol) was kept as a control to understand if activation of UPR is mediated by bacterial effectors or dependent on bacterial invasion. Our results suggest that from as early as post 2h of *Salmonella* infection, both the transcription factors, ATF4 and ATF6 from the cytosol translocated to the nucleus of infected cells as compared to uninfected cells and STM*ΔssaV* infected cells, suggesting that only wild-type *Salmonella* can activate UPR (fig 1A-1D). *Salmonella*-induced ATF4 and ATF6 translocation were retained till   16h post-infection (fig S1A-S1D). However, we

didn't observe any increase in XBP1 recruitment inside the nucleus (fig 1E,1F) upon infection at any time point post-infection (fig S1E, S1F), suggesting that *Salmonella* infection upregulates UPR by PERK and ATF6 pathways in HeLa cells. To alleviate the ER stress, activation of UPR leads to the expansion of the ER, protein folding capacity, and increased phospholipid production [40]. Increased bacterial proliferation requires a continuous source of membrane for the increasing number of SCV, and increased phospholipid production upon UPR could facilitate the SCV proliferation inside the host cell. Therefore, we sought to determine if *Salmonella* infection led to ER expansion during the pathogenesis phase. Our transcriptional analysis suggests that upon infection, there is upregulation at the transcript level of ER shaping proteins such as *Climp63* (responsible for maintaining ER sheets) and *Reticulon-4a* (*Rtn4a*, essential for maintaining ER tubules) (fig 1G, fig 1H). The upregulation was significantly observed at the late time point of infection, i.e., 16h post-infection, suggesting that active proliferation of *Salmonella* and not just invasion (as evident at 2h and 6h post-infection) leads to ER expansion.

Since an increase in the transcript levels of ER shaping proteins was observed at the later time of infection, we tried to quantify the percentage of ER sheets and tubules using the ER marker, RFP-KDEL, upon infection to understand ER expansion. Upon analysis of microscopy images, we observed that *Salmonella*-infected cells showed an increased percentage of ER tubules compared to uninfected cells (fig1I, fig1J). The changes in percent ER tubules were only evident in infected cells as compared to uninfected and bystander cells (neighboring cells that are not infected), suggesting that stabilized SCV and bacterial effectors are responsible for this phenotype (fig S1G and S1H). Together, this data indicates that *Salmonella* infection leads to changes in ER morphology.

Expansion of ER tubules facilitate SCV proliferation and its division

Dynamic ER tubules play an essential role in forming membrane contact sites with several

organelles [41], [42]. Figure 1 indicates that actively proliferating bacteria increased tubular ER

and upregulated the expression of *Rtn4a*. We also tried to understand if there is an advantage of

the increased tubules to bacterial proliferation. To identify the role of ER tubules in mediating

bacterial proliferation, we transiently overexpressed reticulon 4a (Rtn4a) in HeLa cells, which

helps to increase ER tubules [43], and then quantified the proliferation of bacteria by intracellular

survival assay (ICSA). As compared to empty vector and general ER marker sec61β-mCh control,

increased ER tubulation by Rtn4a led to the higher proliferation of *Salmonella* with no significant

change in percent invasion, suggesting an increase in ER tubules helps in bacterial proliferation

(fig2A, figS2A). To validate if this change in proliferation was dependent solely on the function

of Rtn4a, knockdown of Rtn4a via shRNA was performed (figS2B). Knockdown of Rtn4a (˜40%

reduction at transcript level) led to reduced proliferation of bacteria as compared to scrambled

control, hence strengthening the crucial role of ER tubules in bacterial proliferation (fig2B).

Climp63 overexpression led to an increase in ER sheets as opposed to tubules. We quantified the

number of SCV per cell upon transiently overexpressing the Climp63 in HeLa cells. Our data

indicated that the number of SCV present per cell decreases upon overexpressing ER sheets using

Climp63 compared to controls (fig2C, fig2D). Division of *Salmonella* is linked with the fission of

SCV membrane [15]. Since the intricate balance of ER sheets and tubules affects bacterial

proliferation, we hypothesized that ER tubules might facilitate SCV fission. In the case of

mitochondria, membrane-less organelle, and endosomes, ER marks the division site, and later

fission is completed by Drp1 (in mitochondria) and coronin1c and TMCC 1 complex (in

endosome) [18], [24], [44], [45]. To investigate our hypothesis of ER tubules playing a role in SCV

fission, we performed live cell imaging in cells transfected by RFP-KDEL (to mark the ER)

followed by infection with *Salmonella*. Quantification of data suggests that ER tubules mark the

division site precisely at the center of dividing SCV in 78% of events, as observed by image

snapshots and corresponding line scans (fig2E-G, video S1). STM WT infected RAW macrophage

also showed contact of ER on the dividing bacteria (video S2, S2C). This was also evident in fixed

cell images where we could observe Lamp1 enclosed *Salmonella* division in SCV marked by ER

tubules exactly in the center in 56.7±6.61% of fission events, however, in 14±9.07% of events, ER

was seen in proximity to the center and hence mentioned as adjacent while 29.2±8.4% of dividing

SCV did not possess ER tubules on them (fig2H, fig2I). This data suggests the importance of ER

tubules in marking the fission site of SCV. Taken together this data suggests that ER tubules help

in the proliferation and division of SCV by marking the fission site inside the cell and help in

maintaining SCV number within the cell.

***Salmonella* translocated effector SteA is crucial for maintaining SCV contact site with ER,**

resulting in a single bacterium per vacuole

SCV division leads to a single bacterium per vacuole throughout the course of pathogenesis in cell

lines or *in vivo* [14]. Domingues et al., suggested the role of SteA in maintaining the membrane

dynamics of SCV [46]. In accordance with the same, we also observed that *steA* mutant strain of

Salmonella (STM*ΔsteA*) resides as multiple bacteria in one single vacuole (fig 3A). Our data

suggests that compared to STM WT, 30±3% of mutant bacteria reside in one big vacuole,

suggesting that loss of function of *steA* leads to improper SCV fission (fig 3B). Live cell imaging

shows several points of contact between SCV and ER, suggesting that SCV actively makes

membrane contact sites with ER for its survival and proliferation (fig 2E, fig 3C). SteA resides

onto the PI4P-enriched SCV membrane (fig 3C), and hence, we asked if membrane contacts

between ER and SCV are maintained by SteA. Microscopy images show that SteA-enriched SCV

makes contact with ER (fig 3C). Quantification of the data suggests that ER colocalizes more with

the SCV of STM WT than the SCV formed by STM*ΔsteA* (as observed colocalization between

SCV marker Lamp1 and RFP KDEL), suggesting that SteA helps in making contact sites of SCV

with ER (fig 3D, fig 3E). This was also observed upon transfecting HeLa cells with the SteA-eGFP

construct, where SteA-eGFP showed higher colocalization with ER marker RFP-KDEL as

compared to eGFP empty vector control, strengthening the fact that SteA possibly associate with

ER (figS3A, S3B). Overall, this data suggests that SteA might help in making contact sites with

ER and further recruiting ER onto the SCV. These contact sites can be used as a stealthy approach

by *Salmonella* to recruit membrane from ER, and along with other *Salmonella* effector proteins,

facilitating SCV fission.

STM*ΔsteA* shows a defect in proliferation and pathogenicity *in vivo*

Previous reports from our lab suggested that *Salmonella* utilizes the strategy of SCV division to

increase the number of SCV per cell, so it becomes difficult for the cellular system to eradicate the

growing SCV [14]. Since *steA* mutants form a big vacuole inside the cell, we asked how the mutant

strain proliferates in the cell. As compared to STM WT and STM*ΔsteA:steA* (complemented

strain), STM*ΔsteA* proliferated significantly less in the HeLa cell line. (fig4A). Less proliferation

of mutant strain suggests that those bulky vacuoles might be targeted by cellular machinery for

degradation. Quantification of vacuole size suggests that the diameter of SCV formed by

STMΔsteA is around 6.9±3.13µm whereas SCV of STM WT is 0.4±0.11µm (fig4B, fig4C). Using

LC3B as a marker to detect the extent of autophagy, we observed that LC3B colocalized more with

SCV of *steA* mutants as compared to SCV of STM WT at 16h post-infection (fig4D, fig4E).

Actively proliferating *Salmonella* also reduces the number of lysosomes inside the cell to escape

its fusion with acidic lysosomes [14]. In STMΔsteA infected cells, we observed a higher intensity

of Lamp1 similar to uninfected cells, suggesting that STMΔsteA is unable to change the lysosomal

dynamics as opposed to STM WT. A higher flux of lysosomal proteins, along with the recruitment

of LC3B, led to less proliferation of the STMΔsteA inside the cell (figS4A, figS4B). Next, we were

interested in understanding the pathogenicity of STMΔsteA *in vivo*. For this, we intraperitoneally

injected 10^3 CFU of *Salmonella* per C57BL/6 mice, and 3rd-day post-infection mice were

euthanized, and organs such as liver and spleen were harvested to observe organ burden. Our data

suggest that 3rd-day post-infection, STMΔsteA shows less colonization in the spleen and liver as

compared to STM WT and STMΔsteA:steA (fig 4G, fig4H). To further understand the

pathogenicity of the mutant strain, we orally gavaged C57BL/6 mice with 10^7 CFU of STM WT,

STMΔsteA, and STMΔsteA:steA, and 5 days post-infection, mice were euthanized, and organs such

as liver, spleen, and mesenteric lymph nodes were harvested since they act as a secondary site of

infection (fig4J-4L). We observed no significant change in the colonization of bacteria in these

organs, suggesting STMΔsteA doesn't lose its capability to cross the gut barrier and colonize and

disseminate in distal organs and blood if injected through the oral route (figS4C). However, upon

performing a survival assay of mice upon infection through oral gavage, we noticed the delayed

onset of mice death in the STMΔsteA infected cohort (fig4M), where STM WT infected mice

started succumbing to infection from 5th day post-infection, STMΔsteA infected cohort started

succumbing to infection at 7th day post-infection, and at 8th-day post-infection, all the mice

succumbed to death. Taken together, this data suggests that if the fission of SCV is altered,

Salmonella resides as multiple bacteria in one big vacuole and proliferates less. Those big/bulky

vacuoles are further targeted by autophagic machinery and cleared by active lysosomes in the cell

culture system. During *in vivo* infection through the intraperitoneal route, STM*ΔsteA* shows

defects in colonization in the spleen and liver and also affects the initial survival rate of mice,

suggesting that SteA acts as an essential effector for pathogenicity in mice.

Discussion:

Bacterial pathogens invade several cell types and reside in endocytic or phagocytic vacuoles [47].

Cell uses their extraordinary machinery to clear the pathogens, but by using several effector

proteins, bacterial pathogens modulate cellular machinery and prolong their survival. A few

strategies that several bacterial pathogens utilize are: modification of the membrane of the

pathogen-containing vacuole, blocking the activation of xenophagy, and escaping its fusion with

acidic lysosome [48], [49], [50], [51], [52], [53], [54]. Bacterial pathogens such as *Legionella,*

Brucella, Chlamydia, and *Salmonella* are known to reside in vacuoles, but the morphology and

membrane source of the vacuole are quite different. *Legionella* and *Chlamydia* reside in a spacious

vacuole that comprises multiple bacteria in a single vacuole [55], contrary *Salmonella* resides in a

tight-fitting vacuole as a single bacterium per vacuole [14]. The biogenesis of the SCV membrane

is not very well known, but proteome analysis of SCV suggests that the SCV membrane is highly

modified and contains proteins from the lysosome, Golgi, and ER [12]. Several effectors of

Salmonella help in recruiting membrane components, inhibiting the transport of M6PR, and

making contact sites with organelles for its proliferation [56], [57], [58]. Our results suggest that

Salmonella infection induces ER stress inside the cell and activates the UPR. A few ways by which UPR helps in alleviating stress are 1) by increasing the protein folding capacity of cells and 2) by expanding the ER. Our data suggests that wild-type STM infection leads to UPR activation while STM*ΔssaV* is unable to activate the UPR arm. This suggests that bacterial effectors are crucial for initiating ER stress, as it was evident as early as 2h post-infection. Our studies support the previous findings, which suggest *Salmonella*-induced ER stress is crucial for bacterial proliferation [59], [60], [61]. Further, UPR activation leads to an increase in phospholipid production [40], which could be useful for the bacteria as growing SCV requires a continuous source of membrane. *Salmonella*-infected cells show an increase in expression of *Reticulon-4a* and *Climp63* with an increase in tubular ER, suggesting this is an active strategy employed by bacteria for its survival. Exogenously overexpression of Rtn4a provides an advantage in the proliferation of *Salmonella*; however, reduction in ER tubules by knockdown of Rtn4a or overexpression of ER sheets using Climp63 negatively affects the bacterial proliferation and SCV number inside the cell, indicating ER tubules plays a crucial role in bacterial proliferation. The intricate balance of ER sheets and tubules has been recently shown to affect the dynamics of membrane-less organelle [42]. Our study also highlights the importance of ER in the successful proliferation of *Salmonella*. Our live cell data showed the recruitment of ER tubules at the center of dividing SCV. This was evident in HeLa as well as RAW 264.7 macrophage cell lines, suggesting that ER tubule recruitment marks the division site of SCV and leads to a single bacterium per vacuole. ER is very well known for marking the division site of organelles such as mitochondria, membrane-less organelle, and endosomes [18], [42]. Our study suggests that *Salmonella* hijacks the strategy of ER recruitment for its survival and SCV division. We have identified the role of bacterial effector SteA in maintaining the contact of SCV with ER in the formation of possible membrane contact sites.

Earlier studies have shown the presence of SteA on SCV and its colocalization with Golgi [62],

[63], [64]. In our study, we have partially observed a similar phenotype and have observed a new

role of SteA in interaction with ER. Inside the infected cell, the expression of SteA is highly

regulated as it can be mostly seen around the bacteria onto SCV. To investigate the role of SteA in

making contact sites, we looked for the colocalization of ER with SCV of STM WT and

STM*ΔsteA*. We observed the difference in colocalization of ER with SCV, highlighting the

importance of *the Salmonella* effector SteA in mediating the contact site between SCV and ER.

Bacterial pathogens such as *Chlamydia* use its effector protein for forming contacts of its inclusion

bodies with ER [25] and it also targets host cellular protein to facilitate this interaction [65].

Salmonella effectors SseJ and SseL have previously been reported to recruit OSBP1 onto SCV,

which helps in making contact with VapA and VapB (conserved ER proteins known for tethering

of ER with cellular organelles) and maintain SCV vacuolar integrity [66]. Our data suggests the

potential role of SteA in forming contact sites of SCV with ER as a strategy to facilitate its

proliferation.

The loss of function of SteA led to a bulky vacuole carrying multiple bacteria inside and with

defects in proliferation. Evidence from McQuate *et al*., also suggests the defect in the proliferation

of STM*ΔsteA* in epithelial cells as well as macrophages [62]. To understand the difference in

proliferation of STM*ΔsteA* compared to STM WT, we have observed SCV formed by STM*ΔsteA*

colocalizes more with autophagy marker LC3B, suggesting big bulky vacuoles are cleared by the

cellular system and hence lead to less proliferation inside the cell. STM WT also actively reduces

lysosome number inside the cell as an active strategy to block its fusion with the acidic lysosome

[14]. But in the case of STM*ΔsteA* infected cell, we observed a higher mean intensity of LAMP1,

which was similar to the intensity observed in uninfected cells, suggesting that somehow

STMΔsteA could not modulate lysosomal dynamics and, as a result, active lysosome could hinder the bacterial proliferation in cell line model. Similarly, Liu *et al*.,2021 have highlighted the importance of SteA in bacterial proliferation inside cells by impairing the fusion of SCV with lysosomes [67]. Taken together, this data suggests the importance of SteA in bacterial proliferation inside infected cells. STMΔsteA also showed defects in colonization in the spleen and liver if infected through the intraperitoneal route, suggesting SteA is crucial for *in vivo* pathogenicity. However, to our surprise, we did not observe any defect in the *in vivo* colonization of mouse organs such as the liver, spleen, MLN, and blood if infected through the oral gavage route. Survival analysis of mice suggests late onset of mice death in STMΔsteA infected cohort upon infection as compared to STM WT. We have also observed less weight reduction in STMΔsteA infected mice cohort as compared to STM WT, and this difference in weight reduction could provide an advantage in the initial survival rate of STMΔsteA infected mice. (figS4D). Alongside, Geddes *et al*., 2005 have reported the importance of SteA for efficient colonization in mouse spleen using the Balb/c mice model infecting via the intraperitoneal route [63]. We have also observed similar defects in the spleen using C57BL/6 mice. However, upon infection through oral gavage, there are no defects in the colonization of STMΔsteA in organs. This observation suggests that the route of infection can severely affect bacterial colonization and pathogenicity. Infection through the intraperitoneal route bypasses the gut-intestinal barrier, while during oral gavage, *Salmonella* faces several challenges, from the acidic pH of the stomach to several metabolites in the intestinal lumen. More studies are needed to understand this difference in colonization rate depending on the route of infection. Alongside, further investigations are needed to understand the vacuolar status of SCV during *in vivo* infection if infected through oral or intraperitoneal route to understand the difference

in pathogenicity. Taken together, our study suggests the importance of SteA in bacterial

pathogenicity and proliferation.

Our study provides new insights into how bacteria use a strategy of remodeling the ER dynamics

for its benefit. Domingues et al.,2014 and Chen et al.,2021 have highlighted the importance of

SteA for the active division of SCV [46], [68]. Our data supports their findings and suggests the

possible role of SteA in recruiting the ER and facilitating SCV fission. Interestingly, where wild-

type STM can easily divide along with SCV, while loss of function of SteA hampers 30% SCV

division with marked less colocalization with ER, suggesting there can be several other effector

proteins that might facilitate the contacts between ER and SCV. As reported by Chen et al.,2021

about the coordinated role of SteA, SopD2, and SifA in successful SCV division, our data explains

the newly identified role of SteA in the process by making contact sites with ER. Since these

effectors have coordinated roles, it will be interesting to understand how individual bacterial

effectors are crucial in the complete fission of SCV. Several host proteins have also been linked

with SCV division as the absence of those host proteins, such as PLEKHM1 and ESCRT-3, leads

to SCV carrying multiple bacteria [57], [69]. Within the cell, SCV acts as an autonomous organelle

that has acquired a vacuolar membrane from the host system, which also synchronously divides

along with bacterial division. The machinery regulating the SCV division could be difficult to

identify because these fission events are very transient in nature. As per literature evidence, both

host and bacterial effectors could be equally involved in this process to execute successful fission.

A complete knowledge of fission machinery can be used as a targeted approach to inhibit bacterial

proliferation during infection.

Acknowledgements

We acknowledge the Departmental and Divisional Confocal Microscopy Facility and Central

Animal Facility at IISc for providing us with an opportunity for experimentation. Mr. Sumitlal K

and Ms. Navya have helped in image acquisition in Confocal Microscopy.

Funding information

We express our heartfelt gratitude to the financial support from the Department of Biotechnology

(DBT), Ministry of Science and Technology; Department of Science and Technology (DST),

Ministry of Science and Technology. DC acknowledges DAE for the SRC Outstanding

Investigator Award and funds, ASTRA Chair Professorship, and TATA Innovation fellowship

funds. The authors jointly acknowledge the DBT-IISc Partnership Program. Infrastructure support

from ICMR (Center for Advanced Study in Molecular Medicine), DST (FIST), and UGC-CAS

(special assistance) is acknowledged. UC acknowledges the IISc fellowship. The funders had no

role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript. This work was also supported by the Department of Biotechnology

(BT/PR32489/BRB/10/1786/2019), DBT-NBACD (BT/HRD-NBA-NWB/38/2019-20) and

Science and Engineering Research Board (CRG/2019/000281) to S.R.G.S. DBT-JRF

(DBT/2016/IISc/717) to P.B.

Author Contributions

467 UC and DC have conceived the study and designed the experiments. UC has performed all the

468 experiments and participated in acquiring the data. UC and PB has analyzed the data, constructed

469 the figures, and wrote the original draft of the manuscript. UC. PB, DC and GSR have participated

470 in the proofreading and editing of the manuscript. DC has supervised the study and helped in fund

471 acquisition. All the authors have read and approved the manuscript.

References:

[1]   A. Haraga, M. B. Ohlson, and S. I. Miller, "Salmonellae interplay with host cells," Jan. 2008. doi: 10.1038/nrmicro1788.

[2]   F. W. Brenner, R. G. Villar, F. J. Angulo, R. Tauxe, and B. Swaminathan, "GUEST COMMENTARY Salmonella Nomenclature," 2000.

[3]   D. Pradhan and V. Devi Negi, "Stress-induced adaptations in Salmonella: A ground for shaping its pathogenesis," Dec. 01, 2019, *Elsevier GmbH*. doi: 10.1016/j.micres.2019.126311.

[4]   S. Ferguson *et al.*, "Coordinated Actions of Actin and BAR Proteins Upstream of Dynamin at Endocytic Clathrin-Coated Pits," *Dev Cell*, vol. 17, no. 6, pp. 811–822, Dec. 2009, doi: 10.1016/j.devcel.2009.11.005.

[5]   D. Zhou, L. M. Chen, L. Hernandez, S. B. Shears, and J. E. Galán, "A Salmonella inositol polyphosphatase acts in conjunction with other bacterial effectors to promote host cell actin cytoskeleton rearrangements and bacterial internalization," *Mol Microbiol*, vol. 39, no. 2, pp. 248–260, 2001, doi: 10.1046/j.1365-2958.2001.02230.x.

[6]   C. S. Bakshi, V. P. Singh, M. W. Wood, P. W. Jones, T. S. Wallis, and E. E. Galyov, "Identification of SopE2, a Salmonella Secreted Protein Which Is Highly Homologous to SopE and Involved in Bacterial Invasion of Epithelial Cells," 2000. [Online]. Available: http://genome.wustl.edu/gsc/bacterial/salmonella.shtml

[7]   W.-D. Hardt, L.-M. Chen, K. E. Schuebel, X. R. Bustelo, and J. E. Galá, "S. typhimurium Encodes an Activator of Rho GTPases that Induces Membrane Ruffling and Nuclear Responses in Host Cells," 1998.

[8]   K. C. Boddy *et al.*, "Salmonella effector SopD promotes plasma membrane scission by inhibiting Rab10," *Nat Commun*, vol. 12, no. 1, Dec. 2021, doi: 10.1038/s41467-021-24983-z.

[9]   J.-P. Gorvel and S. Méresse, "Maturation steps of the Salmonella-containing vacuole," 2001.

[10]   D. Drecktrah *et al.*, "Dynamic behavior of salmonella-induced membrane tubules in epithelial cells," *Traffic*, vol. 9, no. 12, pp. 2117–2129, 2008, doi: 10.1111/j.1600-0854.2008.00830.x.

[11]   Fran. Garcia-del Portillo, M. B. Zwick, K. Yin Leung, and B. Brett Finlay, "Salmonella induces the formation of filamentous structures containing lysosomal membrane glycoproteins in epithelial cells," 1993. [Online]. Available: https://www.pnas.org

502   [12]   J. C. Santos *et al.*, "The COPII complex and lysosomal VAMP7 determine intracellular Salmonella
503   localization and growth," *Cell Microbiol*, vol. 17, no. 12, pp. 1699–1720, Dec. 2015, doi:
504   10.1111/cmi.12475.

505   [13]   T. Reuter, S. Vorwerk, V. Liss, T. C. Chao, M. Hensel, and N. Hansmeier, "Proteomic Analysis of
506   Salmonella-modified Membranes Reveals Adaptations to Macrophage Hosts," *Molecular and
507   Cellular Proteomics*, vol. 19, no. 5, pp. 900–912, May 2020, doi: 10.1074/mcp.RA119.001841.

508   [14]   S. M. Eswarappa, V. D. Negi, S. Chakraborty, B. K. C. Sagar, and D. Chakravortty, "Division of the
509   Salmonella-containing vacuole and depletion of acidic lysosomes in Salmonella-infected host cells
510   are novel strategies of Salmonella enterica to avoid lysosomes," *Infect Immun*, vol. 78, no. 1, pp.
511   68–79, Jan. 2010, doi: 10.1128/IAI.00668-09.

512   [15]   N. A. Buchmeiert and F. Heffront, "Inhibition of Macrophage Phagosome-Lysosome Fusion by
513   Salmonella typhimurium," 1991. [Online]. Available: https://journals.asm.org/journal/iai

514   [16]   "R. Beuzon *et al.*, Salmonella maintains the integrity of its intracellular vacuole through the action
515   of SifA" The EMBO journal, 2000.

516   [17]   D. W. Holden, "Trafficking of the Salmonella vacuole in macrophages," 2002. doi: 10.1034/j.1600-
517   0854.2002.030301.x.

518   [18]   Friedman *et al.*, ER tubules mark site of mitochondrial division, 2011, Science

519   [19]   R. Fernández-Busnadiego, Y. Saheki, and P. De Camilli, "Three-dimensional architecture of
520   extended synaptotagmin-mediated endoplasmic reticulum-plasma membrane contact sites," *Proc
521   Natl Acad Sci U S A*, vol. 112, no. 16, pp. E2004–E2013, Apr. 2015, doi: 10.1073/pnas.1503191112.

522   [20]   Y. C. Wong, D. Ysselstein, and D. Krainc, "Mitochondria-lysosome contacts regulate mitochondrial
523   fission via RAB7 GTP hydrolysis," *Nature*, vol. 554, no. 7692, pp. 382–386, Feb. 2018, doi:
524   10.1038/nature25486.

525   [21]   H.-J. Wang, G. Guay, L. Pogan, R. Sauvé, and I. R. Nabi, "JCB Report Calcium Regulates the
526   Association between Mitochondria and a Smooth Subdomain of the Endoplasmic Reticulum,"
527   2000. [Online]. Available: http://www.jcb.org

528   [22]   A. A. Rowland, P. J. Chitwood, M. J. Phillips, and G. K. Voeltz, "ER contact sites define the position
529   and timing of endosome fission," *Cell*, vol. 159, no. 5, pp. 1027–1041, Nov. 2014, doi:
530   10.1016/j.cell.2014.10.023.

531   [23]   W. K. Ji, R. Chakrabarti, X. Fan, L. Schoenfeld, S. Strack, and H. N. Higgs, "Receptor-mediated Drp1
532   oligomerization on endoplasmic reticulum," *Receptor-mediated Drp1 oligomerization on
533   endoplasmic reticulum*, vol. 216, no. 12, p. 190538, 2017, doi: 10.1101/190538.

534   [24]   E. Smirnova, L. Griparic, D.-L. Shurland, and A. M. Van Der Bliek, "Dynamin-related Protein Drp1 Is
535   Required for Mitochondrial Division in Mammalian Cells," 2001.

[25] Rebecca Stanhope et al., IncV, a FFAT motif containing Chlamydia protein, tethers the
endoplasmic reticulum to the pathogen containing vacuole, 2017, PNAS

[26] G. Boncompain, C. Müller, V. Meas-Yedid, P. Schmitt-Kopplin, P. B. Lazarow, and A. Subtil, "The
intracellular bacteria Chlamydia hijack peroxisomes and utilize their enzymatic capacity to
produce bacteria-specific phospholipids," *PLoS One*, vol. 9, no. 1, Jan. 2014, doi:
10.1371/journal.pone.0086196.

[27] J. Sedzicki *et al.*, "3D correlative electron microscopy reveals continuity of Brucellacontaining
vacuoles with the endoplasmic reticulum," *J Cell Sci*, vol. 131, no. 4, Feb. 2018, doi:
10.1242/jcs.210799.

[28] M. Li, I. Tripathi-Giesgen, B. A. Schulman, W. Baumeister, and F. Wilfling, "In situ snapshots along
a mammalian selective autophagy pathway," *Proc Natl Acad Sci U S A*, vol. 120, no. 12, Mar. 2023,
doi: 10.1073/pnas.2221712120.

[29] K. A. Datsenko and B. L. Wanner, "One-step inactivation of chromosomal genes in Escherichia coli
K-12 using PCR products," 2000. [Online]. Available: www.pnas.orgcgidoi10.1073pnas.120163297

[30] S. Wang, H. Tukachinsky, F. B. Romano, and T. A. Rapoport, "Cooperation of the ER-shaping
proteins atlastin, lunapark, and reticulons to generate a tubular membrane network," 2016, doi:
10.7554/eLife.18605.001.

[31] Y. Shibata *et al.*, "The reticulon and Dp1/Yop1p proteins form immobile oligomers in the tubular
endoplasmic reticulum," *Journal of Biological Chemistry*, vol. 283, no. 27, pp. 18892–18904, Jul.
2008, doi: 10.1074/jbc.M800986200.

[32] N. Zurek, L. Sparks, and G. Voeltz, "Reticulon short hairpin transmembrane domains are used to
shape ER tubules," *Traffic*, vol. 12, no. 1, pp. 28–41, Jan. 2011, doi: 10.1111/j.1600-
0854.2010.01134.x.

[33] V. Singh, C. Erady, and N. Balasubramanian, "Cell-matrix adhesion controls Golgi organization and
function through Arf1 activation in anchorage-dependent cells," *J Cell Sci*, vol. 131, no. 16, Aug.
2018, doi: 10.1242/jcs.215855.

[34] R. Marwaha, S. B. Arya, D. Jagga, H. Kaur, A. Tuli, and M. Sharma, "The Rab7 effector PLE KHM1
binds Arl8b to promote cargo traffic to lysosomes," *Journal of Cell Biology*, vol. 216, no. 4, pp.
1051–1070, Apr. 2017, doi: 10.1083/jcb.201607085.

[35] Y. Adachi *et al.*, "Drp1 Tubulates the ER in a GTPase-Independent Manner," *Mol Cell*, vol. 80, no. 4,
pp. 621-632.e6, Nov. 2020, doi: 10.1016/j.molcel.2020.10.013.

[36] M. H. Alshareef, E. L. Hartland, and K. McCaffrey, "Effectors targeting the unfolded protein
response during intracellular bacterial infection," Apr. 01, 2021, *MDPI AG*. doi:
10.3390/microorganisms9040705.

[37]   M. Cirone, "Er stress, upr activation and the inflammatory response to viral infection," *Viruses*,
571   vol. 13, no. 5, May 2021, doi: 10.3390/v13050798.

572   [38]   K. Yamamoto *et al.*, "Transcriptional Induction of Mammalian ER Quality Control Proteins Is
573   Mediated by Single or Combined Action of ATF6α and XBP1," *Dev Cell*, vol. 13, no. 3, pp. 365–376,
574   Sep. 2007, doi: 10.1016/j.devcel.2007.07.018.

575   [39]   H. Yoshida *et al.*, "ATF6 Activated by Proteolysis Binds in the Presence of NF-Y (CBF) Directly to the
576   cis-Acting Element Responsible for the Mammalian Unfolded Protein Response," 2000.

577   [40]   R. Sriburi, S. Jackowski, K. Mori, and J. W. Brewer, "XBP1: A link between the unfolded protein
578   response, lipid biosynthesis, and biogenesis of the endoplasmic reticulum," *Journal of Cell
579   Biology*, vol. 167, no. 1, pp. 35–41, Oct. 2004, doi: 10.1083/jcb.200406136.

580   [41]   H. Wu, P. Carvalho, and G. K. Voeltz, "Here, there, and everywhere: The importance of ER
581   membrane contact sites," *Science (1979)*, vol. 361, no. 6401, Aug. 2018, doi:
582   10.1126/science.aan5835.

583   [42]   J. E. Lee, P. I. Cathey, H. Wu, R. Parker, and G. K. Voeltz, "Endoplasmic reticulum contact sites
584   regulate the dynamics of membraneless organelles," *Science (1979)*, vol. 367, no. 6477, Jan. 2020,
585   doi: 10.1126/science.aay7108.

586   [43]   G. K. Voeltz, W. A. Prinz, Y. Shibata, J. M. Rist, and T. A. Rapoport, "A Class of Membrane Proteins
587   Shaping the Tubular Endoplasmic Reticulum," *Cell*, vol. 124, no. 3, pp. 573–586, Feb. 2006, doi:
588   10.1016/J.CELL.2005.11.047.

589   [44]   M. J. Hoyer *et al.*, "A Novel Class of ER Membrane Proteins Regulates ER-Associated Endosome
590   Fission," *Cell*, vol. 175, no. 1, pp. 254-265.e14, Sep. 2018, doi: 10.1016/j.cell.2018.08.030.

591   [45]   W.-K. Ji, A. L. Hatch, R. A. Merrill, S. Strack, and H. N. Higgs, "Actin filaments target the oligomeric
592   maturation of the dynamin GTPase Drp1 to mitochondrial fission sites", doi:
593   10.7554/eLife.11553.001.

594   [46]   L. Domingues, D. W. Holden, and L. J. Mota, "The Salmonella effector SteA contributes to the
595   control of membrane dynamics of Salmonella-containing vacuoles," *Infect Immun*, vol. 82, no. 7,
596   pp. 2923–2934, 2014, doi: 10.1128/IAI.01385-13.

597   [47]   E. D. R. Case, J. A. Smith, T. A. Ficht, J. E. Samuel, and P. De Figueiredo, "Space: A Final Frontier for
598   Vacuolar Pathogens," May 01, 2016, *Blackwell Munksgaard*. doi: 10.1111/tra.12382.

599   [48]   C. L. Birmingham, A. C. Smith, M. A. Bakowski, T. Yoshimori, and J. H. Brumell, "Autophagy
600   controls Salmonella infection in response to damage to the Salmonella-containing vacuole,"
601   *Journal of Biological Chemistry*, vol. 281, no. 16, pp. 11374–11383, Apr. 2006, doi:
602   10.1074/jbc.M509157200.

603   [49]   Tilney *et al.*, How the parasitic bacterium Legionella pneumophila modifies its phagosome and
604   transforms it into rough ER implications for conversion of plasma membrane to the ER
605   membrane, 2001, Journal of Cell Science.

606   [50]   J. M. Kimmey and C. L. Stallings, "Bacterial Pathogens versus Autophagy: Implications for
607   Therapeutic Interventions," Dec. 01, 2016, *Elsevier Ltd*. doi: 10.1016/j.molmed.2016.10.008.

608   [51]   C. Pujol *et al.*, "Yersinia pestis can reside in autophagosomes and avoid xenophagy in murine
609   macrophages by preventing vacuole acidification," *Infect Immun*, vol. 77, no. 6, pp. 2251–2261,
610   Jun. 2009, doi: 10.1128/IAI.00068-09.

611   [52]   K. A. Rzomp, L. D. Scholtes, B. J. Briggs, G. R. Whittaker, and M. A. Scidmore, "Rab GTPases are
612   recruited to chlamydial inclusions in both a species-dependent and species-independent
613   manner," *Infect Immun*, vol. 71, no. 10, pp. 5855–5870, Oct. 2003, doi: 10.1128/IAI.71.10.5855-
614   5870.2003.

615   [53]   F. S. Mesquita, M. Thomas, M. Sachse, A. J. M. Santos, R. Figueira, and D. W. Holden, "The
616   Salmonella deubiquitinase SseI inhibits selective autophagy of cytosolic aggregates," *PLoS Pathog*,
617   vol. 8, no. 6, Jun. 2012, doi: 10.1371/journal.ppat.1002743.

618   [54]   L. Zhou *et al.*, "Salmonella spvC Gene Inhibits Autophagy of Host Cells and Suppresses NLRP3 as
619   Well as NLRC4," *Front Immunol*, vol. 12, Jul. 2021, doi: 10.3389/fimmu.2021.639019.

620   [55]   R. J. Bastidas, C. A. Elwell, J. N. Engel, and R. H. Valdivia, "Chlamydial intracellular survival
621   strategies," *Cold Spring Harb Perspect Med*, vol. 3, no. 5, May 2013, doi:
622   10.1101/cshperspect.a010256.

623   [56]   K. McGourty, T. L. Thurston, S. A. Matthews, L. Pinaud, L. J. Mota, and D. W. Holden, "Salmonella
624   inhibits retrograde trafficking of mannose-6-phosphate receptors and lysosome function," *Science*
625   *(1979)*, vol. 338, no. 6109, pp. 963–967, Nov. 2012, doi: 10.1126/science.1227037.

626   [57]   D. G. McEwan *et al.*, "PLEKHM1 regulates salmonella-containing vacuole biogenesis and
627   infection," *Cell Host Microbe*, vol. 17, no. 1, pp. 58–71, Jan. 2015, doi:
628   10.1016/j.chom.2014.11.011.

629   [58]   A. Sindhwani, S. B. Arya, H. Kaur, D. Jagga, A. Tuli, and M. Sharma, "Salmonella exploits the host
630   endolysosomal tethering factor HOPS complex to promote its intravacuolar replication," *PLoS*
631   *Pathog*, vol. 13, no. 10, Oct. 2017, doi: 10.1371/journal.ppat.1006700.

632   [59]   J. Bernal-Bayard, E. Cardenal-Muñoz, and F. Ramos-Morales, "The salmonella type III secretion
633   effector, Salmonella Leucine-rich Repeat Protein (SlrP), targets the human chaperone ERdj3,"
634   *Journal of Biological Chemistry*, vol. 285, no. 21, pp. 16360–16368, May 2010, doi:
635   10.1074/jbc.M110.100669.

[60] A. N. Antoniou *et al.*, "Salmonella exploits HLA-B27 and host unfolded protein responses to
promote intracellular replication," *Ann Rheum Dis*, vol. 78, no. 1, pp. 74–82, Jan. 2019, doi:
10.1136/annrheumdis-2018-213532.

[61] J. Kriston-Vizi *et al.*, "Salmonella Exhibit Altered Cellular Localization in the Presence of HLA-B27
and Codistribute with Endo-Reticular Membrane," *J Immunol Res*, vol. 2022, 2022, doi:
10.1155/2022/9493019.

[62] S. E. McQuate *et al.*, "Long-term live-cell imaging reveals new roles for Salmonella effector
proteins SseG and SteA," *Cell Microbiol*, vol. 19, no. 1, Jan. 2017, doi: 10.1111/cmi.12641.

[63] K. Geddes, M. Worley, G. Niemann, and F. Heffron, "Identification of new secreted effectors in
Salmonella enterica serovar typhimurium," *Infect Immun*, vol. 73, no. 10, pp. 6260–6271, Oct.
2005, doi: 10.1128/IAI.73.10.6260-6271.2005.

[64] L. Domingues *et al.*, "The Salmonella effector SteA binds phosphatidylinositol 4-phosphate for
subcellular targeting within host cells," *Cell Microbiol*, vol. 18, no. 7, pp. 949–969, Jul. 2016, doi:
10.1111/cmi.12558.

[65] I. Derré, R. Swiss, and H. Agaisse, "The lipid transfer protein CERT interacts with the Chlamydia
inclusion protein IncD and participates to ER-Chlamydia inclusion membrane contact sites," *PLoS
Pathog*, vol. 7, no. 6, Jun. 2011, doi: 10.1371/journal.ppat.1002092.

[66] A. M. Kolodziejek *et al.*, "Salmonella Translocated Effectors Recruit OSBP1 to the Phagosome to
Promote Vacuolar Membrane Integrity," *Cell Rep*, vol. 27, no. 7, pp. 2147-2156.e5, May 2019, doi:
10.1016/j.celrep.2019.04.021.

[67] K. Liu, L. Kong, D. B. Graham, K. L. Carey, and R. J. Xavier, "SAC1 regulates autophagosomal
phosphatidylinositol-4-phosphate for xenophagy-directed bacterial clearance," *Cell Rep*, vol. 36,
no. 4, Jul. 2021, doi: 10.1016/j.celrep.2021.109434.

[68] D. Chen *et al.*, "Systematic reconstruction of an effector-gene network reveals determinants of
Salmonella cellular and tissue tropism," *Cell Host Microbe*, vol. 29, no. 10, pp. 1531-1544.e9, Oct.
2021, doi: 10.1016/j.chom.2021.08.012.

[69] V. Göser, A. Kehl, J. Röder, and M. Hensel, "Role of the ESCRT-III complex in controlling integrity of
the Salmonella-containing vacuole," *Cell Microbiol*, vol. 22, no. 6, Jun. 2020, doi:
10.1111/cmi.13176.

Legends to the Figures

Fig:1-*Salmonella* infection leads to the activation of UPR and expansion of the

Endoplasmic Reticulum

(A) Representative confocal microscopic images of uninfected and infected HeLa cells

with STM WT mCh or STM*ΔssaV* mCh at MOI 10, 16h post-infection and immuno-

stained against ATF4, LAMP1 and DAPI. (B) Represents the quantification of Cell Total

Corrected Fluorescence (CTCF) intensity of ATF4 inside the nucleus, data is

representative of n=100-120 cells, from 4 independent experiments, mean±SD (C)

Representative confocal microscopic images of uninfected and infected HeLa cells with

STM WT mCh or STM*ΔssaV* mCh at MOI 10, 16h post infection and immuno-stained

against ATF6, LAMP1 and DAPI. (D) Represents the quantification of Cell Total

Corrected Fluorescence (CTCF) intensity of ATF6 inside the nucleus, data is

representative of n=60-90 cells from 3 independent experiments, mean±SD (E)

Representative confocal microscopic images of uninfected and infected HeLa cells with

STM WT mCh or STM*ΔssaV* mCh at MOI 10, 16-hour post-infection and immuno-

stained against XBP1, LAMP1, and DAPI (F) Represents the quantification of Cell Total

Corrected Fluorescence (CTCF) intensity of XBP1 inside the nucleus, data is

representative of n=60-90 cells from 3 independent experiments, mean±SD (G)

Quantitative real-time PCR of *Rtn4a* (H) and *Climp63*, N=3, n=3, mean±SEM (I)

Confocal microscopic images of Un-infected and infected HeLa cell showing ER

expansion (J) Represents the quantification of tubular ER, data is representative of n=100-

150 cells from 4 independent experiments, mean±SD. Student's t-test was used to analyze

the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig-2: Expansion of ER tubules facilitates SCV proliferation and its division

(A) Represents the ICSA upon overexpression of Rtn4a, data is representative of N=3,

n=3 mean± SEM (B) ICSA upon Rtn4a knockdown in HeLa cells, data is representative of

N=3, n=3 mean± SEM (C) Representative confocal microscopy images of SCV number

per cell upon overexpression of mCh-EV, sec61β-mCh, and mCh-Climp63, 16h post

infection (D) Quantification of the number of SCV per cell, 16h post infection, data is

representative of n=40-60 cells from 2 independent experiments (E) Snapshot of live cell

imaging of *Salmonella* infected HeLa cells, White arrowhead represent the ER overlap at

dividing bacteria. White line along the length of bacteria represent the portion used for

line scan. (F) Showing the line scan of images. Blue arrows represent the snapshot for

which the line scan is analyzed (snapshots 3 and 4) (G) Quantification of SCV fission

events marked by ER, data is representative of n=11 SCV fission events from 4

independent experiments (H) Confocal microscopic images representing SCV fission

events marked by ER, at center or adjacent (I) Quantification of fission events in fixed

images marked by ER, data is representative of n=49 SCV fission events from 4

independent experiments. White arrowhead represent the ER overlap at dividing SCV.

Student's t-test was used to analyze the data ****<0.0001, ***<0.001, **<0.01, *<0.05.

Fig-3: *Salmonella* translocated effector SteA is crucial for maintaining SCV contacts

with ER, resulting in a single bacterium per vacuole

(A) Represents confocal microscopy images showing SCV formed by STM WT and

STM*ΔsteA,* 16h post-infection (B) Quantification of the number of cells showing multiple

bacteria in one vacuole, 16h post-infection, data is representative of n=120-150 cells from

3 independent experiments, mean±SEM (C) Representative confocal image of infected

HeLa cell showing contact between SteA-HA and ER, arrowheads represent interaction

between ER and SteA-HA (D) Confocal images represents the colocalization of ER with

SCV formed by either STM WT or STM*ΔsteA,* 16h post-infection (E) Quantification of

colocalization coefficient of ER marker RFP KDEL with SCV marker LAMP1, Data is

representative of n=80-100 cells from 2 independent experiments, 16h post infection,

mean±SD. Student's t test was used to analyse the data ****<0.0001, ***<0.001,

**<0.01, *<0.05.

Fig-4: STM*ΔsteA* shows a defect in proliferation and pathogenicity *in vivo*

(A) Intracellular survival assay of STM WT, STM*ΔsteA*, and STM*ΔsteA:steA* in HeLa

cells, data is representative of N=3,n=3, mean±SEM. (B) Line representation to quantify

SCV diameter where red line indicates the length of SCV used for quantification (C)

Quantification of SCV diameter, data is representative of 100-150 cells from 3

independent experiments, 16h post-infection, mean±SD (D) Representative microscopy

images showing colocalization of LC3B with SCV, 16h post-infection(E) Quantification

of colocalization coefficient of LC3B with SCV, data is representative of n=100-130 cells

from 3 independent experiments, mean±SD (F) outline of experimental procedure of mice

infection through intraperitoneal route (G) CFU enumeration in Spleen and (H) liver, data

is representative of N=2, n=5 mice per cohort (I) Outline of experimental procedure of

mice infection through oral gavage and dissected on day 5 post infection (J) CFU

enumeration in organs like Spleen (K) Liver (L) and MLN, data is representative of N=3,

n=5 mice per cohort, mean±SD (M) Survival curve of cohort of mice infected with STM

WT and STM*ΔsteA,* n=8 mice per cohort. Mann-Whitney test was performed to obtain

statistical significance for *in vivo* data. Student's t-test was used to analyze the data from

fig 4A-4E. ****<0.0001, ***<0.001, **<0.01, *<0.05.

Graphical Abstract

(A) SCV makes contact sites with ER using effector SteA

(B) ER marks the SCV division site

 • *Salmonella* infection induces ER stress

 • SCV of STM WT displaying SteA onto SCV and one bacterium per vacuole

 • SCV of STM*ΔsteA*

 • SCV comprises multiple bacteria in single vacuole

(A) Less contact sites between ER and SCV in absence of SteA

(B) STM*ΔsteA* resides as multiple bacteria in one vacuole

Fig:1 *Salmonella* infection leads to the activation of UPR and expansion of the Endoplasmic Reticulum

G

Fold change of *Rtn4a* over β-actin

*

*

UI 2h 6h 16h

Post Infection

H

Fold change of *Climp63* over β-actin

*

UI 2h 6h 16h

Post Infection

I

LAMP1
RFP KDEL
Uninfected

10 µm

1   5 µm

RFP KDEL   5 µm

2   5 µm

5 µm

LAMP1
RFP KDEL
STM WT GFP

10 µm

1   5 µm

RFP KDEL   5 µm

2   5 µm

5 µm

J

Tubular ER (%)

***

UI   STM WT

Fig 2: Expansion of ER tubules facilitates SCV proliferation and its division

H

I

% of SCV fission event

- No ER (29.2±8.4%)
- ER crossing at adjacent (14±9.07%)
- ER crossing at center (56.71±6.61%)

Fig 3: *Salmonella* translocated effector steA is crucial for maintaining SCV contacts with ER resulting in a single bacterium per vacuole state

Fig 4: STM*ΔsteA* shows a defect in proliferation and pathogenicity *in vivo*